

## 1 General comments

In terms of significance, the data set presented and the corresponding article can be rated “1” (excellent) because (i) I regard the data as useful for future investigations of the thermal field, the mechanical state and the thermomechanical behavior of the crust all across Germany; (ii) the presented model cannot be built on a routine basis but relies on the efficiency of the new method; (iii) the data set is complete in the sense that a much larger number of input models has been combined than in previous models and the entire crust all over Germany is covered.

To improve the quality of the final model – now rated “2” – a more detailed description of the criteria (i) for choosing the 27 input models (while potentially disregarding alternatives) and (ii) for giving different priorities to spatially overlapping input models is required.

Concerning the presentation quality, the contribution can also be rated “2” since (i) the manuscript is clear and well structured, (ii) the figures additionally provided to illustrate all subsets of the data are very useful and efficient, and (iii) the manuscript in combination with the landing page for the dataset enables users to understand and (re-)use the data set. Some directions for improvements are given in the following, with specific comments provided separately for the manuscript (2.1) and for the data set (2.2).

## 2 Specific comments

### 2.1 Manuscript

The language of the manuscript is mostly consistent and precise, but an overall check should be done, e.g., (i) regarding comma placement, (ii) changing tenses when describing the work done (e.g., “we decided...”, “we use...”), and (iii) simple typos, some of which but certainly not all are listed below.

The length of the article is appropriate. Its structure is clear except for some statements in the results chapter that should be moved to the discussion (see below). The figures and tables are clear and help to follow the lines of thoughts.

The method applied to build the model has not been used before and is well described. The approach of keeping a minimum thickness of 0.1 meters wherever a model unit is present is indeed similar to an approach used before (e.g., Anikiev et al. 2019), though in the latter case the minimum thickness of 0.1 meters for a unit was applied to the whole lateral extent of the model (including locations where a model unit is in fact not existing). Please add this to the discussion of advantages of the modelling approach.

### Abstract

“[...]However, up to now only one attempt has been made to combine several of them to a Germany-wide model. We present a new Germany-wide 3D geological model [...]” → Already here it should be stated why a new model is required (New input data and models available? Higher resolution required?).

### 1 Introduction

Line 19: 3D subsurface models are introduced as “showing lithostratigraphic horizons”, which would restrict them to the domains above the crystalline basement. The presented model includes

crystalline crustal rocks as well. Their lateral juxtaposition would exclude them from a horizon / layer type model. Use “geological units” instead.

Line 22: The list of 3D models referred to seems to be arbitrary. What was the criterion for this choice of 3D models? If they have something particular in common, it should be stated.

Line 23: Suggestion: use “natural” instead of “undisturbed”. Temperatures can be regarded as “disturbed” by natural processes as groundwater flow.

Line 39: “In the following we first present [...]” → “In the following, we first shortly introduce [...]”

## *2 Model set up*

### *2.1 Data base*

Line 48: It remains unclear which criteria have been applied to choose the 27 models used for model building. Please explain and potentially comment on models that have been disregarded.

Line 54: better “...depending on the position and the corresponding input model at place”.

### *2.2 Model correlation*

This chapter raises an important question: how do the authors deal with inconsistent input information from the different models used? For example, how to deal with the same stratigraphic interface shown at different depth levels? As this point is described in the following chapter (2.3), but the question comes up here already, the reader should at least be referred to the later description.

### *2.3 Point set approach*

Line 106: “[...] which is tolerable for the model scale and resolution. The major advantages of the point set model approach [...]” These parts are actually evaluative statements which should be moved to and deepened in the chapter “4. Discussion”.

Line 100-111: “The order of projection is determined according to various criteria, e.g., year of publication, model resolution, etc.” → This is an important point worth to be described in more detail. Please describe the philosophy behind this approach in more detail. The example given is not enough for transferring this modelling approach to another region, respectively another modelling campaign. For example, does the criterion “year of publication” mean that younger models are always given higher priority than older models, even if an older one covers a larger area or gives a higher resolution? To clarify this, a figure illustrating the decision-making process as a workflow would be helpful. Then the method can be re-used and referred to more easily.

Line 122: “The biggest advantage of the point set model, however, [...]” → Again, this is part of an evaluation of the method and should be moved to the discussion.

In general, this chapter does not provide a detailed description of the computer programs developed to create an initial point set and multiply it several times for projecting it onto input data of various formats. How did the authors deal with input data of different formats?

We agree with the authors that an important advantage of the “point set approach” is that it can be used directly for the fast creation of discretized models with ApplePY (Ahlers et al. 2022; Ziegler et al. 2019) while no specific software is necessary. This should be emphasized more in the Discussion chapter.

### *3 Results*

#### *3.1 Stratigraphic correlation*

[no comments]

#### *3.2 Model units*

[no comments]

#### *3.3 Presentation of results*

Line 164: Please clarify what your definition of “high stratigraphic resolution models” is?

#### *3.4 Generation of a discretized model*

Line 174-175: Please refer to a more detailed description of the format of an Abaqus \*.inp file or provide the description here (as this would help re-using the approach).

##### *3.4.1 Worked example*

Line 187: “[...] whereby the element thickness increases with depth.” → Please explain why and – if this helps to understand the reason – also how the increase of element thickness was designed.

Fig. 6, caption: For readers that are new to the field, it could be interesting to learn which software has been used for the visualization of the model.

### *4. Discussion*

Line 232: “manageability of the final model” → Please clarify in which case(s) a model might not be manageable and in terms of what (computational times for the model production, visualization, ...?).

Line 237: “Overall, the loss of information and vertical uncertainties are acceptable, especially considering the size and purpose of the model.” → This statement is too general and thus requires some reasoning or at least a restriction to the applications intended by the authors.

Lines 248-271: These paragraphs include descriptions of results with some explanations of outcomes that could raise questions on the user’s side. It’s a list of examples the completeness level of which cannot be estimated by the reader. I suggest moving both paragraphs to chapter “3.3 Presentation of results”.

In general, the discussion chapter should be used to deepen some thoughts on (i) the advantages and disadvantages of the modelling approach and the (ii) possibilities and limitations of using the new model, in particular by comparison to some already existing ones. On the methodological side, which problems could be expected if one wants to generate a triangulated mesh from the original model to better represent units that taper out? In terms of the usage of the geological model envisaged by the authors (i.e., geomechanical-numerical modelling), which changes and possibly improvements in the results can be expected compared to previous models (Ahlers et al. 2021, 2022a)?

### *5. Conclusions*

The contents of this chapter actually is a summary of the manuscript and corresponding data set rather than a conclusion (the latter generally reflecting an opinion or a decision for action after considering all given information about something).

### Data availability

It would be a great service to the community to share not only the final model but also the scripts, respectively applications, for creating a point set, multiplying it and projecting it onto input data of various formats.

### Supplement

Supplementary material is provided as three tables (folder “essd-2025-320-supplement”). The 27 individual models used for building the new Germany 3D model are listed with their references in the file 2025\_05\_28\_Table\_S3.docx.

## 2.2 Data set

The data set is accessible via the given weblink (<https://tudatalib.ulb.tu-darmstadt.de/handle/tudatalib/4615>).

The data set is complete. The downloadable folder (Ahlers\_2025\_surfaces) contains 147 \*.xyz-files according to the number of interfaces that define the 146 model units. Files can be opened with standard text editors without any problems. It should be added in the “Beschreibung” that “negative Z-values are in meters below sea level”.

The downloadable folder “Ahlers\_2025\_units\_figures” contains very useful figures for each of the model 146 units, providing easily accessible information (\*.jpg format) about the spatial extent, geometry and input data distribution. Files can be opened with standard graphic viewers without any problems. The additional 11 files of combined plots are useful as well as they represent major stratigraphic units that have been differentiated also in previously published 3D models. The readme.txt file informs about the individual model units (their id’s) used to build the combined plots.

The error of interface depths resulting from implementing a minimum layer thickness of 0.1 m is shortly discussed in the article (4. Discussion). Otherwise, there is no quantification of the model uncertainties done.

The final structural model of Germany and adjacent areas is new in terms of its large extent combined with a relatively high geological and stratigraphic resolution. The model will certainly be useful in the future as a basis for running process simulations on various scales.

### Technical corrections

Line 57: ~~limits~~ → limitations

Line 58: ~~for all kind of~~ → for many types of

Line 65: Germany-wide

Line 132: either “stratigraphically independent” or “stratigraphy independent”

Line 162: indicates

Lines 184: the Netherlands

Line 268: shows