

Subsets of geostationary satellite data over international observing network sites for studying the diurnal dynamics of energy, carbon, and water cycles

5 Hirofumi Hashimoto^{1,2}, Weile Wang¹, Taejin Park^{1,3}, Sepideh Khajehei^{1,3}, Kazuhito Ichii⁴, Andrew R. Michaelis¹, Alberto Guzman^{1,2}, Ramakrishna R. Nemani³, Margaret Torn⁵, Koong Yi⁵, Ian G. Brosnan¹

¹ NASA Ames Research Center, Moffett Field, CA 94035, USA

²Department of Applied Environmental Science, California State University – Monterey Bay, Seaside, CA 93955, USA

³Bay Area Environmental Research Institute, Moffett Field, CA 94035, USA

⁴Center for Environmental Remote Sensing, Chiba University, Chiba-shi, Chiba 263-8522, Japan

10 ⁵Earth and Environmental Sciences Area, Lawrence Berkeley National Laboratory, Berkeley, CA 94705, USA

Correspondence to: Hirofumi Hashimoto (hirofumi.hashimoto-1@nasa.gov)

Abstract. The latest generation of geostationary satellites provide Earth observations similar to widely used polar orbiting sensors but at intervals as frequently as every 5-10 minutes, making them ideal for studying diurnal dynamics of land-atmosphere interactions. The NASA Earth Exchange (NEX) group created the GeoNEX datasets by collating data from several 15 geostationary platforms, including GOES-16/17/18, Himawari-8/9, and GK-2A, and placing them on a common grid to facilitate use by the Earth science community. Here, we document the GeoNEX Coincident Ground Observations dataset (GeCGO) for terrestrial ecosystems studies, and provide examples for its use. Currently, GeCGO provides GOES-16 Advanced Baseline Imager (ABI) data over a 10 x 10 km area surrounding 1586 network sites across Americas. GeCGO make it easy to 20 compare the time series of geostationary data with the diurnal ground observations including carbon/water fluxes and aerosol optical depth, and is extensible to other regions. We also developed GeoNEXTools to facilitate analyses that require both GeoNEX data and other NASA satellite data. The objectives of this paper are to introduce GeCGO and GeoNEXTools, and demonstrate their applications. First, we describe the details of GeCGO and GeoNEXTools. Second, we explain how GeCGO can be integrated with other satellite data. Finally, we showcase comparisons between GeCGO and observations from three ground-based networks.

25 1 Introduction

Satellites monitor the Earth's surface using sensors with a variety of spatial-spectral-temporal characteristics. Sensors on geostationary satellites have unique characteristics, including high frequency observations with constant view zenith angle. Geostationary satellite data have been regarded as less effective for monitoring the Earth's environment due to their low spatial resolution and wide spectral bands. However, the latest generation of geostationary satellites have advanced imaging sensors, 30 e.g. GOES-16/ Advanced Baseline Imager (ABI) and Himawari-8/ Himawari Advanced Imager (AHI), which can provide spatial resolution (i.e., 1 x 1 km) comparable with MODIS (Schmit et al., 2017). MODIS has been the most frequently used sensor for monitoring the global land surface for decades. The high temporal resolution data of the new geostationary satellites

enable us to scrutinize sub-daily/sub-hourly processes on the Earth's surface, which polar-orbiting sensors with low temporal resolution like MODIS are unable to capture (Xiao et al., 2021; Yi et al., 2024). Therefore, the new generation geostationary satellites have the potential to complement the polar orbiting sensors for global and continental-scale research.

Recent studies have demonstrated the effectiveness of using the new generation geostationary satellite data for terrestrial ecosystem modeling. For example, Li et al., (2023) used GOES ABI data to show that suppression of photosynthesis in the afternoon is caused by high vapor pressure deficits (VPD) in the **Western United States**. Hashimoto et al. (2021) used the greater number of clear sky observations from GOES ABI to analyse the leaf phenology over the Amazon and identify seasonal patterns in greenness (**removed**). Others have shown that ABI data can be used to estimate sub-daily Gross Primary Production (GPP) through a vegetation index (Khan et al., 2022), a machine learning technique (Stoy et al., 2024), or a light use efficiency model (Xiao et al., 2021). These efforts reveal the potential of using GEOS ABI for monitoring and modeling the dynamics of land surface vegetation. However, there remain difficulties for scientists in using geostationary satellite data due to their large volume, inconsistent file formats across sensors, and insufficient documentation or software to handle the data.

45




Figure 1: Flowchart showing how GeCGO is created. Geostationary satellite L1B data were used to produce the suite of GeoNEX datasets, then observations over the network site were extracted to create GeCGO (a-c). Users can download and analyze GeCGO data using GeoNEXTools for multi-sensor analysis (d), spatial analysis (e), and time series analysis (f).

50

Scientists faced similar challenges with MODIS data during the early days of the Earth Observing System (EOS) era. This led to the development of MODIS subset data (ORNL DAAC, 2017) and MODIS subset data have since been extensively used by the land surface monitoring and modeling communities. The NASA Earth Exchange (NEX) recognized the importance of data compactness from the EOS experience and leveraged the GeoNEX project (Wang et al., 2020) to create the GeoNEX

55 Coincident Ground Observations dataset (GeCGO) and facilitate the use of data from geostationary satellites. GeCGO is focused on widely monitored field and flux tower sites across the Americas and is accompanied by the GeoNEXTools to help users retrieve GeCGO data, similar to the manner in which MODISTools supported the retrieval of MODIS subset data. The data processing flow is summarized in Fig. 1 and is implemented with Ziggy, an automated processing software for Science Data Analysis Pipelines (Tenenbaum and Wohler, 2024).

60 The objectives of this paper are to introduce GeCGO and the GeoNEXTools, and demonstrate their applications. First, we describe the details of GeCGO and GeoNEXTools. Second, we explain how GeCGO can be integrated with other satellite data. Finally, we showcase comparisons between GeCGO and observations from three ground-based networks.

2 Description of the GeoNEX data, GeCGO, and GeoNEXTools

2.1 GeoNEX dataset

65 The GeoNEX data is a collection of land surface images collected by new generation geostationary weather satellites, including Himawari-8/9 Advanced Himawari Imager (AHI), GOES-16/17/18 ABI, Geo-KOMPSAT-2A (GK-2A) Advanced Meteorological Imager (AMI). Although intended for weather observations, the quality of the data from these satellites are now suitable for studying land surface dynamics. NEX has been producing geostationary data for land surface research communities from Level-1B full-disk scenes (Wang et al., 2020). The GeoNEX data are tiled into 6° by 6° with 0.005° - 0.02° 70 spatial resolutions in geographic projection. Each sensor covers a square region approximately $\pm 60^{\circ}$ from its nadir point to remove the edge pixels with the large view-angle. Tiles including only oceans are not processed. Because the GeoNEX data were made from full-disk images, the temporal resolution is between 10 and 15 minutes.

We first georectified Level-1B data using information provided by institutions that operate the geostationary satellites. Although each sensor has its own state-of-art on-board georeferencing system, there are still residual pixel shifts in each image. 75 For example, 1-pixel shifts for 500 m band data were observed almost every day in AHI data, while ABI shifts were well under ± 0.5 pixel (Wang et al., 2020). The Phase-only Correction (POC) method was employed by matching Shuttle Radar Topography Mission (SRTM) Digital Elevation Model (DEM) based coastlines with the geostationary satellite images (Takenaka et al., 2020). We also corrected the relief displacement caused by the large viewing angle and high-elevation terrain, which is critical for comparing the satellite images with ground observation data.

80 The lowest level GeoNEX data is Level-1G, which contains Top-Of-the-Atmosphere (TOA) reflectance and/or brightness temperature. The Level-2 data is surface reflectance, which is retrieved by applying the Multi-Angle Implementation of Atmospheric Correction (MAIAC) (Lyapustin et al., 2011). Other higher-level data are also available, i.e., Land Surface

Temperature (LST) (Jia et al., 2022) and solar radiation data (Li et al., 2023a). Currently, the Level-2 LST data is available only for North America (i.e., Canada, U.S., and Mexico). The GeoNEX data are provided in the HDF-EOS2 format and 85 available from the NASA Advanced Supercomputing Data Portal (<https://data.nas.nasa.gov/geonex>).

NOAA also provides the ABI land Level 2 products, such as TOA reflectance, TOA brightness temperature, Bidirectional Reflectance Distribution Function (BRDF), and fire classification (Losos et al., 2024). The differences between GeoNEX data and NOAA Level 2 products are that the GeoNEX data are (1) geo-corrected for each image, (2) reprojected to the geographic projection, (3) tiled for each area covering 6° by 6°, and (4) include international geostationary satellite data (see Wang et al., 90 2020 for more details).

2.2 GeCGO

To make it convenient for users to get time series of GeoNEX data where ground-based observations are collected, we created GeCGO by extracting observations from the GeoNEX data over 1586 field or flux tower sites in various observation networks.

To further facilitate its use, GeCGO is provided in same file format as the Oak Ridge National Laboratory (ORNL) Terrestrial 95 Ecology Subsetting & Visualization Services (TESViS) fixed sites subset data (<https://modis.ornl.gov/sites/>) (ORNL DAAC, 2017), formerly known as MODIS/VIIRS Subset Tools, which will be familiar to most scientists. Specifically, GeCGO data are available in Comma-Separated Values (CSV) and JavaScript Object Notation (JSON) formats for each year (see the example of Fig. 2 and Table 1 for metadata descriptions). The data values were organized as 17x17, 9x9, and 5x5 grids for 0.005°, 0.01°, and 0.02° resolution, respectively. For instance, the TOA band 1 reflectance data has 0.01° spatial resolution at

100 nadir, and thus its GeCGO data consists of 9x9 pixel value data. The data values represent sequences of pixels from northwest corner to southeast corner in row-major order (Fig. 2). The data are organized into directories named after the site IDs used in the TESViS Subset. Currently, we provide the Level-1G and Level-2 data within the GeCGO products, available at the NASA GeoNEX data portal (<https://data.nas.nasa.gov/gecko/data.php>). The available GeoNEX data are summarized in Table 2.

105




Figure 2. Example of JSON file and CSV file. The number in the 5x5 grid indicates the pixel position in the grid to clear the order in JSON and CSV file.

110 **Table 1. The metadata in the GeCGO JSON file.**

Key	Value
xllcorner	longitude at the lower left corner (degree)
yllcorner	latitude at the lower left corner (degree)
cellsize	pixel size in north-south and west-east direction (degree)
nrows	number of pixels in north-south direction
ncols	number of pixels in west-east direction
band	name of the band
units	physical unit of data values
latitude	latitude of the flux tower
longitude	longitude of the flux tower
subset	time in UTC and data values

Table 2. The product summary of the GeCGO. Each product has several bands with different spatial resolutions and grid sizes.

Product name	Band name	Spatial resolution (°)	Grid size
Level 1G	band 1-6 TOA reflectance	0.005-0.02	17x17, 9x9, 5x5
	band 7-16 TOA brightness temperature	0.02	5x5
Level 2	band 1-6 surface reflectance	0.005-0.02	17x17, 9x9, 5x5
	solar zenith angle	0.01	9x9
	solar azimuth angle	0.01	9x9
	status QA	0.01	9x9
	AOD	0.01	9x9
	land surface temperature	0.02	5x5
	solar radiation	0.01	9x9

115

GeCGO has the same ground network information as the TESViS Subset and currently includes 1,586 sites across 14 ground networks in the Americas. The GeCGO sites were entirely distributed over the continents in GOES-16 coverage (Fig. 3).
 120 Approximately 76% of the sites are in the AmeriFlux (Chu et al., 2023; Novick et al., 2018), AERONET (Aerosol Robotic Network) (Holben et al., 1998), and **PhenoCam** (Richardson et al., 2018a) networks (Table 3). We excluded the island AERONET sites where the island area is less than 1,000 km².

Table 3. Network names and number of sites of each network in the current GeCGO. Networks that had less than 10 sites were categorized in others. Each site was assigned only a single network based on TESViS data, though there were 125 many sites that were involved in multiple networks.

Network name	Number of stations
AMERIFLUX	540
AERONET	425
PHENOCAM	247
USCRN	170
NEON	54
FLUXNET Canada	22
FLUXNET	21
unaffiliated	18
FORESTGEO	17
URBANFLUXNET	13
LTER	12
others	47
Total	1586




Figure 3: Spatial map of locations in the GOES16 coverage for each network used in GeCGO. The colors of dots represent various network names. Networks with less than 10 sites were categorized as “others”.

There is another existing subset of geostationary satellite data products over AmeriFlux sites (Losos et al., 2024). This subset data provides half-hourly ABI fixed grid products for the AmeriFlux sites including TOA reflectance, surface reflectance, cloud mask, aerosol, solar radiation, etc. Those data were single-pixel time series derived mainly from NOAA high-level


products with the terrain correction. The ABI fixed grid products have different algorithms and procedures from GeCGO. For instance, GeoNEX and NOAA use different atmospheric correction (MAIAC (Wang et al., 2022) for GeoNEX and 6S (He et al., 2019) for NOAA) and geolocation correction algorithms (further orthorectification and geolocation correction in GeoNEX (Wang et al., 2020)). The different processing algorithms and procedures also make spatial resolution and time-step different.

There are many ongoing international efforts to develop geostationary satellite products for various applications. We believe that the subset datasets, including GeCGO and the ABI fixed grid products, available through global ground networks, provide ready-to-use hypertemporal Earth observations and inter-comparison data that can advance modeling and address important scientific questions.

2.3 GeoNEXTools

We developed GeoNEX Subset Tools, GeoNEXTools (<https://github.com/nasa/GeoNEXTools>), to facilitate downloading and manipulating GeoNEX data for specific ground observation sites and time ranges. Although the GeCGO data volume is smaller than the original GeoNEX full-disk and tiled data, handling GeCGO data remains challenging due to the length of the high-frequency time series. The MODIS science community solved a similar issue by developing MODISTools (The MODISTools package: an interface to the MODIS Land Products Subsets Web Services, 2024; Tuck et al., 2014), which is an open-source R package that helps users download and read the MODIS Subset data (<https://github.com/ropensci/MODISTools>). The MODISTools package has become one of the most widely used tools for handling TESViS Subset data. Since many scientists are accustomed to using MODISTools, we developed GeoNEXTools to provide similar functionality as MODISTools. The function names in GeoNEXTools are the same as MODISTools except for the prefix (e.g., `mt_<function name>` for MODISTools and `gt_<function name>` for GeoNEXTools). A full list of the function names and descriptions is provided in **Appendix A**.

Figure 4 illustrates how GeCGO and GeoNEXTools can be used to analyze the diurnal variation in biophysical and meteorological variables. The figure includes data from the Harvard Forest EMS Tower (US-Ha1) from July 23rd to August 5th, 2020 and clearly illustrates the diurnal variability in NDVI, LST, downward shortwave radiation, and AOD. Preliminary examination of the data reveals several other insights. A cloudy day, July 30th to August 1st, is clearly evident in the diurnal variation in LST, which is smaller than on the other days. Figure 4a shows inverse patterns in diurnal surface and TOA NDVI in clear days (July 23rd to July 26th), which indicates the importance of atmospheric correction study in diurnal cycle studies. The clear day NDVI calculated using TOA data show peak values at noon, while the diurnal variation of the surface NDVI exhibits a u-shaped curve. Even though each of the GeoNEX products contains a large volume of data, the GeoNEXTools significantly simplified the process of retrieving and examining single-point time series.



165

Figure 4: Time series from GeCGO at Harvard Forest EMS Tower (US-Ha1) from 2020-07-23 to 2020-08-05. (a) NDVI calculated from band 2 and band 3 at 10 minutes interval. Blue and purple dots were Top-Of-Atmosphere (TOA) (level 1G dataset) and surface (level 2 dataset) NDVI, respectively. (b) Land Surface Temperature (LST) at hourly interval. (c) downward shortwave radiation at hourly interval. (d) Aerosol Optical Depth at 550 nm at 10 minutes interval.

170

3 Use Cases

3.1 Pairing GeCGO data with TESViS Subset data for scientific insight

One of the key advantages of GeCGO is its ability to be combined with data from polar-orbiting satellites. Creating GeCGO in the same format as the TESViS Subset makes it easy to integrate and analyze the two data sets together. Figure 5 illustrates

175 the approach, combining multiple satellite data from the TESViS Subset with GeCGO. Satellite observations are usually made at different times of the day. For example, Terra and Aqua MODIS observe their target around 10:30 AM and 1:30 PM local time, respectively. The observation time of sensors onboard ISS (e.g., ECOsystem Spaceborne Thermal Radiometer Experiment on Space Station (ECOSTRESS)) varies due to changing altitude and orbital inclination of the station. As a result, the phenomena in ecosystem processes observed by different satellite sensors could be specific to the observation time. Thus, 180 using single satellite data may lead to biased results. Using multiple satellites can overcome the observation-time issue, and the geostationary satellite data can serve as a bridge between the instantaneous observations from multiple satellites. For example, the drought impact on vegetation quantified by ECOSTRESS's instantaneous Evaporative Stress Index (ESI) can be temporally extended to an entire day using radiometric temperatures from geostationary satellite data. In addition, the spatial pattern of drought impacts can be scrutinized using reflectance and radiometric temperature from MODIS or VIIRS finer 185 spatial resolution data (Xiao et al., 2021). Such analyses can be easily implemented using the GeCGO and TESViS Subset data (Fig. 5). Beyond the analysis of diurnal changes, GeCGO can be useful for BRDF modeling which requires observations from multiple sun-target-satellite geometries (e.g. Adachi et al., (2019)). Combining GeCGO with the TESViS Subset can help users find the scenes that meet their research needs.

190 Spatial resolution and projection differences between geostationary satellite and other low earth orbit satellites (e.g., MODIS, VIIRS) are challenging issues for users interested in using them together for various scientific applications. We thus chose to convert the geostationary satellite view projection to geographic projection for its easy useability. The original Level-1b data of GOES ABI is skewed at the edge of the coverage where the view angle is large. Meanwhile, the TESViS Subset data has different projections for each product. For example, MODIS data in the TESViS Subset were reprojected from swath images to Sinusoidal projection. As a result, the MODIS subset data show the skew cutout as shown in Fig. 5g,i. Therefore, matching 195 all the pixels between the subset data is not straightforward. Using only the center pixels where the ground observation sites exist is the easiest way to avoid such complex projection conversion processes.




Figure 5: Example of instantaneous satellite observations in a single day (2020/06/07) at the Bondville, Illinois flux tower site. The yellow arrow represents time of the day in local time (Central Daylight Time, UTC-5). The two-dimensional images were obtained from GeCGO or the TESViS subset. The ABI band2 reflectance (0.59 - 0.69 μm) subset data are shown in the upper panel (a-c). The ABI band14 radiometric temperature subset data are shown in the upper panel (d-f). The lower panels show the low Earth orbit satellite data: Terra MODIS reflectance (MOD09A1) band 2 (0.84 - 0.88 μm) (g), Ecostress Evaporative Stress Index (ESI) (h), Aqua MODIS reflectance (MYD09A1) band 2 (0.84 - 0.88 μm) (i), and VIIRS reflectance (VNP09A1) band I1 (0.60 - 0.68 μm) (j). The outside boundaries of the black line box for each satellite image have the same spatial extent, 40.06°N - 39.96°N and 88.3811°W - 88.1970°W. The red cross shows the location of Bondville flux tower (40.0062°N, 88.2904°W). The light-blue triangles indicate the approximate observation time of each satellite image.

3.2 Examples of Using GeCGO with Ground Observation Network Data

3.2.1 AmeriFlux: Which NDVI or NIRv can represents annual GPP?

AmeriFlux is a network of sites that use the eddy covariance technique to measure ecosystem CO₂, water, and energy fluxes across North, Central, and South America (Chu et al., 2023). Due to its wide distribution across diverse ecosystems and climate gradients, AmeriFlux data has been used for validating satellite products (Baldocchi et al., 2001) and for upscaling site-specific observations from AmeriFlux sites to regional and global scale using remote sensing data (Running et al., 2004). However,

since each MODIS on TERRA or AQUA provides only one observation at a specific time of the day in the daytime, converting

MODIS observations to representative daily values requires various assumptions or models to align with the daily average of the half-hourly observations provided by AmeriFlux products. In contrast, the geostationary satellite data can be directly compared with the half-hourly AmeriFlux data. Several data-driven models using geostationary satellite data have been proposed to estimate carbon and water fluxes (Khan et al., 2022; Li et al., 2023b; Stoy et al., 2024; Xiao et al., 2021) and 220 vegetation indices for tracking the diurnal cycle of GPP (Jeong et al., 2023). Given the dynamic nature of fluxes and the environmental factors that influence them throughout the day, combining hyper-temporal geostationary satellite data with AmeriFlux data provides a more accurate representation of diurnal variations in fluxes compared to using polar-orbiting satellite data. GeCGO further enables users to work with specific datasets without needing to download the entire GeoNEX data, making it especially valuable for model development.

225 As an example of using AmeriFlux data, we compared annual GPP with mean NDVI and NIRv (Badgley et al., 2019) to analyze the spatial representation of vegetation indices from ABI data in 2018 (Fig. 6). The annual GPP were obtained from AmeriFlux FLUXNET data product (Pastorello et al., 2020) for each site. The mean annual NDVI and NIRv were derived from daily values of NDVI and NIRv, and the daily NDVI and NIRv were calculated from the median of 90 percentile data for each day. Unlike TOA products (i.e. L1G), L2 data contain significant gaps due to the cloud mask derived from the MAIAC 230 algorithm. Therefore, we excluded the AmeriFlux sites with fewer than 100 days of available daily data.

We found significant relationships across all comparisons between mean annual GPP and vegetation indices (VIs). Even TOA VIs, which did not undergo cloud screening, had high correlations ($r = 0.73$ for NDVI and $r = 0.64$ for NIRv) with annual GPP (Fig. 6a,b). Moreover, surface VIs showed strong correlation with GPP compared to TOA VIs ($r = 0.83$ for NDVI, and $r = 0.81$ for NIRv) (Fig. 6c,d), suggesting that surface reflectance VIs better represented spatial patterns of annual GPP despite 235 the limited data availability. For both L1G and L2, NDVI showed a slightly stronger correlation with annual GPP than NIRv.




Figure 6: Scatter plot (blue dots) of annual GPP ($\text{kgC/m}^2/\text{year}$) at AmeriFlux sites against mean L1G NDVI (a), mean L1G NIRv (b), mean L2 NDVI(c), and mean L2 NIRv(d). The solid line is a linear regression line. The dashed lines represent 95% confidence intervals of the regression.

240

245

250

However, these results do not imply that NDVI is inherently better than NIRv for representing spatial variability of annual GPP. The resolutions of the GeoNEX datasets are greater than 0.005 degree (approximately 500 m at the nadir). While spatial resolution of 500 m may be comparable to the footprint of many flux sites, this resolution only applies to a red visible band (0.64 μm) of ABI at the nadir. As a result, high spatial heterogeneity within the pixels can lead to a scale mismatch between the flux data and the GeoNEX data footprints, although this footprint scale mismatch issue is not only limited to GeoNEX datasets. Chu et al., (2021) provides representativeness information of tower footprint for all AmeriFlux sites. We encourage users to consult with the footprint information to select appropriate sites for directly analyzing the relationship between flux data and GeCGO. Additionally, in this example, we did not account for other factors such as BRDF dependency on sun-target-satellite geometry (Morton et al., 2014). **[removed]** Further analysis, considering these factors, would provide more appropriate conclusion regarding the spatial representation of vegetation indices.


To validate and calibrate VIs derived from GeCGO, high-frequency vegetation indices estimated at the ground-level can be valuable. For example, some scientific instrument manufactures produce sensors capable of measuring NDVI at sub-hourly

intervals (e.g., Apogee Instruments, Holland Scientific). While these sensors can be installed on flux towers and are used at some sites, they are not standard components of flux towers. More notably, a group of flux and remote sensing scientists recently developed a method to derive sub-hourly NDVI and NIRv using quantum sensors and pyranometers, which are commonly installed on flux towers (Mallick et al., 2024). This approach can be used to validate and calibrate VI derived from geostationary satellites, addressing footprint mismatches, improving accuracy, and reducing uncertainties.

3.2.2 AERONET: validation of the Level-2 AOD products using AERONET AOD

AERONET is a global aerosol monitoring network, which monitors the aerosol optical depth using sun photometers. The time interval of observation is less than hour. **The Aerosol Optical Depth (AOD) has large diurnal variability**. Therefore, the sub-hourly observations of GOES data are suitable for large scale AOD estimation compared to once-a-day observation of polar orbiting satellites (Sorek-Hamer et al., 2020). AOD in the GeoNEX Level-2 data is a product of estimating surface reflectance using MAIAC (Lyapustin et al., 2011b).

To assess the accuracy of GeoNEX Level-2 AOD estimates, we compared the AERONET version 3 data (Sinyuk et al., 2020) to the GeCGO AOD data. AERONET provides the AOD data estimated from sun photometers radiance measured approximately every three minutes. We plotted the GeCGO AOD data against AERONET version 3 data for matching times within two minutes in 2019 (Fig. 7). In this example, the GeCGO AOD overestimated the AERONET AOD, especially for the low AOD less than 0.3. GeCGO will facilitate the development of AOD estimation algorithm and will contribute to improving high-temporal AOD estimation algorithms for the geostationary satellite mission focusing on air quality monitoring (i.e. Tropospheric Emissions: Monitoring of Pollution (TEMPO), Global Environmental Monitoring System (GEMS), and Sentinel-4).



275 **Figure 7: Density plot of the GeCGO AOD at 550 nm against the AERONET AOD at 500nm. The white line represents linear regression line. The plot includes only 2019 data over the GOES 16 coverage area.**

3.2.3 PhenoCam: Time-series comparison between L1-G NDVI and PhenoCam greenness

PhenoCam is the phenology ground observation network using time-lapse cameras. Each site provides time-series of digital

280 camera images, with the objective of capturing the seasonal cycle of vegetation phenology (Richardson et al., 2018b).

PhenoCam provides daily statistics of the image values including the daily Green Chromatic Coordinate (GCC). The GCC

was calculated by dividing the Green Digital Number (DN) by the sum of Red, Green, and Blue DNs. The DNs of the

PhenoCam were extracted from Regions Of Interest (ROI), which focus on dominant vegetation types in the images. The GCC

time-series aligned well with polar orbiting satellite NDVI (Richardson et al., 2018a). Miura et al. (2019) reported that the

285 geostationary satellite data can track the detailed phenological and environmental change captured by time-series in camera

images.

To demonstrate usefulness of GeCGO for analyzing the time-series difference between geostationary data and PhenoCam data,

we conducted six comparisons between the GeCGO VIs and PhenoCam Dataset V2 data (Seyednasrollah et al., 2019) (Fig.

8). Note that there is an intrinsic scale mismatch between PhenoCam and GOES data as the target area captured by PhenoCam

290 is much smaller than the spatial resolution of GOES ABI visible bands (1x1 km).

In Fig. 8a and 8b, the PhenoCam sites were located within the same GOES target pixel. The GCC time series of Harvard Farm

North (Fig. 8a) aligned well with GOES NDVI, while Harvard Farm South (Fig. 8b) did not. This difference may be attributed

to the fact that the PhenoCam ROI at Harvard Farm South was covered by heterogeneous landscapes including vegetation

types (grass in this case), leading to an undesirable condition for this comparison the GOES pixel. Similar mismatch issues

295 have been observed in comparisons between GCC and MODIS NDVI (Richardson et al., 2018a).

Figure 8 c-f provides additional examples that show a high correlation between seasonal GCC and GOES NDVI time series.

At the deciduous vegetation sites (Fig. 8 c-e), mismatch between GCC and GOES NDVI was observed early in the growing

season. Specifically, GCC peaked in the early growing season, while the peak of GOES NDVI was delayed. This discrepancy


can be explained by the emergence of new bright leaves or understory greening (Ryu et al., 2014). The detailed observations

300 and analysis at Harvard Forest Environment Measurement Site (EMS) revealed that the GCC peak can be explained only by

combination of change in leaf traits and canopy structure (Keenan et al., 2014). Meanwhile, the GCC and GOES NDVI aligned

well at evergreen forest sites, where seasonal canopy changes are less pronounced than in deciduous trees (Fig. 8f). These

examples demonstrate that the GeoNEX can be a convenient tool for analyzing leaf phenology.



305 **Figure 8: Examples of time-series of L1G 1-km NDVI (black dot) and Green Chromatic Coordinate (GCC) (green line) at six**
PhenoCam sites in 2019. (a) Harvard Farm, Petersham, MA (ID: harvardfarmnorth, Lat: 42.5205°N, Lon: 72.1822°W, grassland).
(b) Harvard Farm South, Petersham, MA (ID: harvardfarmsouth, Lat: 42.5225°N, Lon: 72.1823°W, grassland). (c) Lost Creek, WI
(ID: lostcreek, Lat: 46.0827°N, Lon: 89.9792°W, deciduous shrub wetland). (d) Russell Sage State Wildlife Management Area, LA
(ID: russellsage, Lat: 32.4570°N, Lon: 91.9743°W, deciduous broadleaf). (e) Willow Creek, Chequamegon-Nicolet National Forest,
WI (ID: willowcreek, Lat: 45.8060°N, Lon: 90.0791°W, deciduous broadleaf). (f) Hemlock Tower, Harvard Forest, Petersham, MA
(ID: harvardsage, Lat: 42.5394°N, Lon: 72.1780°W, evergreen needleleaf).

4 Data Availability

310 DOI of GeCGO is <https://doi.org/10.25966/y5pe-xp41> and available at the NASA Ames Data Portal
[\(https://data.nas.nasa.gov/gecgo/data.php\)](https://data.nas.nasa.gov/gecgo/data.php) (Hashimoto et al., 2025). We will add more products on the data portal and
 announce the updates on the NEX website at <https://www.nasa.gov/nasa-earth-exchange-nex/>

5 Code Availability

GeoNEXTools is available at GeoNEXTools GitHub site (<https://github.com/nasa/GeoNEXTools>).

6 Conclusion

320 In contrast to the MODIS satellites, which will soon reach the end of their respective missions, the operational geostationary satellites are expected to provide a long-term data record due to their essential role in weather forecasting. Since the sensors on the latest geostationary satellites are equivalent to the MODIS sensors and thus suitable for observing the land surface at global scale, we anticipate that the GeoNEX data and GeCGO will have a valuable role to play in research continuity at the end of the MODIS era.

325 The GeoNEX project aims to develop key Earth observation and science products using data from a global constellation of geostationary satellite sensors. Initially focusing on the GOES domain (covering North and South America), the GeoNEX dataset is expanding to include additional geographic regions served by other geostationary satellites (i.e., Himawari, GK2A, MTG-I, etc.). NEX has already obtained L1B data of Himawari AHI (84°E to 156°W) and GK2A AMI (72°E to 168°W) and created L1G data. We've also initiated importing and processing MTG-I FCI data to cover Europe and Africa. This extended
330 spatial coverage of GeoNEX will further support global ground networks that are not currently included in GeCGO. Furthermore, in response to the needs of the scientific community, we plan to incorporate additional datasets, such as cloud cover. To increase data accessibility, GeoNEX and the AmeriFlux Management Project have closely collaborated to make GeCGO available through the AmeriFlux website. This collaboration enables users to download and visualize time series of various GeCGO products including spectral vegetation index, land surface temperature, and downwelling shortwave solar
335 radiation. We hope this collaborative effort fosters interdisciplinary research among the flux, remote sensing, and modeling communities to better understand Earth systems and address critical environmental challenges.

In conclusion, we described the details of GeCGO and GeoNEXTools in this paper. GeCGO is convenient for users who want to analyze the large volume of GeoNEX data for land ecosystem monitoring and modeling. GeCGO can help users for synergistical use of the GeoNEX data with other satellite sensor data. GeCGO is also useful to analyze the relationship between
340 the GeoNEX data and ground observation network data. We showed three examples to demonstrate how we can relate the GeCGO with ground observation networks. The first example analyzed the relationship between annual GPP from AmeriFlux and the summation of VIs. The linear relationship showed the possibility of the GeCGO VIs to estimate annual GPP and develop algorithms for spatial variability of annual GPP. The second example uses the geostationary data to track phenological changes observed in PhenoCam data. They highlight the value of frequent observations from geostationary satellites in helping
345 mitigate cloud cover problems and for capturing quick responses of vegetation to environmental changes. As such, these examples demonstrate the value of ready-to-use GeCGO in terrestrial ecophysiology research.

Author Contribution

HH wrote the draft. HH, WW, and TP designed the methodology and investigated the research. SK curated the data and 350 software for public users. KI and RRN, MT, and KY supervised the projects. ARM and AG managed the computer resources. IGB administered the project.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgements

355 We would like to thank Dr. Koen Hufkens (BlueGreen Labs) for allowing us to use his code to develop GeoNEXTTools. This study was funded by National Aeronautics and Space Administration (NASA) Earth Exchange and NASA NNH19ZDA001N-ESROGSS. The AmeriFlux data portal and processing pipeline are maintained by the AmeriFlux Management Project, supported by the U. S. Department of Energy Office of Science, Office of Biological and Environmental Research, under contract number DE-AC02-05CH11231. We collaborated with Chiba University on this study, supported by the Japan Society 360 for the Promotion of Science (JSPS), KAKENHI JP22H05004, and JSPS Core-to-Core Program (JPJSCCA20220008).

References

Adachi, Y., Kikuchi, R., Obata, K., and Yoshioka, H.: Relative Azimuthal-Angle Matching (RAM): A screening method for GEO-LEO reflectance comparison in middle latitude forests, *Remote Sens* (Basel), 11, 1095, <https://doi.org/10.3390/rs11091095>, 2019.

365 Badgley, G., Anderegg, L. D. L., Berry, J. A., and Field, C. B.: Terrestrial gross primary production: Using NIRv to scale from site to globe, *Glob Chang Biol*, 25, 3731–3740, <https://doi.org/10.1111/gcb.14729>, 2019.

Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw, U. K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A new 370 tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *Bull Am Meteorol Soc*, 82, 2415–2434, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.

Chu, H., Luo, X., Ouyang, Z., Chan, W. S., Dengel, S., Biraud, S. C., Torn, M. S., Metzger, S., Kumar, J., Arain, M. A., Arkebauer, T. J., Baldocchi, D., Bernacchi, C., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Bracho, R., 375 Brown, S., Brunsell, N. A., Chen, J., Chen, X., Clark, K., Desai, A. R., Duman, T., Durden, D., Fares, S., Forbrich, I., Gamon, J. A., Gough, C. M., Griffis, T., Helbig, M., Hollinger, D., Humphreys, E., Ikawa, H., Iwata, H., Ju, Y., Knowles, J. F., Knox, S. H., Kobayashi, H., Kolb, T., Law, B., Lee, X., Litvak, M., Liu, H., Munger, J. W., Noormets, A., Novick, K., Oberbauer, S. F., Oechel, W., Oikawa, P., Papuga, S. A., Pendall, E., Prajapati, P., Prueger, J., Quinton, W. L., Richardson, A. D., Russell, E. S., Scott, R. L., Starr, G., Staebler, R., Stoy, P. C., Stuart-Haëntjens, 380 E., Sonnentag, O., Sullivan, R. C., Suyker, A., Ueyama, M., Vargas, R., Wood, J. D., and Zona, D.: Representativeness of Eddy-Covariance flux footprints for areas surrounding AmeriFlux sites, *Agric For Meteorol*, 301–302, 108350, <https://doi.org/10.1016/J.AGRFORMAT.2021.108350>, 2021.

Chu, H., Christianson, D. S., Cheah, Y. W., Pastorello, G., O'Brien, F., Geden, J., Ngo, S. T., Hollowgrass, R., Leibowitz, K., Beekwilder, N. F., Sandesh, M., Dengel, S., Chan, S. W., Santos, A., Delwiche, K., Yi, K., Buechner, C., Baldocchi, 385 D., Papale, D., Keenan, T. F., Biraud, S. C., Agarwal, D. A., and Torn, M. S.: AmeriFlux BASE data pipeline to support network growth and data sharing, *Scientific Data* 2023 10:1, 10, 1–13, <https://doi.org/10.1038/s41597-023-02531-2>, 2023.

Hashimoto, H., Wang, W., Dungan, J. L., Li, S., Michaelis, A. R., Takenaka, H., Higuchi, A., Myneni, R. B., and Nemani, R. R.: New generation geostationary satellite observations support seasonality in greenness of the Amazon evergreen 390 forests, *Nat Commun*, 12, 684, <https://doi.org/10.1038/s41467-021-20994-y>, 2021.

He, T., Zhang, Y., Liang, S., Yu, Y., and Wang, D.: Developing Land Surface Directional Reflectance and Albedo Products from Geostationary GOES-R and Himawari Data: Theoretical Basis, Operational Implementation, and Validation, *Remote Sensing* 2019, Vol. 11, Page 2655, 11, 2655, <https://doi.org/10.3390/RS11222655>, 2019.

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, 395 T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A federated instrument network and data archive for aerosol characterization, *Remote Sens Environ*, 66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.

Jeong, S., Ryu, Y., Dechant, B., Li, X., Kong, J., Choi, W., Kang, M., Yeom, J., Lim, J., Jang, K., and Chun, J.: Tracking diurnal to seasonal variations of gross primary productivity using a geostationary satellite, GK-2A advanced meteorological imager, *Remote Sens Environ*, 284, 113365, <https://doi.org/10.1016/J.RSE.2022.113365>, 2023.

Jia, A., Liang, S., and Wang, D.: Generating a 2-km, all-sky, hourly land surface temperature product from Advanced Baseline 400 Imager data, *Remote Sens Environ*, 278, 113105, <https://doi.org/10.1016/J.RSE.2022.113105>, 2022.

Keenan, T. F., Darby, B., Felts, E., Sonnentag, O., Friedl, M. A., Hufkens, K., O'Keefe, J., Klosterman, S., Munger, J. W., Toomey, M., and Richardson, A. D.: Tracking forest phenology and seasonal physiology using digital repeat photography: A critical assessment, *Ecological Applications*, 24, 1478–1489, <https://doi.org/10.1890/13-0652.1>; PAGE:STRING:ARTICLE/CHAPTER, 2014.

Khan, A. M., Stoy, P. C., Joiner, J., Baldocchi, D., Verfaillie, J., Chen, M., and Otkin, J. A.: The Diurnal Dynamics of Gross Primary Productivity Using Observations From the Advanced Baseline Imager on the Geostationary Operational Environmental Satellite-R Series at an Oak Savanna Ecosystem, *J Geophys Res Biogeosci*, 127, e2021JG006701, <https://doi.org/10.1029/2021JG006701>, 2022.

410 The MODISTools package: an interface to the MODIS Land Products Subsets Web Services: <https://github.com/ropensci/MODISTools>, last access: 11 July 2024.

Li, R., Wang, D., Wang, W., and Nemani, R.: A GeoNEX-based high-spatiotemporal-resolution product of land surface downward shortwave radiation and photosynthetically active radiation, *Earth Syst Sci Data*, 15, 1419–1436, <https://doi.org/10.5194/ESSD-15-1419-2023>, 2023a.

415 Li, X., Ryu, Y., Xiao, J., Dechant, B., Liu, J., Li, B., Jeong, S., and Gentine, P.: New-generation geostationary satellite reveals widespread midday depression in dryland photosynthesis during 2020 western U.S. heatwave, *Sci Adv*, 9, https://doi.org/10.1126/SCIADV.ADI0775/SUPPL_FILE/SCIADV.ADI0775_SM.PDF, 2023b.

Losos, D., Hoffman, S., and Stoy, P. C.: GOES-R land surface products at Western Hemisphere eddy covariance tower locations, *Sci Data*, 11, 1–19, <https://doi.org/10.1038/s41597-024-03071-z>, 2024.

420 Lyapustin, A., Martonchik, J., Wang, Y., Laszlo, I., and Korkin, S.: Multiangle implementation of atmospheric correction (MAIAC): 1. Radiative transfer basis and look-up tables, *J Geophys Res*, 116, D03210, <https://doi.org/10.1029/2010JD014985>, 2011a.

Lyapustin, A., Wang, Y., Laszlo, I., Kahn, R., Korkin, S., Remer, L., Levy, R., and Reid, J. S.: Multiangle implementation of atmospheric correction (MAIAC): 2. Aerosol algorithm, *J Geophys Res*, 116, D03211, <https://doi.org/10.1029/2010JD014986>, 2011b.

425 Mallick, K., Verfaillie, J., Wang, T., Ortiz, A. A., Szutu, D., Yi, K., Kang, Y., Shortt, R., Hu, T., Sulis, M., Szantoi, Z., Boulet, G., Fisher, J. B., and Baldocchi, D.: Net fluxes of broadband shortwave and photosynthetically active radiation complement NDVI and near infrared reflectance of vegetation to explain gross photosynthesis variability across ecosystems and climate, *Remote Sens Environ*, 307, 114123, <https://doi.org/10.1016/J.RSE.2024.114123>, 2024.

430 Miura, T., Nagai, S., Takeuchi, M., Ichii, K., and Yoshioka, H.: Improved characterisation of vegetation and land surface seasonal dynamics in central Japan with Himawari-8 hypertemporal data, *Sci Rep*, 9, 1–12, <https://doi.org/10.1038/s41598-019-52076-x>, 2019.

Morton, D. C., Nagol, J., Carabajal, C. C., Rosette, J., Palace, M., Cook, B. D., Vermote, E. F., Harding, D. J., and North, P. R. J.: Amazon forests maintain consistent canopy structure and greenness during the dry season, *Nature*, 506, 221–224, <https://doi.org/10.1038/nature13006>, 2014.

435 Novick, K. A., Biederman, J. A., Desai, A. R., Litvak, M. E., Moore, D. J. P., Scott, R. L., and Torn, M. S.: The AmeriFlux network: A coalition of the willing, *Agric For Meteorol*, 249, 444–456, <https://doi.org/10.1016/J.AGRFORMAT.2017.10.009>, 2018.

440 ORNL DAAC: Terrestrial Ecology Subsetting & Visualization Services (TESViS) Fixed Sites Subsets,
<https://doi.org/10.3334/ORNLDAAAC/1567>, 2017.

445 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y. W., Poindexter, C., Chen, J., Elbashandy, A.,
Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A.,
Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E.,
Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P.
450 D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J. M., Bowling, D. R., Bracho, R., Brodeur, J.,
Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini,
I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E.,
Curtis, P. S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De Ligne, A., De
Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G.,
Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W.,
Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M.,
Frank, J., Galvagno, M., Gharun, M., Gianelle, D., et al.: The FLUXNET2015 dataset and the ONEFlux processing
455 pipeline for eddy covariance data, *Sci Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.

Richardson, A. D., Hufkens, K., Milliman, T., and Frolking, S.: Intercomparison of phenological transition dates derived from
460 the PhenoCam Dataset V1.0 and MODIS satellite remote sensing, *Sci Rep*, 8, 5679, <https://doi.org/10.1038/s41598-018-23804-6>, 2018a.

Richardson, A. D., Hufkens, K., Milliman, T., Aubrecht, D. M., Chen, M., Gray, J. M., Johnston, M. R., Keenan, T. F.,
Klosterman, S. T., Kosmala, M., Melaas, E. K., Friedl, M. A., and Frolking, S.: Tracking vegetation phenology across
465 diverse North American biomes using PhenoCam imagery, *Sci Data*, 5, 180028,
<https://doi.org/10.1038/sdata.2018.28>, 2018b.

Running, S. W., Nemani, R. R., Heinsch, F. A., Zhao, M., Reeves, M., and Hashimoto, H.: A continuous satellite-derived
measure of global terrestrial primary production, *Bioscience*, 54, 547–560, [https://doi.org/10.1641/0006-3568\(2004\)054\[0547:ACSMOG\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0547:ACSMOG]2.0.CO;2), 2004.

Ryu, Y., Lee, G., Jeon, S., Song, Y., and Kimm, H.: Monitoring multi-layer canopy spring phenology of temperate deciduous
470 and evergreen forests using low-cost spectral sensors, *Remote Sens Environ*, 149, 227–238,
<https://doi.org/10.1016/J.RSE.2014.04.015>, 2014.

Schmit, T. J., Griffith, P., Gunshor, M. M., Daniels, J. M., Goodman, S. J., and Lebair, W. J.: A closer look at the ABI on the
GOES-R series, *Bull Am Meteorol Soc*, 98, 681–698, <https://doi.org/10.1175/BAMS-D-15-00230.1>, 2017.

Seyednasrollah, B., Young, A. M., Hufkens, K., Milliman, T., Friedl, M. A., Frolking, S., and Richardson, A. D.: Tracking
475 vegetation phenology across diverse biomes using Version 2.0 of the PhenoCam Dataset, *Sci Data*, 6, 222,
<https://doi.org/10.1038/S41597-019-0229->

9;SUBJMETA=158,172,2165,2445,2457,4081,631,704;KWRD=ECOSYSTEM+ECOLOGY,ENVIRONMENTAL
+IMPACT,PHENOLOGY, 2019.

Sinyuk, A., Holben, B., Eck, T., Giles, D., Slutsker, I., Korkin, S., Schafer, J., Smirnov, A., Sorokin, M., and Lyapustin, A.:
475 The AERONET Version 3 aerosol retrieval algorithm, associated uncertainties and comparisons to Version 2, *Atmos
Meas Tech*, 13, 3375–3411, <https://doi.org/10.5194/amt-13-3375-2020>, 2020.

Sorek-Hamer, M., Chatfield, R., and Liu, Y.: Review: Strategies for using satellite-based products in modeling PM2.5 and
short-term pollution episodes, *Environ Int*, 144, 106057, <https://doi.org/10.1016/J.ENVINT.2020.106057>, 2020.

Stoy, P., Ranjbar, S., Hoffman, S., and Losos, D.: Estimating terrestrial carbon dioxide and water vapor fluxes from
480 geostationary satellites in near-real time: the ALIVE framework, *EGU24*, <https://doi.org/10.5194/EGUSPHERE-EGU24-4743>, 2024.

Takenaka, H., Sakashita, T., Higuchi, A., and Nakajima, T.: Geolocation correction for geostationary satellite observations by
a phase-only correlation method using a visible channel, *Remote Sens (Basel)*, 12, 2472,
<https://doi.org/10.3390/rs12152472>, 2020.

485 Tenenbaum, P. and Wohler, B.: Ziggy, <https://doi.org/10.5281/zenodo.7859503>, 2024.

Tuck, S. L., Phillips, H. R. P., Hintzen, R. E., Scharlemann, J. P. W., Purvis, A., and Hudson, L. N.: MODISTools –
downloading and processing MODIS remotely sensed data in R, *Ecol Evol*, 4, 4658–4668,
<https://doi.org/10.1002/ECE3.1273>, 2014.

490 Wang, W., Li, S., Hashimoto, H., Takenaka, H., Higuchi, A., Kalluri, S., and Nemani, R.: An introduction to the Geostationary-
NASA Earth Exchange (GeoNEX) Products: 1. Top-of-atmosphere reflectance and brightness temperature, *Remote
Sens (Basel)*, 12, 1267, <https://doi.org/10.3390/RS12081267>, 2020.

Wang, W., Wang, Y., Lyapustin, A., Hashimoto, H., Park, T., Michaelis, A., and Nemani, R.: A novel atmospheric correction
algorithm to exploit the diurnal variability in hypertextural geostationary observations, *Remote Sens (Basel)*, 14,
964, <https://doi.org/10.3390/RS14040964>, 2022.

495 Xiao, J., Fisher, J. B., Hashimoto, H., Ichii, K., and Parazoo, N. C.: Emerging satellite observations for diurnal cycling of
ecosystem processes, *Nat Plants*, 7, 877–887, <https://doi.org/10.1038/s41477-021-00952-8>, 2021.

Yi, K., Senay, G. B., Fisher, J. B., Wang, L., Suvocarev, K., Chu, H., Moore, G. W., Novick, K. A., Barnes, M. L., Keenan,
T. F., Mallick, K., Luo, X., Missik, J. E. C., Delwiche, K. B., Nelson, J. A., Good, S. P., Xiao, X., Kannenberg, S. A.,
500 Ahmadi, A., Wang, T., Bohrer, G., Litvak, M. E., Reed, D. E., Oishi, A. C., Torn, M. S., and Baldocchi, D.: Challenges
and Future Directions in Quantifying Terrestrial Evapotranspiration, *Water Resour Res*, 60, e2024WR037622,
<https://doi.org/10.1029/2024WR037622>, 2024.

Appendix A: Function names and description of the GeoNEXTools

Function name	Description
gt_products()	Lists all available GeoNEX Products Subset products
gt_bands()	Lists all available bands for a GeoNEX Products Subset product
gt_sites()	Lists all available GeoNEX Products Subset pre-processed sites
gt_subset()	Download a GeoNEX Products Subset product for a given GeoNEX subset site
505 <u>gt_batch_subset()</u>	Download a GeoNEX Products Subset product for given multiple GeoNEX subset site