

An Accurate 10 m Annual Crop Map Product of Maize and Soybean Across the United States

Haijun Li¹, Xiao-Peng Song^{1, *}, Bernard Adusei¹, Jeffrey Pickering¹, Andre Lima¹, Andrew Poulson¹, Antoine Baggett¹, Peter Potapov^{1, 2}, Ahmad Khan¹, Viviana Zalles^{1, 2}, Andres Hernandez-Serna¹, Samuel M. Jantz^{1, 3}, Amy H. Pickens¹, Carolina Ortiz-Dominguez¹, Xinyuan Li¹, Theodore Kerr¹, Zhen Song¹, Svetlana Turubanova¹, Eddy Bongwele¹, Heritier Koy Kondjo¹, Anna Komarova¹, Stephen V. Stehman⁴, Matthew C. Hansen¹

¹Department of Geographical Sciences, University of Maryland, College Park, MD 20742, United States

²World Resources Institute, Washington, DC, 20002, United States

³National Institute for Modeling Biological Systems, University of Tennessee, Knoxville, TN 37996, United States

⁴Department of Sustainable Resources Management, SUNY College of Environmental Science and Forestry, Syracuse, NY 13210, United States

Correspondence to: Xiao-Peng Song (xpsong@umd.edu)

Abstract High-resolution crop maps over large spatial extents are fundamental to many agricultural applications; however, generating high-quality crop maps consistently across space and time remains a challenge. In this study, we improved a workflow for crop mapping and developed an openly available, annual, 10-m spatial resolution maize and soybean map product over the Contiguous United States (CONUS) from 2019 to 2022 (available at <https://glad.umd.edu/dataset/mapping-crops-10-m-resolution-united-states>). We obtained all available Sentinel-2 surface reflectance data between May and October for every year, applied quality assurance, corrected the bidirectional reflectance distribution function (BRDF) effects, and generated 10-day analysis ready data (ARD) composites. We then derived multi-temporal metrics from the 10-day ARD as training features for the national-scale wall-to-wall mapping. We implemented a stratified, two-stage cluster sampling, and then conducted annual field surveys and collected ground data. Utilizing the training data with Sentinel-2 multi-temporal metrics and topographic factors, we trained random forest models generalized for annual maize and soybean classification separately. Validated using field data from the two-stage cluster sample, our annual maps achieved consistent overall accuracies (OA) greater than 95% with standard errors of less than 1%. User's accuracies (UAs) and producer's accuracies (PAs) for maize were higher than 91% and 84% across the years, and UAs and PAs for soybean were greater than 88% and 82%, respectively. To illustrate the substantial improvement of the 10-m map over existing datasets, e.g., the 30-m Cropland Data Layer (CDL), we aggregated the 10-m maps to 30-m spatial resolution and quantified the number of mixed pixels that can be reduced by improving the mapping from 30 m to 10 m. The counties with the most maize and soybean production in Iowa, Illinois and Nebraska had the lowest reduction in mixed pixels, ranging from 1% to 7%, whereas southern counties had a higher reduction in mixed pixels. Overall, the median percentages of mixed maize and soybean pixels reduction across all counties were 8% and 9%, respectively. With more Sentinel-2-like data available from continuous observations and incoming satellite missions,

we anticipate that 10-m crop maps will greatly benefit long-term monitoring for agricultural practices from the field to global scales. The dataset is also available at <https://doi.org/10.6084/m9.figshare.28934993.v2> (Li et al., 2025)

1 Introduction

Satellite-derived crop maps are essential to many agricultural applications, such as crop yield prediction (Bolton and Friedl, 2013; Song et al., 2022; Wang et al., 2024), food market forecasting (Tanaka et al., 2023), crop area estimation (Khan et al., 2016), conservation policy design (Song et al., 2021b; Zalles et al., 2021), smallholder livelihood evaluation (Lambert et al., 2018), warfare impacts on food security (Li et al., 2022; Lin et al., 2023), and greenhouse gas emissions in agriculture (Escobar et al., 2020; Ouyang et al., 2023). However, along with these benefits are the outstanding challenges to generating high-quality crop maps, including developing consistent ready-to-use satellite datasets, collecting representative field data, and building classification algorithms robust to phenological variations.

Dense time series of satellite observations with complete spatial coverage is essential to mapping crops at broad scales. With global coverage and daily revisit frequency, the Moderate Resolution Imaging Spectroradiometer (MODIS) data are often used for crop mapping in early studies (Wardlow et al., 2007; Wardlow and Egbert, 2008). However, spatial details within individual small fields can rarely be depicted at 250-m resolution (Fritz et al., 2015), especially for more than 475 million smallholder and family farms accounting for 12% of the world's agricultural land (Lowder et al., 2016). Since the opening of the Landsat archive in 2008 (Woodcock et al., 2008), Landsat data have been extensively used to generate 30-m crop maps in many parts of the world, such as in North America (Boryan et al., 2011; Fiset et al., 2013; Johnson and Mueller, 2021; Song et al., 2017; Wang et al., 2020), Europe (Foerster et al., 2012), South America (Song et al., 2021b), and Asia (Dong et al., 2016; Khan et al., 2021; Remelgado et al., 2020). However, Landsat-based crop mapping is hampered by the relatively sparse 16-day temporal frequency (8 days with two satellites), especially when cloudy weather persists. Compared to Landsat, Sentinel-2A and -2B together have a revisit frequency of 5 days and provide 10-m, 20-m and 60-m spectral bands including red edge bands that are particularly useful for crop identification (Immitzer et al., 2016; Song et al., 2021a). These advantages make Sentinel-2 data one of the best publicly accessible data sources for crop mapping (Ghassemi et al., 2022; Han et al., 2021; Luo et al., 2022; You et al., 2021).

Crop classification from satellite imagery is usually implemented by relating specific crop types to remotely sensed features, using reference data and classification algorithms such as conventional machine learning or advanced deep learning (e.g., Alami Machichi et al., 2023; Joshi et al., 2023). Therefore, *in situ* data can serve as critical references to annotate satellite imagery for supervised classifications, although field surveys over large areas require extensive time and labor resources. Currently, the US Department of Agriculture (USDA) National Agricultural Statistics Service (NASS) collects periodic field data across the US and produces the Cropland Data Layer (CDL) annually based on a large amount of ground data and supervised algorithms (Boryan et al., 2011). When current-year labels are unavailable, some researchers have explored

transferring pre-trained models to target regions or years (Luo et al., 2022; Wang et al., 2019), or generating labels with knowledge-guided approaches (Lin et al., 2022; You et al., 2023). However, these approaches are limited to experiments at small spatial scales, such as the US Midwest and Northeast China, and thus the efficiency of national-scale crop classification over large countries with more challenging environments remains to be explored. In cases where reference data are entirely unavailable, unsupervised classifications are used first to cluster satellite-derived features and then assign crop labels to the clusters to generate approximate crop maps (Konduri et al., 2020; Xiong et al., 2017). They, however, are vulnerable to outliers and noisy features and require intensive visual inspections (Wang et al., 2019). In summary, collecting representative ground data is a critical yet challenging component for large-area crop mapping.

Spatiotemporal consistency in crop classifications is necessary to make annual crop maps comparable and thus allow long-term crop monitoring and change analysis. Yet this is undermined partly due to crop phenology variations across large extents, depending on soil properties, planting dates, and weather conditions, among other factors (Deines et al., 2023; Yang et al., 2017). On one hand, within a calendar year, crop progress is regionally different. To address this issue, some studies trained regional models through agroecological zoning, which requires zone-specific training and validation (de Aballeyra et al., 2020; Wardlow and Egbert, 2008). On the other hand, yearly unaligned phenological profiles can jeopardize the classification consistency across years, especially when extreme weather events occur (Manoochehr et al., 2021). Given these interannual variations, classifiers that accurately identify crops in average normal growing seasons using single-date or time-series satellite imagery may perform poorly for abnormal years. To this end, researchers proposed yearly specific classifications (Massey et al., 2017; Som-ard et al., 2022). However, these annual models need fine-tuning based on reference data from each corresponding year especially when encountering unseen growing trends, and thus cannot be generalized for long-term periods.

Multi-temporal metrics are statistical transformations of temporal profiles of satellite observations that can improve spatial and temporal consistency and facilitate land cover mapping for large areas. In the mid-1980s, researchers derived phenological features from pixel time series from the Advanced Very High Resolution Radiometer (AVHRR) for vegetation monitoring (Malingreau, 1986) and from Landsat Multispectral Scanner (MSS) for crop classification (Badhwar, 1984). This metrics method was then widely used for land cover mapping and change analysis from regional to global scales using AVHRR, MODIS and Landsat data (DeFries et al., 1995; Hansen et al., 2013; Potapov et al., 2021b; Song et al., 2018). For crop mapping over continental scales, the metrics method was used to generalize classification models robust to interannual phenological variations (Song et al., 2021b). Many studies are adopting similar concepts for regional crop mapping (Kerner et al., 2022; Konduri et al., 2020; Yang et al., 2023; Zhong et al., 2014).

In the US, the CDL has been used widely for many applications (Bolton and Friedl, 2013; Gao et al., 2017; Lobell et al., 2020; Wright and Wimberly, 2013; Yan and Roy, 2016). Recently, the 2024 CDL has been successfully released with the spatial resolution increased from 30 m to 10 m (https://www.nass.usda.gov/Research_and_Science/Cropland/SARS1a.php, accessed 26 December 2025). However, the previous years of CDL are at 30-m resolution, and have inconsistent accuracies depending on the location, and inaccurate classifications are observed in sparse or complex agricultural regions (Larsen et al., 2015). The

30-m spatial resolution can lead to substantial mixed pixels, obscuring incremental or pixel-level changes, particularly along field boundaries. In comparison, 10-m maps with a higher spatial resolution can improve the delineation of precise field boundaries, reducing mixed pixels in individual fields, as well as lowering uncertainties of area estimation. Global 10-m crop mapping efforts are rare, although the WorldCereal provides an example (Van Tricht et al., 2023). In Europe, recent 10-m crop mapping efforts include the Crop Map of England (CROME) (CROME, 2024), the parcel-level crop maps in the Netherlands (ESA, 2024), the crop maps produced by the Sentinel-2 for Agriculture (Sen2-Agri) (Defourny et al., 2019; Inglada et al., 2015), and the more recent High Resolution Layer Crop Types (CTY) (EU, 2024). In Asia, large-area crop-specific maps have been generated recently (Han et al., 2021; Li et al., 2023; Mei et al., 2024). In the US, the potential of national-scale 10-m crop mapping has rarely been explored, although a recent prototyping effort has been reported (Huang et al., 2024).

The objective of this study is to develop annual 10-m crop maps with Sentinel-2 time series. We also quantify the benefits of 10-m maps compared to existing 30-m products. In this study, we generated annual maize and soybean maps at 10-m spatial resolution over the entire Contiguous US (CONUS), from 2019 to 2022. We also quantified the benefits of our 10-m crop maps in mixed pixel reduction compared to 30-m maps, at field, regional and national scales. We improved a workflow developed in previous studies (Li et al., 2023; Song et al., 2017) by combining satellite analysis ready data (ARD) generation, field survey design, and machine learning. An overview workflow for annual crop map production is presented in Figure 1.




Figure 1: Overview of the workflow for large-area annual crop map production.

2 Materials and methods

2.1 Satellite analysis ready data (ARD) generation

Operational crop mapping over large areas relies on satellite data that are geometrically and radiometrically consistent with quality assessment (e.g., Boryan et al., 2011; Fiset et al., 2013; Song et al., 2021b). Analysis ready data (ARD), defined by the Committee on Earth Observation Satellites (CEOS), meet such criteria as “have been processed to a minimum set of requirements and organized into a form that allows immediate analysis with a minimum of additional user effort and interoperability both through time and with other datasets” (<https://ceos.org/ard/>, accessed 11 November 2024). To support annual wall-to-wall crop mapping over the CONUS, we obtained all available Sentinel-2 data between May and October, applied quality assurance, corrected the bidirectional reflectance distribution function (BRDF) effects, and generated 10-day ARD composites.

We downloaded Sentinel-2A and -2B Level-2A Bottom of the Atmosphere reflectance (S2 L2A) images from Google Cloud, including the 10-m blue, green, red and near-infrared (NIR) bands, the 20-m red edge (RE1, RE2, RE3), narrow near-infrared (NNIR) and shortwave infrared (SWIR1, SWIR2) bands. We selected images acquired between May 1 and October 31 after filtering out images with > 80% cloud cover. We then processed all available Sentinel-2 data by utilizing the GLAD and Zaratan high-performance computing clusters at the University of Maryland and generated Sentinel-2 ARD for the wall-to-wall crop mapping. Details of the ARD generation are described in the following sections.

2.1.1 Quality assurance

Based on the S2 scene classification (SCL) layer, we generated the cloud mask by merging categories of cloud shadow, thin cirrus, snow, cloud with low, medium and high probability into cloudy pixels. We also produced an additional cloud mask layer derived from the Fmask algorithms (Zhu et al., 2015) and the Cloud Displacement Index (Frantz et al., 2018). We combined the SCL-derived cloud mask with the additional cloud mask as the final quality assurance (QA) layer.

2.1.2 Bidirectional Reflectance Distribution Function (BRDF) correction

We corrected the BRDF effects using the c-factor method to derive nadir BRDF-adjusted reflectance (NBAR) (Roy et al., 2017a, b). The S2 L2A product provides solar and view geometry metadata in 23×23 grids at 5-km spatial resolution. For each multi-spectral instrument (MSI) detector in each spectral band, the solar zenith and azimuth angles remain consistent; the view zenith and azimuth angles, however, vary from one detector to another, and from band to band. We calculated the mean value of the view zenith and azimuth angle for each 5-km grid across all detectors and all spectral bands. Per-pixel solar and view angles at 10-m resolution were derived by nearest neighbor interpolation of the 5-km grid values. As a result, the 10-m angle layers were used to generate NBAR images using the global spectral BRDF model parameters (Roy et al., 2017a, b). This process reduced the BRDF effects and improved the spatial coherence compared to the surface reflectance without BRDF correction (Figure 2).




Figure 2: Sentinel-2 false color composites (R: NIR, G: SWIR1, B: SWIR2) over a selected UTM tile 14TMP centered at (97.131° W, 41.942° N). (a-b) surface reflectance. (c-d) nadir BRDF-adjusted reflectance (NBAR). Two overlapping Sentinel-2B swaths acquired from orbit R112 on July 18, 2022 (backscattering direction) and orbit R012 on July 21, 2022 (forward scattering direction) were used. The orbit R112 data were overlaid on the orbit R012 data where they overlapped. All composites are displayed with the same stretch parameters.

2.1.3 Temporal composition and tiling

We resampled the 20-m bands to 10-m using the nearest neighbor method, applied the QA layer, and created 10-day median composites. For each NBAR band in a given 10-day interval, the median value of all clear-sky observations and the corresponding day of the year (DOY) were selected. We also implemented temporal linear interpolation on a per-pixel basis to fill the data gaps (Griffiths et al., 2019). For a missing value in a 10-day interval, the gap-filled value was calculated from the preceding and subsequent valid observations. A maximum of six 10-day intervals (i.e., 60 days or 2 months) was used to limit the period so that the interpolation was temporally relevant. For cases in which cloud-free observations are unavailable for two months, we did not conduct interpolation.

We divided the entire study area into $1^\circ \times 1^\circ$ non-overlapping tiles in geographic latitude/longitude projection with WGS84 datum. Each tile was named by the latitude and longitude coordinates of the lower-left corner, with $0.0001^\circ \times 0.0001^\circ$ spatial

161 resolution to approximately match a 10-m pixel of Sentinel-2. We reprojected 1,028 Sentinel-2 Universal Transverse Mercator
162 (UTM) tiles into 939 $1^\circ \times 1^\circ$ tiles over the United States.

163 **2.1.4 Multi-temporal metrics**

164 The 10-day S2 ARD may have inconsistent observational frequencies across space and time depending on the geographical
165 location and cloud condition. Generating multi-temporal metrics from ARD can improve data consistency, and thus enable
166 large-area land cover mapping, which has been demonstrated in various applications at continental and global scales (Hansen
167 et al., 2013; Potapov et al., 2021; Song et al., 2021a).

168 Following the method in Potapov et al. (2020), we generated multi-temporal metrics from the 10-day S2 ARD (see Table S1).
169 First, we derived the Normalized Difference Vegetation Index (NDVI, $(\text{NIR} - \text{Red})/(\text{NIR} + \text{Red})$) (Tucker, 1979) and the
170 normalized ratio between shortwave infrared bands (SWSW, $(\text{SWIR1} - \text{SWIR2})/(\text{SWIR1} + \text{SWIR2})$) from corresponding
171 NBAR bands. Second, we ranked time-series observations by each NBAR band or index individually. We then selected the
172 second maximum, the second minimum, and median values per pixel, and calculated the 10th, 25th, 75th, and 90th percentiles.
173 We also calculated the average, standard deviation, and amplitude between these percentiles and the second maximum, the
174 second minimum values. Third, we ranked the observation day of year (DOYs) according to the time-series NDVI, and derived
175 values on the DOYs corresponding to the second maximum, the second minimum, and median, as well as the 10th, 25th, 75th,
176 and 90th percentiles of NDVI values. The average, standard deviation and amplitude were also calculated from these extracted
177 values.

178 In total, we calculated 621 metrics. The NBAR averages between the 25th and 75th percentiles from observations ranked by
179 individual bands are illustrated in Figure 3 and Figure 4a. The NBAR amplitudes reveal land surface phenology and thus
180 simplify visual interpretation of general land cover types such as cropland, open water, forest, and wetland (Figure 4b). When
181 the averages are calculated from observations with the highest NDVI values (between 90th percentile and the second maximum
182 NDVI value), the composite shows surface reflectance during the peak growing season, improving the identification of
183 multiple crop types (Figure 4c and Figure 4d).




Figure 3: Sentinel-2 composites over the United States in 2022. The composites were created using the average value of nadir Bidirectional Reflectance Distribution Function (BRDF)-adjusted reflectance (NBAR) between the 25th and 75th percentiles from observations ranked by individual bands (R: SWIR1, G: NIR, B: Red). The original 10-m data are resampled to 250 m using the nearest neighbor for visualization purposes. The ESRI map is used as background.




Figure 4: Composites of Sentinel-2 multi-temporal metrics in the Mississippi Valley. (a) SWIR1-NIR-Red composite of NBAR average between the 25th and 75th percentiles from observations ranked by individual bands; (b) SWIR1-NIR-Red composite of NBAR amplitude between the second maximum and the second minimum values; (c) NIR-SWIR1-SWIR2 composites of average NBAR between the 90th percentile and the second maximum values from observations ranked by NDVI; (d) 2022 Cropland Data Layer. The coordinate of the center point is (91.312° W, 33.665° N). All panels are displayed in the same scale at 10-m resolution.


2.2 Sampling design and field survey

To support the 10-m crop mapping, we conducted extensive field surveys for *in situ* data collection, based on a two-stage cluster sampling design following Song et al. (2017). This approach has been demonstrated to be effective for agricultural applications in which ground reference data are collected at regional (Khan et al., 2018), national (King et al., 2017; Li et al., 2023), and continental (Song et al., 2021b) scales.

2.2.1 Sampling design

Following previous research, we divided the study area into 20 km × 20 km equal-area blocks and designed the two-stage cluster sampling to target fields to visit. We first derived the per-block maize and soybean area fractions from the previous

203 year's crop map, sorted all blocks from the highest to the lowest fraction, and then stratified the ranked blocks into high,
 204 medium and low strata. Following previous studies (King et al., 2017; Song et al., 2017), we selected a simple random sample
 205 of blocks from each stratum as the primary sampling units (PSUs) and selected a simple random sample of 10 m × 10 m pixels
 206 in each PSU as the secondary sampling units (SSUs) (Figure 5a) (see Table S2).



207 **Figure 5: 2022 stratified sampling design for field survey. (a) stratified sampling design. 20 km × 20 km equal-area blocks were**
 208 **stratified into high, medium and low strata. (b) crop calendar for maize and soybean over the US. Google Earth imagery is used as**
 209 **background.**
 210

211 **2.2.2 Field data collection**

212 The typical planting season of the US maize starts in April while soybean planting starts in May; the harvesting season starts
 213 in September and ends in November for maize and in October for soybean (see Figure 5b above). We conducted the field
 214 survey during the peak growing season in July and August. Consistent with previous research (Li et al., 2023; Song et al.
 215 2021a), we collected two types of datasets during the field survey: 1) ground reference data over the probability sample of
 216 SSUs for map evaluation and crop area estimation; and 2) “windshield survey” reference data for model training. These

windshield survey data were collected along the driving routes between the SSUs, and were only used to train models for classification and not for validation, whereas the probability sample was exclusively used for validation.

For each year from 2019 to 2022, we selected a separate stratified two-stage cluster sample following the general sampling framework and collected in-season ground data (Figure 6, Table S2-S4).




Figure 6: Annual primary sampling unit (PSU) blocks from 2019 to 2022. The ESRI map is used as background.


2.3 Crop classification

We conducted crop classifications in two stages: 1) at the PSU level, we mapped maize and soybean over all sample PSUs using field data, Sentinel-2 time-series imagery, and decision tree classifiers; and 2) at the national scale, we employed random forest classifiers to map maize and soybean using the PSU-maps as training, multi-temporal metrics derived from Sentinel-2 ARD as well as the topographic features derived from TanDEM-X (DLR, 2024) as input. We evaluated the accuracy of the national crop map using the field data from the SSUs to determine the reference class label.

2.3.1 PSU-level crop mapping

We processed all available Sentinel-2 data over the PSUs from May 1 to October 31 to produce in-season PSU-level maize and soybean maps. We trained two decision tree classifiers separately for maize and soybean classification by using all the bands and normalized ratios of any two bands, as well as the “windshield survey” points as training (Figure 7b). Applying the trained models to time-series images, we created a binary maize/non-maize map and a binary soybean/non-soybean map at 10-

234 m resolution for each PSU (Figure 7c). These in-season PSU maps were then pooled as training labels for national-scale wall-
235 to-wall mapping.



236
237 **Figure 7: An example of primary sampling unit (PSU) block-level crop mapping using field data. (a) a representative sample block**
238 **in Illinois with center coordinates shown on the Google Earth imagery. (b) field data collection in the PSU. The secondary sampling**
239 **units (SSUs) of pixels are shown as yellow diamonds. The “windshield survey” points are shown as white dots. The driving routes**
240 **are shown in pink tracks. (c) PSU-level crop maps.**

241 2.3.2 Wall-to-wall crop classification

242 The multi-temporal metrics derived from the Sentinel-2 ARD were the main input for national mapping. In addition, we
243 downloaded the nominal 12-m TanDEM-X data from the German Aerospace Center (DLR, 2024), and derived 10-m spatial
244 resolution elevation, slope, and aspect using nearest neighbor resampling. These topographic data were combined with the
245 multi-temporal metrics (see Section 2.1.4 above) as inputs for supervised classification. We generated training labels from the
246 10-m maize and soybean PSU maps. We randomly selected 0.2% of maize (soybean) and 0.8% of non-maize (non-soybean)
247 pixels from each PSU as training labels. Conflict classification pixels from the binary maize and soybean maps were excluded
248 in the training dataset.

249 To conduct crop classifications, we employed Random Forest (RF), a widely adopted ensemble machine learning algorithm
250 in remote sensing due to its accuracy, computational efficiency, and robustness to noise (Belgiu and Drăguț, 2016; Breiman,
251 2001). Following the approach detailed in Li et al. (2023), we tailored RF binary classifiers separately for maize (RF-Maize)
252 and soybean (RF-Soybean). The models were fine-tuned using a random search followed by a grid search (Probst et al., 2019),
253 on a randomly selected subset of 1% of the training dataset, and subsequently re-trained with optimal hyperparameters on the
254 entire training dataset (see more technical details in Figure S1, Figure S2 and Table S5).

255 We aggregated the per-pixel class probability layers from RF-Maize and RF-Soybean by selecting the highest probability
256 (maize vs. soybean) and derived the aggregated probability layer and corresponding crop mask layer (see Figure S3 for an
257 example). We then applied a 5×5 pixel kernel opening followed by a 10×10 pixel kernel closing, to eliminate scattered

pixels and fill holes within large homogeneous fields. The kernel sizes were selected based on tests and visual assessments to balance noise removal while preserving fine details. We generated the final maize and soybean map using the aggregated probability layer following the area-matching approach reported by Song et al (2017), Song et al. (2021b) and Li et al. (2023).

2.4 Map evaluation

2.4.1 Accuracy assessment

Utilizing the annually field-visited SSUs, we validated the annual maps from 2019 to 2022. Overall accuracy (OA), user's accuracy (UA) and producer's accuracy (PA) with associated uncertainty estimates were estimated using a ratio estimator for two-stage cluster sampling within a stratified design, following good practices (Olofsson et al., 2013; Stehman, 2014). The formulas for accuracy estimation are found in Song et al. (2017, Appendix A.)

2.4.2 Crop area comparison with official statistics

We derived the pixel-counting-based crop areas for maize and soybean from the annual crop maps, for each year from 2019 to 2022. We compared these crop areas with the official statistical crop areas from the USDA NASS at the county and state levels. We then calculated root-mean-square-difference (RMSD) and r^2 between the mapped crop areas and the statistical areas.

3 Results

3.1 Visual assessment

Our 10-m crop map reveals well-known spatial patterns of maize and soybean cultivation in the United States (Figure 8). The dominant soybean cultivation is shown in the Midwest states, the Great Plains states, the Mississippi Valley and the eastern coast, whereas maize is widely distributed across the country.




Figure 8: The 10-m maize and soybean map for 2022. The ESRI map is used as background.

Specifically, our 10-m crop map delineated more field-scale details compared to the 30-m CDL (Figure 9). Midwest states such as Illinois typically have rectangular crop fields, and our 10-m map generated homogeneous fields with clearer boundaries (Figure 9a). Our map also captured more landscape fragmentation, such as smaller fields with greater crop diversity in the Mississippi Valley (Figure 9b) and the agriculture/wetland mosaic in North Dakota (Figure 9c).




Figure 9: Maize and soybean classification in 2022 over selected regions. Rows (a-c) are representative sites in Illinois, Mississippi, and North Dakota. All panels are displayed at the same scale (10 km × 10 km). The coordinates of the center points are shown on the Google Earth imagery. The 10-day composite periods are shown on the Sentinel-2 image (R: NIR, G: SWIR1, B: SWIR2). Maize and soybean are shown in yellow and green colors, respectively.

3.2 Quantitative accuracy assessment

We conducted an accuracy assessment for annual maps using the annual SSUs as references (Table 1). All maps achieved OAs greater than 95% with standard errors less than 1%. UAs and PAs for maize were higher than 91% and 84%, respectively, while UAs and PAs for soybean were higher than 89% and 82%, respectively.

296

297

Table 1: Accuracy assessment for maize and soybean maps from 2019 to 2022. Cell entries in the confusion matrices represent area proportions. Reference data were derived from probability samples of secondary sampling units (SSUs).

Year	Class	Reference				Users'	Producers'	Overall
		Maize	Soybean	Others	Total	accuracy % (SE)	accuracy % (SE)	accuracy % (SE)
2019	Maize	0.1111	0.0017	0.0053	0.1181	94.0 (1.5)	85.6 (2.6)	95.4 (0.5)
	Soybean	0.0003	0.0868	0.0056	0.0927	93.7 (1.8)	83.8 (2.4)	
	Others	0.0183	0.0150	0.7558	0.7891	95.8 (0.6)	98.6 (0.3)	
	Total	0.1297	0.1036	0.7667	1			
2020	Maize	0.1073	0.0031	0.0045	0.1149	93.4 (1.6)	91.0 (1.8)	95.9 (0.5)
	Soybean	0.0012	0.0941	0.0086	0.1039	90.6 (1.8)	84.5 (2.7)	
	Others	0.0095	0.0141	0.7576	0.7812	97.0 (0.5)	98.3 (0.3)	
	Total	0.1180	0.1113	0.7707	1			
2021	Maize	0.1021	0.0044	0.0053	0.1118	91.2 (1.8)	92.8 (1.5)	95.3 (0.6)
	Soybean	0.0012	0.0967	0.0109	0.1088	89.3 (2.5)	82.1 (2.5)	
	Others	0.0066	0.0168	0.7560	0.7793	96.8 (0.5)	97.8 (0.4)	
	Total	0.1098	0.1179	0.7723	1			
2022	Maize	0.0884	0.0024	0.0055	0.0963	91.8 (2.0)	84.0 (3.5)	95.3 (0.7)
	Soybean	0.0019	0.0904	0.0095	0.1018	88.8 (4.0)	85.8 (2.5)	
	Others	0.0150	0.0126	0.7744	0.8020	96.6 (0.6)	98.1 (0.6)	
	Total	0.1052	0.1054	0.7894	1			

298

3.3 Comparison between the crop maps and agricultural statistics

299

300

301

302

303

We compared our map-based area estimates with agricultural statistics reported by the NASS at state and county scales. The state-level area comparisons between our mapped areas and the NASS statistics showed close agreements, with r^2 greater than 0.99 and root-mean-square-difference (RMSDs) less than 900 km² for maize, and RMSDs less than 1,800 km² for soybean (Figure 10). At the county level (Figure 11), our mapped maize and soybean areas also matched the NASS statistics well with r^2 greater than 0.97 and RMSDs between 30 km² and 50 km².




Figure 10: State-level comparison between mapped maize and soybean areas and NASS statistics.




Figure 11: County-level comparisons between mapped maize and soybean areas and NASS statistics.

308 **4 Discussion**

309 **4.1 The benefits of 10-m crop maps in mixed pixel reduction**

310 Using the 2022 10-m crop map as an example, we conducted a quantitative data analysis to illustrate the benefits of 10-m crop
311 mapping over 30-m mapping. We first removed small fields less than 9 10-m pixels (one 30-m pixel), then spatially aggregated
312 the 10-m map to 30-m resolution and derived the maize and soybean cover fraction for each 30-m pixel. We defined pure
313 pixels as 100% cover and anything below as mixed pixels. We applied a 50% cover threshold to determine the dominant crop
314 type within mixed pixels. Pixels where neither maize nor soybean cover reached 50% were ignored. Rather than assessing
315 accuracies for the aggregated 30-m maps, our objective was to compare the 10-m versus 30-m resolution by quantifying
316 changes in mixed pixels and analyzing the spatial patterns.

317 Unsurprisingly, the aggregated 30-m map showed that pure pixels are clustered in large-size homogeneous fields (Figure 12a).
318 Mixed pixels occurred in small, fragmented fields, on field edges, or along the road networks, where crops coexisted with
319 other land cover (e.g., other crops, pasture, built-up, etc.) (Figure 12b, c). Our 10-m maps showed clear advantages over the
320 30-m CDL in mixed pixel reduction in various landscapes (Figure 13). In North Dakota where numerous fields are fragmented,
321 the 10-m map presents more homogeneous fields and captures within-field patterns of water ponds (Figure 13a); for center-
322 pivot irrigated fields in Nebraska, the 10-m map delineates cleaner circular patterns (Figure 13b); in Appalachian Pennsylvania
323 where many fields are in narrow strips, the 10-m map distinguishes neighboring strip cropping fields better than the 30-m CDL
324 in which the fields are mapped with a large amount of mixed pixels (Figure 13c).




Figure 12: The 2022 aggregated 30-m maize and soybean map overlaid with road and rail networks. Columns (a-c) are selected sites in Nebraska, Minnesota, and Ohio. The 30-m map was derived by spatially aggregating the 10-m map by calculating the fractional cover and categorized as pure pixels with 100% cover or mixed pixels with <100% cover. All panels are displayed at the same scale (10 km × 10 km). The coordinates of the center points are shown on the Google Earth imagery. The rail and road networks are obtained from the US TIGER database.



Figure 13: The 2022 aggregated 30-m maize and soybean map and CDL show mixed pixels in various landscapes. (a) wetland/agriculture mosaics in North Dakota; (b) center-pivot irrigated fields in Nebraska; (c) strip fields in Pennsylvania. The coordinates of the center points are shown on the Google Earth imagery.

We obtained the percentage of mixed maize and soybean pixels at the county level to examine the spatial distribution of mixed pixel reduction from 30 m to 10 m (Figure 14). The counties with the highest maize and soybean production, such as those in Iowa, Illinois, and Nebraska, had the least mixed pixel percentages ranging from 1% to 10%, while counties in the upper Midwest, the North and South Plains, the northeast and eastern coast had more mixed pixels (Figure 14a, Figure 14c). Overall, the median percentages of mixed maize and soybean pixels in all counties were 8% and 9%, respectively (Figure 14b, d). Our results show that increasing the spatial resolution of crop mapping from 30 m to 10 m would reduce the number of mixed pixels by 8-9% at the county scale, and substantially benefit many states outside of the Midwest region.




Figure 14: The percentages of 30-m mixed maize and soybean pixels at the county level derived from the 10-m map. (a) the spatial distribution of mixed maize pixels; (b) the statistical distribution of mixed maize pixels; (c-d) the same as (a-b) but for soybean. Counties accounting for 99.9% coverage of the national maize and soybean cultivation derived from the 2022 NASS statistics are shown.

We conducted additional analysis by comparing our aggregated 30-m map with the 30-m CDL for the maize and soybean classes. We applied a 50% threshold to convert our aggregated 30-m map to binary maize and soybean maps and conducted a per-pixel comparison. We then aggregated the per-pixel results to the county level based on the ratio of the difference of maize or soybean pixels between our map and CDL divided by maize or soybean pixels in CDL. Positive values indicate more maize or soybean pixels in our map, whereas negative values indicate more pixels in CDL (Figure 15). In several regions, our map presented more maize pixels, such as in northern Texas and southern Louisiana (Figure 15a), and more soybean pixels, such as in western North Dakota and Georgia (Figure 15c). In general, CDL reported more maize and soybean pixels in most counties compared to our aggregated 30-m map (Figure 15b, d), which is likely due to the exclusion of 10-m maize and soybean pixels below the 50% threshold during aggregation.




Figure 15: Comparison between our aggregated 30-m map with the 2022 30-m CDL at the county level. (a) normalized difference of maize pixel; (b) histogram of the normalized difference of maize pixel; (c-d) the same as (a-b) but for soybean. Counties accounting for 99.9% coverage of the national maize and soybean cultivation derived from the 2022 NASS statistics are shown.

To estimate the number of mixed pixels that might be reduced by improving the CDL's spatial resolution from 30 m to 10 m, we overlaid the 2022 30-m CDL on our 30-m fractional cover map, and computed the number and proportion of CDL maize and soybean pixels that were mixed pixels at the county level (Figure 16). The mixed pixel derived from CDL showed similar spatial distribution patterns to results derived from the aggregated 30-m map (see Figure 14 above). The median percentage of mixed pixels across all counties for both maize and soybean was 8% (Figure 16b, d), which is close to the values of 8% (9% for soybean) derived from our aggregated 30-m map (see Figure 14 above). These consistent mixed pixel estimates from our map and the CDL indicate substantial benefits of 10-m crop maps in reducing mixed pixels over existing 30-m products, especially for regions outside of Midwest, as illustrated in our analyses.




Figure 16: The percentages of 30-m mixed maize and soybean pixels at the county level by comparing our aggregated 30-m map and the 2022 30-m CDL. (a) the spatial distribution of mixed maize pixels; (b) the statistical distribution of mixed maize pixels; (c-d) the same as (a-b) but for soybean. Counties accounting for 99.9% coverage of the national maize and soybean cultivation derived from the 2022 NASS statistics are shown.

4.2 The potential of 10-m crop maps in finer-scale agricultural monitoring

Higher-resolution crop maps have great potential to facilitate remote-sensing-based agricultural applications at finer scales. For example, the Crop Sequence Boundaries (CSB), which delineate polygons of homogeneous cropping sequences with 8-year moving windows, have been developed based on the CDL by the USDA (Hunt et al., 2023). The 30-m CDL was resampled to 10-m resolution to improve the masking of road networks as the roads and rails in rural areas are typically less than 30 m in width. Consequently, the resampled 10-m maps may delineate inaccurate field boundaries due to mixed pixels (Figure 17). The CSB delineated large homogeneous fields well (Figure 17a) but showed more fragments when encountering within-field cropping variations (Figure 17b). The misalignments between the delineated field edges and pixel boundaries are extensive in heterogeneous landscapes and small fields (Figure 17c, d), and thus the polygon-based crop acreage derived from CSB layers may be biased. Alternatively, using originally produced higher-resolution (e.g., 10-m) maps can yield more accurate field

delineation, cropping sequences, and crop area estimates with smaller uncertainties (Duveiller and Defourny, 2010; Ozdogan and Woodcock, 2006; Yan and Roy, 2014).





Figure 17: The aggregated 30-m maize and soybean map overlaid with Crop Sequence Boundaries. Columns (a-d) are selected sites in Illinois, Mississippi, North Dakota, and Iowa. All panels are displayed at the same scale (10 km × 10 km). The coordinates of the center points are shown. The 2015-2022 Crop Sequence Boundaries are obtained from the USDA NASS.

With higher-resolution satellite imagery available from continuous observations (e.g., Sentinel-1 and Sentinel-2) and upcoming missions (e.g., Landsat Next, NASA-ISRO SAR Mission (NISAR)), we anticipate that 10-m crop maps will play a more critical role in agricultural monitoring from the field to global scales.

4.3 The robustness of temporal metrics for annual crop map production

Stacking satellite-derived time-series maps is one of the most common practices to investigate long-term agriculture-related land cover and land use change, such as cropping history (Blickensdörfer et al., 2022; Johnson, 2019), crop and cropland expansion (Lark et al., 2020; Potapov et al., 2021a; Song et al., 2021b; Zalles et al., 2019), and cropland intensification (Kehoe et al., 2017; Marin et al., 2022). However, in large-extent countries such as the US, the spatiotemporal consistency in multi-year crop classifications can be impacted by both intra-annual and interannual variations in crop phenology (Figure 18). For example, the 2022 NDVI time series for maize and soybean showed noticeably different crop progress across the CONUS

399 (Figure 18a, Figure 18b). In Arkansas, maize growth peaked in mid-May and started senescence in early August, whereas
 400 maize in the Midwest states was at early growing stages in mid-May and at the peak growing season in early August. Soybean
 401 also showed noticeable disparities in crop progress across the US states. On the other hand, interannual phenological shifts
 402 also impede the classification consistency (Figure 18c, Figure 18d). In Illinois, similar NDVI profiles between 2020 and 2022
 403 suggested overall consistent growing progress, while the patterns in 2019 and 2021 showed higher interannual variations. In
 404 2021, Illinois experienced an earlier planting pace for maize and soybean partly due to the favorable spring weather conditions
 405 and soybean varieties adapted to early plantation (Nafziger, 2024). In 2019, crop phenology shifted substantially as a result of
 406 planting delays caused by extremely heavy precipitation in the spring (Manoochehr et al., 2021). Consequently, at the state
 407 level in Illinois, maize was planted at only 24% compared to the previous year's 95% and the five-year average of 49% by the
 408 end of May 2019; soybean was planted at 9% compared to the previous year's 79% and the five-year average of 51% (NASS
 409 CPR, 2024).



410
 411 **Figure 18: NDVI time series for maize and soybean from representative sites. (a) 2022 maize NDVI in Arkansas (AR), Illinois (IL),**
 412 **Kansas (KS), Minnesota (MN), South Dakota (SD); (b) the same as (a) but for soybean; (c) 2019-2022 interannual NDVI variations**
 413 **for maize in Illinois; (d) the same as (c) but for soybean. The details about the sites are shown in Figure S4 and Table S6.**

414 Utilizing the multi-temporal metrics to relatively normalize crop phenological variations, our approach can be applied to
 415 generate annual crop maps over large areas, as also illustrated for South America in Song et al. (2021b). Our four-year sampling

416 designs generated large field samples, allowing us to collect representative training data from various growing conditions and
417 geographical regions. Our workflow generated consistently accurate maize and soybean maps over the entire CONUS, from
418 2019 to 2022. The map accuracies for 2019 (an abnormally wet year), and 2020 and 2021 (both years with normal weather)
419 are consistent with those of 2022 (see Table 1 above).

420 5 Data availability

421 The annual 10-m maize and soybean maps over the CONUS from 2019 to 2022 are openly accessible at the website of the
422 Global Land Analysis and Discovery (GLAD) team at the University of Maryland ([https://glad.umd.edu/dataset/mapping-](https://glad.umd.edu/dataset/mapping-crops-10-m-resolution-united-states)
423 [crops-10-m-resolution-united-states](https://glad.umd.edu/dataset/mapping-crops-10-m-resolution-united-states)). The dataset is also available at <https://doi.org/10.6084/m9.figshare.28934993.v2> (Li et
424 al., 2025). The dataset includes a set of GeoTIFF images in the EPSG:4236 spatial reference system. The values 0, 1, 2, 255
425 represent other, maize, soybean, and no data, respectively. External data used in this study are openly accessible online: 1) the
426 Sentinel-2 data were downloaded from Google Cloud Platform ([https://console.cloud.google.com/marketplace/product/esa-](https://console.cloud.google.com/marketplace/product/esa-public-data/sentinel2)
427 [public-data/sentinel2](https://console.cloud.google.com/marketplace/product/esa-public-data/sentinel2)); 2) the Cropland Data Layer were downloaded from the US Department of Agriculture (USDA) National
428 Agricultural Statistics Service (NASS) (https://www.nass.usda.gov/Research_and_Science/Cropland/Release/index.php); 3)
429 the TanDEM-X was downloaded from the German Aerospace Center (<https://tandemx-science.dlr.de>); 4) the agricultural
430 statistics for CONUS were retrieved from the USDA NASS (https://www.nass.usda.gov/Quick_Stats/index.php); 5) the Crop
431 Sequence Boundaries were derived from the USDA NASS ([https://www.nass.usda.gov/Research_and_Science/Crop-](https://www.nass.usda.gov/Research_and_Science/Crop-Sequence-Boundaries/index.php)
432 [Sequence-Boundaries/index.php](https://www.nass.usda.gov/Research_and_Science/Crop-Sequence-Boundaries/index.php)); 6) the road and rail networks were downloaded from the US TIGER database
433 (<https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html>).

434 6 Conclusions

435 Crop maps at 10-m spatial resolution bring substantial benefits for agricultural applications compared to 30-m products for
436 smallholder as well as industrial agricultural countries. In this study, we developed 10-m maize and soybean maps over the
437 Contiguous US (CONUS) from 2019 to 2022, using all available Sentinel-2 observations and field surveys, with overall
438 accuracies consistently greater than 95%. We explicitly examined the benefits of improving the spatial resolution from 30 m
439 to 10 m by quantifying the reduction in mixed pixels. Our analysis showed that, across all counties in the US, the 10-m maps
440 reduced mixed pixels by a median of 8% for maize and 9% for soybean compared to the aggregated 30-m maps, with most
441 mixed pixels occurring along field edges, road networks, and in heterogeneous fields. Our workflow generated annual maps
442 with consistency across space and over time. Our 10-m crop maps were produced at the end of the growing season, around
443 3~4 months earlier than the official 30-m Cropland Data Layer. As more Sentinel-2-like data become accessible from current
444 observations and planned missions such as Landsat Next, 10-m crop maps presented in this study will greatly benefit

agricultural applications including field boundary extraction, crop sequence delineation, crop condition monitoring, precision fertilization and irrigation, from field to global scales.

Author contributions

HL: Software, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization. **XPS:** Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Resources, Funding acquisition. **BA:** Formal analysis, Investigation. **JP:** Formal analysis, Investigation, Data Curation. **AL:** Formal analysis, Investigation. **AP:** Formal analysis, Investigation, Data Curation, Writing - Review & Editing. **AB:** Investigation. **PP:** Methodology, Investigation, Software, Writing - Review & Editing. **AK:** Methodology, Investigation, Writing - Review & Editing. **VZ:** Methodology, Investigation. **AHS:** Investigation. **SMJ:** Investigation, Writing - Review & Editing. **AHP:** Investigation. **COD:** Investigation. **XL:** Investigation, Writing - Review & Editing. **TK:** Investigation, Writing - Review & Editing. **ZS:** Investigation. **ST:** Investigation. **EB:** Investigation. **HKK:** Investigation. **AK:** Investigation. **SVS:** Methodology, Writing - Review & Editing. **MCH:** Conceptualization, Methodology, Resources, Investigation, Funding Acquisition.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Financial support

This research is supported by the Bezos Earth Fund, Google Academic Research Awards, NASA Land-Cover and Land-Use Change Program, NASA Acres Program, and the University of Maryland. The authors acknowledge the University of Maryland supercomputing resources (<http://hpcc.umd.edu>) made available for conducting the research reported in this paper.

References

- de Abelleira, D., Veron, S., Banchemo, S., Mosciaro, M. J., Propato, T., Ferraina, A., Taffarel, M. C. G., Dacunto, L., Franzoni, A., and Volante, J.: First large extent and high resolution cropland and crop type map of Argentina, in: 2020 IEEE Latin American GRSS & ISPRS Remote Sensing Conference (LAGIRS), 392–396, <https://doi.org/10.1109/LAGIRS48042.2020.9165610>, 2020.
- Alami Machichi, M., mansouri, loubna E., imani, yasmima, Bourja, O., Lahlou, O., Zennayi, Y., Bourzeix, F., Hanadé Houmma, I., and Hadria, R.: Crop mapping using supervised machine learning and deep learning: a systematic literature review, International Journal of Remote Sensing, 44, 2717–2753, <https://doi.org/10.1080/01431161.2023.2205984>, 2023.

473 Badhwar, G. D.: Classification of corn and soybeans using multitemporal thematic mapper data, *Remote Sensing of*
474 *Environment*, 16, 175–181, [https://doi.org/10.1016/0034-4257\(84\)90061-0](https://doi.org/10.1016/0034-4257(84)90061-0), 1984.

475 Belgiu, M. and Drăguț, L.: Random forest in remote sensing: A review of applications and future directions, *ISPRS Journal of*
476 *Photogrammetry and Remote Sensing*, 114, 24–31, <https://doi.org/10.1016/j.isprsjprs.2016.01.011>, 2016.

477 Blickensdörfer, L., Schwieder, M., Pflugmacher, D., Nendel, C., Erasmí, S., and Hostert, P.: Mapping of crop types and crop
478 sequences with combined time series of Sentinel-1, Sentinel-2 and Landsat 8 data for Germany, *Remote Sensing of*
479 *Environment*, 269, 112831, <https://doi.org/10.1016/j.rse.2021.112831>, 2022.

480 Bolton, D. K. and Friedl, M. A.: Forecasting crop yield using remotely sensed vegetation indices and crop phenology metrics,
481 *Agricultural and Forest Meteorology*, 173, 74–84, <https://doi.org/10.1016/j.agrformet.2013.01.007>, 2013.

482 Boryan, C., Yang, Z., Mueller, R., and Craig, M.: Monitoring US agriculture: the US Department of Agriculture, National
483 Agricultural Statistics Service, Cropland Data Layer Program, Geocarto International, 26, 341–358,
484 <https://doi.org/10.1080/10106049.2011.562309>, 2011.

485 Breiman, L.: Random forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/a:1010933404324>, 2001.

486 CROME, Crop Map of England: <https://environment.data.gov.uk/dataset/cc389fe9-f026-4b20-a80f-f424ee833ea6>, last access:
487 28 May 2024.

488 Defourny, P., Bontemps, S., Bellemans, N., Cara, C., Dedieu, G., Guzzonato, E., Hagolle, O., Inglada, J., Nicola, L., Rabaute,
489 T., Savinaud, M., Udriou, C., Valero, S., Bégué, A., Dejoux, J.-F., El Harti, A., Ezzahar, J., Kussul, N., Labbassi, K.,
490 Lebourgeois, V., Miao, Z., Newby, T., Nyamugama, A., Salh, N., Shelestov, A., Simonneaux, V., Traore, P. S., Traore, S. S.,
491 and Koetz, B.: Near real-time agriculture monitoring at national scale at parcel resolution: Performance assessment of the
492 Sen2-Agri automated system in various cropping systems around the world, *Remote Sensing of Environment*, 221, 551–568,
493 <https://doi.org/10.1016/j.rse.2018.11.007>, 2019.

494 DeFries, R., Hansen, M., and Townshend, J.: Global discrimination of land cover types from metrics derived from AVHRR
495 pathfinder data, *Remote Sensing of Environment*, 54, 209–222, [https://doi.org/10.1016/0034-4257\(95\)00142-5](https://doi.org/10.1016/0034-4257(95)00142-5), 1995.

496 Deines, J. M., Swatantran, A., Ye, D., Myers, B., Archontoulis, S., and Lobell, D. B.: Field-scale dynamics of planting dates
497 in the US Corn Belt from 2000 to 2020, *Remote Sensing of Environment*, 291, <https://doi.org/10.1016/j.rse.2023.113551>,
498 2023.

499 DLR, TanDEM-X - Digital Elevation Model (DEM) - Global, 12m: <https://tandemx-science.dlr.de/>, last access: 20 June 2024.

500 Dong, J., Xiao, X., Menarguez, M. A., Zhang, G., Qin, Y., Thau, D., Biradar, C., and Moore, B.: Mapping paddy rice planting
501 area in northeastern Asia with Landsat 8 images, phenology-based algorithm and Google Earth Engine, *Remote Sensing of*
502 *Environment*, 185, 142–154, <https://doi.org/10.1016/j.rse.2016.02.016>, 2016.

503 Duveiller, G. and Defourny, P.: A conceptual framework to define the spatial resolution requirements for agricultural
504 monitoring using remote sensing, *Remote Sensing of Environment*, 114, <https://doi.org/10.1016/j.rse.2010.06.001>, 2010.

505 EU, 2024. The High Resolution Layer Crop Types (CTY). EEA geospatial data catalogue. URL
506 <https://sdi.eea.europa.eu/catalogue/srv/api/records/9db29b07-5968-4ce0-8351-1e356b3d7d47>, last access: 21 November
507 2025.

508 ESA, Monitoring crop health across the Netherlands:
509 [https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-](https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-1/Monitoring_crop_health_across_the_Netherlands)
510 [1/Monitoring_crop_health_across_the_Netherlands](https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-1/Monitoring_crop_health_across_the_Netherlands), last access: 20 June 2024.

511 Escobar, N., Tizado, E. J., zu Ermgassen, E. K. H. J., Löfgren, P., Börner, J., and Godar, J.: Spatially-explicit footprints of
512 agricultural commodities: Mapping carbon emissions embodied in Brazil's soy exports, *Global Environmental Change*, 62,
513 102067, <https://doi.org/10.1016/j.gloenvcha.2020.102067>, 2020.

514 Fisette, T., Rollin, P., Aly, Z., Campbell, L., Daneshfar, B., Filyer, P., Smith, A., Davidson, A., Shang, J., and Jarvis, I.: AAFC
515 annual crop inventory, 2013 Second International Conference on Agro-Geoinformatics (Agro-Geoinformatics), 270–274,
516 <https://doi.org/10.1109/Argo-Geoinformatics.2013.6621920>, 2013.

517 Foerster, S., Kaden, K., Foerster, M., and Itzerott, S.: Crop type mapping using spectral–temporal profiles and phenological
518 information, *Computers and Electronics in Agriculture*, 89, 30–40, <https://doi.org/10.1016/j.compag.2012.07.015>, 2012.

519 Frantz, D., Haß, E., Uhl, A., Stoffels, J., and Hill, J.: Improvement of the Fmask algorithm for Sentinel-2 images: Separating
520 clouds from bright surfaces based on parallax effects, *Remote Sensing of Environment*, 215, 471–481,
521 <https://doi.org/10.1016/j.rse.2018.04.046>, 2018.

522 Fritz, S., See, L., McCallum, I., You, L., Bun, A., Moltchanova, E., Duerauer, M., Albrecht, F., Schill, C., Perger, C., Havlik,
523 P., Mosnier, A., Thornton, P., Wood-Sichra, U., Herrero, M., Becker-Reshef, I., Justice, C., Hansen, M., Gong, P., Abdel Aziz,
524 S., Cipriani, A., Cumani, R., Cecchi, G., Conchedda, G., Ferreira, S., Gomez, A., Haffani, M., Kayitakire, F., Malanding, J.,
525 Mueller, R., Newby, T., Nonguierma, A., Olusegun, A., Ortner, S., Rajak, D. R., Rocha, J., Schepaschenko, D.,
526 Schepaschenko, M., Terekhov, A., Tiangwa, A., Vancutsem, C., Vintrou, E., Wenbin, W., van der Velde, M., Dunwoody, A.,
527 Kraxner, F., and Obersteiner, M.: Mapping global cropland and field size, *Global Change Biology*, 21, 1980–92,
528 <https://doi.org/10.1111/gcb.12838>, 2015.

529 Gao, F., Anderson, M. C., Zhang, X., Yang, Z., Alfieri, J. G., Kustas, W. P., Mueller, R., Johnson, D. M., and Prueger, J. H.:
530 Toward mapping crop progress at field scales through fusion of Landsat and MODIS imagery, *Remote Sensing of*
531 *Environment*, 188, 9–25, <https://doi.org/10.1016/j.rse.2016.11.004>, 2017.

532 Ghassemi, B., Dujakovic, A., Żółtak, M., Immitzer, M., Atzberger, C., and Vuolo, F.: Designing a European-wide crop type
533 mapping approach based on machine learning algorithms using LUCAS field survey and Sentinel-2 data, *Remote Sensing*, 14,
534 <https://doi.org/10.3390/rs14030541>, 2022.

535 Griffiths, P., Nendel, C., and Hostert, P.: Intra-annual reflectance composites from Sentinel-2 and Landsat for national-scale
536 crop and land cover mapping, *Remote Sensing of Environment*, 220, 135–151, <https://doi.org/10.1016/j.rse.2018.10.031>, 2019.

537 Han, J., Zhang, Z., Luo, Y., Cao, J., Zhang, L., Zhang, J., and Li, Z.: The RapeseedMap10 database: annual maps of rapeseed
538 at a spatial resolution of 10 m based on multi-source data, *Earth System Science Data*, 13, 2857–2874,
539 <https://doi.org/10.5194/essd-13-2857-2021>, 2021.

540 Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz,
541 S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R.: High-resolution global
542 maps of 21st-century forest cover change, *Science*, 342, 850–3, <https://doi.org/10.1126/science.1244693>, 2013.

543 Huang, Y., Qiu, B., Yang, P., Wu, W., Chen, X., Zhu, X., Xu, S., Wang, L., Dong, Z., Zhang, J., Berry, J., Tang, Z., Tan, J.,
544 Duan, D., Peng, Y., Lin, D., Cheng, F., Liang, J., Huang, H., and Chen, C.: National-scale 10 m annual maize maps for China
545 and the contiguous United States using a robust index from Sentinel-2 time series, *Computers and Electronics in Agriculture*,
546 221, 109018, <https://doi.org/10.1016/j.compag.2024.109018>, 2024.

547 Hunt, K. A., Abernethy, J., Bowman, M., Wallander, S., and Williams, R.: Crop Sequence Boundaries (CSB): Delineated
548 fields using remotely sensed crop rotations, USDA-NASS, Washington, DC, USA, 2023.

549 Immitzer, M., Vuolo, F., and Atzberger, C.: First experience with Sentinel-2 data for crop and tree species classifications in
550 Central Europe, *Remote Sensing*, 8, 166, <https://doi.org/10.3390/rs8030166>, 2016.

551 Inglada, J., Arias, M., Tardy, B., Hagolle, O., Valero, S., Morin, D., Dedieu, G., Sepulcre, G., Bontemps, S., Defourny, P., and
552 Koetz, B.: Assessment of an operational system for crop type map production using high temporal and spatial resolution
553 satellite optical imagery, *Remote Sensing*, 7, 12356–12379, <https://doi.org/10.3390/rs70912356>, 2015.

554 Johnson, D. M.: Using the Landsat archive to map crop cover history across the United States, *Remote Sensing of Environment*,
555 232, 111286, <https://doi.org/10.1016/j.rse.2019.111286>, 2019.

556 Johnson, D. M. and Mueller, R.: Pre- and within-season crop type classification trained with archival land cover information,
557 *Remote Sensing of Environment*, 264, <https://doi.org/10.1016/j.rse.2021.112576>, 2021.

558 Joshi, A., Pradhan, B., Gite, S., and Chakraborty, S.: Remote-sensing data and deep-learning techniques in crop mapping and
559 yield prediction: A systematic review, *Remote Sensing*, 15, 2014, <https://doi.org/10.3390/rs15082014>, 2023.

560 Kehoe, L., Romero-Munoz, A., Polaina, E., Estes, L., Kreft, H., and Kuemmerle, T.: Biodiversity at risk under future cropland
561 expansion and intensification, *Nature Ecology & Evolution*, 1, 1129–1135, <https://doi.org/10.1038/s41559-017-0234-3>, 2017.

562 Kerner, H. R., Sahajpal, R., Pai, D. B., Skakun, S., Puricelli, E., Hosseini, M., Meyer, S., and Becker-Reshef, I.: Phenological
563 normalization can improve in-season classification of maize and soybean: A case study in the central US Corn Belt, *Science
564 of Remote Sensing*, <https://doi.org/10.1016/j.srs.2022.100059>, 2022.

565 Khan, A., Hansen, M. C., Potapov, P., Stehman, S. V., and Chatta, A. A.: Landsat-based wheat mapping in the heterogeneous
566 cropping system of Punjab, Pakistan, *International Journal of Remote Sensing*, 37, 1391–1410,
567 <https://doi.org/10.1080/01431161.2016.1151572>, 2016.

568 Khan, A., Hansen, M., Potapov, P., Adusei, B., Pickens, A., Krylov, A., and Stehman, S.: Evaluating Landsat and RapidEye
569 data for winter wheat mapping and area estimation in Punjab, Pakistan, *Remote Sensing*, 10, 489,
570 <https://doi.org/10.3390/rs10040489>, 2018.

571 Khan, A., Hansen, M. C., Potapov, P., Adusei, B., Stehman, S. V., and Steininger, M. K.: An operational automated mapping
572 algorithm for in-season estimation of wheat area for Punjab, Pakistan, *International Journal of Remote Sensing*, 42, 3833–
573 3849, <https://doi.org/10.1080/01431161.2021.1883200>, 2021.

574 King, L., Adusei, B., Stehman, S. V., Potapov, P. V., Song, X.-P., Krylov, A., Di Bella, C., Loveland, T. R., Johnson, D. M.,
575 and Hansen, M. C.: A multi-resolution approach to national-scale cultivated area estimation of soybean, *Remote Sensing of
576 Environment*, 195, 13–29, <https://doi.org/10.1016/j.rse.2017.03.047>, 2017.

577 Konduri, V. S., Kumar, J., Hargrove, W. W., Hoffman, F. M., and Ganguly, A. R.: Mapping crops within the growing season
578 across the United States, *Remote Sensing of Environment*, 251, <https://doi.org/10.1016/j.rse.2020.112048>, 2020.

579 Lambert, M.-J., Traoré, P. C. S., Blaes, X., Baret, P., and Defourny, P.: Estimating smallholder crops production at village
580 level from Sentinel-2 time series in Mali's cotton belt, *Remote Sensing of Environment*, 216, 647–657,
581 <https://doi.org/10.1016/j.rse.2018.06.036>, 2018.

582 Lark, T. J., Spawn, S. A., Bougie, M., Gibbs, H. K., Lark, T. J., Spawn, S. A., Bougie, M., and Gibbs, H. K.: Cropland
583 expansion in the United States produces marginal yields at high costs to wildlife, *Nature Communications*, 11,
584 <https://doi.org/10.1038/s41467-020-18045-z>, 2020.

585 Larsen, A. E., Hendrickson, B. T., Dedeic, N., and MacDonald, A. J.: Taken as a given: Evaluating the accuracy of remotely
586 sensed crop data in the USA, *Agricultural Systems*, 141, 121–125, <https://doi.org/10.1016/j.agry.2015.10.008>, 2015.

587 Li, H., Song, X.-P., Hansen, M. C., Becker-Reshef, I., Adusei, B., Pickering, J., Wang, L., Wang, L., Lin, Z., Zalles, V.,
588 Potapov, P., Stehman, S. V., and Justice, C.: Development of a 10-m resolution maize and soybean map over China: Matching
589 satellite-based crop classification with sample-based area estimation, *Remote Sensing of Environment*, 294,
590 <https://doi.org/10.1016/j.rse.2023.113623>, 2023.

591 Li, H., Song, X.-P., Adusei, B., Pickering, J., Lima, A., Poulson, A., Baggett, A., Potapov, P., Khan, A., Zalles, V., Hernandez-
592 Serna, A., Jantz, S. M., Pickens, A. H., Ortiz-Dominguez, C., Li, X., Kerr, T., Song, Z., Turubanova, S., Bongwele, E., Koy
593 Kondjo, H., Komarova, A., Stehman, S. V., and Hansen, M. C.: 2019-2022 10-m maize and soybean maps over the United
594 States, <https://doi.org/10.6084/m9.figshare.28934993.v1>, 2025.

595 Li, X.-Y., Li, X., Fan, Z., Mi, L., Kandakji, T., Song, Z., Li, D., and Song, X.-P.: Civil war hinders crop production and
596 threatens food security in Syria, *Nature Food*, 3, 38–46, <https://doi.org/10.1038/s43016-021-00432-4>, 2022.

597 Lin, C., Zhong, L., Song, X.-P., Dong, J., Lobell, D. B., and Jin, Z.: Early- and in-season crop type mapping without current-
598 year ground truth: Generating labels from historical information via a topology-based approach, *Remote Sensing of*
599 *Environment*, 274, <https://doi.org/10.1016/j.rse.2022.112994>, 2022.

600 Lin, F., Li, X., Jia, N., Feng, F., Huang, H., Huang, J., Fan, S., Ciais, P., and Song, X.-P.: The impact of Russia-Ukraine
601 conflict on global food security, *Global Food Security*, 36, 100661, <https://doi.org/10.1016/j.gfs.2022.100661>, 2023.

602 Lobell, D. B., Deines, J. M., and Tommaso, S. D.: Changes in the drought sensitivity of US maize yields, *Nature Food*, 1, 729–
603 735, <https://doi.org/10.1038/s43016-020-00165-w>, 2020.

604 Lowder, S. K., Skoet, J., and Raney, T.: The number, size, and distribution of farms, smallholder farms, and family farms
605 worldwide, *World Development*, 87, 16–29, <https://doi.org/10.1016/j.worlddev.2015.10.041>, 2016.

606 Luo, Y., Zhang, Z., Zhang, L., Han, J., Cao, J., and Zhang, J.: Developing high-resolution crop maps for major crops in the
607 European Union based on transductive transfer learning and limited ground data, *Remote Sensing*, 14, 1809,
608 <https://doi.org/10.3390/rs14081809>, 2022.

609 Malingreau, J.-P.: Global vegetation dynamics: Satellite observations over Asia, *International Journal of Remote Sensing*, 7,
610 1121–1146, <https://doi.org/10.1080/01431168608948914>, 1986.

611 Manoochehr, S., Khoshmanesh, M., Ojha, C., Werth, S., Kerner, H., Grace Carlson, Sonam Futi Sherpa, Guang Zhai, and Jui-
612 Chi Lee: Persistent impact of spring floods on crop loss in U.S. Midwest, *Weather and Climate Extremes*, 34,
613 <https://doi.org/10.1016/j.wace.2021.100392>, 2021.

614 Marin, F. R., Zanon, A. J., Monzon, J. P., Andrade, J. F., Silva, E. H. F. M., Richter, G. L., Antolin, L. A. S., Ribeiro, B. S.
615 M. R., Ribas, G. G., Battisti, R., Heinemann, A. B., and Grassini, P.: Protecting the Amazon forest and reducing global
616 warming via agricultural intensification, *Nature Sustainability*, 1–9, <https://doi.org/10.1038/s41893-022-00968-8>, 2022.

617 Massey, R., Sankey, T. T., Congalton, R. G., Yadav, K., Thenkabail, P. S., Ozdogan, M., and Sánchez Meador, A. J.: MODIS
618 phenology-derived, multi-year distribution of conterminous U.S. crop types, *Remote Sensing of Environment*, 198, 490–503,
619 <https://doi.org/10.1016/j.rse.2017.06.033>, 2017.

620 Mei, Q., Zhang, Z., Han, J., Song, J., Dong, J., Wu, H., Xu, J., and Tao, F.: ChinaSoyArea10m: A dataset of soybean-planting
621 areas with a spatial resolution of 10 m across China from 2017 to 2021, *Earth System Science Data*, 16, 3213–3231,
622 <https://doi.org/10.5194/essd-16-3213-2024>, 2024.

623 Nafziger, E., Early-season soybean management for 2019: [https://farmdoc.illinois.edu/field-crop-](https://farmdoc.illinois.edu/field-crop-production/crop_production/early-season-soybean-management-for-2019.html)
624 [production/crop_production/early-season-soybean-management-for-2019.html](https://farmdoc.illinois.edu/field-crop-production/crop_production/early-season-soybean-management-for-2019.html), last access: 20 June 2024.

625 NASS CPR, Crop Progress Report: https://www.nass.usda.gov/Publications/National_Crop_Progress/, last access: 20 June
626 2024.

627 Olofsson, P., Foody, G. M., Stehman, S. V., and Woodcock, C. E.: Making better use of accuracy data in land change studies:
628 Estimating accuracy and area and quantifying uncertainty using stratified estimation, *Remote Sensing of Environment*, 129,
629 122–131, <https://doi.org/10.1016/j.rse.2012.10.031>, 2013.

630 Ouyang, Z., Jackson, R. B., McNicol, G., Fluet-Chouinard, E., Runkle, B. R. K., Papale, D., Knox, S. H., Cooley, S., Delwiche,
631 K. B., Feron, S., Irvin, J. A., Malhotra, A., Muddasir, M., Sabbatini, S., Alberto, M. C. R., Cescatti, A., Chen, C.-L., Dong, J.,
632 Fong, B. N., Guo, H., Hao, L., Iwata, H., Jia, Q., Ju, W., Kang, M., Li, H., Kim, J., Reba, M. L., Nayak, A. K., Roberti, D. R.,
633 Ryu, Y., Swain, C. K., Tsuang, B., Xiao, X., Yuan, W., Zhang, G., and Zhang, Y.: Paddy rice methane emissions across
634 Monsoon Asia, *Remote Sensing of Environment*, 284, 113335, <https://doi.org/10.1016/j.rse.2022.113335>, 2023.

635 Ozdogan, M. and Woodcock, C. E.: Resolution dependent errors in remote sensing of cultivated areas, *Remote Sensing of*
636 *Environment*, 103, 203–217, <https://doi.org/10.1016/j.rse.2006.04.004>, 2006.

637 Potapov, P., Hansen, M. C., Kommareddy, I., Kommareddy, A., Turubanova, S., Pickens, A., Adusei, B., Tyukavina, A., and
638 Ying, Q.: Landsat analysis ready data for global land cover and land cover change mapping, *Remote Sensing*, 12, 426,
639 <https://doi.org/10.3390/rs12030426>, 2020.

640 Potapov, P., Turubanova, S., Hansen, M. C., Tyukavina, A., Zalles, V., Khan, A., Song, X.-P., Pickens, A., Shen, Q., and
641 Cortez, J.: Global maps of cropland extent and change show accelerated cropland expansion in the twenty-first century, *Nature*
642 *Food*, 3, 19–28, <https://doi.org/10.1038/s43016-021-00429-z>, 2021a.

643 Potapov, P., Li, X., Hernandez-Serna, A., Tyukavina, A., Hansen, M. C., Kommareddy, A., Pickens, A., Turubanova, S., Tang,
644 H., Silva, C. E., Armston, J., Dubayah, R., Blair, J. B., and Hofton, M.: Mapping global forest canopy height through
645 integration of GEDI and Landsat data, *Remote Sensing of Environment*, 253, <https://doi.org/10.1016/j.rse.2020.112165>,
646 2021b.

647 Probst, P., Wright, M. N., and Boulesteix, A.-L.: Hyperparameters and tuning strategies for random forest, *Wiley*
648 *Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, 9, e1301, <https://doi.org/10.1002/widm.1301>, 2019.

649 Remelgado, R., Zaitov, S., Kenjabaev, S., Stulina, G., Sultanov, M., Ibrakhimov, M., Akhmedov, M., Dukhovny, V., and
650 Conrad, C.: A crop type dataset for consistent land cover classification in Central Asia, *Scientific Data*, 7, 250,
651 <https://doi.org/10.1038/s41597-020-00591-2>, 2020.

652 Roy, D. P., Li, Z., and K. Zhang, H.: Adjustment of Sentinel-2 Multi-Spectral Instrument (MSI) red-edge band reflectance to
653 nadir BRDF adjusted reflectance (NBAR) and quantification of red-edge band BRDF Effects, *Remote Sensing*, 9,
654 <https://doi.org/10.3390/rs9121325>, 2017a.

655 Roy, D. P., Li, J., Zhang, H. K., Yan, L., Huang, H., and Li, Z.: Examination of Sentinel-2A multi-spectral instrument (MSI)
656 reflectance anisotropy and the suitability of a general method to normalize MSI reflectance to nadir BRDF adjusted reflectance,
657 *Remote Sensing of Environment*, 199, 25–38, <https://doi.org/10.1016/j.rse.2017.06.019>, 2017b.

658 Som-ard, J., Immitzer, M., Vuolo, F., Ninsawat, S., and Atzberger, C.: Mapping of crop types in 1989, 1999, 2009 and 2019
659 to assess major land cover trends of the Udon Thani Province, Thailand, *Computers and Electronics in Agriculture*, 198,
660 107083, <https://doi.org/10.1016/j.compag.2022.107083>, 2022.

661 Song, X. P., Hansen, M. C., Stehman, S. V., Potapov, P. V., Tyukavina, A., Vermote, E. F., and Townshend, J. R.: Global land
662 change from 1982 to 2016, *Nature*, 560, 639–643, <https://doi.org/10.1038/s41586-018-0411-9>, 2018.

663 Song, X.-P., Potapov, P. V., Krylov, A., King, L., Di Bella, C. M., Hudson, A., Khan, A., Adusei, B., Stehman, S. V., and
664 Hansen, M. C.: National-scale soybean mapping and area estimation in the United States using medium resolution satellite
665 imagery and field survey, *Remote Sensing of Environment*, 190, 383–395, <https://doi.org/10.1016/j.rse.2017.01.008>, 2017.

666 Song, X.-P., Huang, W., Hansen, M. C., and Potapov, P.: An evaluation of Landsat, Sentinel-2, Sentinel-1 and MODIS data
667 for crop type mapping, *Science of Remote Sensing*, 3, 100018, <https://doi.org/10.1016/j.srs.2021.100018>, 2021a.

668 Song, X.-P., Hansen, M. C., Potapov, P., Adusei, B., Pickering, J., Adami, M., Lima, A., Zalles, V., Stehman, S. V., Di Bella,
669 C. M., Conde, M. C., Copati, E. J., Fernandes, L. B., Hernandez-Serna, A., Jantz, S. M., Pickens, A. H., Turubanova, S., and
670 Tyukavina, A.: Massive soybean expansion in South America since 2000 and implications for conservation, *Nature*
671 *Sustainability*, 4, 784–792, <https://doi.org/10.1038/s41893-021-00729-z>, 2021b.

672 Song, X.-P., Li, H., Potapov, P., and Hansen, M. C.: Annual 30 m soybean yield mapping in Brazil using long-term satellite
673 observations, climate data and machine learning, *Agricultural and Forest Meteorology*, 326,
674 <https://doi.org/10.1016/j.agrformet.2022.109186>, 2022.

675 Stehman, S. V.: Estimating area and map accuracy for stratified random sampling when the strata are different from the map
676 classes, *International Journal of Remote Sensing*, 35, 4923–4939, <https://doi.org/10.1080/01431161.2014.930207>, 2014.

677 Tanaka, T., Sun, L., Becker-Reshef, I., Song, X.-P., and Puricelli, E.: Satellite forecasting of crop harvest can trigger a cross-
678 hemispheric production response and improve global food security, *Communications Earth & Environment*, 4, 334,
679 <https://doi.org/10.1038/s43247-023-00992-2>, 2023.

680 Van Tricht, K., Degerickx, J., Gilliams, S., Zanaga, D., Battude, M., Grosu, A., Brombacher, J., Lesiv, M., Bayas, J. C. L.,
681 Karanam, S., Fritz, S., Becker-Reshef, I., Franch, B., Mollà-Bononad, B., Boogaard, H., Pratihast, A. K., Koetz, B., and
682 Szantoi, Z.: WorldCereal: A dynamic open-source system for global-scale, seasonal, and reproducible crop and irrigation
683 mapping, *Earth System Science Data*, 15, 5491–5515, <https://doi.org/10.5194/essd-15-5491-2023>, 2023.

684 Wang, S., Azzari, G., and Lobell, D. B.: Crop type mapping without field-level labels: Random forest transfer and unsupervised
685 clustering techniques, *Remote Sensing of Environment*, 222, 303–317, <https://doi.org/10.1016/j.rse.2018.12.026>, 2019.

686 Wang, S., Di Tommaso, S., Deines, J. M., and Lobell, D. B.: Mapping twenty years of corn and soybean across the US Midwest
687 using the Landsat archive, *Scientific Data*, 7, 307, <https://doi.org/10.1038/s41597-020-00646-4>, 2020.

688 Wang, Y., Feng, K., Sun, L., Xie, Y., and Song, X.-P.: Satellite-based soybean yield prediction in Argentina: A comparison
689 between panel regression and deep learning methods, *Computers and Electronics in Agriculture*, 221, 108978,
690 <https://doi.org/10.1016/j.compag.2024.108978>, 2024.

691 Wardlow, B. D. and Egbert, S. L.: Large-area crop mapping using time-series MODIS 250 m NDVI data: An assessment for
692 the U.S. Central Great Plains, *Remote Sensing of Environment*, 112, 1096–1116, <https://doi.org/10.1016/j.rse.2007.07.019>,
693 2008.

694 Wardlow, B. D., Egbert, S. L., and Kastens, J. H.: Analysis of time-series MODIS 250 m vegetation index data for crop
695 classification in the U.S. Central Great Plains, *Remote Sensing of Environment*, 108, 290–310,
696 <https://doi.org/10.1016/j.rse.2006.11.021>, 2007.

697 Woodcock, C. E., Allen, R., Anderson, M., Belward, A., Bindschadler, R., Cohen, W., Gao, F., Goward, S. N., Helder, D.,
698 Helmer, E., Nemani, R., Oreopoulos, L., Schott, J., Thenkabail, P. S., Vermote, E. F., Vogelmann, J., Wulder, M. A., and
699 Wynne, R.: Free access to Landsat imagery, *Science*, 320, 1011, <https://doi.org/10.1126/science.320.5879.1011a>, 2008.

700 Wright, C. K. and Wimberly, M. C.: Recent land use change in the Western Corn Belt threatens grasslands and wetlands,
701 *Proceedings of the National Academy of Sciences*, 110, 4134–4139, <https://doi.org/10.1073/pnas.1215404110>, 2013.

702 Xiong, J., Thenkabail, P. S., Gumma, M. K., Teluguntla, P., Poehnelt, J., Congalton, R. G., Yadav, K., and Thau, D.:
703 Automated cropland mapping of continental Africa using Google Earth Engine cloud computing, *ISPRS Journal of*
704 *Photogrammetry and Remote Sensing*, 126, 225–244, <https://doi.org/10.1016/j.isprsjprs.2017.01.019>, 2017.

705 Yan, L. and Roy, D. P.: Automated crop field extraction from multi-temporal Web Enabled Landsat Data, *Remote Sensing of*
706 *Environment*, 144, 42–64, <https://doi.org/10.1016/j.rse.2014.01.006>, 2014.

707 Yan, L. and Roy, D. P.: Conterminous United States crop field size quantification from multi-temporal Landsat data, *Remote*
708 *Sensing of Environment*, 172, 67–86, <https://doi.org/10.1016/j.rse.2015.10.034>, 2016.

709 Yang, Y., Wilson, L. T., and Wang, J.: A spatially explicit crop planting initiation and progression model for the conterminous
710 United States, *European Journal of Agronomy*, 90, 184–197, <https://doi.org/10.1016/j.eja.2017.08.004>, 2017.

711 Yang, Z., Diao, C., and Gao, F.: Towards scalable within-season crop mapping with phenology normalization and deep
712 learning, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 16, 1390–1402,
713 <https://doi.org/10.1109/JSTARS.2023.3237500>, 2023.

714 You, N., Dong, J., Huang, J., Du, G., Zhang, G., He, Y., Yang, T., Di, Y., and Xiao, X.: The 10-m crop type maps in Northeast
715 China during 2017–2019, *Scientific Data*, 8, 41, <https://doi.org/10.1038/s41597-021-00827-9>, 2021.

716 You, N., Dong, J., Li, J., Huang, J., and Jin, Z.: Rapid early-season maize mapping without crop labels, *Remote Sensing of*
717 *Environment*, 290, <https://doi.org/10.1016/j.rse.2023.113496>, 2023.

718 Zalles, V., Hansen, M. C., Potapov, P. V., Stehman, S. V., Tyukavina, A., Pickens, A., Song, X. P., Adusei, B., Okpa, C.,
719 Aguilar, R., John, N., and Chavez, S.: Near doubling of Brazil’s intensive row crop area since 2000, *Proceedings of the National*
720 *Academy of Sciences*, 116, 428–435, <https://doi.org/10.1073/pnas.1810301115>, 2019.

721 Zalles, V., Hansen, M. C., Potapov, P. V., Parker, D., Stehman, S. V., Pickens, A. H., Parente, L. L., Ferreira, L. G., Song, X.-
722 P., Hernandez-Serna, A., and Kommareddy, I.: Rapid expansion of human impact on natural land in South America since
723 1985, *Science Advances*, 7, eabg1620, <https://doi.org/10.1126/sciadv.abg1620>, 2021.

724 Zhong, L., Gong, P., and Biging, G. S.: Efficient corn and soybean mapping with temporal extendability: A multi-year
725 experiment using Landsat imagery, *Remote Sensing of Environment*, 140, 1–13, <https://doi.org/10.1016/j.rse.2013.08.023>,
726 2014.

727 Zhu, Z., Wang, S., and Woodcock, C. E.: Improvement and expansion of the Fmask algorithm: Cloud, cloud shadow, and
728 snow detection for Landsats 4–7, 8, and Sentinel 2 images, *Remote Sensing of Environment*, 159, 269–277,
729 <https://doi.org/10.1016/j.rse.2014.12.014>, 2015.