

# Global inventory of doubly substituted isotopologues of methane ( $\Delta^{13}\text{CH}_3\text{D}$ and $\Delta^{12}\text{CH}_2\text{D}_2$ )

**Authors:** Sara M. Defratyka<sup>1,2</sup>, Julianne M. Fernandez<sup>3,\*</sup>, Getachew A Adnew<sup>4</sup>, Guannan Dong<sup>5</sup>, Peter M.J. Douglas<sup>6</sup>, Daniel L. Eldridge<sup>7</sup>, Giuseppe Etiope<sup>8,9</sup>, Thomas Giunta<sup>10</sup>, Mojghan A. Haghnegahdar<sup>3</sup>, Alexander N. Hristov<sup>11</sup>, Nicole Hultquist<sup>3</sup>, Iñaki Vadillo<sup>12</sup>, Josue Jautzy<sup>13</sup>, Ji-Hyun Kim<sup>14</sup>, Jabrane Labidi<sup>15</sup>, Ellen Lalk<sup>16,\*\*</sup>, Wil Leavitt<sup>17,18,19</sup>, Jiawen Li<sup>18</sup>, Li-Hung Lin<sup>20</sup>, Jiarui Liu<sup>21,22</sup>, Lucía Ojeda<sup>12</sup>, Shuhei Ono<sup>16</sup>, Jeemin H. Rhim<sup>23</sup>, Thomas Röckmann<sup>24</sup>, Barbara Sherwood Lollar<sup>25</sup>, Malavika Sivan<sup>24</sup>, Jiayang Sun<sup>3,26</sup>, Gregory T. Ventura<sup>27</sup>, David T. Wang<sup>16,\*\*\*</sup>, Edward D. Young<sup>22</sup>, Naizhong Zhang<sup>28</sup> and Tim Arnold<sup>29,1</sup>

<sup>1</sup> School of GeoSciences, University of Edinburgh, Edinburgh, UK

<sup>2</sup> National Physical Laboratory, Hampton Road, London, UK

<sup>3</sup> Department of Geology, University of Maryland, College Park; College Park, Maryland, USA

<sup>4</sup> Terrestrial Ecosystem Analysis Group, Institute of Geography, People and Processes, Department of Geosciences and Resource management, Science Faculty, University of Copenhagen, Øster Voldgade 10 1350 Copenhagen K, Copenhagen, Denmark

<sup>5</sup> Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA

<sup>6</sup> Earth and Planetary Sciences and Geotop Research Centre, McGill University, Montreal, QC, Canada

<sup>7</sup> Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

<sup>8</sup> Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, Rome, Italy

<sup>9</sup> Faculty of Environmental Science and Engineering, Babes-Bolyai University, Cluj-Napoca, Romania

<sup>10</sup> Geo-Ocean (UMR6538), University Brest, CNRS, Ifremer, Plouzané, France

<sup>11</sup> Department of Animal Science, The Pennsylvania State University

<sup>12</sup> Department of Geology, Faculty of Science, University of Malaga, Malaga, Spain

<sup>13</sup> Geological Survey of Canada - Natural Resources Canada

<sup>14</sup> Marine Geology & Energy Division, Korea Institute of Geoscience and Mineral Resources, Daejeon, South Korea

<sup>15</sup> Université de Paris, Institut de Physique du Globe de Paris, CNRS, F-75005 Paris, France

<sup>16</sup> Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139 USA

<sup>17</sup> Department of Geology & Geophysics, University of Utah, Salt Lake City, UT, USA

<sup>18</sup> Department of Earth Sciences, Dartmouth College, Hanover, NH, USA

<sup>19</sup> Department of Chemistry, Dartmouth College, Hanover, NH, USA

<sup>20</sup> Department of Geosciences, National Taiwan University, Taipei 106, Taiwan

<sup>21</sup> Marine Science Institute, University of California, Santa Barbara, California 93106, USA

<sup>22</sup> Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles, California 90095, USA

<sup>23</sup> Department of Ecology, Evolution, and Marine Biology University of California, Santa Barbara

<sup>24</sup> Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University, Utrecht, the Netherlands

<sup>25</sup> Department of Earth Sciences, 22 Ursula Franklin St. University of Toronto, Toronto, ON Canada M5S 3B1

<sup>26</sup> Air Resources Laboratory, National Oceanic and Atmospheric Administration, College Park, Maryland 20740, USA

<sup>27</sup> Department of Geology, Saint Mary's University, Halifax, NS B3H 3C3, Canada

<sup>28</sup> Laboratory for Air Pollution / Environmental Technology, Empa, 8600 Dübendorf, Switzerland

<sup>29</sup> Department of Physical Geography and Ecosystem Science, Lund University, Sweden

\* Now at: Global Monitoring Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA,

\*\* Now at: U.S. Geological Survey, Woods Hole Coastal and Marine Science Center, Woods Hole, MA 02543, USA

\*\*\* Now at: U.S. Department of Energy, 1000 Independence Ave SW, Washington, DC 20585 USA

*Correspondence to:* Sara M. Defratyka (sara.defratyka@ed.ac.uk)

**Short summary (up to 500 characters).** Measurement of methane's doubly substituted isotopologues at natural abundances holds promise for better constraining the Earth's atmospheric CH<sub>4</sub> budget. We compiled 1475 measurements from field samples and laboratory experiments, conducted since 2014, to facilitate the differentiation of CH<sub>4</sub> formation pathways and processes, to identify existing gaps limiting application of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ , and to develop isotope ratio source signature inputs for global CH<sub>4</sub> flux modelling.

## **Abstract:**

Measurements of methane (CH<sub>4</sub>) molecules containing two rare isotopes ( $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$ ), also termed doubly substituted or 'clumped' isotopologues, have the potential to provide two additional isotopic dimensions to help investigate the mechanisms underlying global atmospheric trends in CH<sub>4</sub>. In this work, we summarise the current state of research on doubly substituted CH<sub>4</sub> isotopologues, with an emphasis on compiling results of all relevant work. The database comprises 1475 records compiled from the literature published until April 2025 (<https://dx.doi.org/10.5285/51ae627da5fb41b8a767ee6c653f83e6>). For field samples, 40% of records were sourced from natural gas reservoirs, while microbial terrestrial (e.g., agriculture, lake, wetland) samples account only for 12.5%. Lakes samples contribute 75% to collected microbial terrestrial samples. There is limited or no representation of samples coming from significant microbial CH<sub>4</sub> sources to the atmosphere, like wetlands, agricultural practices and landfills. To date, laboratory experiments were mostly focused on microbial (28% of samples from laboratory experiments) and pyrogenic (15%) methanogenesis or anaerobic (16%), and aerobic (8%) CH<sub>4</sub> oxidation, with only single study of photochemical oxidation via OH and Cl, which constitutes 5% of the laboratory experiments entries. The distinct ranges of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  values measured in these studies suggests their potential to improve our understanding of atmospheric CH<sub>4</sub>. This work provides an overview of the major gaps in measurements and identifies where further studies should be focussed to enable the highest impact on understanding global CH<sub>4</sub>.

## **1. Introduction**

Methane's bulk isotopic signatures (in particular  $\delta^{13}\text{C}-\text{CH}_4$ ), have been commonly used to constrain CH<sub>4</sub> emissions sources and budget changes (Basu et al., 2022; Lan et al., 2021; Menoud et al., 2022a; Sherwood et al., 2017; Turner et al., 2019). While the observed recent negative trend in  $\delta^{13}\text{C}-\text{CH}_4$  with increasing CH<sub>4</sub> mole fraction in the atmosphere implies a shift towards increasing microbial sources, the magnitude of this shift is difficult to quantify owing to the uncertainty in the isotopic source terms (Nisbet et al., 2019). Thus, additional independent tracers of CH<sub>4</sub> fluxes to the atmosphere would be useful to improve the understanding of global CH<sub>4</sub> changes.

The isotopologues  $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$ , referred to as doubly-substituted or "clumped" isotopologues, are thermodynamically more stable than the more abundant singly substituted CH<sub>4</sub> (i.e.,  $^{13}\text{CH}_4$  and  $^{12}\text{CH}_3\text{D}$ ). High precision measurements of the ratios of these rarer isotopologues present new tracer capabilities to quantify CH<sub>4</sub> sources and sinks (e.g., Douglas et al., 2017; Eiler, 2007; Haghnegahdar et al., 2017; Sivan et al., 2024; Stolper et al., 2014b; Young et al., 2017). The reported values,  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ , represent the measured isotopologue ratios ( $^{13}\text{CH}_3\text{D}/^{12}\text{CH}_4$  and  $^{12}\text{CH}_2\text{D}_2/^{12}\text{CH}_4$ , respectively) relative to their calculated values that assumes a random distribution of isotopes amongst the CH<sub>4</sub> isotopologues. This parameterization proves beneficial, as at thermodynamic isotopic equilibrium, the deviation in these isotopologue ratios from a purely random distribution is solely a function of temperature and it is independent from the bulk isotopic contents. Therefore, measurements of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  can constrain CH<sub>4</sub> formation temperatures, if the CH<sub>4</sub> has formed in thermodynamic isotopic equilibrium. An important aspect of this parameterization is that at

sufficiently high temperatures under thermodynamic isotopic equilibrium (where exchange of isotopes between isotopologues is fully reversible) the doubly substituted isotopic signature tends towards zero. At low temperatures, however, the abundance of clumped isotopes is much higher than expected from random distribution (e.g., Eldridge et al., 2019; Stolper et al., 2014a; Young et al., 2016).

When CH<sub>4</sub> is not in thermodynamic isotopic equilibrium, values of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  can reflect other physicochemical processes, such as their formation and consumption reactions (kinetic isotope effects, combinatorial effects, etc.), mixing of different sources, and physical transport processes such as molecular diffusion (e.g., Douglas et al., 2017; Gonzalez et al., 2019; Ono et al., 2014; Röckmann et al., 2016; Stolper et al., 2014b; Wang et al., 2024; Yeung, 2016; Young, 2019; Young et al., 2017). Therefore, measurements of doubly substituted isotopologues can provide additional analytical dimensions to distinguish between atmospheric sources (e.g., microbial, thermogenic, and abiotic CH<sub>4</sub>) and sinks (Chung and Arnold, 2021; Douglas et al., 2017; Haghnegahdar et al., 2017; Stolper et al., 2014a; Whitehill et al., 2017; Young, 2019). For example, it is currently understood that the  $\Delta^{13}\text{CH}_3\text{D}$  of atmospheric CH<sub>4</sub> is more sensitive to sources than sinks because it does not appear to be strongly affected by currently known sink reactions, while  $\Delta^{12}\text{CH}_2\text{D}_2$  is currently understood to be sensitive to both atmospheric CH<sub>4</sub> sources and sinks (Chung and Arnold, 2021; Haghnegahdar et al., 2017, 2023, 2024; Sivan et al., 2024; Whitehill et al., 2017). Thus, the atmospheric monitoring of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  has the potential to yield novel and unique insights into the temporal and spatial variations in atmospheric CH<sub>4</sub> source and sink reactions.

The first attempt to measure the rare CH<sub>4</sub> isotopologues from the ambient air was presented by Mroz et al. (1989), with further methods development refined by Ma et al. (2008) and Tsuji et al. (2012). The first precise measurements of doubly substituted CH<sub>4</sub>, specifically  $\Delta_{18}$  (combined  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ ) or  $\Delta^{13}\text{CH}_3\text{D}$  were published in 2014 (Ono et al., 2014; Stolper et al., 2014a, b). Young et al., (2017) reported on the first  $^{12}\text{CH}_2\text{D}_2$  data from laboratory and natural CH<sub>4</sub> sources. Since then, these measurements have become more relevant, particularly within the isotope geochemistry community. Measuring  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  from ambient air samples, however, is more challenging as it requires the collection and quantitative extraction of CH<sub>4</sub> from about 1000 L of air (1 m<sup>3</sup>). The first  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  measurements from the atmosphere, based on ambient air collections in Maryland (USA) and Utrecht (Netherlands), differed from model predictions of the atmosphere based on certain assumptions of source and sink reaction signatures (Chung and Arnold, 2021; Haghnegahdar et al., 2017, 2023; Sivan et al., 2024). The discrepancy could therefore come from either incorrectly assigned kinetic isotope effects associated with sink reactions or the assumptions regarding source signatures, or both (Haghnegahdar et al., 2023; Sivan et al., 2024; Wang et al., 2023b). This underscores the importance of obtaining improved constraints on source signatures and the isotope effects associated with sink reactions for improving the utility of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  in the study of atmospheric CH<sub>4</sub>.

For this study, we have compiled an open-source database (Defratyka et al., 2025) (<https://dx.doi.org/10.5285/51ae627da5fb41b8a767ee6c653f83e6>) of existing measurements of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ , including studies where only  $\Delta^{13}\text{CH}_3\text{D}$  was measured, from peer-reviewed scientific journal publications. The database contains almost 1500 values of doubly substituted isotope ratio measurements, from about 75 peer-reviewed scientific publications. The database is designed for utilization by the geochemistry and atmospheric science communities. This paper describes the collected  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  values that are included in the database. Our purpose is to present the current knowledge of doubly substituted isotopologues of CH<sub>4</sub> and identify existing gaps that presently limit our ability to apply  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  to understanding of atmospheric CH<sub>4</sub>.

## 2. CH<sub>4</sub> doubly substituted isotopologue ratios

### 2.1. $\Delta^{13}\text{CH}_3\text{D}$ and $\Delta^{12}\text{CH}_2\text{D}_2$ notations and calibration

A comprehensive review of the theory and nomenclature of doubly substituted isotopologue geochemistry is detailed in Eiler (2007, 2013), Wang et al., (2004) and Young et al. (2016, 2017). Briefly, doubly substituted isotopologue ratios of CH<sub>4</sub> are reported and parameterized as Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> values, defined to quantify a measured difference in the isotopologue ratios relative to a random distribution:

$$\Delta^{13}\text{CH}_3\text{D} = \frac{R_{\text{sample}}^{13}\text{CH}_3\text{D}}{R_{\text{stochastic}}^{13}\text{CH}_3\text{D}} - 1 \quad (1),$$

$$\Delta^{12}\text{CH}_2\text{D}_2 = \frac{R_{\text{sample}}^{12}\text{CH}_2\text{D}_2}{R_{\text{stochastic}}^{12}\text{CH}_2\text{D}_2} - 1 \quad (2).$$

Where:

$R_{\text{sample}}^{13}\text{CH}_3\text{D}$  and  $R_{\text{sample}}^{12}\text{CH}_2\text{D}_2$  are the measured isotopologue ratios of  $^{13}\text{CH}_3\text{D}/^{12}\text{CH}_4$  and  $^{12}\text{CH}_2\text{D}_2/^{12}\text{CH}_4$ , respectively,

and

$R_{\text{stochastic}}^{13}\text{CH}_3\text{D}$  and  $R_{\text{stochastic}}^{12}\text{CH}_2\text{D}_2$  are the calculated isotopologue ratios of  $^{13}\text{CH}_3\text{D}/^{12}\text{CH}_4$  and  $^{12}\text{CH}_2\text{D}_2/^{12}\text{CH}_4$ , respectively, based on the assumption of a random distribution of isotopes amongst all stable isotopologues.

As an effect, the isotopologue ratio approaches that based on a random distribution under high-temperature equilibrium conditions, which by definition results in Δ<sup>13</sup>CH<sub>3</sub>D or Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> values of zero (e.g., Douglas et al., 2016; Eiler, 2007, 2013; Stolper et al., 2014a; Young, 2019). It should be noted that non-zero values of Δ<sup>13</sup>CH<sub>3</sub>D or Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> can result from the simple mixing of two separate CH<sub>4</sub> pools with distinct bulk isotopic compositions, without any chemical or physical processes inducing isotopic fractionation (e.g., Young et al., 2016).

In this paper, the terms ‘enriched’ and ‘depleted’ refer to comparative values of Δ<sup>13</sup>CH<sub>3</sub>D or Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> – higher numbers as enriched and lower numbers as depleted – for example when comparing samples of CH<sub>4</sub>, products and reactants of a chemical reaction, or the evolution of CH<sub>4</sub> in a physical process.

## 2.2. Existing instrumentation

The measurement of Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> is resource intensive, requiring specialised facilities that are currently not widely available (e.g., Eiler, 2007; Liu et al., 2024b; Ono et al., 2014a; Sivan et al., 2024; Stolper et al., 2014a; Young et al., 2017). Magnetic sector High Resolution Isotope Ratio Mass Spectrometry (HR-IRMS) is the most common method to measure Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> (Dong et al., 2020; Eldridge et al., 2019; Haghnegahdar et al., 2023; Liu et al., 2024b; Sivan et al., 2024; Stolper et al., 2014a; Sun et al., 2023; Thiagarajan et al., 2020; Wang et al., 2023a; Young et al., 2016, 2025; Zhang et al., 2021). The first magnetic sector HR-IRMS instrument developed for this purpose was the non-commercial prototype model of the Thermo Scientific 253 Ultra HR-IRMS (developed and installed solely at the California Institute of Technology) that was able to measure a value of the combined <sup>13</sup>CH<sub>3</sub>D and <sup>12</sup>CH<sub>2</sub>D<sub>2</sub> abundances via a parameter defined as Δ<sub>18</sub> (Eiler et al., 2013; Stolper et al., 2014 a,b; Stolper et al., 2015). A large-radius gas-source multiple-collector isotope ratio mass spectrometer capable of operating up to a mass resolving power (MRP) of 80,000 (Panorama, Nu Instrument) was the first developed HR-IRMS to measure separately Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> (Young et al., 2016, 2017). This was followed by the commercially-available production model of the Thermo Scientific Ultra HR-

IRMS that can also measure  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  and routinely achieves a MRP of 30-35,000 (e.g., Eldridge et al., 2019; Thiagarajan et al., 2020; Zhang et al., 2021; Wang et al. 2023a; Sivan et al., 2024). The obtained MRP allows to achieve precise measurements for sample of >2 mL STP (standard temperature and pressure) of  $\text{CH}_4$  ( $\sim 80 \mu\text{mol}$ ) for Panorama (e.g., Labidi et al., 2020) and  $3 \pm 1$  mL STP for Ultra (Sivan et al., 2024). Measurements of smaller volume of  $\text{CH}_4$  sample result in larger uncertainties caused by degraded counting statistic. The detailed description of the performance of these instruments and measurement protocols for different laboratories can be found in the cited references above.

Distinct from mass spectrometry, measurements of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  are also possible owing to developments in infrared absorption spectroscopy using quantum cascade lasers (TILDAS, Aerodyne Research) operated in near room temperature with narrow line widths and high power (Chen et al., 2022; Gonzalez et al., 2019; Ono et al., 2014; Prokhorov and Mohn, 2022; Zhang et al., 2025). The first TILDAS instrument to achieve high precision  $\Delta^{13}\text{CH}_3\text{D}$  measurements was demonstrated at the Massachusetts Institute of Technology in 2014 (Ono et al., 2014).  $\Delta^{13}\text{CH}_3\text{D}$  measurement by the TILDAS instrument are achieved using the absorption line in a spectral region around  $8.6 \mu\text{m}$ , as there are fewer interferences from hot bands (Ono et al., 2014). Gonzalez et al. (2019) presented a possibility to implement TILDAS to measure  $\Delta^{12}\text{CH}_2\text{D}_2$  with precision of 0.5 ‰. Routinely, TILDAS measurements requires 10 mL of  $\text{CH}_4$  for  $\Delta^{13}\text{CH}_3\text{D}$  measurements and 20 mL for  $\Delta^{12}\text{CH}_2\text{D}_2$  (e.g., Gonzalez et al., 2019; Ono et al., 2014). Recently, Zhang et al. (2025) were able to reduce the required volume of  $\text{CH}_4$  to 3-7 mL STP for  $\Delta^{13}\text{CH}_3\text{D}$  and to 10 mL STP for  $\Delta^{12}\text{CH}_2\text{D}_2$ , via further instrument optimization.

HR-IRMS signal stability of the detected ions at very low ion currents is key to enable precise isotope ratio measurement through signal acquisition over several hours or even days (e.g., Sivan et al., 2024; Stolper et al., 2014a; Young et al., 2016). Across instrumentation, internal precision and external reproducibility are comparable between laboratories and instruments, achieving down to 0.35 ‰ for  $\Delta^{13}\text{CH}_3\text{D}$  and 1.35 ‰ for  $\Delta^{12}\text{CH}_2\text{D}_2$ , depending on the measurement technique. The TILDAS and Panorama systems were cross-calibrated on the same set of carbon and hydrogen isotopically characterised laboratory working standards for  $\text{CH}_4$  to ensure accuracy between different analytical systems (Ono et al., 2014; Young et al., 2017; Zhang et al., 2025).

At thermodynamic isotopic equilibrium,  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  values can be linked to a  $\text{CH}_4$  formation temperature via monotonic functions, presented in Table S1 (Beaudry et al., 2021; Douglas et al., 2017; Eldridge et al., 2019; Gruen et al., 2018; Liu and Liu, 2016; Ono et al., 2014; Stolper et al., 2014a; Thiagarajan et al., 2020; Webb and Miller, 2014; Young et al., 2017; Zhang et al., 2021). Different theoretical calculations have been used to obtain these relationships but discrepancies among them are smaller than the current analytical uncertainties. Currently, equilibrated gas experiments along with these theoretical calculations are the basis for calibrating  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  measurements via either magnetic sector HR-IRMS or laser spectroscopy (Eldridge et al., 2019; Liu et al., 2024b; Ono et al., 2014; Sivan et al., 2024; Stolper et al., 2014a; Wang et al., 2015).

### 2.2.1. Samples extraction and purification


Quantitative extraction and complete purification of  $\text{CH}_4$  from natural samples is currently necessary to attain the required precision and accuracy to detect differences in clumped isotopic composition (Eiler, 2007; Prokhorov and Mohn, 2022; Safi et al., 2024; Sivan et al., 2024; Sun et al., 2023; Young et al., 2017). Two main methods have been applied so far across laboratories. One employs cryogenic trapping at near absolute zero temperature using a Helium cryostat (Stolper et al., 2014a) and the other have used chromatographic separations techniques (Young et al., 2017).

Measuring doubly substituted isotopologues in ambient air is a major analytical challenge. Since krypton has a similar concentration in the atmosphere and boiling point as CH<sub>4</sub> (Kr: 1.14 ppm in the atmosphere, -153.4 °C boiling point; CH<sub>4</sub>: 1.93 ppm, -161.5 °C), it makes separation by fractional distillation alone impossible. Recently, combined gas chromatography and cryogenic methods were successfully implemented to purify CH<sub>4</sub> from 10<sup>2</sup>-10<sup>3</sup> litres of ambient air to measure both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ . These approaches generally involve the pumping of large volumes of air through sequential cryogenic traps that selectively isolate CH<sub>4</sub> from other contaminants using established absorbents (Haghnegahdar et al., 2023; Sivan et al., 2024).

### 3. Database methods and description

#### 3.1. Data gathering

The compilation of this doubly substituted CH<sub>4</sub> isotopologues database is inspired by similar efforts of existing databases for bulk isotopes of CH<sub>4</sub> (Lan et al., 2021; Menoud et al., 2022a, b; Sherwood et al., 2017, 2021). To verify if the compiled data compares well with previous studies, figure 1 and table 1 present bulk isotopes from this database in the reference to previously reported  $\delta^{13}\text{C-CH}_4$  and  $\delta\text{D-CH}_4$  (Menoud et al., 2022a; Sherwood et al., 2021). Across compared group types, our additional bulk isotope ratio data fall within the established ranges. Fossil fuel and thermogenic source signatures overlap, however, they are not strictly equivalent. Thermogenic CH<sub>4</sub> in our dataset is slightly enriched ( $\delta^{13}\text{C-CH}_4$ :  $-39.0 \pm 9.6\text{‰}$ ;  $\delta\text{D-CH}_4$ :  $-169.2 \pm 41.9\text{‰}$ ), compared to fossil fuel. For the comparison, only terrestrial microbial (e.g., agriculture, lakes, wetlands) from this database is compared with previously compiled data and shows strong agreement with the range of previous microbial samples, with depleted  $\delta^{13}\text{C-CH}_4$  and  $\delta\text{D-CH}_4$  values ( $\delta^{13}\text{C-CH}_4$ :  $-62.9 \pm 13.2\text{‰}$ ;  $\delta\text{D-CH}_4$ :  $-298.1 \pm 47.7\text{‰}$ ). Pyrogenic methane, though represented by only two samples in the new database, shows  $\delta^{13}\text{C-CH}_4$  and  $\delta\text{D-CH}_4$  values consistent with previous studies. This alignment supports the representativeness of our inferred doubly substituted CH<sub>4</sub> isotopologues ratio source signatures for use alongside the bulk isotope ratios in global modelling of the CH<sub>4</sub> budget. Our database also provides further additional measurements of the bulk isotopes to aid in further work to refine the source signatures  $\delta^{13}\text{C-CH}_4$  and  $\delta\text{D-CH}_4$ .



**Figure 1.** Database entries plotted as  $\delta^{13}\text{C-CH}_4$  versus  $\delta\text{D-CH}_4$  alongside the Menoud et al., 2022a database. Error bars are taken from original studies. ff: fossil fuels, Tr: thermogenic, M: microbial, Pr: pyrogenic.

**Table 1.** Comparison across the three databases of  $\delta^{13}\text{C-CH}_4$  and  $\delta\text{D-CH}_4$  by group type. The mean value is reported with  $\pm 1$  standard deviation, and minimum and maximum values in brackets.

Group type	$\delta^{13}\text{C-CH}_4$			$\delta\text{D-CH}_4$		
	samples	median (‰)	mean (‰)	samples	median (‰)	mean (‰)

fossil fuels			-45.5 ± 9.1			-185.5 ± 38.7
Menoud et al. 2022	707	-44.2	[-82.1; -18.3]	394	-185.3	[-355.0; -63.8]
fossil fuels			-44.9 ± 10.6			-196.1 ± 48.6
Sherwood et al. 2021	9477	-43.0	[-87.0; -14.8]	3371	-191.7	[-415.0; -62.0]
thermogenic			-39.0 ± 9.6			-169.2 ± 41.9
Defratyka. et al. 2025	309	-38.3	[-73.0; -21.6]	309	-159.7	[-300.2; -100.8]
microbial			-58.5 ± 8.5			-309.7 ± 50.4
Menoud et al. 2022	471	-58	[-96.1; -36.5]	187	-307.1	[-472.0; -93.2]
microbial			-61.6 ± 6.9			-304.0 ± 36.6
Sherwood et al. 2021	131	-62.4	[-79.6; -45.5]	20	-304.0	[-358.0; -205.0]
microbial			-62.9 ± 13.2			-298.1 ± 47.7
Defratyka. et al. 2025	120	-66.8	[-77.7; 4.2]	120	-294.7	[-383.5; -90.5]
pyrogenic			-25.9 ± 7.7			-176.7 ± 59.0
Menoud et al. 2022	42	-27.2	[-42.7; -9.0]	11	-192.0	[-285.0; -81.0]
pyrogenic			-26.0 ± 5.3			-21.8 ± 15.5
Sherwood et al. 2021	29	-26.9	[-33.4; -12.5]	4	-208.0	[-232.0; -195.0]
pyrogenic			-27.7 ± 1.6			-248.6 ± 10.7
Defratyka. et al. 2025	2	-27.7	[-28.8; -26.5]	2	-248.6	[-256.1; -241.0]

The references included in the database of doubly substituted CH<sub>4</sub> isotopologues comprise mostly peer-reviewed articles, with a smaller percentage from conference papers. The aggregated studies were carried out between 2014 and 2025 across 10 laboratories worldwide. As the aim of this study is to include all existing studies of doubly substituted isotopologue ratios, we also incorporated results from laboratory experiments, and of CH<sub>4</sub> dissolved in water (i.e. in oceans, wetlands, and inland waters), which were not included in bulk isotopes databases.

### 3.1.1. The structure of the database

For efficient utilization of the database, we start with parameters (column names) from the databases of Sherwood et al., (2017, 2021) and Menoud et al., (2022). Then, we added the parameters to better represent the characteristics of doubly substituted isotope ratio measurements. Selected parameters are described in the metadata of the database (<https://dx.doi.org/10.5285/51ae627da5fb41b8a767ee6c653f83e6>). Collection and analysis dates, along with instrument and measurement laboratory are included to facilitate comparison between studies. For each entry of  $\Delta_{18}$ ,  $\Delta^{13}\text{CH}_3\text{D}$  or  $\Delta^{12}\text{CH}_2\text{D}_2$ , the number of samples, measured value, uncertainty, and type of uncertainty are provided. The parameter “other tracers” was added to include information about other tracers collected alongside doubly substituted isotopologues and bulk isotope ratio measurements of CH<sub>4</sub>. This parameter can be used to filter and group data for the further processing by database users. We also added the “lab field” parameter to make it easier to filter the database based on whether the sample was collected in the field or obtained from a laboratory experiment.

For samples collected from the field, we provided exact location (latitude and longitude), coming from the original article or approximate location, estimated based on geographical information in the article. The parameter “coordinates from primary source” was added to indicate if sampling location was given in the original article. We used the parameters documented by Menoud et al., (2022a) to describe the CH<sub>4</sub> source for field samples: group type, group, category and subcategory but with modifications to better reflect properties of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  studies conducted so far (table 2). For example, in group type, we divided microbial sources into three categories: microbial terrestrial, microbial fossil fuels (microbial ff) and microbial marine. Additionally, we incorporated a parameter “sources specification” to add any information coming from the primary studies’ publications that did not match the already included source parameters (e.g., thermodynamic disequilibrium or equilibrium, natural gas maturity, sources mixture). Parameters: “sample type”, “reservoir type”, “depth type” (i.e., unit of

reservoir depth from original paper) and “depth” were included for the description of field sampling conditions.

Whenever possible, we connected these groups and categories to the broadly used Selected Nomenclature for Air Pollution (SNAP) and Intergovernmental Panel on Climate Changes (IPCC, guidelines 2006) emissions categories for field samples (table 2). The Emissions Database for Global Atmospheric Research (EDGAR) inventories are compatible with IPCC nomenclatures, which facilitates implementation of the database and comparison with existing emissions inventories (details in section 4.3.1). In the database, samples from laboratory experiments, ambient air, and volcano (both mud volcano and steam volcano) measurements are not linked to SNAP and IPCC categories. Also, the SNAPP and IPCC categories were not allocated to groundwater nor deep marine samples (i.e., marine seeps, sediments, and pore fluid), as they represent insignificant sources of CH<sub>4</sub> to the atmosphere.

**Table 2.** Group type, group, category, and subcategory of CH<sub>4</sub> sources for field samples with SNAP and IPCC categories, based on source categories from Menoud et al. (2022).

GROUP TYPE	GROUP	CATEGORY	SUB_CATEGORY	SNAP	IPCC 2006
abiotic	exploitation	oil non-associated	natural gas	5	1B2
		metal mine	groundwaters	-	-
	seeps	marine; temperate; volcanoes	hydrothermal vent, marine seep; hyperalkaline spring, hot spring, spring; mud volcano	-	-
ambient air	urban background	-	-	-	-
	mixture with CH <sub>4</sub> source	-	-	-	-
	clean background	-	-	-	-
microbial terrestrial	agriculture	rice paddies	rice paddies	10	3C7
		ruminants	dairy cow	10	3A1
	exploitation	metal mine	groundwater	-	-
	seeps	temperate; volcanoes	groundwater, spring; mud volcano	-	-
	wetlands	polar (incl. boreal), temperate	lake, pond, swamp	11	3B4
microbial fossil fuel (microbial ff)	exploitation	coal	coal seam gas	5	1B2
		biodegradation of oil, conventional	gas installation, natural gas, oil field	5	1B2
microbial marine	sediment	marine	marine sediment, pore fluid	-	-
	seeps	marine	cold seep, marine seep, pockmark	-	-
mixture	exploitation	conventional, unconventional, shale, oil non-associated, oil associated, coal	gas installation, natural gas, oil field, coal seam gas	5	1B2
		metal mine	groundwater	-	-
	sediment	marine	marine sediment	-	-
		marine	marine seep	-	-
	seeps	temperate	groundwater, hyperalkaline spring	-	-
		volcanoes	mud volcano, steam volcano	-	-
		polar (incl. boreal)	lake	11	3B4
	wetlands				
others	exploitation	conventional	gas installation, natural gas	5	1B2
		metal mine	groundwater	-	-
	sediment	marine	marine sediment	-	-



	seeps	temperate; volcanoes	groundwater; hydrothermal,	-	-
	vehicle exhaust	-	steam vent, mud volcano, spring		
			-	7	1A3
	wetlands	polar (incl. boreal)	lake	11	3B4
pyrogenic	fossil fuel burning, biomass burning	charcoal, oak logs	biomass burning	11	3C1
thermogenic	exploitation	conventional, conventional oil associated, conventional oil non-associated, unconventional oil associated, unconventional oil non-associated, oil associated, oil non-associated, shale, unconventional shale	gas installation, natural gas	5	1B2
	sediment	marine; quartz-hosted inclusions	marine sediment; natural gas	-	-
	seeps	marine volcanoes	hydrothermal vent, marine seep	-	-
			hydrothermal, steam vent, mud volcano	-	-
	wetlands	polar (incl. boreal)	lake	11	3B4

For samples coming from laboratory experiments, we added a specification of the type of laboratory experiment (e.g., abiotic or microbial methanogenesis, pyrolysis experiment, AOM or AeOM methanotrophy) in the group type column (table 3). Also, parameters “lab experiment type” and “lab experiment detail” were added to include details of conducted experiments. “Catalytic equilibration” experiments are focused on defining the thermal equilibration curve, used for the instruments calibration (Eldridge et al., 2019; Liu et al., 2024b; Ono et al., 2014; Wang et al., 2019; Young et al., 2017).

**Table 3.** Group type and laboratory experiment type for laboratory experiment samples

GROUP TYPE	LAB EXPERIMENT TYPE
abiotic methanogenesis	high temperature abiotic low temperature abiotic
microbial methanogenesis	acetoclastic hydrogenotrophic methoxydotrophic methylotrophic methylphosphonate
pyrogenic methanogenesis	alkane pyrolysis coal pyrolysis ethane pyrolysis hydrous pyrolysis nonhydrous pyrolysis propane pyrolysis shale pyrolysis
AeOM methanotrophy	pure culture
AOM methanotrophy	enrichment culture field samples incubation
photochemical oxidation	control sample Cl oxidation OH oxidation

catalytic equilibration	bracketing calibration
mixing experiment	-
sediment incubation	aquatic environment sediment microbially enhanced coal bed CH <sub>4</sub> wetland soil

Due to variations in measurement protocols across laboratories, uncertainties are reported in different ways and therefore we reported uncertainty per entry as described in the database. Most of the laboratories report one or two internal standard errors (int SE) to reflect precision based on HR-IRMS counting statistics (e.g., Ash et al., 2019; Douglas et al., 2016, 2017; Thiagarajan et al., 2020; Wang et al., 2023a; Young et al., 2017). Others use external reproducibility, expressed as one or two external standard deviations (ext SD) (Eldridge et al., 2019; Giunta et al., 2021; Wang et al., 2024a; Warr et al., 2021a). When one sample is measured more than once, one SE or two SE are reported as uncertainty in the database (Stolper et al., 2015; Wang et al., 2018; Wang et al., 2024). For some studies, uncertainty is reported as 95% confidence intervals (95% CI) (e.g., Beaudry et al., 2021; Lalk et al., 2024; Ono et al., 2014).


As studies were made over time by different laboratories, not all required database parameters were included in existing peer-reviewed articles. For the future, proposed parameters should ideally be published with data. Additionally, a consistent description of CH<sub>4</sub> sources, group type, group, category, subcategory and laboratory experiment type, using the parameters proposed in table 2 and 3 is encouraged to facilitate interpretation and intercomparison between laboratories and methods.

## 4. Results and discussion

### 4.1. Data summary


Out of all data entries, field samples comprise 958 entries, while 517 entries come from laboratory experiments. Of these, 53% of entries report only  $\Delta_{18}$  or  $\Delta^{13}\text{CH}_3\text{D}$ . Potentially, the lack of  $^{12}\text{CH}_2\text{D}_2$  measurements can hinder data interpretation, especially for microbial, abiotic or mixed samples, where  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  can be modified differently (e.g., Douglas et al., 2017; Giunta et al., 2019; Gruen et al., 2018; Thiagarajan et al., 2020; Warr et al., 2021; Young et al., 2016, 2017). To avoid data misinterpretation, other tracers, for example radiocarbon or seismic reflection data, must be measured alongside to  $\Delta^{13}\text{CH}_3\text{D}$  (e.g., Chowdhury et al., 2024; Douglas et al., 2020).

Regarding the parameter “group type”, thermogenic samples contribute 32% to the field samples, while there is low representation of pyrogenic samples (0.21% of field samples) (figure 2). “Others” is a broad group type of field samples with ambiguous origin from various sources (e.g., natural gas, groundwaters from metal mines, marine and mud volcano samples), where it was not possible to clearly determine the group type based on isotopes and other tracers. Hypothesized origins of these samples are given as ‘source specification’ parameter in the database. Also, vehicle exhaust samples are classified as “others”, as different processes can cause CH<sub>4</sub> emissions from the exhaust (Sun et al., 2025b). Additionally, for samples where two different sources of CH<sub>4</sub> were mixed, indicated as group type ‘mixture’, more information on the type of mixture is added under the parameter “source specification” in the database. For ambient air “group type”, distinction between background samples and mixture of ambient air and gas coming from CH<sub>4</sub> source (e.g., gas sample collected above wetland, (Haghnegahdar et al., 2024; Sun et al., 2025b)) was made using the “group” parameter.



**Figure 2.** Frequency of entries for field samples categorised by a) group type and b) subcategories.


The distribution of measurements in  $\Delta^{13}\text{CH}_3\text{D}$  versus  $\Delta^{12}\text{CH}_2\text{D}_2$  space is presented in figure 3, both for field samples and laboratory experiments. To simplify data interpretation, field samples categorized as “others” or “mixture” are omitted. Also, samples where ambient air is mixed with the gas from  $\text{CH}_4$  source are omitted. The majority of thermogenic samples fall close to the thermodynamic isotopic equilibrium curve, with a few samples having more depleted  $\Delta^{12}\text{CH}_2\text{D}_2$  than predicted (details in section 4.2.). Microbial marine and microbial ff samples are near or at thermodynamic isotopic equilibrium but with some enrichment relative to equilibrium observed. Most of the microbial terrestrial samples (e.g., lakes, wetlands or agriculture) are clearly depleted in both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ , relative to the equilibrium. Different ratios for microbial terrestrial compared to microbial ff and microbial marine suggests different methanogenesis reactions or additional processes, such as methanotrophy or mixed patterns of microbial carbon cycling within in these environments (details in section 4.2.). Regarding abiotic  $\text{CH}_4$ , most of the samples are out of thermodynamic isotopic equilibrium (e.g., Douglas et al., 2020; Labidi et al., 2020; Lin et al., 2023; Young et al., 2017). It must be noted, that abiotic  $\text{CH}_4$  is empirically one of the least well characterized endmembers, both in terms of field and laboratory studies.



**Figure 3.** Database entries plotted as  $\Delta^{13}\text{CH}_3\text{D}$  versus  $\Delta^{12}\text{CH}_2\text{D}_2$ . Error bars are taken from original studies (details in section 3.1.1). a) fields samples based on 247 entries, where samples categorized as “others”, “mixture” and “ambient air mixed with  $\text{CH}_4$  source” are omitted for simplicity. b) laboratory experiments based on 210 entries. Laboratory experiments with deuterium-enriched water substrate (Gruen et al., 2018; Li et al., 2024, 2025a; Taenzer et al., 2020) are not included as they do not appear under normal incubation or environmental conditions. A solid black line represents the thermodynamic isotopic equilibrium curve, using equations from Young et al., (2017).


For laboratory experiments, the deviation from thermodynamic isotopic equilibrium depends on the studied methanogenesis pathway or the type of methanotrophy (aerobic (AeOM) versus anaerobic (AOM) CH<sub>4</sub> oxidation) (details in section 4.2.). For example, AOM methanotrophy experiments show a large enrichment for both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  (Liu et al., 2023; Ono et al., 2021). Notably, Gruen et al., (2018), Li et al., (2024, 2025a), and Taenzer et al., (2020), carried out incubations with deuterium-enriched substrate to explore mechanisms behind combinatorial effects. Thus, observed clumped isotopologues do not represent the isotopic values of natural-occurring microbial CH<sub>4</sub> and should be carefully re-interpreted.

Regarding pyrogenic methanogenesis, some samples have doubly-substituted isotope ratio compositions consistent with thermodynamic isotopic equilibrium, while others create more depleted values, due to a combination of kinetic isotope effects, combinatorial effects, and varying degrees of hydrogen isotope exchange (Dong et al., 2021; Eldridge et al., 2023; Shuai et al., 2018a). The abiotic synthesis of CH<sub>4</sub> in laboratory-controlled experiments shows enriched  $\Delta^{13}\text{CH}_3\text{D}$ , consistent with thermodynamic isotopic equilibrium, associated with systematically depleted  $\Delta^{12}\text{CH}_2\text{D}_2$ , due to combinatorial effects (Young et al., 2017, Labidi et al., 2024).



**Figure 4.** Frequency of entries for laboratory experiments categorised by group type

About 27% of the laboratory experimental entries come from studies on microbial methanogenesis, focused on various pure cultures of methanogenic archaea (e.g., acetoclastic, hydrogenotrophic and methylotrophic methanogenesis) (figure 4) (Douglas et al., 2016, 2020; Giunta et al., 2019; Gruen et al., 2018; Rhim and Ono, 2022; Stolper et al., 2015; Warr et al., 2021a; Young et al., 2017). Notably, Li et al. (2025a) conducted methanogenesis experiment where few data points come from extremely deuterium-enriched water ( $\delta\text{D}$  of water about 3000 ‰ and 8000 ‰). Such high  $\delta^2\text{H}$  of water cannot be found in the nature, thus obtained CH<sub>4</sub> has very atypical isotopic values (figure 5).



**Figure 5.**  $\Delta^{13}\text{CH}_3\text{D}$  versus  $\Delta^{12}\text{CH}_2\text{D}_2$  for microbial methanogenesis laboratory experiments. Laboratory experiments with deuterium-enriched water substrate (Gruen et al., 2018; Li et al., 2024, 2025a; Taenzer et al., 2020) are not included as they do not appear under normal incubation or environmental conditions.

4.4% of laboratory samples, classified as sediment incubation, were collected in the field and incubated in controlled laboratory conditions (Douglas et al., 2020; Haghnegahdar et al., 2023, 2024; Ijiri et al., 2018; Wang et al., 2024a). A single laboratory experiment focused on photochemical oxidation by OH and Cl was also conducted, however, only  $\Delta^{13}\text{CH}_3\text{D}$  was measured (Whitehill et al., 2017). A laboratory experiment focused on mixing of two  $\text{CH}_4$  sources, containing different bulk isotopic compositions, was conducted to confirm mixing curve delivered from theoretical calculation, related to the definition of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  (Young et al., 2016).

#### 4.2. State of knowledge about $\text{CH}_4$ doubly-substituted isotopologue ratios

Methane is produced at the surface and in subsurface environments via biogenic (microbial), thermogenic, or abiotic processes, while the majority of the  $\text{CH}_4$  emitted to the atmosphere comes from microbial, thermogenic, and pyrolytic (biomass and biofuel burning) sources (e.g., Saunio et al., 2025; Schoell, 1988; Stolper et al., 2018). Thermogenic  $\text{CH}_4$  forms by the thermally-activated breakdown of organic molecules, where ‘primary thermogenic’ is a term used to describe  $\text{CH}_4$  produced from kerogen and ‘secondary thermogenic’ is used to describe the breakdown of long-chain hydrocarbons (e.g., Lalk et al., 2023; Stolper et al., 2018). Stolper et al. (2014b) proposed that thermogenic  $\text{CH}_4$  is predominantly in thermodynamic isotopic equilibrium at its formation temperature, which was supported by studies focused on natural gas or volcanic samples (Beaudry et al., 2021; Douglas et al., 2016, 2017; Jiang et al., 2024; Kim et al., 2023; Rumble et al., 2018; Stolper et al., 2014b, 2015, 2018; Thiagarajan et al., 2020; Wang et al., 2015; Xie et al., 2021; Young et al., 2017). Formation temperatures calculated from doubly substituted isotope ratio measurements can help to determine the natural gas maturity and distinguish “atypical” thermogenic gas (from shallow or immature systems to deep or over-mature systems) from abiotic  $\text{CH}_4$  (Jiang et al., 2024; Kim et al., 2023; Li et al., 2025b; Shuai et al., 2018b). Some exceptions of doubly substituted isotope ratios deviating from thermodynamic isotopic equilibrium were observed from unconventional, oil-non-associated or oil-associated gas reservoirs (figure 6) (Douglas et al., 2017; Kim et al., 2023; Lalk et al., 2022; Stolper et al., 2018; Xie et al., 2021), which is consistent with laboratory pyrolysis experiments and gas generation models implying at least partly kinetically-driven signatures (Dong et al., 2021; Eldridge et al., 2023; Shuai et al., 2018a; Xia and Gao, 2019). For low maturity or oil-associated natural gas, a contribution from microbial sources can occur, for example due to  $\text{CH}_4$  generation during oil biodegradation (e.g., secondary microbial  $\text{CH}_4$ ). However, the likelihood that microbial  $\text{CH}_4$  has both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  within the thermogenic range remains low (Giunta et al., 2019; Lalk et al., 2022; Thiagarajan et al., 2020; Xie et al., 2021).

Microbial  $\text{CH}_4$  is produced by microorganisms via three main pathways: hydrogenotrophic, acetoclastic, and methylotrophic methanogenesis, with the first two being the predominant (Conrad, 2005; Thauer, 1998). Typically, subsurface microbial  $\text{CH}_4$  from geological basins is mostly generated through the hydrogenotrophic pathway, where doubly substituted isotope ratios tend towards thermodynamic isotopic equilibrium (figure 3 and 5) (Ash et al., 2019; Douglas et al., 2016, 2017, 2020; Giunta et al., 2019; Shuai et al., 2021; Stolper et al., 2015; Thiagarajan et al., 2020; Wang et al., 2024a; Warr et al., 2021a; Young et al., 2017). Studies of pore water from the Michigan Basin, showed that deep subsurface  $\text{CH}_4$  can also be generated by acetoclastic methanogenesis at thermodynamic isotopic equilibrium for  $^{13}\text{CH}_3\text{D}$  but at substantial disequilibrium for  $^{12}\text{CH}_2\text{D}_2$  (Jautzy et al., 2021). The majority of microbial  $\text{CH}_4$  from shallow freshwater environments is generated during acetoclastic methanogenesis, which can result in strong depletion for both  $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$  (figure 3 and 5)

(Conrad, 2005; Douglas et al., 2016, 2017, 2020; Haghnegahdar et al., 2024; Li et al., 2025a; Stolper et al., 2014b; Wang et al., 2015; Whiticar, 1999; Young et al., 2017). In systems with presumed slow CH<sub>4</sub> generation rates, favouring enzymatic isotopic reversibility, microbial CH<sub>4</sub> likely can form at or near thermodynamic isotopic equilibrium, while in systems with rapid CH<sub>4</sub> formations, microbial CH<sub>4</sub> tends to depart from thermodynamic isotopic equilibrium (Douglas et al., 2020; Shuai et al., 2021; Stolper et al., 2015; Wang et al., 2015).

Methane can also be produced abiotically, for example via Sabatier reactions linked to hydrogen production from serpentinization in hydrothermal systems (Cumming et al., 2019; Douglas et al., 2017; Labidi et al., 2020; Nothaft et al., 2021; Ojeda et al., 2023; Suda et al., 2022; Wang et al., 2018; Young et al., 2017). It has been observed from deep groundwater seeps accessed via or within deep subsurface layers, for instance in metal mines, where it can also mix with microbial CH<sub>4</sub> followed by re-equilibration (Nothaft et al., 2021; Warr et al., 2021a; Young et al., 2017). Typically, abiotic CH<sub>4</sub> is produced at temperatures exceeding 250 °C in seafloor hydrothermal fluids or in the continental seeps, springs and fracture waters at temperatures lower than 100 °C (Etiope and Sherwood Lollar, 2013; Labidi et al., 2024; Young et al., 2017). During controlled laboratory synthesis under hydrothermal conditions, the majority of the  $\Delta^{13}\text{CH}_3\text{D}$  measurements closely reflect the temperature of abiotic CH<sub>4</sub> generation (based on thermodynamic isotopic equilibrium).  $\Delta^{12}\text{CH}_2\text{D}_2$  was observed with depletions down to -40‰, which can be attributed to a D/H combinatorial effect associated with the various steps of hydrogen addition to carbon occurring during CH<sub>4</sub> formation (Labidi et al., 2024).

Using doubly substituted isotope ratio measurements, the mixed thermogenic-microbial origin of CH<sub>4</sub> was observed in marine environments, including CH<sub>4</sub> clathrates (Giunta et al., 2021; Zhang et al., 2021), lakes (Douglas et al., 2016), mud volcanoes (Lin et al., 2023; Liu et al., 2024a; Rumble et al., 2018), oil fields (Tyne et al., 2021) and natural gas (Douglas et al., 2017; Giunta et al., 2019; Kim et al., 2023; Lalk et al., 2022; Stolper et al., 2014b, 2015; Thiagarajan et al., 2020, 2022). Mixing between different CH<sub>4</sub> sources (containing different bulk isotopic compositions) in different proportions creates a non-linear relationship in  $\Delta^{12}\text{CH}_2\text{D}_2$  vs  $\Delta^{13}\text{CH}_3\text{D}$  space. Measurement of both doubly-substituted isotope ratios therefore provides additional information to help define the mixed end members and understand if physical or chemical transformation processes have taken place (e.g., Douglas et al., 2016; Young et al., 2016; Zhang et al., 2021).

Notably, existing studies showed a range of doubly-substituted isotope ratios for mud volcano samples, suggesting their different origins (thermogenic, microbial, abiotic or mixed) and potentially reflecting subsequent alteration processes such as AOM (Ijiri et al., 2018; Lalk et al., 2022; Lin et al., 2023; Liu et al., 2023, 2024a; Rumble et al., 2018). Additionally,  $\Delta^{13}\text{CH}_3\text{D}$  was used to demonstrate a microbial origin of CH<sub>4</sub> in deep subsurface coal beds in the north-western Pacific (Inagaki et al., 2015) and shallow subsurface mud volcano in the Nankai accretionary complex (Ijiri et al., 2018), which could otherwise be incorrectly identified as thermogenic sources. Also,  $\Delta^{12}\text{CH}_2\text{D}_2$  vs  $\Delta^{13}\text{CH}_3\text{D}$  suggested mixing of thermogenic and microbial CH<sub>4</sub> in coal bed reservoirs (Wang et al., 2024b, c).

Combinatorial effects occur when a molecule contains indistinguishable atoms of the same element derived from pools with different isotope ratios. This purely mathematical phenomenon comes from the definition of doubly-substituted isotope ratio in reference to the stochastic distribution and has been predicted theoretically (Röckmann et al., 2016; Yeung, 2016) and demonstrated experimentally for CH<sub>4</sub> (Labidi et al., 2024; Taenzer et al., 2020; Wang et al., 2024a). Among the two mass-18 isotopologues of CH<sub>4</sub>, only  $\Delta^{12}\text{CH}_2\text{D}_2$  can be influenced by combinatorial effects, as it features two indistinguishable deuterium substitutions for hydrogen. Combinatorial effects for  $\Delta^{12}\text{CH}_2\text{D}_2$  values must be taken into account in low-temperature abiotic or biotic systems where the hydrogen atoms of CH<sub>4</sub> originates from multiple reservoirs, which has been observed in microbial samples (Giunta et al., 2019; Jautzy et al., 2021; Young et al., 2017), mud volcanos (Liu et al., 2024a), natural gas (Shuai et al.,

2021; Xie et al., 2021), or during abiotic, microbial and pyrogenic methanogenesis experiments (Dong et al., 2021; Eldridge et al., 2023; Labidi et al., 2024; Li et al., 2025a). Notably, Eldridge et al., (2023) showed that combinatorial effects alone cannot explain the non-equilibrium of  $\Delta^{12}\text{CH}_2\text{D}_2$ , observed in their pyrogenic methanogenesis experiments focused on  $\text{CH}_4$  formation from methyl precursors (i.e. ethane). They pointed out the role of other important processes such as the influence of kinetic isotope effects and inheritance reactions (i.e., inheriting 'clumps' from methyl groups in the precursor molecule), in addition to combinatorial effects.

Before emission to the atmosphere,  $\text{CH}_4$  can be consumed through aerobic oxidation (AeOM) or anaerobic oxidation (AOM). In terrestrial ecosystems (e.g., wetlands) and oxygenated marine water columns, AeOM plays a crucial role, while in gas seeps and sulphate-rich marine sediments, AOM likely dominate causing inhibition of  $\text{CH}_4$  emissions to the atmosphere (e.g., Wang et al., 2016 and references therein). Minor depletions in  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  were observed in AeOM-dominated systems, but low-temperature equilibrium or significant enrichments in  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  were observed in the case of AOM (figures 3 and 7) (Giunta et al., 2022; Kim et al., 2023; Liu et al., 2023; Ono et al., 2021). One hypothesis states that the reversibility of initial steps of AOM promotes thermodynamic equilibration (Ash et al., 2019; Giunta et al., 2022; Ono et al., 2021; Zhang et al., 2021). Alternatively, another hypothesis proposes that near-thermodynamic equilibrium of doubly substituted isotope ratios in marine sediments can be attained via a slow rate of methanogenesis, with reversible enzymatic reaction steps (Douglas et al., 2020; Shuai et al., 2021; Stolper et al., 2015; Wang et al., 2015). As AeOM and AOM have distinctive kinetic isotope effects in natural settings, doubly-substituted isotope ratios may be used to track and differentiate both AeOM and AOM in nature (Adnew et al., 2025; Ash et al., 2019; Giunta et al., 2019, 2022; Krause et al., 2022; Li et al., 2024; Tyne et al., 2021; Warr et al., 2021b; Zhang et al., 2021).


In the troposphere, reaction with OH is the primary removal mechanism of  $\text{CH}_4$  (90%), with other minor contributions from microbial oxidation in soils and vegetation, loss to the stratosphere, and reactions with tropospheric Cl (e.g., Saunio et al., 2025). Overall, isotopologues containing bonds of lighter isotopes are preferentially removed through photochemical oxidation, leading to an enrichment in heavier isotopologues of the remaining  $\text{CH}_4$  pool (table S2) (e.g., Haghnegahdar et al., 2017; Whitehill et al., 2017). Laboratory experiments showed that photochemical oxidation by OH has only a minor impact on  $\Delta^{13}\text{CH}_3\text{D}$  of tropospheric  $\text{CH}_4$  (i.e. the  $^{13}\text{C}$ -D bond does not react significantly slower than that calculated based on equivalent singly substituted reactants) (Whitehill et al., 2017). Thus, measurements of  $\Delta^{13}\text{CH}_3\text{D}$  in the atmosphere can provide constraints on  $\text{CH}_4$  source strengths, while  $\Delta^{12}\text{CH}_2\text{D}_2$  is predicted to provide information on  $\text{CH}_4$  sink strength, as implemented in global scale atmospheric models (Chung and Arnold, 2021; Haghnegahdar et al., 2017; Whitehill et al., 2017). Aside from the atmospheric models, Wang et al. (2023b) used machine learning incorporated with a random forest model to predict steady-state atmospheric  $\text{CH}_4$  doubly substituted isotope ratios. The first measurements of the doubly substituted isotope ratio of  $\text{CH}_4$  in the atmosphere were more depleted for both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  than predicted by atmospheric models and available source signature information (Chung and Arnold, 2021; Haghnegahdar et al., 2017, 2023; Sivan et al., 2024). Haghnegahdar et al. (2023) proposed that differences between measurements and predictions required depleted doubly substituted isotopic signature values for the (total) source flux than previously assumed. On the other hand, Sivan et al. (2024) highlighted that the observed discrepancy could also be caused by inaccuracy in the theoretical values of the kinetic isotopic effect (KIE) of  $\text{CH}_4$  reactions with OH, Cl and soils sinks. They indicated that a small adjustment in the sink KIE, along with slightly lower source mixture than previously assumed, could align atmospheric and source doubly substituted isotopic signatures (Sivan et al., 2024).

### 4.3. Data representatives and importance for atmospheric sciences

The distribution of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  derived from field samples per simplified subcategory is plotted on figure 2b, while figures 6 and 7 present box plots for measured doubly substituted isotopes

from field samples and laboratory experiments, respectively. For simplicity, in figures 2b and 6, and thereafter, some subcategories are merged. Gas installation and natural gas subcategories are merged into natural gas. Hot spring, spring, and hyperalkaline spring are unified as spring. Marine sediment, marine seep, pore fluid and pockmark are grouped as a marine subcategory. Hydrothermal and volcano steam samples are unified as volcano. Finally, swamp and ponds are merged as wetlands, while lakes are in a separate subcategory. Around 40% of field samples were collected from reservoirs of natural gas (figure 2b). About 3% of field samples come from coal seam gas and 12.5% come from microbial terrestrial sources. There is a significant representation of marine (12.5% of field samples) and volcano mud samples (6% of field samples), although, their emissions to the atmosphere are negligible. For samples categorized as microbial terrestrial, the majority of entries come from lakes (75% of microbial terrestrial), with a small contribution from agriculture (6%) or wetland (12%) samples, which are significant  $\text{CH}_4$  emitters to the atmosphere. Only  $\Delta^{13}\text{CH}_3\text{D}$  was measured for four ruminants samples (Lopes et al., 2016; Wang et al., 2015). Only three samples from rice paddies have so far been collected, where both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  were measured (Haghnegahdar et al., 2023; Wang et al., 2023a). So far, no waste samples have been collected directly from the source for studies of doubly substituted isotope ratios. The recent studies of Sun et al. (2025a) focused on collection of big volume ambient air samples, where background air was mixed with gas coming from microbial  $\text{CH}_4$  sources, like wetlands and landfills. Application of a Keeling plot method (Pataki, 2003), allowed determination of targeted sources (Sun et al., 2025a).


Published  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  for natural gas are consistent with a thermogenic origin (figure 3 and 6, table S3 and S4). Observed outliers come from low maturity or oil-associated natural gas where a microbial contribution could be significant (Kim et al., 2023; Lalk et al., 2022; Thiagarajan et al., 2020; Xie et al., 2021). No significant variation has been observed in measurements made of biomass burning, dairy cows (ruminants), or rice paddies within the available, limited dataset but this may not reflect the variation within the true population (table S3 and S4). Significant variation in both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  is observed for spring and mud volcano subcategories, as these samples have varying microbial, thermogenic, abiotic, or mixed origins. Finally, a wide distribution is observed for lake samples, potentially originating from seasonal variation in  $\text{CH}_4$  production, oxidation in the lake subsurface or methanogenic metabolisms involved (Lalk et al., 2024).



**Figure 6.** Summary of the distribution of measurement results, a)  $\Delta^{13}\text{CH}_3\text{D}$  and b)  $\Delta^{12}\text{CH}_2\text{D}_2$  from field studies based on simplified subcategories as described in section 4.3.

For the laboratory experiments, culturing of different strains of archaea and wide variations in experimental parameters resulted in a wide distribution of observed doubly substituted isotopic compositions, especially for  $\Delta^{12}\text{CH}_2\text{D}_2$  (figure 7, table S5 and S6). AOM methanotrophy experiments show significant enrichment in both  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  relative to the other categories.





**Figure 7.** Summary of the distribution of measurement results, a)  $\Delta^{13}\text{CH}_3\text{D}$  and b)  $\Delta^{12}\text{CH}_2\text{D}_2$  from laboratory studies based on group types as described in section 4.1. The outliers for catalytic equilibration come from the sample measured at the beginning of the experiment, when equilibration on the catalyst did not start yet.

#### 4.3.1. Evaluation of the database in relation to emission to the atmosphere

On a global scale, using a bottom-up approach (e.g., using data-driven and process based models for natural sources and inventories for anthropogenic sources) for the year 2020, anthropogenic emissions contribute about 54% of the  $\text{CH}_4$  emissions to the atmosphere, originating from agriculture (40%), fossil fuel extraction and use (34%), waste (19%) and anthropogenic biomass burning (7%) (Saunois et al., 2025). Wetlands account for most of the natural  $\text{CH}_4$  emissions (51%), with a significant contribution from inland freshwaters (35%) and remaining emission coming from other sources, including onshore and offshore geological emissions (e.g., mud volcanoes, volcanoes, vents, seepages) (Saunois et al., 2025). Regarding the main  $\text{CH}_4$  emitters to the atmosphere, natural gas and oil are the most represented emission category in the doubly substituted  $\text{CH}_4$  isotopologue database (39% of field samples), while coal seams gas samples represent 4% of the field samples in the database. There are no reported measurements of  $\Delta^{12}\text{CH}_2\text{D}_2$  for ruminants (4 samples for  $\Delta^{13}\text{CH}_3\text{D}$  values), and no records of either  $\Delta^{13}\text{CH}_3\text{D}$  or  $\Delta^{12}\text{CH}_2\text{D}_2$  from directly sampled waste. Additionally, there is a very limited sample size for some important emissions subcategories such as biomass burning (0.2%) and rice paddies (0.3%). As field sampling is time consuming and location-constrained, measurements made this far do not reflect a realistic spatio-temporal variation of doubly substituted isotope ratios, both for anthropogenic and natural  $\text{CH}_4$  sources. With such limited studies, the current estimated  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  source signatures may not be representative. Thus, some assumptions on the source signature inputs to global scale models of double substituted isotope ratios have to be made (table 4). To better reflect  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  of  $\text{CH}_4$  emission sectors, further sampling should be focused on underrepresented  $\text{CH}_4$  sources and on numerous conditions affecting emissions from individual sectors, for example impact of reservoir depth and coal type for coal seam gas or impact of diet and living conditions for rumen (table 4). An effort should be made to measure  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  from thawing permafrost, as it may be a significant source of  $\text{CH}_4$  to the atmosphere in the future (Douglas et al., 2020; Ellenbogen et al., 2024; Walter Anthony et al., 2024).

**Table 4.** Global  $\text{CH}_4$  emissions and inferred doubly substituted  $\text{CH}_4$  isotope ratio signatures with remarks on the current representativeness of main  $\text{CH}_4$  sources to the atmosphere and requirements for future studies. Uncertainties of global emissions are reported as [min-max] range.

Group	Category	Global flux	$\Delta^{13}\text{CH}_3\text{D}$ [‰]			$\Delta^{12}\text{CH}_2\text{D}_2$ [‰]			Remarks		
			Average signature	Range	Samples number	Average signature	Range	Samples number	Representativeness	Existing assumptions	models

		[Tg CH <sub>4</sub> yr <sup>-1</sup> ] Bottom- up <sup>1)</sup>								
fossil fuels	coal seam gas	41 [38-43]	3.77	2.16; 5.87	35	10.20	4.25; 20.05	32	coal samples collected for sediment incubation experiments; no samples from mine ventilation; no information about impact of depth of coal seams or type of coal extraction	Whitehill et al. (2017): only $\Delta^{13}\text{CH}_3\text{D}$ , a common signature for lakes, landfill, all fossil fuels and biomass burning, estimated based on Wang et al. (2015);
	natural gas	74 [67-80] <sup>2)</sup>	3.36	0.30; 10.22	381	3.79	-23.13; 19.15	114	Emission from natural gas and oil merged in models and inventories; the best representation in the database; samples taken from sources with or without thermodynamic equilibrium; samples taken from different extraction regions; future sampling should be focused on underrepresented regions and various oil and gas infrastructure	Haghnegahdar et al (2017): assumed a common signature for all fossil fuels and biomass burning;
	oil field		3.98	2.90; 5.66	14	6.13	0.01; 12.46	13		Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al (2017), $\Delta^{13}\text{CH}_3\text{D}$ different to Haghnegahdar et al (2017) but common to all fossil categories; Haghnegahdar et al (2023): a common signature for all fossil fuels
microbial (except microbial fossil fuels)	dairy cow	117 [114-124] <sup>3)</sup>	1.32	0.76; 1.66	4	N/A	N/A	N/A	only $\Delta^{13}\text{CH}_3\text{D}$ measured; uncertain if dairy cow isotope ratio is representative for all ruminants and manure; critical demand of more sampling (type of rumen, impact of diet and living conditions, regional variation, different manure management systems), demand for $\Delta^{12}\text{CH}_2\text{D}_2$ measurements	Whitehill et al. (2017): only $\Delta^{13}\text{CH}_3\text{D}$ , a common signature for ruminants and rice paddies, estimated based on Wang et al. (2015); Haghnegahdar et al (2017): different signature using three different scenarios; Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al. (2017), $\Delta^{13}\text{CH}_3\text{D}$ based on cow rumen measurements. Haghnegahdar et al (2023): signatures based on interpretation of their wetland measurements.
	lake	112 [49-202] <sup>4)</sup>	3.35	-0.48; 9.60	91	-20.97	-37.76; 8.55	13	Samples taken mostly from lakes in the US with some	Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al

									contribution from European and Chinese lakes; only one study focused on seasonal variation, but no $\Delta^{12}\text{CH}_2\text{D}_2$ measurement (Lalk et al. 2024)	(2017), $\Delta^{13}\text{CH}_3\text{D}$ based on freshwater measurements
	rice paddies	32 [29-37]	-1.36	-1.80; -1.02	3	-36.04	-43.17; -31.11	3	Three field samples over two studies (two samples from China and one from the US), demand of increased spatial representation and samples from different rice paddies management systems (e.g., flooding, soil, rice variety)	Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al (2017), $\Delta^{13}\text{CH}_3\text{D}$ based on cow rumen measurements Haghnegahdar et al (2023): signatures based on interpretation of their wetland measurements
	waste <sup>6)</sup>	71 [60-84]	-1.3	N/A	N/A	-38.8	N/A	N/A	One of the main sources of $\text{CH}_4$ to the atmosphere; no representation of direct samples in the database; one study of mixed ambient air and landfill air (Sun et al. 20125), critical demand of samples from solid landfill, wastewater treatment and biogas, including sampling in different regions and seasons	Haghnegahdar et al. (2017): different signature using three different scenarios; Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al (2017), $\Delta^{13}\text{CH}_3\text{D}$ based on cow rumen measurements Haghnegahdar et al (2023): signatures based on interpretation of their wetland measurements
	wetlands	161 [131-198]	-0.49	-2.16; 1.08	14	-45.61	-57.16; -31.02	10	Samples taken only from wetlands in the US; demand for samples from different wetland regions, including tropical (significant $\text{CH}_4$ emitter) and polar wetlands and permafrost	Haghnegahdar et al (2017): category divided into boreal and tropical wetlands Chung and Arnold (2021): $\Delta^{12}\text{CH}_2\text{D}_2$ as in Haghnegahdar et al (2017), $\Delta^{13}\text{CH}_3\text{D}$ based on freshwater measurements
pyrogenic	biomass burning	27 [20-41] <sup>5)</sup>	2.16	1.63; 2.69	2	-16.31	-18.12; -14.49	2	Demand for samples from different type of biomass and biofuel; need for examination of the impact of burning conditions on isotope ratios (few laboratory	Haghnegahdar et al (2017): assumed thermodynamic equilibrium, common signature for all fossil fuels and biomass burning;

									experiments conducted)	
--	--	--	--	--	--	--	--	--	---------------------------	--

- 600
- 1) CH<sub>4</sub> global flux from Saunois et al. 2025 for the year 2020

2) CH<sub>4</sub> global flux for natural gas and oil merged into one category in Saunois et al. 2025

3) enteric fermentation & manure category in Saunois et al. 2025

4) inland freshwater category in Saunois et al. 2025

5) biomass and biofuel burning together from Saunois et al. 2025
- 605
- 6) Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> of waste sector from indirect measurement (e.g., ambient air mixed with gas from landfill) from Sun et al. 2025

610


In addition to increasing the sampling frequency for the main CH<sub>4</sub> sources, an effort should also be made to extend sampling to other areas with significant CH<sub>4</sub> emissions to the atmosphere, including super-emitters. Using TROPOMI (TROPOspheric Monitoring Instrument) satellite data, super-emitters were detected for coal mining, oil and gas production regions, and along the major gas transmission pipelines (Schuit et al., 2023). The majority of detected super-emitters is related to urban areas (35% of detected super-emitters), with a possible large contribution from landfills (Schuit et al., 2023), where no direct samples of Δ<sup>13</sup>CH<sub>3</sub>D and Δ<sup>12</sup>CH<sub>2</sub>D<sub>2</sub> have been taken so far.

615

Comparing locations of field samples and a map of anthropogenic CH<sub>4</sub> emissions, based on EDGAR v8.0 inventories, there is a considerable deficiency in measurements of doubly substituted isotope ratios from numerous locations with elevated CH<sub>4</sub> emissions (figure 8). No samples have been analysed from regions with significant CH<sub>4</sub> emissions, like Central Africa, southwestern South America, India, Pakistan, western China, New Zealand, and Indonesia. There is no data from the EDGAR database for certain areas, such as Siberia and Canada, where increased anthropogenic emissions can occur as well.

620

Furthermore, sampling should be conducted in regions with notable natural emissions, such as wetlands and internal freshwaters, including thawing permafrost.



**Figure 8.** Global locations of collected field samples for doubly substituted isotope measurement (blue and green circles) overlaid on an estimate of the total CH<sub>4</sub> annual emission rates for year 2022 from the EDGAR v8.0 inventory.

625

### 5. Data availability

Data may be accessed from the following DOI:  
 Defratyka, S. M., Fernandez, J. M., and Arnold, T.: Methane doubly substituted (clumped) isotopologues database, CEDA Arch., <https://dx.doi.org/10.5285/51ae627da5fb41b8a767ee6c653f83e6>, 2025.

## 6. Conclusions

This study presents a compilation of  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  measurements from field samples and laboratory experiments, from results published between 2014 and 2025, by numerous laboratories. The database is designed for utilization by the geochemistry and atmospheric science communities. The database of doubly substituted isotope ratios comprises 1475 data records from 75 peer-reviewed articles (figure 2a and 4). Of this data, 53% of the database entries report only  $\Delta_{18}$  or  $\Delta^{13}\text{CH}_3\text{D}$ , which can hinder data interpretation, especially for microbial, abiotic or mixed samples, when used without any additional tracer (Chowdhury et al., 2024; Douglas et al., 2017; Giunta et al., 2019; Gruen et al., 2018; Thiagarajan et al., 2020; Warr et al., 2021a; Young et al., 2016, 2017). For field samples, 40% of the data records come from natural gas, mostly from the basins in the US and China. Samples collected from lakes contribute 75% of microbial terrestrial samples. At the current state, there is a limited representation of samples coming from wetlands and agriculture sources and there is no representation of directly sampled waste sector (figure 2b).

As our ability to measure doubly substituted isotopologues of  $\text{CH}_4$  in the atmosphere improves, a commensurate effort to improve our understanding of source signatures is needed in order to make the very most of these measurements in understanding the global atmospheric  $\text{CH}_4$  budget. Studies should focus on the main emission sectors to the atmosphere, in particular on underrepresented sectors such as agriculture (e.g., ruminants, manure, rice cultivation), wetlands (including polar), waste and biomass burning. Also new field campaigns should focus on areas with increased  $\text{CH}_4$  emissions, including super-emitters. An additional effort is also required to provide more ambient air background samples, ideally from remote, clean air sites. To better understand  $\text{CH}_4$  sinks, more experiments focused on photochemical oxidation by OH and Cl must also be conducted.

**Supplement link** (link for excel spreadsheet with Tables S1-S6, given by editor of ESSD)

### Author contributions

**Conceptualization:** SMD, JMF, TA; **Investigation and data curation:** GAA, GD, PMJD, DLE, GE, TG, MAH, ANH, NH, VI, JJ, JHK, JL, EL, WL, JL, LHL, JL, LO, SO, JR, TR, BSL, MS, JS, GTV, DTW, EDY, NZ; **Formal analysis:** SMD; **Visualization:** SMD; **Writing (original draft preparation):** SMD; **Writing (review and editing):** SMD, JMF, TA, GAA, GD, PMJD, DLE, GE, TG, MAH, ANH, NH, VI, JJ, JHK, JL, EL, WL, JL, LHL, JL, LO, SO, JR, TR, BSL, MS, JS, GTV, DTW, EDY, NZ.

### Competing interests

The authors declare that they have no conflict of interest.

### Acknowledgements

For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission. We gratefully acknowledge the authors and researchers whose previously published work was used for this data aggregation. The data compiled herein are derived from and built upon findings reported in peer-reviewed scientific literature. All original sources have been cited appropriately in the accompanying references.

Funding for this work came from the UKRI NERC POLYGRAM project NE/V007149/1 ([www.polygram.ac.uk](http://www.polygram.ac.uk)), the EURAMET 21GRD04 isoMET project and the NPL Director's Fund. The 21GRD04 isoMET project has received funding from the European Partnership on Metrology, co-financed from the European Union's Horizon Europe Research and Innovation Programme and by the Participating States. This article is not a product of the U.S. Department of Energy. Views and opinions expressed in this article are the authors' own and do not represent those of the United States Government.

## References

- 675 Adnew, G. A., Rockmann, T., Blunier, T., Jørgensen, C. J., Sapper, S. E., Veen, C. van der, Sivan, M., Popa, M. E., and Christiansen, J. R.: Clumped isotope measurements reveal aerobic oxidation of methane below the Greenland ice sheet, *Geochim. Cosmochim. Acta*, 389, 249–264, <https://doi.org/doi.org/10.1016/j.gca.2024.11.009>, 2025.
- Ash, J. L., Egger, M., Treude, T., Kohl, I., Cragg, B., Parkes, R. J., Slomp, C. P., Sherwood Lollar, B., and Young, E. D.: Exchange catalysis during anaerobic methanotrophy revealed by  $^{12}\text{CH}_2\text{D}_2$  and  $^{13}\text{CH}_3\text{D}$  in methane, *Geochem. Perspect. Lett.*, 26–30, <https://doi.org/10.7185/geochemlet.1910>, 2019.
- 680 Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of methane and  $\delta^{13}\text{C}$  of methane, *Atmospheric Chem. Phys.*, 22, 15351–15377, <https://doi.org/10.5194/acp-22-15351-2022>, 2022.
- 685 Beaudry, P., Stefansson, A., Fiebig, J., Rhim, J. H., and Ono, S.: High temperature generation and equilibration of methane in terrestrial geothermal systems: Evidence from clumped isotopologues, *Geochim. Cosmochim. Acta*, 309, 209–234, <https://doi.org/10.1016/j.gca.2021.06.034>, 2021.
- Chen, C., Qin, S., Wang, Y., Holland, G., Wynn, P., Zhong, W., and Zhou, Z.: High temperature methane emissions from Large Igneous Provinces as contributors to late Permian mass extinctions, *Nat. Commun.*, 13, <https://doi.org/10.1038/s41467-022-34645-3>, 2022.
- 690 Chowdhury, A., Ventura, G. T., Owino, Y., Lalk, E. J., MacAdam, N., Dooma, J. M., Ono, S., Fowler, M., MacDonald, A., Bennett, R., MacRae, R. A., Hubert, C. R. J., Bentley, J. N., and Kerr, M. J.: Cold seep formation from salt diapir-controlled deep biosphere oases, *PNAS*, 121, <https://doi.org/10.1073/pnas.2316878121>, 2024.
- Chung, E. and Arnold, T.: Potential of Clumped Isotopes in Constraining the Global Atmospheric Methane Budget, *Glob. Biogeochem. Cycles*, 35, <https://doi.org/10.1029/2020GB006883>, 2021.
- 695 Conrad, R.: Quantification of methanogenic pathways using stable carbon isotopic signatures: a review and a proposal, *Org. Geochem.*, 36, 739–752, 2005.
- Cumming, E. A., Rietze, A., Morrissey, L. S., Cook, M. C., Rhim, J. H., Ono, S., and Morrill, P. L.: Potential sources of dissolved methane at the Tablelands, Gros Morne National Park, NL, CAN: A terrestrial site of serpentinization, *Chem. Geol.*, 514, 42–53, <https://doi.org/10.1016/j.chemgeo.2019.03.019>, 2019.
- 700 Defratyka, S., Fernandez, J., and Arnold, T.: Methane doubly substituted (clumped) isotopologues database, NERC EDS Cent. Environ. Data Anal., <https://dx.doi.org/10.5285/51ae627da5fb41b8a767ee6c653f83e6>, 2025.
- Dong, G., Xie, H., Thiagarajan, N., Eiler, J., Zhang, N., Nakagawa, M., Yoshida, N., Eldridge, D., Stolper, D., Albrecht, N., and Kohl, I. E.: Clumped isotope analysis of methane using HR-IRMS: New insights into origin and formation mechanisms of natural gases and a potential geothermometer, 2020.
- 705 Dong, G., Xie, H., Formolo, M., Lawson, M., Sessions, A., and Eiler, J.: Clumped isotope effects of thermogenic methane formation: Insights from pyrolysis of hydrocarbons, *Geochim. Cosmochim. Acta*, 303, 159–183, <https://doi.org/10.1016/j.gca.2021.03.009>, 2021.

- 710 Douglas, P. M. J., Stolper, D. A., Smith, D. A., Walter Anthony, K. M., Paull, C. K., Dallimore, S., Wik, M., Crill, P. M., Winterdahl, M., Eiler, J. M., and Sessions, A. L.: Diverse origins of Arctic and Subarctic methane point source emissions identified with multiply-substituted isotopologues, *Geochim. Cosmochim. Acta*, 188, 163–188, <https://doi.org/10.1016/j.gca.2016.05.031>, 2016.
- 715 Douglas, P. M. J., Stolper, D. A., Eiler, J. M., Sessions, A. L., Lawson, M., Shuai, Y., Bishop, A., Podlaha, O. G., Ferreira, A. A., Santos Neto, E. V., Niemann, M., Steen, A. S., Huang, L., Chimiak, L., Valentine, D. L., Fiebig, J., Luhmann, A. J., Seyfried, W. E., Etiope, G., Schoell, M., Inskeep, W. P., Moran, J. J., and Kitchen, N.: Methane clumped isotopes: Progress and potential for a new isotopic tracer, *Org. Geochem.*, 113, 262–282, <https://doi.org/10.1016/j.orggeochem.2017.07.016>, 2017.
- 720 Douglas, P. M. J., Moguel, R. G., Anthony, K. M. W., Wik, M., Crill, P. M., Dawson, K. S., Smith, D. A., Yanay, E., Lloyd, M. K., Stolper, D. A., Eiler, J. M., and Sessions, A. L.: Clumped Isotopes Link Older Carbon Substrates With Slower Rates of Methanogenesis in Northern Lakes, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL086756>, 2020.
- Eiler, J. M.: “Clumped-isotope” geochemistry—The study of naturally-occurring, multiply-substituted isotopologues, *Earth Planet. Sci. Lett.*, 262, 309–327, <https://doi.org/10.1016/j.epsl.2007.08.020>, 2007.
- 725 Eiler, J. M.: The Isotopic Anatomies of Molecules and Minerals, *Annu. Rev. Earth Planet. Sci.*, 41, 411–441, <https://doi.org/10.1146/annurev-earth-042711-105348>, 2013.
- Eldridge, D. L., Korol, R., Lloyd, M. K., Turner, A. C., Webb, M. A., Miller, T. F., and Stolper, D. A.: Comparison of Experimental vs Theoretical Abundances of  $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$  for Isotopically Equilibrated Systems from 1 to 500 °C, *ACS Earth Space Chem.*, 3, 2747–2764, <https://doi.org/10.1021/acsearthspacechem.9b00244>, 2019.
- 730 Eldridge, D. L., Turner, A. C., Bill, M., Conrad, M. E., and Stolper, D. A.: Experimental determinations of carbon and hydrogen isotope fractionations and methane clumped isotope compositions associated with ethane pyrolysis from 550 to 600 °C, *Geochim. Cosmochim. Acta*, 355, 235–265, <https://doi.org/10.1016/j.gca.2023.06.006>, 2023.
- 735 Ellenbogen, J. B., Borton, M. A., McGivern, B. B., Cronin, D. R., Hoyt, D. W., Freire-Zapata, V., McCalley, C. K., Varner, R. K., Crill, P. M., Wehr, R. A., Chanton, J. P., Woodcroft, B. J., Tfaily, M. M., Tyson, G. W., Rich, V. I., and Wrighton, K. C.: Methylophily in the Mire: direct and indirect routes for methane production in thawing permafrost, *mSystems*, 9, <https://doi.org/doi.org/10.1128/msystems.00698-23>, 2024.
- Etiope, G. and Sherwood Lollar, B.: ABIOTIC METHANE ON EARTH, *Rev. Geophys.*, 51, 276–299, <https://doi.org/10.1002/rog.20011>, 2013.
- 740 Giunta, T., Young, E. D., Warr, O., Kohl, I., Ash, J. L., Martini, A., Mundle, S. O. C., Rumble, D., Pérez-Rodríguez, I., Wasley, M., LaRowe, D. E., Gilbert, A., and Sherwood Lollar, B.: Methane sources and sinks in continental sedimentary systems: New insights from paired clumped isotopologues  $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$ , *Geochim. Cosmochim. Acta*, 245, 327–351, <https://doi.org/10.1016/j.gca.2018.10.030>, 2019.
- 745 Giunta, T., Labidi, J., Kohl, I. E., Ruffine, L., Donval, J. P., Géli, L., Çağatay, M. N., Lu, H., and Young, E. D.: Evidence for methane isotopic bond re-ordering in gas reservoirs sourcing cold seeps from the Sea of Marmara, *Earth Planet. Sci. Lett.*, 553, 116619, <https://doi.org/10.1016/j.epsl.2020.116619>, 2021.
- Giunta, T., Young, E. D., Labidi, J., Sansjofre, P., Jézéquel, D., Donval, J.-P., Brandily, C., and Ruffine, L.: Extreme methane clumped isotopologue bio-signatures of aerobic and anaerobic methanotrophy: Insights from the Lake Pavin and the Black Sea sediments, *Geochim. Cosmochim. Acta*, 338, 34–53, <https://doi.org/10.1016/j.gca.2022.09.034>, 2022.
- 750 Gonzalez, Y., Nelson, D. D., Shorter, J. H., McManus, J. B., Dyroff, C., Formolo, M., Wang, D. T., Western, C. M., and Ono, S.: Precise Measurements of  $^{12}\text{CH}_2\text{D}_2$  by Tunable Infrared Laser Direct Absorption Spectroscopy, *Anal. Chem.*, 91, 14967–14974, <https://doi.org/10.1021/acs.analchem.9b03412>, 2019.

- Gruen, D. S., Wang, D. T., Könneke, M., Topçuoğlu, B. D., Stewart, L. C., Goldhammer, T., Holden, J. F., Hinrichs, K.-U., and Ono, S.: Experimental investigation on the controls of clumped isotopologue and hydrogen isotope ratios in microbial methane, *Geochim. Cosmochim. Acta*, 237, 339–356, <https://doi.org/10.1016/j.gca.2018.06.029>, 2018.
- Haghnegahdar, M. A., Schauble, E. A., and Young, E. D.: A model for  $^{12}\text{CH}_2\text{D}_2$  and  $^{13}\text{CH}_3\text{D}$  as complementary tracers for the budget of atmospheric  $\text{CH}_4$ , *Glob. Biogeochem. Cycles*, 31, 1387–1407, <https://doi.org/10.1002/2017GB005655>, 2017.
- Haghnegahdar, M. A., Sun, J., Hultquist, N., Hamovit, N. D., Kitchen, N., Eiler, J., Ono, S., Yarwood, S. A., Kaufman, A. J., Dickerson, R. R., Bouyon, A., Magen, C., and Farquhar, J.: Tracing sources of atmospheric methane using clumped isotopes, *PNAS*, 120, <https://doi.org/10.1073/pnas.2305574120>, 2023.
- Haghnegahdar, M. A., Hultquist, N., Hamovit, N. D., Yarwood, S. A., Bouyon, A., Kaufman, A. J., Sun, J., Magen, C., and Farquhar, J.: A Better Understanding of Atmospheric Methane Sources Using  $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$  Clumped Isotopes, *J. Geophys. Res.*, 129, <https://doi.org/doi.org/10.1029/2024JG00817>, 2024.
- Ijiri, A., Inagaki, F., Kubo, Y., Adhikari, R. R., Hattori, S., Hoshino, T., Imachi, H., Kawagucci, S., Morono, Y., Ohtomo, Y., Ono, S., Sakai, S., Takai, K., Toki, T., Wang, D. T., Yoshinaga, M. Y., Arnold, G. L., Ashi, J., Case, D. H., Feseker, T., Hinrichs, K.-U., Ikegawa, Y., Ikehara, M., Kallmeyer, J., Kumagai, H., Lever, M. A., Morita, S., Nakamura, K., Nakamura, Y., Nishizawa, M., Orphan, V. J., Røy, H., Schmidt, F., Tani, A., Tanikawa, W., Terada, T., Tomaru, H., Tsuji, T., Tsunogai, U., Yamaguchi, Y. T., and Yoshida, N.: Deep-biosphere methane production stimulated by geofluids in the Nankai accretionary complex, *Sci. Adv.*, 4, eaao4631, <https://doi.org/10.1126/sciadv.aao4631>, 2018.
- Inagaki, F., Hinrichs, K.-U., Kubo, Y., Bowles, M. W., Heuer, V. B., Hong, W.-L., Hoshino, T., Ijiri, A., Imachi, H., Ito, M., Kaneko, M., Lever, M. A., Lin, Y.-S., Methé, B. A., Morita, S., Morono, Y., Tanikawa, W., Bihan, M., Bowden, S. A., Elvert, M., Glombitza, C., Gross, D., Harrington, G. J., Hori, T., Li, K., Limmer, D., Liu, C.-H., Murayama, M., Ohkouchi, N., Ono, S., Park, Y.-S., Phillips, S. C., Prieto-Mollar, X., Purkey, M., Riedinger, N., Sanada, Y., Sauvage, J., Snyder, G., Susilawati, R., Takano, Y., Tasumi, E., Terada, T., Tomaru, H., Trembath-Reichert, E., Wang, D. T., and Yamada, Y.: Exploring deep microbial life in coal-bearing sediment down to ~2.5 km below the ocean floor, *Science*, 349, 420–424, <https://doi.org/10.1126/science.aaa6882>, 2015.
- Jautzy, J. J., M.J.Douglas, P., Xie, H., M.Eiler, J., and D.Clark, I.:  $\text{CH}_4$  isotopic ordering records ultra-slow hydrocarbon biodegradation in the deep subsurface, *Earth Planet. Sci. Lett.*, 562, <https://doi.org/10.1016/j.epsl.2021.116841>, 2021.
- Jiang, W., Qingmei Liu, Jiacheng Li, Yun Li, Wen Liu, Xian Liu, Haizu Zhang, Pingan Peng, and Yongqiang Xiong: Deciphering the origin and secondary alteration of deep natural gas in the Tarim basin through paired methane clumped isotopes, *Mar. Pet. Geol.*, 160, <https://doi.org/10.1016/j.marpetgeo.2023.106614>, 2024.
- Kim, J.-H., Anna M. Martini, Shuhei Ono, Ellen Lalk, Grant Ferguson, and Jennifer C. McIntosh: Clumped and conventional isotopes of natural gas reveal basin burial, denudation, and biodegradation history, *Geochim. Cosmochim. Acta*, 361, 133–151, <https://doi.org/10.1016/j.gca.2023.10.017>, 2023.
- Krause, S. J. E., Liu, J., D.Young, E., and Treude, T.:  $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$  signatures of methane aerobically oxidized by *Methylosinus trichosporium* with implications for deciphering the provenance of methane gases, *Earth Planet. Sci. Lett.*, 593, <https://doi.org/10.1016/j.epsl.2022.117681>, 2022.
- Labidi, J., Young, E. D., Giunta, T., Kohl, I. E., Seewald, J., Tang, H., Lilley, M. D., and Früh-Green, G. L.: Methane thermometry in deep-sea hydrothermal systems: Evidence for re-ordering of doubly-substituted isotopologues during fluid cooling, *Geochim. Cosmochim. Acta*, 288, 248–261, <https://doi.org/10.1016/j.gca.2020.08.013>, 2020.
- Labidi, J., McCollom, T. M., Giunta, T., Sherwood Lollar, B., Leavitt, W. D., and Young, E. D.: Clumped Isotope Signatures of Abiotic Methane: The Role of the Combinatorial Isotope Effect, *J. Geophys. Res.*, 129, <https://doi.org/10.1029/2023JB028194>, 2024.



- Lalk, E., Thomas Pape, Danielle S. Gruen, Norbert Kaul, Jennifer S. Karolewski, Gerhard Bohrmann, and Shuhei Ono: Clumped methane isotopologue-based temperature estimates for sources of methane in marine gas hydrates and associated vent gases, *Geochim. Cosmochim. Acta*, 327, 276–297, <https://doi.org/10.1016/j.gca.2022.04.013>, 2022.
- Lalk, E., Seewald, J. S., Bryndzia, L. T., and Ono, S.: Kilometer-scale  $\Delta^{13}\text{CH}_3\text{D}$  profiles distinguish end-member mixing from methane production in deep marine sediments, *Org. Geochem.*, 181, <https://doi.org/10.1016/j.orggeochem.2023.104630>, 2023.
- Lalk, E., Velez, A., and Ono, S.: Methane Clumped Isotopologue Variability from Ebullition in a Mid-latitude Lake, *ACS Earth Space Chem.*, <https://doi.org/10.1021/acsearthspacechem.3c00282>, 2024.
- Lan, X., Basu, S., Schwietzke, S., Bruhwiler, L. M. P., Dlugokencky, E. J., Michel, S. E., Sherwood, O. A., Tans, P. P., Thoning, K., Etiope, G., Zhuang, Q., Liu, L., Oh, Y., Miller, J. B., Pétron, G., Vaughn, B. H., and Crippa, M.: Improved Constraints on Global Methane Emissions and Sinks Using  $\delta^{13}\text{C}-\text{CH}_4$ , *Glob. Biogeochem. Cycles*, 35, <https://doi.org/10.1029/2021GB007000>, 2021.
- Li, J., Chiu, B. K., Piasecki, A. M., Feng, X., Landis, J. D., Marcum, S., Young, E. D., and Leavitt, W. D.: The evolution of multiply substituted isotopologues of methane during microbial aerobic oxidation, *Geochim. Cosmochim. Acta*, 381, 223–238, <https://doi.org/10.1016/j.gca.2024.06.032>, 2024.
- Li, J., Ash, J. L., Cobban, A., Kubik, B. C., Rizzo, G., Thompson, M., Guibourdenche, L., Berger, S., Morra, K., Lin, Y., Mueller, E. P., Masterson, A. L., Stein, R., Fogel, M., Torres, M. A., Feng, X., Holden, J. F., Martini, A., Welte, C. U., Jetten, M., Young, E. D., and Leavitt, W. D.: The clumped isotope signatures of multiple methanogenesis metabolisms, *Env. Sci Technol*, 59, 13798–13810, <https://doi.org/10.1021/acs.est.5c03255>, 2025a.
- Li, J., Liu, Q., Jiang, W., Li, Y., Shuai, Y., and Xiong, Y.: Tracing the Contribution of Abiotic Methane in Deep Natural Gases From the Songliao Basin, China Using Bulk Isotopes and Methane Clumped Isotopologue  $^{12}\text{CH}_2\text{D}_2$ , *Geochem. Geophys. Geosystems*, 26, e2024GC011705, <https://doi.org/10.1029/2024GC011705>, 2025b.
- Lin, Y.-T., Rumble, D., Young, E. D., Labidi, J., Tu, T.-H., Chen, J.-N., Pape, T., Bohrmann, G., Lin, S., Lin, L.-H., and Wang, P.-L.: Diverse Origins of Gases From Mud Volcanoes and Seeps in Tectonically Fragmented Terrane, *Geochem. Geophys. Geosystems*, 24, <https://doi.org/10.1029/2022GC010791>, 2023.
- Liu, J., Harris, R. L., Ash, J. L., Ferry, J. G., Krause, S. J. E., Labidi, J., Prakash, D., Sherwood Lollar, B., Treude, T., Warr, O., and Young, E. D.: Reversibility controls on extreme methane clumped isotope signatures from anaerobic oxidation of methane, *Geochim. Cosmochim. Acta*, 348, 165–186, <https://doi.org/10.1016/j.gca.2023.02.022>, 2023.
- Liu, J., Treude, T., Abbasov, O. R., Baloglanov, E. E., Aliyev, A. A., Harris, C. M., Leavitt, W. D., and Young, E. D.: Clumped isotope evidence for microbial alteration of thermogenic methane in terrestrial mud volcanoes, *Geology*, 52, 22–26, <https://doi.org/10.1130/G51667.1>, 2024a.
- Liu, Q. and Liu, Y.: Clumped-isotope signatures at equilibrium of  $\text{CH}_4$ ,  $\text{NH}_3$ ,  $\text{H}_2\text{O}$ ,  $\text{H}_2\text{S}$  and  $\text{SO}_2$ , *Geochim. Cosmochim. Acta*, 175, 252–270, <https://doi.org/10.1016/j.gca.2015.11.040>, 2016.
- Liu, Q., Li, J., Jiang, W., Li, Y., Lin, M., Liu, W., Shuai, Y., Zhang, H., Peng, P., and Xiong, Y.: Application of an absolute reference frame for methane clumped-isotope analyses, *Chem. Geol.*, 646, <https://doi.org/10.1016/j.chemgeo.2024.121922>, 2024b.
- Lopes, J. C., Matos, L. F. de, Harper, M. T., Giallongo, F., Oh, J., Gruen, D., Ono, S., Kindermann, M., Duval, S., and Hristov, A. N.: Effect of 3-nitrooxypropanol on methane and hydrogen emissions, methane isotopic signature, and ruminal fermentation in dairy cows, *J Dairy Sci*, 99, 5335–5344, <http://dx.doi.org/10.3168/jds.2015-10832>, 2016.
- Ma, Q., Wu, S., and Tang, Y.: Formation and abundance of doubly-substituted methane isotopologues ( $^{13}\text{CH}_3\text{D}$ ) in natural gas systems, *Geochim. Cosmochim. Acta*, 72, 5446–5456, <https://doi.org/10.1016/j.gca.2008.08.014>, 2008.

- Menoud, M., Veen, C. van der, Fernandez, J., Semra, B., Lowry, D., France, J., Fisher, R., Maazallahi, H., Korbeń, P., Schmidt, M., Stanisavljević, M., Necki, J., Łakomiec, P., Rinne, J., Defratyka, S., Yver-Kwok, C., Vinkovic, K., Andersen, T., Chen, H., and Röckmann, T.: European Methane Isotope Database Coupled with a Global Inventory of Fossil and Non-Fossil  $\delta^{13}\text{C}$ - and  $\delta^2\text{H}$ -CH<sub>4</sub> Source Signature Measurements: V2.0.0, Utrecht Univ. Data Set, <https://doi.org/10.24416/UU01-YP43IN>, 2022a.
- Menoud, M., van der Veen, C., Lowry, D., Fernandez, J. M., Bakkaloglu, S., France, J. L., Fisher, R. E., Maazallahi, H., Stanisavljević, M., Nećki, J., Vinkovic, K., Łakomiec, P., Rinne, J., Korbeń, P., Schmidt, M., Defratyka, S., Yver-Kwok, C., Andersen, T., Chen, H., and Röckmann, T.: New contributions of measurements in Europe to the global inventory of the stable isotopic composition of methane, *Earth Syst. Sci. Data*, 14, 4365–4386, <https://doi.org/10.5194/essd-14-4365-2022>, 2022b.
- Mroz, E. J., Alei, M., Capps, J. H., Guthals, P. R., Mason, A. S., and Rokop, D. J.: Detection of multiply deuterated methane in the atmosphere, *Geophys. Res. Lett.*, 16, 677–678, <https://doi.org/10.1029/GL016i007p00677>, 1989.
- Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Fisher, R. E., Lowry, D., Michel, S. E., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., Brownlow, R., Cain, M., France, J. L., Hermansen, O., Hossaini, R., Jones, A. E., Levin, I., Manning, A. C., Myhre, G., Pyle, J. A., Vaughn, B. H., Warwick, N. J., and White, J. W. C.: Very Strong Atmospheric Methane Growth in the 4 Years 2014–2017: Implications for the Paris Agreement, *Glob. Biogeochem. Cycles*, 33, 318–342, <https://doi.org/10.1029/2018GB006009>, 2019.
- Nothaft, D. B., Templeton, A. S., Rhim, J. H., Wang, D. T., Labidi, J., Miller, H. M., Boyd, E. S., Matter, J. M., Ono, S., Young, E. D., Kopf, S. H., Kelemen, P. B., and Conrad, M. E.: Geochemical, Biological, and Clumped Isotopologue Evidence for Substantial Microbial Methane Production Under Carbon Limitation in Serpentinites of the Samail Ophiolite, Oman, *J. Geophys. Res.*, 126, <https://doi.org/10.1029/2020JG006025>, 2021.
- Ojeda, L., Etiope, G., Jimenez-Gavilan, P., Martonos, I. M., Rockmann, T., Popa, M. E., Sivan, M., Castro-Gamez, A. F., Benavente, J., and Vadillo, I.: Combining methane clumped and bulk isotopes, temporal variations in molecular and isotopic composition, and hydrochemical and geological proxies to understand methane's origin in the Ronda peridotite massifs (Spain), *Chem. Geol.*, 642, <https://doi.org/10.1016/j.chemgeo.2023.121799>, 2023.
- Ono, S., Wang, D. T., Gruen, D. S., Sherwood Lollar, B., Zahniser, M. S., McManus, B. J., and Nelson, D. D.: Measurement of a Doubly Substituted Methane Isotopologue,  $^{13}\text{CH}_3\text{D}$ , by Tunable Infrared Laser Direct Absorption Spectroscopy, *Anal. Chem.*, 86, 6487–6494, <https://doi.org/10.1021/acs.chem.2c01949>, 2022.
- Ono, S., Jeemin H. Rhim, Danielle S. Gruen, Heidi Taubner, Martin Kolling, and Gunter Wegener: Clumped isotopologue fractionation by microbial cultures performing the anaerobic oxidation of methane, *Geochim. Cosmochim. Acta*, 293, 70–85, <https://doi.org/10.1016/j.gca.2020.10.015>, 2021.
- Pataki, D. E.: Seasonal cycle of carbon dioxide and its isotopic composition in an urban atmosphere: Anthropogenic and biogenic effects, *J. Geophys. Res.*, 108, 4735, <https://doi.org/10.1029/2003JD003865>, 2003.
- Prokhorov, I. and Mohn, J.: CleanEx: A Versatile Automated Methane Preconcentration Device for High-Precision Analysis of  $^{13}\text{CH}_4$ ,  $^{12}\text{CH}_3\text{D}$ , and  $^{13}\text{CH}_3\text{D}$ , *Anal. Chem.*, 94, 9981–9986, <https://doi.org/10.1021/acs.analchem.2c01949>, 2022.
- Rhim, J. H. and Ono, S.: Combined carbon, hydrogen, and clumped isotope fractionations reveal differential reversibility of hydrogenotrophic methanogenesis in laboratory cultures, *Geochim. Cosmochim. Acta*, 335, 383–399, <https://doi.org/10.1016/j.gca.2022.07.027>, 2022.
- Röckmann, T., Popa, M. E., Krol, M. C., and Hofmann, M. E. G.: Statistical clumped isotope signatures, *Sci. Rep.*, 6, <https://doi.org/10.1038/srep31947>, 2016.
- Rumble, D., Ash, J. L., Wang, P.-L., Lin, L.-H., Lin, Y.-T., and Tu, T.-H.: Resolved measurements of  $^{13}\text{CDH}_3$  and  $^{12}\text{CD}_2\text{H}_2$  from a mud volcano in Taiwan, *J. Asian Earth Sci.*, 167, 218–221, <https://doi.org/10.1016/j.jseaes.2018.03.007>, 2018.

- Safi, E., Arnold, T., and Rennick, C.: Fractionation of Methane Isotopologues during Preparation for Analysis from Ambient Air, *Anal. Chem.*, 96, 6139–6147, <https://doi.org/10.1021/acs.analchem.3c04891>, 2024.
- 890 Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguía-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, *Earth Syst. Sci. Data*, 17, 1873–1958, <https://doi.org/10.5194/essd-17-1873-2025>, 2025.
- 895 Schoell, M.: MULTIPLE ORIGINS OF METHANE IN THE EARTH, *Chem. Geol.*, 71, 1–10, [https://doi.org/doi.org/10.1016/0009-2541\(88\)90101-5](https://doi.org/doi.org/10.1016/0009-2541(88)90101-5), 1988.
- 900 Schuit, B. J., Maasakkers, J. D., Bijl, P., and Mahapatra, G.: Automated detection and monitoring of methane super-emitters using satellite data, *Atmos Chem Phys*, 23, 9071–9098, <https://doi.org/10.5194/acp-23-9071-2023>, 2023.
- Sherwood, O. A., Schwietzke, S., Arling, V. A., and Etiope, G.: Global Inventory of Gas Geochemistry Data from Fossil Fuel, Microbial and Burning Sources, version 2017, *Earth Syst. Sci. Data*, 9, 639–656, <https://doi.org/10.5194/essd-9-639-2017>, 2017.
- 905 Sherwood, O. A., Schwietzke, S., and Lan, X.: Global Inventory of Fossil and Non-fossil  $\delta^{13}\text{C}$ -CH<sub>4</sub> Source Signature Measurements for Improved Atmospheric Modeling, version 2020, *Earth Syst. Res. Lab.*, <https://doi.org/10.15138/qn55-e011>, 2021.
- 910 Shuai, Y., Douglas, P. M. J., Zhang, S., Stolper, D. A., Ellis, G. S., Lawson, M., Lewan, M. D., Formolo, M., Mi, J., He, K., Hu, G., and Eiler, J. M.: Equilibrium and non-equilibrium controls on the abundances of clumped isotopologues of methane during thermogenic formation in laboratory experiments: Implications for the chemistry of pyrolysis and the origins of natural gases, *Geochim. Cosmochim. Acta*, 223, 159–174, <https://doi.org/10.1016/j.gca.2017.11.024>, 2018a.
- 915 Shuai, Y., Etiope, G., Zhang, S., Douglas, P. M. J., Huang, L., and Eiler, J. M.: Methane clumped isotopes in the Songliao Basin (China): New insights into abiotic vs. biotic hydrocarbon formation, *Earth Planet. Sci. Lett.*, 482, 213–221, <https://doi.org/10.1016/j.epsl.2017.10.057>, 2018b.
- 920 Shuai, Y., Xie, H., Zhang, S., Zhang, Y., and Eiler, J.: Recognizing the pathways of microbial methanogenesis through methane isotopologues in the subsurface biosphere, *Earth Planet. Sci. Lett.*, 566, <https://doi.org/10.1016/j.epsl.2021.116960>, 2021.
- Sivan, M., Röckmann, T., van der Veen, Carina, and Popa, Maria Elena: Extraction, purification, and clumped isotope analysis of methane ( $^{13}\text{CDH}_3$  and  $^{12}\text{CD}_2\text{H}_2$ ) from sources and the atmosphere, *Atmos Meas Tech*, 17, 2687–2705, <https://doi.org/10.5194/amt-17-2687-2024>, 2024.
- 925 Stolper, D. A., Sessions, A. L., Ferreira, A. A., Santos Neto, E. V., Schimmelmann, A., Shusta, S. S., Valentine, D. L., and Eiler, J. M.: Combined  $^{13}\text{C}$ -D and D-D clumping in methane: Methods and preliminary results, *Geochim. Cosmochim. Acta*, 126, 169–191, <https://doi.org/10.1016/j.gca.2013.10.045>, 2014a.
- Stolper, D. A., Lawson, M., Davis, C. L., Ferreira, A. A., Neto, E. V. S., Ellis, G. S., Lewan, M. D., Martini, A. M., Tang, Y., Schoell, M., Sessions, A. L., and Eiler, J. M.: Formation temperatures of thermogenic and biogenic methane, *Science*, 344, 1500–1503, <https://doi.org/doi.10.1126/science.1254509>, 2014b.
- 930 Stolper, D. A., Martini, A. M., Clog, M., Douglas, P. M., Shusta, S. S., Valentine, D. L., Sessions, A. L., and Eiler, J. M.: Distinguishing and understanding thermogenic and biogenic sources of methane using multiply substituted isotopologues, *Geochim. Cosmochim. Acta*, 161, 219–247, <https://doi.org/10.1016/j.gca.2015.04.015>, 2015.

- Stolper, D. A., Lawson, M., Formolo, M. J., Davis, C. L., Douglas, P. M. J., and Eiler, J. M.: The utility of methane clumped isotopes to constrain the origins of methane in natural gas accumulations, *Geol. Soc. Lond. Spec. Publ.*, 468, 23–52, <https://doi.org/10.1144/SP468.3>, 2018.
- 935 Suda, K., Aze, T., Miyairi, Y., Yokoyama, Y., Matsui, Y., Ueda, H., Saito, T., Sato, T., Sawaki, Y., Nakai, R., Tamaki, H., Takahashi, H. A., Morikawa, N., and Ono, S.: The origin of methane in serpentinite-hosted hyperalkaline hot spring at Hakuba Happo, Japan: Radiocarbon, methane isotopologue and noble gas isotope approaches, *Earth Planet. Sci. Lett.*, 585, <https://doi.org/10.1016/j.epsl.2022.117510>, 2022.
- 940 Sun, J., Magen, C., Haghnegahdar, M. A., Liu, J., Fernandez, J. M., and Farquhar, J.: Constraining Wetland and Landfill Methane Emission Signatures Through Atmospheric Methane Clumped Isotopologue Measurements, *J. Geophys. Res.*, 130, <https://doi.org/10.1029/2024JG008249>, 2025a.
- 945 Sun, J., Haghnegahdar, M. A., Fernandez, J. M., Magen, C., and Farquhar, J.: Controls on concentrations and clumped isotopologues of vehicle exhaust methane, *PLOS ONE*, 20, e0315304, <https://doi.org/10.1371/journal.pone.0315304>, 2025b.
- Sun, T., Cao, J., Qiu, H., Fu, P., Lu, H., Ning, Z., Chen, D., Deng, Y., and Yang, S.: Investigation and optimization of methane purification method for natural gas by two-column gas chromatography: A preliminary test for doubly substituted isotopologue ( $^{13}\text{CH}_3\text{D}$ ) measurements, *Front. Mar. Sci.*, 10, <https://doi.org/10.3389/fmars.2023.969567>, 2023.
- 950 Taenzer, L., Labidi, J., Masterson, A. L., Feng, X., Rumble, D., Young, E. D., and Leavitt, W. D.: Low  $\Delta^{12}\text{CH}_2\text{D}_2$  values in microbialgenic methane result from combinatorial isotope effects, *Geochim. Cosmochim. Acta*, 285, 225–236, <https://doi.org/10.1016/j.gca.2020.06.026>, 2020.
- Thauer, R. K.: Biochemistry of methanogenesis: a tribute to Marjory Stephenson, *Microbiology*, 144, 2377–2406, 1998.
- 955 Thiagarajan, N., Kitchen, N., Xie, H., Ponton, C., Lawson, M., Formolo, M., and Eiler, J.: Identifying thermogenic and microbial methane in deep water Gulf of Mexico Reservoirs, *Geochim. Cosmochim. Acta*, 275, 188–208, <https://doi.org/10.1016/j.gca.2020.02.016>, 2020.
- Thiagarajan, N., Pedersen, J. H., Brunstad, H., Rinna, J., Lepland, A., and Eiler, J.: Clumped isotope constraints on the origins of reservoir methane from the Barents Sea, *Pet. Geosci.*, 28, <https://doi.org/10.1144/petgeo2021-037>, 2022.
- 960 Tsuji, K., Hiroaki Teshima, Hiroyuki Sasada, and Naohiro Yoshida: Spectroscopic isotope ratio measurement of doubly-substituted methane, *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.*, 98, 43–46, <http://dx.doi.org/10.1016/j.saa.2012.08.028>, 2012.
- Turner, A. J., Frankenberg, C., and Kort, E. A.: Interpreting contemporary trends in atmospheric methane, *Proc. Natl. Acad. Sci.*, 116, 2805–2813, <https://doi.org/10.1073/pnas.1814297116>, 2019.
- 965 Tyne, R. L., Barry, P. H., Lawson, M., Byrne, D. J., Warr, O., Xie, H., Hillegonds, D. J., Formolo, M., Summers, Z. M., Skinner, B., Eiler, J. M., and Ballentine, C. J.: Rapid microbial methanogenesis during CO<sub>2</sub> storage in hydrocarbon reservoirs, *Nature*, 600, 670–674, <https://doi.org/10.1038/s41586-021-04153-3>, 2021.
- Walter Anthony, K. M., Anthony, P., and Hasson, N.: Upland Yedoma taliks are an unpredicted source of atmospheric methane, *Nat. Commun.*, 15, <https://doi.org/10.1038/s41467-024-50346-5>, 2024.
- 970 Wang, D. T., Gruen, D. S., Sherwood Lollar, B., Hinrichs, K.-U., Stewart, L. C., Holden, J. F., Hristov, A. N., Pohlman, J. W., Morrill, P. L., Könneke, M., Delwiche, K. B., Reeves, E. P., Sutcliffe, C. N., Ritter, D. J., Seewald, J. S., McIntosh, J. C., Hemond, H. F., Kubo, M. D., Cardace, D., Hoehler, T. M., and Ono, S.: Nonequilibrium clumped isotope signals in microbial methane, *Science*, 348, 428–431, <https://doi.org/10.1126/science.aaa4326>, 2015.

- 975 Wang, D. T., Welander, P. V., and Ono, S.: Fractionation of the methane isotopologues  $^{13}\text{CH}_4$ ,  $^{12}\text{CH}_3\text{D}$ , and  $^{13}\text{CH}_3\text{D}$  during aerobic oxidation of methane by *Methylococcus capsulatus* (Bath), *Geochim. Cosmochim. Acta*, 192, 186–202, <http://dx.doi.org/10.1016/j.gca.2016.07.031>, 2016.
- Wang, D. T., Reeves, E. P., McDermott, J. M., Seewald, J. S., and Ono, S.: Clumped isotopologue constraints on the origin of methane at seafloor hot springs, *Geochim. Cosmochim. Acta*, 223, 141–158, 980 <https://doi.org/10.1016/j.gca.2017.11.030>, 2018.
- Wang, D. T., Sattler, A., Paccagnini, M., and Chen, F. G.: Method for calibrating methane clumped isotope measurements via catalytic equilibration of methane isotopologues on  $\gamma$ -alumina, *Rapid Commun. Mass Spectrom.*, 34, <https://doi.org/10.1002/rcm.8555>, 2019.
- Wang, X., Cong-Qiang Liu, Naizhong Zhang, Sheng Xu, Zhiyong Pang, Si-Liang Li, Hu Ding, Jianfa Chen, Zengye Xie, 985 and Rob M. Ellama: Clumped methane isotopologues ( $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$ ) of natural samples measured using a high-resolution mass spectrometer with an improved pretreatment system, *J. Anal. At. Spectrom.*, <https://doi.org/DOI: 10.1039/d2ja00315e>, 2023a.
- Wang, X., Liu, C.-Q., Yi, Y., Zeng, M., Li, S.-L., and Niu, X.: Machine Learning Predicts the Methane Clumped Isotopologue ( $^{12}\text{CH}_2\text{D}_2$ ) Distributions Constrain Biogeochemical Processes and Estimates the Potential Budget, 990 *Environ. Sci.*, 57, 17876–17888, <https://doi.org/10.1021/acs.est.3c00184>, 2023b.
- Wang, X., Chen, B., Chen, L., Dong, G., Csernica, T., Zhang, N., Liu, J., Shuai, Y., Liu, C.-Q., Xu, Z., Li, S.-L., and Xu, S.: Biogenic methane clumped isotope signatures: Insights from microbially enhanced coal bed methane, *Fuel*, 365, <https://doi.org/10.1016/j.fuel.2024.131307>, 2024a.
- Wang, X., Biying Chen, Hui Nai, Cong-Qiang Liu, Guannan Dong, Naizhong Zhang, Si-Liang Li, Jonathan Gropp, 995 Jennifer McIntosh, Rob M. Ellam, John M. Eiler, and Sheng Xu: Clumped isotopes constrain thermogenic and secondary microbial methane origins in coal bed methane, *Earth Planet. Sci. Lett.*, 647, 119023, <https://doi.org/10.1016/j.epsl.2024.119023>, 2024b.
- Wang, X., Chen, B., Dong, G., Zhang, N., Liu, W., Han, J., Liu, C.-Q., Li, S.-L., Eiler, J. M., and Xu, S.: Microbial contribution estimated by clumped isotopologues ( $^{13}\text{CH}_3\text{D}$  and  $^{12}\text{CH}_2\text{D}_2$ ) characteristics in a  $\text{CO}_2$  enhanced 1000 coal bed methane reservoir, *Sci. Total Environ.*, 922, 170926, <https://doi.org/10.1016/j.scitotenv.2024.170926>, 2024c.
- Wang, Z., Schauble, E. A., and Eiler, J. M.: Equilibrium thermodynamics of multiply substituted isotopologues of molecular gases, *Geochim. Cosmochim. Acta*, 68, 4779–4797, <https://doi.org/10.1016/j.gca.2004.05.039>, 2004.
- Warr, O., Young, E. D., Giunta, T., Kohl, I. E., Ash, J. L., and Sherwood Lollar, B.: High-resolution, long-term isotopic and isotopologue variation identifies the sources and sinks of methane in a deep subsurface carbon cycle, 1005 *Geochim. Cosmochim. Acta*, 294, 315–334, <https://doi.org/10.1016/j.gca.2020.12.002>, 2021a.
- Warr, O., Young, E. D., Giunta, T., Kohl, I. E., Ash, J. L., and Sherwood Lollar, B.: High-resolution, long-term isotopic and isotopologue variation identifies the sources and sinks of methane in a deep subsurface carbon cycle, *Geochim. Cosmochim. Acta*, 294, 315–334, <https://doi.org/10.1016/j.gca.2020.12.002>, 2021b.
- 1010 Webb, M. A. and Miller, T. F.: Position-Specific and Clumped Stable Isotope Studies: Comparison of the Urey and Path-Integral Approaches for Carbon Dioxide, Nitrous Oxide, Methane, and Propane, *J. Phys. Chem. A*, 118, 467–474, <https://doi.org/10.1021/jp411134v>, 2014.
- Whitehill, A. R., Joelsson, L. M. T., Schmidt, J. A., Wang, D. T., Johnson, M. S., and Ono, S.: Clumped isotope effects during OH and Cl oxidation of methane, *Geochim. Cosmochim. Acta*, 196, 307–325, 1015 <https://doi.org/10.1016/j.gca.2016.09.012>, 2017.
- Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem. Geol.*, 161, 291–314, [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3), 1999.

- Xia, X. and Gao, Y.: Kinetic clumped isotope fractionation during the thermal generation and hydrogen exchange of methane, *Geochim. Cosmochim. Acta*, 248, 252–273, <https://doi.org/10.1016/j.gca.2019.01.004>, 2019.
- 1020 Xie, H., Dong, G., Formolo, M., Lawson, M., Liu, J., Cong, F., Mangenot, X., Shuai, Y., Ponton, C., and Eiler, J.: The evolution of intra- and inter-molecular isotope equilibria in natural gases with thermal maturation, *Geochim. Cosmochim. Acta*, 307, 22–41, <https://doi.org/10.1016/j.gca.2021.05.012>, 2021.
- Yeung, L. Y.: Combinatorial effects on clumped isotopes and their significance in biogeochemistry, *Geochim. Cosmochim. Acta*, 172, 22–38, <https://doi.org/10.1016/j.gca.2015.09.020>, 2016.
- 1025 Young, E. D.: A Two-Dimensional Perspective on CH<sub>4</sub> Isotope Clumping: Distinguishing Process from Source, in: *Deep Carbon*, edited by: Orcutt, B. N., Daniel, I., and Dasgupta, R., Cambridge University Press, 388–414, <https://doi.org/10.1017/9781108677950.013>, 2019.
- Young, E. D., Rumble, D., Freedman, P., and Mills, M.: A large-radius high-mass-resolution multiple-collector isotope ratio mass spectrometer for analysis of rare isotopologues of O<sub>2</sub>, N<sub>2</sub>, CH<sub>4</sub> and other gases, *Int. J. Mass Spectrom.*, 401, 1–10, <https://doi.org/10.1016/j.ijms.2016.01.006>, 2016.
- 1030 Young, E. D., Kohl, I. E., Sherwood Lollar, B., Etiope, G., Rumble, D., Li (李姝宁), S., Haghnegahdar, M. A., Schauble, E. A., McCain, K. A., Foustoukos, D. I., Sutcliffe, C., Warr, O., Ballentine, C. J., Onstott, T. C., Hosgormez, H., Neubeck, A., Marques, J. M., Pérez-Rodríguez, I., Rowe, A. R., LaRowe, D. E., Magnabosco, C., Yeung, L. Y., Ash, J. L., and Bryndzia, L. T.: The relative abundances of resolved <sup>12</sup>CH<sub>2</sub>D<sub>2</sub> and <sup>13</sup>CH<sub>3</sub>D and mechanisms controlling isotopic bond ordering in abiotic and biotic methane gases, *Geochim. Cosmochim. Acta*, 203, 235–264, <https://doi.org/10.1016/j.gca.2016.12.041>, 2017.
- 1035 Young, E. D., Labidi, J., and Kohl, I. E.: Advances in measuring multiply-substituted isotopologues of gas molecules with geochemical applications, in: *Treatise on Geochemistry*, Elsevier, 645–670, 2025.
- 1040 Zhang, N., Snyder, G. T., Lin, M., Nakagawa, M., Gilbert, A., Yoshida, N., Matsumoto, R., and Sekine, Y.: Doubly substituted isotopologues of methane hydrate (<sup>13</sup>CH<sub>3</sub>D and <sup>12</sup>CH<sub>2</sub>D<sub>2</sub>): Implications for methane clumped isotope effects, source apportionments and global hydrate reservoirs, *Geochim. Cosmochim. Acta*, 315, 127–151, <https://doi.org/10.1016/j.gca.2021.08.027>, 2021.
- 1045 Zhang, N., Prokhorov, I., Kueter, N., Li, G., Tuzson, B., Magyar, P. M., Ebert, V., Sivan, M., Nakagawa, M., Gilbert, A., Ueno, Y., Yoshida, N., Röckmann, T., Bernasconi, S. M., Emmenegger, L., and Mohn, J.: Rapid High-Sensitivity Analysis of Methane Clumped Isotopes ( $\Delta^{13}\text{CH}_3\text{D}$  and  $\Delta^{12}\text{CH}_2\text{D}_2$ ) Using Mid-Infrared Laser Spectroscopy, *Anal. Chem.*, <https://doi.org/10.1021/acs.analchem.4c05406>, 2025.