

Recent summer soil moisture drying in Switzerland based on measurements from the SwissSMEX network

Martin Hirschi¹, Dominik Michel¹, Dominik L. Schumacher¹, Wolfgang Preimesberger², Sonia I. Seneviratne¹

¹ETH Zurich, Institute for Atmospheric and Climate Science, Universitätstrasse 16, 8092 Zürich, Switzerland

²TU Wien, Department of Geodesy and Geoinformation, Wiedner Hauptstrasse 8-10, 1040 Vienna, Austria

Correspondence to: Martin Hirschi (martin.hirschi@env.ethz.ch)

Abstract. We present curated time series of in situ soil moisture measurements from the Swiss Soil Moisture Experiment (SwissSMEX) network, covering the period 2010–2025. At ten grassland stations on the Swiss Plateau, SwissSMEX provides volumetric soil water content (VWC) data at multiple depths in the soil profile, typically down to a maximum depth of 120 cm. Measurements are obtained from capacitance sensors, as well as from time domain reflectometry (TDR) sensors. The VWC measurements are used to compute integrated soil water content (IWC) down to 50 cm depth as an indicator of root-zone water. We document recent measures that have been taken to secure the SwissSMEX network and to ensure the continuity of its long-term IWC time series. The IWC and the underlying VWC time series are available at <https://doi.org/10.3929/ethz-b-000743711> (Hirschi et al., 2025a).

As a use case for the SwissSMEX network, the data are used to investigate drying trends on the Swiss Plateau, where notably drier summers and more frequent droughts have been reported in recent decades. This application of the in situ data is based on summer and summer-half-year anomalies of IWC. Furthermore, the SwissSMEX-based IWC anomalies and trends therein are compared with those derived from soil moisture of a widely used land reanalysis product (ERA5-Land) and a merged passive microwave remote-sensing product (C3S SM PASSIVE). The in situ IWC time series from stations with best temporal coverage agree well with ERA5-Land and C3S SM PASSIVE soil moisture from the corresponding grid cells, with correlations of 0.79 or higher for the median time series calculated across these stations. The ~~non-significant~~ drying over the common period 2010–2025 amounts to -1.4 mm yr^{-1} for SwissSMEX, ~~to -1.2 mm yr^{-1} for ERA5-Land, and to -1.1 mm yr^{-1} for C3S SM PASSIVE~~ in summer. Although the SwissSMEX network indicates that summer soil drying has increased in recent years, the 16 years of in situ data currently available are not yet sufficient to robustly estimate a significant trend. This highlights the importance of sustaining ongoing measurements to ensure a seamless continuation of soil moisture monitoring in Switzerland.

Deleted: and

Deleted: 1

1 Introduction

Droughts in Switzerland have become a significant concern due to their impacts on agriculture, water resources management, energy production, shipping and ecology. After a prolonged period without severe droughts, Switzerland has experienced increasingly frequent events in the last two decades, namely in the summers of 2003, 2015, 2018, and 2022, and in the
35 springs/early summers of 2011 and 2020 (Calanca, 2007; Brunner et al., 2019; Orth et al., 2016; Schumacher et al., 2024; MeteoSchweiz, 2018, 2012; BAFU, 2016). As a result, notable trends towards drier summer half-years were observed in Switzerland from 1981 to 2020 based on the climatic water balance calculated from meteorological station observations and the soil water content from reanalyses (Scherrer et al., 2022; hereafter referred to as S22). Furthermore, S22 showed that this
40 drying trend is primarily driven by a non-significant decrease in precipitation and a significant increase in evapotranspiration due to rising temperatures. However, the considered reanalysis products (ERA5-Land and ERA5) differed considerably in their representation of summer-half-year soil moisture and evapotranspiration in Switzerland. Both showed a significant increase in evapotranspiration from 1980 to 2020, but the trend is about twice as strong in ERA5-Land as in ERA5. The latter product already indicated a soil moisture limitation of evapotranspiration in dry years, whereas the former shows an evapotranspiration surplus in these years. Consequently, dry anomalies and the drying trend in soil moisture were more
45 pronounced in ERA5 than in ERA5-Land, with the latter agreeing better with a limited set of in situ stations.

Given these existing uncertainties in the reanalysis data, we present curated in situ time series from the Swiss Soil Moisture Experiment (SwissSMEX) soil moisture network (Mittelbach and Seneviratne, 2012). Although in situ soil moisture measurements have their own inherent problems and challenges (e.g. due to soil heterogeneity, limited representativeness of
50 point measurements, and sensor calibration issues), they are still the only available ground information to contrast with land reanalysis or remotely sensed soil moisture. The SwissSMEX network was installed as a research network between 2008 and 2010. Since then, it has measured volumetric water content (VWC) and soil temperature in profiles from 5 cm down to a maximum depth of typically 120 cm using time domain reflectometry (TDR) and capacitance sensors (Mittelbach and Seneviratne, 2012; Mittelbach, 2011). The number of depth levels varies across the network, and redundant profiles of the
55 same sensor type are available at some stations. The VWC measurements are also used to calculate integrated soil water content (IWC, in mm) down to 50 cm depth as an indicator of root-zone water, which is relevant for plant growth in agriculture and natural vegetation, and is directly related to land water-balance fluxes. While root depths vary greatly in space and depend on plant species, the choice of 50 cm depth covers shallow-rooted plants such as lettuce, onions, potatoes, various vegetables and most native grass species (Brouwer et al., 1989). In addition, this depth is covered by all stations of the network.

60 Ten SwissSMEX grassland sites on the Swiss Plateau are still operational (Fig. 1), but in recent years sensors have started to fail frequently, leading to temporal interruptions or the complete termination of individual measurement series (Michel et al., 2022b). This prompted an intervention to preserve the measurement network for the coming years. These sensor failures are

Deleted: T

Deleted: S22 showed, h

Deleted: that

Deleted: s

Deleted: are

70 particularly problematic for deriving IWC, which relies on the simultaneous availability of VWC measurements from sensors
at multiple depths. They also impair the capacity of the network for real-time monitoring and trend detection. This is especially
relevant as the increasing frequency of droughts in recent years has prompted the Swiss Federal Council to initiate a programme
in May 2022 to establish and expand a national early detection and warning system for droughts, including a national in situ
soil moisture network (BAFU, 2024). The implementation of the national network at existing SwissSMEX sites will benefit
from the secured long-term records.

75 This data paper presents and documents the SwissSMEX long-term soil moisture time series record and recent measures that
have been taken to secure its operational status. It revisits the documented summer drying trends on the Swiss Plateau to
demonstrate the validity and applicability of the presented in situ measurements and compares SwissSMEX-based trends with
those derived from alternate land reanalysis and remote sensing soil moisture products, extending on the analysis of S22. The
80 in situ time series presented here also contributed to the CH2025 national climate scenarios (MeteoSwiss and ETH Zurich,
2025).




Figure 1 Locations of the SwissSMEX stations on the Swiss Plateau, along with the nearest grid cells of ERA5-Land and C3S SM PASSIVE (see Sect. 2.2). Station abbreviations are provided in Table 1.

90 **2 Data and methods**

2.1 SwissSMEX

2.1.1 Status of the network and new installations

The standard instrumentation of SwissSMEX, initiated in 2008, consisted of low-cost capacitance soil moisture sensors (Decagon 10HS) at all measurement depths of 5, 10, 30, 50, 80, and 120 cm, and of TDR soil moisture sensors (IMKO TRIME-
95 PICO) at 10 and 80 cm depth (Mittelbach, 2011). Both sensor types provide volumetric water content (VWC, in Vol.% or m³ m⁻³). In the follow-up project SwissSMEX-*Veg*, selected stations were additionally equipped with TDR sensors at 5, 30 and 50 cm depth in 2010. Ten SwissSMEX grassland sites on the Swiss Plateau are still operational (Fig. 1, Table 1). The installed low-cost 10HS sensors are prone to higher measurement errors when only the factory calibration is used (e.g., Domínguez-Niño et al., 2019; Mittelbach et al., 2012). Therefore, site-specific exponential calibration functions were derived
100 from the parallel 10HS and TDR measurements and applied to the raw 10HS readings. These site-specific calibrations allow to reduce the absolute error to about 2 Vol.% for most 10HS measurements, including under dry and moist soil conditions, and extend the effective measurement range of the sensors to up to 50 Vol.% (Mittelbach et al., 2011).


105 **Table 1 Overview of the ten SwissSMEX stations on the Swiss Plateau (see also Fig. 1). For more detailed information on soil properties, we refer to Mittelbach (2011, Appendix B therein). The crosses indicate the different station combinations used to calculate the Swiss Plateau median summer and summer-half-year soil moisture anomalies (see Sect. 2.3), and “Missing years” indicates the absence of these anomalies for specific years.**

Station	Full name	Coordinates (lat, lon)	Soil texture	Best coverage ¹	S22	S22 + best coverage	Missing years	
							JJA	AMJJAS
BAS	Basel	47.54, 7.58	silt loam		X	X	2014–2022	2014–2022
BER	Bern	46.99, 7.46	loam	X	X	X	2011	none
CHM	Chamau	47.21, 8.41	sandy loam	X		X	2021	none
CHN	Changins	46.40, 6.23	loam		X	X	2010, 2017, 2018, 2020	2010, 2017, 2018
PAY	Payerne	46.81, 6.94	loam				2021, 2022	2021, 2022
PLA	Plaffeien	46.75, 7.27	sandy loam	X		X	none	none
REC	Reckenholz	47.43, 8.52	loam	X	X	X	2021, 2024	2021, 2024
RHB	Rietholzbach	47.38, 8.99	loam				2014, 2015	2014, 2015
TAE	Tänikon	47.48, 8.90	loam	X		X	2023–2025	2023–2025
WYN	Wynau	47.26, 7.79	silt loam				2010–2012, 2014, 2015, 2021, 2022	2010–2012, 2014, 2021, 2022

¹ maximum one summer/summer half-year missing in the historical 2010–2022 period (see Sect. 2.1.2)

The VWC is measured as 10-minute instantaneous values, which are then further aggregated to hourly and daily values. As
 110 part of the regular processing of the SwissSMEX measurements, the individual sensor series undergo automatic range and
 outlier checks, as well as visual inspection. The valid data range is defined as 0–0.7 m³ m⁻³, and spikes in the ten-minute values
 are removed by applying a one-day moving average and a threshold of ± two standard deviations. A comprehensive quality
 assessment and intercomparison of the existing soil moisture time series from the SwissSMEX network revealed a degradation
 in data availability due to sensor failures over time, compromising the long-term continuity of the time series (Michel et al.,
 115 2022b). This analysis showed that, as of 2022, more than one-third of all sensors had ceased to function since their installation
between 2008 and 2010 (Figs. 2 and 3). This sensor failure rate is higher for the 10HS capacitance sensors (43 %) than for the
 TDR sensors (36 %; cf. Appendix A of Michel et al., 2022b).

Deleted: had stopped working since the start of the installation in 2008



120 **Figure 2** Time series diagram of the IMKO TRIME-PICO TDR (TD) sensor availability at the SwissSMEX grassland stations on the Swiss Plateau. Abbreviations for station names are given in Table 1; numbers alongside the sensor type indicate the installation depth in cm. This is a condensed view that combines all redundant sensor profiles. Note that station RHB includes an additional sensor at 25 cm depth, and that the sensors at the other depths are partly shifted by 5 cm in this profile. For simplicity, the sensors installed at depths of 15 cm, 35 cm, and 55 cm are shown as TD_010, TD_030, and TD_050, respectively. The TDR installations in 2022 are highlighted by red rectangles.




Figure 3 As Fig. 2, but for the Decagon 10HS (HS) sensors. In addition to the note on station RHB, REC has a 10HS sensor at 150 cm depth instead of 120 cm, which is shown as HS_120 for simplicity.

In light of this degrading data availability, and to secure the network for the coming years, defective TDR sensors at 10 and 30 cm soil depth were replaced with new IMKO TRIME-PICO64 TDR sensors at the SwissSMEX grassland stations during summer 2022, or, where not already present, additional TDR sensors were installed at these depths in the existing profiles (Fig. 2; Michel et al., 2022a). This intervention at easily accessible depths, carried out without disturbing the ongoing operations, ensured the continuity of the measurement series at these depths.

2.1.2 Vertically integrated soil water content calculation

The degradation of sensor availability is particularly relevant for deriving the 0–0.5 m depth integrated soil water content (IWC, in mm). This integration is performed using the trapezoidal method (e.g. Hupet et al., 2004) and relies on the simultaneous availability of VWC measurements from sensors at multiple depths. The trapezoidal integration approximates the integral by assuming a linear variation between the measurements at the individual depths and was computed using the R-function trapz from the caTools package (Tuszynski, 2021; version 1.18.2). Historically, the integration was performed based

on the n VWC measurements (VWC_i) at depths z_i of 0.05, 0.10, 0.30 and 0.50 m (Mittelbach and Seneviratne, 2012), plus an additional value of VWC at the surface, which is set equal to the measurement at 0.05 m depth (see Eq. 1). These depths were originally fully equipped with 10HS sensors, whereas TDR sensors were predominantly installed at 10 and 80 cm depth (Mittelbach et al., 2011; see also Figs. 2 and 3). Over time, at some stations, already existing TDR sensors were incorporated into the integration to replace defective 10HS sensors (e.g. station Changins in Fig. 5), or the number of sensors (i.e. depths) included in the integration had to be reduced while ensuring the representativeness of the remaining sensor data (e.g. station Chamau in Fig. 4). This set of sensors used for the historical IWC time series is referred to as the original sensor configuration (IWC_{orig} , see also Table S1).

$$IWC_{\text{orig}} \text{ (in mm)} = \left(VWC_{0.05\text{m}} \cdot 0.05 \text{ m} + \sum_{i=1}^{n-1} \frac{VWC_{i+1} + VWC_i}{2} \cdot (z_{i+1} - z_i) \right) \cdot 1000 \quad (1)$$

With the increasing number of sensor failures in recent years (Figs. 2 and 3), however, the calculation of IWC_{orig} based on the original sensor configurations has become impossible at a growing number of sites. With the upgrade of the network in 2022, all grassland stations are now equipped with at least two TDR sensors in the 0–0.5 m soil column, i.e. at 10 and 30 cm depth. These sensors are used as an alternative for deriving the IWC when coverage with the original sensor configuration is lacking. For this TDR-only based IWC_{TDRonly} , the 10 cm measurement is used to represent the 5 cm and 10 cm depths, and the 30 cm measurement is used to represent the 30 cm and 50 cm depths (Eq. 2).

$$IWC_{\text{TDRonly}} \text{ (in mm)} = \left(VWC_{0.10\text{m}} \cdot 0.1 \text{ m} + \frac{VWC_{0.3\text{m}} + VWC_{0.1\text{m}}}{2} \cdot 0.2 \text{ m} + VWC_{0.30\text{m}} \cdot 0.2 \text{ m} \right) \cdot 1000 \quad (2)$$

The historical hourly IWC time series (IWC_{orig}) and the 10 and 30 cm TDR-only based series (IWC_{TDRonly}) are then merged by correcting the mean offset and difference in standard deviation of the latter with respect to the former to account for the difference in the representative depth (Eq. 3a,b).

$$IWC_{\text{TDRonly,corrected}} = \left(IWC_{\text{TDRonly}} - \overline{IWC_{\text{TDRonly}}} \right) / \text{relsd} + \overline{IWC_{\text{orig}}} \quad (3a)$$

$$\text{relsd} = \text{sd}(IWC_{\text{TDRonly}}) / \text{sd}(IWC_{\text{orig}}) \quad (3b)$$

The overbar denotes the temporal average, sd the standard deviation, and relsd the ratio of the standard deviations. The offset and relative standard deviation between the two series are calculated from the overlapping hourly time steps (except for BAS and WYN, where there is no overlap and all time steps are therefore used to calculate the scaling parameters). This correction assures that the data distribution of $IWC_{\text{TDRonly,corrected}}$ matches that of IWC_{orig} (in terms of mean and standard deviation). Merging IWC_{orig} and $IWC_{\text{TDRonly,corrected}}$ substantially increases recent station coverage with IWC. In some cases, historical

gaps can also be filled in this way for stations that already had TDR sensors at 10 and 30 cm depth prior to 2022. This is illustrated in Figs. 4 and 5 at the examples of the stations Chamau (CHM) and Changins (CHN), respectively. At Chamau, the historical IWC time series could be extended to the present using the new TDR sensors, while at Changins, historical gaps in IWC could additionally be filled. Corresponding plots for the other stations can be found in Figs. S1–S4.





Figure 4 Daily time series of volumetric water content (VWC, in $\text{m}^3 \text{m}^{-3}$) and vertically integrated soil water content (IWC, in mm) at the station Chamau (CHM), where IWC_{orig} could be extended to the present. (Top) VWC from the 10HS sensors contributing to IWC_{orig} , based on the original sensor configuration (see also Table S1), and from the two new TDRs installed in 2022 that are used to calculate $\text{IWC}_{\text{TDRonly}}$. (Bottom) IWC_{orig} and the 10 and 30 cm TDR-only based $\text{IWC}_{\text{TDRonly}}$ and $\text{IWC}_{\text{TDRonly_corrected}}$ (see Eq. 3a,b). The original offset of $\text{IWC}_{\text{TDRonly}}$ is also noted, as well as its relative standard deviation and RMSD compared with IWC_{orig} (all metrics are based on the overlapping hourly time steps). See Figs. S1–S4 for similar plots for the other stations.



185 **Figure 5** As Fig. 4 but for Changins (CHN), where historical gaps in IWC_{orig} could be filled. Note that $IWC_{TDOnly_corrected}$ is only shown when IWC_{orig} is not available.

2.2 Comparison datasets

2.2.1 ERA5-Land

190 The land component of the ERA5 reanalysis (Hersbach et al., 2020) provides global, hourly, high-resolution information of the water and energy cycles over land in a consistent representation (Muñoz-Sabater et al., 2021). ERA5-Land is a single simulation based on the land-surface model HTESSEL (Hydrology-Tiled ECMWF Scheme for Surface Exchanges over Land, Balsamo et al., 2009) forced by ERA5 near-surface atmospheric reanalysis fields, with additional lapse-rate correction of temperature. HTESSEL distinguishes between four different soil layers with the following layer depths: layer 1 at 0–7 cm; layer 2 at 7–28 cm; layer 3 at 28–100 cm; and layer 4 at 100–289 cm. ERA5-Land soil moisture is regularly used in the analyses
195 of the Copernicus European State of the Climate Reports. The data have been extracted on a $0.1^\circ \times 0.1^\circ$ latitude–longitude grid with monthly temporal resolution from the Copernicus Climate Data Store (CDS; C3S, 2022). For the comparison with SwissSMEX, the closest grid cells were used for each station location (see [Figure 1](#)).

Deleted: Table

200 The IWC in the 0–0.5 m soil layer, as available from SwissSMEX, has been calculated based on VWC of the soil layers 1–3:

$$IWC_{0-0.5m} \text{ (in mm)} = (VWC_{0-7cm} \cdot 0.07 \text{ m} + VWC_{7-28cm} \cdot 0.21 \text{ m} + VWC_{28-100cm} \cdot 0.22 \text{ m}) \cdot 1000 \quad (4)$$

The assumption is that the $VWC_{28-100cm}$ is representative for the 22 cm between 28 and 50 cm soil depth.

2.2.2 C3S satellite soil moisture

205 The Copernicus Climate Change Service (C3S) provides up-to-date global satellite surface soil moisture data through the CDS every 10 days (C3S SM; Dorigo et al., 2025). The algorithm for v202505 is the same one as in version 9 of the European Space Agency Climate Change Initiative (ESA CCI) soil moisture products (Dorigo et al., 2024; Dorigo et al., 2017; Gruber et al., 2019). The PASSIVE product (representative at ~2–5 cm depth) is purely satellite (radiometer)-based and therefore preferred for our study over COMBINED, which may adopt certain model features (Dorigo et al., 2017). The C3S SM PASSIVE data were gap-filled using a stand-alone 3D-interpolation framework, independent of ancillary variables (Preimesberger et al., 210 2025). Root-zone soil moisture was derived by extrapolating gap-filled surface estimates to deeper soil layers using an exponential filter (Wagner et al., 1999; Albergel et al., 2008), calibrated for four layers with a large number of in situ time series (Pasik et al., 2023): layer 1 at 0–10 cm; layer 2 at 10–40 cm; layer 3 at 40–100 cm; and layer 4 at 100–200 cm. These data are also being used in the Copernicus European State of the Climate Reports (C3S and WMO, 2025) and available on a 0.25° x 0.25° latitude–longitude grid with daily temporal resolution. The closest grid cells for each station location (see 215 [Figure 1](#)) were used for comparison with SwissSMEX, and we restrict the usage of C3S SM PASSIVE to data from 1991 onwards due to larger data gaps prior to this date in the non-gap-filled product, and the thereby caused high uncertainties associated with the interpolated values (Preimesberger et al., 2025).

Deleted: Table

The IWC in the 0–0.5 m soil layer has been calculated based on VWC of the soil layers 1–3:

$$220 \quad IWC_{0-0.5m} \text{ (in mm)} = (VWC_{0-10cm} \cdot 0.1 \text{ m} + VWC_{10-40cm} \cdot 0.3 \text{ m} + VWC_{40-100cm} \cdot 0.1 \text{ m}) \cdot 1000 \quad (5)$$

The assumption is that the $VWC_{40-100cm}$ is representative for the 10 cm between 40 and 50 cm soil depth.

2.3 Calculation of summer and summer-half-year anomalies and trends therein

To demonstrate the utility of the SwissSMEX data, it is applied to investigate the temporal evolution and drying trends in mean summer (June to August, JJA) and summer-half-year (April to September, AMJJAS) soil moisture anomalies. Calculated based 225 on the daily IWC time series (i.e. the merged IWC_{orig} and $IWC_{TDRonly_corrected}$; see Sect. 2.1.2), the anomalies are expressed as absolute or percentage deviations from the corresponding summer ([Eq. 6a,b](#)) or summer-half-year mean for 2010–2025 (i.e. in mm or % relative to the 2010–2025 mean, respectively).

Deleted: 0

Deleted: 0

$$IWC_{mm_anomaly} \text{ (in mm)} = (IWC_{JJA} - \overline{IWC_{JJA}}) \quad (6a)$$

$$IWC_{\%_anomaly} \text{ (in \%)} = \frac{IWC_{JJA} - \overline{IWC_{JJA}}}{\overline{IWC_{JJA}}} * 100 \quad (6b)$$

IWC_{JA} are the annual mean summer values calculated from the daily IWC time series, and the overbar denotes the temporal average over 2010–2025. The summer-half-year anomalies are calculated analogously. Note that the percentage anomalies should not be confused with the Vol.% units of the VWC sensor readings. We deliberately allow up to 35 % missing days when calculating the mean anomalies for each summer and summer half-year, respectively. This choice represents a compromise between the number of stations available for the summer drying analysis (see Table 1) and ensuring reasonable annual data coverage. We also note that certain use cases might warrant stricter data availability thresholds.

Based on the soil moisture anomalies, we test different combinations of stations to investigate the median evolution (and trends therein) of summer and summer-half-year soil moisture on the Swiss Plateau. “Best coverage” includes stations with at most one summer/summer half-year missing within the historical 2010–2022 period (i.e. until the new TDR sensor installations); “S22” includes the stations used in S22 (representative of an average elevation on the Swiss Plateau); “S22 + best coverage” combines both sets; and we also consider the combination of “all stations” listed in Table 1.

The robust Theil-Sen trend estimator (Theil, 1950; Sen, 1968) is used to compute trends, while the non-parametric Mann-Kendall trend test is used to determine trend significance (Mann, 1945; Kendall, 1975). These allow the detection of monotonic changes over time and are robust to non-normally distributed data. Statistical significance is indicated by a p-value below 0.05. The R-functions sens.slope and mk.test from the trend package (Pohlert, 2020; version 1.1.4) are used for the calculations. Trends are computed based on the median time series of the respective station combinations.

3 Application of SwissSMEX for investigating summer soil moisture anomalies and trends

3.1 Swiss Plateau station combinations

Figure 6 presents the individual soil moisture anomaly time series of the summer IWC at SwissSMEX stations (2010–2025) for each station combination considered (see Sect. 2.3, Table 1), along with the respective median anomalies. While there is some variation in the anomalies across the individual stations, the station medians of the four station combinations exhibit a consistent temporal evolution. This is confirmed by the pairwise Pearson correlations between the median time series, which are all at least 0.8₉, and by the corresponding root-mean-square deviations (RMSD), which reach at most 7.0 mm (Fig. 7, values in the red dashed rectangle). Similar values of agreement are obtained when summer-half-year instead of summer soil moisture is considered (Fig. S5). Thus, the median evolution of IWC on the Swiss Plateau, based on different combinations of stations, is largely consistent.

Deleted: these

Deleted: values

Deleted: 9

Deleted: 6.9







Figure 6 Soil moisture anomaly time series of summer IWC of different SwissSMEX station combinations (see Sect. 2.3, Table 1) and respective station medians. (a) Stations with best temporal coverage, (b) station selection as in S22, (c) S22 and best coverage stations combined, (d) all Swiss Plateau stations. Anomalies are expressed as the absolute deviations from the summer mean over 2010–2025. The straight lines show the Theil-Sen trend slopes for the period 2010–2025 based on the respective station median time series (solid lines indicate a significant trend with $p < 0.05$; dashed lines indicate a non-significant trend).




Deleted:



Deleted:



Deleted:



Deleted:




Figure 7 Pearson correlation and RMSD matrices from the pairwise comparison of SwissSMEX, ERA5-Land and C3S SM PASSIVE soil moisture, considering the different station combinations for the Swiss Plateau median time series of summer soil moisture anomalies (expressed as the absolute deviations from the 2010–2025 summer mean). Values are calculated for the common period 2010–2025. All correlations are significant ($p < 0.05$). The red dashed rectangle highlights the comparison of the different station combinations based on SwissSMEX, while the red solid rectangle highlights the comparison of the gridded products and SwissSMEX based on the best coverage station combination. See Fig. S5 for the corresponding summer-half-year correlations and RMSDs.

Deleted:

Deleted: 0

295 All SwissSMEX station combinations also show a (non-significant) tendency towards drying (see straight lines in Fig. 6). The
respective Theil-Sen trend slopes based on the period 2010–2025 range from -1.4 mm yr⁻¹ to -0.7 mm yr⁻¹ for summer, and from
-0.6 mm yr⁻¹ to -0.2 mm yr⁻¹ for summer half-year, when trends tend to be lower (Table 2). The respective trends for the
percentage anomalies amount to -0.9 % yr⁻¹ to -0.5 % yr⁻¹ for summer, and to -0.4 % yr⁻¹ to -0.1 % yr⁻¹ for summer half-year.
Thus, the recent soil drying tendency on the Swiss Plateau is noticeable in all combinations of SwissSMEX stations used for
the median time series (see also Fig. 5). However, the trends are not significant and show some degree of variation in their
magnitude between the different station combinations.

300

Table 2 Theil-Sen trend slopes of summer and summer-half-year IWC anomalies based on the median of different station combinations (i.e. best coverage, S22, S22 + best coverage, all stations). Anomalies are expressed as the absolute deviations from the summer resp. summer-half-year mean over 2010–2025, and as percentage deviations from this mean. Trend slopes are based on the period 2010–2025, and trend significance is determined using the non-parametric Mann-Kendall trend test.

	Dataset	Best coverage		S22		S22 + best coverage		All stations	
		mm yr ⁻¹	% yr ⁻¹	mm yr ⁻¹	% yr ⁻¹	mm yr ⁻¹	% yr ⁻¹	mm yr ⁻¹	% yr ⁻¹
summer	SwissSMEX	-1.4	-0.9	-0.7	-0.5	-1.2	-0.7	-1.1	-0.6
	ERA5-Land	-1.2	-0.7	-1.3	-0.7	-1.1	-0.7	-1.0	-0.6
	C3S SM PASSIVE	-1.1	-0.7 ^(*)	-0.8	-0.5	-0.9	-0.6 ^(*)	-0.8	-0.6
summer half-year	SwissSMEX	-0.6	-0.4	-0.2	-0.1	-0.6	-0.3	-0.5	-0.3
	ERA5-Land	-0.5	-0.3	-0.7	-0.4	-0.6	-0.3	-0.6	-0.3
	C3S SM PASSIVE	-0.9 ^(*)	-0.5 ^(*)	-0.5	-0.3	-0.7	-0.5	-0.7	-0.5

^(*) denotes p < 0.05

305 3.2 Comparison with alternate long-term reanalysis and remote sensing soil moisture

The agreement between the different station combinations regarding the temporal evolution of the Swiss Plateau median soil moisture is also evident in Fig. 8, which, in addition to SwissSMEX, includes the ERA5-Land and the C3S SM PASSIVE soil moisture. The three independent data sources show good agreement in the inter-annual variations of the summer and summer-half-year soil moisture anomalies over the period 2010–2025, with correlations of 0.80 or higher between the gridded products and SwissSMEX based on the median of the best coverage stations (see also Fig. 7 and Fig. S5, values within the red solid rectangles). This holds for both the absolute soil moisture anomalies, as well as the percentage anomalies (Fig. S6). The RMSD from SwissSMEX amounts to 6.9 mm for the absolute anomalies (or 4.0 % for the percentage anomalies) for both C3S SM PASSIVE and ERA5-Land in summer (Fig. 8, Fig. S6). For the summer half-year, these values are lower due to the lower year-to-year variability (5.5 mm or 3.1 % for C3S SM PASSIVE, and 5.3 mm or 3.0 % for ERA5-Land). The pairwise correlations between the three data sources are also significant for the other station combinations, both for summer (Fig. 7) and the summer-half-year (Fig. S5).

315

Deleted: 7

Deleted: 0

Deleted: 8

Deleted: 4

Deleted: 2

Deleted: 1

Deleted: 6

Deleted: 2

Deleted: 0.9

Deleted: 5

Deleted: ^(*)

Deleted: ·

Deleted: ⁽ⁱ⁾

Deleted: 5⁽ⁱ⁾

Deleted: 7

Deleted: 3

Deleted: 7

Deleted: 4

Deleted: 4

Deleted: 2

Deleted: 5

Deleted: 5

Deleted: ·

Deleted: 6

Deleted: 8^(*)

Deleted: ⁽ⁱ⁾

Deleted: ⁽ⁱ⁾

Deleted: 4⁽ⁱ⁾

Deleted: ⁽ⁱ⁾ denotes p < 0.1, and

Deleted: 79






Figure 8 Evolution of (a) summer and (b) summer-half-year soil moisture anomalies based on SwissSMEX, ERA5-Land and C3S SM PASSIVE. Anomalies are expressed as absolute deviations from the summer or summer-half-year mean IWC over the period 2010–2025, respectively (see Fig. S6 for corresponding anomalies as percentage deviations from this mean). Shown are the median time series based on the best coverage stations (solid lines), as well as the ones based on the alternative station combinations presented in Section 2.3 (dotted lines). For the gridded products, the nearest grid cells to the stations are considered as the basis for the median. Straight lines show the Theil-Sen trend slopes for the period 2010–2025 (based on the best coverage stations time series), which are also indicated to the right of the plots (solid lines indicate a significant trend with $p < 0.05$; dashed lines indicate a non-significant trend). Also noted are the Pearson correlation r and RMSD of the gridded products with respect to SwissSMEX, and known drought summers are indicated by vertical lines.

The Theil-Sen trend slopes for the median of the best coverage stations amount to -1.4 mm yr^{-1} (or $-0.9\% \text{ yr}^{-1}$) for SwissSMEX, to -1.2 mm yr^{-1} (or $-0.7\% \text{ yr}^{-1}$) for ERA5-Land, and to -1.1 mm yr^{-1} (or $-0.7\% \text{ yr}^{-1}$) for C3S SM PASSIVE for summer (Fig. 8 and Fig. S6, Table 2). The summer-half-year trends based on the best coverage stations are about half of the summer values for SwissSMEX (-0.6 mm yr^{-1}) and ERA5-Land (-0.5 mm yr^{-1}), whereas C3S SM PASSIVE shows similar trends for summer and the summer half-year (-0.9 mm yr^{-1} for the latter). Over the period 2010–2025, only the C3S SM PASSIVE summer-half-year trends are significant for the absolute anomalies. The same holds for the percentage anomalies of this product for both summer and the summer half-year. This is because the product exhibits lower year-to-year variability and less pronounced negative summer and summer-half-year anomalies compared with SwissSMEX and ERA5-Land (Fig. 8, Fig. S6). The drying tendencies for all station combinations are listed in Table 2.

ERA5-Land and C3S SM PASSIVE also show reasonable agreement in the inter-annual variations of Swiss Plateau soil moisture since 1991 (Fig. A1, Fig. S6), with correlations of 0.68 for summer and 0.63 for summer half-year ($p < 0.05$ for both) based on the median of the best coverage stations. Even though the sign of the anomalies does not agree in some cases, the dry anomalies of 2003, 2015, 2018, 2022 and 2023 are apparent in both products, with ERA5-Land displaying more negative anomalies compared to C3S SM PASSIVE. This is consistent with previously reported differences in the drought detection capacity of these gridded products (Hirschi et al., 2025b). As in the period 2010–2025, the different station combinations also show largely consistent temporal evolutions since 1991. The long-term trend of ERA5-Land becomes significant when considering summer-half-year anomalies since 1980 (Figs. A1 and S6, see also Appendix B), in agreement with the analysis



Deleted:



Deleted:

Deleted: 0

Deleted: 8

Deleted: 1

Deleted: 6

Deleted: 7

Deleted: 7

Deleted: 1

of S22. Similarly, as in the period 2010–2025, C3S SM PASSIVE trends are significant when considering summer-half-year anomalies since 1991. This temporally extended perspective highlights that the recent evolution of SwissSMEX aligns well with the longer-term evolution of soil moisture from the two gridded products, all of which indicate an increase in summer soil drying in recent years.

Deleted: summer and

3.3 Differences to S22

Compared with S22, the IWC of SwissSMEX presented here relies more heavily on TDR sensors, which have proven to be more robust in operation, and which have been partly renewed or additionally installed in 2022 (see Section 2.1.1). Despite the applied site-specific calibration of the 10HS sensors, the temporal variability of the TDR sensors tends to be partly larger than that of the 10HS sensors. This may explain part of the larger temporal variability and stronger anomalies observed in the presented SwissSMEX time series of the Swiss Plateau compared with S22 (cf. Fig. S3 therein). In particular, the IWC at the station CHN was based on 10HS sensors in S22, whereas it is now based on TDR sensors and extended with 10 and 30 cm TDR-based integration (see Fig. 5). Similarly, the station BER now includes a 10 cm TDR sensor in the integration, replacing the 10HS sensor at the same depth, which stopped working in 2021 (see Table S1 for an overview of the sensors used). Given the limited number of stations in S22, these changes in sensor configuration result in the observed differences in the temporal variability of the SwissSMEX IWC time series. Consequently, the agreement between SwissSMEX and ERA5-Land improves considerably compared with S22, in which SwissSMEX showed lower year-to-year variations in IWC than ERA5-Land.

4 Data availability

The hourly and daily IWC time series of SwissSMEX (Swiss Plateau stations), i.e. the historical IWC_{orig} based on the original sensor configuration, the TDR-only based IWC_{TDRonly} and IWC_{TDRonly_corrected}, and the merged IWC will be made available from the ETH Research Collection at <https://doi.org/10.3929/ethz-b-000743711> (Hirschi et al., 2025a) (for the review, data is available at <https://polybox.ethz.ch/index.php/s/Xt4Zr9sN5PePyKM>). Additionally, the VWC of the individual sensors and depths is available from the same location. ERA5-Land is available from the CDS at <https://doi.org/10.24381/cds.68d2bb30> (Muñoz-Sabater et al., 2021). The C3S SM PASSIVE gap-filled root-zone soil moisture is available at <https://doi.org/10.48436/9j8ad-z2q11> (Preimesberger et al., 2026).

5 Summary and conclusion

We present a curated and comprehensive set of in situ soil moisture time series from the SwissSMEX stations on the Swiss Plateau, covering the period 2010–2025. SwissSMEX provides data on volumetric soil water content (VWC) at various depths in the soil profile, which are used to calculate integrated soil water content (IWC) down to 50 cm depth as an indicator of root-zone water. We document the steps that have recently been taken to secure the network and its long-term IWC time series,

Deleted: ed

Deleted: ed

460 which have been affected by an increasing number of sensor failures in recent years. In summer 2022, defective TDR sensors
at 10 and 30 cm depth were replaced, and additional ones were installed where they had not previously been present in the
profile at these depths. These new TDR measurements are used as an alternative for the IWC calculation when coverage with
the original sensors is lacking. We merge the original IWC with the IWC based on the new TDR sensors by correcting the
mean offset and the difference in standard deviation of the latter relative to the former to account for the difference in the
representative depth.

Deleted: d

465 As a validation use case for the network, we demonstrate the utility of SwissSMEX and its secured IWC time series for
analysing documented summer drying trends on the Swiss Plateau. We focus on summer and summer-half-year IWC anomalies
and trends therein, and compare these with ERA5-Land reanalysis and C3S SM PASSIVE remote-sensing soil moisture. We
find good agreement in the temporal evolution of SwissSMEX in situ soil moisture across different combinations of stations
on the Swiss Plateau. All station combinations share a drying tendency over the period 2010–2025. Comparing the in situ time
series with ERA5-Land and C3S SM PASSIVE soil moisture also reveals a good agreement between the three independent
data sources, with correlations of 0.80 or higher for the median time series of the stations with best temporal coverage. The
non-significant drying over the period 2010–2025, based on the median of these stations, amounts to -1.4 mm yr^{-1} for
SwissSMEX, -1.2 mm yr^{-1} for ERA5-Land, and -1.1 mm yr^{-1} for C3S SM PASSIVE for summer.

Deleted: d

Deleted: ed

Deleted: d

Deleted: ed

Deleted: 79

Deleted: and

Deleted: 1

475 While the SwissSMEX network suggests that summer soil drying has increased in recent years, the 16 years of in situ data
currently available are not yet sufficient to robustly estimate a significant trend. The projected increases in air temperature and
evaporative demand, combined with a decline in summer precipitation, are expected to lead to further summer drying in
Switzerland in the coming years (CH2018, 2018; MeteoSwiss and ETH Zurich, 2025). In this context, the continuation of the
480 SwissSMEX measurements and the ongoing efforts at the federal level to establish a national soil moisture measurement
network are crucial for ensuring a seamless continuation of soil moisture monitoring in Switzerland.





Figure A1 As Fig. 8 but extended with the long-term evolution of (a) summer and (b) summer-half-year soil moisture anomalies based on ERA5-Land (starting in 1980) and C3S SM PASSIVE (starting in 1991). For both long-term products, the Theil-Sen trend slopes for the periods 1980–2025 and 1991–2025, respectively, are indicated by straight lines (solid lines indicate a significant trend with $p < 0.05$; dashed lines indicate no significant trend).

Appendix B

Temporal robustness of trend


To investigate the impact of the shortness of the SwissSMEX time series on the significance of the trend, Fig. B1 shows the p-value of the Mann-Kendall trend test based on ERA5-Land soil moisture as a function of the starting year used to calculate the trend (i.e. the length of the time series). A starting year of 2010 corresponds to the 2010–2025 trend (16 years of data, i.e. the SwissSMEX period), while a starting year of 1980 corresponds to the 1980–2025 trend (46 years of data). In addition to the summer and summer-half-year trends, trends based on annual anomalies are also considered

Due to the larger inter-annual variability, summer trends tend to be overall less significant (i.e. have higher p-values) than trends based on annual and summer-half-year soil moisture anomalies, even though the absolute trend magnitude is larger for summer than for the summer half-year (see Table 2). For the short period 2010–2025, trends in SwissSMEX and ERA5-Land soil moisture anomalies are not significant (circles in Fig. B1, see also Table 2). Trends in summer-half-year soil moisture anomalies become significant for certain periods when extending the starting year of the trend test calculation. Particularly, summer-half-year trends that start before 1983 become significant, consistent with the results of S22. The same partly applies to annual and summer-half-year trends from 1993 to 1996. Trends in summer anomalies are, however, non-significant ($p > 0.05$) for all starting years, and the corresponding p-values vary considerably over time. Higher p-values are observed when the starting year has a comparably dry summer compared to surrounding years (e.g. as 2003, 1989).



Deleted: starting
Deleted: between
Deleted: and

520 This analysis shows that the significance of the trend depends on both the time frame of the anomalies (annual, summer half-year, or summer) and the period used for the trend calculation. Thus, while the SwissSMEX network indicates that soils have become drier in recent years, the 16 years of in situ data currently available are not yet sufficient to robustly estimate significant trends. Extending the period used to calculate the trend yields significant results, particularly for summer-half-year soil moisture anomalies, although the significance of the trend may still be impacted by anomalous soil moisture conditions in the starting year.



530 **Figure B1** Significance of the Theil-Sen trend slope (p-value of the Mann-Kendall trend test) based on ERA5-Land soil moisture as a function of the starting year for calculating the trend (i.e. the length of the time series). The trends are calculated based on the median time series of the absolute soil moisture anomalies from the best coverage stations (solid lines in Fig. 8). In addition to the summer and summer-half-year trends, annual trends are also considered. Starting year 2010 refers to the 2010–2025 trend (or 16 years of data), starting year 1980 refers to the 1980–2025 trend (46 years of data). The p-values for SwissSMEX based on the 16-year time series are shown as circles at starting year 2010. The horizontal lines indicate the $p=0.05$ and $p=0.1$ thresholds.

Author contribution


MH, DM, DLS and SIS: Conceptualization; MH, DM: Formal analysis, Investigation, Methodology, Visualisation; MH, DM, WP: Data curation; MH: Writing – original draft preparation; all authors: Writing – review & editing

535 **Competing interests**

The authors declare that they have no conflict of interest.

Acknowledgements

We acknowledge partial financial support from GCOS Switzerland for the update of the SwissSMEX network in 2022. We thank Michael Rösch for his technical support and maintenance of the measurement sites.



Deleted:

Deleted: aining

References

Albergel, C., Rüdiger, C., Pellarin, T., Calvet, J. C., Fritz, N., Froissard, F., Suquia, D., Petitpa, A., Piguet, B., and Martin, E.: From near-surface to root-zone soil moisture using an exponential filter: an assessment of the method based on in-situ observations and model simulations, *Hydrology and Earth System Sciences*, 12, 1323-1337, 10.5194/hess-12-1323-2008, 2008.

BAFU: Hitze und Trockenheit im Sommer 2015: Auswirkungen auf Mensch und Umwelt, Bundesamt für Umwelt BAFU, Berne, Umwelt-Zustand 1629, 108 pp., 2016.

BAFU: Drought: monitoring, forecasting and warning: <https://www.bafu.admin.ch/bafu/en/home/topics/water/extreme-events/drought-monitoring-forecasting-and-warning.html>, last access: 31-03-2025, 2024.

Balsamo, G., Viterbo, P., Beljaars, A., van den Hurk, B., Hirschi, M., Betts, A. K., and Scipal, K.: A Revised Hydrology for the ECMWF Model: Verification from Field Site to Terrestrial Water Storage and Impact in the Integrated Forecast System, *Journal of Hydrometeorology*, 10, 623-643, 10.1175/2008jhm1068.1, 2009.

Brouwer, C., Prins, K., and Heibloem, M.: Irrigation Water Management: Irrigation Scheduling, FAO, <https://www.fao.org/4/t7202e/t7202e00.htm>, 1989.

Brunner, M. I., Liechti, K., and Zappa, M.: Extremeness of recent drought events in Switzerland: dependence on variable and return period choice, *Nat Hazard Earth Sys*, 19, 2311-2323, 10.5194/nhess-19-2311-2019, 2019.

C3S: ERA5-Land monthly averaged data from 1950 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [dataset], 10.24381/cds.68d2bb30, 2022.

C3S and WMO: Soil moisture, in: European State of the Climate 2024, 39-41, 10.24381/14j9-s541, 2025.

Calanca, P.: Climate change and drought occurrence in the Alpine region: How severe are becoming the extremes?, *Global Planet Change*, 57, 151-160, 10.1016/j.gloplacha.2006.11.001, 2007.

CH2018: CH2018 – Climate Scenarios for Switzerland, National Centre for Climate Services, Zürich, Technical Report, 271 pp., <https://www.nccs.admin.ch/nccs/de/home/klimawandel-und-auswirkungen/schweizer-klimaszenarien/technical-report.html>, 2018.

Domínguez-Niño, J. M., Bogen, H. R., Huisman, J. A., Schilling, B., and Casadesús, J.: On the Accuracy of Factory-Calibrated Low-Cost Soil Water Content Sensors, *Sensors*, 19, ARTN 3101 10.3390/s19143101, 2019.

Dorigo, W., Preimesberger, W., Lems, J., Frederikse, T., and Dostalova, A.: Soil moisture gridded data from 1978 to present (v202505) [dataset], 10.24381/cds.d7782f18, 2025.

Dorigo, W., Preimesberger, W. H., S., Van der Schalie, R., De Jeu, R., Kidd, R., Rodriguez-Fernandez, N., Hirschi, M., Stradiotti, P., Frederikse, T., Gruber, A., and Duchemin, D.: ESA Soil Moisture Climate Change Initiative (Soil_Moisture_cci): PASSIVE product, Version 09.1, NERC EDS Centre for Environmental Data Analysis [dataset], 10.5285/ca55ac11fc814b0d95e68a34a10539c1, 2024.

Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M., Forkel, M., Gruber, A., Haas, E., Hamer, P. D., Hirschi, M., Ikonen, J., de Jeu, R., Kidd, R., Lahoz, W., Liu, Y. Y., Miralles, D., Mistelbauer, T., Nicolai-Shaw, N., Parinussa, R., Pratola, C., Reimer, C., van der Schalie, R., Seneviratne, S. I., Smolander, T., and Lecomte, P.: ESA CCI Soil Moisture for improved Earth system understanding: State-of-the art and future directions, *Remote Sensing of Environment*, 203, 185-215, 10.1016/j.rse.2017.07.001, 2017.

Gruber, A., Scanlon, T., van der Schalie, R., Wagner, W., and Dorigo, W.: Evolution of the ESA CCI Soil Moisture climate data records and their underlying merging methodology, *Earth System Science Data*, 11, 717-739, 10.5194/essd-11-717-2019, 2019.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 1999-2049, 10.1002/qj.3803, 2020.

Hirschi, M., Michel, D., Mittelbach, H., Rösch, M., and Seneviratne, S. I.: SwissSMEX soil moisture data (up to 2024), ETH Zurich [dataset], 10.3929/ethz-b-000743711, 2025a.

Field Code Changed

Field Code Changed

Field Code Changed

- Hirschi, M., Stradiotti, P., Crezee, B., Dorigo, W., and Seneviratne, S. I.: Potential of long-term satellite observations and reanalysis products for characterising soil drying: trends and drought events, *Hydrol. Earth Syst. Sci.*, 29, 397-425, 10.5194/hess-29-397-2025, 2025b.
- 595 Hupet, F., Bogaert, P., and Vanclooster, M.: Quantifying the local-scale uncertainty of estimated actual evapotranspiration, *Hydrological Processes*, 18, 3415-3434, 10.1002/hyp.1504, 2004.
- Kendall, M. G.: *Rank Correlation Methods*, Griffin, London, 1975.
- Mann, H. B.: Nonparametric Tests against Trend, *Econometrica*, 13, 245-259, 10.2307/1907187, 1945.
- MeteoSchweiz: Klimabulletin Jahr 2011, MeteoSwiss, Zürich, 12 pp., <https://www.meteoschweiz.admin.ch/service-und-publikationen/publikationen/berichte-und-bulletins/2012/klimabulletin-jahr-2011.html>, 2012.
- 600 MeteoSchweiz: Hitze und Trockenheit im Sommerhalbjahr 2018 - eine klimatologische Übersicht, MeteoSwiss, Zürich, Fachbericht MeteoSchweiz 272, 38 pp., <https://www.meteoschweiz.admin.ch/service-und-publikationen/publikationen/berichte-und-bulletins/2018/hitze-und-trockenheit-im-sommerhalbjahr-2018-eine-klimatologische-uebersicht.html>, 2018.
- 605 MeteoSwiss and ETH Zurich: Climate CH2025 - Scientific Report, Federal Office of Meteorology and Climatology MeteoSwiss, Zurich, 10.18751/climate/scenarios/ch2025/sr/1.0/, 2025.
- Michel, D., Hirschi, M., Buri, R., and Seneviratne, S. I.: Updating the SwissSMEX soil moisture monitoring network infrastructure to secure the continuity of the long-term data series and ensure the data quality, Final report GCOS-CH supported activities, 10.3929/ethz-b-000713939, 2022a.
- 610 Michel, D., Hirschi, M., Buri, R., Wittneben, B., and Seneviratne, S. I.: Enhancing cross-disciplinary process understanding and scientific exploitation of soil moisture and land evaporation ECV measurements in Switzerland (SMiLE-ECV-CH), Final report GCOS-CH supported activities, 10.3929/ethz-b-000713937, 2022b.
- Mittelbach, H.: Soil moisture in Switzerland: Analyses from the Swiss Soil Moisture Experiment, ETH Zurich, 133 pp., 10.3929/ethz-a-007142621, 2011.
- 615 Mittelbach, H. and Seneviratne, S. I.: A new perspective on the spatio-temporal variability of soil moisture: temporal dynamics versus time-invariant contributions, *Hydrology and Earth System Sciences*, 16, 2169--2179, 10.5194/hess-16-2169-2012, 2012.
- Mittelbach, H., Lehner, I., and Seneviratne, S. I.: Comparison of four soil moisture sensor types under field conditions in Switzerland, *Journal of Hydrology*, 430--431, 39--49, 10.1016/j.jhydrol.2012.01.041, 2012.
- 620 Mittelbach, H., Casini, F., Lehner, I., Teuling, A. J., and Seneviratne, S. I.: Soil moisture monitoring for climate research: Evaluation of a low-cost sensor in the framework of the Swiss Soil Moisture Experiment (SwissSMEX) campaign, *Journal of Geophysical Research*, 116, D05111, 10.1029/2010JD014907, 2011.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349-4383, 10.5194/essd-13-4349-2021, 2021.
- 625 Orth, R., Zscheischler, J., and Seneviratne, S. I.: Record dry summer in 2015 challenges precipitation projections in Central Europe, *Sci Rep-Uk*, 6, ARTN 28334 10.1038/srep28334, 2016.
- 630 Pasik, A., Gruber, A., Preimesberger, W., De Santis, D., and Dorigo, W.: Uncertainty estimation for a new exponential-filter-based long-term root-zone soil moisture dataset from Copernicus Climate Change Service (C3S) surface observations, *Geoscientific Model Development*, 16, 4957-4976, 10.5194/gmd-16-4957-2023, 2023.
- Pohlert, T.: trend: Non-Parametric Trend Tests and Change-Point Detection, 10.32614/CRAN.package.trend, 2020.
- Preimesberger, W., Stradiotti, P., and Dorigo, W.: ESA CCI Soil Moisture GAPPILLED: An independent global gap-free satellite climate data record with uncertainty estimates, *Earth Syst. Sci. Data Discuss.*, 2025, 1-40, 10.5194/essd-2024-610, 2025.
- 635 Preimesberger, W., Lems, J., Hirschi, M., and Dorigo, W. A.: C3S SM PASSIVE Daily Gap-filled Root-Zone Soil Moisture from merged multi-satellite observations (Version v2), TU Wien [dataset], 10.48436/9j8ad-z2q11, 2026.
- Scherrer, S. C., Hirschi, M., Spirig, C., Maurer, F., and Kotlarski, S.: Trends and drivers of recent summer drying in Switzerland, *Environmental Research Communications*, 4, 10.1088/2515-7620/ac4fb9, 2022.

- 640 Schumacher, D. L., Zachariah, M., Otto, F., Barnes, C., Philip, S., Kew, S., Vahlberg, M., Singh, R., Heinrich, D., Arrighi, J.,
van Aalst, M., Hauser, M., Hirschi, M., Bessenbacher, V., Gudmundsson, L., Beaudoin, H. K., Rodell, M., Li, S. H., Yang,
W. C., Vecchi, G. A., Harrington, L. J., Lehner, F., Balsamo, G., and Seneviratne, S. I.: Detecting the human fingerprint in the
summer 2022 western-central European soil drought, *Earth System Dynamics*, 15, 131-154, 10.5194/esd-15-131-2024, 2024.
- 645 Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau, *Journal of the American Statistical Association*,
63, 1379-1389, 10.2307/2285891, 1968.
- Theil, H.: A rank-invariant method of linear and polynomial regression analysis, *Proc Konink Nederl. Akad. Wetensch. Ser.
A Math. Sci.*, 53, 386-392 (part I), 521-525 (part II), 1397-1412 (part III), 1950.
- Tuszynski, J.: caTools: Tools: Moving Window Statistics, GIF, Base64, ROC AUC, etc, 10.32614/CRAN.package.caTools,
2021.
- 650 Wagner, W., Lemoine, G., and Rott, H.: A method for estimating soil moisture from ERS scatterometer and soil data, *Remote
Sensing of Environment*, 70, 191-207, Doi 10.1016/S0034-4257(99)00036-X, 1999.