



Multi-year observations of BVOC and ozone: concentrations and fluxes measured above and below the canopy in a mixed temperate forest

Clément Dumont¹, Bert W. D. Verreyken^{1,2}, Niels Schoon², Benjamin Bergmans⁴, Crist Amelynck^{2,3}, and Bernard Heinesch¹

¹Gembloux Agro-Bio Tech, University of Liège, Belgium

²Belgian Institute for Space Aeronomy, Belgium

³Department of Chemistry, Ghent University, Belgium

⁴Institut Scientifique de Service Public, Liège, Belgium

Correspondence: Clément Dumont (clement.dumont@uliege.be)

Abstract.

Volatile organic compounds (VOCs) and ozone (O₃) are key constituents of tropospheric chemistry, affecting both air quality and climate. Forests are major emitters of biogenic VOCs (BVOCs), yet large uncertainties remain regarding the diversity of exchanged compounds, the drivers of their bidirectional fluxes, and their in-canopy chemistry. Long-term and comprehensive in situ datasets remain scarce, limiting our understanding of these complex processes. We conducted a three-year field campaign (2022—2024) at the Integrated Carbon Observation System forest station of Vielsalm (BE-Vie), combining vertical concentration profile and eddy covariance flux measurements above and below the canopy. Using a PTR-ToF-MS and a dedicated open-source processing pipeline, we identified 51 significantly exchanged VOCs. The vertical and diurnal gradients of the mixing ratios reflected the interplay between emission, deposition, chemistry, and transport. Combined with a profile of turbulence statistics, these observations offer an opportunity to investigate their behaviour within the canopy. The forest acted as a net VOC source in summer ($\sim 1.25 \mu\text{g m}^{-2} \text{s}^{-1}$), while deposition dominated in autumn. Many oxygenated VOCs displayed bidirectional exchange. Monoterpenes, isoprene, and methanol were the most abundant flux contributors, but 15—30 (30—43) compounds were needed to account for 90% of total emissions (depositions), depending on the season. Below-canopy BVOC and O₃ fluxes reached $\sim 10\%$ of above-canopy ones, with proportionally enhanced below-canopy ozone uptake at night. This study provides one of the most detailed long-term datasets of VOC and O₃ exchange in a temperate forest and serves as a key reference for improving process-based models of biogenic, physical, and chemical exchange in forest ecosystems.

1 Introduction

Volatile organic compounds (VOCs) form a heterogeneous group of reactive trace gases that play a key role in atmospheric chemistry. Together with nitrogen oxides (NO_x), they contribute to both the formation and destruction of tropospheric ozone (Stockwell and Forkel, 2002), influencing regional ozone pollution. By reacting with hydroxyl and peroxy radicals, VOCs also affect the atmospheric oxidizing capacity and the lifetime of methane (Yoon et al., 2025), the second most important



radiatively active gas after carbon dioxide (Canadell et al., 2021). In addition, VOCs are precursors in the formation and growth of secondary organic aerosols (SOA) (Mahilang et al., 2021), thereby degrading air quality, impacting human health, and influencing climate through interactions with radiation and serving as cloud condensation nuclei.

25 Each year, terrestrial ecosystems emit over 1000 Tg of organic carbon in the form of biogenic VOCs (BVOCs), excluding methane (Guenther et al., 2012). This exceeds global emissions of methane ($\sim 550 \text{ Tg yr}^{-1}$) (Saunio et al., 2016) and anthropogenic VOCs ($\sim 200 \text{ Tg yr}^{-1}$) (Huang et al., 2017). Forests are the dominant BVOC emitters among terrestrial ecosystems (Isidorov et al., 2022, and references therein) and have therefore long been a major focus of atmospheric research.

Ozone (O_3), in addition to its role in radiative forcing as a potent greenhouse gas (Rowlinson et al., 2020), is toxic to both humans and plants. In terrestrial ecosystems, it undergoes various dry deposition pathways—including stomatal and non-stomatal processes such as uptake by leaf cuticles, soil, snow, or water surfaces (Clifton et al., 2020). Ozone deposition contributes significantly to reducing tropospheric ozone levels. However, when ozone enters leaves through stomata, it generates reactive oxygen species (ROS), which can damage plant cells, accelerate senescence, and reduce carbon assimilation via photosynthesis (Fiscus et al., 2005).

35 Despite decades of research into ozone uptake partitioning (Padro, 1996; Cieslik, 1998; Lamaud et al., 2002; Zhang et al., 2002; Mikkelsen et al., 2004; Fares et al., 2014; Horváth et al., 2017; Zhou et al., 2017b; Finco et al., 2018; Gerosa et al., 2022b), significant uncertainties remain. One such uncertainty lies in the reactions of ozone with BVOCs, which constitute an increasingly recognized atmospheric sink. This process, however, remains highly variable and difficult to quantify (Kurpius and Goldstein, 2003; Goldstein et al., 2004; Wolfe et al., 2011; Vermeuel et al., 2021).

40 Unlike ozone, BVOCs can undergo both emission and deposition. Early research primarily focused on isoprene and monoterpenes—the two most widely emitted isoprenoids globally (Guenther et al., 2012)—whose fluxes were long considered exclusively upward, driven by light and temperature (Guenther et al., 1995). Later studies extended global emission inventories to include oxygenated VOCs (OVOCs, e.g., methanol, ethanol, acetone, acetaldehyde, formaldehyde, formic and acetic acids), sesquiterpenes, and isoprenoid oxidation products (Niinemets et al., 2014; Guenther et al., 2012), but largely continued to focus on emissions rather than deposition. Yet, in situ studies have highlighted non-negligible deposition fluxes of BVOCs (Karl et al., 2010; Bamberger et al., 2011; Jardine et al., 2011; Ruuskanen et al., 2011; Laffineur et al., 2012; Park et al., 2013a; Nguyen et al., 2015; Wohlfahrt et al., 2015; Zhou et al., 2017b), which are now recognized as significant in global budgets. For instance, Safieddine et al. (2017) estimated that 460 Tg C yr^{-1} of reactive organic carbon is removed from the atmosphere via physical deposition. A regional simulation over the U.S. further showed that dry and wet deposition of organic vapours accounted for 50 60–75% of the removal of tropospheric SOA burden in summer 2010 (Hodzic et al., 2014).

It is now well established that many BVOCs undergo bidirectional exchange. Yet, emissions and depositions are generally modelled separately in chemical transport models (CTMs), despite being driven by similar environmental variables (Forkel et al., 2015). Emissions are typically simulated using the MEGAN model, which combines species-specific emission potentials with responses to light, temperature, CO_2 inhibition, and leaf age (Guenther et al., 2012). Deposition, in contrast, is usually 55 calculated as the product of ambient concentration and deposition velocity, using an Ohm's law analogy introduced by Wesely (1989). This approach depends on deposition velocity estimates, which are still scarce, uncertain, and rarely incorporate



seasonal variation (Niinemets et al., 2014). Other frameworks based on compensation points require precise knowledge of intercellular VOC concentrations and stomatal behaviour, and are difficult to upscale to the ecosystem level (Niinemets et al., 2014).

60 An additional process often overlooked in surface exchange models is in-canopy chemistry, which can significantly alter fluxes of highly reactive compounds such as sesquiterpenes (Forkel et al., 2015). VOCs can be lost between their point of release and the atmosphere above the canopy (Rinne et al., 2007; Zhou et al., 2017b), affecting CTM predictions of atmospheric concentrations in ways that are not yet adequately represented (Vermeuel et al., 2024; Link et al., 2024). Accounting for these reactions also requires identifying the vertical distribution of sources and sinks within the canopy. Although several studies have
 65 applied inverse modelling to approach this question, few ecosystems and tracers have been characterized to date (Karl et al., 2004b; Tiwary et al., 2007; Wada et al., 2020; Petersen et al., 2023). Overall, there is a growing call for a unified framework to model bidirectional BVOC exchange, integrating altogether dry and wet deposition, direct emissions, and in-canopy chemistry (Pleim and Ran, 2011; Niinemets et al., 2014; Forkel et al., 2015).

While emission models such as MEGAN have become increasingly sophisticated, they still represent only a fraction of
 70 the chemical diversity of atmospheric VOCs. For example, MEGAN2.1 includes 147 compounds (Guenther et al., 2012), whereas most CTMs simulate only 30–45 species (Millet et al., 2018). This limited representation contrasts starkly with field observations: Goldstein and Galbally (2007) estimated that between 10^4 and 10^5 distinct organic compounds have been detected in the atmosphere. Flux measurements have also revealed remarkable chemical diversity, with over 500 VOC-related ions observed over an orange orchard (Park et al., 2013a) and 377 over a temperate forest (Millet et al., 2018).

75 Improving VOC exchange parametrizations requires in situ, ecosystem-scale flux measurements across a diversity of ecosystems and environmental conditions. Traditionally, such measurements relied on disjunct eddy covariance with PTR-Quad-MS instruments (Müller et al., 2010; Laffineur et al., 2011, 2012, 2013; Rinne et al., 2007; Yang et al., 2013; Seco et al., 2015; Bachy et al., 2016, 2018; Seco et al., 2017). This method allowed the monitoring of a limited number of pre-selected compounds and could not resolve isobaric interferences. As a consequence, this technique was mostly used to investigate well-known BVOCs
 80 (Schallhart et al., 2016). More recently, the advent of PTR-ToF-MS coupled with eddy covariance enabled the simultaneous detection of fluxes for hundreds of VOCs with improved sensitivity and mass resolution, allowing isobar separation (Ruuskanen et al., 2011; Kaser et al., 2013; Park et al., 2013a, b; Brilli et al., 2014, 2016; Schallhart et al., 2016; Jurán et al., 2017; Millet et al., 2018; Schallhart et al., 2018; Jensen et al., 2018; Fischer et al., 2021; Manco et al., 2021; Loubet et al., 2022; Petersen et al., 2023; Vermeuel et al., 2023b). However, many studies remained compound-targeted, and the technical complexity of
 85 PTR-ToF-MS operation and data processing limited their temporal coverage (Rinne et al., 2016). As a result, long-term flux measurements across seasons and under variable conditions remain scarce.

In this study, we revisit a temperate mixed forest in southern Belgium (Vielsalm station, BE-Vie), more than ten years after the first PTR-Quad-MS flux measurements (Laffineur et al., 2011, 2012, 2013), with the aim of expanding the existing dataset. Using a PTR-ToF-MS and ozone analysers, we conducted a three-year measurement campaign to obtain:

90 (i) 1-minute concentrations of BVOCs and O_3 above the canopy and at ground level,



- (ii) 30-minute fluxes of BVOCs and O₃ at the same two levels,
- (iii) vertical profiles of BVOC and O₃ concentrations at seven levels from the trunk space to above the canopy, and
- (iv) vertical profiles of sonic anemometers at eight levels to characterize in-canopy turbulence.

We provide a detailed description of the VOC data processing workflow—including newly developed and openly available processing tools—from raw PTR-ToF-MS signal treatment to compound identification, quantification, flux calculation, and data flagging, following a non-targeted and comprehensive approach. Ozone flux data processing is also described in detail, and methodological issues raised in earlier studies (Muller et al., 2010; Zhu et al., 2015) are discussed.

To our knowledge, this study presents the longest combined VOC and ozone flux dataset publicly available to date, spanning three years and covering seasonal transitions from spring to summer and summer to autumn. Combined with additional open-access measurements at the site (extensive meteorological and phenological characterization, BTEX, HAPs, NO_x, fine particles, and aerosols), this structured database offers a unique opportunity to address key knowledge gaps. These include quantifying the magnitude and diversity of BVOC exchange in temperate forests, their influence on ozone levels, the dynamics of bidirectional fluxes, the spatial distribution of sources and sinks, and the role of in-canopy chemistry. As a result, it provides a strong basis for improving current parametrizations in surface exchange models and CTMs, and to advancing our understanding of VOC and ozone interactions in the Earth system.

2 Material & Methods

2.1 Study site

The study site (BE-Vie) is a mature mixed forest ecosystem, located in Vielsalm in the Belgian Ardenne region (50°18'N, 6°00'E; altitude 470 m a.s.l.). This site is part of several gas measurement networks: ICOS (class 2 ecosystem station) (<https://www.icos-cp.eu/>), ACTRIS (for particle and gas in situ) (<https://www.actris.eu/>), and EMEP (level 1 station) (<https://www.emep.int/>). Its climate is temperate maritime and the 50-100 cm deep soil is classified as a dystric cambisol. The vegetation within the target area (that is, the area that contributes the most to fluxes measured at the flux tower) is a mixture of coniferous species, mainly Norway spruce (*Picea abies* [L.] Karst.), Douglas fir (*Pseudotsuga menziesii* [Mirb.] Franco) and Silver fir (*Abies alba* Miller); and deciduous species, mainly beeches (*Fagus sylvatica* L.). The understory vegetation is very sparse or absent in some areas. Fig. 1 gives a distribution of dominant tree species in most of the target area. This classification is based on multispectral images acquired with a drone in 2018 (Lanssens, 2019). The dominant winds blow from the south-west (sector dominated by *Fagus sylvatica*) and the north-east (sector dominated by *Picea abies* and *Pseudotsuga menziesii*). The footprint climatology was obtained using the model by (Kljun et al., 2015) and is shown in Fig. 1.

The study site is located in a rural area with low anthropogenic activity, except for a sawmill 3 km south-west of the flux tower and two small industrial hubs around 7 km north-east and south-east. ICOS-related measurements are carried out on the 51 m high flux tower and its close surroundings. Another measurement station, operated by the Walloon Scientific Institute




Figure 1. Map of the study site with details about (i) dominant tree distribution, (ii) footprint climatology (50, 70 and 80% isopleths) and (iii) research infrastructures (flux tower and ISSeP station). The grey line represents the border of the ICOS target area.

for Public Service (ISSeP)—the regional environmental authority—is located in the same forest, approximately 250 m from the flux tower. Measurements carried out at the ISSeP station are detailed in Sect. 2.5.1. More information about the study site are available in Aubinet et al. (2001, 2018). Note that all times reported in this paper are given in local time (LT), defined as
 125 Central European Time (UTC+1) without daylight saving time (DST).

2.2 Experimental setup

Three setups were operated at BE-Vie from 2022 to 2024: the TOP system, the TRUNK system, and the PROFILE system. These are described below, and details about the related analytical techniques are provided in Sect. 2.3, while flux computations are discussed in Sect. 2.3.3 and 2.4.




Figure 2. Schematic representation of the concentration and sonic anemometer profile setup, including a zoom on the pneumatical system. Instruments include: FOS – fast ozone analyser; T400 – slow ozone analyser; PTR-ToF-MS – fast VOC analyser; HS-50 & R3-50 – sonic anemometers; CB – calibration bottle; MFC – mass flow controller; CC – catalytic converter; MN – manometer; MP – membrane pump; MFM – mass flow meter; BV – buffer volume; NV – needle valve. Line colours indicate function: purple – ozone line, red – ozone/VOC line, blue – background line, orange – calibration line, brown – exhaust line. The positions of the solenoid valves are indicated by NO (normally open, i.e. open without electrical activation) and NC (normally closed, i.e. open only when electrically activated).

130 2.2.1 TOP system

To characterize the bidirectional fluxes of BVOCs and O_3 above the canopy, a first setup was installed at the top of the 51 m high flux tower (level H7 in Fig. 2). Flux measurements were performed using the eddy covariance (EC) technique, which is based on fast measurements of the vertical wind velocity and the mixing ratio of the tracer of interest.

At the top of the tower, a 3D HS-50 sonic anemometer (*Gill Instrument Ltd, UK*) was used to record 3D wind components
 135 at a frequency of 10 Hz (see Table 1 for a summary of the instruments used). To measure BVOC and O_3 mixing ratios, ambient air was sampled at level H7 close to the HS-50 measurement volume and pumped to a wooden shelter at ground level through
 60 m long and 6.4 mm inner-diameter PFA (perfluoroalkoxy alkane) tubing (*Fluortechnik-Deutschland, Germany*) at a flow rate of about 10 SLM (litre per minute, under standard conditions of pressure (100 kPa) and temperature (273.15 K)). This line was heated and thermally insulated to prevent condensation. A PFA raincap and a PTFE (polytetrafluoroethylene) particulate
 140 filter (2 μm pore size, *Pall Corp, Port Washington, NY, USA*) were placed at the tubing inlet. The filter was replaced monthly.

In the shelter, ambient air was subsampled towards a PTR-ToF-MS 4000 (*Ionicon Analytik GmbH, Innsbruck, Austria*) instrument for fast measurements (10 Hz) of VOC mixing ratios. On the same line, air was also subsampled towards a T400



145 The T400 instrument was used to obtain absolute and stable measurements of ozone, but its measurement frequency was not
 suitable for flux computation. Therefore, an additional Fast Ozone Analyser (*FOS, Sextant Technology Ltd, Wellington, New
 Zealand*) was installed at the top of the tower. Ambient air was sampled close to the sampling point of the VOC/O₃ line and
 brought to the FOS analyser through a 6.5 m long and 6.4 mm inner-diameter PFA tube with a flow rate of about 3.5 SLM.
 This line was also heated, thermally insulated, and protected by a PFA raincap and a PTFE particulate filter. The FOS signal
 150 was recorded at a frequency of 10 Hz.

Table 1: Overview of instruments and related measurements. The access to the four distinct datasets (groups i, ii, iii and iv) are
 given in Sect. *Code and data availability*.

Compound/ Parameter	Instrument (manufacturer)	Measurement principle	Acquisition frequency (aggregation frequency)	Variable type	Heights(s) [m a.g.l.]
<i>(i) Main measurements (present study)</i>					
VOC	PTR-ToF-4000 (Ionicon Analytik)	proton-transfer mass spectro.	10 Hz	Conc.	3-11-19-27-
			(35 min)	profile	35-43-51
			10 Hz	Conc.	51
			(1 min or 30 min)	& Flux	
O ₃	T400 (Teledyne)	UV absorption	10 Hz	Conc.	3 ^a
			(1 min or 30 min)	& Flux	
			6 s	Conc.	3-11-19-27-
			(35 min)	profile	35-43-51
O ₃	FOS (Sextant)	chemi- luminescence	6 s	Conc.	51
			(1 min or 30 min)	& Flux	
			6 s	Conc.	3 ^a
			(1 min or 30 min)	& Flux	
O ₃	FOS (Sextant)	chemi- luminescence	10 Hz (30 min)	Flux	51
O ₃	FOS (Sextant)	chemi- luminescence	10 Hz (30 min)	Flux	3
3D wind	HS-50 (Gill)	ultrasonic time-of-flight	10 Hz (30 min)	-	6-18-51

Table 1 (continued)



Compound/ Parameter	Instrument (manufacturer)	Measurement principle	Acquisition frequency (aggregation frequency)	Variable type	Heights(s) [m a.g.l.]
3D wind	R3 (Gill)	ultrasonic time-of-flight	10 Hz (30 min)	-	3-12-24-30-36
<i>(ii) ICOS measurements</i>					
Comprehensive set of meteorological and phenological variables					
A complete description of ICOS variables is available along with the downloaded dataset.					
<i>(iii) ISSeP measurements</i>					
Set of atmospheric pollutants: VOC, PAH, O ₃ , NO, NO ₂ , CO, SO ₂ , Hg, PM _{2.5} , PM ₁₀ , ultrafine particles, aerosols					
For the complete list of compounds, instruments, acquisition frequency and heights, see Table S1.					
<i>(iv) Remote-sensing measurements</i>					
Tropospheric column + vertical profile of HCHO, NO ₂ , aerosols					
For the complete list of compounds, instruments, acquisition frequency and heights, see Table S1.					
^a Not available in 2022					

2.2.2 TRUNK system

The lowest level of the VOC/O₃ profile (H1, see Sect. 2.2.3) was located on a 3 m high mast installed in 2022 in the beech sector, approximately 10 m west of the tower. Ambient air was sampled through a 60 m long tube leading to the shelter. In 2023, a second FOS instrument (5 m long and 6.4 mm inner-diameter PFA tube, flowrate of 3.5 SLM) and a 3D R3-50 sonic anemometer (*Gill Instrument Ltd, UK*) were added to the 3-m mast to establish a system compatible with EC measurements. This configuration, referred to as the TRUNK system, samples air from within the trunk space of the forest, primarily to characterize the VOC and ozone fluxes exchanged by the soil. However, a small part of this space is occupied by young woody vegetation, as shown in Fig. 3.

2.2.3 PROFILE system

The characterization of exchange processes inside the canopy is based on two types of profiles: (i) a VOC and O₃ concentration profile and (ii) a turbulence profile measured using 3D sonic anemometers. The two 60 m long lines installed for the TRUNK and TOP systems were supplemented by five additional tubes of the same length, material and diameter, resulting in a concentration profile with seven evenly distributed sampling points. The measurement heights for levels H7–H1 are given in Table 1. All of these lines were made of PFA, heated, and connected to a manifold inside the shelter. The TOP and TRUNK sonic anemometers were supplemented by an additional HS-50 and R3-50 sonic anemometers, respectively mounted on 3 m and 2 m long aluminium arms attached to the tower. These sonic anemometers were rotated over the three-year measurement period to characterize turbulence at as many heights as possible. The corresponding heights are listed in Table 1.



Figure 3. Panoramic view (360°) taken from the mast of the TRUNK system in summer 2023.

2.2.4 Pneumatical setup

In the shelter, a manifold system made of seven three-way PFA solenoid valves was used to sequentially sample ambient air from levels H7 to H1. During TOP (H7) and TRUNK (H1) system measurements, air was sampled for at least 30 minutes to apply the EC method. In PROFILE mode, air was sequentially sampled from levels H6 to H1, for respective periods of 5 minutes. To obtain a complete concentration profile, the last 5 minutes of a preceding TOP mode were included in the PROFILE cycle.

Downstream of the manifold, air was subsampled towards the PTR-ToF-MS and T400 instruments. Due to the pressure drop along the 60 m sampling line, the inlet pressure of the T400 becomes too low for proper operation. To compensate for this, a membrane pump (*KNF N86KT.18, KNF Neuberger GmbH, Freiburg, Germany*) was installed upstream of the T400 to bring the sampled air to atmospheric pressure. The PTR-ToF-MS requires regular calibration and background measurements. For background measurements, the ambient air subsampled from the H7 line was sent through a catalytic converter (*Parker, type HPZA-3500, Haverhill, MA, USA*) to remove VOC before being analysed. For calibration, a user-defined flow rate of a VOC standard (*Apel-Riemer Environmental Inc., FL, USA*) was mixed with zero-VOC air from the background line. More details on VOC quantification are given in Sect. 2.3.2. A total flow of about 10 SLM was pumped through the selected sampling line by a membrane pump (*MD12C NT, Vacuubrand, Germany*) and manually controlled by a needle valve. A buffer volume was added between the needle valve and the flow meter to dampen the flow and pressure variations caused by the pump. The non-selected sampling lines were continuously flushed with ambient air to allow for fast switching between lines.

185 2.2.5 Data coverage

During the three years of measurements, the VOC and O₃ analytical instruments were operated as simultaneously as possible, with particular emphasis on the spring, summer, and autumn periods. Figure 4 summarizes the monthly percentages of half-hour intervals that were successfully measured by each system.




Figure 4. Percentage of half-hours covered per month and per year for measurements shared between the TOP, TRUNK and PROFILE systems.

In 2022, the PTR-ToF-MS and the T400 instruments sequentially measured 30-minute periods at the TOP level and in PROFILE mode. On average, TOP measurements were performed 3.5 times more frequently than PROFILE measurements. Background VOC measurements were systematically performed for 30 minutes every 4 hours, and single-point VOC calibrations were performed every 3 to 4 days. These procedures limited data coverage for other measurement systems.

In 2023 and 2024, TRUNK measurements were added. In this updated sequence, TOP measurements occurred on average 2.5 times more frequently than PROFILE or TRUNK measurements.

Overall, ozone flux measurements at the TOP and TRUNK levels were more frequent than those for VOCs, since the two FOS analysers operated continuously, while the PTR-ToF-MS had to alternate between levels. This will be discussed further in Sect. 2.4.2.

2.3 VOC measurements

2.3.1 PTR-ToF-MS settings

The PTR-ToF-MS analyser was operated in the H₃O⁺ mode. Air was subsampled by the instrument through a 1.2 m long silcosteel capillary (1 mm inner diameter). The drift tube reactor was operated at a pressure of 3.3 hPa and a reduced electric field (E/N) of 135 Td (1 Townsend = 10⁻²¹ V m²). Both the inlet line and the drift tube reactor were kept at 353 K. The inner part of the drift tube has an EVR (Extended Volatility Range) coating to minimize losses of sticky compounds to the surfaces of the reactor. A hexapole ion guide is used to efficiently transport ions from the drift tube to the mass analyser section.

The Time-of-Flight mass analyser was operated at an extraction time and frequency of 2 μs and 40 kHz, respectively, allowing for a mass-to-charge (m/z) range of 7–392. A small flow of diiodobenzene, provided by a built-in permeation device (PerMaSCal) was constantly added to the drift tube for frequent mass scale calibration based upon the position of protonated diiodobenzene (at m/z 330.848), its fragment (at m/z 203.943), and H₃¹⁸O⁺ (at m/z 21.022) on the detector. Time-of-flight mass



spectra were integrated over 100 ms to allow for EC flux calculations. A maximal mass resolution ($m/\Delta m$) of 4200 was obtained
 210 for the high mass range. At m/z 59.049 (protonated acetone) and m/z 137.132 (protonated monoterpenes), the mass resolution
 was about 3400 and 3800, respectively. The peak area related to primary ion signals ranged from 8766–25554 (m/z 21.022
 related to H_3O^+) to 11–619 (m/z 38.033; $H_3O^+(H_2O)$) counts per second (cps) over the three years of measurements. By
 considering the relative transmission of the instrument at m/z 21.022 and 38.33 (respectively 9–15% and 29–43% relative to
 the transmission at m/z 180.937), together with the relative abundance of the $H_3^{18}O^+$ and $H_5^{17}O^{16}O^+ + DH_4^{16}O_2^+$ isotopes
 215 (respectively 488 and 669), this results in a transmission-corrected source ion count rate of $40 \cdot 10^6 - 110 \cdot 10^6$ cps.

2.3.2 Signal processing and quantification

Peak identification and integration of recorded spectra were performed on a near-daily basis with the Ionicon Data Analyzer
 software (IDA; 2022, v 1.0.0.2; 2023, v 2.0.1.2; 2024, v 2.2.1.1). Subsequent data processing was performed in a python-based
 framework referred to as the Peak Area Processing software (PAP) which was used for non-targeted peak selection, instrument
 220 characterization, and quantification of mixing ratios with related uncertainties at both 10 Hz and 1 min time resolution. A
 general overview of these processes is given below. An extended version of the discussion on how VOCs were quantified and
 details on the configuration of the PAP software and instrument operations can be found in Sect. S2.

Mass selection

The mass scale was calibrated in the IDA software every 60 s and peak-shapes were considered stable over the course of
 225 1 hour. Independent peak identification was performed for each IDA analysis with up to 8 m/z values identified per peak system
 (collection of isobaric peaks with the same nominal mass). A peak system was only considered for analysis if its maximum ion
 signal intensity exceeded 0.1 cps. To obtain a list of non-targeted m/z ratios, Density-Based Spatial Clustering of Applications
 with Noise (Ester et al., 1996), or DBSCAN, was employed to identify mass-to-charge ratios where peaks were regularly
 observed. An automated selection of ion m/z values has been performed based on (i) the width of the DBSCAN intervals
 230 centred around the mass (stability of peak identification by IDA), (ii) the fraction of data above the limit of quantification
 (significance of concentrations), and (iii) the m/z localization (interpretability of the data). Afterwards, a manual selection to
 discard m/z values related to isotopes or hydrated ion species was done. Compound attribution was performed by making use
 of PTR-MS databases (Pagonis et al., 2019; Yáñez-Serrano et al., 2021) and measurements reported at ecosystem sites (Kim
 et al., 2010; Hellén et al., 2018; Schallhart et al., 2018; Pfannerstill et al., 2021) to identify the compounds most likely to
 235 contribute to the observed signal.

Instrument characterization

Calibrations were performed every 3–4 days to characterize the instrument transmission and calculate calibration factors for
 compounds included in the calibration bottle (Table S2). Instrument transmission (relative to the one at m/z 21.022) was
 calculated using a subset of compounds (associated to m/z 33.033, 42.034, 45.033, 59.049, 79.054, 93.070, 107.086, and
 240 180.937) included in the calibration bottle. The transmission curve between m/z 21.022 and 180.937 was defined through
 linear interpolation of transmissions obtained during calibrations. At high m/z , the transmission curve becomes more stable
 and we assumed a constant behaviour ($Tr_{m/z > 180.937} \equiv Tr_{180.937}$).



Mixing ratio calculation

The mixing ratio of compounds was calculated using either the kinetic or the calibration approach, similar to the discussion
 245 in the ACTRIS measurement guidelines for VOC analysis with PTR-MS instruments (Dusanter S. et al., 2025). In both cases,
 integrated peak areas (cps) obtained from IDA were corrected for ion transmission, and normalized with respect to a source
 ion peak area of 10^6 cps to account for variations in the source ion production. VOC volume mixing ratios (VMR) were
 calculated by dividing the background-subtracted transmission-corrected normalized peak areas (tc-ncps) by the compound-
 dependent sensitivities (tc-ncps ppbv⁻¹). For compounds present in the calibration standard, sensitivities were calculated at
 250 every calibration. At m/z values not related to compounds available in the standard, sensitivities were derived from calculated
 H_3O^+ /VOC reaction rate constants and product ion distributions, as well as drift tube reactor conditions, as explained in more
 detail in Sect. S2.1.3.

Uncertainty quantification

The expanded or combined uncertainty (with a coverage factor of 2) of the calculated VOC mixing ratios combines both the
 255 precision of the peak areas and systematic errors (accuracy) associated with instrument calibration or sensitivity calculations:

$$\text{expanded uncertainty} = 2 \cdot \sqrt{\text{precision}^2 + \text{accuracy}^2} \quad (1)$$

The precision of the peak areas can be determined from Poisson statistics and approximated by the square root of the
 integrated counts over the measurement interval. However, peak areas obtained with IDA are inherently baseline-subtracted
 and can be negative for small peaks, hereby invalidating the above-mentioned approach. For 1-min averaged data, the precision
 260 was therefore approximated by the precision of the mean 100 ms peak area over the averaging interval, assuming a normal
 distribution.

The systematic uncertainty (accuracy) of the volume mixing ratios for compounds included in the calibration standard
 was obtained by combining the stated uncertainty on the mixing ratios of those compounds in the standard ($\sim 2.5\%$) with
 the uncertainty of the dilution factor ($\sim 1.5\%$). A conservative systematic uncertainty of $\sim 56\%$ was used for mixing ratios
 265 quantified using the kinetic approach, due to large uncertainties on the calculated H_3O^+ /VOC rate constants ($\sim 25\%$) and on
 the product ion distributions ($\sim 50\%$). This accuracy is close to results by Sekimoto et al. (2017) which showed that measured
 sensitivities agreed within 20–50% with theoretical sensitivities calculated using molecular mass, elemental composition, and
 functional group of the analyte.

For the combined uncertainty, Simon et al. (2023) included an additional 5% uncertainty due to relative humidity effects.
 270 However, as the $H_3O^+(H_2O)$ signal was generally less than 3% that of $H_3O^+ + H_3O^+(H_2O)$ (Fig. S4), the impact of relative
 humidity was assumed to be small and this additional uncertainty is not considered here.

The limit of detection (quantification) for concentrations is defined as three (ten) times the precision of the associated
 background measurement, of which we only considered the last 5 min to assure equilibrium. The mean and standard deviation
 of the measurement distribution during these 5 min were used to define the background value and its precision, respectively.

275 More details on uncertainty characterization are given in Sect. S2.1.4.



2.3.3 Flux computation

Half-hourly VOC fluxes were computed following the general procedure commonly used for eddy covariance flux calculation. This included raw data despiking and gap filling, block averaging, coordinate rotation of wind components, time-lag estimation by maximizing covariance and data shifting, covariance calculation and correction for high-frequency spectral losses.

280 EC fluxes are commonly filtered under low turbulence conditions, particularly for CO₂ fluxes, using a friction velocity (u^*) threshold. However, given the different nature of VOC exchange processes compared to CO₂, no u^* filtering was applied in this study. Additional information and justification for this choice are provided in Sect. S3.1.

All flux computations, from raw sonic anemometer data and processed VOC concentrations, were performed using the GEDdySoft ("Gembloux Eddy-covariance Software") software. GEDdySoft was initially developed from the open source InnFlux
 285 software (Striednig et al., 2020), which supports both conventional and disjunct eddy covariance analyses. The original MATLAB® code was converted to Python, and the software has been continuously expanded with additional data handling and flux processing features. For this study, version 4.0 of GEDdySoft was used and is publicly available (see Code and data availability section).

Unless explicitly stated, the same procedure was applied to both the TOP and TRUNK systems. One exception concerns the
 290 tilt correction of the sonic anemometers: the double rotation method was used for the TOP system, while the sector-wise planar fit method was applied to the TRUNK system. Below-canopy turbulence was sometimes low, so applying the double rotation method at the half-hourly scale could result in high rotation angle values and erroneous fluxes (notably artificially high absolute fluxes; see Fig. S6, where the effect is most visible for O₃ fluxes). The sector-wise planar fit method, which computes a single set of tilt angles based on long-term time series, was able to solve this issue and was therefore applied to TRUNK fluxes (with
 295 tilt angles determined for 2023 and applied to both 2023 and 2024). Details about both methods are given in Wilczak et al. (2001). The structure at the back of the sonic anemometer used for the TOP system (HS-50) can act as an obstacle to airflow. Consequently, fluxes coming from a $\pm 10^\circ$ wind sector around the sonic anemometer arm were discarded, following the ICOS protocol (Sabbatini et al., 2018). No filtering was applied to TRUNK fluxes, which were measured with an omnidirectional sonic anemometer.

300 The main steps of flux computation described above were also applied to ozone fluxes. However, considering the differences in the measurement setup, several methodological aspects diverged between the two flux calculations. These aspects are detailed in the present Section for VOCs and in Sect. 2.4.1 for ozone.

Lag-time determination

The lag time between the vertical wind component fluctuations w' and the VOC concentration fluctuations c' —resulting from
 305 inlet-to-detector transit time and instrument response—was determined by maximizing their covariance within a physically plausible window (11–15 s) for each flux window and each m/z. If the maximum covariance occurred at the edge of the window, a fixed lag of 13 s was assigned. This default value corresponds to both the centre of the window and the median lag time observed for monoterpenes (VOC with the highest signal-to-noise ratio, SNR) across the entire measurement campaign.



It is also consistent with expectations based on tube dimensions and flow rates (first-order estimate of the residence time, ~ 10 s).

To refine lag estimation and reduce spurious detections under low SNR conditions, the covariance function was smoothed using a 15-sample (1.5 s) moving average prior to the search. This approach, as recommended by Taipale et al. (2010) and Langford et al. (2015), effectively mitigates “mirroring” effects associated with low fluxes while preserving sensitivity to time-varying lags.

The stability of lag times was assessed visually across all campaigns and for all m/z values. Under high SNR conditions, lag times were found to be highly stable, with the exception of two ions species: 33.033 (protonated methanol) and 47.049 (protonated ethanol) (as well as H_3O^+ and its hydrates, which were excluded from the analysis). For these two m/z , a small but systematic dependence on air relative humidity (RH) was observed. When RH exceeded 95%, the lag time tended to increase in absolute value, shifting from -13 s up to -20 s (data not shown). Consequently, the centre of the lag-search window was parametrized as a function of RH for these two ions.

Correction for flux high-frequency losses

An experimental co-spectral approach was adopted to correct for high-frequency attenuation, and was applied in three steps. First, a transfer function was derived by dividing the normalized mean co-spectrum of wc (attenuated) by that of wT (considered ideal), and was then fitted using a Lorentzian function to obtain a single cut-off frequency. Second, to derive a flux correction factor at the half-hour scale, the area under the wT co-spectrum (representing the non-attenuated flux) was divided by the area under the same co-spectrum multiplied by the transfer function (representing the attenuated flux). Third, these correction factors, obtained for a limited number of half-hours after drastic data filtering (see Sect. S3.3), were sorted and averaged into five wind speed classes for unstable and stable atmospheric conditions separately. These two lookup tables were then used to assign a correction factor for each individual half-hour, depending on its wind speed and atmospheric stability. The procedure was applied to monoterpenes to benefit from their high SNRs. Since the correction characterizes the experimental setup in combination with the flow characteristics, without any expected compound-specific aspects, it was then applied to all other compounds.

The choice of a co-spectral, rather than spectral, approach was motivated by its capacity to empirically integrate all causes of signal attenuation—tube damping, instrumental response, and sensor separation—without relying on a theoretical transfer function for sensor separation. Furthermore, our method does not require a reference co-spectrum, whose formulation and ability to represent real conditions at the site are often subject to discussion. It should be noted that, following the recommendations of Peltola et al. (2021), the square root of the transfer function was used rather than the transfer function itself to derive both the cut-off frequency (step one) and the correction factor (step two), in order to correctly account for the phase shift introduced by our lag-time determination method.

For the TRUNK system, step one could not be applied due to generally low fluxes and associated low SNRs across all compounds. However, given that the cut-off frequency reflects the setup alone and that the TRUNK and TOP configurations were identical, the cut-off frequency derived from the TOP system was applied. Steps two and three were executed independently for the TRUNK level, as atmospheric turbulence is location-specific.



This procedure led to correction factors ranging from 1.10 to 1.26 for the TOP system, and from 1.10 to 1.40 for the TRUNK
 one. More details are provided in Sect. S3.3.

Flux quality control

Flux quality was assessed at two levels. First, the high-frequency inputs used for flux computation—VOC mixing ratios and sonic anemometer data—were (in)validated based on statistical tests developed by Vitale et al. (2020). These tests have
 the advantage, compared to the more traditional tests of Vickers and Mahrt (1997), of being completely automatable, not
 requiring user choices on, for example, threshold values. Adapted from the open-source RFlux toolbox (<https://github.com/icos-etc/RFlux>), they were integrated in our GEDdySoft processing pipeline. The combination of the tests present in RFlux and
 described in (Vitale et al., 2020) (HF₅, HF₁, HD₅, KID) led to the creation of an instrumental flag, hereafter named *flag_instr*.
 Only a few half-hour periods were flagged for instrumental issues (Fig. S12), with a maximum of 12% of fluxes flagged for
 m/z 80.058 in the TOP system.

Other instrumental tests available in the RFlux toolbox—including AL1, DIP, and DDI—although implemented in the ICOS
 standard processing pipeline, were not applied here. These tests tended to flag excessively long periods, including VOC fluxes
 with low SNR, as well as those displaying clear and reasonable diurnal dynamics. Furthermore, they have not yet been described
 in peer-reviewed literature and have never been validated for compounds with inherently lower SNR such as VOCs. For these
 reasons, they were considered unsuitable for our dataset and were excluded from the present analysis.

Second, a quality flag was assigned to the computed fluxes based on steady-state (stationarity) and integral turbulence char-
 acteristic tests, following the methodology developed by Mauder and Foken (2004). This flag, named *flag_MF* (for Mauder
 and Foken), can take three values: 0 indicates high-quality fluxes; 1 corresponds to fluxes suitable for general analysis; and 2
 denotes poor-quality fluxes. For stationarity, we preferred the test proposed by Mahrt (1998) over that of Foken and Wichura
 (1996), as it proved more effective at flagging anthropogenic influences (see Sect. 2.3.4). Fluxes flagged at level 2 were dis-
 carded for the remainder of this study. The percentage of half-hours discarded according to *flag_MF* reached up to 20% for the
 TOP system and 33% for the TRUNK system, both cases for m/z 137.132. This filtering is primarily due to the influence of
 anthropogenic sources (see Sect. 2.3.4), an effect further described in Sect. S3.4.

A combined flag (*flag_tot*) was created by integrating *flag_instr* and *flag_MF*, as well as an additional flag (*flag_plume*)
 designed to filter out periods affected by anthropogenic emissions (see Sect. 2.3.4). Only fluxes passing all three flag criteria
 were assigned a value of 0 for *flag_tot* and retained for further analysis in this study.

Correction for density fluctuation

VOC fluxes can be biased by fluctuations in temperature and water vapour, which affect air density. These effects can be ac-
 counted for using the Webb–Pearman–Leuning (WPL) corrections (Webb et al., 1980). In our setup, temperature fluctuations
 were considered attenuated due to the length of the sampling lines, which largely exceeded the 3.20 m minimum recommended
 by Rannik et al. (1997). Moreover, during flux measurements using a PTR-ToF-MS, Loubet et al. (2022) reported that correc-
 tions for water vapour fluctuations were very small (less than 2% for 75% of the time). For these reasons, WPL corrections
 were not applied to VOC fluxes in this study.



Flux storage

380 To obtain the net exchange of VOCs, a storage term should be added to the turbulent flux. The storage flux represents the temporal variation of the mixing ratio of the compound below the EC measurement height, and is expected to be non-negligible for tall towers and under low turbulence conditions, such as at night. The storage term was calculated following the procedure described by Montagnani et al. (2018):

$$S_i = \sum_{j=1}^N \rho_{d_i} \left(\frac{\Delta c_i}{\Delta t} \right)_j \Delta z_j \quad (2)$$

385 where S_i is the storage flux for tracer i ($\text{nmol m}^{-2} \text{s}^{-1}$), ρ_{d_i} is the dry air density of layer j , $\left(\frac{\Delta c_i}{\Delta t} \right)_j$ is the temporal variation of mixing ratio for tracer i (nmol mol^{-1}) at height j over a time period Δt (s), and Δz_j is the thickness of layer j (m). Temporal changes in mixing ratios were derived from concentration profiles measured every 2 to 2.5 hours.

The resulting storage fluxes exhibited erratic temporal dynamics and, for many VOCs, their magnitudes often exceeded those of the turbulent flux. We consider that calculating temporal variations in concentrations over 2 to 2.5 hour intervals is unlikely
 390 to accurately capture true storage processes within the canopy, particularly for reactive compounds with short atmospheric lifetimes. This limitation likely results in unreliable storage flux estimates. To avoid degrading the overall data quality, storage fluxes were therefore not included in the net flux calculation.

2.3.4 Filtering of anthropogenic sources

Over the three years of measurements, we occasionally observed abnormally high VOC mixing ratios under winds from the
 395 230–270° sector. This phenomenon had already been reported for monoterpenes and methanol during earlier campaigns in 2009 and 2010 (Laffineur et al., 2011, 2012), and was attributed to a wood panel factory located 3 km south-west of the tower. Although outside of the flux footprint, this facility was deemed a dominant VOC source compared to surrounding biogenic emissions, with magnitude sufficient to influence local measurements.

In the present study, this anthropogenic influence was mostly observed at night, yet occurred irregularly and did not af-
 400 fect all VOCs uniformly, with many of the detected ions not being impacted at all. The most affected ions included protonated monoterpenes (e.g. m/z 137.132), as well as fragments and isotopes, sesquiterpenes, and potential derivatives such as m/z 135.117 (likely p-cymene). A few low molecular weight oxygenated compounds—methanol, formaldehyde, and acetic acid—were also affected, though to a lesser extent. During plume events, the affected VOCs exhibited elevated mixing ratios and mixing ratio variances, and occasionally large negative fluxes (i.e. apparent deposition), which are not expected for
 405 compounds like monoterpenes.

To identify such episodes, Laffineur et al. (2011) proposed an upper threshold on the 30-min variance of monoterpene mixing ratios in the 230–270° wind sector, assuming that incomplete mixing between anthropogenic and biogenic sources led to higher variability in mixing ratios. However, extending this method would require manually identifying all impacted ions and setting compound-specific thresholds, which is subjective and impractical.



410 To address this limitation, we developed an automatic flagging procedure. For each VOC, we computed q_{99} , the 99th percentile of its 30-min mixing ratio variance calculated across all wind directions except 180–300°, which is subject to sawmill influence. Then, any half-hour within the 180–300° sector with a variance exceeding q_{99} was flagged as potentially affected (Fig. S11). This approach successfully identified affected VOCs (7% of half-hours flagged for monoterpenes or methanol, see Fig. S12) while leaving unaffected compounds mostly untouched (e.g. only 0.3% flagged for isoprene). Although many spoiled
 415 half-hours had already been identified by the stationarity test ($flag_{MF}$), this new $flag_{plume}$ completed the identification of such events.

While some fluxes exhibited a well-defined lag time during plume events—suggesting real turbulent deposition, as also observed by Bamberger et al. (2011)—these episodes showed erratic dynamics and no consistent relationship with environmental drivers. Therefore, to focus the analysis on biogenic VOC exchange, we excluded flagged fluxes from further processing.
 420 Similarly, concentrations flagged by $flag_{plume}$ were not considered in the rest of this study.

2.3.5 Identification of significant exchanges

Among the m/z retained by the DBSCAN algorithm, not all ions exhibited significant fluxes. To determine whether a compound is significantly exchanged, it is common practice to compare half-hourly fluxes with their corresponding flux limit of detection (LOD_f). In this study, LOD_f was estimated at the 99% confidence level as three times the random flux error, calculated in
 425 GEDdySoft following the method described in Finkelstein and Sims (2001).

Extrapolating such comparisons from the half-hourly scale to longer periods, in order to assess whether a compound is consistently exchanged, is not straightforward. Loubet et al. (2022) addressed this by comparing the mean flux of VOCs with an averaged LOD_f , itself calculated as the square root of the sum of squared individual LOD_f values, divided by the number of records (Langford et al., 2015). While this approach allows a simple comparison over an entire campaign, it may fail to
 430 detect compounds significantly exchanged only during short periods.

To overcome this limitation, we developed a novel three-step methodology, applied individually to all m/z values. First, each half-hour period was flagged as significant if the absolute flux exceeded its LOD_f . Second, a day was considered significant if at least 25% of the daytime half-hourly periods (defined as 08:00–20:00 LT) were significant. Finally, we considered the number of consecutive significant days. Given that the mechanisms responsible for VOC emission or deposition are unlikely to
 435 occur on a single day, we selected m/z values for which at least three consecutive days of significant exchange were observed during the measurement period.

2.4 Ozone measurements

2.4.1 Raw ozone fluxes

Ozone flux measurements were performed with a fast ozone analyser (FOS). In the instrument, O_3 molecules present in ambient
 440 air react with a coumarin disc (solid or dry phase), thereby producing a light signal (chemiluminescence reaction) recorded in units of Volts (V) at a high frequency (10 Hz). Coumarin discs were produced from silica plates impregnated with a solution of



coumarin, methanol and ethylene glycol. Since the efficiency of the chemiluminescence reaction decreases over time, coumarin discs were replaced on average once a week during the 2022 and 2023 campaigns. In 2024, disc replacements were less frequent considering the important workload (up to 36 days between disc replacements).

445 Raw ozone fluxes were computed using the EddyPro® software, applying the same processing steps as described in Sect. 2.3.3. As for VOCs, no u^* filtering was applied. Given the differences in the measurement setup, certain aspects of the flux computation differ and are detailed below.

Lag time determination

450 As for VOCs, the physical lag time between w' and c' was determined by maximizing the covariance function within a predefined lag window. When no clear extremum was found within this window, a nominal lag value was applied. Due to the higher SNR of O_3 fluxes compared to many VOCs, no smoothing of the covariance function was necessary. Across the different measurement campaigns, the median lag time was close for the O_3 TOP and TRUNK systems, with values of 2.2 s and 2.0 s, respectively.

Correction for flux high-frequency losses

455 The three-step experimental co-spectral approach applied to VOCs was also used for ozone. The only difference is that, due to more pronounced ozone fluxes measured at the TRUNK level compared to VOCs, it was possible to determine a cut-off frequency for the TRUNK system.

Correction factors ranged from 1.12 to 1.37 for the TOP system, and from 1.03 to 1.28 for the TRUNK one. Further details are provided in Sect. S3.3.

460 Flux quality control

An instrumental flag was generated by combining the individual tests developed by Vickers and Mahrt (1997) and implemented in EddyPro®. The attack angle and non-steady wind tests were excluded, as they tended to over-flag data that did not appear problematic upon visual inspection. Very few half-hour periods (<1%) were affected by instrumental issues (Fig. S12).

465 EddyPro® also computes a flux quality flag similar to the previously described *flag_{MF}*, based on the stationarity test of Foken and Wichura (1996). Over the three-year period, 9% of the fluxes were flagged as level 2 (poor quality) for the TOP system, and 17% for the TRUNK system (Fig. S12).

No influence from the wood factory was detected in the ozone data; therefore, no *flag_{plume}* was applied. Data flagged by *flag_{tot}* were excluded from further analysis in this study.

Correction for density fluctuation

470 From the original WPL equation (Webb et al., 1980), we neglected temperature fluctuations—assumed to be attenuated through the sampling tubing, as for VOCs—and only assessed the impact of water vapour fluctuations. The latter resulted in a correction of less than 1% of the raw fluxes on average. This value is much lower than the one reported in Gerosa et al. (2022a)—where temperature-related corrections could not be neglected considering the large-diameter tubing—and also well below our flux random uncertainty estimated according to Finkelstein and Sims (2001). For these reasons, no WPL correction was applied.

475 Flux storage

The ozone storage was computed with Eq. 2. For TOP storage, 7 layers were considered (H1 to H7, see Fig. 2). Profile



measurements were taken every 2 or 2.5 hours, so half-hourly storage values were linearly interpolated between adjacent measurements. Since profiles were not continuously scanned throughout the measurement campaigns, a gap-filling approach was required. For TOP storage, we used the storage computed from TOP mixing ratio (which was available at least once every three half-hours) as a proxy for the actual storage term. In combination with other predictors (friction velocity u^* , air temperature, incoming PPFD at the top of the canopy, hour of the day, and month number), we trained a Random Forest regression model to estimate and gap-fill storage fluxes derived from the concentration profile. This procedure represents an improvement over the methodology used by Aubinet et al. (2018), who employed look-up tables to gap-fill missing storage data.

No model was required to gap-fill TRUNK storage fluxes. Since only one layer was considered, the storage flux was computed directly when the T400 was in PROFILE mode (using level H1) or TRUNK mode. The missing half-hour values were linearly interpolated between adjacent data points, provided that gaps were shorter than 3 hours.

Unlike for VOCs, ozone storage showed a consistent diel pattern (Fig. S14, similar to that observed by (Finco et al., 2018)). We therefore decided to add the storage flux to the calibrated turbulent flux (calibration described in Sect. 2.4.2) to obtain the net ozone flux.

2.4.2 Ozone flux calibration

Since the FOS analyser provides only relative ozone concentrations, the raw fluxes were calibrated using a T400 UV absorption analyser, which delivers absolute and stable O_3 mixing ratios in ppbv (nmol mol^{-1}). The T400 instrument was calibrated annually before each measurement campaign, and automatic zero and span checks were performed nightly to monitor potential drifts.

As noted in Muller et al. (2010) and Zhu et al. (2015), calibration procedures for FOS-based ozone fluxes are rarely detailed, despite their significant impact on data quality. The FOS analyser's sensitivity decreases over time due to the consumption of coumarin molecules in the chemiluminescence reaction with ozone. This sensitivity loss is therefore non-linear (Güsten and Heinrich, 1996), varies from one disc to another, and may include an offset between fast and slow signals.

Muller et al. (2010) compared three post-processing calibration methods (see Table 2) by comparing fast and slow ozone signals (1) at the half-hourly scale without considering an offset (Ratio Method, RM), (2) at the half-hourly scale with an offset determined for each coumarin disc (Ratio Offset Method, ROM) and (3) over the lifetime of a coumarin disc using a single regression (Disc Calibration Method, DCM). The RM captures changes in FOS sensitivity more effectively than the DCM, which assumes a stable response throughout the disc lifetime. The ROM represents a compromise; although it applies a half-hourly ratio, the offset remains fixed over the disc's life and may not follow short-term sensitivity shifts.

Zhu et al. (2015) later introduced the Regression Variance Method (RVM), which identifies an optimal averaging window between the FOS and T400 signals using Allan-Werle variance analysis. A 3-minute period was found to best track sensitivity changes, but it is also more susceptible to white noise than RM and ROM (30-min windows), and especially DCM (multi-day windows). Both RM and RVM assume zero offset between fast and slow signals—an assumption invalidated by observations from Muller et al. (2010) and Zhu et al. (2015).



An important practical constraint in our study was the limited temporal overlap between fast and slow ozone measurements. The T400 analyser alternated between TOP, TRUNK, and PROFILE modes, preventing continuous half-hourly reference data at any one level. Therefore, we developed a new approach: the Period Calibration Method (PCM). Inspired by DCM, PCM accounts for potential offsets and reduces sensitivity to high-frequency noise. However, unlike DCM, it identifies optimal sub-periods during which the FOS sensitivity remained consistent, allowing more accurate tracking of temporal variability in analyser performance.

With the PCM (Period Calibration Method), calibrated fluxes were computed as:

$$F_{cal} = \frac{\beta_1 F_{raw}}{v_{mol}} \quad (3)$$

where F_{cal} is the calibrated flux ($\text{nmol m}^{-2} \text{s}^{-1}$), F_{raw} is the raw flux obtained from FOS measurements (V m s^{-1}), β_1 is the slope of the regression curve established between T400 and the FOS measurements ($\text{nmol mol}^{-1} \text{V}^{-1}$), and v_{mol} is the air molar volume ($\text{m}^3 \text{mol}^{-1}$). Note that while regressions between the T400 and FOS measurements were established with an intercept, this constant term does not intervene in Eq. 3 because it does not correlate with w' .

The overall uncertainty associated with the calibrated flux was estimated from Eq. 3:

$$\varepsilon_{F,cal} = \frac{\sqrt{(\beta_1 \cdot \varepsilon_{F,raw})^2 + (F_{raw} \cdot \varepsilon_{\beta_1})^2}}{v_{mol}} \quad (4)$$

where $\varepsilon_{F,cal}$ is the calibrated flux uncertainty ($\text{nmol m}^{-2} \text{s}^{-1}$), $\varepsilon_{F,raw}$ is the raw flux random error as estimated by Finkelstein and Sims (2001) (V m s^{-1}), and ε_{β_1} is the standard error of the slope estimate ($\text{nmol mol}^{-1} \text{V}^{-1}$). The first term under the square root represents the random flux error and the second term represents the calibration error. On average, the calibration error contributed less than 3% of the total uncertainty. This is in accordance with the observations made by (Horváth et al., 2017).

The procedure to establish the regressions between FOS and T400 data comprised four steps:

1. Data validation

FOS and T400 measurements were automatically validated or invalidated based on instrumental parameters such as pressure, flow rate, temperature, and UV lamp status (for the T400). Measurements from the T400 were discarded during the first 3 minutes following a mode switch (e.g., PROFILE or TRUNK to TOP, PROFILE to TRUNK) due to sensitivity to pressure variations. This delay allowed for pressure stabilization.

2. Data averaging

The FOS signal was averaged to match the T400's resolution at a 6 s timestamp. Unlike the 15-minute averaging used by Muller et al. (2010), this finer resolution captured more variability and increased the number of data points for regression. Both T400 and FOS time series were visually inspected to ensure data quality.

3. Data shifting



Table 2. Comparison of O₃ fluxes calibration methods.

	Calibration method				
	Ratio (RM)	Ratio Offset (ROM)	Ratio Variation (RVM)	Disc Calibration (DCM)	Period Calibration (PCM)
Authors	Müller 2010 Zhu 2015	Müller 2010 Zhu 2015	Zhu 2015	Müller 2010	This study
Period of comparison	30 min	30 min + disc	3 min	disc	user choice (1h up to disc lifetime)
Level of calibration	30 min	30 min	10 Hz	30 min	30 min
Offset considered	False	True	False	True	True
Shift of sensitivity considered	True	True & False	True	False	True
Sensitivity to white noise	Moderate	Moderate	High	Very low	Low

A time lag was observed between the two 6 s time series, attributed to the difference in tubing lengths (approximately 6 m for the FOS versus 60 m for the T400). To enhance regression quality, the FOS data was shifted by a time lag determined for each regression using covariance maximization. If a clear covariance maximum was not observed, a nominal time lag estimated for the entire measurement campaign was applied.

545 4. Regression optimization

Linear regressions were established using least-squares fits. Equation 3 was applied only when a satisfactory correlation ($R^2 > 0.5$) between relative (FOS) and absolute (T400) O₃ concentrations was achieved, as in Muller et al. (2010). Selecting the optimal regression period was a trade-off between maximizing the number of half-hour intervals meeting the R^2 criterion and minimizing calibration-associated uncertainty (Eq. 4). Initially, increasing the regression period
 550 (from 1 h up to 24 h) enhanced the R^2 values and reduced slope errors, as the larger dataset countered random fluctuations (white noise). However, beyond a certain period, shifts in the T400-FOS relationship over time reduced R^2 , while slope errors plateaued or increased. To balance these behaviours, different regression windows were tested, and for each year and system, an optimal period was selected (Sect. S3.5). Final regression slopes (β_1) and their associated uncertainty were then used to calibrate raw fluxes and compute the associated uncertainties.

555 2.5 Ancillary measurements

2.5.1 Other trace gas and particle measurements

Besides the high temporal resolution VOC and O₃ concentration and flux measurements, which are the main focus of this paper, numerous other in situ—both online and offline—measurements of short-lived climate forcers were conducted at the



nearby ISSeP air quality station. In addition, remote sensing measurements of NO₂, HCHO, and aerosols were carried out from the top of the tower. For completeness, these variables and their associated instrumentation are listed in Table 1.

Furthermore, during the period covered by the high-resolution VOC and ozone datasets, two pan-European Intensive Measurement Campaigns on VOCs and O₃ were organized.

The first campaign, coordinated by the EMEP Task Force on Measurements and Modelling (TFMM), took place from 12 to 19 July 2022 and focused on ozone formation under heatwave conditions. It was conducted in close collaboration with the ACTRIS and RI-URBANS European research infrastructures. Among other objectives, this campaign aimed at intensified VOC measurements at selected sites, including the BE-Vie site (BE0007R). In addition to continuous online PTR-ToF-MS measurements, ambient air was sampled daily around noon or in the early afternoon at the ISSeP air quality station for subsequent offline analysis in reference laboratories. Air collected in Silcosteel canisters was analysed by GC-MS, while VOCs adsorbed on TENAX-TA or HPLC cartridges were analysed by GC-MS and HPLC, respectively. More details on the analytical methods, as well as some preliminary data on NMHCs, OVOCs, and terpene concentrations at BE-Vie during this campaign, can be found in Fagerli et al. (2023).

A second EMEP/ACTRIS/RI-URBANS campaign, the “EUROpe-wide Intensive Campaign on Volatile Organic Compounds” (EUROVOC), was held in September 2024. This campaign aimed at gaining deeper insights into VOC emissions, including their temporal variability and chemical speciation, by deploying high-resolution measurements at sites located near emission sources (e.g., urban, industrial, traffic, harbour, and forest areas), including the Vielsalm site.

2.5.2 Sonic anemometer profiles

Raw data from the sonic anemometer profiles were processed using the EddyPro® software to compute key turbulence-related variables along the soil–canopy–atmosphere continuum. These include wind speed, wind direction, friction velocity (u^*), the standard deviation of the vertical wind component (σ_w), turbulent kinetic energy (TKE), Monin–Obukhov length, and atmospheric stability. The Lagrangian integral time scale (T_L), which is not included in EddyPro®, was calculated separately as the time integral of the autocorrelation function of $w(t)$, where w is the vertical wind component (Raupach, 1989). These variables—particularly σ_w and T_L —may be especially relevant for inferring the vertical positioning of sources and sinks from concentration profiles (Karl et al., 2004a; Leuning, 2000; Nemitz et al., 2000; Tiwary et al., 2007; Wada et al., 2020; Petersen et al., 2023).

2.5.3 ICOS data

As part of the ICOS network, a series of additional measurements were available at BE-Vie during the three measurement campaigns (Vincke et al., 2025). Turbulent fluxes of CO₂ and H₂O were measured at 51 m a.g.l. using an infrared gas analyser (LI-7200, LI-COR, Lincoln, NE, USA) coupled with the HS-50 sonic anemometer described previously.

Following the ICOS protocol, a comprehensive set of meteorological measurements was collected at BE-Vie, both above and within the canopy. These included air temperature and humidity, atmospheric pressure, incoming and outgoing longwave



and shortwave radiation, incoming and outgoing photosynthetic photon flux density (PPFD), soil temperature and moisture, precipitation, and CO₂ and H₂O mixing ratio profiles along the tower.

Vegetation development was monitored using images captured every three days by a StarDot NetCam SC (PhenoCam) installed at the top of the flux tower and oriented towards the north-west sector (mixture of beech and Norway spruce). These images were processed using the vegindex Python package (<https://python-vegindex.readthedocs.io/en/latest/>) to extract the Green Chromatic Coordinate (GCC), also known as the "greenness index." Seasonal variations in the GCC provide a reliable proxy for vegetation onset and senescence. Further methodological details are provided in Seyednasrollah et al. (2019).

3 Results and discussion

3.1 Meteorological and phenological conditions

Throughout the three measurement campaigns, certain meteorological and phenological events stand out. While mean diel evolutions appear broadly similar across the years, some distinct year-to-year features are also apparent (Fig. 5).

In 2022, a heatwave that lasted eight consecutive days was observed from August 9 to August 16. This period was characterized by higher temperatures and radiation compared to the other years, and by lower relative humidity, partly due to very low precipitation (data not shown). From early December of the same year, temperatures were consistently below 0 °C for an extended period of days—a phenomenon not observed in the other years.

The year 2023 was marked by the two hottest months ever recorded in Belgium (June and September). At the end of May and beginning of June, a month-long period of anticyclonic conditions brought warm and dry air from the north-east sector.

Latent heat and carbon fluxes followed the expected patterns for a mixed temperate forest. Photosynthesis (Fig 5f) and transpiration (Fig 5e) were enhanced by the increase in temperature and radiation during the transition from spring to summer. The evolution of GCC (Fig 5g) served as a proxy for leaf expansion in deciduous trees (*Fagus sylvatica*). A strong development of vegetation was observed between April 24 and May 8. Visual inspection of images captured from the top of the flux tower indicated that budburst occurred around April 28, with little variation between years. In 2022, the onset of leaf fall was estimated between October 20 and October 26. Warmer conditions in 2023 delayed leaf fall compared to 2022, with observations indicating a period between November 2 and November 8. Phenological images were not available for 2024 to determine the leaf fall period.

3.2 Diversity of VOCs

3.2.1 Number of detected VOCs

In total, 76 m/z values were retained by the DBSCAN algorithm. These ion species are listed in Table S3, along with their chemical formula, tentative compound identification, and parameters related to sensitivity computation.

It is important to note that compounds can give rise to signals at multiple ion masses, which may correspond either to isotopes associated with the chemical formula or to ion species resulting from the fragmentation of nascent, excited protonated




Figure 5. Average diel patterns (in LT) of meteorological and phenological conditions for each two-week period. Solid lines represent hourly means, and shaded areas indicate the standard deviation around the mean. **(a)** Air temperature. **(b)** Incoming global radiation. **(c)** Relative humidity. **(d)** Wind direction. **(e)** Latent heat flux. **(f)** Net CO₂ flux. **(g)** Greenness index. All variables were measured at the top of the flux tower (TOP system). The greenness index (GCC) was measured once per day, so no diel pattern is shown for this variable and a single mean is provided for each two-week period.

molecules (i.e., fragment ions). When reporting total VOC fluxes or concentrations, it is crucial to consider only one ion per compound to avoid double counting.

In general, isotopes were not taken into account and have been excluded when compiling the list of ion masses considered (Table S2). One notable exception is benzene (m/z 79.054), whose peak is entangled with that of protonated hydrated acetic acid (m/z 79.039). At their respective isotope masses (m/z 80.042 and 80.058), this entanglement is reduced due to the relatively higher abundance of the benzene isotope. As a result, only the benzene isotope signal was considered for benzene quantification, eliminating any risk of double counting.

In contrast, both protonated molecules and their fragment ions are listed in Table S2 and were included when representing the diversity of detected ion species (Fig. 6). However, for the summed fluxes shown in Figs. 8 and 9, only one signal was taken into account for each compound. To assure the best possible quantification, we used the mass related to the highest PT/fragment yields (Table S3), ending up with 65 ions used for the quantification of total VOC exchange.

Among the 76 detected m/z values, 57 ion species were identified as significantly exchanged at the top of the tower, based on the three-step comparison of fluxes with LOD_f described in Sect. 2.3.5. The algorithm performed well in identifying ions with clear flux dynamics, with only a few exceptions: two m/z values were found by the algorithm but displayed erratic flux



patterns (m/z 97.028 ($C_5H_4O_2H^+$) and m/z 107.086 ($C_8H_{10}H^+$)), while five others exhibited consistent diurnal trends but were not detected by the algorithm (m/z 97.065 ($C_6H_8OH^+$), 99.044 ($C_5H_6O_2H^+$), 101.023 ($C_4H_4O_3H^+$), 103.075 ($C_5H_{10}O_2H^+$), and 115.075 ($C_6H_{10}O_2H^+$)). The latter were manually included, and the former excluded, resulting in a final list of 60 m/z values considered significantly exchanged for the TOP system. When removing multiples ions related to a same compound
 640 (fragments and protonated compounds), we ended up with a list of 51 significantly exchanged compounds.

This number is intermediate in comparison to other studies using a PTR-ToF-MS instrument. Some reported hundreds of ions with significant fluxes: 494 over an orange orchard in California (Park et al., 2013a); 377 above a mixed temperate forest in the USA (Millet et al., 2018); around 200 with a PTR3-ToF-MS above a boreal forest in Hyytiälä, Finland (Fischer et al., 2021); and 123 over a winter wheat field in France (Loubet et al., 2022). In contrast, other studies reported lower numbers
 645 than in the present work: 29 in a deciduous forest in Northern Italy (Schallhart et al., 2016); between 10 and 20 over a similar forest in the same region (Jensen et al., 2018); 25 above a boreal forest in Finland (Schallhart et al., 2018); and 18 compounds distributed over 43 ion species at a grassland site in Austria during harvest (Ruuskanen et al., 2011). These large differences can be explained by the sensitivity of the PTR-ToF-MS instrument but also by the ecosystem type—for example, an orange orchard may emit substantially more VOCs than a temperate forest (Loubet et al., 2022). However, even for similar ecosystems
 650 and climatic conditions, Millet et al. (2018) reported six times more significant compounds than in the present study. This raises questions not only about instrumental capabilities, but also about the criteria used to determine whether a flux is considered significant or not (Schallhart et al., 2016; Jensen et al., 2018).

For the TRUNK fluxes, only 31 m/z values were identified as significantly exchanged. No visual inspection was applied to adjust this list, as the TRUNK flux dynamics were more difficult to assess visually.

655 3.2.2 Order of magnitudes

The mass range of the 76 detected ions, as well as the range of average mixing ratios and mean absolute fluxes of the tentatively associated compounds, are illustrated in Fig. 6. It must be noted that in this study, VOC fluxes are reported in mass units ($\mu g\ m^{-2}\ s^{-1}$), allowing for straightforward aggregation and carbon budget assessments.

The PTR-ToF-MS instrument was able to detect and quantify protonated VOCs with mass-to-charge ratios ranging from
 660 31.018 (formaldehyde) to 205.195 (sesquiterpenes). In terms of average mixing ratios, concentrations span several orders of magnitude from approximately 3 ppbv for methanol down to pptv level for the least abundant compounds. Mixing ratios measured at the TRUNK level remain in the same range, although somewhat smaller than the ones acquired at the top of the flux tower. The top-left area of Fig. 6 is occupied by low molecular weight oxygenated compounds which are listed in Table 3: methanol (m/z 33.033), acetic acid and its fragment (m/z 61.028 and 43.018 respectively), acetone (m/z 59.049), ethanol
 665 (m/z 47.049), formaldehyde (m/z 31.018), formic acid (m/z 47.013), and acetaldehyde (m/z 45.033). These present the highest mixing ratios, across all seasons, although lower in autumn.

VOC concentrations show a clear decreasing trend with increasing m/z , both above the canopy (TOP system, panel a) and near ground level (TRUNK system, panel b). This pattern largely reflects the lower volatility of heavier compounds, which




Figure 6. Mean VOC mixing ratio over the whole campaign (in log scale) as a function of mass-to-charge ratio m/z for the 76 detected ions. **(a)** TOP system. **(b)** TRUNK system. For each m/z , the circle size is weighted by the corresponding absolute mean flux. The eight most exchanged ions (in terms of absolute mean flux) are highlighted on the graph.

limits their ability to remain in the gas phase and accumulate in the atmosphere. Additionally, higher reaction rate constants
 670 with atmospheric oxidants contribute to the reduced concentrations of heavier BVOCs.

Local emissions also influence this pattern, as suggested by the circle sizes. Most of the strongest emissions correspond
 to ions with m/z values below 80, while heavier compounds tend to show weaker emissions. Despite the overall trend, some
 compounds deviate from the expected decrease, exhibiting higher concentrations than those associated with neighbouring m/z
 values. Notable examples include m/z 137.132 (sum of monoterpenes) and m/z 135.115 ($\text{C}_{10}\text{H}_{14}\text{H}^+$, potentially p-cymene).
 675 These anomalies are associated with high fluxes, indicating an effect of local emissions on ambient VOC concentrations.

The seasonal dynamics of VOC ambient concentrations and fluxes will be further described in Sect. 3.3 and 3.4, respectively.



Table 3: Seasonal mean fluxes and concentrations of VOCs and ozone measured at the TOP system during 2022, 2023, and 2024. Only VOC ions exhibiting significant exchange are reported and listed in descending order of flux magnitude. A tentative compound attribution is provided for ions with relatively high confidence. For others, candidates are listed in Table S3. Note that ozone fluxes are reported in $\text{nmol m}^{-2} \text{s}^{-1}$ as in the rest of the manuscript, while VOC fluxes were transformed from $\mu\text{g m}^{-2} \text{s}^{-1}$ to $\text{ng m}^{-2} \text{s}^{-1}$ to improve scaling.

Ion m/z (Da)	Tentative attribution	Mean flux 24h (day/night)			Mean mixing ratio 24h		
		Spring	Summer	Autumn	Spring	Summer	Autumn
				nmol m ⁻² s ⁻¹	ppbv		
	O ₃ ozone	-6.33 (-8.96/-3.69)	-6.55 (-9.42/-3.77)	-5.43 (-6.48/-4.26)	37.8	40.5	30.4
				ng m ⁻² s ⁻¹	ppbv		
81.070	C ₆ H ₈ H ⁺ fragm. monoterpene	58.7 (86.1/27.8)	134 (211/50.1)	31.6 (42.8/19.7)	0.219	0.353	0.187
137.132	C ₁₀ H ₁₆ H ⁺ monoterpenes	43.5 (63.1/21.0)	115 (183/39.9)	23.7 (32.0/14.8)	0.149	0.257	0.134
69.070	C ₅ H ₈ H ⁺ isoprene	28.3 (49.4/6.31)	75.3 (136/14.4)	13.0 (24.8/1.17)	0.128	0.272	0.0773
33.033	CH ₄ OH ⁺ methanol	34.5 (56.7/11.5)	11.6 (23.4/-0.487)	-12.1 (-9.86/-14.5)	4.88	4.47	1.50
41.039	C ₃ H ₄ H ⁺ fragm. isoprene	19.7 (33.6/5.15)	46.9 (84.6/9.72)	7.17 (15.0/-0.776)	0.404	0.532	0.288
47.049	C ₂ H ₆ OH ⁺ ethanol	8.61 (14.8/2.15)	5.25 (10.1/0.416)	-11.1 (-9.08/-13.3)	1.55	1.46	0.973
135.117	C ₁₀ H ₁₄ H ⁺ p-cymene	8.98 (14.1/3.88)	20.5 (32.8/8.01)	6.33 (8.35/4.29)	0.156	0.222	0.152
59.049	C ₃ H ₆ OH ⁺ acetone	13.9 (21.0/6.33)	15.4 (24.0/6.57)	-4.19 (-1.84/-6.60)	1.61	2.10	1.12
43.018	C ₂ H ₂ OH ⁺ fragm. acetic acid	4.13 (7.33/0.786)	5.46 (10.3/0.701)	-5.93 (-5.97/-5.90)	1.60	2.39	0.813
61.028	C ₂ H ₄ O ₂ H ⁺	5.05	3.53	-3.60	1.50	1.76	0.577

Table 3 (continued)



Ion m/z (Da)	Tentative attribution	Mean flux 24h (day/night)			Mean mixing ratio 24h		
		Spring	Summer	Autumn	Spring	Summer	Autumn
		ng m ⁻² s ⁻¹			ppbv		
80.058	acetic acid	(6.97/3.00)	(5.04/2.04)	(-4.07/-3.12)			
	¹³ C ¹² C ₅ H ₆ H ⁺	1.82	5.81	1.19	0.120	0.144	0.152
31.018	benzene isotope	(3.46/0.215)	(9.53/2.06)	(1.52/0.857)			
	CH ₂ OH ⁺	-5.33	-6.63	-6.75	1.33	1.48	0.641
45.033	formaldehyde	(-7.56/-2.98)	(-9.36/-3.87)	(-8.25/-5.20)			
	C ₂ H ₄ OH ⁺	6.03	4.77	-2.42	0.651	0.707	0.379
149.132	acetaldehyde	(9.77/2.05)	(7.66/1.87)	(-1.87/-2.99)			
	C ₁₁ H ₁₆ H ⁺	2.35	2.89	1.12	0.0154	0.0149	0.0146
93.070	fragm. SQT	(3.56/1.02)	(4.64/1.10)	(1.40/0.821)			
	C ₇ H ₈ H ⁺	3.39	5.77	1.76	0.0854	0.0971	0.0845
95.086		(4.80/1.94)	(8.74/2.72)	(2.30/1.21)			
	C ₇ H ₁₀ H ⁺	3.22	7.43	1.81	0.0225	0.0375	0.0181
47.013		(4.80/1.52)	(12.1/2.60)	(2.44/1.15)			
	CH ₂ O ₂ H ⁺	0.863	0.957	-0.342	0.636	0.773	0.279
83.086	formic acid	(1.76/-0.00507)	(2.14/0.000928)	(-0.616/-0.0860)			
	C ₆ H ₁₀ H ⁺	5.38	3.94	0.882	0.0568	0.0570	0.0271
77.023	fragm. C ₆ H ₁₂ OH ⁺	(7.90/2.74)	(5.87/1.97)	(1.26/0.498)			
	C ₂ H ₄ O ₃ H ⁺	-2.94	-3.40	-1.78	0.119	0.162	0.0511
77.060		(-4.25/-1.61)	(-4.73/-2.07)	(-2.25/-1.30)			
	C ₃ H ₈ O ₂ H ⁺	2.32	2.53	-0.0964	0.109	0.131	0.0654
67.054		(3.79/0.787)	(4.19/0.893)	(0.298/-0.490)			
	C ₅ H ₆ H ⁺	1.72	3.89	0.836	0.0196	0.0325	0.0137
57.070		(2.75/0.616)	(6.59/1.06)	(1.35/0.316)			
	C ₄ H ₈ H ⁺	1.70	2.04	0.505	0.165	0.169	0.139
73.065		(2.67/0.705)	(3.34/0.754)	(0.876/0.133)			
	C ₄ H ₈ OH ⁺	0.933	1.26	-1.14	0.153	0.159	0.0967
205.195	MEK	(1.56/0.284)	(2.10/0.430)	(-0.991/-1.29)			
	C ₁₅ H ₂₄ H ⁺	1.36	1.67	0.757	0.00632	0.00842	0.00696

Table 3 (continued)



Ion m/z (Da)	Tentative attribution	Mean flux 24h (day/night)			Mean mixing ratio 24h		
		Spring	Summer	Autumn	Spring	Summer	Autumn
		ng m ⁻² s ⁻¹			ppbv		
	sesquiterpenes	(2.06/0.550)	(2.52/0.754)	(0.952/0.557)			
71.049	C ₄ H ₆ OH ⁺	1.26	0.930	-0.390	0.0845	0.253	0.0638
	MVK + MACR	(2.11/0.394)	(1.98/-0.0362)	(-0.0627/-0.719)			
87.044	C ₄ H ₆ O ₂ H ⁺	0.951	1.07	-0.157	0.0825	0.119	0.0411
		(1.51/0.362)	(1.63/0.489)	(0.0321/-0.351)			
87.080	C ₅ H ₁₀ OH ⁺	1.88	2.08	0.0286	0.0547	0.0543	0.0318
		(3.00/0.726)	(3.31/0.840)	(0.246/-0.190)			
151.112	C ₁₀ H ₁₄ OH ⁺	0.480	1.03	0.280	0.0284	0.0377	0.0172
	fragm. C ₁₀ H ₁₆ O ₂ H ⁺	(0.784/0.147)	(1.51/0.502)	(0.334/0.224)			
85.101	C ₆ H ₁₂ H ⁺	0.828	0.792	0.260	0.0255	0.0265	0.0210
	fragm. C ₆ H ₁₄ OH ⁺	(1.24/0.395)	(1.16/0.417)	(0.353/0.165)			
101.060	C ₅ H ₈ O ₂ H ⁺	1.23	1.39	0.121	0.0648	0.0808	0.0238
		(2.05/0.368)	(2.23/0.536)	(0.322/-0.0829)			
89.060	C ₄ H ₈ O ₂ H ⁺	0.502	0.550	-0.529	0.0833	0.0962	0.0434
		(0.804/0.184)	(1.02/0.0741)	(-0.410/-0.648)			
137.060	C ₈ H ₈ O ₂ H ⁺	0.128	0.643	0.152	0.00868	0.00947	0.00474
		(0.248/0.00346)	(1.18/0.0849)	(0.181/0.121)			
65.023	CH ₄ O ₃ H ⁺	0.531	0.623	0.0751	0.0498	0.0621	0.0210
		(0.989/0.0649)	(1.14/0.120)	(0.154/-0.00441)			
115.075	C ₆ H ₁₀ O ₂ H ⁺	0.305	0.608	0.0859	0.0313	0.0391	0.0133
		(0.583/0.0217)	(1.01/0.201)	(0.151/0.0203)			
99.080	C ₆ H ₁₀ OH ⁺	0.920	0.860	0.103	0.0308	0.0352	0.0150
		(1.39/0.422)	(1.34/0.365)	(0.227/-0.0231)			
101.023	C ₄ H ₄ O ₃ H ⁺	-0.0604	0.110	-0.00878	0.0297	0.0338	0.0112
		(-0.122/0.00301)	(0.253/-0.0330)	(0.00323/-0.0209)			
123.117	C ₉ H ₁₄ H ⁺	1.49	0.841	0.233	0.0117	0.0145	0.00606
		(2.33/0.629)	(1.32/0.347)	(0.306/0.159)			
75.044	C ₃ H ₆ O ₂ H ⁺	0.370	0.336	-0.348	0.0812	0.123	0.0380

Table 3 (continued)



Ion m/z (Da)	Tentative attribution	Mean flux 24h (day/night)			Mean mixing ratio 24h		
		Spring	Summer	Autumn	Spring	Summer	Autumn
		ng m ⁻² s ⁻¹			ppbv		
71.086	hydroxyacetone	(0.566/0.164)	(0.593/0.0781)	(-0.312/-0.386)			
	C ₅ H ₁₀ H ⁺	0.493	0.732	0.163	0.0288	0.0314	0.0240
		(0.769/0.207)	(1.12/0.330)	(0.248/0.0772)			
121.101	C ₉ H ₁₂ H ⁺	0.454	0.466	0.162	0.0188	0.0237	0.0209
		(0.708/0.194)	(0.769/0.162)	(0.232/0.0922)			
97.065	C ₆ H ₈ OH ⁺	0.303	0.488	0.0780	0.0172	0.0229	0.00966
		(0.520/0.0776)	(0.761/0.212)	(0.126/0.0284)			
101.096	C ₆ H ₁₂ OH ⁺	0.924	0.704	0.178	0.0143	0.0153	0.00827
		(1.33/0.504)	(0.977/0.424)	(0.254/0.102)			
83.049	C ₅ H ₆ OH ⁺	0.615	0.867	0.228	0.0267	0.0489	0.0144
	3-methylfuran	(0.999/0.211)	(1.37/0.359)	(0.340/0.113)			
42.034	C ₂ H ₃ NH ⁺	-0.0605	-0.131	-0.774	0.0801	0.129	0.104
	acetonitrile	(-0.0549/-0.0664)	(-0.0615/-0.201)	(-0.805/-0.743)			
111.117	C ₈ H ₁₄ H ⁺	0.343	0.390	0.128	0.00395	0.00434	0.00224
	fragm. C ₈ H ₁₆ OH ⁺	(0.525/0.157)	(0.627/0.153)	(0.179/0.0767)			
133.101	C ₁₀ H ₁₂ H ⁺	0.298	0.537	0.172	0.00637	0.00829	0.00540
	p-cymenene	(0.447/0.140)	(0.876/0.181)	(0.232/0.110)			
141.127	C ₉ H ₁₆ OH ⁺	0.745	0.308	0.0723	0.00618	0.00629	0.00234
		(1.19/0.300)	(0.513/0.104)	(0.103/0.0402)			
139.112	C ₉ H ₁₄ OH ⁺	0.393	0.760	0.126	0.0174	0.0407	0.0110
		(0.655/0.121)	(1.33/0.191)	(0.193/0.0588)			
49.028	CH ₄ O ₂ H ⁺	-0.765	-0.692	-0.492	0.0265	0.0251	0.0115
		(-1.16/-0.361)	(-1.02/-0.366)	(-0.633/-0.347)			
143.143	C ₉ H ₁₈ OH ⁺	0.926	0.498	0.0485	0.00648	0.00534	0.00157
	nonanal	(1.54/0.320)	(0.783/0.213)	(0.0871/0.00944)			
97.101	C ₇ H ₁₂ H ⁺	0.700	0.617	0.141	0.00791	0.00891	0.00460
	fragm. C ₇ H ₁₄ OH ⁺	(1.08/0.303)	(0.922/0.311)	(0.218/0.0631)			
53.040	C ₄ H ₄ H ⁺	0.252	0.477	0.107	0.0103	0.0143	0.00852

Table 3 (continued)



Ion m/z (Da)	Tentative attribution	Mean flux 24h (day/night)			Mean mixing ratio 24h		
		Spring	Summer	Autumn	Spring	Summer	Autumn
		ng m ⁻² s ⁻¹			ppbv		
99.044	C ₅ H ₆ O ₂ H ⁺	(0.410/0.0883)	(0.827/0.124)	(0.188/0.0243)	0.0188	0.0337	0.00984
		0.290	0.410	0.0286			
117.091	C ₆ H ₁₂ O ₂ H ⁺	(0.454/0.122)	(0.677/0.142)	(0.0751/-0.0188)	0.0104	0.0106	0.00458
		0.268	0.285	0.00694			
153.127	C ₁₀ H ₁₆ OH ⁺ camphor	(0.419/0.114)	(0.434/0.134)	(0.0495/-0.0360)	0.00767	0.0124	0.00774
		0.366	0.677	0.227			
153.055	C ₈ H ₈ O ₃ H ⁺ methyl salicylate	(0.549/0.169)	(1.08/0.264)	(0.312/0.140)	0.0124	0.0105	0.00174
		0.727	0.648	0.0327			
103.075	C ₅ H ₁₀ O ₂ H ⁺	(1.23/0.216)	(1.12/0.177)	(0.0661/-0.000800)	0.0141	0.0127	0.00481
		0.250	0.230	-0.00872			
63.044	C ₂ H ₆ O ₂ H ⁺ 1,2-ethanediol	(0.364/0.134)	(0.350/0.111)	(0.0190/-0.0368)	0.0246	0.0296	0.0123
		0.534	0.496	-0.0562			
115.112	C ₇ H ₁₄ OH ⁺	(0.862/0.190)	(0.816/0.167)	(0.0122/-0.125)	0.00620	0.00701	0.00307
		0.306	0.298	0.0408			
85.065	C ₅ H ₈ OH ⁺	(0.481/0.124)	(0.442/0.152)	(0.0751/0.00609)	0.0146	0.0178	0.00706
		0.320	0.381	0.0230			
		(0.507/0.126)	(0.619/0.139)	(0.0890/-0.0436)			

3.3 Ambient concentrations

Although the present study focuses on biogenic VOC and ozone exchange, the analysis of ambient concentrations provides valuable insights into some of the mechanisms driving the observed fluxes. Among the many measured compounds, a selection of chemical species with characteristic diel patterns is presented in Fig. 7. These exhibit pronounced cycles, albeit with differing phases and amplitudes, likely reflecting distinct sources, sinks, and reactivity. Additionally, their vertical concentration profiles may help identify sources and sinks at different heights.

Across seasons, average ozone concentrations ranged between 20 and 60 ppbv (Fig.7b), consistent with values reported by Vermeuel et al. (2021) in a mixed temperate forest. Concentrations peaked in summer (June) and reached a minimum in autumn. This seasonal behaviour, along with the intra-daily variations described below, confirms the photochemical origin of ozone, which depends on solar radiation and temperature (Pellegrini et al., 2007). During the day, O₃ mixing ratios increase progressively after sunrise and peak in mid-afternoon (typically 16:00–18:00 LT), driven by photochemical production in-



volving NO_2 photolysis and reactions with biogenic VOCs (Stockwell and Forkel, 2002; Faiola et al., 2024). After peaking, concentrations decline steadily into the evening, due to the cessation of photochemistry and titration by NO, along with ongoing deposition. In a maritime pine forest, Kammer et al. (2018) observed nighttime ozone levels dropping below the detection limit (0.5 ppbv), suggesting efficient removal by highly reactive and abundant BVOCs. A similar phenomenon was reported by Andersen et al. (2024), but attributed to thermal inversion, which limited ozone replenishment from above; in this case, chemical loss alone was considered insufficient to explain the low nighttime ozone levels. In the present study, however, nighttime O_3 levels remained relatively high (on average between 10 and 20 ppbv).

Ozone concentrations were generally lowest near the ground. Vertical stratification was especially evident in the morning and evening, when atmospheric mixing was reduced. The decrease near ground level may be attributed to continuous deposition from the canopy top downward (Zhou et al., 2017a), which was also observed at night in this study (Sect. 3.5). Furthermore, reduced photochemical production due to radiation attenuation within the canopy likely contributes to lower ozone levels close to the ground during the day.

In contrast, NO_x concentrations (Fig. 7a) showed a nocturnal pattern, peaking at night and in early morning. This accumulation likely results from soil NO emissions (Min et al., 2014) in combination with a stable nocturnal boundary layer that limits vertical mixing. As NO_x measurements were conducted at the ISSeP station, vertical concentration profiles are not available in this dataset, preventing direct verification of this hypothesis. After sunrise, NO_x levels decrease rapidly due to oxidation and dilution, occurring concurrently with ozone production. The amplitude of average NO_x concentrations ranged from 0 to a maximum of 5 ppbv, confirming the rural character of the site, with limited seasonal variation except in December, when peak values were observed—possibly due to reduced NO_2 photolysis or increased wintertime anthropogenic NO emissions.

Although chemically distinct from NO_x , many VOCs displayed similar diel behaviour, characterized by nighttime accumulation followed by a daytime minimum. These include ethanol (m/z 47.049) as well as heavier compounds, potentially including xylenes (m/z 107.086), trimethylbenzene (m/z 121.101), p-cymene (m/z 135.117), nonanal (m/z 143.143), and camphor (m/z 153.127). This pattern is illustrated for ethanol in Fig. 7d. The observed nighttime increase may reflect emissions under limited mixing conditions, supported by the stratification observed in vertical profiles. The daytime decrease, however, is more difficult to interpret, especially given the upward fluxes of ethanol measured during the day (Fig. 8d) and its relatively low reactivity (estimated atmospheric lifetime of 2.8 days, (Naik et al., 2010)). One possible explanation is that background concentrations are sufficiently high that local emissions do not result in noticeable concentration increases.

In contrast to ethanol, methanol concentrations increased after sunrise (around 06:00 LT) and peaked in the evening before declining at night (Fig. 8c). The diel patterns of both OVOCs are broadly similar, but offset in time. This timing difference may arise from differing emission dynamics—methanol being among the most strongly exchanged VOCs (Table 3)—and differences in atmospheric reactivity (methanol's estimated lifetime of 4.7 days (Millet et al., 2008)). Both compounds also showed decreasing vertical concentrations from above-canopy to ground level at night.

Methanol and ethanol concentrations were highest in May, whereas isoprene (Fig. 8e), methyl vinyl ketone (MVK) and methacrolein (MACR) (Fig. 8f), and monoterpenes (Fig. 8g) peaked in June, July, and August. These differences likely reflect distinct emission processes, further discussed in Sect. 3.4.



Isoprene and its oxidation products (MVK + MACR) exhibited diurnal profiles similar to that of ozone. Isoprene concentrations followed the expected light- and temperature-dependent emission patterns (Guenther et al., 2012), but showed a slight
 725 delay compared to solar radiation and isoprene fluxes. Peak concentrations typically occurred in the early afternoon (between 13:00 and 16:00 LT), whereas incoming radiation peaked around noon, isoprene fluxes near 13:00 LT, and air temperature somewhat later (15:30–16:00 LT). This pattern suggests that isoprene emissions (fluxes) closely track solar radiation, while atmospheric concentrations respond with a delay of a few hours following emissions. Vertically, maximum isoprene concentrations were not observed at the top of the flux tower (51 m a.g.l.) but rather near the canopy top, around 31 m a.g.l., indicating
 730 that emissions were primarily localized within or just above the canopy. From November onward, isoprene concentrations declined sharply, consistent with the strong seasonality of its emissions.

MVK + MACR differed from isoprene in timing and vertical distribution. Their concentrations peaked approximately 0 to 3 hours after those of isoprene, consistent with their secondary formation through isoprene oxidation (estimated isoprene lifetime of ~ 1.4 h, based on an average daytime OH concentration of $2 \cdot 10^6$ molec. cm^{-3} (Jensen et al., 2018)). While isoprene
 735 concentrations were highest near the top of the canopy, MVK and MACR peaked slightly higher, around 51 m a.g.l., near the top of the measurement profile. This vertical offset may reflect enhanced isoprene oxidation above rather than inside the canopy.

Finally, monoterpenes displayed less pronounced diel patterns compared to the other species. Their emissions, which are predominantly temperature-driven (Guenther et al., 2012), also occurred at night. This may contribute to the lower diel variability in their mixing ratios relative to isoprene. Uniquely, monoterpenes reached maximum concentrations near ground level,
 740 a pattern also observed in a boreal forest by Petersen et al. (2023). This could reflect continuous nighttime emissions that accumulate within a shallow nocturnal boundary layer. Around sunrise (06:00–07:00 LT), concentrations increased at higher levels, possibly due to the onset of turbulent mixing.

While the above interpretations offer plausible explanations for the observed diel and vertical patterns, they remain limited by the complexity of the processes involved. In particular, ambient concentration dynamics results from the interplay between
 745 emissions, depositions, atmospheric transport, and chemical transformations. Disentangling the different factors affecting this type of dataset require either methods to isolate specific features or more integrative approaches. For instance, to isolate where sources/sinks are located within the forest canopy, one can use Lagrangian modelling in combination with vertical concentration and turbulence profile measurements (Karl et al., 2004b; Tiwary et al., 2007; Wada et al., 2020; Petersen et al., 2023). Source apportionment studies can identify the impact of specific sources on local concentrations by using positive matrix
 750 factorisation (Song et al., 2024), by looking at meso-scale transport affect the concentration dataset (Simon et al., 2023), or by using a combination of both (Verreyken et al., 2021). The disadvantage of matrix factorisation is that chemical degradation is not explicitly taken into account. To overcome this, chemically aged masses related to a specific source can be designated as separate factors and ageing regarded as a linear combination between both (Verreyken et al., 2021; Song et al., 2024). To identify the impact of chemistry on trace gas concentrations, one can use atmospheric chemistry models in various setups.
 755 Simple 0-D chemistry models can quantify chemical losses occurring at a specific site (Andersen et al., 2024). However, when characterizing a forest canopy, more integrative 1-D canopy exchange models (Ashworth et al., 2016; Zhou et al., 2017b) better represent the various processes occurring in the ecosystem. Unfortunately, 1-D canopy exchange models often neglect




Figure 7. Average diel evolution (in LT) of vertical concentration profiles of selected trace gases for each month of the 2022, 2023, and 2024 field campaigns. (a) Nitrogen oxides ($\text{NO} + \text{NO}_2$). (b) Ozone. (c) Methanol. (d) Ethanol. (e) Isoprene. (f) MVK + MACR. (g) Sum of monoterpenes. Median values of 1-hour aggregated data were computed, then interpolated horizontally (over time) and vertically (over height). The dashed line represents the median concentration measured at the TOP system.

or strongly simplify the effect of advection of plumes originating from sources outside the forest. To get a broader picture of chemistry processes, the use of 3-D chemistry and transport models (Poraicu et al., 2025) are valuable tools for constraining the regional BVOC budget. However, they often lack a detailed representation of forest canopy structure, which can limit their accuracy in simulating in-canopy mechanisms. Altogether, combining multiple approaches remains essential to fully capture the complexity of trace gas dynamics within and above forest canopies.

3.4 VOC fluxes

3.4.1 Flux balance

BVOCs can both be emitted by the ecosystem (positive fluxes) and removed from the atmosphere through deposition (negative fluxes) via different biological, physical and chemical pathways. The total ecosystem exchange resulting from the balance between these two processes is illustrated in Fig. 8, across eight representative two-week periods.

Note that only the 65 unique ions, representing each a single compound, were considered in this figure to avoid skewing the mass balance. The description below focuses first on the year 2023, for which the measurement coverage was the most continuous.

During spring and summer, the net BVOC exchange was largely dominated by emissions (panels f, g, and h), with a peak in mid-June (panel h), reaching a maximum mean value of $\sim 1.25 \mu\text{g m}^{-2} \text{s}^{-1}$ —more than three times higher than in May.



Similar magnitudes were observed in July and August (not shown, but comparable to panel h). These maxima correspond to $\sim 0.9 \mu\text{g C m}^{-2} \text{ s}^{-1}$ (carbon mass basis) or $\sim 18 \text{ nmol m}^{-2} \text{ s}^{-1}$ (molar basis). This value is twice as high as the summed
 775 flux of 494 detected ions in an orange orchard in summer (Park et al., 2013a). This difference is attributed to isoprene and monoterpenes, which contributed to most of the exchange in the present study while being very low in the orange orchard. Our maximum total flux of $18 \text{ nmol m}^{-2} \text{ s}^{-1}$ is closer to ranges reported by (Schallhart et al., 2016) in a Mediterranean oak-hornbeam forest (between 25 and $30 \text{ nmol m}^{-2} \text{ s}^{-1}$), where isoprene emissions were dominant and much higher.

Expressing VOC fluxes on a carbon mass basis can be useful for assessing their contribution to atmospheric carbon. For
 780 comparison, daytime CO_2 uptake was about $20 \mu\text{mol m}^{-2} \text{ s}^{-1}$ which is equivalent to $240 \mu\text{g C m}^{-2} \text{ s}^{-1}$, such that the total carbon emitted as VOCs represents only 0.37% of the carbon captured by the ecosystem as CO_2 .

From September onward, the observed net fluxes decreased and approached a balance between emissions and depositions (panels i and j). This seasonal transition is attributed both to meteorological factors—namely decreasing temperature and solar radiation, which are key drivers of BVOC production—and to phenological changes such as vegetation senescence. Although
 785 Fig. 8 might suggest that downward fluxes were enhanced in autumn, they varied little across seasons and remained close to $\sim -0.1 \mu\text{g m}^{-2} \text{ s}^{-1}$ or $\sim -0.05 \mu\text{g C m}^{-2} \text{ s}^{-1}$ or $\sim -2 \text{ nmol m}^{-2} \text{ s}^{-1}$. These seemingly constant depositions are probably the result of compensating mechanisms. Lower mixing ratios and reduced stomatal aperture in autumn compared to spring (Fig. 7) would be expected to decrease stomatal VOC uptake (Niinemets et al., 2014). However, higher autumn humidity may enhance the deposition of water-soluble VOCs onto canopy water films followed by a (bacterial) consumption process (Laffineur et al.,
 790 2012), potentially compensating for the reduced stomatal uptake. This implies that downward VOC fluxes are less sensitive to seasonal variation than emissions, though a comprehensive modelling approach is needed to disentangle these effects.

At the diurnal scale, transitions from net uptake during the night to net release during the day are also evident. This diel pattern was particularly clear at the start of the 2023 campaign (panel f). During the night and early morning, depositions were observed for small oxygenated compounds such as methanol (m/z 33.033), ethanol (47.049), acetone (59.049), acetic
 795 acid (61.028), acetaldehyde (45.033), and formic acid (47.013). As shown in Fig. 5c, humidity was higher at night, creating conditions favouring the adsorption of such compounds on water films. Higher concentrations found at night (Fig. 7) could also have induced downward fluxes, although stomatal closure exerts a strong control on the stomatal sink. At night, isoprene flux (m/z 69.070), which is light-dependent, was not detected in the absence of solar radiation. In contrast, temperature-driven compounds such as monoterpenes (m/z 137.132) and m/z 135.117 ($\text{C}_{10}\text{H}_{14}\text{H}^+$, potentially p-cymene) were still emitted at
 800 night, as illustrated in panels i and j.

From around 07:00 LT, total emissions increased, primarily driven by monoterpenes and isoprene, which were the dominant daytime BVOC sources in this ecosystem. Concurrently, net fluxes of small oxygenated BVOCs switched from negative to positive shortly after sunset. Niinemets et al. (2014) describes this transition as the result of the outgassing of VOCs from water films to the gas phase and stomatal opening. However, it is important to note here that the eddy covariance technique provides
 805 net exchange measurements; hence, this shift does not imply the absence of deposition, but rather that emission rates surpass deposition. This bidirectional behaviour—nighttime net uptake followed by daytime net release—has been previously documented (Millet et al., 2018; Schallhart et al., 2016) and confirms the existence of two-way exchange mechanisms. Compounds



exhibiting this pattern are referred to as “bidirectional BVOCs.” Their relative contribution was particularly high in May, following the canopy leaf-out. In panel g, methanol emissions even exceeded those of isoprene and monoterpenes, which may reflect leaf expansion, as methanol is known to be produced by pectin demethylation during cell wall expansion (Kreuzwieser, 2002; Wohlfahrt et al., 2015). After the summer, these compounds generally reverted to net deposition over the full day.

Among all of the detected VOCs, formaldehyde was the only compound presenting downward fluxes at all seasons and every hour. However, one should note that the quantification of formaldehyde concentrations and fluxes comes with high challenges and uncertainties. Protonated formaldehyde can undergo back reactions with water vapour in the PTR-ToF-MS instrument (Inomata et al., 2008), making its quantification sensitive to relative humidity. In the present study, the sensitivity of our PTR-ToF-MS to formaldehyde was estimated from a comparison with HPLC analyses performed during the EMEP2022 campaign (Fagerli et al. (2023) and Sect. 2.5.1), which resulted in a very good correlation. However, this comparison was made over samples taken during 6 warm days in July. The estimated sensitivity does therefore not take different classes of relative humidities into account. This humidity-dependent sensitivity can lead to fluctuations in high-frequency data which skew fluxes computed with the EC technique. Loubet et al. (2022) estimated that the impact of water vapour variations on fluxes was negligible for most VOCs except formaldehyde, which should be considered with caution. The same advice is given in the present study.

BVOCs not individually represented in Fig. 8 still contributed substantially to the overall flux (see the “other > 0” and “other < 0” categories). In summer, between 15 and 20 species were required to capture 90% of the total upward flux expressed in carbon mass units. In comparison, Millet et al. (2018) found that only 5 ions were sufficient to represent 90% of the summed emissions in July, despite considering 377 ions in the total flux. In both cases, this indicates that strong emitters are followed by a long tail of VOCs with very low upward fluxes, although this phenomenon is more pronounced in Millet et al. (2018). In autumn, 30 compounds were needed to capture 90% of the total upward flux, as the relative importance of the top emitters (isoprene, monoterpenes) decreased. To account for 90% of the summed depositions, between 30 and 43 compounds were necessary. This confirms the observation by Millet et al. (2018) that the total deposition flux is composed of a larger number of contributing species.

Vermeuel et al. (2023a) expanded the suite of VOCs measured over a coniferous forest by combining PTR-ToF-MS with iodide chemical ionization mass spectrometry (ICIMS) flux measurements. While upward fluxes of VOC species detected with PTR-ToF-MS and included in CTMs were accurately simulated by GEOS-Chem, many additional species not represented in CTMs contributed significantly to the total downward VOC flux. This led to an underestimation of the reactive carbon sink by GEOS-Chem. Notably, downward fluxes of VOCs detected by PTR-ToF-MS alone accounted for only 63% of the total deposition, indicating that 37% would have been missed without the inclusion of ICIMS measurements. The compounds missed without ICIMS include oxidation products of isoprene and of 232-MBO, as well as >C₂ organic acids. These findings suggest that the results presented in the present study may underestimate part of the total VOC deposition.

The seasonal and diel trends described for 2023 were also observed in 2022 and 2024, with some differences that can be attributed to varying meteorological conditions and the differing contributions of tree species to the measured fluxes (discussed in Sect. 3.4.2). In early May 2022, warmer and sunnier weather favoured stronger emissions compared to the same period



in 2023 (panels a–b vs. f–g). Conversely, in June 2024, emission levels were much lower (panel l) than those in June 2022 (panel c) and 2023 (panel h)—a difference of nearly a factor of three—further highlighting the high sensitivity of BVOC
 845 fluxes to light and temperature. Notably, the relatively warm autumn conditions in 2024 sustained active emissions into early November (panel o), in contrast to 2023 where emissions did not show any clear dynamics (panel k).

The maximum monoterpene flux observed in the present study ($1.63 \mu\text{g m}^{-2} \text{s}^{-1}$) exceeded the value previously reported at the same site in 2009 using a PTR-Quad-MS ($0.69 \mu\text{g m}^{-2} \text{s}^{-1}$) (Laffineur et al., 2011). In both cases, these maxima occurred under air temperatures exceeding 30°C . From 2022 to 2024, only 0.1% of the monoterpene fluxes exceeded the 2009 value
 850 of $0.69 \mu\text{g m}^{-2} \text{s}^{-1}$. In contrast, the maximum isoprene flux in the present study was lower ($1.18 \mu\text{g m}^{-2} \text{s}^{-1}$) than in 2009 ($1.96 \mu\text{g m}^{-2} \text{s}^{-1}$). Methanol was another compound reported in 2009 and 2010 (Laffineur et al., 2012). While the 24-hour average maximum emissions were similar in both studies (approximately $0.07 \mu\text{g m}^{-2} \text{s}^{-1}$ in spring and $0.02 \mu\text{g m}^{-2} \text{s}^{-1}$ in summer), methanol deposition fluxes reported by Laffineur et al. (2012) were roughly twice those measured in the present dataset. Methanol concentrations, however, were comparable in both studies, and no unusual meteorological event was reported
 855 in 2009 or 2010 that could explain these differences. Given the many factors that may have varied between the two measurement periods—including vegetation type and cover, climatic conditions, and instrumental characteristics—it is difficult to draw firm conclusions based solely on the observed differences in monoterpene, isoprene, and methanol fluxes.

Below-canopy fluxes measured by the TRUNK system exhibited a seasonal transition similar to that observed at the TOP level (Fig. 9). However, the net balance was already dominated by deposition at the beginning of the campaign (panel a). The
 860 dominant emissions observed in late spring and early summer (panels b and c) declined earlier and more steeply, resulting in a faster shift towards net deposition in subsequent periods (panels d, e, and f).

Flux magnitudes (maximum diel values) were at least ten times lower than those observed at the top of the canopy, and across all periods. Maximum emission values rarely exceeded $0.01 \mu\text{g m}^{-2} \text{s}^{-1}$, reflecting the more limited production capacity below the canopy. Importantly, no additional significant BVOC signals appeared in the TRUNK data that were not already detected
 865 in the TOP dataset. This suggests that the same suite of compounds is involved in the exchange processes, albeit at reduced intensity.

The diurnal flux dynamics observed above the canopy were much less pronounced in the TRUNK system. A clear daytime emission peak was only visible in panels b and c, corresponding to late May and mid-June. During the remainder of the year, no systematic diurnal pattern emerged, suggesting that the below-canopy environment is governed by mechanisms other than those
 870 controlling foliar emissions. Isidorov and Zaitsev (2022) list a variety of biotic and abiotic drivers of leaf litter decomposition and soil-atmosphere VOC exchanges, many of which remain poorly characterized. Given the diversity of processes involved, a large variability in VOC exchanges measured near the ground is expected. Nonetheless, owing to its continuity and coverage of a wide range of environmental conditions, the present dataset may offer an opportunity to disentangle some of these (a)biotic controls.

875 Finally, we emphasize that TRUNK measurements do not solely represent soil and litter emissions, as some understorey vegetation (including young trees) was present below 3 m a.g.l. (Fig. 3). This vegetation may explain the daytime upward fluxes observed in panels b and c, which follow the evolution of temperature and light.




Figure 8. Seasonal evolution of VOC fluxes measured by the TOP system over the years 2022, 2023, and 2024. For each m/z value and hour of the day (in LT), mean fluxes are calculated only when at least 25% of the expected half-hourly data are available within the corresponding period. The 10 most exchanged m/z values are shown individually (see legend), while the remaining compounds are grouped into emission ("other > 0") and deposition ("other < 0") categories. Subplots aligned vertically correspond to the same seasonal period and share identical y-axis limits to allow for direct visual comparison. The same figure with variable y-axis is available in Sect. S3.7 for a distinct type of visualization.




Figure 9. Same as Fig. 8, but for TRUNK fluxes.



3.4.2 Flux dynamics

Figures 8 and 9 provide an overview of total BVOC exchange but do not highlight the specific temporal dynamics that could shed light on production and deposition mechanisms. A more detailed analysis of the flux time series allows for the identification of different categories of BVOCs based on their flux temporal behaviour. To illustrate these patterns, a selection of five m/z values is presented in Fig. 10.

Isoprene and monoterpenes (panels a and b) exhibit the clearest and least noisy flux signals. Another m/z value associated with well-defined emissions is 153.055 ($C_8H_8O_3H^+$, potentially methyl salicylate), although its flux magnitude is approximately 100 times lower (data not shown). As discussed previously, the emission of isoprene and monoterpenes peaked in summer, between June 5 and August 14. Isoprene and monoterpenes showed similar flux magnitudes, but some differences are noteworthy. For instance, from June 5 to June 19, fluxes of monoterpenes were higher in 2022 than in 2023, whereas the opposite was true for isoprene. Temperature and radiation were comparable during this period in both years (Fig. 5), however, differences in the prevailing wind direction may partly explain the observed patterns. In 2022, the winds predominantly originated from the south-west sector, where *Fagus sylvatica* is the dominant tree species (Fig. 1). Measurements conducted in 2009 at the same site by Laffineur et al. (2011) identified *Fagus sylvatica* as the main monoterpenes emitter among the species present at BE-Vie. The higher monoterpenes emissions in 2022 appear to support this finding. Conversely, in 2023, winds mainly originated from the north-east sector, dominated by *Picea abies*. The increased isoprene fluxes during that period suggest that *Picea abies* may act as an isoprene emitter at BE-Vie, in line with the findings of Filella et al. (2007), although this could not be confirmed by Laffineur et al. (2011).

Another possible explanation for the increased emissions of m/z 69.070 in 2023 compared to 2022 is provided by the signal at m/z 87.080 (panel d), to which 2-methyl-3-buten-2-ol (MBO) may contribute. Between May 22 and June 19, m/z 87.080 also showed higher upward fluxes in 2023. MBO emissions have been reported from Norway spruce forests by Juráň et al. (2017), and since this compound is known to fragment at m/z 69.070, the enhanced fluxes observed in panel b under winds from the *Picea abies* sector may reflect MBO emissions. Interestingly, emissions at m/z 87.080 developed earlier in the season than those of isoprene, and showed only limited variation throughout spring and summer. If MBO is indeed contributing to the signal at m/z 87.080, this observation contrasts with the estimates from Guenther et al. (2012), which suggest similar light and temperature dependencies for both isoprene and MBO.

The differences in emission patterns observed between the two years, and their apparent link to prevailing wind directions and stand composition, illustrate the challenges and opportunities of working in a heterogeneous forest ecosystem. While such heterogeneity complicates the attribution of observed fluxes to specific sources, it reflects the reality of many temperate forests and is likely to become even more common as forest management moves away from monocultures towards more resilient, mixed-species stands (Seliger et al., 2023, and references therein). At BE-Vie, this heterogeneity is not evenly spread but rather organized in species-dominated patches. Future efforts to refine flux interpretation could build on the existing high-resolution tree species distribution map (Fig. 1) combined with footprint-weighted flux disaggregation approaches, to better attribute observed fluxes to their biogenic sources.




Figure 10. Average diel patterns (in LT) of BVOC TOP fluxes for each two-week period. (a) Sum of monoterpenes. (b) Isoprene. (c) Methanol. (d) MBO. (e) $C_6H_{10}H^+$. The five m/z values shown exhibit distinct exchange behaviours. Solid lines represent hourly mean fluxes, and shaded areas indicate the standard deviation around the mean. For each two-week period, the diel pattern is displayed only if data were available for at least 25% of the expected hours.

The methanol flux (panel c) is a clear example of the bidirectional behaviour typical of some BVOCs, with weak nighttime deposition followed by daytime emissions. Methanol fluxes rise earlier and more rapidly than those of isoprene or monoterpenes, peaking between May 22 and June 5. This early increase and subsequent decline after June 19 likely reflect leaf development and cell expansion processes (Wohlfahrt et al., 2015).

In panel e, m/z 83.086 ($C_6H_{10}H^+$, potentially a fragment of hexenol or hexanal) exhibited an unexpected emission peak in 2023, between May 8 and May 22. This peak resulted from exceptionally high upward fluxes recorded over two days (May 9–11) and was also observed for other C6–C9 aldehydes and alcohols: m/z 101.096 ($C_6H_{12}OH^+$, likely protonated hexenol or hexanal), m/z 143.143 ($C_9H_{18}OH^+$, potentially nonanal), and two additional ions probably related to nonenal, m/z 123.117 ($C_9H_{14}H^+$) and m/z 141.127 ($C_9H_{16}OH^+$). These emission bursts were likely triggered by a hailstorm that occurred during this period in 2023, which caused significant canopy damage and left many broken branches and leaves on the forest floor. The above-mentioned ion species are commonly referred to as green leaf volatiles (GLVs) (Dudareva et al., 2013) and may have originated from wounded plant tissues—a well-documented source of GLV emissions (Scala et al., 2013; Niinemets et al., 2014). During this burst, the GLVs reached peak fluxes ranging from 18 times (for m/z 101.096) to 37 times (for m/z 123.117) their median positive fluxes, highlighting the substantial enhancement of their emissions under stress conditions.




Figure 11. Same as Fig. 10 but for O₃ fluxes. **(a)** TOP system. **(b)** TRUNK system.

3.5 Ozone fluxes

As expected, ozone fluxes were always negative, representing a net deposition into the ecosystem (Fig. 11). Unlike for VOCs, O₃ fluxes were expressed in molar units, which is a common practice, to directly reflect the number of molecules involved in atmospheric chemical reaction.

930 For the TOP system, O₃ depositions showed a clear diurnal pattern, with a maximum absolute flux found mostly around 11:00 LT and 12:00 LT. This evolution from nighttime to daytime depositions can be attributed to the stomatal aperture and subsequent ozone uptake, which is known as a major O₃ deposition pathway (Clifton et al., 2020). Average daytime ozone uptake rapidly increased from spring (8.96 nmol m⁻² s⁻¹) until reaching maximum values in summer (9.42 nmol m⁻² s⁻¹, Table 3). This reflects elevated O₃ concentrations (Fig. 7) and leaf development among deciduous species, which began in
 935 late April/early May. During summer, O₃ depositions varied to a small extent and started decreasing from October to reach 24h-averaged fluxes of -5.43 nmol m⁻² s⁻¹ in autumn. Again, this is primarily due to the onset of leaf senescence—with leaf fall typically occurring in late November—combined with lower ozone levels in ambient air.

Nighttime depositions showed less variation than daytime ones (in terms of mean values and standard deviations). The non-zero values indicate a continuous uptake, even when stomata were closed. These nighttime fluxes remained relatively stable
 940 throughout the seasons compared to daytime depositions, although the phenology of the forest did evolve. This supports the idea of non-stomatal deposition pathways, such as uptake by leaf cuticles, soil and water (Clifton et al., 2020).

Overall, the inter-annual variability of TOP fluxes was small. Daytime ozone uptake reached a maximum in 2022 in the fortnight from July 31 to August 14 with a maximum mean value of 20 nmol m⁻² s⁻¹. This can be directly attributed to the heatwave encountered at that time—as ozone reached concentrations above 70 ppbv—, despite the fact that stomatal conduc-
 945 tance must have been reduced under these conditions.

Below-canopy fluxes also exhibited a diurnal pattern, with an increase from non-zero nighttime depositions to maximum daytime depositions around 14:00 LT on average—later than observed for the TOP system. This delay may be attributed to



ozone concentrations taking longer to rise after sunrise at the TRUNK level compared to above the canopy (Fig. 7), possibly due to delayed photochemical activity near the forest floor.

950 Both in 2023 and 2024, TRUNK fluxes rapidly declined from July 3 and remained in the same order of magnitude for the rest of the campaign. This contrasts with above canopy fluxes which maintained high deposition values and seemed more sensitive to meteorological and phenological conditions. The explanation for this decrease remains unclear. As a reminder, TRUNK fluxes were measured 3 m above ground level, and therefore primarily reflect non-stomatal deposition pathways, such as uptake by soil, litter, or water surfaces (Clifton et al., 2020). These processes would be expected to persist even during
955 summer.

TRUNK fluxes represented on average approximately 10% of the above-canopy fluxes, although large variations were observed—from 0 to 50%. This is broadly consistent with the range of 5 to 39% reported by Fares et al. (2014), and references therein. The relative contribution of TRUNK fluxes was often higher at night, in agreement with observations from the Landes forest by Lamaud et al. (2002), likely due to the reduced importance of stomatal deposition in the total ozone sink
960 during the night.

In 2023, we observe a little boost of below-canopy O₃ uptake from November 6 until November 20. This period exactly coincides with the fall of dead leaves from deciduous trees. An explanation of this increased consumption might be linked to a reaction of ozone with compounds present on the surface of senescent and/or dead leaves. This phenomenon was observed both in situ and in vitro by Potier et al. (2015, 2017) and was attributed to ascorbate leaching out from senescent and dead
965 leaves.

4 Conclusions

This study presents a three-year field campaign (2022–2024) of VOC and ozone dynamics at the extensively instrumented ICOS forest station of Vielsalm (BE-Vie), combining a vertical profile with eddy covariance fluxes above and below the canopy. A new open-source processing pipeline (PAP and GEddySoft) was developed to quantify VOC mixing ratios and
970 fluxes using a PTR-ToF-MS instrument. Various critical steps of the eddy covariance methodology applied to VOC and ozone flux measurements were addressed, including data quality flagging, storage corrections, sonic anemometer rotation within the trunk space, and ozone flux calibration.

The processing of VOC data resulted in the identification of 60 significantly exchanged ions, among 76 VOC-related signals. These included both well-known biogenic VOCs and lesser-studied compounds, reflecting the chemical and functional diversity
975 of VOC exchange in forest ecosystems. The number of significantly exchanged ions is intermediate in comparison to other studies using a PTR-ToF-MS instrument—while Park et al. (2013a); Millet et al. (2018); Fischer et al. (2021); Loubet et al. (2022) identified hundreds of compounds, others found tens of VOCs exhibiting significant fluxes (Ruuskanen et al., 2011; Jensen et al., 2018; Schallhart et al., 2018). Despite potential differences in ecosystem signatures and instrumental capabilities, this raises questions about the criteria used to determine whether a flux is considered significant or not.



980 Ambient mixing ratios revealed clear vertical and diurnal gradients shaped by the interplay of emission, deposition, photochemistry, and atmospheric transport. Ozone and NO_x exhibited tightly coupled patterns, with daytime O_3 formation and nighttime NO_x accumulation. VOCs showed compound-specific behaviour: ethanol remained relatively constant due to high background levels, methanol followed a daytime emission pattern, and isoprene peaked near the canopy with oxidation products (MVK + MACR) accumulating at higher altitudes. At night, monoterpenes accumulated near the forest floor, indicating
 985 sustained temperature-driven emissions and limited mixing. These observations highlight the value of high-resolution datasets for exploring the spatial and temporal dynamics of reactive trace gases within a forest ecosystem.

The forest acted as a net source of VOCs for most of the measurement period, with peak emissions in summer exceeding $1.25 \mu\text{g m}^{-2} \text{s}^{-1}$. Monoterpenes, isoprene, and methanol dominated the fluxes. However, between 15 and 30 compounds were needed to capture 90% of the total emissions (depending on the season), and between 30 and 43 for deposition, confirming the
 990 importance of considering a broad range of compounds beyond a few dominant species. According to Vermeuel et al. (2023a), up to 37% of the total downward flux of VOCs would be missed when relying on PTR-ToF-MS measurements alone. Therefore, while the upward fluxes reported in this study are likely representative of total ecosystem emissions, the total deposition should be regarded as a lower limit of the actual ecosystem sink.

The total VOC deposition remained relatively stable across seasons ($\sim -0.1 \mu\text{g m}^{-2} \text{s}^{-1}$), while emissions showed stronger
 995 seasonal variability. Many oxygenated VOCs exhibited bidirectional behaviour, switching from nighttime deposition to daytime emission—especially in spring after budburst—underlining the importance of biological processes and environmental conditions.

The interannual variability in flux magnitudes—such as the markedly lower emissions observed in June 2024 compared to previous years—further emphasizes the sensitivity of VOC exchange to meteorological and phenological drivers. Fluxes
 1000 measured near the forest floor were generally close to equilibrium between emissions and depositions, except in late spring and early summer, when upward fluxes increased.

Additional insights were gained into species-specific emission patterns, with wind direction modulating isoprene and monoterpene fluxes. For instance, *Picea abies* was identified as a likely isoprene and MBO emitter at this site. A short-lived burst of green leaf volatiles following a hailstorm also illustrated the ecosystem's capacity for rapid VOC responses to disturbance. The
 1005 spatial heterogeneity of the forest, while challenging for flux attribution, also offers an opportunity to improve our understanding of species-specific emissions in realistic, mixed-species stands.

Ozone fluxes remained consistently downward, with the strongest uptake in spring and early summer. Below-canopy deposition accounted for $\sim 10\%$ of the above-canopy flux on average, although this ratio was variable. These patterns point to persistent non-stomatal uptake processes, both above and below the canopy, and suggest opportunities for further partitioning
 1010 of ozone sinks.

Overall, this study provides one of the most detailed long-term characterizations of trace gas exchange in a temperate mixed forest. The diversity of behaviours observed across compounds, seasons, and canopy levels offers a unique dataset for the development and calibration of process-based models, including bidirectional biogenic, physical and chemical exchange



schemes. Such models are essential for improving predictions of VOC–ozone–aerosol interactions and their feedbacks on air
1015 quality and climate.

5 Code and data availability

Four distinct groups of datasets are available for the BE-Vie site during the 2022-2023-2024 period:

- (i) Main measurements (present study). 1-min concentrations of VOCs and O₃ are available at <https://doi.org/10.18758/NVFBA74V> (Verreyken et al., 2025); 30-min concentrations and fluxes (TOP & TRUNK) of VOCs and O₃ are available
1020 at <https://doi.org/10.18758/KHV8ZXU2> (Dumont et al., 2025a); vertical profiles of VOC and O₃ concentrations and turbulence are available at <https://doi.org/10.18758/BED4Q2VY> (Dumont et al., 2025b).
- (ii) ICOS measurements. Flux and meteorological data were downloaded from the ETC L2 ARCHIVE on the ICOS Carbon portal <https://doi.org/10.18160/S6HM-CP8Q> (Vincke et al., 2025).
- (iii) ISSeP measurements. Data from the ISSeP air quality station were downloaded from the wallonair platform <https://data.wallonair.be/>. More information available upon request from b.bergmans@issep.be.
1025
- (iv) Remote-sensing measurements. Data available upon request from alexis.merlaud@aeronomie.be.

Results from the EMEP campaigns (VOC analysis by GC-MS and HPLC) will be made available through the EBAS data repository (<http://ebas.nilu.no/>).

The Peak Area Processing software (PAP, used for non-targeted peak selection, instrument characterization, and quantifica-
1030 tion of mixing ratio's with related uncertainties) is available here: <https://github.com/bverreyk/PeakAreaProcessing>. GEddySoft (software used to process VOC fluxes with the eddy covariance method) is available here: <https://github.com/BernardHeinesch/GEddySoft>.

Author contributions. This study was conceptualized by CA, NS, BH, CD and BV. VOC measurements were supervised by CA, NS, and BV, and ozone measurements by CD. PTR-ToF-MS data were curated by BV, CA, and NS, and mixing ratios were computed by BV and
1035 CA (formal analysis). VOC fluxes were calculated and post-processed by BH and CD (formal analysis). Data curation and formal analysis of ozone concentrations and fluxes, as well as of the turbulence profile, were carried out by CD. CD and BV performed the formal analysis and visualization of the final datasets, and wrote the first draft of the manuscript. All authors participated in the review and editing process of the final paper version. BH, CA, and NS contributed to funding acquisition. BH and CA supervised the research activities.

Competing interests. The contact author has declared that none of the authors has any competing interests.



1040 *Acknowledgements.* The authors gratefully acknowledge the Belgian Federal Science Policy Office (BELSPO) for supporting the ACTRIS-
BE project (FSIRI/00/AC1) and the BIRA/ULiège FED-tWIN project BERTRAC (Prf-2021-034_BERTRAC#2), as well as the Public Ser-
vice of Wallonia for their support to the ICOS-Wallonia project (1217769). The authors thank the Carbon Portal (CP) for providing access
to the ETC L2 ARCHIVE, and Caroline Vincke (PI of the BE-Vie station) for facilitating access to the Vielsalm site. We also thank ISSeP
for providing access to their NO_x measurements. We thank the technical team at the Royal Belgian Institute for Space Aeronomy (Sam De
1045 Pauw, Tom Egerickx, Gerry Pieck and Octavian Miron) for designing and installing the concentration profile setup. Finally, we thank Henri
Chopin for his support with the technical aspects of ozone measurements, and Ariane Faurès for her advice on spectral corrections and for
sharing the corresponding data management code. The authors also acknowledge the use of the AI language model ChatGPT-4 for assistance
with Python coding during data analysis and for language refinement of the manuscript. All scientific interpretations and conclusions remain
those of the authors.



1050 References

- Andersen, S. T., McGillen, M. R., Xue, C., Seubert, T., Dewald, P., Türk, G. N., Schuladen, J., Denjean, C., Etienne, J. C., Garrouste, O., Jamar, M., Harb, S., Cirtog, M., Michoud, V., Cazaunau, M., Bergé, A., Cantrell, C., Dusanter, S., Picquet-Varrault, B., Kukui, A., Mellouki, A., Carpenter, L. J., Lelieveld, J., and Crowley, J. N.: Measurement report: Sources, sinks, and lifetime of NO_x in a suburban temperate forest at night, *Atmospheric Chemistry and Physics*, 24, 11 603–11 618, <https://doi.org/10.5194/acp-24-11603-2024>, 2024.
- 1055 Ashworth, K., Chung, S. H., McKinney, K. A., Liu, Y., Munger, J. W., Martin, S. T., and Steiner, A. L.: Modelling bidirectional fluxes of methanol and acetaldehyde with the FORCAsT canopy exchange model, *Atmospheric Chemistry and Physics*, 16, 15 461–15 484, <https://doi.org/10.5194/acp-16-15461-2016>, 2016.
- Aubinet, M., Chermanne, B., Vandenhaute, M., Longdoz, B., Yernaux, M., and Laitat, E.: Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes, *Agricultural and Forest Meteorology*, 108, 293–315, [https://doi.org/10.1016/S0168-1923\(01\)00244-1](https://doi.org/10.1016/S0168-1923(01)00244-1),
1060 2001.
- Aubinet, M., Hurdebise, Q., Chopin, H., Debacq, A., De Ligne, A., Heinesch, B., Manise, T., and Vincke, C.: Inter-annual variability of Net Ecosystem Productivity for a temperate mixed forest: A predominance of carry-over effects?, *Agricultural and Forest Meteorology*, 262, 340–353, <https://doi.org/10.1016/j.agrformet.2018.07.024>, 2018.
- Bachy, A., Aubinet, M., Schoon, N., Amelynck, C., Bodson, B., Moureaux, C., and Heinesch, B.: Are BVOC exchanges in agricultural
1065 ecosystems overestimated? Insights from fluxes measured in a maize field over a whole growing season, *Atmospheric Chemistry and Physics*, 16, 5343–5356, <https://doi.org/10.5194/acp-16-5343-2016>, 2016.
- Bachy, A., Aubinet, M., Amelynck, C., Schoon, N., Bodson, B., Moureaux, C., Delaplace, P., De Ligne, A., and Heinesch, B.: Methanol exchange dynamics between a temperate cropland soil and the atmosphere, *Atmospheric Environment*, 176, 229–239, <https://doi.org/10.1016/j.atmosenv.2017.12.016>, 2018.
- 1070 Bamberger, I., Hörtnagl, L., Ruuskanen, T. M., Schnitzhofer, R., Müller, M., Graus, M., Karl, T., Wohlfahrt, G., and Hansel, A.: Deposition fluxes of terpenes over grassland, *Journal of Geophysical Research Atmospheres*, 116, 1–13, <https://doi.org/10.1029/2010JD015457>, 2011.
- Brilli, F., Gioli, B., Zona, D., Pallozzi, E., Zenone, T., Fratini, G., Calfapietra, C., Loreto, F., Janssens, I. A., and Ceulemans, R.: Simultaneous leaf- and ecosystem-level fluxes of volatile organic compounds from a poplar-based SRC plantation, *Agricultural and Forest Meteorology*, 187, 22–35, <https://doi.org/10.1016/j.agrformet.2013.11.006>, 2014.
- 1075 Brilli, F., Gioli, B., Fares, S., Terenzio, Z., Zona, D., Gielen, B., Loreto, F., Janssens, I. A., and Ceulemans, R.: Rapid leaf development drives the seasonal pattern of volatile organic compound (VOC) fluxes in a 'coppiced' bioenergy poplar plantation, *Plant Cell and Environment*, 39, 539–555, <https://doi.org/10.1111/pce.12638>, 2016.
- Canadell, J., Monteiro, P., Costa, M., Cotrim da Cunha, L., Cox, P., Eliseev, A., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lo-hila, A., Patra, P., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: *Global Carbon and other Biogeochemical Cycles and Feedbacks*, p. 673–816, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.007>, 2021.
- 1080 Cieslik, S.: Energy and ozone fluxes in the atmospheric surface layer observed in Southern Germany highlands, *Atmospheric Environment*, 32, 1273–1281, [https://doi.org/https://doi.org/10.1016/S1352-2310\(97\)00190-8](https://doi.org/https://doi.org/10.1016/S1352-2310(97)00190-8), 1998.
- Clifton, O. E., Fiore, A. M., Massman, W. J., Baublitz, C. B., Coyle, M., Emberson, L., Fares, S., Farmer, D. K., Gentine, P., Gerosa, G.,
1085 Guenther, A. B., Helmig, D., Lombardozzi, D. L., Munger, J. W., Patton, E. G., Pusede, S. E., Schwede, D. B., Silva, S. J., Sörgel, M.,



- Steiner, A. L., and Tai, A. P. K.: Dry Deposition of Ozone Over Land: Processes, Measurement, and Modeling, *Reviews of Geophysics*, 58, <https://doi.org/10.1029/2019RG000670>, 2020.
- Dudareva, N., Klempien, A., Muhlemann, J. K., and Kaplan, I.: Biosynthesis, function and metabolic engineering of plant volatile organic compounds, *New Phytologist*, 198, 16–32, <https://doi.org/10.1111/nph.12145>, 2013.
- 1090 Dumont, C., Verreyken, B. W. D., Schoon, N., Amelynck, C., and Heinesch, B.: 3-years of (O)VOC and ozone flux measurements at a mixed temperate forest at the Vielsalm ICOS ecosystem station (Belgium), <https://doi.org/10.18758/KHV8ZXU2>, 2025a.
- Dumont, C., Verreyken, B. W. D., Schoon, N., Heinesch, B., and Amelynck, C.: 3-years of (O)VOC, ozone, and turbulence profile measurements at a mixed temperate forest at the Vielsalm ICOS ecosystem station (Belgium), <https://doi.org/10.18758/BED4Q2VY>, 2025b.
- Dusanter S., Holzinger R., Klein F., Salameh T., and Jamar M.: Measurement Guidelines for VOC Analysis by PTR-MS, <https://www.actris.eu/sites/default/files/inline-files/PTRMS%20SOP%20%28April2025%29.pdf>, 2025.
- 1095 Ester, M., Kriegel, H.-P., Sander, J., and Xu, X.: A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise, www.aaai.org, 1996.
- Fagerli, H., Benedictow, A., Van Caspel, W., Gauss, M., Ge, Y., Jonson, J. E., Klein, H., Nyíri, Á., Simpson, D., Tsyro, S., Valdebenito, Á., Wind, P., Aas, W., Hjellbrekke, A., Solberg, S., Tørseth, K., Espen Yttri, K., Matthews, B., Schindlbacher, S., Ullrich, B., Wankmüller, R., Klimont, Z., Scheuschner, T., Kuenen, J. J. P., Hellén, H., Jaffrezo, J.-L., Tusha, D., Mothes, F., Salameh, T., van Drooge, B. L., and Wegener, R.: EMEP Status report: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Tech. rep., Norwegian Meteorological Institute, ISSN 1504-6192, https://emep.int/publ/reports/2023/EMEP_Status_Report_1_2023.pdf, 2023.
- 1100 Faiola, C., Ossola, R., and McGlynn, D.: Chapter 5 - The role of biogenic volatile organic compounds and plant surfaces in the formation and scavenging of ozone and particulate matter, including secondary organic aerosol, in: *Biogenic Volatile Organic Compounds and Climate Change*, edited by Brilli, F. and Decesari, S., pp. 165–195, Elsevier, ISBN 978-0-12-821076-5, <https://doi.org/10.1016/B978-0-12-821076-5.00004-5>, 2024.
- 1105 Fares, S., Savi, F., Muller, J., Matteucci, G., and Paoletti, E.: Simultaneous measurements of above and below canopy ozone fluxes help partitioning ozone deposition between its various sinks in a Mediterranean Oak Forest, *Agricultural and Forest Meteorology*, 198–199, 181–191, <https://doi.org/10.1016/j.agrformet.2014.08.014>, 2014.
- 1110 Filella, I., Wilkinson, M. J., Llusà, J., Hewitt, C. N., and Peñuelas, J.: Volatile organic compounds emissions in Norway spruce (*Picea abies*) in response to temperature changes, *Physiologia Plantarum*, 130, 58–66, <https://doi.org/10.1111/j.1399-3054.2007.00881.x>, 2007.
- Finco, A., Coyle, M., Nemitz, E., Marzuoli, R., Chiesa, M., Loubet, B., Fares, S., Diaz-Pines, E., Gasche, R., and Gerosa, G.: Characterization of ozone deposition to a mixed oak–hornbeam forest – flux measurements at five levels above and inside the canopy and their interactions with nitric oxide, *Atmospheric Chemistry and Physics*, 18, 17 945–17 961, <https://doi.org/10.5194/acp-18-17945-2018>, 2018.
- 1115 Finkelstein, P. L. and Sims, P. F.: Sampling error in eddy correlation flux measurements, *Journal of Geophysical Research: Atmospheres*, 106, 3503–3509, <https://doi.org/10.1029/2000JD900731>, 2001.
- Fischer, L., Breitenlechner, M., Canaval, E., Scholz, W., Striednig, M., Graus, M., Karl, T. G., Petäjä, T., Kulmala, M., and Hansel, A.: First eddy covariance flux measurements of semi-volatile organic compounds with the PTR3-TOF-MS, *Atmospheric Measurement Techniques*, 14, 8019–8039, <https://doi.org/10.5194/amt-14-8019-2021>, 2021.
- 1120 Fiscus, E. L., Booker, F. L., and Burkey, K. O.: Crop responses to ozone: uptake, modes of action, carbon assimilation and partitioning, *Plant, Cell Environment*, 28, 997–1011, <https://doi.org/10.1111/j.1365-3040.2005.01349.x>, 2005.
- Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agricultural and Forest Meteorology*, 78, 83–105, [https://doi.org/10.1016/0168-1923\(95\)02248-1](https://doi.org/10.1016/0168-1923(95)02248-1), 1996.



- Forkel, R., Guenther, A., Ashworth, K., Bedos, C., Delon, C., Lathiere, J., Noe, S., Potier, E., Rinne, J., Tchepel, O., and Zhang, L.:
 1125 Bi-directional Exchange of Volatile Organic Compounds, pp. 169–179, Springer Netherlands, Dordrecht, ISBN 978-94-017-7285-3,
https://doi.org/10.1007/978-94-017-7285-3_8, 2015.
- Gerosa, G. A., Bignotti, L., and Marzuoli, R.: Effect of corrections for water vapor sensitivity of coumarin targets and for density fluctuations (WPL) on O₃ fluxes measured with the eddy covariance technique, *Bulletin of Atmospheric Science and Technology*, 3,
<https://doi.org/10.1007/s42865-022-00053-0>, 2022a.
- 1130 Gerosa, G. A., Marzuoli, R., and Finco, A.: Interannual variability of ozone fluxes in a broadleaf deciduous forest in Italy, *Elementa: Science of the Anthropocene*, 10, 1–22, <https://doi.org/10.1525/elementa.2021.00105>, 2022b.
- Goldstein, A. H. and Galbally, I. E.: Known and Unexplored Organic Constituents in the Earth's Atmosphere, *Environmental Science Technology*, 41, 1514–1521, <https://doi.org/10.1021/es072476p>, 2007.
- Goldstein, A. H., McKay, M., Kurpius, M. R., Schade, G. W., Lee, A., Holzinger, R., and Rasmussen, R. A.: Forest thinning experiment confirms ozone deposition to forest canopy is dominated by reaction with biogenic VOCs, *Geophysical Research Letters*, 31, 1–4,
 1135 <https://doi.org/10.1029/2004GL021259>, 2004.
- Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research: Atmospheres*, 100, 8873–8892, <https://doi.org/10.1029/94JD02950>, 1995.
- 1140 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Güsten, H. and Heinrich, G.: On-line measurements of ozone surface fluxes: Part I. Methodology and instrumentation, *Atmospheric Environment*, 30, 897–909, [https://doi.org/10.1016/1352-2310\(95\)00269-3](https://doi.org/10.1016/1352-2310(95)00269-3), 1996.
- 1145 Hellén, H., Praplan, A. P., Tykkä, T., Ylivinkka, I., Vakkari, V., Bäck, J., Petäjä, T., Kulmala, M., and Hakola, H.: Long-term measurements of volatile organic compounds highlight the importance of sesquiterpenes for the atmospheric chemistry of a boreal forest, *Atmospheric Chemistry and Physics*, 18, 13 839–13 863, <https://doi.org/10.5194/acp-18-13839-2018>, 2018.
- Hodzic, A., Aumont, B., Knote, C., Lee-Taylor, J., Madronich, S., and Tyndall, G.: Volatility dependence of Henry's law constants of condensable organics: Application to estimate depositional loss of secondary organic aerosols, *Geophysical Research Letters*, 41, 4795–
 1150 4804, <https://doi.org/10.1002/2014GL060649>, 2014.
- Horváth, L., Koncz, P., Móring, A., Nagy, Z., Pintér, K., and Weidinger, T.: An Attempt to Partition Stomatal and Non-stomatal Ozone Deposition Parts on a Short Grassland, *Boundary-Layer Meteorology*, 167, 303–326, <https://doi.org/10.1007/s10546-017-0310-x>, 2017.
- Huang, G., Brook, R., Crippa, M., Janssens-Maenhout, G., Schieberle, C., Dore, C., Guizzardi, D., Muntean, M., Schaaf, E., and Friedrich, R.: Speciation of anthropogenic emissions of non-methane volatile organic compounds: A global gridded data set for 1970–2012, *Atmospheric Chemistry and Physics*, 17, 7683–7701, <https://doi.org/10.5194/acp-17-7683-2017>, 2017.
- 1155 Inomata, S., Tanimoto, H., Kameyama, S., Tsunogai, U., Irie, H., Kanaya, Y., and Wang, Z.: Technical Note: Determination of formaldehyde mixing ratios in air with PTR-MS: laboratory experiments and field measurements, *Atmospheric Chemistry and Physics*, 8, 273–284, <https://doi.org/10.5194/acp-8-273-2008>, 2008.
- Isidorov, V. A. and Zaitsev, A. A.: Reviews and syntheses: VOC emissions from soil cover in boreal and temperate natural ecosystems of the
 1160 Northern Hemisphere, *Biogeosciences*, 19, 4715–4746, <https://doi.org/10.5194/bg-19-4715-2022>, 2022.



- Isidorov, V. A., Pirožnikow, E., Spirina, V. L., Vasyanin, A. N., Kulakova, S. A., Abdulmanova, I. F., and Zaitsev, A. A.: Emission of volatile organic compounds by plants on the floor of boreal and mid-latitude forests, *Journal of Atmospheric Chemistry*, 79, 153–166, <https://doi.org/10.1007/s10874-022-09434-3>, 2022.
- Jardine, K., Yañez Serrano, A., Arneth, A., Abrell, L., Jardine, A., Artaxo, P., Alves, E., Kesselmeier, J., Taylor, T., Saleska, S., and Huxman, T.: Ecosystem-scale compensation points of formic and acetic acid in the central Amazon, *Biogeosciences*, 8, 3709–3720, <https://doi.org/10.5194/bg-8-3709-2011>, 2011.
- Jensen, N. R., Gruening, C., Goded, I., Müller, M., Hjorth, J., and Wisthaler, A.: Eddy-covariance flux measurements in an Italian deciduous forest using PTR-ToF-MS, PTR-QMS and FIS, *International Journal of Environmental Analytical Chemistry*, 98, 758–788, <https://doi.org/10.1080/03067319.2018.1502758>, 2018.
- Juráň, S., Pallozzi, E., Guidolotti, G., Fares, S., Šigut, L., Calfapietra, C., Alivernini, A., Savi, F., Večeřová, K., Křůmal, K., Večeřa, Z., and Urban, O.: Fluxes of biogenic volatile organic compounds above temperate Norway spruce forest of the Czech Republic, *Agricultural and Forest Meteorology*, 232, 500–513, <https://doi.org/10.1016/j.agrformet.2016.10.005>, 2017.
- Kammer, J., Perraudin, E., Flaud, P.-M., Lamaud, E., Bonnefond, J., and Villenave, E.: Observation of nighttime new particle formation over the French Landes forest, *Science of The Total Environment*, 621, 1084–1092, <https://doi.org/10.1016/j.scitotenv.2017.10.118>, 2018.
- Karl, T., Potosnak, M., Guenther, A., Clark, D., Walker, J., Herrick, J. D., and Geron, C.: Exchange processes of volatile organic compounds above a tropical rain forest: Implications for modeling tropospheric chemistry above dense vegetation, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004738>, 2004a.
- Karl, T., Potosnak, M., Guenther, A., Clark, D., Walker, J., Herrick, J. D., and Geron, C.: Exchange processes of volatile organic compounds above a tropical rain forest: Implications for modeling tropospheric chemistry above dense vegetation, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004738>, 2004b.
- Karl, T., Harley, P., Emmons, L., Thornton, B., Guenther, A., Basu, C., Turnipseed, A., and Jardine, K.: Efficient atmospheric cleansing of oxidized organic trace gases by vegetation, *Science*, 330, 816–819, <https://doi.org/10.1126/science.1192534>, 2010.
- Kaser, L., Karl, T., Guenther, A., Graus, M., Schnitzhofer, R., Turnipseed, A., Fischer, L., Harley, P., Madronich, M., Gochis, D., Keutsch, F. N., and Hansel, A.: Undisturbed and disturbed above canopy ponderosa pine emissions: PTR-TOF-MS measurements and MEGAN 2.1 model results, *Atmospheric Chemistry and Physics*, 13, 11 935–11 947, <https://doi.org/10.5194/acp-13-11935-2013>, 2013.
- Kim, S., Karl, T., Guenther, A., Tyndall, G., Orlando, J., Harley, P., Rasmussen, R., and Apel, E.: Emissions and ambient distributions of Biogenic Volatile Organic Compounds (BVOC) in a ponderosa pine ecosystem: interpretation of PTR-MS mass spectra, *Atmospheric Chemistry and Physics*, 10, 1759–1771, <https://doi.org/10.5194/acp-10-1759-2010>, 2010.
- Kljun, N., Calanca, P., Rotach, M. W., and Schmid, H. P.: A simple two-dimensional parameterisation for Flux Footprint Prediction (FFP), *Geoscientific Model Development*, 8, 3695–3713, <https://doi.org/10.5194/gmd-8-3695-2015>, 2015.
- Kreuzwieser, J.: Biosynthesis of aldehydes and organic acids, pp. 101–114, Springer Netherlands, Dordrecht, ISBN 978-94-015-9856-9, https://doi.org/10.1007/978-94-015-9856-9_5, 2002.
- Kurpius, M. R. and Goldstein, A. H.: Gas-phase chemistry dominates O₃ loss to a forest, implying a source of aerosols and hydroxyl radicals to the atmosphere, *Geophysical Research Letters*, 30, 2–5, <https://doi.org/10.1029/2002GL016785>, 2003.
- Laffineur, Q., Aubinet, M., Schoon, N., Amelynck, C., Müller, J. F., Dewulf, J., Van Langenhove, H., Steppe, K., Šimpraga, M., and Heinesch, B.: Isoprene and monoterpene emissions from a mixed temperate forest, *Atmospheric Environment*, 45, 3157–3168, <https://doi.org/10.1016/j.atmosenv.2011.02.054>, 2011.



- Laffineur, Q., Aubinet, M., Schoon, N., Amelynck, C., Müller, J. F., Dewulf, J., Van Langenhove, H., Steppe, K., and Heinesch, B.: Abiotic and biotic control of methanol exchanges in a temperate mixed forest, *Atmospheric Chemistry and Physics*, 12, 577–590, <https://doi.org/10.5194/acp-12-577-2012>, 2012.
- Laffineur, Q., Aubinet, M., Schoon, N., Amelynck, C., Müller, J. F., Dewulf, J., Steppe, K., and Heinesch, B.: Impact of diffuse light on isoprene and monoterpene emissions from a mixed temperate forest, *Atmospheric Environment*, 74, 385–392, <https://doi.org/10.1016/j.atmosenv.2013.04.025>, 2013.
- Lamaud, E., Carrara, A., Brunet, Y., Lopez, A., and Druilhet, A.: Ozone fluxes above and within a pine forest canopy in dry and wet conditions, *Atmospheric Environment*, 36, 77–88, [https://doi.org/10.1016/S1352-2310\(01\)00468-X](https://doi.org/10.1016/S1352-2310(01)00468-X), 2002.
- Langford, B., Acton, W., Ammann, C., Valach, A., and Nemitz, E.: Eddy-covariance data with low signal-to-noise ratio: Time-lag determination, uncertainties and limit of detection, *Atmospheric Measurement Techniques*, <https://doi.org/10.5194/amt-8-4197-2015>, 2015.
- Lanssens, B.: Estimation de la biomasse par espèce forestière par télédétection : application à la validation d’un modèle dynamique de végétation, Master’s thesis, University of Liège, <http://hdl.handle.net/2268.2/8298>, 2019.
- Leuning, R.: Estimation of Scalar Source/Sink Distributions in Plant Canopies Using Lagrangian Dispersion Analysis: Corrections for Atmospheric Stability and Comparison with a Multilayer Canopy Model, *Boundary-Layer Meteorology*, 96, 293–314, <https://doi.org/10.1023/A:1002449700617>, 2000.
- Link, M. F., Pothier, M. A., Vermeuel, M. P., Riches, M., Millet, D. B., and Farmer, D. K.: In-Canopy Chemistry, Emissions, Deposition, and Surface Reactivity Compete to Drive Bidirectional Forest-Atmosphere Exchange of VOC Oxidation Products, *ACS EST Air*, 1, 305–315, <https://doi.org/10.1021/acsestair.3c00074>, 2024.
- Loubet, B., Buysse, P., Gonzaga-Gomez, L., Lafouge, F., Ciuraru, R., Decuq, C., Kammer, J., Bsaibes, S., Boissard, C., Durand, B., Gueudet, J.-C., Fanucci, O., Zurfluh, O., Abis, L., Zannoni, N., Truong, F., Baisnée, D., Sarda-Estève, R., Staudt, M., and Gros, V.: Volatile organic compound fluxes over a winter wheat field by PTR-Qi-TOF-MS and eddy covariance, *Atmospheric Chemistry and Physics*, 22, 2817–2842, <https://doi.org/10.5194/acp-22-2817-2022>, 2022.
- Mahilang, M., Deb, M. K., and Pervez, S.: Biogenic secondary organic aerosols: A review on formation mechanism, analytical challenges and environmental impacts, *Chemosphere*, 262, 127 771, <https://doi.org/10.1016/j.chemosphere.2020.127771>, 2021.
- Mahrt, L.: Flux Sampling Errors for Aircraft and Towers, *Journal of Atmospheric and Oceanic Technology*, 15, 416–429, [https://doi.org/10.1175/1520-0426\(1998\)015<0416:FSEFAA>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<0416:FSEFAA>2.0.CO;2), 1998.
- Manco, A., Brilli, F., Famulari, D., Gasbarra, D., Gioli, B., Vitale, L., di Tommasi, P., Loubet, B., Arena, C., and Magliulo, V.: Cross-correlations of Biogenic Volatile Organic Compounds (BVOC) emissions typify different phenological stages and stressful events in a Mediterranean Sorghum plantation, *Agricultural and Forest Meteorology*, 303, <https://doi.org/10.1016/j.agrformet.2021.108380>, 2021.
- Mauder, M. and Foken, T.: Documentation and Instruction Manual of the Eddy-Covariance Software Package TK2, Abt. Mikrometeorologie, pp. 60, ISSN 1614–8916, <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:bvb:703-opus-8665%5Cnhttp://opus4.kobv.de/opus4-ubbayreuth/frontdoor/index/index/docId/681>, 2004.
- Mikkelsen, T., Ro-Poulsen, H., Hovmand, M., Jensen, N., Pilegaard, K., and Egeløv, A.: Five-year measurements of ozone fluxes to a Danish Norway spruce canopy, *Atmospheric Environment*, 38, 2361–2371, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2003.12.036>, new methods of risk assessment for ozone impacts on vegetation, 2004.
- Millet, D. B., Jacob, D. J., Custer, T. G., de Gouw, J. A., Goldstein, A. H., Karl, T., Singh, H. B., Sive, B. C., Talbot, R. W., Warneke, C., and Williams, J.: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmospheric Chemistry and Physics*, 8, 6887–6905, <https://doi.org/10.5194/acp-8-6887-2008>, 2008.



- Millet, D. B., Alwe, H. D., Chen, X., Deventer, M. J., Griffis, T. J., Holzinger, R., Bertman, S. B., Rickly, P. S., Stevens, P. S., Léonardis, T., Locoge, N., Dusanter, S., Tyndall, G. S., Alvarez, S. L., Erickson, M. H., and Flynn, J. H.: Bidirectional Ecosystem–Atmosphere Fluxes of Volatile Organic Compounds Across the Mass Spectrum: How Many Matter?, *ACS Earth and Space Chemistry*, 2, 764–777, <https://doi.org/10.1021/acsearthspacechem.8b00061>, 2018.
- 1240 Min, K.-E., Pusede, S. E., Browne, E. C., LaFranchi, B. W., and Cohen, R. C.: Eddy covariance fluxes and vertical concentration gradient measurements of NO and NO₂ over a ponderosa pine ecosystem: observational evidence for within-canopy chemical removal of NO_x, *Atmospheric Chemistry and Physics*, 14, 5495–5512, <https://doi.org/10.5194/acp-14-5495-2014>, 2014.
- Montagnani, L., Grünwald, T., Kowalski, A., Mammarella, I., Merbold, L., Metzger, S., Sedláč, P., and Siebicke, L.: Estimating the storage term in eddy covariance measurements: the ICOS methodology, *International Agrophysics*, 32, 551–567, [https://doi.org/10.1515/intag-](https://doi.org/10.1515/intag-2017-0037)
1245 2017-0037, 2018.
- Muller, J. B., Percival, C. J., Gallagher, M. W., Fowler, D., Coyle, M., and Nemitz, E.: Sources of uncertainty in eddy covariance ozone flux measurements made by dry chemiluminescence fast response analysers, *Atmospheric Measurement Techniques*, 3, 163–176, <https://doi.org/10.5194/amt-3-163-2010>, 2010.
- Müller, M., Graus, M., Ruuskanen, T. M., Schnitzhofer, R., Bamberger, I., Kaser, L., Titzmann, T., Hörtnagl, L., Wohlfahrt, G., Karl,
1250 T., and Hansel, A.: First eddy covariance flux measurements by PTR-TOF, *Atmospheric Measurement Techniques*, 3, 387–395, <https://doi.org/10.5194/amt-3-387-2010>, 2010.
- Naik, V., Fiore, A. M., Horowitz, L. W., Singh, H. B., Wiedinmyer, C., Guenther, A., de Gouw, J. A., Millet, D. B., Goldan, P. D., Kuster, W. C., and Goldstein, A.: Observational constraints on the global atmospheric budget of ethanol, *Atmospheric Chemistry and Physics*, 10, 5361–5370, <https://doi.org/10.5194/acp-10-5361-2010>, 2010.
- 1255 Nemitz, E., Sutton, M. A., Gut, A., San José, R., Husted, S., and Schjoerring, J. K.: Sources and sinks of ammonia within an oilseed rape canopy, *Agricultural and Forest Meteorology*, 105, 385–404, [https://doi.org/10.1016/S0168-1923\(00\)00205-7](https://doi.org/10.1016/S0168-1923(00)00205-7), 2000.
- Nguyen, T. B., Crounse, J. D., Teng, A. P., Clair, J. M., Paulot, F., Wolfe, G. M., and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proceedings of the National Academy of Sciences of the United States of America*, 112, E392–E401, <https://doi.org/10.1073/pnas.1418702112>, 2015.
- 1260 Niinemets, Ü., Fares, S., Harley, P., and Jardine, K. J.: Bidirectional exchange of biogenic volatiles with vegetation: Emission sources, reactions, breakdown and deposition, *Plant, Cell and Environment*, 37, 1790–1809, <https://doi.org/10.1111/pce.12322>, 2014.
- Padro, J.: Summary of ozone dry deposition velocity measurements and model estimates over vineyard, cotton, grass and deciduous forest in summer, *Atmospheric Environment*, 30, 2363–2369, [https://doi.org/https://doi.org/10.1016/1352-2310\(95\)00352-5](https://doi.org/10.1016/1352-2310(95)00352-5), fourth International Conference on Atmospheric Science and Applications to Air Quality, 1996.
- 1265 Pagonis, D., Sekimoto, K., and de Gouw, J.: A Library of Proton-Transfer Reactions of H₃O⁺ Ions Used for Trace Gas Detection, *Journal of the American Society for Mass Spectrometry*, 30, 1330–1335, <https://doi.org/10.1007/s13361-019-02209-3>, 2019.
- Park, J.-H., Goldstein, A. H., Timkovsky, J., Fares, S., Weber, R., Karlik, J., and Holzinger, R.: Active Atmosphere-Ecosystem Exchange of the Vast Majority of Detected Volatile Organic Compounds, *Science*, 341, 643–647, <https://doi.org/10.1126/science.1235053>, 2013a.
- Park, J. H., Goldstein, A. H., Timkovsky, J., Fares, S., Weber, R., Karlik, J., and Holzinger, R.: Eddy covariance emission and deposition flux
1270 measurements using proton transfer reaction ndash; Time of flight ndash; Mass spectrometry (PTR-TOF-MS): Comparison with PTR-MS measured vertical gradients and fluxes, *Atmospheric Chemistry and Physics*, 13, 1439–1456, <https://doi.org/10.5194/acp-13-1439-2013>, 2013b.



- Pellegrini, E., Lorenzini, G., and Nali, C.: The 2003 European Heat Wave: Which Role for Ozone? Some Data from Tuscany, Central Italy, Water, Air, and Soil Pollution, 181, 401–408, <https://doi.org/10.1007/s11270-006-9310-z>, 2007.
- 1275 Peltola, O., Aslan, T., Ibrom, A., Nemitz, E., Rannik, U., and Mammarella, I.: The high-frequency response correction of eddy covariance fluxes - Part 1: An experimental approach and its interdependence with the time-lag estimation, Atmospheric Measurement Techniques, 14, 5071–5088, <https://doi.org/10.5194/amt-14-5071-2021>, 2021.
- Petersen, R., Holst, T., Mölder, M., Kljun, N., and Rinne, J.: Vertical distribution of sources and sinks of volatile organic compounds within a boreal forest canopy, Atmospheric Chemistry and Physics, 23, 7839–7858, <https://doi.org/10.5194/acp-23-7839-2023>, 2023.
- 1280 Pfannerstill, E. Y., Reijrink, N. G., Edtbauer, A., Ringsdorf, A., Zannoni, N., Araújo, A., Ditas, F., Holanda, B. A., Sá, M. O., Tsokankunku, A., Walter, D., Wolff, S., Lavrič, J. V., Pöhlker, C., Sörgel, M., and Williams, J.: Total OH reactivity over the Amazon rainforest: variability with temperature, wind, rain, altitude, time of day, season, and an overall budget closure, Atmospheric Chemistry and Physics, 21, 6231–6256, <https://doi.org/10.5194/acp-21-6231-2021>, 2021.
- Pleim, J. and Ran, L.: Surface Flux Modeling for Air Quality Applications, Atmosphere, 2, 271–302, <https://doi.org/10.3390/atmos2030271>,
1285 2011.
- Poraicu, C., Müller, J.-F., Stavrakou, T., Amelynck, C., Verreyken, B. W. D., Schoon, N., Vigouroux, C., Kumps, N., Brioude, J., Tulet, P., and Mouchel-Vallon, C.: Constraining the budget of NO_x and volatile organic compounds at a remote tropical island using multi-platform observations and WRF-Chem model simulations, Atmospheric Chemistry and Physics, 25, 6903–6941, <https://doi.org/10.5194/acp-25-6903-2025>, publisher: Copernicus GmbH, 2025.
- 1290 Potier, E., Ogée, J., Jouanguy, J., Lamaud, E., Stella, P., Personne, E., Durand, B., Mascher, N., and Loubet, B.: Multilayer modelling of ozone fluxes on winter wheat reveals large deposition on wet senescing leaves, Agricultural and Forest Meteorology, 211–212, 58–71, <https://doi.org/10.1016/j.agrformet.2015.05.006>, 2015.
- Potier, E., Loubet, B., Durand, B., Flura, D., Bourdat-Deschamps, M., Ciuraru, R., and Ogée, J.: Chemical reaction rates of ozone in water infusions of wheat, beech, oak and pine leaves of different ages, Atmospheric Environment, 151, 176–187,
1295 <https://doi.org/10.1016/j.atmosenv.2016.11.069>, 2017.
- Rannik, Ü., Vesala, T., and Keskinen, R.: On the damping of temperature fluctuations in a circular tube relevant to the eddy covariance measurement technique, Journal of Geophysical Research: Atmospheres, 102, 12 789–12 794, <https://doi.org/10.1029/97JD00362>, 1997.
- Raupach, M.: Applying Lagrangian fluid mechanics to infer scalar source distributions from concentration profiles in plant canopies, Agricultural and Forest Meteorology, 47, 85–108, [https://doi.org/10.1016/0168-1923\(89\)90089-0](https://doi.org/10.1016/0168-1923(89)90089-0), 1989.
- 1300 Rinne, J., Taipale, R., Markkanen, T., Ruuskanen, T. M., Hellén, H., Kajos, M. K., Vesala, T., and Kulmala, M.: Hydrocarbon fluxes above a Scots pine forest canopy: Measurements and modeling, Atmospheric Chemistry and Physics, 7, 3361–3372, <https://doi.org/10.5194/acp-7-3361-2007>, 2007.
- Rinne, J., Karl, T., and Guenther, A.: Simple, stable, and affordable: Towards long-term ecosystem scale flux measurements of VOCs, Atmospheric Environment, 131, 225–227, <https://doi.org/10.1016/j.atmosenv.2016.02.005>, 2016.
- 1305 Rowlinson, M. J., Rap, A., Hamilton, D. S., Pope, R. J., Hantson, S., Arnold, S. R., Kaplan, J. O., Arneth, A., Chipperfield, M. P., Forster, P. M., and Nieradzik, L.: Tropospheric ozone radiative forcing uncertainty due to pre-industrial fire and biogenic emissions, Atmospheric Chemistry and Physics, 20, 10 937–10 951, <https://doi.org/10.5194/acp-20-10937-2020>, 2020.
- Ruuskanen, T. M., Müller, M., Schnitzhofer, R., Karl, T., Graus, M., Bamberger, I., Hörtnagl, L., Brilli, F., Wohlfahrt, G., and Hansel, A.: Eddy covariance VOC emission and deposition fluxes above grassland using PTR-TOF, Atmospheric Chemistry and Physics, 11, 611–625,
1310 <https://doi.org/10.5194/acp-11-611-2011>, 2011.



- Sabbatini, S., Mammarella, I., Arriga, N., Fratini, G., Graf, A., Hörtnagl, L., Ibrom, A., Longdoz, B., Mauder, M., Merbold, L., Metzger, S., Montagnani, L., Pitacco, A., Rebmann, C., Sedláč, P., Šigut, L., Vitale, D., and Papale, D.: Eddy covariance raw data processing for CO₂ and energy fluxes calculation at ICOS ecosystem stations, *International Agrophysics*, 32, 495–515, <https://doi.org/10.1515/intag-2017-0043>, 2018.
- 1315 Safieddine, S. A., Heald, C. L., and Henderson, B. H.: The global nonmethane reactive organic carbon budget: A modeling perspective, *Geophysical Research Letters*, 44, 3897–3906, <https://doi.org/10.1002/2017GL072602>, 2017.
- Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry, C., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H. S., Kleinen, T., Krummel, P., Lamarque, J. F., Langenfelds, R., Locatelli, R., Machida, T., 1320 Maksyutov, S., McDonald, K. C., Marshall, J., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F. J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I., Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Steele, P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., Van Weele, M., Van Der Werf, G. R., Weiss, R., Wiedinmyer, C., Wilton, D. J., Wiltshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu, 1325 Q.: The global methane budget 2000–2012, *Earth System Science Data*, 8, 697–751, <https://doi.org/10.5194/essd-8-697-2016>, 2016.
- Scala, A., Allmann, S., Mirabella, R., Haring, M. A., and Schuurink, R. C.: Green leaf volatiles: A plant's multifunctional weapon against herbivores and pathogens, *International Journal of Molecular Sciences*, 14, 17 781–17 811, <https://doi.org/10.3390/ijms140917781>, 2013.
- Schallhart, S., Rantala, P., Nemitz, E., Taipale, D., Tillmann, R., Mentel, T. F., Loubet, B., Gerosa, G., Finco, A., Rinne, J., and Ruuskanen, T. M.: Characterization of total ecosystem-scale biogenic VOC exchange at a Mediterranean oak–hornbeam forest, *Atmospheric Chemistry and Physics*, 16, 7171–7194, <https://doi.org/10.5194/acp-16-7171-2016>, 2016.
- 1330 Schallhart, S., Rantala, P., Kajos, M. K., Aalto, J., Mammarella, I., Ruuskanen, T. M., and Kulmala, M.: Temporal variation of VOC fluxes measured with PTR-TOF above a boreal forest, *Atmospheric Chemistry and Physics*, 18, 815–832, <https://doi.org/10.5194/acp-18-815-2018>, 2018.
- Seco, R., Karl, T., Guenther, A., Hosman, K. P., Pallardy, S. G., Gu, L., Geron, C., Harley, P., and Kim, S.: Ecosystem-scale volatile organic compound fluxes during an extreme drought in a broadleaf temperate forest of the Missouri Ozarks (central USA), *Global Change Biology*, 21, 3657–3674, <https://doi.org/10.1111/gcb.12980>, 2015.
- 1335 Seco, R., Karl, T., Turnipseed, A., Greenberg, J., Guenther, A., Llusia, J., Peñuelas, J., Dicken, U., Rotenberg, E., Kim, S., and Yakir, D.: Springtime ecosystem-scale monoterpene fluxes from Mediterranean pine forests across a precipitation gradient, *Agricultural and Forest Meteorology*, 237–238, 150–159, <https://doi.org/10.1016/j.agrformet.2017.02.007>, 2017.
- 1340 Sekimoto, K., Li, S.-M., Yuan, B., Koss, A., Coggon, M., Warneke, C., and de Gouw, J.: Calculation of the sensitivity of proton-transfer-reaction mass spectrometry (PTR-MS) for organic trace gases using molecular properties, *International Journal of Mass Spectrometry*, 421, 71–94, <https://doi.org/10.1016/j.ijms.2017.04.006>, 2017.
- Seliger, A., Ammer, C., Kreft, H., and Zerbe, S.: Diversification of coniferous monocultures in the last 30 years and implications for forest restoration: a case study from temperate lower montane forests in Central Europe, *European Journal of Forest Research*, 142, 1353–1368, <https://doi.org/10.1007/s10342-023-01595-4>, 2023.
- 1345 Seyednasrollah, B., Young, A. M., Hufkens, K., Milliman, T., Friedl, M. A., Frolking, S., and Richardson, A. D.: Tracking vegetation phenology across diverse biomes using Version 2.0 of the PhenoCam Dataset, *Scientific Data*, 6, 1–11, <https://doi.org/10.1038/s41597-019-0229-9>, 2019.



- Simon, L., Gros, V., Petit, J.-E., Truong, F., Sarda-Estève, R., Kalalian, C., Baudic, A., Marchand, C., and Favez, O.: Two years of volatile
 1350 organic compound online in situ measurements at the Site Instrumental de Recherche par Télédétection Atmosphérique (Paris region,
 France) using proton-transfer-reaction mass spectrometry, *Earth System Science Data*, 15, 1947–1968, <https://doi.org/10.5194/essd-15-1947-2023>, 2023.
- Song, J., Gkatzelis, G. I., Tillmann, R., Brüggemann, N., Leisner, T., and Saathoff, H.: Characterization of biogenic volatile organic com-
 pounds and their oxidation products in a stressed spruce-dominated forest close to a biogas power plant, *Atmospheric Chemistry and*
 1355 *Physics*, 24, 13 199–13 217, <https://doi.org/10.5194/acp-24-13199-2024>, publisher: Copernicus GmbH, 2024.
- Stockwell, W. R. and Forkel, R.: Ozone and volatile organic compounds: isoprene, terpenes, aldehydes, and organic acids, pp. 257–276,
 Springer Netherlands, Dordrecht, ISBN 978-94-015-9856-9, https://doi.org/10.1007/978-94-015-9856-9_14, 2002.
- Striednig, M., Graus, M., Märk, T., and Karl, T.: InnFLUX-an open-source code for conventional and disjunct eddy covariance analysis
 of trace gas measurements: An urban test case, *Atmospheric Measurement Techniques*, 13, 1447–1465, [https://doi.org/10.5194/amt-13-
 1360 *1447-2020*, 2020.](https://doi.org/10.5194/amt-13-1447-2020)
- Taipale, R., Ruuskanen, T. M., and Rinne, J.: Lag time determination in DEC measurements with PTR-MS, *Atmospheric Measurement*
Techniques, 3, 853–862, <https://doi.org/10.5194/amt-3-853-2010>, taipale, R. Ruuskanen, T. M. Rinne, J., 2010.
- Tiwary, A., Fuentes, J. D., Barr, J. G., Wang, D., and Colls, J. J.: Inferring the source strength of isoprene from ambient concentrations,
Environmental Modelling Software, 22, 1281–1293, <https://doi.org/10.1016/j.envsoft.2006.09.002>, 2007.
- 1365 Vermeuel, M. P., Cleary, P. A., Desai, A. R., and Bertram, T. H.: Simultaneous Measurements of O₃ and HCOOH Vertical Fluxes Indicate
 Rapid In-Canopy Terpene Chemistry Enhances O₃ Removal Over Mixed Temperate Forests, *Geophysical Research Letters*, 48, 1–15,
<https://doi.org/10.1029/2020GL090996>, 2021.
- Vermeuel, M. P., Millet, D. B., Farmer, D. K., Pothier, M. A., Link, M. F., Riches, M., Williams, S., and Garofalo, L. A.: Closing the
 Reactive Carbon Flux Budget: Observations From Dual Mass Spectrometers Over a Coniferous Forest, *Journal of Geophysical Research:*
 1370 *Atmospheres*, 128, 1–20, <https://doi.org/10.1029/2023JD038753>, 2023a.
- Vermeuel, M. P., Novak, G. A., Kilgour, D. B., Clafin, M. S., Lerner, B. M., Trowbridge, A. M., Thom, J., Cleary, P. A., Desai, A. R.,
 and Bertram, T. H.: Observations of biogenic volatile organic compounds over a mixed temperate forest during the summer to autumn
 transition, *Atmospheric Chemistry and Physics*, 23, 4123–4148, <https://doi.org/10.5194/acp-23-4123-2023>, 2023b.
- Vermeuel, M. P., Millet, D. B., Farmer, D. K., Ganzeveld, L. N., Visser, A. J., Alwe, H. D., Bertram, T. H., Cleary, P. A., Desai, A. R.,
 1375 Helmig, D., Kavassalis, S. C., Link, M. F., Pothier, M. A., Riches, M., Wang, W., and Williams, S.: A Vertically Resolved Canopy
 Improves Chemical Transport Model Predictions of Ozone Deposition to North Temperate Forests, *Journal of Geophysical Research:*
Atmospheres, 129, <https://doi.org/10.1029/2024jd042092>, publisher: American Geophysical Union (AGU), 2024.
- Verreyken, B., Amelynck, C., Schoon, N., Müller, J.-F., Brioude, J., Kumps, N., Hermans, C., Metzger, J.-M., Colomb, A., and Stavrakou, T.:
 Measurement report: Source apportionment of volatile organic compounds at the remote high-altitude Maïdo observatory, *Atmospheric*
 1380 *Chemistry and Physics*, 21, 12 965–12 988, <https://doi.org/10.5194/acp-21-12965-2021>, 2021.
- Verreyken, B. W. D., Dumont, C., Schoon, N., Heinesch, B., and Amelynck, C.: 3-years of (O)VOC and ozone concentration measurements
 at a mixed temperate forest at the Vielsalm ICOS ecosystem station (Belgium), <https://doi.org/10.18758/NVFBA74V>, 2025.
- Vickers, D. and Mahrt, L.: Quality control and flux sampling problems for tower and aircraft data, *Journal of Atmospheric and Oceanic*
Technology, 14, 512–526, [https://doi.org/10.1175/1520-0426\(1997\)014<0512:QCAFSP>2.0.CO;2](https://doi.org/10.1175/1520-0426(1997)014<0512:QCAFSP>2.0.CO;2), 1997.



- 1385 Vincke, C., Bogaerts, G., Chebbi, W., Chebbi, W., Chopin, H., Demoulin, L., Douxfls, B., Engelmann, T., Faurès, A., Heinesch, B., Manise, T., Orgun, A., Piret, A., and Thyron, T.: ETC L2 ARCHIVE from Vielsalm, 2020–2024, <https://hdl.handle.net/11676/HaUqvLvNC5h4zUO2UY4CMIfR>, 2025.
- Vitale, D., Fratini, G., Bilancia, M., Nicolini, G., Sabbatini, S., and Papale, D.: A robust data cleaning procedure for eddy covariance flux measurements, *Biogeosciences*, 17, 1367–1391, <https://doi.org/10.5194/bg-17-1367-2020>, 2020.
- 1390 Wada, R., Ueyama, M., Tani, A., Mochizuki, T., Miyazaki, Y., Kawamura, K., Takahashi, Y., Saigusa, N., Takanashi, S., Miyama, T., Nakano, T., Yonemura, S., Matsumi, Y., and Kataka, G.: Observation of vertical profiles of NO, O₃, and VOCs to estimate their sources and sinks by inverse modeling in a Japanese larch forest, *Journal of Agricultural Meteorology*, 76, 1–10, <https://doi.org/10.2480/agrmet.D-18-00029>, 2020.
- Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour transfer, *Quarterly Journal of the Royal Meteorological Society*, 106, 85–100, <https://doi.org/10.1002/qj.49710644707>, 1980.
- 1395 Wesely, M.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmospheric Environment* (1967), 23, 1293–1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.
- Wilczak, J. M., Oncley, S. P., and Stage, S. A.: Sonic anemometer tilt correction algorithms, *Boundary-Layer Meteorology*, 99, 127–150, <https://doi.org/10.1023/A:1018966204465>, 2001.
- 1400 Wohlfahrt, G., Amelynck, C., Ammann, C., Arneth, A., Bamberger, I., Goldstein, A. H., Gu, L., Guenther, A., Hansel, A., Heinesch, B., Holst, T., Hörtnagl, L., Karl, T., Laffineur, Q., Neftel, A., McKinney, K., Munger, J. W., Pallardy, S. G., Schade, G. W., Seco, R., and Schoon, N.: An ecosystem-scale perspective of the net land methanol flux: synthesis of micrometeorological flux measurements, *Atmospheric Chemistry and Physics*, 15, 7413–7427, <https://doi.org/10.5194/acp-15-7413-2015>, 2015.
- Wolfe, G. M., Thornton, J. A., McKay, M., and Goldstein, A. H.: Forest-atmosphere exchange of ozone: Sensitivity to very reactive biogenic VOC emissions and implications for in-canopy photochemistry, *Atmospheric Chemistry and Physics*, 11, 7875–7891, <https://doi.org/10.5194/acp-11-7875-2011>, 2011.
- 1405 Yáñez-Serrano, A., Filella, I., LLusià, J., Gargallo-Garriga, A., Granda, V., Bourtsoukidis, E., Williams, J., Seco, R., Cappellin, L., Werner, C., de Gouw, J., and Peñuelas, J.: GLOVOCS - Master compound assignment guide for proton transfer reaction mass spectrometry users, *Atmospheric Environment*, 244, 117 929, <https://doi.org/10.1016/j.atmosenv.2020.117929>, 2021.
- 1410 Yang, M., Beale, R., Smyth, T., and Blomquist, B.: Measurements of OVOC fluxes by eddy covariance using a proton-transfer- reaction mass spectrometer-method development at a coastal site, *Atmospheric Chemistry and Physics*, 13, 6165–6184, <https://doi.org/10.5194/acp-13-6165-2013>, 2013.
- Yoon, J. Y. S., Wells, K. C., Millet, D. B., Swann, A. L. S., Thornton, J., and Turner, A. J.: Impacts of Interannual Isoprene Variations on Methane Lifetimes and Trends, *Geophysical Research Letters*, 52, <https://doi.org/10.1029/2025GL114712>, 2025.
- 1415 Zhang, L., Brook, J. R., and Vet, R.: On ozone dry deposition—with emphasis on non-stomatal uptake and wet canopies, *Atmospheric Environment*, 36, 4787–4799, [https://doi.org/10.1016/S1352-2310\(02\)00567-8](https://doi.org/10.1016/S1352-2310(02)00567-8), 2002.
- Zhou, P., Ganzeveld, L., Rannik, Ü., Zhou, L., Gierens, R., Taipale, D., Mammarella, I., and Boy, M.: Simulating ozone dry deposition at a boreal forest with a multi-layer canopy deposition model, *Atmospheric Chemistry and Physics*, 17, 1361–1379, <https://doi.org/10.5194/acp-17-1361-2017>, 2017a.
- 1420 Zhou, P., Ganzeveld, L., Taipale, D., Rannik, , Rantala, P., Rissanen, M. P., Chen, D., and Boy, M.: Boreal forest BVOC exchange: emissions versus in-canopy sinks, *Atmospheric Chemistry and Physics*, 17, 14 309–14 332, <https://doi.org/10.5194/acp-17-14309-2017>, 2017b.



Zhu, Z., Zhao, F., Voss, L., Xu, L., Sun, X., Yu, G., and Meixner, F. X.: The effects of different calibration and frequency response correction methods on eddy covariance ozone flux measured with a dry chemiluminescence analyzer, *Agricultural and Forest Meteorology*, 213, 114–125, <https://doi.org/10.1016/j.agrformet.2015.06.016>, 2015.