



Two Biogenic Volatile Organic Compound Emission Datasets over Europe Based on Land Surface Modelling and Satellite Data Assimilation

Paul D. Hamer¹, Miha Markelj¹, Oscar Rojas-Munoz², Bertrand Bonan², Jean-Christophe Calvet²,
5 Virginie Marécal², Alex Guenther³, Heidi Trimmel^{4,5}, Islen Vallejo¹, Sabine Eckhardt¹, Gabriela Sousa
Santos¹, Katerina Sindelarova⁶, David Simpson^{7,8}, Norbert Schmidbauer¹, Leonor Tarrasón¹

¹NILU, 2027 Kjeller, Norway

²Centre National de Recherches Météorologiques, Université de Toulouse, Météo-France, CNRS, Toulouse, France

³Department of Earth System Science, University of California at Irvine, Irvine, CA 92697, USA

10 ⁴International Institute for Applied Systems Analysis (IIASA), Schloßplatz 1, 2361, Laxenburg, Austria

⁵University of Natural Resources and Life Sciences, Vienna, Department of Water, Atmosphere and Environment, Institute
of Meteorology and Climatology, Gregor-Mendel-Straße 33, 1180, Vienna, Austria

⁶Department of Atmospheric Physics, Faculty of Mathematics and Physics, Charles University, Prague, V Holešovičkách 2,
18000 Prague 8, Czechia

15 ⁷Division for Climate Modelling and Air Pollution, Norwegian Meteorological Institute, 0313 Oslo, Norway

⁸Department of Space, Earth and Environment, Chalmers University of Technology, 412 96 Gothenburg, Sweden

Correspondence to: Paul D. Hamer (paul.d.hamer@gmail.com)

Abstract. Biogenic volatile organic compound (BVOC) emissions from vegetation represent a major source of volatile
20 compounds globally and play an important role as precursors for tropospheric ozone. Understanding their emissions is
therefore crucial for quantifying the impact of ozone on air quality. We present two datasets of biogenic volatile organic
compound emissions that cover the European modelling domain of the Copernicus Atmospheric Monitoring Service at a
resolution of $0.1^\circ \times 0.1^\circ$ to support the study of European scale air quality. The compounds included in the dataset follow the
VOCs included in the regional atmospheric chemistry model mechanism (RACM). The datasets were produced within the
25 framework of the EU's SEEDS project. We produced each dataset by coupling modelling output variables from the
SURFEX land surface model with the MEGAN3.0 BVOC emission model. In one instance, the SURFEX model was run in
free-running mode, which we term the open-loop (OL) and in the other case we assimilated satellite observations of leaf area
index (LAI), which we term the analysis. The OL and analysis land surface model outputs form the basis for each emission
dataset that are called SURFEX-MEGAN3.0 OL and SURFEX-MEGAN3.0 analysis, respectively. The OL dataset is
30 available over a five-year period from 2018-2022 and the analysis dataset is available over the three-year period 2018-2020.
SURFEX was run for both the OL and analysis simulations in a configuration that allowed simulated vegetation to respond
to variations in meteorology over time to more realistically track vegetation phenology. Evaluation of the land surface model
output LAI and root-zone soil moisture (RZSM) showed that the OL and analysis simulations had good skill at tracking
temporal changes in both variables, with the analysis performing better in each instance. We perform a variety of evaluations



on the isoprene emissions specifically given the importance of this compound for atmospheric chemistry. We evaluated the temporal variability of isoprene emissions in both datasets and found that the majority of the interannual and monthly variability was linked to variability in LAI that in specific cases, like the summer of 2019, could be linked to drought impacts on vegetation growth simulated by SURFEX. We evaluated the daily temporal variability of the OL and analysis isoprene emission datasets against in-situ online observations of isoprene concentrations at 8 sites in western Europe and found moderate to strong correlation between the emissions and observations in almost all location-year pairings. We also evaluated the OL and analysis emission datasets against other published bottom-up isoprene emission datasets over the same European domain used in this study. We found that the SURFEX-MEGAN3.0 OL and analysis isoprene emission datasets lie between the minimum (CAMSGLOB-BIOv3.1) and maximum (MEGAN-MACC) published emission datasets based on bottom-up approaches. Furthermore, we were able to attribute differences in seasonality between SURFEX-MEGAN3.0 and other emission inventories to differences in the temporal variability of the underlying LAI dataset used to compile them. Overall, our findings show the importance of variability in LAI in controlling isoprene emissions on monthly to annual timescales. Combining this with the demonstrated skill of the emissions in evaluation with independent data, this points towards the value of an Earth-system approach to BVOC emission modelling.

1 Introduction

Vegetation represents a major source of volatile organic compounds (VOCs) to Earth's atmosphere with estimates indicating that biogenic VOCs (so-called, BVOCs) contribute to approximately 90% of all the VOCs emitted to the atmosphere (Guenther et al., 1995, 2012; Lin et al., 2021). The remaining portion is anthropogenic in origin. VOCs play a critical role in atmospheric chemistry and mediate photochemical ozone formation (itself a greenhouse gas and air pollutant at the surface) as ozone precursors, and impact the oxidation capacity of the atmosphere, which can, in turn, impact the levels of the greenhouse gas methane (Thornhill et al., 2021). VOCs (including the BVOCs) also contribute to secondary organic aerosol formation (Carlton et al., 2009; Donahue et al., 2009), which also creates a radiative forcing effect (Sporre et al., 2019). Through these means, BVOCs, as the dominant component of VOCs at the global scale, have an important influence on atmospheric composition, surface air quality, and climate.

Plants produce and release BVOCs during normal aspects of their growth, development, and senescence (Laothawornkitkul et al., 2009). Plants also deliberately release BVOCs for airborne signalling between individual plants for elicitation or priming of plant defences, signalling via attractor molecules to pollinators and seed dispersers, and for protection against pathogens (Laothawornkitkul et al., 2009), e.g., by direct deterrence or via attraction of the predators of herbivores. Lastly, damage to plants carried out either by herbivores or occurring from environmental factors such as ozone damage, high wind damage, or extreme temperature stress can lead to the release of BVOCs from within the stressed, exposed, or damaged plant tissue.



Plant emission of BVOCs varies in time according to changing environmental conditions, which can either result in the stimulation of plant growth or lead to biotic and abiotic stresses. For instance, diurnal variations in solar radiation and ambient temperature as well as soil moisture availability affect the photosynthetic activity and in turn impact emissions. For example, (Guenther et al., (1993) shows that for a variety of species, isoprene emissions increase as ambient and leaf

70 temperatures increase until a certain point is reached over $\sim 40^{\circ}\text{C}$ at which point emissions of isoprene, and other de novo BVOC lacking a storage reservoir, begin to decline. In fact, leaf temperature is the most important variable since it has a more direct control and link to the plant physiological processes and underlying enzymatic activity that controls their metabolism. Thus, direct and indirect radiation play an important role in controlling BVOC emissions. As another consequence of this, shading of leaves within the canopy is another important process that impacts BVOC emissions. Given

75 the importance of temperature for controlling emissions, there is a strong connection between BVOC emissions and heat waves whereby the hot weather can stimulate emissions of BVOCs over wide areas that can then in turn lead to significant photochemical ozone formation. Heatwaves can also interact with drought as the hot weather can act to increase the rate at which soils dry out and dry soils also increase the proportion of absorbed solar insolation that gets converted to sensible heat. Thus, droughts and heatwaves often occur concurrently and so the combined effects of both on BVOC emissions must be

80 considered. Indeed, droughts can lead to reductions in BVOC emissions even during heatwaves as the water stress impairs photosynthesis and plant physiology. Drought stress can also lead to secondary effects that reduce BVOC emissions as the water stress can lead to browning and die-off of leaves, which can be observed as reductions in leaf area index (LAI) even before the growing season has culminated. In addition, phenological vegetation changes during the growing season, e.g., leaf surface area and leaf age, as well as longer-term changes associated with overall plant age can also impact emissions.

85 BVOC fluxes can be measured for individual plant species at leaf level in field or laboratory conditions or for whole ecosystem types using field eddy covariance flux tower measurements. A key limitation of emission monitoring is the lack of spatial coverage of the observing network of BVOC fluxes. Furthermore, the implied heterogeneity in BVOC emissions arising from the varying spatial distribution of emitting species coupled with the spatio-temporal variability in driving environmental conditions strongly limits the spatial representativeness of BVOC flux observations at specific locations. This

90 problem places limitations on attempts to directly extrapolate the spatially-sparse observations of BVOC emissions to estimate emissions at the global scale. Similarly, this problem also limits attempts to spatially interpolate the observed emission data between study sites in order to spatially map emissions. These challenges create the need for modelling approaches that can estimate BVOC emissions with spatial continuity over the Earth's surface while considering the spatial distribution of emitting species and the spatio-temporal variability of driving environmental conditions. Such modelling approaches could thus deliver spatial gap-filling in regions lacking observations, which would have scientific value for studying air quality, atmospheric composition, and climate. Given this need, there have naturally been different efforts and approaches developed to model BVOC emissions from vegetation.

Modelling approaches that have been used to estimate BVOC emissions can be separated into three broad categories: 1) empirically-based bottom-up models, e.g., different versions of MEGAN (Guenther et al., 1995, 2020, 2012), IBIS (Naik et



100 al., 2004), and EMEP (Simpson et al., 1999, 2012); 2) bottom-up models that attempt to simulate plant physiological leaf
 processes associated with photosynthesis, e.g., LPJ-GUESS (Arneth et al., 2007, 2011; Pacifico et al., 2011; Young et al.,
 2009); and 3) top-down models that use observations of the ambient concentrations of the BVOC oxidation product, HCHO,
 to constrain emissions using inversion methods (Millet et al., 2006; Oomen et al., 2024; Palmer et al., 2003; Stavrakou et al.,
 2009). Guenther et al. (1993) and references therein document a much more extensive history of modelling isoprene
 105 emissions that use empirically based approaches.

Here we focus on the bottom-up BVOC model MEGAN, which has become a widely used tool in the atmospheric science
 community and in particular in the field of air quality modelling and forecasting. Sindelarova et al. (2014) and references
 therein provide a useful overview of some of the applications of MEGAN and includes a compilation of the range in
 emissions estimated at the global scale as well as an example application of its own. Different efforts to estimate BVOC
 110 emissions using the bottom-up approach result in global isoprene emission estimates ranging between 410 and 680 Tg yr⁻¹
 (Arneth et al., 2007, 2011; Emmons et al., 2010; Guenther et al., 2006, 2012; Lathière et al., 2006, 2010; Levis et al., 2003;
 Müller et al., 2008; Naik et al., 2004; Pacifico et al., 2011; Pfister et al., 2008; Potter et al., 2001; Shim et al., 2005;
 Stavrakou et al., 2009; Tao and Jain, 2005; Wiedinmyer et al., 2006; Young et al., 2009).

The large range in global BVOC emission estimates from MEGAN results from a significant uncertainty in the input data
 115 and the underlying mechanisms that drive BVOC emission estimation. Guenther et al. (2006) and references therein provide
 an excellent summary of some of the sources of uncertainty that arise from input data to the MEGAN model. For instance,
 use of differing LAI datasets can lead to relative changes in global isoprene emission estimates from -11% up to +29%
 (Guenther et al., 2006). Meanwhile, the use of different meteorological input datasets has a somewhat smaller relative impact
 on isoprene emission estimates with a range from -11% up to +15%. Guenther et al. (2006) also estimate that variations in
 120 plant function type mapping can impact isoprene emissions by up to -13% to +18%. Emission factors, also known as
 emission potentials, represent the emission from a unit leaf area or unit land surface area under a standard set of ambient
 conditions, which in turn get scaled upwards or downwards depending on the meteorological conditions at any given point in
 time. Emission factors have to be estimated from maps of vegetation type, climatic zone, and known emission rates from
 different plant species. Emission factors therefore represent another significant source of uncertainty (Langford et al., 2017),
 125 with the use of differing emissions factor datasets having a significant impact. A recent study (Sindelarova et al., 2022)
 showed that using different emission factors can lead to changes in global isoprene emission estimates of up to 33%. Lastly,
 there is no current consensus on how to account for drought effects on isoprene emissions using MEGAN. For instance,
 Sindelarova et al. (2014, 2022) both choose not to use the MEGAN soil moisture algorithm in their reference dataset and
 found that using the algorithm can reduce global isoprene emission estimates by up to 50%. Furthermore, estimation of soil
 130 moisture itself is also likely to be a source of uncertainty.

Our motivation in this present study is to present two new BVOC emission datasets for use in the study of European air
 quality developed with the specific aim to address key uncertainties in BVOC emissions arising from the representation of
 vegetation phenology. We try to address this uncertainty by coupling MEGAN with the outputs (soil and vegetation



variables) from a well-validated land surface model (Masson et al., 2013) capable of detailed representations of vegetation dynamics (Calvet et al., 1998; Gibelin et al., 2006) and soil moisture (Decharme et al., 2011). We try to further address this uncertainty by assimilating satellite observations of LAI in the land surface model. Thus, we create two distinct BVOC emission datasets, one based on the free-running land surface model run, or so-called open-loop (2018-2022), and another based on the assimilation analysis (2018-2022) made by assimilating satellite LAI observations. Note, that the assimilation-based dataset has a shorter timeframe due to the limited availability of suitable LAI observations. Both datasets can be used to allow an Earth-systems approach to permit feedbacks and interactions between meteorology/climate, vegetation, and atmospheric composition.

Since our motivation was to develop BVOC emission datasets that are relevant for current-day European air quality modelling, we quickly highlight the current state of the art of the BVOC modelling over Europe using MEGAN for cases that have a focus on air quality. MEGAN has been used in some prominent examples of open-source BVOC emission datasets e.g., MEGAN-MACC (Sindelarova et al., 2014) and CAMS-GLOB-BIO (Sindelarova et al., 2022) that are designed for use in applications related to air quality and atmospheric composition simulation. The more recent example, which is CAMS-GLOB-BIOv3.1, had a spatial resolution of $0.25^\circ \times 0.25^\circ$, was developed as part of the Copernicus Atmospheric Monitoring Service (CAMS), and was aimed at delivering improved estimates of BVOC emissions to support air quality modelling both in Europe and at the global scale. CAMS produces operational air quality forecasts and reanalyses at the global and regional scales and the activity within CAMS to produce the CAMS-GLOB-BIO emissions can be seen as a supporting activity both to the CAMS air quality modelling and air quality modelling activities external to CAMS itself. The emission datasets presented within this paper were produced within the frame of the EU funded Sentinel EO-based Emission and Deposition Service (SEEDS) project (<https://www.seedsproject.eu/>, last access: 17 July 2025). One aim of SEEDS was to produce emission datasets of pollutant emissions based on Earth observation data in support and development of the CAMS European air quality forecasting. Therefore, one aim of this present study is to develop a new methodology that can be used to create a BVOC emission dataset specifically for use in the CAMS European regional modelling activities.

In addition to the targeted advancements in phenological modelling and data assimilation, we also implemented some further advancements relative to previous work (e.g., Sindelarova et al., 2022) and detail these below:

- i) Estimation of BVOC emissions at higher spatial resolution ($0.1^\circ \times 0.1^\circ$) and temporal resolution (1-hour) compared to existing publicly available BVOC emission datasets over Europe, e.g., CAMS-GLOB-BIO (Sindelarova et al., 2022).
- ii) Using the state-of-the-art land cover maps over Europe, ECOCLIMAP-II (Faroux et al., 2013), which contain specific adaptations of the CORINE land cover to European conditions.
- iii) Updated MEGAN version to v3.0 (Guenther et al., 2020; Zhang et al., 2021). Many existing publicly available BVOC emission datasets and examples of applications in the literature use MEGAN 2.1.

In Section 2 we present the methodology including the emission model, the land surface model, and the other datasets used in the production of the emission data including the meteorological forcing. In Section 3 we present the results including



spatiotemporal analysis of the BVOC emission inventories and a comparison of the BVOC emissions with other emission datasets and observations of isoprene emissions. In Section 4 we describe how to access the datasets. In Section 5 we present the conclusions.

2 Methodology

2.1 BVOC emission modelling system overview

We first present an overview of the entire modelling system that we use to estimate BVOC emissions and create the two datasets presented in this paper (Hamer et al., 2025a, Hamer et al., 2025b). This is in order to highlight that our method consists of a unique framework of different modelling elements and data sources and to properly explain the various connections and workflow between each of these components. Figure 1 shows a schematic flow diagram of our methodological framework that highlights each modelling component and how they connect together. The modelling components (shown in black boxes) consist of the SURFEX land surface model and its data assimilation system, the MEGAN emission factor processor (MEGAN EFP), and the MEGAN3.0 BVOC emission model. We use a variety of data sources (shown in grey boxes) that fit into each modelling component. The first step in this framework involves running the SURFEX land surface model, which primarily uses the ECMWF HRES meteorology and the ECOCLIMAP-II land cover as inputs. In addition, when SURFEX is run using the LDAS-Monde data assimilation algorithm it also ingests LAI satellite observations to produce the assimilation analysis output. When SURFEX is run without the assimilation step in free-running mode this output is termed the open-loop. These two separate data flows from SURFEX (highlighted in blue boxes within Figure 1) are fed separately into the MEGAN3.0 model, and along with the emission factors from the MEGAN EFP. Each of these produce a separate corresponding output BVOC emission dataset (i.e., open-loop and assimilation analysis) shown in the green boxes. Further details of the MEGAN3.0 emission model and the MEGAN EFP are discussed in the following section (Section 2.2) and are shown in Figure 2.




Figure 1. A flow diagram showing how the different modelling elements within this BVOC modelling framework link together. Model components are shown in the black boxes, external input data sources are shown in grey boxes, intermediate data sources are shown in blue boxes, and output data are shown in green boxes. There are three data flows (ECMWF HRES meteorology, ECOCLIMAP-II land surface cover, and satellite observations of LAI based on PROBA-V and AVHRR data) into the SURFEX LDAS-Monde land surface modelling and data assimilation system. The output from SURFEX serves as the inputs into MEGAN3.0. There are two additional dataflows into the MEGAN3.0 model, which are emission factors calculated by the MEGAN EFP and the ECMWF HRES meteorology. We produce two differing BVOC emission datasets (open-loop and assimilation analysis) based on the corresponding output datasets from SURFEX of each type.

2.2 The MEGAN 3.0 Emission model

2.2.1 Model overview

We first provide an overview of the basic conceptual framework built within MEGAN3.0 (Guenther et al., 2020). This concept can be described in a simple equation $F_i = \gamma_i \times \sum_{k=1}^n (\varepsilon_{i,k} \times \chi_k)$ (1) defining the net emission flux into the above-canopy atmosphere of species, i as, F_i .

$$F_i = \gamma_i \times \sum_{k=1}^n (\varepsilon_{i,k} \times \chi_k) \quad (1)$$

Here, γ_i is the chemical species-specific activity, $\varepsilon_{i,k}$ is the emission factor at standard conditions for vegetation type k , and χ_k is the fractional grid box areal coverage. The activity represents changing environmental conditions (e.g., air temperature and radiation) and vegetation properties (e.g., LAI) that impact BVOC emissions. Activity is calculated by multiplying a set of different γ activity parameters together in series (explained in more detail in Sect. 2.1.2) where each parameter represents a different process impacting activity. The emission factors are a measure of the potential emissions of a particular compound from a specific plant type. As a whole, this conceptual framework is expressed via algorithms within a software package containing the MEGAN3.0 model (consisting of five sub-components) and as well a separate MEGAN emission factor pre-



processor (MEGAN EFP). We provide a schematic diagram in Figure 2 giving an overview of MEGAN3.0 and its five sub-components, the MEGAN EFP, the dependencies on external data sources at each step, and the sequential steps used in the production of the intermediate datasets that all eventually lead to the final output BVOC emissions. These intermediate
 215 datasets are associated with specific environmental processes that impact activity (e.g., soil moisture) as well as physical properties affecting leaf physiology, e.g., temperature and radiation.

We now describe the five sub-components of MEGAN3.0 in turn. The first is DAYMET, which calculates daily meteorological parameters using ECMWF (in our case) wind speed and accumulated photosynthetically active radiation (PAR), and 2m-temperature from the SURFEX land surface model. DAYMET calculates the daily mean, minimum, and
 220 maximum temperature, the daily mean photosynthetic photon flux density (PPFD), and the maximum daily wind speed.


The second sub-component is CANMET, which is a canopy meteorological model that calculates shaded and sunny leaf temperatures and PPFDs for different vertical layers in the vegetation canopy as well as the fraction of sunny leaves at each canopy layer height. CANMET is important for calculating the transmission of solar radiation into the lower layers of the of dense canopy when LAI is high, i.e., $> 3 \text{ m}^2.\text{m}^{-2}$. In the configuration used in the SEEDS project, CANMET uses wind
 225 speed, surface pressure and accumulated PAR from ECMWF's IFS HRES data, and 2m-temperature and 2m-specific humidity calculated within SURFEX.

The third sub-component within MEGAN3.0 is MEGSEA, which is responsible for calculating the effects of soil physical properties estimated by the SURFEX land surface model (root zone soil moisture and root zone soil temperature) on BVOC and soil NO_x emissions. The outputs from MEGSEA are the soil moisture activity parameter, γ_{SM} , and the soil NO_x activity
 230 parameter, which are both dependent on soil moisture and are sensitive to drought effects.

The fourth sub-component is MEGVEA, which uses parameters calculated by the preceding sub-components (DAYMET, CANMET, and MEGSEA) to calculate all of the remaining activity parameters (more details in Sect. 2.2.2) associated with different physical and biological processes. The code then calculates the product of all of these parameters to then calculate and output the activity for a range of aggregated BVOC species.

Prior to describing the final sub-component of MEGAN3.0 it is appropriate to briefly describe MEGAN EFP. MEGAN EFP calculates the emission factors and light-dependent factors over the chosen spatial grid using a set of maps of plant ecotype and growth form as input. The calculation of the emission factors and the MEGAN EFP software package are described in more detail in Sect. 2.2.3.

The final sub-component with MEGAN3.0 is MGN2MECH. MGN2MECH performs a dual function. First, it calculates the emissions of BVOCs by multiplying the emission factors and activity according to Eq. (1). Second, it performs a
 240 reaggregation of the BVOC emissions to derive an output emissions dataset that is compatible with the selected chemical mechanism. In our example case this is for the RACM chemical mechanism (Stockwell et al., 1997). The chemical species associated with RACM mechanism that are included in the datasets are described in detail in Table S1 in the supplement.



245 **Figure 2. A detailed flow diagram of the MEGAN3.0 BVOC emission model as used in our application. The five modelling components of MEGAN3.0 are shown in the coloured boxes (DAYMET in dark blue, CANMET in light green, MEGSEA in orange, MEGVEA in dark green, MGN2MECH in red). Note that MEGAN3.0 has to be run sequentially in this order. The MEGAN emission factor pre-processor shown in a light blue box must be run prior to the final MEGAN3.0 step (MGN2MECH). Note that the input and output flows for each model sub-component are represented using corresponding colour coded arrows.**
 250 **The various intermediate inputs and outputs are shown in light grey boxes. Lastly, the output BVOC emissions calculated by MEGAN3.0 from the MGN2MECH routine are shown in the grey-filled box.**

2.2.2 Emission Activity

The activity, γ_i , represents the vegetation BVOC emission response to changing environmental conditions. γ_i is calculated by multiplying together a series of other parameters that represent vegetation properties and responses to changing environmental conditions. We describe how γ_i is calculated for different chemical species and then describe how the parameters most relevant to this study (g_{TP} and g_{SM}) are calculated.

We define the standard activity that applies to the generic chemical species defined within MEGAN3.0 (i.e., this excludes isoprene, ethanol, acetaldehyde, and carbon monoxide) as:

$$\gamma_i = LAI \times \gamma_{TP} \times \gamma_{LA} \times \gamma_{HW} \times \gamma_{AQ} \times \gamma_{HT} \times \gamma_{LT} \times \gamma_{SM} \times LDF$$



260 Here, the activity used for BVOC chemical species that have a generic response are γ_{TP} , which is the canopy average temperature and radiation response parameter, γ_{LA} is the leaf age response parameter, γ_{HW} is the response to high windstorms parameter, γ_{AQ} is the response to air pollution (ozone) parameter, γ_{HT} is the response to high temperature parameter, γ_{LT} is the response to low temperature parameter, γ_{SM} is the soil moisture response parameter, and LDF is the light dependent fraction (the LDF is explained further in Sect. 2.2.3).

265 Next, we first define the emission activity for isoprene as:

$$g_{isoprene} = LAI \times \gamma_{TP} \times \gamma_{CO_2} \times \gamma_{LA} \times \gamma_{HW} \times \gamma_{AQ} \times \gamma_{HT} \times \gamma_{LT} \times \gamma_{SM} \times LDF$$

Where γ_{CO_2} defines the carbon dioxide response, which is only relevant for isoprene and thus makes the isoprene emission activity slightly different from all of the other chemical species. Next, we define the activity for ethanol and acetaldehyde:

$$\gamma_{ethanol \text{ or } acetaldehyde} = LAI \times \gamma_{TP} \times \gamma_{BD} \times \gamma_{LA} \times \gamma_{HW} \times \gamma_{AQ} \times \gamma_{HT} \times \gamma_{LT} \times \gamma_{SM}$$

270 Where γ_{BD} is the bidirectional exchange LAI response parameter, which is only relevant for both ethanol and acetaldehyde. The definition of the emission activity for carbon monoxide also differs from that of the other chemical species:

$$\gamma_{CO} = LAI \times \gamma_{TP} \times \gamma_{LA} \times \gamma_{HW} \times \gamma_{AQ} \times \gamma_{HT} \times \gamma_{LT} \times \gamma_{SM}$$

We now describe the equations defining the two activity parameters of interest to this work in turn. The canopy average temperature and radiation response parameter, γ_{TP} , is calculated via:

275

$$\gamma_{TP} = \sum_{j=1}^{j=5} (\gamma_{TP_{LDF}}^j \times LDF + \gamma_{TP_{LIF}}^j \times (1 - LDF)) \times W^j$$

Where W^j are the predefined layer-specific weights distributing the potential emission across the different canopy layers and are defined as follows: 0.119, 0.239, 0.284, 0.239, and 0.119. These weights implicitly assume the vertical structure of the canopy follows this prescribed definition. The light-dependent fraction temperature and radiation response g_{TP} is calculated for each layer, j , within the canopy via:

280

$$\gamma_{TP_{LDF}}^j = (\gamma_{CD}^j \times \gamma_{TLD_{sun}}^j \times \gamma_{P_{sun}}^j \times f_{sun}^j) + (\gamma_{TLD_{shade}}^j \times (\gamma_{P_{shade}}^j \times (1 - f_{sun}^j)))$$

Here, γ_P is the light response parameter, which is calculated independently for sunlit and shaded leaf areas depending on the photon flux within the canopy (calculated by CANMET – see Sect. 2.2.1) for both sunlit and shaded areas. γ_{TLD} is the light dependent temperature response parameter, which is also calculated independently for sunlit and shaded leaf areas. f_{sun}^j is the fraction of sunlit area at each canopy layer height and g_{CD} is the canopy depth parameter.

285 The light-independent fraction temperature and radiation response parameter for each canopy layer is defined as:

$$\gamma_{TP_{LIF}}^j = (\gamma_{TLI_{sun}}^j \times f_{sun}^j) + (\gamma_{TLI_{shade}}^j (1 - f_{sun}^j))$$

γ_{TLI} is the light independent temperature response activity parameter, which is calculated separately for sunlit and shaded areas to take into account the differences in leaf surface temperature in both regions of the canopy at a particular layer height.



290 The soil moisture activity factor, g_{SM} , is calculated differently according to whether the soil moisture, q , satisfies one of three different conditions:

$$\begin{aligned} \gamma_{SM} &= 0 & q < q_{wilt} \\ \gamma_{SM} &= q - q_{wilt} / 0.04 & q_{wilt} \leq q \leq q_{wilt} + 0.04 \\ \gamma_{SM} &= 1 & q_{wilt} < q \end{aligned}$$

295 Where q_{wilt} is the wilting point taken from the SURFEX physio-geographic maps, which is volumetric soil moisture at which plants are defined to wilt in SURFEX for a particular location.

The γ_{HW} , γ_{AQ} , γ_{HT} , and γ_{LT} activity factors are calculated with an approach that is similar to the above equation for the soil moisture activity factor. For the remaining activity parameters we refer readers to Guenther et al. (2012), which describe the details of how γ_{LA} and γ_{CO2} are calculated.

300 2.2.3 Emission Factors

The MEGAN EFP is built on the Python programming language with the SQLite database system as an opensource program that generates the emission factor and light dependence factors (EF/LDFs) required to drive MEGAN3.0. The program first generates EF/LDFs for individual plant types and then integrates them with plant type distribution data to calculate landscape-average EF/LDFs for a modelling domain.

305 To generate EF/LDFs for individual plant types, emission factor measurements compiled in the MEGAN EFP database are assigned a number from 0 to 4, called the J-rating, to indicate the quality of the data. A J-rating of 0 indicates the lowest quality including qualitative measurements and measurements conducted with methods that have high uncertainties and potentially strong bias. A J-rating of 4 is the highest quality data indicating that the data were obtained using methods that meet the recommendations of the BVOC emission measurement community (e.g., Niinemets et al., 2011). The MEGAN EFP
 310 allows users to choose to use all data or just measurements higher than a specified minimum J-value. The MEGAN EFP database also contains specific leaf area data (SLA) to convert EF measurements reported in terms of emissions per unit leaf mass to emissions per unit leaf area which is used by MEGAN.

Landscape average EF/LDFs are calculated with the MEGAN EFP by synthesizing leaf level plant trait data, including BVOC emission factors, specific leaf area and emission light dependence factor, with landcover data (described in more
 315 detail shortly), including ecotype and growth-form fractions for each location in a modelling domain. Additional information in the MEGAN EFP database includes descriptions of biogenic compounds, emission classes, publication references, vegetation types, and canopy vertical distribution characteristics. These data allow users to identify the emissions data that were used to drive the MEGAN3.0 model emission inputs.

We now describe the landcover data used by MEGAN EFP in more detail. The MEGAN3.0 emission factor distributions are
 320 based on four landcover input types: growth form fractions, ecotypes, plant type composition, and plant type emission factors. The growth form fractions, and ecotype inputs are $\sim 1 \text{ km}^2$ (30 second latitude \times 30 second longitude) resolution



global maps. The growth forms include trees, which are further divided into broadleaf vs needleleaf trees and tropical and extratropical trees, shrubs, grass and other herbaceous plants and crops. The total vegetation cover was based on a twelve-year climatology (Broxton et al., 2014) and the relative fractions from global datasets of tree cover (Hansen et al., 2003), shrub and grass cover (Tuanmu and Jetz, 2014) and crop cover (Latham et al., 2014) that were merged with regional data including the NLCD 30-meter landcover data for the U.S. (Homer et al., 2015). The ecotypes include ~900 global ecotypes and ~1200 regionally specific ecotypes for the U.S. and Australia (Guenther et al., 2012) and ~400 locally specific ecotypes representing specific U.S. cities and urban neighborhoods.

The plant type composition data is specified for each growth form and ecotype. For example, the plant type composition for a pine-oak forest ecotype includes the plant type composition for each growth form including broadleaf trees (dominated by oaks), needleleaf trees (dominated by pines), grass (temperate woodland grass), shrub (temperate woodland grass) and crop (generic crop). The plant types range from very specific, such as a plant species or even subspecies, to general types such as broadleaf tropical rainforest tree or arctic grass. The final database contains emission factors for each of the 20 MEGAN3.0 emission categories for each plant type. The four input data types were used to estimate weighted average emission factors at each location. The emission factors were based on data compiled for MEGAN2.1 (Guenther et al., 2012) with updates for vegetation in parts of the U.S. and Australia based on locally specific data.

2.3 Land surface model data

We use ISBA (Interactions Between Soil, Biosphere, Atmosphere) within the SURFEX (SURface EXternalisée) modelling platform (Masson et al., 2013) in this work to provide some of the key land surface variables (i.e., 2m-specific humidity, 2m-temperature, root zone soil temperature, root zone soil moisture, LAI) used in MEGAN. SURFEX simulates heat, moisture, and gas fluxes at the atmosphere-surface boundary and is designed to be coupled with meteorological models and atmospheric forcing both online and offline, respectively. SURFEX has been used successfully in a wide range of applications, e.g., river discharge prediction (Fairbairn et al., 2017), drought monitoring (Albergel et al., 2019), and urban climate studies (Schoetter et al., 2020). Regarding the soil-plant system, ISBA can dynamically simulate changes in Leaf Area Index (LAI) according to how meteorology impacts on the growing season. It has been shown to be skilful at estimating phenological changes and soil moisture on seasonal timescales when forced with state-of-the-art meteorological atmospheric forcing (Albergel et al., 2019; Szczypka et al., 2014).

SURFEX simulates four broad land use classes, nature, town, fresh water (lakes, rivers, and lagoons), and sea (see schematic embedded within Figure 1). The nature type is represented within SURFEX by 12 sub-classes of land surface class that represent different types of biomes and agricultural land use. These different land surface types and sub-classes are defined by ECOCLIMAP-II (Faroux et al., 2013). These 12 sub-classes are listed in Table 1. SURFEX was run using 12 sub-classes (termed patches) of the nature land type because some of the model options (detailed below in Table 2) that are required to



run simulations of dynamic vegetation can only be run with these 12 patch types. Table 2 summarises all of the relevant model options used.

SURFEX was configured to run on the same $0.1^\circ \times 0.1^\circ$ spatial grid we use for MEGAN3.0, which is the same spatial domain as used by the CAMS regional air quality models and the ECMWF HRES meteorology. The meteorological parameters required by SURFEX were retrieved from ECMWF onto this spatial grid. While this increased high spatial resolution has been already used in applications for specific isolated domains within Europe (Albergel et al., 2019), this is the first time SURFEX LDAS-MONDE will be run for the whole of Europe at this high spatial resolution.

SURFEX also includes a capability to perform data assimilation of satellite observations of land surface variables (soil moisture and LAI) using the Simplified Extended Kalman Filter (see e.g., Albergel et al., 2017). In this configuration SURFEX is termed the SURFEX LDAS-Monde (Land Data Assimilation System-World/Global), which can be applied with relative ease to study any area of the world. Assimilation of satellite observations of LAI and soil moisture in SURFEX LDAS-Monde has been shown to improve estimates of the soil moisture content and of phenological changes (Albergel et al., 2017) with LAI having a more significant and beneficial effect for the estimation of root zone soil moisture. For the purposes of the production of the BVOC emission datasets presented in this article we produce two different sets of land surface data for use in MEGAN. The first is just SURFEX run in a free-running mode with no data assimilation, which we term the open-loop dataset. The second is a SURFEX simulation run created by running SURFEX with data assimilation of satellite LAI, which we term the analysis. The analysis is run with the aim to improve estimation of LAI and the root zone soil variables that are relevant for vegetation growth and function that are used in MEGAN, i.e., root zone soil moisture and root zone soil temperature.

To produce the assimilation-based analysis we assimilate LAI satellite data products from the Copernicus Land Monitoring Service (CLMS, <https://land.copernicus.eu/global/products/lc>, last access: 17 July 2025). Specifically, this includes assimilation of the PROBA-V LAI (Verger et al., 2014) product for years 2018 and 2019, using the GEOV1 product of CGLS. Since PROBA-V was decommissioned in 2020, we used the THEIA AVHRR-derived LAI (<https://www.theia-land.fr/product/serie-de-variables-vegetales-avhrr-fr/>, last access: 17 July 2025) and performed a seasonal cumulative density function (CDF) matching of the latter from 1999 to 2019 (21 years) in order to use the CDF-matched THEIA LAI as a proxy of GEOV1 for the whole 2020. This AVHRR LAI product is derived from the Land Long Term Data Record (LTDR) AVHRR data (Vermotte 2021) (https://landweb.modaps.eosdis.nasa.gov/data/userguide/LTDR_Ver5_Products_UserGuide_v1.0.pdf, last access: 17 July 2025) using another version of the GEOV2-AVHRR algorithm described in Pacholczyk and Verger (2020) (https://www.theia-land.fr/wp-content/uploads/2022/03/THEIA-MU-44-0369-CNES-GEOV2-AVHRR-Product-User-Manual-V2_AV.pdf, last access: 17 July 2025).

We did not use the CLMS LAI products for LAI from Sentinel-2 or from Sentinel-3 because we did not need the higher resolution of Sentinel-2 in this application, and we found a discontinuity between the PROBA-V and the Sentinel-3 data version that was available at the time of study.



Table 1. List of SURFEX nature tile patch numbers and corresponding ISBA vegetation type.

Patch number	ISBA vegetation type
1	Bare soil
2	Bare rock
3	Permanent snow
4	Deciduous broadleaf
5	Needle leaf
6	Evergreen broadleaf
7	C3 crops
8	C4 crops
9	Irrigated crops
10	Temperate grassland
11	Tropical grassland
12	Wetlands, parks and gardens

Table 2. Model options used in SURFEX

Model option	Selected model setting
ISBA photosynthetic scheme	Nitrogen dilution scheme – NIT
Soil moisture scheme	Diffusion scheme
Number of soil layers	14
Town model	No town model – All urban areas represented as solid rock
Lake model	Water flux model following Charnock formula
Sea model	Sea flux model following Charnock formula

2.4 Meteorology

Both the SURFEX and MEGAN3.0 model algorithms rely on the use of meteorological data from the European Centre for Medium-Range Weather Forecasts (ECMWF) HRES operational forecast (Owens, R G, Hewson, 2018) for describing the meteorological conditions. The data was extracted via the MARS service (ECMWF's Meteorological Archival and Retrieval System) on a $0.1^{\circ} \times 0.1^{\circ}$ grid with hourly resolution for the surface levels. The forecast at 12:00 was used and the following 36 hours have been retrieved and then post-processed to create a single contiguous daily time series running from the 13th hour of the forecast (00:00 UTC) to the 36th hour (23:00 UTC). We then combined each individual daily time series together



to make a continuous running time series spanning the full dataset time period, i.e., 2018-2022. Table 3 shows which meteorological variables were extracted for use in which of the two models (SURFEX and MEGAN3.0). In the case of MEGAN3.0 we used two of the variables (2m air temperature and 2m specific humidity) calculated by SURFEX. There is a specific advantage to using these variables from SURFEX as opposed to the ECMWF HRES data directly because SURFEX can account for surface effects like surface fluxes of latent and specific heat and surface driven turbulence arising from different vegetation canopy heights and densities and different transpiration rates. In the case of all other meteorological variables the models used ECMWF HRES data.

Table 3. Summary of meteorological variables used as inputs by the SURFEX and MEGAN3.0 algorithms.

ECMWF HRES Meteorology	Calculated within SURFEX	Meteorological Variables used in SURFEX	Meteorological Variables used in MEGAN3.0
Surface Pressure		✓	✓
2m Air Temperature		✓	-
	2m Air Temperature	-	✓
Wind Speed		✓	✓
Air Specific Humidity		✓	-
	2m Specific Humidity		✓
Liquid Precipitation		✓	-
Solid Precipitation		✓	-
Incoming Shortwave Radiation		✓	-
Incoming Longwave Radiation		✓	-
PAR			✓

2.5 Isoprene Observations

We perform an evaluation of the temporal variability of the BVOC emissions using in-situ observations of isoprene collected during the period of our datasets (2018-2022). We obtained the isoprene observations from the EBAS database (Europe-wide observations excluding UK; [EBAS home – ebas homepage](#), last access: 17 July 2025) and from UK-air (specific to UK observations; [Data Archive - Defra, UK](#), last access: 17 July 2025). The EBAS database contains a wide variety of isoprene observations collected using different monitoring techniques. We select only observations made using the online gas chromatograph mass spectrometry method and we made a further selection to exclude observations made in dense urban



415 settings, e.g., Marylebone Road, London, to avoid sites with a strong influence from anthropogenic isoprene sources (Khan et al., 2018).

3 Results and Discussion

We describe different aspects of the SURFEX-MEGAN3.0 datasets within the following three sub-sections. First, in section 3.1, we perform an evaluation of the underlying LAI data produced in the course of running the SURFEX-MEGAN3.0 production chain. Then, in section 3.2, we describe the spatiotemporal characteristics of the datasets and we describe the impact of the LAI data assimilation on the results. Lastly, we evaluate the SURFEX-MEGAN3.0 emission inventory for isoprene against other published emission inventories in section 3.3. Throughout these discussions we focus on isoprene emissions. Isoprene is one of the most important BVOCs because its emissions are larger than other emitted BVOC species and it is one of the most reactive BVOCs and it therefore has an important influence on atmospheric composition.


425 3.1 Performance of SURFEX land surface model

One important conceptual basis for the creation of the BVOC emission datasets presented in this paper was that we could improve BVOC emission modelling by focusing on improvements of the input data used by the MEGAN emission model. Indeed, we hypothesize that BVOC emission modelling can be advanced by using the SURFEX land surface model to provide improved estimates of LAI and soil moisture based on dynamic, realistic vegetation phenology. Furthermore, we posit that the inclusion of the methodological step to assimilate satellite observations of LAI further improves the model representation of LAI used for BVOC estimation. Thus, we now attempt to evaluate the performance of the SURFEX land surface model for these key input variables used by MEGAN. We perform this evaluation using three approaches: i) use of TROPOMI satellite observations of solar induced fluorescence (SIF) to evaluate LAI, ii) in-situ observations of soil moisture, iii) a review and discussion of literature covering existing case studies evaluating the performance of the SURFEX land surface model, and iv) a brief review of the data quality of the LAI satellite observations used. In addition, we refer readers to Sect. 3.2.2 that covers a discussion of the impacts of the assimilation of satellite observations of LAI as this gives some further insight into the relative performance of the OL and analysis model runs.

To assess the correlation between the SIF observations and both the LAI open-loop and LAI analysis, temporal correlation coefficients with SIF and LAI are calculated over the entire domain from 1 May 2018 to 31 December 2020. The results, shown in Figure 3, show robust correlations over much of the domain with correlation coefficients (R) greater than 0.7 for both the LAI open-loop and LAI analysis. Semi-arid regions such as parts of the Iberian Peninsula and the Middle-East, together with Nordic regions at high latitudes, show weaker correlation values. These regions present sparse vegetation and the SIF signal is weak, which likely degrades the correlations by making the SIF observations more subject to random and systematic errors in the retrieval. Figure 3 also shows the difference in correlation coefficients between the analysis and the open-loop LAI. When comparing the analysis with the OL (Figure 3b), improvements can be seen over almost the whole



study area. In particular, it can be seen that the assimilation improves the correlation between LAI and SIF over Germany and the Czech Republic. It should be noted, however, that the OL LAI does still correlate with SIF in a reasonable way, but the areas where it shows weaker correlation are more widespread.



450 **Figure 3. Maps showing (a) the Pearson correlation in time between the LAI analysis and the observed TROPOMI SIF (re-gridded to the CAMS spatial gridding) for each $0.1^\circ \times 0.1^\circ$ grid cell and (b) the difference in correlation between the LAI analysis and SIF and the LAI open-loop and SIF.**

We now analyse the soil moisture represented by the SURFEX model. The SURFEX model soil layer corresponding to the SMOSMANIA in situ measurements at 0.3 m depth is layer 5 (0.2-0.4 m soil layer). This is also the layer that we use to
 455 create the root-zone soil moisture (RSZM) dataset used by MEGAN. The OL and analysis root zone soil moisture (RSZM) simulations for layer 5 are compared with observations from the Saint-Felix de Lauragais (SFL) station of the SMOSMANIA network for the study period (1 May 2018 to 31 December 2020) located in the south-west of France. The open-loop, analysis and in situ time series of RSZM are shown in Figure 4. The temporal patterns of the open-loop and analysis results clearly correlate with the observed seasonal variability of the RSZM, with similar times of rewetting events.
 460 The correlation scores are 0.92 and 0.93 for the OL and analysis (all seasons), respectively, and 0.74 and 0.77 (summertime only) for both model versions. The observed RSZM values range between 0.16 and 0.37 $\text{m}^3 \text{m}^{-3}$, while the corresponding variations in the open-loop and analysis simulations show values between 0.25 and 0.46 $\text{m}^3 \text{m}^{-3}$. The larger simulated RSZM values can be explained by differences between the soil properties used in the model, such as porosity, and the local soil properties around the soil moisture probe. Differences between the open-loop and analysis simulations occur during certain
 465 periods of the year, such as the autumn of 2019. The vegetation model is sensitive to soil moisture deficit through a soil wetness index corresponding to rescaled volumetric soil moisture between field capacity and wilting point. In this way, a large absolute bias for volumetric soil moisture has little effect on the vegetation response to drought.




Figure 4. Time series of observed in-situ RSZM at the Saint-Felix de Lauragais SMOSMANIA monitoring station (green), the RSZM from the SURFEX open-loop simulation (yellow), and the RSZM from the SURFEX analysis simulation (red).

Unlike other land surface models, phenology in ISBA is entirely driven by photosynthesis and LAI responds to environmental factors such as drought, temperature, or solar radiation through photosynthesis (Calvet et al. 1998, Gibelin et al. 2006). ISBA is not calibrated and the vegetation parameters that drive photosynthesis and plant growth are derived from the literature (Delire et al. 2020). Despite this unique way of representing phenology, ISBA can achieve good skill compared to other LSMs (Peano et al. 2021, Friedlingstein et al. 2022).

The CLMS LAI data assimilated by SURFEX has been evaluated extensively (Brown et al., 2020; Verger et al., 2023) demonstrating its consistency with MODIS products, temporal consistency, and highest accuracy when compared to reference data. Thus, the LAI data ingested into the SURFEX assimilation algorithm has an established track record of being of high quality.

In summary, the simulated LAI and RZSM show robust correlations with independent data, and prior work establishes the skill of SURFEX and representing vegetation phenology relative to other LSMs. For LAI, the correlation with SIF is better than 0.7 over much of the domain. However, areas covered by sparse vegetation, semi-arid regions such as parts of Spain, together with Nordic regions at high latitudes, show weaker correlation values.



3.2 Characteristics of SURFEX-MEGAN BVOC dataset

3.2.1 Spatiotemporal distribution of SURFEX-MEGAN BVOC emissions over Europe

Annual Mean Isoprene Emissions Over the CAMS European Domain for 2018-2022 - Open Loop




Figure 5. Maps of annual mean isoprene emissions (units of $\text{kg.m}^{-2}.\text{s}^{-1}$) calculated using the SURFEX-MEGAN3.0 algorithms over the period 2018-2022 using the open-loop configuration of SURFEX. The panels show each year over this time period moving sequentially from 2018 to 2022 from left to right. The colour bars indicate increasing mean isoprene emissions as the colours transition from pale yellow to dark green. Note that the colour bar has a quasi-logarithmic scale. The annual total mass of isoprene emissions (units of Tg) are shown above each plot.

We first describe the temporal and spatial characteristics of the datasets on an annual basis. The annual mean isoprene emissions are displayed as maps over the CAMS European domain in Figure 5 for the period 2018-2022. The spatial distribution of the isoprene emissions on an annual scale is dependent on the general distribution of isoprene emitting vegetation as represented in the distribution of the isoprene emission factors. Thus, we see the regions with the highest year-round emissions over areas with dense forest, and the regions with the lowest emissions (near-zero) over desert regions, high mountains, and permanent ice and snow.



Standard Deviation of Isoprene Emissions Over the CAMS European Domain for 2018-2022 - Open Loop




Figure 6. The standard deviation of isoprene emissions over the CAMS European domain for the 2018-2022 period based on the SURFEX model open-loop simulations. Note that the standard deviation is expressed in the units of the emissions ($\text{kg.m}^{-2}\text{s}^{-1}$).

The total annual mass of isoprene emissions is shown in Tg above each panel in Figure 5. The average annual emitted mass of isoprene for the open-loop emissions over the time period 2018-2022 was calculated to be 7.20 Tg yr^{-1} with a standard deviation of $\pm 0.28 \text{ Tg yr}^{-1}$. The emissions were estimated to be at their highest in 2018 and their lowest in 2020 and the difference in annual emissions between these two extremes was 0.82 Tg yr^{-1} , which represents 11.4% of the average annual emission of isoprene in this time period. While the standard deviation on the total annual emission over this period is 0.28 Tg yr^{-1} , this temporal variability is not evenly distributed spatially as can be seen within Figure 6, which shows the standard deviation in isoprene emissions calculated on an inter-annual basis. The areas of highest annual variability over the 2018-2022 period coincide with the co-location of forests, emitting vegetation species, and meteorologically induced variability resulting from temperature, radiation, and soil moisture forcing. Indeed, the effect of these last variables can be seen in more detail within Figure 7 which shows the emission factors over the CAMS domain and the standard deviation of the inter-annual variability for the gamma parameters for soil moisture, LAI, and radiation-temperature. Some of the areas with the highest year-to-year variability in emissions correspond to regions over where the emission factors are highest. Since the



emission factors are invariant in time, the standard deviation of the emissions will naturally be larger over regions with a higher emission factor as this acts to inflate the variability applied by the parameters used to calculate the activity. Similarly, LAI can vary quite significantly in some regions and where this is co-located with higher emission factors, this can lead to LAI having an important role in modulating emissions, e.g., over France, the Carpathian Mountains, Dinaric Alps, and the Caucasus.




Figure 7. Maps of the MEGAN3.0 EFP isoprene emission factors in units of nano moles $\text{m}^{-2} \text{s}^{-1}$ (top left), the LAI gamma in units of $\text{m}^{-2} \cdot \text{m}^{-2}$ (top right), the soil moisture gamma (unitless) (bottom left), and the temperature-radiation gamma in the bottom right.

In addition to analysing the standard deviation of the gamma parameters and emissions, we also analyse the correlation of the emissions to the different gamma parameters over different temporal timescales (yearly, monthly, daily, and hourly). We calculate the correlation in time at each grid cell of the soil moisture gamma, the temperature-radiation gamma, and LAI to the isoprene emissions calculated by MEGAN3.0. Thus, for each spatial grid cell and each pair of variables we derived three



different R^2 values for each of the four selected timescales we averaged over. In order to support the interpretation of the inter-annual variability in isoprene represented in Figure 5 and Figure 6 we first analyse the correlation between isoprene emissions and the most influential gamma parameters over a yearly timescale and present these results in Figure 8. Within Figure 8, we represent the different R^2 values for each grid cell as a series of statistical distributions using violin plots. These plots show that the correlation between the isoprene emissions and the different gamma parameters is largest, both in terms of being positive and magnitude (median of $\sim 0.75 R^2$), for LAI. This suggests that LAI is the dominant driver of isoprene emission inter-annual variability within our modelling framework. It also further highlights the advantage of taking an Earth system approach by coupling dynamic vegetation modelling to BVOC emission modelling. The two remaining gamma parameters, i.e., the soil moisture and radiation-temperature gammas, have much lower median correlation to the emissions on a yearly timescale. Indeed, at times both show significant negative correlations. Even though the soil moisture gamma shows slightly higher overall correlation to the emissions, this difference probably has little physical meaning. Despite both the soil moisture and radiation-temperature gammas showing low median correlations and some negative correlations, there are still grid cells that show large positive correlation for both parameters indicating that these parameters play a role within some specific regions.





Figure 8. Violin plots showing the statistical distribution of R^2 correlation values between different activity parameters and the isoprene emissions calculated on a per grid cell basis averaged on an annual basis for each year from 2018-2022. The radiation-temperature gamma (GAMTP) is shown in dark green, the soil moisture gamma (GAMSM) in light blue, and LAI in turquoise. The pink horizontal bar in each plot represents the median R^2 value.



Monthly Mean Isoprene Emissions Over the CAMS European Domain averaged over 2018-2022 - Open Loop



545 **Figure 9.** Maps of monthly mean isoprene emissions (units of $\text{kg.m}^{-2}.\text{s}^{-1}$) calculated using the SURFEX-MEGAN3.0 algorithms and averaged over 2018-2022 using the open-loop configuration of SURFEX. The colour bars indicate increasing mean isoprene emissions as the colours transition from pale yellow to dark green. Note that the colour bar has a quasi-logarithmic scale. The monthly total mass of isoprene emissions (units of Tg) are shown above each plot.




Figure 10. Violin plots showing the statistical distribution of R^2 correlation values between different activity parameters and the isoprene emissions calculated on a per grid cell basis over different averaging periods (monthly in plot (a), daily in plot (b), and hourly in plot (c)) for each year from 2018–2022. The radiation-temperature gamma (GAMTP) is shown in dark green, the soil moisture gamma (GAMSM) in light blue, and LAI in turquoise. The pink horizontal bar in each plot represents the median R^2 value.

We now examine the monthly and seasonal variability in the isoprene emissions. For this purpose, we present the monthly mean emissions from the OL-based dataset averaged over the 2018–2022 time period. This way we avoid presenting single



years that can be affected by extremes in meteorology that lead to large increases or changes in isoprene emissions. For reference, however, the monthly means for all five years are presented in the supplement.

560 The monthly mean isoprene emissions over the CAMS European domain are shown in Figure 9. First and foremost, we see the combined effects of the growing season, leading to changes in leaf area, and the annual cycle of sunlight and temperature during the year in this plot, with emissions at a minimum in the winter months and maximum in the summer months. Not only are the effects of the growing season visible from month to month, but we also see the progression of the growing season geographically. More southern regions have a growing season that starts earlier in the year compared to northern
 565 regions, but southern regions also show signs of reduced vegetation activity during summer as temperatures increase beyond optimal growing conditions. Similarly, western regions with milder maritime climates have increased emissions in the late winter and early spring compared to areas of eastern Europe with harsher continental-type winter conditions. Furthermore, the emissions peak in regions with higher densities of forest and thus emitting species and that coincide with regions with high levels of LAI (see Figure 7).

570 There is a clear pattern presented in Figure 10 showing how the different MEGAN activity parameters drive variability in the emissions over different timescales. According to Figure 10(a), the monthly variability in emissions is driven by variability in the radiation-temperature gamma and LAI, with radiation-temperature being the dominant of the two parameters influencing isoprene emissions. At the shorter timescales, i.e., daily and hourly shown in Figure 10(b) and Figure 10(c), respectively, the radiation-temperature gamma increasingly dominates the variability in isoprene emissions compared to the
 575 other two parameters. Indeed, the correlation of the radiation-temperature gamma to emissions actually increases as the timescale shortens while the correlation of LAI to the emissions decreases markedly from monthly to hourly timescales. Figure 10 also shows that there is a remarkable consistency of the correlations of each parameter across each year that the dataset covers, i.e., 2018-2022. Indeed, there are only small variations in the median R^2 value across each year for each parameter and for each timescale. Furthermore, the shape of the distribution of the R^2 values for the radiation-temperature
 580 gamma is very consistent over all timescales and years. The R^2 values for LAI and the soil moisture gamma do differ, however, in this regard with each pairing of year and timescale displaying markedly different distribution from each other. The distribution of the R^2 of the soil moisture gamma varies widely, which is indicative of the varying influence this parameter has on controlling isoprene emissions.

We now examine the variability in the simulated isoprene emissions during the summer of 2019. We select data from 2019
 585 to study and evaluate because 2019 presented some extreme and unseasonably hot conditions during the early summer that led to very large isoprene emissions. The results for the monthly mean isoprene emissions for the summer of 2019 are presented in Figure 11 covering the June, July, and August period along with the monthly mean LAI from the SURFEX open-loop simulation. Later in the summer, hot, dry weather caused the dynamic vegetation scheme within SURFEX to simulate strong reductions in leaf cover that led to reduced isoprene emissions. Indeed, isoprene emissions in 2019 peaked in
 590 June according to the SURFEX-MEGAN3.0 dataset, which makes 2019 unique among the five-year dataset since normally emissions peak in July within these data. Thus, 2019 presents an example of the impact extreme weather can have on



isoprene emissions via its effect on vegetation represented by the dynamic vegetation scheme. Visible within the maps, there are sharp declines in isoprene emissions within specific areas of Europe over this period, which can also be seen in the total emitted isoprene mass. To further investigate this link, we plot the temporal correlations (on a daily-averaged timescale) of the isoprene emissions to LAI over this time period in Figure 12. Figure 12 (a) shows large areas of Europe with high correlation between LAI and the isoprene emissions, and it is possible to see that some of the areas with high correlation correspond to areas with large declines in LAI from June to August in Figure 11. Furthermore, Figure 12 (b) shows the correlation between LAI and isoprene emissions only in areas where the LAI declined from June to August, and it is possible to see that the application of this mask to the dataset removed most of the areas with negative correlation between LAI and isoprene emissions. This evidence strongly suggests that the decline in LAI, which is driven by drought and heat stress on plants, is an important causative agent behind the reductions in isoprene emissions in the data during the summer of 2019.





Figure 11. Maps of monthly mean isoprene emissions (units of $\text{kg.m}^{-2}.\text{s}^{-1}$) calculated using the SURFEX-MEGAN3.0 algorithms using the LAI analysis configuration of SURFEX (top-row) and LAI (units $\text{m}^{-2}.\text{m}^{-2}$) calculated by SURFEX in the open-loop configuration over the period June to August 2019. The panels sequentially show the monthly means for June, July, and August. The top row colour bars indicate increasing mean isoprene emissions as the colours transition from pale yellow to dark green. The bottom-row colour bars show increasing mean LAI as the colours transition from dark blue to yellow. The annual total mass of isoprene emissions (units of Tg) are shown above each plot in the top-row and the differences in annual total mass of isoprene emissions are shown above each plot in the top-row.



615 **Figure 12. Maps of the correlation between LAI and the daily mean isoprene emissions over the CAMS European domain for the June to August 2019 time period. The left-hand panel shows the correlations for all grid cells within Europe while the right-hand panel shows the correlations only in locations where the LAI decreased between June 1st and August 31st. The colour bar shows green areas where the correlation was positive while the red areas shows where the correlation was negative.**

This analysis proves that modelling the vegetation dynamically for MEGAN3.0 has a direct impact on the isoprene emissions it calculates during drought and heatwave conditions. This approach moves beyond using either a climatology or more simplistic phenological model to calculate LAI. Furthermore, since we have established the skill of the LAI simulation within prior discussions in Sect. 3.1, we can have some confidence that the representation of these changes in LAI and the consequent declines in isoprene are realistic.

3.2.2 Impact of LAI data assimilation

625 We next evaluate the impact that the LAI data assimilation step has on the estimation of isoprene emissions within the SURFEX-MEGAN3.0 system. The LAI assimilation analysis data from SURFEX is based on the data assimilation of LAI satellite observations. Suitable satellite observations were only available during the 2018-2020 period, so the evaluation of the use of the LAI analysis dataset only covers this 3-year period. Prior to analysing the effect of the LAI data assimilation step on isoprene emissions, we first evaluate the effect on LAI itself of performing the data assimilation step.


630 To evaluate the effect of the assimilation we plot in Figure 13 the annual mean LAI analysis minus OL LAI difference for the three years (2018-2020) when we have coverage of both datasets. We can see that the LAI analysis is consistently higher than the OL over France, the UK, and Ireland and consistently lower over northern Italy, areas of the Iberian Peninsula, and eastern Europe. The absolute differences in LAI between the two datasets peak at around $0.4 \text{ m}^2.\text{m}^{-2}$ over these highlighted regions and the absolute differences generally stay below $0.2 \text{ m}^2.\text{m}^{-2}$. The difference between the analysis and OL calculated relative to the LAI analysis similarly shows peaks of ~40% (both positive and negative) over specific regions (e.g., regions of Spain, Italy, north Africa, and the Middle East) across the three years when these datasets are compared. Of the three years, 2019 is the year with the lowest relative differences and 2020 shows the largest relative differences.

We briefly discuss a further implication of this evaluation of the LAI data assimilation step. We can consider that the LAI assimilation step provides an indirect evaluation of the free-running model LAI from the OL simulation. This is because the




640 LAI data assimilation step corrects the OL LAI simulation in regions where the OL model minus observations errors are large, which in turn leads to larger assimilation increments (either positive or negative) in these locations. Conversely, in areas where the increments are low, we can determine that the OL LAI estimates are already in closer agreement with the satellite observations of LAI.

Annual Mean Difference in LAI Over the CAMS European Domain for 2018-2020 - LAI Analysis minus OL



Annual Mean Relative Difference in LAI Over the CAMS European Domain for 2018-2020 - LAI Analysis minus OL



645 **Figure 13. Maps of the absolute and relative difference in annual mean LAI (units of m².m⁻²) between the LAI analysis minus the open-loop configurations over the period 2018-2020. The panels show each year over this time period moving sequentially from 2018 to 2022 from left to right. The colour bar indicates the difference in annual mean LAI between the LAI-analysis and open-loop configurations with red colours indicating larger annual mean open-loop LAI and green indicating larger LAI-analysis based LAI. The difference in annual mean LAI is shown above each plot.**


650 Next, we analyse the impact that the LAI data assimilation has on the isoprene emissions both on an annual and monthly average basis. Our aim here is to directly evaluate the differences between the two emissions datasets (OL-based emissions and LAI-analysis based emissions) presented in this study. We first look at the annual averages and the results of this comparison are presented in Figure 14. We have already presented the OL-based emissions in Sect. 3.2.1, so we only show the annual averages of the analysis-based emission data for 2018-2020 in Figure 14 along with the analysis minus OL differences and the percentage difference in isoprene emissions between the two datasets. The annual average isoprene emission total calculated from the analysis-based emissions (calculated over 3 years 2018-2020) is 7.19 Tg yr⁻¹ with a standard deviation of 0.40 Tg yr⁻¹. Note that the average annual emitted isoprene from the OL-based emissions over this




660 same time period were 7.14 Tg yr^{-1} with a standard deviation of 0.44 Tg yr^{-1} . We identify four main findings from the analysis of these data. First, the impact of the data assimilation upon the overall annual emission totals is relatively small as we can see from the differences noted above the plots in the second row of Figure 14. Second, while the impact on the sum of the emissions is minimal at continental scale (noting the annual average difference in emission totals), we see much stronger variations in the emissions within specific regions, e.g., the Iberian Peninsula in 2018, northern Italy, north Africa, and Croatia in 2020. Third, when we look at the percentage differences between the OL- and analysis-based datasets, we can see regions with large relative differences. Finally, the CANMET routine in MEGAN3.0 calculates the penetration of solar radiation through the vertical column of the canopy, and it can be considered that as LAI increases above $3 \text{ m}^2\text{.m}^{-2}$ (see 665 Figure 11 for reference), solar radiation becomes significantly attenuated within the canopy. As such, increases in LAI beyond this threshold do not lead to large increases in isoprene emissions. This means that regions with lower levels of LAI (i.e., $< 3 \text{ m}^2\text{.m}^{-2}$) are more sensitive to changes in LAI that result from the assimilation step. The effects of this are visible, for example, areas of the Iberian Peninsula and north Africa during 2018 and in northern Italy from 2018-2020.



Annual Mean Isoprene Emissions Over the CAMS European Domain for 2018-2020 - LAI Analysis



Annual Mean Isoprene Emissions Over the CAMS European Domain for 2018-2020 - LAI Analysis minus Open Loop



Relative Difference in Annual Mean Isoprene Emissions Over the CAMS European Domain - LAI Analysis minus Open Loop




Figure 14. Maps of annual mean isoprene emissions (units of $\text{kg.m}^{-2}.\text{s}^{-1}$) calculated using the SURFEX-MEGAN3.0 LAI analysis configuration of SURFEX (top-row), the difference between the LAI analysis minus the open-loop configurations (middle-row), and the percentage difference between the LAI analysis minus open-loop over the period 2018-2020. The panels show each year over this time period moving sequentially from 2018 to 2020 from left to right. The top-row colour bars indicate increasing mean isoprene emissions as the colours transition from pale yellow to dark green. The middle-row colour bars indicate the difference in annual mean isoprene emissions between the LAI-analysis and open-loop configurations with red colours indicating larger annual mean open-loop emissions and green indicating larger LAI-analysis based emissions. The bottom-row colour bar indicates the percentage difference in mean annual isoprene emissions between the LAI-analysis and open-loop with green indicating larger LAI-analysis based emissions and red indicating larger open-loop based emissions. The annual total mass of isoprene emissions (units of Tg) are shown above each plot in the top-row and the differences in annual total mass of isoprene emissions are shown above each plot in the middle-row.



We present the absolute monthly mean differences in analysis- and OL-based isoprene emissions averaged over the 2018-2020 time period in Figure 15. When we analyse the impacts of the LAI assimilation on a monthly average basis, the absolute differences in isoprene emissions visible at regional scale resulting from the LAI data assimilation are larger than those seen in the annual averages (Figure 14). Note that we have plotted the monthly mean analysis-based isoprene emissions averaged over 2018-2020 and these are shown in the supplement. In addition, the monthly mean emissions for the assimilation analysis are also plotted for each year and shown in the supplement.

The magnitude of the absolute differences between the analysis- and OL-based emissions shown in Figure 15 follows the growing cycle such that the size of the difference in emissions peaks when the emissions are at their highest for May through August. There are a few notable differences in emissions between the OL- and analysis-based emissions. First, the analysis-based emissions are larger in central and eastern Europe in April and May. Second, the picture becomes more complex in June, while Russia and the areas of higher altitude woodland across Europe (Carpathian mountains, Jura, Dinaric Alps, and Black Forest) show increased emissions in the analysis, areas of France, Turkey, the Iberian Peninsula, and north Africa show lower emissions compared to the OL. Lastly, in July and August, the OL emissions are larger than the analysis-based emissions over most of Europe, with France and Germany showing the largest differences between the two datasets.



Monthly Mean Isoprene Emissions Over the CAMS European Domain averaged over 2018-2020 - LAI Analysis minus Open Loop




Figure 15. Maps of monthly mean (units of $\text{kg.m}^{-2}.\text{s}^{-1}$) difference in isoprene emissions between the LAI analysis minus the open-loop configurations averaged over 2018 to 2020. The panels show each month during 2019 moving sequentially from January to December first from left to right and then from top to bottom. The colour bars indicate the difference in monthly mean isoprene emissions between the LAI-analysis and open-loop configurations with red colours indicating larger monthly mean open-loop emissions and green indicating larger LAI-analysis based emissions. The monthly total mass difference of isoprene emissions (units of Tg) are shown above each plot.



705

3.3 Comparison of SURFEX-MEGAN BVOC emissions with other datasets

We now perform an evaluation of the two BVOC emission datasets presented in this study. We perform this evaluation in two different ways. First, we evaluate the isoprene emission datasets against in-situ observations of isoprene from monitoring sites across Europe. And the second, we evaluate the isoprene emissions with respect to other emission datasets.

710 3.3.1 Evaluation using in-situ observations

We now evaluate the isoprene emissions using in-situ observations of isoprene concentrations obtained using the online gas-chromatograph (GC) technique. Note that we have excluded observations from other observation techniques, e.g., adsorption tube and canister sampling combined with the offline GC technique due to known issues involving systematic biases with observations obtained using these methods (Plass-Dülmer et al., 2006). In addition, these other observation techniques offer lower time resolution and sometimes do not provide continuous sampling. The emissions are extracted for the geographical location of the in-situ observations for this comparison. Since we are not comparing like with like, we cannot perform a direct evaluation of the absolute values, and so we focus on performing an analysis of the temporal correlation between the emissions and ambient isoprene concentrations. To facilitate this comparison, we normalise the emissions and isoprene concentrations onto a common scale running from 0 to 1. This allows us to carry out an evaluation of the temporal variability of the emissions and to derive the temporal correlation in terms of R^2 between the simulated emissions and observations. We perform this analysis each year over the period 1st April though to 1st October. Although limiting the evaluation time period to the approximate duration of the growing season reduces the overall quality of the correlations leading to reduced R^2 scores, this mostly arises due to the removal of lower observed isoprene concentrations and emissions that occur during the wintertime. Including the lower values of both concentrations and emissions out of the growing season artificially boosts the scores. However, our objective here is to evaluate the higher temporal frequency of the day-to-day correlations. We also filter the data to remove all cases where the observations lay below the limit of detection (LOD). In addition, we remove data for particular years, for sites where there was either a lack of data, or there were apparent issues with the quality of the observations, i.e., large populations of data points near to the instrument LOD showing apparent systematic bias. The years we exclude are noted in relation to the compiled statistics shown in Table 4. Furthermore, we also excluded observations from very polluted urban sites in London, Paris, and Marseille due to concerns about contributions to ambient isoprene concentrations from anthropogenic sources (Khan et al., 2018).

We calculated the daily mean emissions from both emission datasets and daily mean isoprene concentrations at the location of each observation site. We then calculated the Pearson correlation coefficients between the emission-observation pairings for each year and site. We collate all the correlation coefficients for each site, and for each year of the dataset, and present them in Table 4.



Table 4. Correlation coefficients showing the correlation in time between daily mean isoprene emissions and daily mean isoprene concentrations calculated for a variety of monitoring sites across Europe. We calculate the correlation coefficients individually for each year for the OL (2018-2022 shown as O) and for the analysis (2018-2020 shown as A). The monitoring sites are distributed across France, the UK, Switzerland, Germany, and Finland. The stations Feyzin Stade, Vernaion, Eltham, Chilbolton, Auchencorth, Beromünster Hohenpeissenberg, and Pallas are represented by the following three-letter acronyms, respectively, FEY, VER, ELT, CHI, AUC, BER, HOH, and PAL.

	France				United Kingdom				Switzerland				Germany		Finland	
Year	FEY		VER		ELT		AUC		BER		ZÜR		HOH		PAL	
	O	A	O	A	O	A	O	A	O	A	O	A	O	A	O	A
2018	0.56	0.54	0.73	0.72	0.36	0.26	0.80	0.80	0.80		-		0.79	0.80	0.76	0.75
2019	0.74	0.73	0.81	0.80	0.85	0.81	++		0.83		-		0.81	0.81	0.70	0.70
2020	0.63	0.66	0.63	0.65	0.51	0.44	0.72	0.72	0.72		-		0.88	0.88	0.39	0.38
2021	0.72	-	0.88	-	0.87	-	++	-	0.78	-	-	-	0.81	-		-
2022	0.36	-	0.63	-	0.17	-	++	-	0.64	-	0.89	-	0.76	-		-

The correlations shown in Table 4 show that in general the isoprene emissions correlate with the observed isoprene concentrations at a moderate to strong level (22 cases of strong correlation $> 0.7 R^2$, 7 cases of moderate correlation $> 0.4 R^2$, and only 4 cases of weak correlation $< 0.4 R^2$). The weakest correlations only occur at FEY in 2022, at ELT for 2018 and 2022, and at PAL in 2021. The ELT site is an urban background site located inside Greater London, so it is possible it is partially affected by anthropogenic sources of isoprene. In this case of the ELT, the poor correlations could be due to its proximity to London and therefore might experience influence from anthropogenic sources of isoprene (Khan et al., 2018). Note that the R^2 scores indicate there is no consistent improvement in the analysis emissions relative to the open-loop emissions. There are instances (both specific years and locations) where the analysis has both higher and lower correlation than the open-loop but overall there is not a strong indication either way which has consistently better correlation. This is in spite of the fact that the LAI analysis shows higher correlation with the independent TROPOMI SIF observations than the open-loop (see Sect. 3.1).

Overall, this evaluation shows that the two isoprene emission datasets are able to capture the temporal variability of emissions on daily timescales. This is encouraging given that this comparison compares emissions to concentrations and there are several likely sources of error that would obfuscate such a comparison. First, there is the issue of the representativity of the $0.1^\circ \times 0.1^\circ$ grid scale to the observation sites, which is important because of isoprene's very short lifetime. Second, isoprene's ambient concentration can also be affected by the abundance of its sinks, the hydroxyl radical, the nitrate radical, and ozone, which are ignored when only looking at the emissions. Lastly, other meteorological effects such as wind, turbulence, boundary layer height, and isoprene lifetime are implicitly ignored in this comparison. For instance, higher wind speeds and/or boundary layer heights would lead to lower ambient isoprene concentrations for the same emission source strength. The weak to moderate correlations found at some sites and for some years imply that further



study is required to further understand and interpret the potential sources of model error underpinning the poor performance of our datasets in these cases.

3.3.2 Evaluation using other emission datasets

765 A primary motivation of this work was to develop two BVOC datasets that could be used to support the study of European air quality specifically in relation to CAMS. We therefore perform an evaluation of the SURFEX-MEGAN3.0 emission datasets by comparing them to the MEGAN-MACC (Sindelarova et al., 2014) and CAMS-GLOB-BIO (Sindelarova et al., 2022) BVOC emission datasets that were developed within the frame of CAMS. Furthermore, these datasets have received recent attention and represent the current state of the art in terms of emission modelling for Europe using MEGAN. In the case of the CAMS-GLOB-BIO emissions, we include a comparison with the v1.2, v3.0, and v3.1 versions of this dataset.

770 A timeseries of the monthly mean isoprene emissions for each of these datasets is calculated over the CAMS European domain and plotted in Figure 16. In addition, in Table 5, we present a compilation of the annual emission totals for isoprene for each dataset and each year they are available. According to these comparisons, the MEGAN-MACC emission inventory is the dataset with the highest overall emissions over the period 2018-2020 over the CAMS domain. Meanwhile, the two SURFEX-MEGAN3.0 emission datasets are very similar to one another, and it is possible to see only minor differences during the periods with the highest isoprene emissions in the summertime. The SURFEX-MEGAN3.0 datasets are also very similar to the CAMS-GLOB-BIOv1.2 dataset, which appears to be coincidental given the large difference in input data and MEGAN model version used (i.e., version 2.1 in the case of CAMS-GLOB-BIOv1.2). On the other hand, both the CAMS-GLOB-BIOv3.0 and CAMS-GLOB-BIOv3.1 dataset emissions are both lower overall compared to the SURFEX-MEGAN3.0 datasets.

780 Aside from differences in the absolute magnitude of the emissions, Figure 16 shows that the different datasets also have different seasonalities from each other. The SURFEX-MEGAN3.0 and the CAMS-GLOB-BIOv3.1 represent the two most contrasting emission datasets in terms of seasonality. Out of all of the emission datasets, the SURFEX-MEGAN3.0 emissions consistently peaks earliest in the summer season (i.e., June to July) while the CAMS-GLOB-BIOv3.1 always peak the latest in the summer season (i.e., July to August).

785 We only have this comparison for the 2018-2019 period for v3.0, and the 2018-July 2019 period for v1.2, but both of these two CAMS-GLOB-BIO datasets show a similar month-to-month variability to that of the SURFEX-MEGAN3.0 datasets, i.e., peaks in emissions towards the early part of summer on average. Although both datasets do show a more prolonged peak lasting into July in 2019. Meanwhile, the MEGAN-MACC emissions show their highest peak in emissions in June during 2018 and during July during 2019 and 2020 placing it overall somewhat between the SURFEX-MEGAN3.0 and CAMS-GLOB-BIOv3.1 emissions.

We have evaluated the underlying causes of the differences in seasonality highlighted between the different emission datasets. This evaluation included analysing the different meteorological datasets and their effects on the GAMMATP variable (influenced by temperature and solar radiation), LAI, and the effect of soil moisture upon the SURFEX-MEGAN3.0



emissions (note that the other isoprene emissions in this example do not consider soil moisture effects). This evaluation revealed that in general the LAI datasets (based on Yuan et al., 2011) used to calculate the monthly emission factors in the CAMS-GLOB_BIOv3.1 emissions tend to peak later in the summer than the LAI calculated by SURFEX. Furthermore, the CAMS-GLOB-BIOv3.1 emissions used a climatological mean of LAI derived from MODIS (Sindelarova et al., 2022; Yuan et al., 2011) to calculate the LAI used with the MEGAN activity factor calculations. As a result, in years where SURFEX estimated particularly extreme deviations in LAI from the historical climatology with an early peak in LAI, e.g., 2019, the divergent effect on the emission seasonality is even greater. In terms of the overall magnitude, the annual mean of the SURFEX assimilation analysis LAI over the European domain was $1.12 \text{ m}^2.\text{m}^{-2}$, $1.15 \text{ m}^2.\text{m}^{-2}$, and $1.18 \text{ m}^2.\text{m}^{-2}$ in the years 2018, 2019, and 2020, respectively, compared to the mean of the climatological LAI of $1.00 \text{ m}^2.\text{m}^{-2}$ from Yuan et al. (2011) used by CAMS-GLOB-BIOv3.1. The annual mean LAIs from the OL for 2021 and 2022 were $1.24 \text{ m}^2.\text{m}^{-2}$ and $1.17 \text{ m}^2.\text{m}^{-2}$, respectively. Thus, this indicates that the magnitude of the SURFEX LAI is slightly higher than that used by CAMS-GLOB-BIOv3.1




Figure 16. A time series of the monthly mean total emitted isoprene over the 2018-2022 period as calculated the SURFEX-MEGAN3.0 OL (green solid), SURFEX-MEGAN3.0 Analysis (black dot-dash), MEGAN-MACC (blue dash), CAMS-GLOB-BIOv1.2 (orange solid), CAMS-GLOB-BIOv3.0 (purple dash), and CAMS-GLOB-BIOv3.1 (red dash), emission inventories. The units of the isoprene emission totals are in Tg. The SURFEX-MEGAN3.0 OL and CAMS-GLOB-BIOv3.1 emission inventories are available over the full time period of 2018-2022, while the SURFEX-MEGAN3.0 Analysis and MEGAN-MACC emissions are only available over the period 2018-2020, the CAMS-GLOB-BIOv1.2 are only available 2018-July 2019, and the CAMS-GLOB-BIOv3.0 are only available 2018-2019.

Overall, this evaluation shows that the magnitude of the total isoprene emissions within the CAMS European domain of SURFEX-MEGAN3.0 emissions falls well within the bounds set by existing isoprene emission inventories. Indeed, both datasets fall within the upper and lower bounds set by the most recent published isoprene emission datasets published in association with the MACC and CAMS projects. There are some notable differences from other datasets in terms of the seasonal variability. Further study and evaluation will be required to determine if this difference represents a realistic



820 seasonal variability. The evaluation performed in Sect. 3.3.1 provides some support for the variability represented in both the OL and analysis-based datasets.

825 **Table 5. Annual total emitted isoprene in Tg over the CAMS European modelling domain as determined by the SURFEX-MEGAN3.0 OL, SURFEX-MEGAN3.0 Analysis, CAMS-GLOB-BIOv3.1, and SUMO emission inventories. The emission datasets are available over different time periods. The SURFEX-MEGAN3.0 OL and CAMS-GLOB-BIOv3.1 emission inventories are available over the full tie period of 2018-2022, while the SURFEX-MEGAN3.0 Analysis emissions are only available over the period 2018-2020. These annual emission totals are calculated from 1st January to 31st December in all cases except for CAMS-GLOB-BIOv1.2 that for 2019 were calculated January through to July.**

Year	SURFEX- MEGAN3.0 OL	SURFEX- MEGAN3.0 Analysis	MEGAN- MACC	CAMS- GLOB- BIOv1.2	CAMS- GLOB- BIOv3.0	CAMS-GLOB- BIOv3.1
2018	7.70	7.71	10.81	7.27	4.94	4.67
2019	7.01	7.07	10.92	4.67*	4.63	4.61
2020	6.84	6.91	10.87			4.70
2021	7.50					4.72
2022	7.15					5.14

*Only calculated over January through to July 2019

We quickly discuss the magnitude of the global isoprene emissions from the MEGAN-MACC and CAMS-GLOB-BIO datasets. In the case of MEGAN-MACC, the global BVOC emissions were estimated to be 760 Tg (C) yr⁻¹ with 70% of this carbon mass from isoprene (602.9 Tg (isoprene) yr⁻¹) over the 30-year period from 1980 to 2010 that was studied. In the case of CAMS-GLOB-BIO four estimates were made ranging between 424 and 591 Tg (C) yr⁻¹ (299.1 to 440.5 Tg (isoprene) yr⁻¹), with CAMS-GLOB-BIOv3.1 being the recommended data set according to the authors (591 Tg (C) yr⁻¹ and 440.5 Tg (isoprene) yr⁻¹). Overall, the emission totals calculated over the CAMS European domain and presented in Table 5 represent only a small fraction of the global emission totals.

4 Data Availability

Both of the isoprene emission datasets are available as hourly means at a spatial resolution of 0.1° × 0.1° over the CAMS European domain. The data can be accessed from the SEEDS project data portal (<https://www.seedsproject.eu/data>, last access: 17 July 2025) and are available in Zarr format or can alternatively be downloaded in netcdf format using a short python script by following the instructions provided. The other compounds (including isoprene again) are available on the Harvard Dataverse inside the dataverse for the SEEDS project (<https://dataverse.harvard.edu/dataverse/Horizon-SEEDS-project>, last access: 17 July 2025). Two different datasets were created within the SEEDS dataverse, one each for the analysis (Hamer et al., 2025a) and OL (Hamer et al., 2025b) (<https://doi.org/10.7910/DVN/LAUVTU> and <https://doi.org/10.7910/DVN/69G1FX>, respectively, last access: 17 July 2025).



845 5 Conclusions

We have presented two new datasets of BVOC emission inventories that cover the CAMS European domain that are both publicly available and ready for use in applications related to the study of European scale air quality. As per this intention, both datasets were produced at a spatial resolution of $0.1^\circ \times 0.1^\circ$ exactly matching the spatial domain utilised by the CAMS European regional forecast and analyses. The datasets were produced within the framework of the EU's SEEDS project and
 850 are called SURFEX-MEGAN3.0 OL and SURFEX-MEGAN3.0 analysis. The OL dataset is available over a five-year period from 2018-2022 and the analysis dataset is available over the three-year period 2018-2020.

Another motivation for the production of this new dataset was to address key uncertainties in BVOC emission modelling arising from representation of vegetation phenology and to move towards an Earth-systems approach allowing feedbacks and interactions between meteorology/climate, vegetation, and atmospheric composition. To this end, we applied a methodology
 855 based on using a land surface model capable of dynamic vegetation modelling that responds to an externally applied meteorological forcing. To further address uncertainties in the simulation of vegetation phenology, the land surface model is further supported by data assimilation of satellite observations of LAI to create an assimilation analysis, which forms the basis of the analysis emission dataset. In addition, we also used the most recent version (at the beginning of the study) of MEGAN3.0.

860 Since we used SURFEX land surface model within the MEGAN3.0 production chain, we perform an evaluation of the key variables estimated by SURFEX that are used within MEGAN3.0. We evaluated both LAI and RZSM. In the case of LAI, we use observations of SIF from the TROPOMI satellite to establish that the SURFEX LAI datasets have good skill over the main body of Europe, the UK, Italy, Ireland, southern Sweden and into eastern Europe as well and that the analysis for LAI had improved correlation to SIF compared to the OL. The RZSM was evaluated against in-situ observations in south-western
 865 France and was shown to have strong year-round correlation and moderate correlation during the summertime. The RZSM analysis performed better than the RZSM OL.

Analysis of the OL emission dataset over the 5-year period 2018-2022 shows that the average annual isoprene emissions were 7.20 Tg yr^{-1} and had a standard deviation of 0.28 Tg yr^{-1} . The difference between the year with the largest (2018) and smallest (2020) emissions was equivalent to 11.4% of the average annual emission. The analysis-based average (calculated
 870 over 3 years 2018-2020) is 7.19 Tg yr^{-1} with a standard deviation of 0.40 Tg yr^{-1} . Note that the average of the OL-based emissions over this same time period was 7.14 Tg yr^{-1} with a standard deviation of 0.44 Tg yr^{-1} . The areas with the highest inter-annual variability correspond to areas of large standing woodland with the densest levels of emitting plant species. Analysis showed that this variability is induced by (listed with decreasing importance) variability in LAI, meteorology (temperature and radiation), and RZSM. LAI had the consistently highest correlation with isoprene emissions on the inter-
 875 annual timescale indicating the important role that variations in vegetation and any induced effects by meteorology (e.g., affecting the growing season and drought) can play.



Analysis of the monthly variations in the OL dataset averaged over the 5-year period 2018-2022 show that monthly isoprene emissions follow the development of the growing season within different climatic zones. Isoprene emissions peak during the summer period (June through August) and are highest on average in July. An analysis of the underlying factors (listed in decreasing importance) driving the monthly variability showed that meteorology (temperature and radiation), LAI, and RSZM impacted the changes in isoprene emissions. This finding was reproduced consistently for all five years in the OL dataset. The importance of situations where extreme meteorological conditions can impact vegetation were highlighted in the context of the summer of 2019. Here, it was shown that reductions in LAI driven by extreme heat and drought led to decreased isoprene emissions in the month of July compared to June of that same year.

The evaluations and analysis performed on these datasets at annual and monthly timescales highlights the importance of including a dynamic representation of vegetation by using a land surface model in the MEGAN emission production chain. Furthermore, this highlights the potential advantages of an Earth system approach to BVOC emission modelling. Performing the LAI data assimilation step does not impact the average annual LAI levels or the total magnitude of isoprene emissions at the continental scale to a significant degree either on monthly or annual timescales. However, larger absolute and relative variations in emissions and LAI do occur when looking at smaller spatial scales. The impact of the LAI data assimilation step is largest during the summertime.

We evaluate the temporal (daily) correlation of isoprene emissions with in-situ observations of isoprene concentrations from 8 sites across Europe. We find that both the OL and analysis datasets have moderate ($> 0.4 R^2$) to strong ($> 0.7 R^2$) temporal correlation with the in-situ observations for the majority of years and sites analysed with only 4 cases (site-year pairings) showing weak ($< 0.4 R^2$) correlation. This indicates that, overall, this method has good skill at estimating the temporal variability in the emissions. No overall strong difference between the skill of the OL and analysis datasets was found when the emissions were evaluated in this manner using in-situ observations.

Compared to other isoprene emission datasets produced within the frame of MACC and CAMS, the two SURFEX-MEGAN3.0 datasets estimate emission totals lie between the minimum (CAMS-GLOB-BIOv3.1) and maximum (MEGAN-MACC) estimates in isoprene emissions over the CAMS European domain for the 2018-2020 time period. Indeed, the SURFEX-MEGAN3.0 isoprene emissions were very similar in magnitude to the CAMS-GLOB-BIOv1.2 isoprene emissions. We did note large differences in the seasonality of the different datasets whereby the SURFEX-MEGAN3.0, CAMS-GLOB-BIOv1.2, and CAMS-GLOB-BIOv3.0 emissions peaked earlier in summer, the MEGAN-MACC emissions peaked in the middle of summer, and the CAMS-GLOB-BIOv3.1 emissions peaked the latest in summer. An investigation showed that the differences between the earliest (SURFEX-MEGAN3.0) and latest peaking emissions (CAMS-GLOB-BIOv3.1) were due to differences in the seasonality of the LAI datasets used in the two different approaches whereby the LAI estimated by the SURFEX OL and analysis tended to peak earlier than the MODIS based LAI used for CAMS-GLOB-BIOv3.1 (Sindelarova et al., 2022; Yuan et al., 2011). These differences lead to substantial differences between these two emission datasets when compared on a month-by-month basis like this and it highlights the importance of our approach attempting to improve the temporal variability of LAI datasets used in the MEGAN algorithm. Indeed, Guenther et al., 2006



identify that uncertainties in LAI input data could impact global, annual isoprene emissions from -11% to 29%, but we see much larger relative differences between the SURFEX-MEGAN3 and CAMS-GLOB-BIOv3.1 emissions over Europe when compared on a month-to-month basis. Future work aimed at further intercomparisons of LAI input datasets would be valuable to support further advancement on this topic.

915 Guenther et al. (2006) and Sindelarova et al. (2022) point to other sources of uncertainty that we do not directly address in this work, which include mapping and consistency of plant functional types (PFTs) and the manner in which the impacts of soil moisture and drought are treated within the MEGAN algorithm. Addressing these uncertainties in future work will be a priority in order to further improve the quality of emissions produced using the SURFEX-MEGAN algorithm. Indeed, the work of Oumami et al. (2024) already highlights the advantage of harmonizing PFTs between SURFEX and MEGAN, so it would be advantageous to use the SURFEX PFTs derived from either ECOCLIMAP-II (Faroux et al., 2013) or ECOCLIMAP-SG (Druel et al., 2022) as a basis for calculating the EFs used by MEGAN in the future.

Accurately representing the impacts of soil moisture and drought on BVOC emissions is another important area of research. In light of the large uncertainties of the role of soil moisture and drought, some groups (Oomen et al., 2024; Sindelarova et al., 2022) have taken an approach to exclude the influence of soil moisture in the MEGAN algorithm. There is compelling evidence (Bonn et al., 2019; Jiang et al., 2018; Pegoraro et al., 2004; Yuan et al., 2016; Zheng et al., 2017), however, that vegetation response to soil moisture and drought plays an important role in BVOC emission variability, and this remains an open research question of how this should be addressed in a generalized way within the MEGAN model framework. The newly proposed parametrization of soil moisture stress on isoprene described in (Wang et al., 2022) was tested in CAMS-GLOB-BIO and led to a decrease in global isoprene total emissions by 15 %, with even larger reductions in dry regions such as Australia, sub-Saharan Africa or Middle East (K. Sindelarova, personal communication). Future work should therefore address this source of uncertainty by testing different algorithms for different drought situations for varied climatic zones and ecotypes.

Author Contributions

PDH co-developed the software interface between SURFEX and MEGAN3.0, co-designed the experiments, co-wrote the research proposal, carried out most of the technical data processing, led the scientific investigation related to the BVOC emissions, and wrote most of the manuscript. MM contributed significantly to the data visualization and supported the scientific investigation. ORM carried out the analysis using TROPOMI SIF data. BB ran setup and the SURFEX simulations and performed many of the related data processing tasks. JCC led the portion of the scientific investigation related to SURFEX and the TROPOMI SIF data, co-designed the experiments, and co-wrote sections of text related to SURFEX. VM co-wrote the research proposal, co-designed the experiments, contributed to the design and structure of the manuscript, and co-led the scientific investigation. AG contributed with expertise on the MEGAN3.0 model, co-designed the experiments, wrote the section on MEGAN3.0 emission factors, and reviewed all descriptions of MEGAN3.0. HT co-developed the



software interface between SURFEX and MEGAN3.0. IV carried out data processing and software development tasks related to the MEGAN3.0 emission factor code. SE downloaded and processed all data from ECMWF. G. Sousa Santos processed CAMS ozone data to derive the ozone damage gamma in MEGAN3.0. KS contributed to the intercomparison with the CAMS-GLOB-BIO datasets. DS contributed to the intercomparison with the CAMS-GLOB-BIO datasets. NS contributed with expertise related to isoprene observations based on online GC-MS. LT was the coordinator of the SEEDS project, co-designed the experiments, and led and followed up the work towards the main results of the project.

Competing interests

The contact author has declared that none of the authors have any competing interests.

Financial support

This work has been funded by the European Union's Horizon2020 research and innovation programme under grant agreement No 101004318. The contributions from H. Trimmel were supported by Wiener Wissenschafts-, Forschungs-und Technologiefonds (WWTF) within the Environmental System Research Call 2017 [grant number [ESR17-027](#), project UOZONE].

References

- Albergel, C., Munier, S., Leroux, D. J., Dewaele, H., Fairbairn, D., Barbu, A. L., Gelati, E., Dorigo, W., Faroux, S., Meurey, C., Le Moigne, P., Decharme, B., Mahfouf, J.-F., and Calvet, J.-C.: Sequential assimilation of satellite-derived vegetation and soil moisture products using SURFEX_v8.0: LDAS-Monde assessment~over the Euro-Mediterranean area, *Geosci Model Dev*, 10, 3889–3912, <https://doi.org/10.5194/gmd-10-3889-2017>, 2017.
- Albergel, C., Dutra, E., Bonan, B., Zheng, Y., Munier, S., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., and Calvet, J.-C.: Monitoring and Forecasting the Impact of the 2018 Summer Heatwave on Vegetation, *Remote Sens (Basel)*, 11, <https://doi.org/10.3390/rs11050520>, 2019a.
- Albergel, C., Dutra, E., Bonan, B., Zheng, Y., Munier, S., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., and Calvet, J.-C.: Monitoring and Forecasting the Impact of the 2018 Summer Heatwave on Vegetation, *Remote Sens (Basel)*, 11, <https://doi.org/10.3390/rs11050520>, 2019b.
- Arneth, A., Niinemets, Ü., Pressley, S., Bàck, J., Hari, P., Karl, T., Noe, S., Prentice, I. C., Serça, D., Hickler, T., Wolf, A., and Smith, B.: Process-based estimates of terrestrial ecosystem isoprene emissions: Incorporating the effects of a direct CO₂-isoprene interaction, *Atmos Chem Phys*, 7, 31–53, <https://doi.org/10.5194/acp-7-31-2007>, 2007.



- 970 Arneth, A., Schurgers, G., Lathiere, J., Duhl, T., Beerling, D. J., Hewitt, C. N., Martin, M., and Guenther, A.: Global terrestrial isoprene emission models: Sensitivity to variability in climate and vegetation, *Atmos Chem Phys*, 11, 8037–8052, <https://doi.org/10.5194/acp-11-8037-2011>, 2011.
- Bonn, B., Magh, R. K., Rombach, J., and Kreuzwieser, J.: Biogenic isoprenoid emissions under drought stress: Different responses for isoprene and terpenes, *Biogeosciences*, 16, 4627–4645, <https://doi.org/10.5194/bg-16-4627-2019>, 2019.
- 975 Brown, L. A., Meier, C., Morris, H., Pastor-Guzman, J., Bai, G., Lerebourg, C., Gobron, N., Lanconelli, C., Clerici, M., and Dash, J.: Evaluation of global leaf area index and fraction of absorbed photosynthetically active radiation products over North America using Copernicus Ground Based Observations for Validation data, *Remote Sens Environ*, 247, <https://doi.org/10.1016/j.rse.2020.111935>, 2020.
- Broxton, P. D., Zeng, X., Scheftic, W., and Troch, P. A.: A MODIS-based global 1-km maximum green vegetation fraction dataset, *J Appl Meteorol Climatol*, 53, 1996–2004, 2014.
- 980 Calvet, J.-C., Noilhan, J., Roujean, J.-L., Bessemoulin, P., Cabelguenne, M., Olioso, A., and Wigneron, J.-P.: An interactive vegetation SVAT model tested against data from six contrasting sites, *Agric For Meteorol*, 92, 73–95, [https://doi.org/https://doi.org/10.1016/S0168-1923\(98\)00091-4](https://doi.org/https://doi.org/10.1016/S0168-1923(98)00091-4), 1998.
- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos Chem Phys*, 9, 4987–5005, <https://doi.org/10.5194/acp-9-4987-2009>, 2009.
- 985 Decharme, B., Boone, A., Delire, C., and Noilhan, J.: Local evaluation of the Interaction between Soil Biosphere Atmosphere soil multilayer diffusion scheme using four pedotransfer functions, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/https://doi.org/10.1029/2011JD016002>, 2011.
- Donahue, N. M., Robinson, A. L., and Pandis, S. N.: Atmospheric organic particulate matter: From smoke to secondary organic aerosol, *Atmos Environ*, 43, 94–106, 2009.
- 990 Druel, A., Munier, S., Mucia, A., Albergel, C., and Calvet, J. C.: Implementation of a new crop phenology and irrigation scheme in the ISBA land surface model using SURFEX-v8.1, *Geosci Model Dev*, 15, 8453–8471, <https://doi.org/10.5194/gmd-15-8453-2022>, 2022.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci Model Dev*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- Faroux, S., Kaptué Tchuenté, A. T., Roujean, J.-L., Masson, V., Martin, E., and Le Moigne, P.: ECOCLIMAP-II/Europe: a twofold database of ecosystems and surface parameters at 1 km resolution based on satellite information for use in land surface, meteorological and climate models, *Geosci Model Dev*, 6, 563–582, <https://doi.org/10.5194/gmd-6-563-2013>, 2013a.
- 1000 Faroux, S., Kaptué Tchuenté, A. T., Roujean, J.-L., Masson, V., Martin, E., and Le Moigne, P.: ECOCLIMAP-II/Europe: a twofold database of ecosystems and surface parameters at 1 km resolution based on satellite information for use in land



- surface, meteorological and climate models, *Geosci Model Dev*, 6, 563–582, <https://doi.org/10.5194/gmd-6-563-2013>, 2013b.
- Gibelin, A.-L., Calvet, J.-C., Roujean, J.-L., Jarlan, L., and Los, S. O.: Ability of the land surface model ISBA-A-gs to simulate leaf area index at the global scale: Comparison with satellites products, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006691>, 2006.
- Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., and others: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research: Atmospheres*, 100, 8873–8892, 1995.
- Guenther, A., Karl, T., Harley, P., Weidinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature) and Physics Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases an, *Atmos Chem Phys*, 3181–3210, 2006a.
- Guenther, A., Karl, T., Harley, P., Weidinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature) and Physics Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases an, *Atmos Chem Phys*, 3181–3210, 2006b.
- Guenther, A., Jiang, X., Shah, T., Huang, L., Kemball-Cook, S., and Yarwood, G.: Model of emissions of gases and aerosol from nature version 3 (MEGAN3) for estimating biogenic emissions, in: *Air Pollution Modeling and its Application XXVI* 36, 187–192, 2020a.
- Guenther, A., Jiang, X., Shah, T., Huang, L., Kemball-Cook, S., and Yarwood, G.: Model of emissions of gases and aerosol from nature version 3 (MEGAN3) for estimating biogenic emissions, in: *Air Pollution Modeling and its Application XXVI* 36, 187–192, 2020b.
- Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and monoterpene emission rate variability: model evaluations and sensitivity analyses, *Journal of Geophysical Research: Atmospheres*, 98, 12609–12617, 1993.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci Model Dev*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012a.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci Model Dev*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012b.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci Model Dev*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012c.



- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci Model Dev*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012d.
- 1040 Hamer, P., Trimmel, H., Markelj, M., Rojas-Munoz, O., Bonan, B., Calvet, J.-C., Vallejo, I., Eckhardt, S., Sousa-Santos, G., Marecal, V., and Tarrason, L.: SURFEX-MEGAN3.0 analysis based BVOC emissions, <https://doi.org/10.7910/DVN/LAUVTU>, 2025a.
- Hamer, P., Trimmel, H., Markelj, M., Rojas-Munoz, O., Bonan, B., Calvet, J.-C., Vallejo, I., Eckhardt, S., Sousa-Santos, G., Marecal, V., and Tarrason, L.: SURFEX-MEGAN3.0 open-loop based BVOC emissions, <https://doi.org/10.7910/DVN/69G1FX>, 2025b.
- 1045 Hansen, M. C., DeFries, R. S., Townshend, J. R. G., Carroll, M., Dimiceli, C., and Sohlberg, R. A.: Global percent tree cover at a spatial resolution of 500 meters: First results of the MODIS vegetation continuous fields algorithm, *Earth Interact*, 7, 1–15, 2003.
- Homer, C., Dewitz, J., Yang, L., Jin, S., Danielson, P., Xian, G., Coulston, J., Herold, N., Wickham, J., and Megown, K.: Completion of the 2011 National Land Cover Database for the conterminous United States—representing a decade of land
 1050 cover change information, *Photogramm Eng Remote Sensing*, 81, 345–354, 2015.
- Jiang, X., Guenther, A., Potosnak, M., Geron, C., Seco, R., Karl, T., Kim, S., Gu, L., and Pallardy, S.: Isoprene emission response to drought and the impact on global atmospheric chemistry, *Atmos Environ*, 183, 69–83, <https://doi.org/10.1016/j.atmosenv.2018.01.026>, 2018.
- Khan, M. A. H., Schlich, B.-L., Jenkin, M. E., Shallcross, B. M. A., Moseley, K., Walker, C., Morris, W. C., Derwent, R. G.,
 1055 Percival, C. J., and Shallcross, D. E.: A two-decade anthropogenic and biogenic isoprene emissions study in a London urban background and a London urban traffic site, *Atmosphere (Basel)*, 9, 387, 2018.
- Langford, B., Cash, J., Acton, W. J. F., Valach, A. C., Hewitt, C. N., Fares, S., Goded, I., Gruening, C., House, E., Kalogridis, A.-C., Gros, V., Schafers, R., Thomas, R., Broadmeadow, M., and Nemitz, E.: Isoprene emission potentials from European oak forests derived from canopy flux measurements: an assessment of uncertainties and inter-algorithm
 1060 variability, *Biogeosciences*, 14, 5571–5594, <https://doi.org/10.5194/bg-14-5571-2017>, 2017.
- Laothawornkitkul, J., Taylor, J. E., Paul, N. D., and Hewitt, C. N.: Biogenic volatile organic compounds in the Earth system, *New Phytologist*, 183, 27–51, 2009.
- Latham, J., Cumani, R., Rosati, I., and Bloise, M.: Global land cover share (GLC-SHARE) database beta-release version 1.0-2014, FAO: Rome, Italy, 29, 2014.
- 1065 Lathière, J., Hauglustaine, D. A., Friend, A. D., De Noblet-Ducoudré, N., Viovy, N., and Folberth, G. A.: Impact of climate variability and land use changes on global biogenic volatile organic compound emissions, *Atmos Chem Phys*, 6, 2129–2146, <https://doi.org/10.5194/acp-6-2129-2006>, 2006.



- Lathière, J., Hewitt, C. N., and Beerling, D. J.: Sensitivity of isoprene emissions from the terrestrial biosphere to 20th century changes in atmospheric CO₂ concentration, climate, and land use, *Global Biogeochem Cycles*, 24, n/a-n/a, 1070 <https://doi.org/10.1029/2009gb003548>, 2010.
- Levis, S., Wiedinmyer, C., Bonan, G. B., and Guenther, A.: Simulating biogenic volatile organic compound emissions in the Community Climate System Model, *Journal of Geophysical Research: Atmospheres*, 108, 1–9, <https://doi.org/10.1029/2002jd003203>, 2003.
- Lin, Y., Lun, X., Tang, W., Zhang, Z., Jing, X., Fan, C., and Wang, Q.: Characteristics and chemical reactivity of biogenic 1075 volatile organic compounds from dominant forest species in the Jing-Jin-Ji area, China, *For Ecosyst*, 8, 1–14, 2021.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouysse, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., 1080 Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geosci Model Dev*, 6, 929–960, <https://doi.org/10.5194/gmd-6-929-2013>, 2013.
- Millet, D. B., Jacob, D. J., Turquety, S., Hudman, R. C., Wu, S., Fried, A., Walega, J., Heikes, B. G., Blake, D. R., Singh, H. B., and others: Formaldehyde distribution over North America: Implications for satellite retrievals of formaldehyde columns 1085 and isoprene emission, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Müller, J. F., Stavrou, T., Wallens, S., De Smedt, I., Van Roozendaal, M., Potosnak, M. J., Rinne, J., Munger, B., Goldstein, A., and Guenther, A. B.: Global isoprene emissions estimated using MEGAN, ECMWF analyses and a detailed canopy environment model, *Atmos Chem Phys*, 8, 1329–1341, <https://doi.org/10.5194/acp-8-1329-2008>, 2008.
- Naik, V., Delire, C., and Wuebbles, D. J.: Sensitivity of global biogenic isoprenoid emissions to climate variability and 1090 atmospheric CO₂, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003jd004236>, 2004.
- Niinemets, Ü., Kuhn, U., Harley, P. C., Staudt, M., Arneth, A., Cescatti, A., Ciccioli, P., Copolovici, L., Geron, C., Guenther, A., and others: Estimations of isoprenoid emission capacity from enclosure studies: measurements, data processing, quality and standardized measurement protocols, *Biogeosciences*, 8, 2209–2246, 2011.
- Oomen, G. M., Müller, J. F., Stavrou, T., De Smedt, I., Blumenstock, T., Kivi, R., Makarova, M., Palm, M., Röhling, A., 1095 Té, Y., Vigouroux, C., Friedrich, M. M., Frieß, U., Hendrick, F., Merlaud, A., PETERS, A., Richter, A., Van Roozendaal, M., and Wagner, T.: Weekly derived top-down volatile-organic-compound fluxes over Europe from TROPOMI HCHO data from 2018 to 2021, *Atmos Chem Phys*, 24, 449–474, <https://doi.org/10.5194/acp-24-449-2024>, 2024.
- Oumami, S., Arteta, J., Guidard, V., Tulet, P., and Hamer, P. D.: Evaluation of isoprene emissions from the coupled model SURFEX–MEGANv2.1, *Geosci Model Dev*, 17, 3385–3408, <https://doi.org/10.5194/gmd-17-3385-2024>, 2024.
- 1100 Owens, R. G., Hewson, T. D.: ECMWF Forecast User Guide, Reading, <https://doi.org/10.21957/m1cs7h>, 2018.



- Pacifico, F., Harrison, S. P., Jones, C. D., Arneth, A., Sitch, S., Weedon, G. P., Barkley, M. P., Palmer, P. I., Serça, D., Potosnak, M., Fu, T. M., Goldstein, A., Bai, J., and Schurgers, G.: Evaluation of a photosynthesis-based biogenic isoprene emission scheme in JULES and simulation of isoprene emissions under present-day climate conditions, *Atmos Chem Phys*, 11, 4371–4389, <https://doi.org/10.5194/acp-11-4371-2011>, 2011.
- 1105 Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Pegoraro, E., Rey, A., Greenberg, J., Harley, P., Grace, J., Malhi, Y., and Guenther, A.: Effect of drought on isoprene emission rates from leaves of *Quercus virginiana* Mill., *Atmos Environ*, 38, 6149–6156, <https://doi.org/10.1016/j.atmosenv.2004.07.028>, 2004.
- 1110 Pfister, G. G., Emmons, L. K., Hess, P. G., Lamarque, J. F., Orlando, J. J., Walters, S., Guenther, A., Palmer, P. I., and Lawrence, P. J.: Contribution of isoprene to chemical budgets: A model tracer study with the NCAR CTM MOZART-4, *Journal of Geophysical Research Atmospheres*, 113, 1–21, <https://doi.org/10.1029/2007JD008948>, 2008.
- Plass-Dülmer, C., Schmidbauer, N., Slemr, J., Slemr, F., and D’Souza, H.: European hydrocarbon intercomparison experiment AMOHA part 4: Canister sampling of ambient air, *Journal of Geophysical Research Atmospheres*, 111, <https://doi.org/10.1029/2005JD006351>, 2006.
- 1115 Potter, C. S., Alexander, S. E., Coughlan, J. C., and Klooster, S. A.: Modeling biogenic emissions of isoprene: Exploration of model drivers, climate control algorithms, and use of global satellite observations, *Atmos Environ*, 35, 6151–6165, [https://doi.org/10.1016/S1352-2310\(01\)00390-9](https://doi.org/10.1016/S1352-2310(01)00390-9), 2001.
- 1120 Shim, C., Wang, Y., Choi, Y., Palmer, P. I., Abbot, D. S., and Chance, K.: Constraining global isoprene emissions with Global Ozone Monitoring Experiment (GOME) formaldehyde column measurements, *Journal of Geophysical Research Atmospheres*, 110, 1–14, <https://doi.org/10.1029/2004JD005629>, 2005.
- Simpson, D., Winiwarter, W., Börjesson, G., Cinderby, S., Ferreiro, A., Guenther, A., Hewitt, C. N., Janson, R., Khalil, M. A. K., Owen, S., and others: Inventorying emissions from nature in Europe, *Journal of Geophysical Research: Atmospheres*, 104, 8113–8152, 1999.
- 1125 Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J. P., Valdebenito, A., and Wind, P.: The EMEP MSC-W chemical transport model – Technical description, *Atmos Chem Phys*, 12, 7825–7865, <https://doi.org/10.5194/acp-12-7825-2012>, 2012.
- 1130 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrou, T., Müller, J. F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos Chem Phys*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014a.



- Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., Müller, J. F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos Chem Phys*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014b.
- 1135 Sindelarova, K., Markova, J., Simpson, D., Huszar, P., Karlicky, J., Darras, S., and Granier, C.: High-resolution biogenic global emission inventory for the time period 2000–2019 for air quality modelling, *Earth Syst Sci Data*, 14, 251–270, <https://doi.org/10.5194/essd-14-251-2022>, 2022a.
- 1140 Sindelarova, K., Markova, J., Simpson, D., Huszar, P., Karlicky, J., Darras, S., and Granier, C.: High-resolution biogenic global emission inventory for the time period 2000–2019 for air quality modelling, *Earth Syst Sci Data*, 14, 251–270, <https://doi.org/10.5194/essd-14-251-2022>, 2022b.
- Sporre, M. K., Blichner, S. M., Karset, I. H. H., Makkonen, R., and Berntsen, T. K.: BVOC--aerosol--climate feedbacks investigated using NorESM, *Atmos Chem Phys*, 19, 4763–4782, 2019.
- 1145 Stavrakou, T., Müller, J.-F., De Smedt, I., Van Roozendaal, M., Kanakidou, M., Vrekoussis, M., Wittrock, F., Richter, A., and Burrows, J. P.: The continental source of glyoxal estimated by the synergistic use of spaceborne measurements and inverse modelling, *Atmos Chem Phys*, 9, 8431–8446, 2009.
- Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, *Journal of Geophysical Research: Atmospheres*, 102, 25847–25879, 1997.
- Szczypta, C., Calvet, J.-C., Maignan, F., Dorigo, W., Baret, F., and Ciais, P.: Suitability of modelled and remotely sensed essential climate variables for monitoring Euro-Mediterranean droughts, *Geosci Model Dev*, 7, 931–946, <https://doi.org/10.5194/gmd-7-931-2014>, 2014.
- 1150 Tao, Z. and Jain, A. K.: Modeling of global biogenic emissions for key indirect greenhouse gases and their response to atmospheric CO₂ increases and changes in land cover and climate, *Journal of Geophysical Research Atmospheres*, 110, 1–13, <https://doi.org/10.1029/2005JD005874>, 2005.
- 1155 Thornhill, G., Collins, W., Olivié, D., Skeie, R. B., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O'Connor, F. M., Paulot, F., Schulz, M., Scott, C. E., Séférian, R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.: Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models, *Atmos Chem Phys*, 21, 1105–1126, <https://doi.org/10.5194/acp-21-1105-2021>, 2021.
- 1160 Tuanmu, M.-N. and Jetz, W.: A global 1-km consensus land-cover product for biodiversity and ecosystem modelling, *Global Ecology and Biogeography*, 23, 1031–1045, 2014.
- Verger, A., Sánchez-Zapero, J., Weiss, M., Descals, A., Camacho, F., Lacaze, R., and Baret, F.: GEOV2: Improved smoothed and gap filled time series of LAI, FAPAR and FCover 1 km Copernicus Global Land products, *International Journal of Applied Earth Observation and Geoinformation*, 123, <https://doi.org/10.1016/j.jag.2023.103479>, 2023.



- 1165 Wang, H., Lu, X., Seco, R., Stavrakou, T., Karl, T., Jiang, X., Gu, L., and Guenther, A. B.: Modeling Isoprene Emission
 Response to Drought and Heatwaves Within MEGAN Using Evapotranspiration Data and by Coupling With the Community
 Land Model, *J Adv Model Earth Syst*, 14, <https://doi.org/10.1029/2022MS003174>, 2022.
- Wiedinmyer, C., Tie, X., Guenther, A., Neilson, R., and Granier, C.: Future changes in biogenic isoprene emissions: How
 might they affect regional and global atmospheric chemistry?, *Earth Interact*, 10, 1–19, <https://doi.org/10.1175/EI174.1>,
 1170 2006.
- Young, P. J., Arneth, A., Schurgers, G., Zeng, G., and Pyle, J. A.: The CO₂ inhibition of terrestrial isoprene emission
 significantly affects future ozone projections, *Atmos Chem Phys*, 9, 2793–2803, <https://doi.org/10.5194/acp-9-2793-2009>,
 2009.
- Yuan, H., Dai, Y., Xiao, Z., Ji, D., and Shangguan, W.: Reprocessing the MODIS Leaf Area Index products for land surface
 1175 and climate modelling, *Remote Sens Environ*, 115, 1171–1187, <https://doi.org/10.1016/j.rse.2011.01.001>, 2011.
- Yuan, X., Calatayud, V., Gao, F., Fares, S., Paoletti, E., Tian, Y., and Feng, Z.: Interaction of drought and ozone exposure on
 isoprene emission from extensively cultivated poplar, *Plant Cell Environ*, 39, 2276–2287, <https://doi.org/10.1111/pce.12798>,
 2016.
- Zhang, M., Zhao, C., Yang, Y., Du, Q., Shen, Y., Lin, S., Gu, D., Su, W., and Liu, C.: Modeling sensitivities of BVOCs to
 1180 different versions of MEGAN emission schemes in WRF-Chem (v3.6) and its impacts over eastern China, *Geosci Model
 Dev*, 14, 6155–6175, <https://doi.org/10.5194/gmd-14-6155-2021>, 2021.
- Zheng, Y., Unger, N., Tadić, J. M., Seco, R., Guenther, A. B., Barkley, M. P., Potosnak, M. J., Murray, L. T., Michalak, A.
 M., Qiu, X., Kim, S., Karl, T., Gu, L., and Pallardy, S. G.: Drought impacts on photosynthesis, isoprene emission and
 atmospheric formaldehyde in a mid-latitude forest, *Atmos Environ*, 167, 190–201,
 1185 <https://doi.org/10.1016/j.atmosenv.2017.08.017>, 2017.