



UEx-L-Eddies: Decadal and global long-lived mesoscale eddy trajectories with coincident air-sea CO₂ fluxes and biogeochemical conditions

Daniel J. Ford¹, Jamie D. Shutler¹, Katy L. Sheen¹, Gavin H. Tilstone², and Vassilis Kitidis²

¹ Centre for Geography and Environmental Sciences (CGES), University of Exeter, Penryn, UK.

² Plymouth Marine Laboratory, Plymouth, UK

Correspondence to: Daniel J. Ford (d.ford@exeter.ac.uk)

Abstract. Mesoscale eddies are prevalent features within the global ocean that modify the physical, chemical and biological properties as they move and evolve. These modifications can alter the air-sea exchange of CO₂, and therefore these features may be hotspots for enhanced or reduced CO₂ uptake compared to the surrounding environment. The understanding of the global and regional effect of mesoscale eddies on ocean CO₂ uptake is however limited and largely based on single eddies or small regional subsets. Here, we provide a global dataset of 5996 long lived eddies trajectories (lifetimes greater than a year) with corresponding air-sea CO₂ fluxes all tracked using a Lagrangian approach between 1993 to 2022. The trajectories comprise 3244 anticyclonic ('warm core') and 2752 cyclonic ('cold core') eddies and the dataset provides the biogeochemical conditions, including the CO₂ fluxes, within and outside each eddy. The dataset refines a previous regional methodology with a focus on climate quality environmental parameters and uses a global neural network for estimating the fugacity of CO₂ in seawater (fCO₂ (sw)) along with a comprehensive air-sea CO₂ flux uncertainty budget. These refinements provide a robust foundation for studying the modulation of air-sea CO₂ fluxes by mesoscale eddies. As an example use of the dataset, we investigate the role of mesoscale eddies in modifying the global and regional air-sea CO₂ fluxes, by comparing the eddy driven air-sea CO₂ flux to that of the surrounding environment. We find that globally, long-lived anticyclonic eddies enhanced the CO₂ sink by 4.5 ± 2.8 % (95 % confidence), while long-lived cyclonic eddies reduce the CO₂ sink by 0.7 ± 2.6 %. Collectively, the long-lived eddies indicate an enhancement of the ocean CO₂ sink by 2.7 ± 1.1 Tg C yr⁻¹. Propagating the air-sea CO₂ flux uncertainties was found to be a key component needed to fully understand apparent differences between previous regional and global studies. The long lived eddies (UEx-L-Eddies) dataset is available on Zenodo at <https://doi.org/10.5281/ZENODO.16355763> (Ford et al., 2025).



1. Introduction

Mesoscale eddies are known to affect the physical, chemical and biological properties of the oceans (Dufois et al., 2016; Frenger et al., 2013; Laxenaire et al., 2019; Li et al., 2025; Nencioli et al., 2018; Orselli et al., 2019b, a; Pezzi et al., 2021). These rotating bodies of water have radii on the order 100 km, lifetimes from a few days to multiple years, and can transit ocean basins transporting distinct water masses within them (Chelton et al., 2011; Pegliasco et al., 2022b). Eddies generally fall into two categories; (1) anticyclonic and (2) cyclonic. Anticyclonic eddies are associated with high pressure centres, clockwise rotation in the Northern Hemisphere (or anticlockwise in the Southern Hemisphere), warmer sea surface temperatures (SST), and a depression of isopycnals (and downwelling of water within the eddy core). Whereas cyclonic eddies are generally the opposite; low pressure centres, anticlockwise rotation in the Northern Hemisphere (or clockwise in the Southern Hemisphere), cooler SSTs, and an elevation of isopycnals (and upwelling in the eddy core). During their lifetimes, these eddies can alter the air-sea CO₂ exchange through their modification of the ocean and atmospheric properties. As the CO₂ solubility in seawater is highly temperature sensitive, the fCO₂ (sw) in anticyclonic eddies could theoretically be elevated and therefore the features may act as a weaker CO₂ sink or stronger CO₂ source compared to the surrounding environment. Conversely the opposite may be true for cyclonic eddies, with reduced fCO₂ (sw), and increased capacity to act as a stronger ocean CO₂ sink. But mesoscale eddies are complex dynamic features, and these generalisations may not always apply as their response will always be dependent upon the ocean basin conditions where the eddy formed and through which the eddy moves, along with how they evolve and interact with that ocean water and the atmosphere. For example, Chen et al. (2007) identified a cyclonic eddy acting as a weaker CO₂ sink compared to the surrounding environment due to upwelling of CO₂ and nutrients within the eddy core. Orselli et al. (2019b) showed six anticyclonic Agulhas eddies that were acting as a stronger CO₂ sink (than the surrounding water) during Austral winter. Pezzi et al. (2021) identified an anticyclonic eddy acting as a strong CO₂ source in the Southwestern Atlantic. Whereas, through using a biogeochemical model, Song et al. (2016) suggested that these eddy modifications may have seasonal difference, whereby anticyclonic (cyclonic) acting as stronger (weaker) CO₂ sinks in summer, but stronger (weaker) sources in winter.

Despite the abundance of mesoscale eddies, previous studies generally investigate singular eddies (Chen et al., 2007; Jones et al., 2017; Pezzi et al., 2021) or a regional subset of eddies (Ford et al., 2023; Orselli et al., 2019b; Song et al., 2016) and their effect on the air-sea CO₂ flux. Thus, the global cumulative effect of all types of eddies on the air-sea CO₂ flux is still under investigation. Ford et al. (2023), used a Lagrangian tracking approach and suggested that long-lived (lifetimes greater than one year) mesoscale eddies enhanced the air-sea CO₂ flux in the South Atlantic Ocean by ~0.05 Tg C yr⁻¹ (~0.08%). Guo and Timmermans (2024) used a spatial and timeseries decomposition to extract the mesoscale flow impact on the air-sea CO₂ fluxes globally, and estimate a small integrated effect of 0.72 Tg C yr⁻¹ (compared to global ocean uptake of ~2.9 Pg C yr⁻¹). However, this result may include mesoscale signals not related to mesoscale eddies (Guo and Timmermans, 2024).



Li et al. (2025), using a method that tracked individual eddies similar to Ford et al. (2023), showed that mesoscale eddies within the Kuroshio and Gulf Stream western boundary currents could enhance the CO₂ sink by 28.34 ± 9.41 Tg C yr⁻¹.

60

In this paper we produce a global dataset of long lived (defined as lifetimes greater than one year) mesoscale eddies ($N = 5996$; radii > 30 km) and their associated air-sea CO₂ fluxes tracked in a Lagrangian mode between 1993 and 2022. The methodology refines the approach described in Ford et al. (2023), using a global neural network approach and published tools which are also used to generate one ocean carbon sink dataset submission to the annual Global Carbon Budget assessments (Friedlingstein et al., 2025). Following recommendations for global ocean carbon assessments (Shutler et al., 2024) we prioritise the use of climate quality satellite data records (Embury et al., 2024; Sathyendranath et al., 2019) within the analysis. The uncertainties on the air-sea CO₂ fluxes are systematically assessed following the work of Ford et al. (2024a). These refinements provide a robust foundation to studying the modulation of air-sea CO₂ flux by mesoscale eddies, with an uncertainty budget. We demonstrate the use of the global dataset to assess regional and global air-sea CO₂ fluxes of long-lived eddies and to estimate their net impact on CO₂ uptake of the ocean.

70

2. Methods

2.1 Satellite and reanalysis data

The importance of prioritising the use of climate data records to study long time series and the ocean carbon sink was highlighted in Shutler et al. (2024). We used the European Space Agency's climate change initiative (CCI) climate data records SST-CCI (v3; ~ 4 km; 1993 to 2022) for SST (Embury et al., 2024; Good and Embury, 2024) and the Ocean Colour CCI (OC-CCI) for the chlorophyll-a (chl-a) concentrations (v6; ~ 4 km; 1997 to 2022; Sathyendranath et al., 2019, 2023), with their respective per observation uncertainties. The CCI-SST was bias corrected for a cool bias with respect to global SST drifters, representative of SST at 20 cm (~ 0.05 K; Embury, 2023; Embury et al., 2024), which is used to provide an accurate estimation of fCO₂(_{sw}) (in section 2.3), and for the air-sea CO₂ flux calculation (in section 2.4).

We were unable to use the sea surface salinity (SSS) CCI climate data record for our application due to the 8 day temporal resolution of these data. We therefore used the Copernicus Marine Service GLORYS12V1 ocean reanalysis product for SSS (~ 9 km; 1993 to 2022; CMEMS, 2021; Jean-Michel et al., 2021), and the ocean mixed layer depth (MLD) as no climate data record is available for MLD. No climate data record is available for wind speed, therefore the Cross-Calibrated Multi-Platform (CCMP) wind speed dataset (6-hourly; ~ 25 km; 1993 to 2022) was chosen (Mears et al., 2022; Remote Sensing Systems et al., 2022) which is often used for ocean carbon assessments (Ford et al., 2024a).

85



2.2 Eddy Trajectories Atlas

The satellite altimetry based Mesoscale Eddy Product (version META3.2) as described in Pegliasco et al. (2022b, a), and distributed by the Archiving, Validation, and Interpretation of Oceanographic Satellite data (AVISO), was used to identify the trajectories of mesoscale eddies between 1993 and 2022. We extracted the eddy trajectories globally, that had a lifetime greater than one year, which gave 3244 anticyclonic eddies and 2752 cyclonic eddies for further analysis. For each eddy trajectory, a daily position was provided along with a polygon shape that estimates the eddy shape and size from the altimetry-based data. These eddy polygons were used to extract a daily timeseries of the environmental data described in Section 2.1, where the daily conditions within the eddy were calculated (mean, median, standard deviation, interquartile range, maximum number of available data points, number of valid data points). This was repeated for the area surrounding the eddy, where we consider the ‘area outside’ to be a circle centred on the eddy but with three times the mean radius of the eddy and the area inside the eddy polygon itself removed. The chosen radii (of three times the mean radius) was used as Ford et al. (2023) showed that the results of their study were consistent when using a ‘surrounding area criterion’ between two and five radii.

Daily timeseries of conditions within and surrounding the eddy, were then converted to a monthly median timeseries using the daily median values. The daily median was chosen to reduce the impact of any potential outliers caused by any limited data coverage due to cloud cover in the chlorophyll-a record. The daily median and mean were generally consistent for the SST, SSS, MLD and wind speed fields as these are spatially complete fields.

2.3 $f\text{CO}_2$ (sw) neural network (UEXP-FNN-U) and uncertainty

The monthly $f\text{CO}_2$ (sw) and air-sea gas fluxes were estimated using the methods and tools of the University of Exeter Physics Feed Forward neural network with uncertainties (UEXP-FNN-U) which are routinely used to generate ocean sink data for the annual Global Carbon Budget assessments (Friedlingstein et al., 2025), and described in Ford et al. (2024a). The UEXP-FNN-U methodology estimates the $f\text{CO}_2$ (sw) based on in situ data that is considered representative of the subskin layer (~0.2 m water depth), which allows for an accurate air sea CO_2 flux calculation (Woolf et al., 2016; Section 2.4). The methods used are consistent with those in Ford et al (2024a), so only a summary of the method is provided here. The UEXP-FNN-U was trained with in situ $f\text{CO}_2$ (sw) from the Surface Ocean CO_2 Atlas (SOCAT; Bakker et al., 2016) that have been recalculated to a consistent temperature and depth dataset (Ford et al., 2024d), along with predictor variables of SST, SSS, MLD and the atmospheric dry mixing ratio of CO_2 ($x\text{CO}_2$ (atm)), and anomalies of each with respect to a long term monthly climatology.

For each eddy the monthly median timeseries of the SST, SSS, MLD were provided to the UEXP-FNN-U to estimate the $f\text{CO}_2$ (sw). The $x\text{CO}_2$ (atm) was calculated from the National Oceanic and Atmospheric Administration Global Monitoring Laboratory (NOAA-GML) monthly 1 degree fields (Lan et al., 2023) that were used within the neural network training



where a weighted mean of the nearest four pixels taken at the mean (centre) position of each eddy in the month was used. Anomalies in SST, SSS, MLD and $x\text{CO}_2(\text{atm})$ were calculated with respect to a 1 degree monthly climatology.

The uncertainties in the $\text{fCO}_2(\text{sw})$ were calculated as described in Ford et al. (2024a). The $\text{fCO}_2(\text{sw})$ uncertainty has three components: (1) the network uncertainty estimated as the two standard deviation of the 10 neural network ensemble, (2) the parameter uncertainty was the propagated input parameter uncertainties and was estimated using a lookup table and (3) the evaluation uncertainty which was the evaluation with respect to the SOCAT observations (Bakker et al., 2016). All three components are combined in quadrature, assuming they are independent and uncorrelated (Taylor, 1997), to provide a total uncertainty (considered 95% confidence). The uncertainty components were calculated for each $\text{fCO}_2(\text{sw})$ estimate.

Additionally, a second version of the neural network was run. This version included chl-a (and the chl-a anomaly) as a predictor and was used to produce a second estimate of $\text{fCO}_2(\text{sw})$. Ford et al. (2022a) highlighted that the inclusion of more representative biological parameters improved the regional estimation of $\text{fCO}_2(\text{sw})$ in the South Atlantic Ocean. Therefore, this additional neural network output was generated using the same software used to create the UExp-FNN-U estimate of $\text{fCO}_2(\text{sw})$ (Ford et al., 2024c) just with the added chl-a predictor. However, we note the limitation of this second $\text{fCO}_2(\text{sw})$ estimate that uses chl-a. This dependency on optically derived remote sensing data (ie the chl-a data) means that it was limited to producing estimates after October 1997 (as routine ocean colour observations are not available before this date) and it could not provide estimates during polar winter due to missing daily chl-a data (as the low light levels inhibit optical retrievals).

The neural network estimated $\text{fCO}_2(\text{sw})$ were compared to recalculated SOCAT observations (Ford et al., 2024d; Goddijn-Murphy et al., 2015) within eddies to assess the accuracy and precision of the estimates. The individual cruise SOCAT observations are gridded (to monthly 1 degree) to provide the training and independent test data to the UExp-FNN-U, and therefore these $\text{fCO}_2(\text{sw})$ observations are not strictly independent. For each eddy trajectory, the SOCAT observations were collocated with the daily eddy polygon. The daily SOCAT observations that fell within the eddy were then aggregated into monthly mean $\text{fCO}_2(\text{sw})$, which could be compared to the neural network monthly $\text{fCO}_2(\text{sw})$. We calculated a series of statistics including the bias, root mean square difference (RMSD), slope and intercept of a Type II linear regression to characterise the differences between the neural network outputs and monthly mean SOCAT $\text{fCO}_2(\text{sw})$. As in Ford et al. (2021) weighted variants of these statistics were also calculated to capture the uncertainties in both sets of data (neural network output and the SOCAT in situ data), assuming a SOCAT $\text{fCO}_2(\text{sw})$ uncertainty of $5 \mu\text{atm}$ (Bakker et al., 2016) and the calculated total $\text{fCO}_2(\text{sw})$ uncertainty.



2.4 Air-sea CO₂ flux calculations and uncertainties

The CO₂ flux calculations were performed using FluxEngine v4.0.9.1 (Holding et al., 2019; Shutler et al., 2016), using the “rapid” transport approximation (Woolf et al., 2016), at monthly time steps. The evidence continues to grow supporting the calculation of air-sea CO₂ fluxes with consideration of the vertical temperature gradients, which is supported by theoretical (Woolf et al., 2016), observation based (Dong et al., 2022b; Shutler et al., 2020; Watson et al., 2020), modelling (Bellenger et al., 2023), and recently two in situ studies (Dong et al., 2024; Ford et al., 2024b). Therefore, the air-sea CO₂ fluxes were calculated using a bulk formulation that allows for the vertical temperature gradients to be captured. The calculations are consistent with the methods used to create the UExp-FNN-U dataset that is submitted to the annual Global Carbon Budget assessments (Friedlingstein et al., 2025), except here a simplified approach to determine the skin SST value is used.

The air sea CO₂ flux (F) was calculated as:

$$F = k_{600} (S_c/600)^{-0.5} (\alpha_{\text{subskin}} f\text{CO}_2(\text{sw,subskin}) - \alpha_{\text{skin}} f\text{CO}_2(\text{atm})) \quad (1)$$

Where k is the gas transfer velocity estimated from the monthly wind speeds and the Nightingale et al. (2000) gas transfer parameterisation. α_{subskin} and α_{skin} are the solubility of CO₂ at the base, and top of the mass boundary layer respectively, and were calculated as a function of SST and SSS (Weiss, 1974). α_{subskin} was calculated from the bias corrected CCI-SST SST and the CMEMS SSS. α_{skin} was calculated with the same datasets, but with a fixed cool (-0.17K) (Donlon et al., 1999) and salty (+0.1 psu) skin effect. We used a fixed cool skin here, instead of the dynamic cool skin approach (that uses COARE 3.5; Fairall et al., 1996) as used within the UExp-FNN-U Global Carbon Budget submission due to the computation overhead needed to extract the additional environmental fields required for the calculations. This simplified approach has only a small effect on the global scale (Dong et al., 2022b), and therefore we do not see it as a limitation. $f\text{CO}_2(\text{atm})$ was estimated for the NOAA-GML xCO₂(atm), ERA5 sea level pressure (Hersbach et al., 2019) and the CCI-SST with a cool salty skin following Dickson et al. (2007). $f\text{CO}_2(\text{sw,subskin})$ was provided by the neural network $f\text{CO}_2(\text{sw})$. The ERA5 sea level pressure was retrieved from monthly 0.25 deg fields, using a weighted mean of the 4 closest observations to the mean monthly eddy position. None of the eddies considered were under sea ice (as the eddy detection data and algorithm cannot track in areas of ice), and therefore the term “1 – ice” which is generally included within Eq. 1 (to linearly scale the gas fluxes with sea ice concentration) has not been included.

The air-sea CO₂ flux uncertainties were calculated following the methods in Ford et al. (2024a), and consistent literature values for the uncertainties in the wind speed (1.9 ms⁻¹; 95% confidence; Mears et al., 2022a), salinity (0.2 psu; 95% confidence; Jean-Michel et al., 2021), xCO₂(atm) (0.4 µatm; 95% confidence; Lan et al., 2023) and gas transfer parameterisation (20%; 95% confidence; Woolf et al., 2019). The SST uncertainty was extracted from the daily CCI-SST dataset and were converted to monthly uncertainties assuming a five day temporal correlation (Ford et al., 2024a). The uncertainties were calculated at the 95% confidence (or the 2 sigma).



The monthly mean daily flux of CO_2 ($\text{g C m}^{-2} \text{d}^{-1}$) was multiplied by the number of days and the mean area of the eddy as provided by the eddy trajectories, in the respective month. The fluxes (Tg C mon^{-1}) were then added cumulatively to retrieve the net cumulative CO_2 flux for each eddy (Tg C). Collating the combined uncertainties requires careful consideration of their temporal correlations. Some uncertainties will be temporally decorrelated, and others have temporal correlations. We used the assumptions made in Ford et al. (2024a), that the SST, SSS, wind speed, $x\text{CO}_2(\text{atm})$ and $f\text{CO}_2(\text{sw})$, and components dependent on these uncertainties, are temporally uncorrelated and are therefore propagated assuming they are independent (Taylor, 1997). Whereas, the remaining uncertainties that stem from the Schmidt number, solubilities and gas transfer parameterisation algorithm uncertainties are assumed temporally correlated and therefore are summed (Ford et al., 2024a).

The air-sea CO_2 flux calculations and uncertainty estimates were computed for the two variants of $f\text{CO}_2(\text{sw})$. The computations were also applied separately for the eddy and the area outside the eddy, assuming the same area coverage of the eddy for both calculations (i.e allowing the cumulative fluxes to be compared for the same area coverage).

2.5 Modification of air-sea CO_2 fluxes due to the existence of the eddy

As shown in Ford et al. (2023), the air-sea CO_2 flux into an eddy can be considered as two components: (1) the flux that would occur without the presence of the eddy and (2) the mesoscale modification of the flux through both oceanic and atmospheric effects of the eddy presence. The flux that would occur without the eddy being present can be estimated using the conditions that are driving the air-sea CO_2 flux in the environment surrounding the eddy. This reference flux can be removed from the air-sea CO_2 flux calculated for within the eddy to indicate the mesoscale modification of the flux due to the existence of the eddy, which was converted to a percentage change with respect to the surrounding environment CO_2 flux, following Ford et al. (2023).


The eddy modification of the air-sea CO_2 flux was calculated for each individual eddy, and then the median percentage modification was estimated for global and regional subsets, due to the lower sensitivity to outliers. We repeat the percentage change calculations in a Monte Carlo uncertainty propagation approach to evaluate the full extent of the uncertainties, whereby the eddy modification flux was perturbed within their uncertainties (95%) 1000 times independently (i.e., assuming the individual eddy flux modification uncertainties are uncorrelated). The two standard deviation value of the resulting ensemble was taken as the 95% confidence on the median percentage change for the global or regional subsets due to the uncertainties.



3. Results

3.1 Geographical distribution of mesoscale eddy cumulative air-sea CO₂ flux

205 In total 5996 eddies were tracked and their air-sea CO₂ flux estimated, which comprised 3244 anticyclonic and 2752
cyclonic eddies between 1993 and 2022 (Figure 1). The geographical distribution of the cumulative air-sea CO₂ flux into
both eddy types generally followed the global distribution of air-sea CO₂ fluxes. The temperate regions showed eddies with
strong CO₂ sink characteristics over their lifetimes, whereas eddies in the subtropical showed weaker CO₂ sinks, or even CO₂
sources. Regionally the Indian Ocean showed stronger CO₂ sinks associated with anticyclonic eddies when compared to the
210 Atlantic and Pacific Oceans (Figure 1a). The South Pacific showed anticyclonic eddies acting as weaker CO₂ sinks compared
to the North Pacific and had more eddies acting as CO₂ sources. Notable regions where cyclonic eddies were acting as strong
CO₂ sinks are within the Indian Ocean, and Northwestern Atlantic Ocean (Figure 1b). Cyclonic eddies in the South Pacific
tended to act more as CO₂ sources than sinks (Figure 1b). The Southern Ocean showed the anticyclonic and cyclonic eddies
acting as either weak CO₂ sinks or weak CO₂ sources (Figure 1).



215

Figure 1: (a) The cumulative air-sea CO₂ flux into the anticyclonic eddies where the scatter points are plotted at the formation location of each eddy. (b) same as (a) but for cyclonic eddies. Basemap from Natural Earth v4.0.0 (<https://www.naturalearthdata.com/>).



3.2 Example eddy trajectory

Figure 2 shows an example of an eddy trajectory in the North Pacific Ocean that was selected due to the ~3 year lifetime, that highlights the seasonality and variability of the environmental data, the $f\text{CO}_2(\text{sw})$ and associated air-sea CO_2 fluxes with the uncertainties shown. Over the three years the eddy moves around a relatively small region within the subpolar region (Figure 2c). Within the eddy, an expected SST seasonal cycle was present (Figure 2a), along with an interannual variability within the SSS timeseries (Figure 2b). The estimated $f\text{CO}_2(\text{sw})$ also highlighted a clear seasonal cycle with higher $f\text{CO}_2(\text{sw})$ in the winter months, and lower $f\text{CO}_2(\text{sw})$ in the summer (Figure 2d). The eddy exhibited a period of strong CO_2 outgassing during winter, followed by a small CO_2 sink within the summer months (Figure 2e). When cumulatively summed, the air-sea CO_2 fluxes indicate that the eddy outgassed CO_2 over its lifetime, but clearly this outgassing was not year-round (Figure 2f). The example eddy illustrates the available data that could be used to evaluate the driving mechanism that are affecting the $f\text{CO}_2(\text{sw})$ and air-sea CO_2 fluxes over the eddy's lifetime.

3.3 UExp-FNN-U $f\text{CO}_2(\text{sw})$ compared to SOCAT observations within eddies

The UExp-FNN-U was trained on a global dataset of $f\text{CO}_2(\text{sw})$ and so it is important to assess its performance which provides some level of confidence that the eddy variability is being correctly captured. The accuracy and precision estimates between the SOCAT in situ observations and the UExp-FNN-U within the eddies showed good performance (Figure 3) similar to the results for the global scale in Ford et al. (2024a) (weighted bias = $-0.08 \mu\text{atm}$ and $\text{RMSD} = 20.75$). For anticyclonic eddies, we observed a larger weighted RMSD (precision) of $28.49 \mu\text{atm}$ ($N=2726$ monthly matches; Figure 3a). For cyclonic eddies we observed a lower RMSD of $24.31 \mu\text{atm}$ ($N = 1776$; Figure 3c). Both eddy types showed small weighted biases (accuracy) and therefore we consider the UExp-FNN-U generated $f\text{CO}_2(\text{sw})$ to sufficiently represent the eddy $f\text{CO}_2(\text{sw})$. Figures 3b and 3d show that the uncertainties calculated for the $f\text{CO}_2(\text{sw})$ were able to sufficiently represent the differences to the SOCAT observations. Thereby providing validity to the $f\text{CO}_2(\text{sw})$ contribution to the air-sea CO_2 flux uncertainty budgets.




Figure 2: Exemplar eddy trajectory (eddy 194465) in the North Pacific Ocean with calculated air-sea CO_2 fluxes (a) Sea surface temperature (SST) for the example eddy's lifetime. Black line is the daily SST, where dark grey and light grey shading indicates the 1 sigma (~67% confidence) and 2 sigma (~95% confidence) uncertainties. Red line is the median monthly SST. (b) same as (a) for sea surface salinity. (c) Geographical eddy trajectory, where colour indicates the age of eddy (blue is eddy formation and yellow is eddy dissipation). (d) Monthly timeseries of fugacity of CO_2 in seawater ($f\text{CO}_2(\text{sw})$; solid line) and dry mixing ratio of CO_2 in the atmosphere ($x\text{CO}_2(\text{atm})$; dashed line) for the eddy. Dark grey and light grey shading indicates the 1 sigma (~67% confidence) and 2 sigma (~95% confidence) uncertainties on the $f\text{CO}_2(\text{sw})$. Red dots indicate $f\text{CO}_2(\text{sw})$ in situ observations from the Surface Ocean CO_2 Atlas within the eddy. (e) same as (d) but for the air-sea CO_2 flux where a positive flux means CO_2 outgassing. Dashed black line indicates an air-sea CO_2 flux of 0. (f) same as (d) but for the cumulative air-sea CO_2 flux. Red line and banding indicate the cumulative air-sea CO_2 flux for the surrounding environment.




Figure 3: (a) Comparison of the UExP-FNN-U $f\text{CO}_2(\text{sw})$ to in situ SOCAT observations within anticyclonic eddies. Solid black line is the 1:1. Dashed line is the Type II linear regression. In text statistics are root mean square difference (RMSD), bias, slope and intercept of a Type II linear regression and number of matches (N). (b) same as (a) but showing the uncertainty on the $f\text{CO}_2(\text{sw})$ (2 sigma; 95% confidence) as errorbars for anticyclonic eddies. (c and d) same as (a and b) for cyclonic eddies.


3.4 Uncertainty in the mesoscale eddy cumulative air-sea CO_2 flux

Two exemplar eddies, eddy A with a lifetime of 12 months and eddy B with a lifetime of 42 months, are shown in Figure 4. These were selected to highlight the differences in the relative and absolute contributions of each uncertainty component to the total uncertainty, and how these can change over time for eddies of differing lifetimes. The absolute uncertainty magnitudes for eddy B were larger than eddy A (Figure 4b, d), but the relative contributions of each component showed similarities.



For both eddies at the end of their life, the $f\text{CO}_2(\text{sw})$ component was the dominant source to the uncertainty for the whole
265 lifetime, followed by the gas transfer parameterisation uncertainty. For eddy A, wind speed was the next largest contributor
to the uncertainties, whereas for the eddy B, the solubility component uncertainties were larger than the wind speed
uncertainty.

Throughout both eddy lifetimes the dominant uncertainty contributions changed. For eddy A, at formation showed that the
wind speed and solubility components were larger contributors than the gas transfer uncertainty until four months after
270 formation (Figure 4b). Within eddy B, the wind speed was a larger contributor than the solubility components until 12
months after formation, at which time the solubility component becomes a larger contributor (Figure 4d). Uncertainties due
to the Schmidt number and $f\text{CO}_2(\text{atm})$ terms were a small contribution to the uncertainty in both eddies.



275 **Figure 4:** (a) The total cumulative air-sea CO₂ flux uncertainty (2 sigma) for an exemplar anticyclonic eddy, eddy A, (1 year lifetime; eddy 496) split into the relative contributions for the individual components. (b) The total air-sea CO₂ flux uncertainty in absolute terms. Legend in (b) corresponds to colours in (a). (c) and (d) same as (a) and (b) but for eddy B, an anticyclonic eddy (42 months lifetime; eddy 194465). Note different x-axis limits for (a) and (b) compared to (c) and (d).



3.5 Global and regional mesoscale modifications of the air-sea CO₂ flux

280 An example application of the dataset was to assess the modification of the cumulative air-sea CO₂ flux by individual eddies. The analysis indicated that individual eddies could enhance (negative percentage changes) or suppress (positive percentage changes) the CO₂ sink. (Figure 5). Both anticyclonic (Figure 5a) and cyclonic eddies (Figure 5b) showed individual eddies that were either enhancing or suppressing the air-sea CO₂ flux. Regional signatures in the air-sea CO₂ flux modification were apparent, for example anticyclonic eddies in the South Pacific and Southern Ocean had a greater tendency to enhance the

285 CO₂ sink, whereas in the Indian Ocean there was not a discernible tendency. Cyclonic eddies in the Southern Ocean indicated a larger suppression of the CO₂ sink than for example the North Pacific.

Considering all the eddies studied and the calculated uncertainties, anticyclonic eddies were identified to enhance the cumulative CO₂ flux, where these eddies acted as stronger CO₂ sink (weaker CO₂ source) by 4.5 ± 2.8 % (95 % confidence). Cyclonic eddies indicated a slight suppression of the cumulative air-sea CO₂ flux by 0.7 ± 2.6 %, acting overall to weaken

290 the CO₂ sinks (or as stronger CO₂ sources). Here we note, at the 95 % confidence the cumulative CO₂ sink enhancement by anticyclonic eddies was significantly difference from 0 (i.e the confidence interval did not include 0) when uncertainties were accounted for, but this was not significantly different from 0 for cyclonic eddies.

The regional differences can be emphasised by considering median eddy modifications within different regional subsets (Figure 6) instead of globally (Figure 5c, d). The eddy modification of CO₂ fluxes within the regions showed differing

295 magnitudes that fall within different significance bands when the uncertainties are accounted for. For example, the Southern Ocean shows an anticyclonic enhancement of the CO₂ sink of 6.2 ± 5.3 % (significant at 95 % confidence), with cyclonic eddies suppressing the CO₂ sink by 2.5 ± 4.6 %. In the North Pacific, we find similar results where anticyclonic eddies enhance by 5.6 ± 5.2 %, and cyclonic eddies suppress by 1.7 ± 7.1 %. Consistent results were found for the South Pacific but noting the cyclonic eddies showed a larger uncertainty interval of 11.4 %. Interestingly, the South Atlantic Ocean shows the

300 opposite eddy modifications to much of the global ocean where anticyclonic suppressed the sink by 1.8 ± 15.6 % and cyclonic eddies appear to enhance the CO₂ sink by 0.1 ± 14.3 %. The uncertainty intervals on these are however the largest of any region, likely due to the lowest number of eddies considered.




Figure 5: (a) Geographical distribution of the anticyclonic eddies' modification of the air-sea CO₂ flux. Negative values indicate a stronger CO₂ sink (weaker CO₂ source), and positive values indicate a weaker CO₂ sink (stronger CO₂ source). (b) same as (a) for the cyclonic eddies. (c) Box plot showing the anticyclonic eddy modification of the air-sea CO₂ flux. Red line indicates the median, box indicates the 25th and 75th quartiles, whiskers extend from the 25th and 75th quartiles by 1.5 interquartile ranges. Circles indicate data considered outliers. Dark red shading indicates the 1 sigma (~68% confidence) uncertainty on the median by propagating the air-sea CO₂ flux uncertainties using a Monte Carlo uncertainty propagation. Light red shading indicates the 2 sigma uncertainty on the median (~95% confidence). X-axis label shows number of eddies (N), the median modification with the 2 sigma uncertainty. (d) same as (c) but for the cyclonic eddies. Basemap in (a) and (b) from Natural Earth v4.0.0 (<https://www.naturalearthdata.com/>).




Figure 6: (a) Ocean basins considered for further analysis, with a colour for each region. Regions follow the RECCAP2 ocean basin definition, but each basin was split at the Equator into North and South. North Indian Ocean was removed due to low number of eddies analysed. (b) Box plot showing the eddy modification of the air-sea CO₂ flux for the region shown with the arrow. Red line indicates the median, box indicates the 25th and 75th quartiles, whiskers extend from the 25th and 75th quartiles by 1.5 interquartile ranges. Circles indicate data considered outliers (greater than 1.5 interquartile ranges outside the 25th and 75th percentile). Dark red shading indicates the 1 sigma (~68% confidence) uncertainty on the median by propagating the air-sea CO₂ flux uncertainties using a Monte Carlo uncertainty propagation. Light red shading indicates the 2 sigma uncertainty on the median (~95% confidence). X-axis label shows number of eddy (N), the median modification with the 2 sigma uncertainty. (c), (d), (e), (f), (g) same as (b) for their respective regions identified by the arrow. Basemap in (a) from Natural Earth v4.0.0 (<https://www.naturalearthdata.com/>).

4. Discussion

4.1 Mesoscale eddy air-sea CO₂ fluxes and uncertainties

The mesoscale eddy air-sea CO₂ fluxes provide both the CO₂ fluxes for each month with uncertainties and the corresponding environmental data (i.e SST, SSS) within and outside of each eddy (Figure 2). These data allow a range of analyses to be conducted, for example, in this study, we show how the mesoscale modification of the air-sea CO₂ flux can be determined from these data (Figure 5; Figure 6). Other potential applications could include, analysing the thermal and non-thermal components in driving the global eddy modified air-sea CO₂ fluxes (as illustrated by Ford et al. (2023) for the South Atlantic), or for investigating nutrient entrainment within the eddies and how it links to biological variations within the eddy track, or the variability in phytoplankton biomass and / or productivity within the eddies which are important for improving our understanding of carbon rate dynamics, and their impacts on ecology and biodiversity. The dataset presented here



therefore provides the basis for a wide range of studies to assess the evolution of mesoscale eddies and their air-sea CO₂ fluxes alongside understanding the linkages with their localised environmental conditions.

The dataset air-sea CO₂ flux estimates are accompanied by a comprehensive uncertainty budget developed by Ford et al. (2024a) (Figure 2; Figure 3). This is the first dataset of eddy air-sea CO₂ fluxes to include a uncertainty budget that has been built on the principles where all known sources of uncertainty are systematically considered (however small) and propagated to the final uncertainty using standard propagation techniques and a well-established uncertainty framework (BIPM, 2008; Taylor, 1997). The budget therefore provides an uncertainty on each air-sea CO₂ flux estimate, and the fCO₂ (sw), which can be accounted for within further analyses (e.g. as used in Ford et al., 2021, 2022b) and aids in assigning confidence to any results, as demonstrated in the example results that have been presented.

The comparisons to the SOCAT fCO₂ (sw) observations within eddies provide further confidence in the retrieved UExP-FNN-U fCO₂ (sw) and resulting air-sea CO₂ fluxes. We showed that for both the anticyclonic and cyclonic eddies the accuracy (bias) and precision (RMSD) were similar to the global scale performance of these approaches (Ford et al., 2024a). This result was consistent with Ford et al. (2023) for the South Atlantic Ocean, who showed that both eddy types were well represented by the neural network approach (except Ford et al. (2023) determined this from a lower number of crossover data points than presented here). Li et al. (2025) also showed for their neural network approach, similar accuracy and precision results for the fCO₂ (sw) within eddies for four western boundary current regions. Although we did observe a slightly lower precision (higher RMSD), likely due to eddy related processes that are not fully captured within the neural network approach. These results also provide validity to the calculated fCO₂ (sw) uncertainties, which in the majority of cases are dominated by the fCO₂ (sw) evaluation uncertainty component. As the retrieved eddy fCO₂ (sw) bias and RMSD was similar to the global performance (given in Ford et al., 2024), we are confident in the UExP-FNN-U fCO₂ (sw) within the eddies.

Within the UEx-L-Eddies we provide a secondary fCO₂ (sw) estimate (and associated air-sea CO₂ fluxes) from a global fCO₂ (sw) neural network, which included chl-a as a predictor. We include the additional neural network because Ford et al. (2022a) highlighted that the inclusion of more representative biological parameters improved the regional estimation of fCO₂ (sw) in the South Atlantic Ocean, which is likely to be the same for other regions. Previous studies have shown the importance of biological modulation of fCO₂ (sw) within eddies (Orselli et al., 2019b), the resulting CO₂ fluxes, and how the importance changes over the eddy lifetime (Ford et al., 2023). This additional neural network showed similar but slightly improved precision (lower weighted RMSD) when compared to the in situ SOCAT observations, although to a lower number of data points (Figure S1; anticyclonic bias = -0.17 uatm, RMSD = 27.36 µatm; cyclonic bias = 0.74 µatm, RMSD = 23.57 µatm). These estimates are however restricted to regions between 50 °N and 50 °S due to the availability of ocean colour chl-a data in polar winter (i.e for a full eddy timeseries the eddy must remain within the available ocean colour data). We therefore provide the secondary neural network to further aid in understanding the processes that are driving mesoscale eddy modification of the air-sea CO₂ fluxes.



Previous eddy trajectory datasets have been produced, for example Dong et al. (2022a), which include environmental datasets (e.g. SST) that can be used to understand the effects of eddies on physical and biological properties. The UEx-L-Eddies however extends the principles of these datasets to include air-sea CO₂ fluxes but also has a focus on climate quality dataset (i.e the ESA CCI datasets) and provides comprehensive uncertainties. Therefore it provides a robust dataset for understanding long-lived eddy effects on the surface properties and air-sea CO₂ fluxes. In the future, we plan to include in situ observations by Biogeochemical Argo floats (BGC-Argo; Roemmich et al., 2019), which could be used to provide in situ based fCO₂ (sw) and air-sea CO₂ fluxes to further verify the air-sea CO₂ fluxes (e.g., as suggested by Keppler et al. (2024)).

4.2 Comparison to previous global and regional eddy modifications of the air-sea CO₂ fluxes

Previous studies have investigated the effect of mesoscale eddies on global and regional air-sea CO₂ fluxes. Guo and Timmermans (2024) evaluate the cumulative effect of mesoscale variability on the air-sea CO₂ flux globally, which they find enhances the global air-sea CO₂ flux by 0.72 Mt C yr⁻¹, or 0.72 Tg C yr⁻¹. With the UEx-L-Eddies, if the individual eddy air-sea CO₂ flux modifications are summed for the whole dataset, we find a global cumulative enhancement of the ocean CO₂ sink by long-lived mesoscale eddies of 75 ± 33 Tg C between 1993 and 2022. This would be equivalent to 2.7 ± 1.1 Tg C yr⁻¹ (95 % confidence interval). The calculated uncertainties with the UEx-L-Eddies allows robust uncertainty estimates to be provided alongside further analyses of the individual eddies, allowing significance of comparisons to be assessed. Differences here may be due to Guo and Timmermans (2024) including mesoscale variability not associated with mesoscale eddies (such as filaments, and current meanders), as their method does not track individual eddies. It could also be due to the UEx-L-Eddies only covering long-lived eddies, that represent 0.4 % of eddies within the META3.2 trajectories dataset and therefore misses the contribution of smaller eddies (Pegliasco et al., 2022b) that would be included with Guo and Timmermans (2024).

Li et al. (2025) showed for the Kuroshio current that anticyclonic eddies enhanced the CO₂ sink by 15 ± 1.73 %, and cyclonic eddies reduced the CO₂ sink by 5.7 ± 1.5 %. Similar results were also shown for the Gulf Stream. Both the Gulf Stream and the Kuroshio current are dominated by short-lived eddies (e.g., those that survive for less than 1 year) in comparison to the long-lived eddies studied within the UEx-L-Eddies dataset, and therefore comparing these two estimates is inappropriate. However, our regional results for the North Pacific and North Atlantic Oceans do show a consistent direction of change (i.e., an enhanced sink) but with smaller magnitudes (Figure 6).

Keppler et al. (2024) investigate the role of mesoscale eddies in modifying the air-sea CO₂ flux in the Southern Ocean using Biogeochemical Argo profilers between April 2014 to December 2022. They find anticyclonic eddies enhanced the air-sea CO₂ sink by 7 ± 2 % and cyclonic eddies reduced the air-sea CO₂ flux by 2 ± 2 % (1 sigma uncertainties). Within the UEx-



L-Eddies, we found that anticyclonic eddies enhanced the CO₂ sink by 6.2 ± 5.3 % (2 sigma uncertainties), and cyclonic eddies reduced the sink by 2.5 ± 4.6 % between 1993 and 2022 (Figure 6g). These consistent results provide confidence to the air-sea CO₂ flux estimates within the UEx-L-Eddies.

400 Ford et al. (2023) showed that within the South Atlantic Ocean, anticyclonic (N = 36) and cyclonic (N = 31) eddies enhanced the CO₂ sink by 3.7 % and 1.7 %, respectively. In our analysis for the South Atlantic Ocean (Figure 6e) we showed that anticyclonic suppressed the CO₂ sink by 1.8 ± 15.1 (N = 207) and cyclonic eddies enhanced the CO₂ sink by 0.1 ± 14.3 % (N = 155) respectively, where confidence intervals are expressed as 95 % confidence. Within this dataset, we consider ~5 times more eddies than Ford et al. (2023) and find that the air-sea CO₂ flux uncertainties have a large effect on our resulting
405 confidence, making the results indistinguishable at 95 % confidence (even at 67 % confidence the two are indistinguishable). The comparison highlights the importance of the calculated uncertainties and their use within further analyses and comparisons with other air-sea CO₂ fluxes.

4.3 Limitations when using the UEx-L-Eddies

410 For some eddies the daily environmental data can have missing values even for complete coverage data (for example, the CCI-SST). These gaps stem from the META3.2 eddy trajectories dataset where the polygon to define the limits of the eddy does not form correctly, and therefore we were unable to extract values where the polygon was undefined.

5. Summary

The UEx-L-Eddies is a dataset of the air-sea CO₂ fluxes for (N=5996) long lived mesoscale eddies calculated in a
415 Lagrangian mode within the global ocean. We use a global fCO₂ (sw) neural network (as used within one dataset submitted to the Global Carbon Budget called UExP-FNN-U) to estimate the fCO₂ (sw) within the eddies at a monthly resolution. We prioritise the use of climate quality datasets within the analysis. The air-sea CO₂ fluxes (also calculated following the methods of UExP-FNN-U) are accompanied by a comprehensive uncertainty budget (using a published methodology), that considers all known sources of uncertainty. We show for an exemplar eddy that the seasonal cycles of the eddy fCO₂ (sw) and
420 air-sea CO₂ fluxes are captured and can be cumulatively added to assess the CO₂ uptake (or outgassing) of individual eddies. The comprehensive air-sea CO₂ flux uncertainties provide a robust basis for assessing the confidence in the eddy air-sea CO₂ flux estimates and can be propagated to further analysis. This illustrates how the importance of the different uncertainty components can change through time highlighting the shortfall of only quantifying selected contributions to the uncertainties or assuming fixed values.



425 Within the uncertainty assessment, we find that the $f\text{CO}_2(\text{sw})$ in the eddies are estimated with an accuracy (bias) of $\sim 1 \mu\text{atm}$ and a precision (RMSD) of $28.21 \mu\text{atm}$ for anticyclonic, and accuracy of $\sim 1 \mu\text{atm}$ and a precision of $24.04 \mu\text{atm}$ for cyclonic eddies. These accuracy and precision estimates provide validity to the neural network $f\text{CO}_2(\text{sw})$.

We demonstrate a use case of the UEx-L-Eddies dataset to evaluate the air-sea CO_2 flux modification, and resultant integrated net CO_2 sink, by long-lived mesoscale eddies, globally and regionally. We find that anticyclonic eddies enhance
430 the net sink by $4.5 \pm 2.8 \%$ ($N = 3244$), and cyclonic eddies suppress by $0.7 \pm 2.6 \%$ ($N = 2752$) where uncertainties are the 95% confidence interval. Regional differences in the eddy modification are observed, for example within the Southern Ocean, anticyclonic eddies enhanced the CO_2 sink by $6.2 \pm 5.3 \%$, and cyclonic eddies reduced the sink by $2.5 \pm 5.6 \%$. We demonstrate how the use case results are consistent with previous regional analyses. Our example also highlighted the importance of using the accompanying uncertainty information when comparing studies, and caution should be taken in
435 drawing conclusions from small samples or individual eddies, without considering the underlying comprehensive uncertainty budgets for the air-sea CO_2 fluxes. The data presented could now be used to understand the processes occurring within these eddies that are driving these modifications of the air-sea CO_2 fluxes, and how regionally these processes may vary.

Author Contributions

DJF, GHT, VK, KS and JDS conceived the study and the methodology. DJF performed the analysis, testing of the dataset
440 and wrote the original draft. All authors provided input to the final manuscript.

Competing Interests

The authors declare no competing interests.

Data and Code Availability

The code for the analysis is available and version controlled on Github at <https://github.com/JamieLab/pyEddyCO2>. The
445 UEx-L-Eddies dataset are available on Zenodo (<https://doi.org/10.5281/ZENODO.16355763>; Ford et al., 2025). The AVISO+ eddies trajectories data (META 3.2) was retrieved from AVISO+ (<https://doi.org/10.24400/527896/A01-2022.005.220209>; Pegliasco et al., 2022a). The CCI-SST climate record (v3.0) were retrieved from CEDA (<https://doi.org/10.5285/4A9654136A7148E39B7FEB56F8BB02D2>; Good and Embury, 2024). The OC-CCI chl-a (v6) were retrieved from CEDA (<https://doi.org/10.5285/5011D22AAE5A4671B0CBC7D05C56C4F0>; Sathyendranath et al.,
450 2023). The CMEMS GLORYS12V1 SSS and MLD were retrieved from CMEMS (<https://doi.org/10.48670/moi-00021>; CMEMS, 2021). The CCMP wind speeds (v3.1) were retrieved from Remote Sensing Systems (<https://doi.org/10.56236/rss->



uv6h30; Remote Sensing Systems et al., 2022). The $x\text{CO}_2$ (atm) were retrieved from NOAA-GML
 (https://doi.org/10.15138/DVNP-F961; Lan et al., 2023). In situ SOCAT observations that have been recalculated to a
 consistent depth and temperature dataset were retrieved from Zenodo (https://doi.org/10.5281/zenodo.15706025; Ford et al.,
 455 2024d).

Acknowledgements

DJF and JDS were supported by funding from the European Space Agency under the projects ‘Satellite-based observations
 of Carbon in the Ocean: Pools, Fluxes and Exchanges’ (SCOPE; 4000142532/23/I-DT) and ‘Ocean Carbon for Climate’
 (OC4C; 3-18399/24/I-NB). GHT and VK were supported by The Atlantic Meridional Transect is funded by the UK Natural
 460 Environment Research Council through its National Capability Long-term Single Centre Science Programme, Atlantic
 Climate and Environment Strategic Science - AtlantiS (grant number NE/Y005589/1). This study contributes to the
 international IMBeR project and is contribution number 423 of the AMT programme.

The Surface Ocean CO_2 Atlas (SOCAT) is an international effort, endorsed by the International Ocean Carbon Coordination
 465 Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS) and the Integrated Marine Biosphere Research
 (IMBeR) program, to deliver a uniformly quality-controlled surface ocean CO_2 database. The many researchers and funding
 agencies responsible for the collection of data and quality control are thanked for their contributions to SOCAT. For the
 purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted
 Manuscript version arising from this submission.

470 References

- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O’Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S.
 D., Nakaoka, S. I., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof,
 R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F.,
 Cai, W. J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet,
 475 C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M.
 P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A.,
 Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T.,
 Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson,
 K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R.,
 480 Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., Van Heuven, S. M. A.



- C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality fCO₂ data in version 3 of the Surface Ocean CO₂ Atlas (SOCAT), *Earth System Science Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Bellenger, H., Bopp, L., Ethé, C., Ho, D., Duvel, J. P., Flavoni, S., Guez, L., Kataoka, T., Perrot, X., Parc, L., and Watanabe, M.: Sensitivity of the Global Ocean Carbon Sink to the Ocean Skin in a Climate Model, *JGR Oceans*, 128, e2022JC019479, <https://doi.org/10.1029/2022JC019479>, 2023.
- BIPM: Evaluation of measurement data—Guide to the expression of uncertainty in measurement., 2008.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91, 167–216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- Chen, F., Cai, W. J., Benitez-Nelson, C., and Wang, Y.: Sea surface pCO₂-SST relationships across a cold-core cyclonic eddy: Implications for understanding regional variability and air-sea gas exchange, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL028058>, 2007.
- CMEMS: Copernicus Marine Modelling Service global ocean physics reanalysis product (GLORYS12V1), Copernicus Marine Modelling Service [dataset], <https://doi.org/10.48670/moi-00021>, 2021.
- Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to Best Practices for Ocean CO₂ measurements, PICES Special Publication, IOCCP Report No . 8, 2007.
- Dong, C., Liu, L., Nencioli, F., Bethel, B. J., Liu, Y., Xu, G., Ma, J., Ji, J., Sun, W., Shan, H., Lin, X., and Zou, B.: The near-global ocean mesoscale eddy atmospheric-oceanic-biological interaction observational dataset, *Sci Data*, 9, 436, <https://doi.org/10.1038/s41597-022-01550-9>, 2022a.
- Dong, Y., Bakker, D. C. E., Bell, T. G., Huang, B., Landschützer, P., Liss, P. S., and Yang, M.: Update on the Temperature Corrections of Global Air-Sea CO₂ Flux Estimates, *Global Biogeochemical Cycles*, 36, <https://doi.org/10.1029/2022GB007360>, 2022b.
- Dong, Y., Bakker, D. C. E., Bell, T. G., Yang, M., Landschützer, P., Hauck, J., Rödenbeck, C., Kitidis, V., Bushinsky, S. M., and Liss, P. S.: Direct observational evidence of strong CO₂ uptake in the Southern Ocean, *Sci. Adv.*, 10, eadn5781, <https://doi.org/10.1126/sciadv.adn5781>, 2024.
- Donlon, C. J., Nightingale, T. J., Sheasby, T., Turner, J., Robinson, I. S., and Emery, W. J.: Implications of the oceanic thermal skin temperature deviation at high wind speed, *Geophysical Research Letters*, 26, 2505–2508, <https://doi.org/10.1029/1999GL000547>, 1999.
- Dufois, F., Hardman-Mountford, N. J., Greenwood, J., Richardson, A. J., Feng, M., and Matear, R. J.: Anticyclonic eddies are more productive than cyclonic eddies in subtropical gyres because of winter mixing, *Science Advances*, 2, 1–7, <https://doi.org/10.1126/sciadv.1600282>, 2016.
- Embury, O.: SST CCI Product Validation and Intercomparison Report, https://climate.esa.int/documents/2369/SST_CCI_D4.1_PVIR_v2.1-signed.pdf, 2023.



- Embury, O., Merchant, C. J., Good, S. A., Rayner, N. A., Høyer, J. L., Atkinson, C., Block, T., Alerskans, E., Pearson, K. J., Worsfold, M., McCarroll, N., and Donlon, C.: Satellite-based time-series of sea-surface temperature since 1980 for climate applications, *Sci Data*, 11, 326, <https://doi.org/10.1038/s41597-024-03147-w>, 2024.
- Fairall, C. W., Bradley, E. F., Godfrey, J. S., Wick, G. A., Edson, J. B., and Young, G. S.: Cool-skin and warm-layer effects on sea surface temperature, *Journal of Geophysical Research: Oceans*, 101, 1295–1308, <https://doi.org/10.1029/95JC03190>, 1996.
- Ford, D. J., Tilstone, G. H., Shutler, J. D., Kitidis, V., Lobanova, P., Schwarz, J., Poulton, A. J., Serret, P., Lamont, T., Chuqui, M., Barlow, R., Lozano, J., Kampel, M., and Brandini, F.: Wind speed and mesoscale features drive net autotrophy in the South Atlantic Ocean, *Remote Sensing of Environment*, 260, 112435, <https://doi.org/10.1016/j.rse.2021.112435>, 2021.
- Ford, D. J., Tilstone, G. H., Shutler, J. D., and Kitidis, V.: Derivation of seawater pCO₂ from net community production identifies the South Atlantic Ocean as a CO₂ source, *Biogeosciences*, 19, 93–115, <https://doi.org/10.5194/bg-19-93-2022>, 2022a.
- Ford, D. J., Tilstone, G. H., Shutler, J. D., and Kitidis, V.: Identifying the biological control of the annual and multi-year variations in South Atlantic air–sea CO₂ flux, *Biogeosciences*, 19, 4287–4304, <https://doi.org/10.5194/bg-19-4287-2022>, 2022b.
- Ford, D. J., Tilstone, G. H., Shutler, J. D., Kitidis, V., Sheen, K. L., Dall’Olmo, G., and Orselli, I. B. M.: Mesoscale Eddies Enhance the Air-Sea CO₂ Sink in the South Atlantic Ocean, *Geophysical Research Letters*, 50, e2022GL102137, <https://doi.org/10.1029/2022GL102137>, 2023.
- Ford, D. J., Blannin, J., Watts, J., Watson, A. J., Landschützer, P., Jersild, A., and Shutler, J. D.: A Comprehensive Analysis of Air-Sea CO₂ Flux Uncertainties Constructed From Surface Ocean Data Products, *Global Biogeochemical Cycles*, 38, e2024GB008188, <https://doi.org/10.1029/2024GB008188>, 2024a.
- Ford, D. J., Shutler, J. D., Blanco-Sacristán, J., Corrigan, S., Bell, T. G., Yang, M., Kitidis, V., Nightingale, P. D., Brown, I., Wimmer, W., Woolf, D. K., Casal, T., Donlon, C., Tilstone, G. H., and Ashton, I.: Enhanced ocean CO₂ uptake due to near-surface temperature gradients, *Nature Geoscience*, <https://doi.org/10.1038/s41561-024-01570-7>, 2024b.
- Ford, D. J., Blannin, J., Watts, J., Watson, A. J., Landschützer, P., Jersild, A., and Shutler, J. D.: OceanICU Neural Network Framework with per pixel uncertainty propagation (v1.1), [software], <https://doi.org/10.5281/ZENODO.12597803>, 2024c.
- Ford, D. J., Shutler, J. D., Ashton, I., Sims, R. P., and Holding, T.: Reanalysed (depth and temperature consistent) surface ocean CO₂ atlas (SOCAT) version 2024 (v2) (v2), [dataset], <https://doi.org/10.5281/zenodo.15706025>, 2024d.
- Ford, D. J., Shutler, J. D., Sheen, K. L., Tilstone, G. H., and Kitidis, V.: UEx-L-Eddies: decadal and global long-lived mesoscale eddy trajectories with coincident air-sea CO₂ fluxes and biogeochemical conditions (v0-2), [dataset], <https://doi.org/10.5281/ZENODO.16355763>, 2025.



- Frenger, I., Gruber, N., Knutti, R., and Münnich, M.: Imprint of Southern Ocean eddies on winds, clouds and rainfall, *Nature*
 545 *Geoscience*, 6, 608–612, <https://doi.org/10.1038/ngeo1863>, 2013.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H., Luijkx, I.
 T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B.,
 Alin, S. R., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig, H. C., Bopp, L., Cadule, P.,
 Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan, T., Decayeux, J., Djeutchouang, L.
 550 M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A., Ford, D. J., Foster, A., Gasser, T., Gehlen, M.,
 Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina,
 T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Kato, E., Keeling, R. F., Klein Goldewijk, K.,
 Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S. K., Lefèvre, N., Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G.,
 Mayot, N., McGuire, P. C., Metzl, N., Monacchi, N. M., Morgan, E. J., Nakaoka, S.-I., Neill, C., Niwa, Y., Nützel, T.,
 555 Olivier, L., Ono, T., Palmer, P. I., Pierrot, D., Qin, Z., Resplandy, L., Roobaert, A., Rosan, T. M., Rödenbeck, C.,
 Schwinger, J., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Steinhoff, T., et al.: Global Carbon Budget 2024, *Earth*
Syst. Sci. Data, 17, 965–1039, <https://doi.org/10.5194/essd-17-965-2025>, 2025.
- Goddijn-Murphy, L. M., Woolf, D. K., Land, P. E., Shutler, J. D., and Donlon, C.: The OceanFlux Greenhouse Gases
 methodology for deriving a sea surface climatology of CO₂ fugacity in support of air-sea gas flux studies, *Ocean Science*, 11,
 560 519–541, <https://doi.org/10.5194/os-11-519-2015>, 2015.
- Good, S. A. and Embury, O.: ESA Sea Surface Temperature Climate Change Initiative (SST_cci): Level 4 Analysis product,
 version 3.0, [dataset], <https://doi.org/10.5285/4A9654136A7148E39B7FEB56F8BB02D2>, 2024.
- Guo, Y. and Timmermans, M.: The Role of Ocean Mesoscale Variability in Air-Sea CO₂ Exchange: A Global Perspective,
Geophysical Research Letters, 51, e2024GL108373, <https://doi.org/10.1029/2024GL108373>, 2024.
- 565 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum,
 I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 monthly averaged data on single levels from
 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [dataset],
<https://doi.org/10.24381/cds.fl7050d7>, 2019.
- Holding, T., Ashton, I. G., Shutler, J. D., Land, P. E., Nightingale, P. D., Rees, A. P., Brown, I., Piolle, J.-F., Kock, A.,
 570 Bange, H. W., Woolf, D. K., Goddijn-Murphy, L., Pereira, R., Paul, F., Girard-Ardhuin, F., Chapron, B., Rehder, G.,
 Ardhuin, F., and Donlon, C. J.: The FluxEngine air–sea gas flux toolbox: simplified interface and extensions for in situ
 analyses and multiple sparingly soluble gases, *Ocean Science*, 15, 1707–1728, <https://doi.org/10.5194/os-15-1707-2019>,
 2019.
- Jean-Michel, L., Eric, G., Romain, B.-B., Gilles, G., Angélique, M., Marie, D., Clément, B., Mathieu, H., Olivier, L. G.,
 575 Charly, R., Tony, C., Charles-Emmanuel, T., Florent, G., Giovanni, R., Mounir, B., Yann, D., and Pierre-Yves, L. T.: The



- Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis, *Frontiers in Earth Science*, 9, 1–27, <https://doi.org/10.3389/feart.2021.698876>, 2021.
- Jones, E. M., Hoppema, M., Strass, V., Hauck, J., Salt, L., Ossebaar, S., Klaas, C., van Heuven, S. M. A. C., Wolf-Gladrow, D., Stöven, T., and de Baar, H. J. W.: Mesoscale features create hotspots of carbon uptake in the Antarctic Circumpolar Current, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 138, 39–51, <https://doi.org/10.1016/j.dsr2.2015.10.006>, 2017.
- 580 Keppler, L., Eddebbbar, Y. A., Gille, S. T., Guisewhite, N., Mazloff, M. R., Tamsitt, V., Verdy, A., and Talley, L. D.: Effects of Mesoscale Eddies on Southern Ocean Biogeochemistry, *AGU Advances*, 5, e2024AV001355, <https://doi.org/10.1029/2024AV001355>, 2024.
- 585 Lan, X., Tans, P., Thoning, K., and NOAA Global Monitoring Laboratory: NOAA Greenhouse Gas Marine Boundary Layer Reference - CO₂, [dataset], <https://doi.org/10.15138/DVNP-F961>, 2023.
- Laxenaire, R., Speich, S., and Stegner, A.: Evolution of the Thermohaline Structure of One Agulhas Ring Reconstructed from Satellite Altimetry and Argo Floats, *Journal of Geophysical Research: Oceans*, 124, 8969–9003, <https://doi.org/10.1029/2018JC014426>, 2019.
- 590 Li, X., Gan, B., Zhang, Z., Cao, Z., Qiu, B., Chen, Z., and Wu, L.: Oceanic uptake of CO₂ enhanced by mesoscale eddies, *Sci. Adv.*, 11, eadt4195, <https://doi.org/10.1126/sciadv.adt4195>, 2025.
- Mears, C., Lee, T., Ricciardulli, L., Wang, X., and Wentz, F.: Improving the Accuracy of the Cross-Calibrated Multi-Platform (CCMP) Ocean Vector Winds, *Remote Sensing*, 14, 4230, <https://doi.org/10.3390/rs14174230>, 2022.
- Nencioli, F., Dall’Olmo, G., and Quartly, G. D.: Agulhas Ring Transport Efficiency From Combined Satellite Altimetry and Argo Profiles, *Journal of Geophysical Research: Oceans*, 123, 5874–5888, <https://doi.org/10.1029/2018JC013909>, 2018.
- 595 Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochemical Cycles*, 14, 373–387, <https://doi.org/10.1029/1999GB900091>, 2000.
- Orselli, I. B. M., Goyet, C., Kerr, R., de Azevedo, J. L. L., Araujo, M., Galdino, F., Touratier, F., and Garcia, C. A. E.: The effect of Agulhas eddies on absorption and transport of anthropogenic carbon in the South Atlantic Ocean, *Climate*, 7, 1–25, <https://doi.org/10.3390/CL17060084>, 2019a.
- 600 Orselli, I. B. M., Kerr, R., Azevedo, J. L. L. de, Galdino, F., Araujo, M., and Garcia, C. A. E.: The sea-air CO₂ net fluxes in the South Atlantic Ocean and the role played by Agulhas eddies, *Progress in Oceanography*, 170, 40–52, <https://doi.org/10.1016/j.pocean.2018.10.006>, 2019b.
- 605 Pegliasco, C., Busche, C., and Faugere, Y.: Mesoscale Eddy Trajectory Atlas META3.2 Delayed-Time all satellites: version META3.2 DT allsat (3.2 DT allsat 1993-01-01/2022-02-09), [dataset], <https://doi.org/10.24400/527896/A01-2022.005.220209>, 2022a.



- Pegliasco, C., Delepoulle, A., Mason, E., Morrow, R., Faugère, Y., and Dibarboure, G.: META3.1exp: a new global mesoscale eddy trajectory atlas derived from altimetry, *Earth System Science Data*, 14, 1087–1107,
 610 <https://doi.org/10.5194/essd-14-1087-2022>, 2022b.
- Pezzi, L. P., de Souza, R. B., Santini, M. F., Miller, A. J., Carvalho, J. T., Parise, C. K., Quadro, M. F., Rosa, E. B., Justino, F., Sutil, U. A., Cabrera, M. J., Babanin, A. V., Voermans, J., Nascimento, E. L., Alves, R. C. M., Munchow, G. B., and Rubert, J.: Oceanic eddy-induced modifications to air–sea heat and CO₂ fluxes in the Brazil-Malvinas Confluence, *Scientific Reports*, 11, 10648, <https://doi.org/10.1038/s41598-021-89985-9>, 2021.
- 615 Remote Sensing Systems, Mears, C., Lee, T., Ricciardulli, L., Wang, X., and Wentz, F.: RSS Cross-Calibrated Multi-Platform (CCMP) 6-hourly ocean vector wind analysis on 0.25 deg grid, Version 3.0, [dataset], <https://doi.org/10.56236/rss-uv6h30>, 2022.
- Roemmich, D., Alford, M. H., Claustre, H., Johnson, K. S., King, B., Moum, J., Oke, P. R., Owens, W. B., Pouliquen, S., Purkey, S., Scanderbeg, M., Suga, T., Wijffels, S. E., Zilberman, N., Bakker, D., Baringer, M. O., Belbeoch, M., Bittig, H.
 620 C., Boss, E., Calil, P., Carse, F., Carval, T., Chai, F., Conchubhair, D. O., D’Ortenzio, F., Dall’Olmo, G., Desbruyères, D., Fennel, K., Fer, I., Ferrari, R., Forget, G., Freeland, H., Fujiki, T., Gehlen, M., Greenan, B., Hallberg, R., Hibiya, T., Hosoda, S., Jayne, S., Jochum, M., Johnson, G. C., Kang, K. R., Kolodziejczyk, N., Koertzing, A., Le Traon, P. Y., Lenn, Y. D., Maze, G., Mork, K. A., Morris, T., Nagai, T., Nash, J., Garabato, A. N., Olsen, A., Pattabhi, R. R., Prakash, S., Riser, S., Schmechtig, C., Shroyer, E., Sterl, A., Sutton, P., Talley, L., Tanhua, T., Thierry, V., Thomalla, S., Toole, J., Troisi, A.,
 625 Trull, T., Turton, J. D., Velez-Belchi, P. J., Walczowski, W., Wang, H., Wanninkhof, R., Waterhouse, A., Watson, A., Wilson, C., Wong, A. P., Xu, J., and Yasuda, I.: On the future of Argo: A global, full-depth, multi-disciplinary array, *Frontiers in Marine Science*, 6, 1–28, <https://doi.org/10.3389/fmars.2019.00439>, 2019.
- Sathyendranath, S., Brewin, R. J. W., Brockmann, C., Brotas, V., Calton, B., Chuprin, A., Cipollini, P., Couto, A. B., Dingle, J., Doerffer, R., Donlon, C., Dowell, M., Farman, A., Grant, M., Groom, S., Horseman, A., Jackson, T., Krasemann, H.,
 630 Lavender, S., Martinez-Vicente, V., Mazeran, C., Mélin, F., Moore, T. S., Müller, D., Regner, P., Roy, S., Steele, C. J., Steinmetz, F., Swinton, J., Taberner, M., Thompson, A., Valente, A., Zühlke, M., Brando, V. E., Feng, H., Feldman, G., Franz, B. A., Frouin, R., Gould, R. W., Hooker, S. B., Kahru, M., Kratzer, S., Mitchell, B. G., Muller-Karger, F. E., Sosik, H. M., Voss, K. J., Werdell, J., and Platt, T.: An ocean-colour time series for use in climate studies: The experience of the ocean-colour climate change initiative (OC-CCI), *Sensors*, 19, <https://doi.org/10.3390/s19194285>, 2019.
- 635 Sathyendranath, S., Jackson, T., Brockmann, C., Brotas, V., Calton, B., Chuprin, A., Clements, O., Cipollini, P., Danne, O., Dingle, J., Donlon, C., Grant, M., Groom, S., Krasemann, H., Lavender, S., Mazeran, C., Mélin, F., Müller, D., Steinmetz, F., Valente, A., Zühlke, M., Feldman, G., Franz, B., Frouin, R., Werdell, J., and Platt, T.: ESA Ocean Colour Climate Change Initiative (Ocean_Colour_cci): Version 6.0, 4km resolution data, [dataset], <https://doi.org/10.5285/5011D22AAE5A4671B0CBC7D05C56C4F0>, 2023.



- 640 Shutler, J. D., Land, P. E., Piolle, J. F., Woolf, D. K., Goddijn-Murphy, L., Paul, F., Girard-Ardhuin, F., Chapron, B., and
 Donlon, C. J.: FluxEngine: A flexible processing system for calculating atmosphere-ocean carbon dioxide gas fluxes and
 climatologies, *Journal of Atmospheric and Oceanic Technology*, 33, 741–756, <https://doi.org/10.1175/JTECH-D-14-00204.1>, 2016.
- Shutler, J. D., Wanninkhof, R., Nightingale, P. D., Woolf, D. K., Bakker, D. C., Watson, A., Ashton, I., Holding, T.,
 645 Chapron, B., Quilfen, Y., Fairall, C., Schuster, U., Nakajima, M., and Donlon, C. J.: Satellites will address critical science
 priorities for quantifying ocean carbon, *Frontiers in Ecology and the Environment*, 18, 27–35,
<https://doi.org/10.1002/fee.2129>, 2020.
- Shutler, J. D., Gruber, N., Findlay, H. S., Land, P. E., Gregor, L., Holding, T., Sims, R. P., Green, H., Piolle, J.-F., Chapron,
 B., Sathyendranath, S., Rousseaux, C. S., Donlon, C. J., Cooley, S., Turner, J., Valauri-Orton, A., Lowder, K., Widdicombe,
 650 S., Newton, J., Sabia, R., Rio, M.-H., and Gaultier, L.: The increasing importance of satellite observations to assess the
 ocean carbon sink and ocean acidification, *Earth-Science Reviews*, 250, 104682,
<https://doi.org/10.1016/j.earscirev.2024.104682>, 2024.
- Song, H., Marshall, J., Munro, D. R., Dutkiewicz, S., Sweeney, C., McGillicuddy, D. J., and Hausmann, U.: Mesoscale
 modulation of air-sea CO₂ flux in Drake Passage, *Journal of Geophysical Research: Oceans*, 121, 6635–6649,
 655 <https://doi.org/10.1002/2016JC011714>, 2016.
- Taylor, J. R.: An introduction to error analysis, University Science Books, Sausalito, Calif., 1997.
- Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G. C., Landschützer, P., Woolf, D. K., and Goddijn-
 Murphy, L.: Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon inventory, *Nature*
Communications, 11, 1–6, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.
- 660 Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, 2, 203–215,
[https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/10.1016/0304-4203(74)90015-2), 1974.
- Woolf, D. K., Land, P. E., Shutler, J. D., Goddijn-Murphy, L. M., and Donlon, C. J.: On the calculation of air-sea fluxes of
 CO₂ in the presence of temperature and salinity gradients, *Journal of Geophysical Research: Oceans*, 121, 1229–1248,
<https://doi.org/10.1002/2015JC011427>, 2016.
- 665 Woolf, D. K., Shutler, J. D., Goddijn-Murphy, L., Watson, A. J., Chapron, B., Nightingale, P. D., Donlon, C. J., Piskozub, J.,
 Yelland, M. J., Ashton, I., Holding, T., Schuster, U., Girard-Ardhuin, F., Grouazel, A., Piolle, J. F., Warren, M., Wrobel-
 Niedzwiecka, I., Land, P. E., Torres, R., Prytherch, J., Moat, B., Hanafin, J., Ardhuin, F., and Paul, F.: Key Uncertainties in
 the Recent Air-Sea Flux of CO₂, *Global Biogeochemical Cycles*, 33, 1548–1563, <https://doi.org/10.1029/2018GB006041>,
 2019.

670