



The PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database: A FAIR-aligned compilation of seawater $\delta^{18}\text{O}$ data to uncover 'hidden' insights from the global ocean

Alyssa R. Atwood¹, Andrea L. Moore¹, Kristine L. DeLong², Sylvia E. Long¹, Sara C. Sanchez³, Jessica
5 A. Hargreaves⁴, Chandler A. Morris⁵, Raquel E. Pauly¹, Émilie P. Dassié⁶, Thomas Felis⁴, Antje H.L.
Voelker⁷, Sujata A. Murty⁸, Kim M. Cobb⁵

¹Dept. of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee, FL, 32303, USA

²Geography and Anthropology and the Coastal Studies Institute, Baton Rouge, LA, 70803, USA

³Atmospheric and Oceanic Science Department, University of Colorado Boulder, Boulder, CO, 80309, USA

10 ⁴MARUM - Center for Marine Environmental Sciences, University of Bremen, Bremen, 28359, Germany

⁵Department of Earth, Environmental, and Planetary Sciences, Brown Univ., Providence, RI, 02912, USA

⁶UMR EPOC, University of Bordeaux, Pessac, 33615, France

⁷Divisão de Geologia e Georecursos Marinhos, Instituto Português do Mar e da Atmosfera, Alges, 1495-165, Portugal

⁸Atmospheric and Environmental Sciences, University at Albany, State University of New York, Albany, NY, 12226, USA

15 *Correspondence to:* Alyssa R. Atwood (aatwood@fsu.edu)

Abstract. The stable isotope values of seawater ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) provide valuable information on the exchange of water between the ocean, atmosphere, and cryosphere and on ocean mixing processes. As such, observational seawater $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data place powerful constraints on hydrologic changes in the modern ocean. Seawater $\delta^{18}\text{O}$ data are also essential for calibrating paleoclimate proxies based on the $\delta^{18}\text{O}$ of marine carbonates and are an increasingly critical diagnostic tool for assessing model
20 performance and skill in isotope-enabled global climate models. Despite their broad value, no centralized and actively-curated database for this type of data exists, even though a growing number of new seawater $\delta^{18}\text{O}$ datasets have been generated over the last decade. As such, many seawater $\delta^{18}\text{O}$ datasets remain 'hidden'. To improve the accessibility of seawater $\delta^{18}\text{O}$ data for the Earth Science research community, the Past Global Changes (PAGES) CoralHydro2k project has created a new, machine-readable, and metadata-rich database of observational seawater $\delta^{18}\text{O}$ data, paired with seawater $\delta^2\text{H}$ and salinity data, that is
25 compliant with findability, accessibility, interoperability, and reusability (FAIR) standards for digital assets. The data has been collected from public databases and repositories, direct researcher data submissions, scientific papers, and student theses. In total, the PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database contains over 18,600 data points with extensive metadata that makes the database suitable for a myriad of research applications. For hidden data, we searched for and included all datasets within the global ocean. For public data, our data collation efforts were focused on the upper 50 m from 35°N to 35°S (to aid in
30 CoralHydro2k's seawater $\delta^{18}\text{O}$ reconstruction studies using $\delta^{18}\text{O}$ and Sr/Ca in tropical-subtropical coral skeletons). We also provide a set of best practices to the community for reporting seawater isotope data in the future.



Short Summary. The stable isotopic composition of seawater is a valuable tool for studying the global water cycle in the past, present, and future. However, an active repository dedicated to archiving this type of data has been lacking, and many datasets remain hidden from public view. We have created a new database of observational seawater isotope data that is rich in metadata, publicly accessible, and machine readable to increase its availability and usability for a variety of Earth Science applications.

1 Introduction

1.1 Progress and challenges in the synthesis of seawater oxygen isotope data

The stable isotopes of water ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) are powerful tracers of the global water cycle, tracking water as it continuously cycles between the ocean, atmosphere, and land. As water molecules undergo phase changes during this process, the lighter, more abundant isotope (^{16}O and ^1H) is preferentially vaporized during evaporation with respect to the heavier, less abundant isotope (^{18}O and ^2H), while ^{18}O and ^2H are preferentially condensed during precipitation (Dansgaard, 1964). This partitioning of isotopes based on mass allows the isotope values (where $\delta^{18}\text{O} = \left[\frac{\frac{^{18}\text{O}}{^{16}\text{O}}_{\text{sample}}}{\frac{^{18}\text{O}}{^{16}\text{O}}_{\text{standard}}} - 1 \right] * 1000$) of water to be used as a tracer of the hydrologic cycle (Dansgaard, 1954; Galewsky et al., 2016; Gat, 1996). In the ocean, the isotope values of seawater ($\delta^{18}\text{O}_{\text{sw}}$ and $\delta^2\text{H}_{\text{sw}}$) can provide valuable information on an array of processes, including heat and mass exchange with the atmosphere (via precipitation and evaporation), large-scale ocean circulation, and freshwater input from rivers and ice sheets (Akhoudas et al., 2021; Benetti et al., 2016; Benway and Mix, 2004; Biddle et al., 2019; Craig and Gordon, 1965; Dee et al., 2023; Frew et al., 2000; Imbrie et al., 1984; Jacobs et al., 1985; Lisiecki and Raymo, 2005; Meredith et al., 1999; Strain and Tan, 1993). $\delta^{18}\text{O}_{\text{sw}}$ and $\delta^2\text{H}_{\text{sw}}$ values can also provide insight into other ocean tracers such as salinity, since they covary strongly due to the influence of evaporation and precipitation on each of these variables (Craig and Gordon, 1965; LeGrande and Schmidt, 2011). However, because key processes act differentially on salinity as compared to the stable isotope values, $\delta^{18}\text{O}_{\text{sw}}$ and $\delta^2\text{H}_{\text{sw}}$ provide an additional degree of freedom for constraining ocean mixing and the local moisture budget. Furthermore, stable isotope measurements, such as $\delta^{18}\text{O}$ in marine biominerals and $\delta^2\text{H}$ in lipids, can be used to trace plankton and animal movement and provide provenance for ecology, conservation, archaeology, and food forensics studies (Doubleday et al., 2022). Given these wide-ranging applications, seawater isotope data are used in a wide range of fields, including oceanography, atmospheric science, geology, marine biology, archaeology, and geography.

Seawater isotope values also create a common unit that uniquely links paleoclimate reconstructions to modern climate observations and isotope-enabled model simulations. Modern $\delta^{18}\text{O}_{\text{sw}}$ data are essential for the calibration of paleoclimate proxies of past ocean variability based on the $\delta^{18}\text{O}$ of marine carbonates such as corals, foraminifera, mollusks, ostracods, and coralline algae. Recent paleoclimate data assimilation efforts such as the Last Millennium Reanalysis project (e.g., Tardif et al., 2019) would greatly benefit from a spatial network of $\delta^{18}\text{O}_{\text{sw}}$ data to improve quantification of proxy uncertainty and for



training the proxy system models that underlie those efforts. Modern $\delta^2\text{H}_{\text{sw}}$ data are used in the calibration of
 65 paleoceanographic proxies based on the $\delta^2\text{H}$ of alkenones and other lipid biomarkers in marine sediments (e.g., Eglinton and
 Eglinton, 2008). When used in tandem with $\delta^{18}\text{O}$ data (i.e., to calculate d-excess in surface ocean and overlying water vapor),
 these data can be used to constrain evaporation parameters (e.g., Benetti et al., 2014). As such, observational and reconstruction
 efforts based on seawater isotope values enable scientists to better understand the underlying physics that govern the water
 cycle, and to extend hydroclimate records back to the preindustrial era, thus contextualizing anthropogenic climate change and
 70 improving the skill of future climate projections.

Observational $\delta^{18}\text{O}_{\text{sw}}$ data can also be used to provide boundary conditions in climate models and to assess model performance
 and skill. The increasing integration of oxygen isotopes of water in climate models – from models of intermediate complexity
 to fully coupled Earth System Models (e.g., Blossey et al., 2010; Bong et al., 2024; Bony et al., 2008; Brady et al., 2019;
 75 Cauquoin et al., 2019; Dee et al., 2015; Field et al., 2014; Fiorella et al., 2021; Kurita et al., 2011; Lee and Fung, 2008; Noone
 and Simmonds, 2002; Nusbaumer et al., 2017; Risi et al., 2010, 2020, 2021; Schmidt et al., 2007; Tada et al., 2021; Wei et al.,
 2018; Werner et al., 2011; Yoshimura et al., 2008) – bolsters the interpretation of modern and paleoclimate observations, while
 also providing opportunities to test model performance in resolving key features of the hydrologic cycle, e.g., the representation
 of moisture transport, circulation, and surface water fluxes.

80 Paralleling recent advances in the numerical simulation of water isotopes, new analytical capabilities have also developed in
 recent years, including new *in situ* atmospheric measurement techniques and strategies (Finkenbiner et al., 2022; Gupta et al.,
 2009; Henze et al., 2022), and the development of global atmospheric data products from a variety of remote sensors (e.g.,
 Diekmann et al., 2021; Schneider et al., 2022; Worden et al., 2019). As a result, measurements of water isotopes have become
 85 increasingly incorporated in coordinated observing networks and monitoring studies of precipitation and atmospheric water
 vapor, including the Global Network of Isotopes in Precipitation (www.iaea.org/services/networks/gnip) and the National
 Ecological Observatory Network (www.neonscience.org/).

However, no such coordinated observing network for seawater $\delta^{18}\text{O}$ currently exists. Unlike meteorological observations on
 90 land, observations of ocean hydrological properties (e.g., precipitation, evaporation, and salinity) are either limited to the past
 few decades *via* satellite remote sensing and the ARGO program (Wong et al., 2020) or are confined to select coastal and
 island locations that have the necessary infrastructure to support sustained *in situ* measurements of ocean surface properties.
 Furthermore, these ocean observations rarely include $\delta^{18}\text{O}_{\text{sw}}$ because there is currently no cost-effective, easily deployable
 instrumentation to measure seawater isotopes *in situ*. Thus seawater samples must be taken back to a laboratory for isotopic
 95 analysis. Despite these structural challenges, a growing number of $\delta^{18}\text{O}_{\text{sw}}$ datasets have been generated in recent decades due
 to the accelerated collection of $\delta^{18}\text{O}_{\text{sw}}$ samples, new instrumentation such as cavity-ring down isotope analyzers with reduced



analytical costs, and the capability to measure both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ in parallel, and new sampling devices that enable long-term seawater sample collections (e.g., Jannasch et al., 2004; Khare et al., 2021).

100 In recognition of the broad value of $\delta^{18}\text{O}_{\text{sw}}$ data to the Earth Sciences, a major effort to gather $\delta^{18}\text{O}_{\text{sw}}$ data occurred in the 1990s (Bigg and Rohling, 2000; Schmidt, 1999; Schmidt et al., 1999) and resulted in the development of the NASA's Goddard Institute for Space Studies (GISS) Global Seawater Oxygen-18 database (<https://data.giss.nasa.gov/o18data/>), which contains over 26,000 global measurements of $\delta^{18}\text{O}_{\text{sw}}$ (and some $\delta^2\text{H}$ data) from the 1950s to 2000s. In 2006, that database was used to construct a global gridded dataset of $\delta^{18}\text{O}_{\text{sw}}$ and to characterize regional relationships between $\delta^{18}\text{O}_{\text{sw}}$ and salinity (LeGrande and Schmidt, 2006) and it has subsequently been used in a broad range of studies involving $\delta^{18}\text{O}_{\text{sw}}$. However, the NASA GISS database is no longer actively updated, with the last $\delta^{18}\text{O}_{\text{sw}}$ measurement added in 2011. As a result, a growing number of new $\delta^{18}\text{O}_{\text{sw}}$ datasets published since 2011 remain without an active $\delta^{18}\text{O}_{\text{sw}}$ -specific data repository in which to archive the data. Researchers have instead provided the $\delta^{18}\text{O}_{\text{sw}}$ data in the supplemental tables of journal articles, or have archived the $\delta^{18}\text{O}_{\text{sw}}$ data with other geochemical data (e.g., coral $\delta^{18}\text{O}$), in data repositories such as the National Centers for Environmental Information (NCEI) for Paleoclimatology (<https://www.ncei.noaa.gov/products/paleoclimatology>) and Pangaea (<https://www.pangaea.de/>). Because these datasets can be difficult to find, non-machine-readable, and/or decentralized, they are not easily accessible to the wide range of research communities that would benefit from this data (see a related review by Chamberlain et al., 2021). Furthermore, many publishers and several funding agencies now require researchers to archive their data in FAIR and public repositories. For these reasons, a comprehensive database of $\delta^{18}\text{O}_{\text{sw}}$ data that is publicly available and actively maintained is critically needed.

1.2 The PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database

Inspired by the PAGES (Past Global Changes) Hydro2k Workshop in 2016 (PAGES Hydro2k Consortium, 2017), the PAGES CoralHydro2k project was formed in 2017 to investigate the variability of hydrology and temperature in the tropical surface ocean during the past 2000 years based on the combination of coral $\delta^{18}\text{O}$, which varies with temperature and $\delta^{18}\text{O}_{\text{sw}}$, and the strontium-to-calcium ratio (Sr/Ca) in corals, which is a temperature proxy. The CoralHydro2k project was built upon previous PAGES 2k efforts, namely Ocean2k and Iso2k (Konecky et al., 2020, 2023; Tierney et al., 2015), which compiled published coral $\delta^{18}\text{O}$ records and other data into new machine-readable databases to track temperature and hydroclimate changes over the Common Era. To aid in the calibration and interpretation of the paired coral $\delta^{18}\text{O}$ and Sr/Ca records in the database, and derive coral-based reconstructions of seawater $\delta^{18}\text{O}$, the CoralHydro2k project also started to compile $\delta^{18}\text{O}_{\text{sw}}$ data.

125 In recognition of the broad value of $\delta^{18}\text{O}_{\text{sw}}$ data and the growing number of $\delta^{18}\text{O}_{\text{sw}}$ datasets that have been generated during the last two decades, the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database project was launched in 2020 to recover 'hidden' $\delta^{18}\text{O}_{\text{sw}}$ data that were not easily findable. During the past five years, we have integrated these records, along with any associated $\delta^2\text{H}$, salinity, and temperature data, with data from public databases and repositories to create a new, centralized, machine-readable,



130 and metadata-rich database that aligns with findability, accessibility, interoperability, and reusability (FAIR) standards (Wilkinson, 2016). Here we provide a detailed description of the PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database. We highlight the opportunities and limitations of this database, and provide a set of best practices to the community for reporting this type of seawater isotope data in the future.

2 Methods

135 2.1 Collaborative model

CoralHydro2k and its Seawater $\delta^{18}\text{O}$ Database started in 2017 as a project in Phase 3 of the PAGES 2k network, a long-running initiative to study past global changes over the last 2000 years and to compile paleoclimate data in publicly available, machine-readable databases (PAGES 2k Network Coordinators, 2017). CoralHydro2k included team members from the Phase 1 PAGES 2k Ocean2k working group (Tierney et al., 2015) and Phase 2 Iso2k working group (Konecky et al., 2020) and many new members, particularly from the coral paleoclimate community. CoralHydro2k continues into Phase 4 of PAGES 2k, focusing on reconstructing past changes in tropical ocean temperature and hydroclimate using paired Sr/Ca and $\delta^{18}\text{O}$ from coral archives over the last 2000 years (Hargreaves et al., 2020; Walter et al., 2023). Recurring calls went out within the international paleoclimate community for working group members, coral experts, and paleo data assimilation experts to join the effort with monthly teleconference meetings and one in-person meeting in 2019 (Hargreaves et al., 2020). As a result, the CoralHydro2k database was produced, a global, actively curated compilation of coral $\delta^{18}\text{O}$ and Sr/Ca proxy records of tropical ocean hydrology and temperature for the Common Era (Walter et al., 2022, 2023). A number of sub-projects were developed in conjunction with CoralHydro2k, including a project to develop new proxy system models (PSM) for coral $\delta^{18}\text{O}$. The group working on this sub-project realized that the spatial and temporal coverage of observational $\delta^{18}\text{O}_{\text{sw}}$ data were too sparse to integrate into the PSM framework and that many new $\delta^{18}\text{O}_{\text{sw}}$ datasets produced during the last few decades are not easily findable or accessible.

Therefore, CoralHydro2k formed a new sub-project in 2020 to compile existing seawater $\delta^{18}\text{O}$ data with rich metadata following FAIR standards (Atwood et al., 2024; DeLong et al., 2022). Researchers were invited to submit their data to the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database *via* a Qualtrics survey and accompanying YouTube video that provided instructions on how to submit data. Additionally, the team set up a Seawater Oxygen Isotopes Community (<https://www.earthchem.org/communities/seawater-oxygen-isotopes/>) in the EarthChem Library (ECL), a data repository that archives, publishes, and provides access to data in the geosciences. The ECL offers a suite of services for data preservation and access, including long-term archiving and data registration with a Digital Object Identifier (DOI). Through the new Seawater Oxygen Isotopes Community, new seawater $\delta^{18}\text{O}$ (and $\delta^2\text{H}$) datasets can be submitted and assigned a DOI, which allows the datasets to be cited and tracked when used by other researchers. The CoralHydro2k members promoted this new



database at international conferences in the United States and Europe, in the PAGES newsletter (Atwood et al., 2024), and in Eos, the monthly magazine of the American Geophysical Union (AGU) (DeLong et al., 2022).

The workload for assembling the seawater data and metadata was performed by CoralHydro2k members and new members of the Seawater $\delta^{18}\text{O}$ Database sub-project. The team was made of volunteer scientists from all academic levels, including undergraduate and graduate students, postdoctoral researchers, and early- to senior-level scientists from a number of international academic and research institutions. The work was completed remotely in synchronous working sessions and asynchronously across several virtual platforms (Google Suite, Slack, and Zoom). Data discovery, metadata protocols, and compilation were done collaboratively as the project progressed.

2.2 Data aggregation and formatting

The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database was designed to be as inclusive and comprehensive as possible in its record-selection criteria to support the project's goal of developing a FAIR database of global seawater $\delta^{18}\text{O}$ measurements, paired with $\delta^2\text{H}$ and salinity measurements, and to include as much 'hidden' data as possible. Thus, the Seawater $\delta^{18}\text{O}$ Database selection criteria were less restrictive than other PAGES 2k efforts, and the database includes data from peer-reviewed scientific literature, student theses and dissertations, public data repositories, and direct author submission.

In alignment with FAIR data principles, the Seawater $\delta^{18}\text{O}$ Database contains extensive metadata. Where available, the seawater $\delta^{18}\text{O}$ data is paired with seawater $\delta^2\text{H}$, salinity, and temperature data. Eight metadata fields are required, with an additional 44 optional metadata fields that provide supporting information on site information, sample collection and storage notes, the isotope analysis method and instrumentation, and error information. Additionally, a template is provided to assist researchers with future submissions to the database and to establish a set of best practices for reporting seawater isotope data.

For hidden data, we searched for and included datasets spanning all depths and all latitudes across the global ocean. For publicly available data, we focused on including data from the upper 50 m between 35°N to 35°S (to aid in CoralHydro2k's seawater $\delta^{18}\text{O}$ reconstruction studies using $\delta^{18}\text{O}$ and Sr/Ca in tropical-subtropical corals).

2.3 Metadata description and quality control

The metadata is described in this section and Tables 1–2. The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database team implemented several rounds of quality control measures for the data and metadata. Following the Iso2k database procedure (Konecky et al., 2020), each metadata field has an associated quality control certification "Level" from 1 to 6, described below and in Table 1. Level 1 and Level 2 metadata fields constitute 'essential' metadata, and if a dataset lacked one of these fields, it was excluded from the database.



- *Level 1* fields are required for inclusion in the database and they contain standardized vocabularies, according to Table 2. They are recommended as primary fields for filtering and querying records in the database. They were subject to the highest Quality Control (QC) standard. Examples of Level 1 metadata are: “Collection year”, “Collection month”, “Latitude”, “Longitude”, and “Depth”.
- *Level 2* metadata fields are required for inclusion, but they are not generalizable enough to use standardized vocabularies. They were subject to the highest QC standard and the metadata were obtained from the original publication or data source. An example of Level 2 metadata is “Site name or geographic area”.
- *Level 3* metadata fields add important supplementary information related to the seawater $\delta^{18}\text{O}$ measurements. They contain standardized vocabularies and can be used as secondary fields for filtering and querying the database; however, they are generally not available for all records and thus not required for inclusion in the database. They were subject to the highest QC standard. Examples of Level 3 metadata are: “Collection day”, “ $\delta^{18}\text{O}$ error”, “ $\delta^{18}\text{O}$ analysis technique”, “Water isotope analysis date”, “ $\delta^2\text{H}$ value”, “Temperature value”, and “Salinity value”.
- *Level 4* metadata fields also add important supplementary information related to the seawater $\delta^{18}\text{O}$ data, but they are not generalizable enough to use standardized vocabularies. They are also generally not available for all records and thus not required for inclusion in the database. They were subject to the highest QC standard. Examples of Level 4 metadata are: “ $\delta^{18}\text{O}$ correction notes”, “ $\delta^{18}\text{O}$ error notes”, “Sample ID”, “Publication citation”, “Dataset citation”, “Cruise ID”, “ $\delta^{18}\text{O}$ analysis location”, “Sample collection, processing, and storage notes”, and “Water isotope analysis notes”.
- *Level 5* metadata fields may be useful to some users of the database but they are generally not available for all records and thus not required for inclusion in the database. In many cases, these fields contain freeform text with direct quotes from the original publications. During the QC certification process, these fields were checked against the original publication and a quote or summary of the relevant information was provided in the database, but the information provided may not be comprehensive. Examples of Level 5 metadata are: “Location description” and “Location type”.
- *Level 6* metadata fields may be useful to some users of the database. This metadata field was completed when the information was easily accessible from the original publications, but some metadata may be missing. There is only one Level 6 metadata field in the database: “Temperature/salinity notes”.

Table 1: Description of all metadata fields in the PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database. Bold text indicates required fields in the database (Level 1 and 2).

Level #	Metadata field	Metadata field description	Type	Metadata category
1	CoralHydro2kID*	Unique ID for this database	Text	Entity
1	Collection year	Year of sample collection, YYYY	Numeric	Entity
1	Collection month	Month of sample collection, MM	Numeric	Entity
1	Latitude	Latitude of the sampling site in decimal degrees. South is negative. Decimal degrees N, from -180 to 180	Numeric	Entity
1	Longitude	Longitude of the sampling site in decimal degrees.	Numeric	Entity



		West is negative. Decimal degrees E, from -180 to 180		
1	Depth	Sampling depth, in meters (m) below sea level (no minus sign).	Numeric	Entity
1	Depth units	Depth measurement units (meters below sea level)	Text	Entity
1	$\delta^{18}\text{O}$ value	Measured $\delta^{18}\text{O}$ value	Numeric	Seawater Data
1	$\delta^{18}\text{O}$ units	$\delta^{18}\text{O}$ units (per mille)	Text	Seawater Data
1	$\delta^{18}\text{O}$ correction	Indicates whether a correction has been made to the $\delta^{18}\text{O}$ values after the original publication. If the data point has been corrected, "Y" is indicated. If the data has not been corrected, "N" is indicated. If known (e.g., in the data collected from the NASA GISS $\delta^{18}\text{O}_{\text{sw}}$ database), the value of the applied correction is indicated in the " $\delta^{18}\text{O}$ correction notes" metadata field. Additional information about each correction is provided in " $\delta^{18}\text{O}$ correction notes" (Level 4) metadata field.**	Logic	Seawater Data
1	Evaporation flag	Flag indicating the presence of potential evaporation effects on $\delta^{18}\text{O}$ value. "Y" is indicated for cases where authors note that sample evaporation could be a concern or cases where $\delta^{18}\text{O}$ data have been corrected for evaporation; "N" otherwise. Further information is provided in the "Sample collection, processing, and storage notes" (Level 4) metadata field.	Logic	Queryable
1	Reference standard	Reference standard used in reporting the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values (SMOW, VSMOW)	Text	Seawater Data
1	Access date	Date in which the data was downloaded from data repositories, submitted by researchers, or acquired from journal articles (YYYY/MM/DD)	Text	Entity
2	Site name or geographic area	Name of the site or the general area from which the water sample was collected	Text	Entity
3	Collection day	Day of sample collection, DD. In some cases, the collection day was not specified in the original publication, only the collection month or a range of dates. In these cases, the midpoint of the date range	Numeric	Entity



		was selected as the collection day and note is made in the 'Collection date notes' (Level 4) metadata field.		
3	Collection time	Time of sample collection, in Coordinated Universal Time (UTC) 24-hour format, HH:MM:SS	Text	Entity
3	Water isotope analysis date	Date of water isotope analysis (YYYY/MM/DD)	Text	Entity
3	$\delta^{18}\text{O}$ error	Reported error of the $\delta^{18}\text{O}$ value. Because many different types of error are reported in the literature, standardization was impossible; therefore, we report the most comprehensive error provided. The type of error along with any supporting information is provided in the " $\delta^{18}\text{O}$ error notes" (Level 4) metadata field.	Numeric	Seawater Data
3	$\delta^{18}\text{O}$ error units	If $\delta^{18}\text{O}$ error exists, the units are specified (per mille)	Text	Seawater Data
3	$\delta^{18}\text{O}$ analysis technique	Type of instrument used to make the isotope measurement (isotope ratio mass spectrometry (IRMS), Cavity Ring Down Spectroscopy (CRDS), and off-axis integrated cavity output spectroscopy (ICOS)	Text	Entity
3	$\delta^2\text{H}$ value	Measured $\delta^2\text{H}$ value	Numeric	Seawater Data
3	$\delta^2\text{H}$ units	$\delta^2\text{H}$ units (per mille)	Text	Seawater Data
3	$\delta^2\text{H}$ error	Reported error of the $\delta^2\text{H}$ value. Because many different types of error are reported in the literature, standardization was impossible; therefore, we report the most comprehensive error provided.	Numeric	Seawater Data
3	Temperature value	Seawater temperature (degrees Celsius)	Numeric	Seawater Data
3	Temperature units	Units of temperature value (degrees Celsius).	Text	Seawater Data
3	Temperature error	Reported error of the temperature value	Numeric	Seawater Data
3	Salinity value	Seawater salinity	Numeric	Seawater Data
3	Salinity units	Salinity units (typically "PSU" or "parts per thousand"). Retain original units provided in the publication.	Text	Seawater Data



3	Salinity error	Reported error of the salinity value	Numeric	Seawater Data
4	Collection date notes	A note is made here if the collection day was not specified in the original publication, and only the collection month or a range of dates were specified. In this case the midpoint of the date range was selected as the collection day.	Text	Entity
4	Location notes	A note is made here if the latitude and longitude coordinates are not exact (e.g., some of the NASA GISS database entries have notes stating “A: Position was read off a graph of locations and therefore is not exact”). In these cases, the notes are copied to this metadata field.	Text	Entity
4	Depth notes	If only pressure (and not depth) was reported in the original dataset, a note is made here about how the depth conversion was performed. If a range of depths is provided in the original dataset, the midpoint of the depth range is reported in the ‘Depth’ field and the range is stated here.	Text	Entity
4	$\delta^{18}\text{O}$ correction notes	If any $\delta^{18}\text{O}$ correction was made to the $\delta^{18}\text{O}$ values subsequent to the original publication, the value of the correction is reported here, along with any accompanying information about how and why the correction was made.	Text	Seawater Data
4	$\delta^{18}\text{O}$ error notes	Information about the reported error of the $\delta^{18}\text{O}$ value, including the type of error along with any supporting information.	Text	Seawater Data
4	Sample ID	Unique sample ID provided by original authors.	Text	Queryable
4	$\delta^{18}\text{O}$ analysis location	University or institute where the isotope measurements were made.	Text	Entity
4	Publication citation	Citation of the original publication of the data. When the data was obtained from a data repository and no publication citation was found, “NaN” is entered in this field and the relevant citation appears in the “Dataset citation” field.	Text	Entity
4	Publication DOI or URL	DOI or URL of the original publication.	Text	Entity
4	Dataset citation	If the data was obtained from a data repository, the dataset citation is provided here.	Text	Entity



4	Dataset URL	If the data was obtained from a data repository, the dataset URL is provided here.	Text	Entity
4	Dataset ID	If the data was obtained from a data repository, the Dataset ID from that repository is provided here.	Text	Entity
4	Data provenance notes	Indicates the source of the $\delta^{18}\text{O}$ data (and $\delta^2\text{H}$, SST, and SSS data). In some cases, select metadata may have been collected from other data sources, such as the original publication, which is indicated in this field.	Text	Entity
4	Cruise ID	Cruise ID, if applicable.	Text	Entity
4	Station ID	Station ID, if applicable.	Text	Entity
4	Cruise report	Citation of the cruise report, if applicable.	Text	Entity
4	Cruise report URL	Link to the cruise report, if applicable.	Text	Entity
4	Sample collection, processing, and storage notes	Notes on sample collection, sample processing, and sample storage. Sample processing includes any treatments prior to analysis (distillation, filtration, etc). Sample storage includes the type of storage container and any preventative measures taken against evaporation (sample seals, refrigeration, etc).	Text	Entity
4	Water isotope analysis notes	Notes on the isotope analysis methods, including details of any equilibration steps used between the water and CO_2 or H_2 gas, the specific type of instrumentation used for the isotopic analysis, details of the calibration steps and standards used, number of replicate measurements, corrections for instrumental drift and memory effects, and preventative steps taken to minimize salt contamination (e.g., for cavity ring down spectroscopy).	Text	Entity
5	Location type	Type of water body from which the water samples were collected (e.g., open ocean, coastal, bay, lagoon, estuary, enclosed sea, marginal/semi-enclosed sea).	Text	Entity
5	Location description	Description of the sampling location, including a description of the major water masses and currents influencing the region, as well as details on the surface water balance, groundwater or riverine input, upwelling, distance from the coastline, depth and geometry of bay or lagoon, and/or description of co-located coral reef site if applicable.	Text	Entity



6	Temperature/salinity notes	Notes on the analysis of temperature and salinity.	Text	Entity
---	----------------------------	--	------	--------

*There are four cases in which datasets obtained from the GISS database were not clearly associated with a publication, or the provided reference did not match the dataset. In those cases, the Schmidt et al. (1999) database citation is provided in the “Dataset citation” metadata field and the author letters “SC” are used in the unique CoralHydro2k ID to reference that citation (SC99AO0001, SC99PO0001, SC99IO0001, SC99GI0001). Additional details about the citations and data provenance appear in the “Data provenance notes” metadata field.

**Corrections were applied in several datasets in the NASA GISS $\delta^{18}\text{O}_{\text{sw}}$ database to standardize the data based on deep water masses to correct for changes in standards, different analysis techniques, and other systematic errors. All corrections are noted in the “ $\delta^{18}\text{O}$ correction” metadata field, so the user can remove the corrections if desired. Corrections were also applied in some datasets in the Reverdin et al. (2022) LOCEAN database to adjust for minor evaporation biases. In that case, the correction value is unknown.

Table 2: Standardized controlled vocabulary options for metadata fields in the database.

Metadata Field	Standardized Entries
Depth units	m (below sea level)
$\delta^{18}\text{O}$ units	per mille
$\delta^{18}\text{O}$ correction	Y, N
Evaporation flag	Y, N
Reference standard	VSMOW, SMOW
$\delta^{18}\text{O}$ error units	per mille
$\delta^{18}\text{O}$ analysis technique	IRMS, CRDS, ICOS
$\delta^2\text{H}$ units	per mille
Temperature units	°C
Salinity units	PSU, PSS-78, PPT, g/L



235 3 Key characteristics of the seawater $\delta^{18}\text{O}$ data

3.1 Spatial and temporal coverage of the database

The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database contains 18,598 data points from 106 datasets (Fig. 1A,B). 53% of the data is categorized as “hidden” data (i.e., data not currently available in public databases or public repositories; Fig. 1C,D), and the remaining 47% of the data is from public databases or public repositories (Table 3). 10,407 measurements (56%) are from the sea surface (depth ≤ 5 m; Fig. 1A), 3,693 (20%) are from the mixed layer (between 5–50 m), and 4,498 (24%) are below 50 m. The time span of the database covers 1972 to 2021 (Fig. 2) and the depth range covers the surface to 5,797 m below sea level. The earliest data point in the database was collected in September 1972 and the most recent data point was collected on October 8, 2021. 3,480 data points (19%) were collected before the year 2000, and 15,118 data points (81%) were collected on or after the year 2000 (Fig. 2). Because the search for hidden datasets focused on the region between 35°N and 35°S, 75% of the measurements in the database are located within the tropical-subtropical region.




Figure 1: Locations and seawater $\delta^{18}\text{O}$ values of data in the database: (A) all surface ocean data (upper 5 m of the water column), (B) all subsurface ocean data (below 5 m), (C) the hidden surface ocean data only (upper 5 m), (D) the hidden subsurface ocean data only (below 5 m).




Figure 2: Sample collection dates of the seawater $\delta^{18}\text{O}$ data in the database from (A) the surface ocean (upper 5 m) and (B) the subsurface ocean (below 5 m).

255 In addition to $\delta^{18}\text{O}$ measurements, the database also includes paired $\delta^2\text{H}$, salinity, and temperature measurements when available. 16,098 data points (87%) have paired salinity values (Fig. 3A), 13,871 data points (75%) have paired temperature values (Fig. 3B), and 9,769 data points (53%) have paired $\delta^2\text{H}$ measurements (Fig. 3C). 185 measurements have an evaporation flag (Fig. 3D), which allows the user to filter out samples that may be influenced by post-collection evaporation from the database.




Figure 3: Locations and seawater $\delta^{18}\text{O}$ values of the data in the database with (A) paired salinity measurements, (B) paired temperature measurements, (C) paired $\delta^2\text{H}$ measurements, and (D) an evaporation flag.

Compared to the seawater $\delta^{18}\text{O}$ database presented in LeGrande and Schmidt (2006), data coverage in the surface ocean (upper 5 m) is substantially improved in the tropics and subtropics, particularly in the northern Indian Ocean, the eastern Atlantic Ocean, the northeast coast of South America, the Mediterranean Sea, and the equatorial Pacific Ocean. However, poor data coverage still exists in the western Pacific Ocean and Maritime Continent region, the southeastern Indian Ocean, and the subtropical Pacific Ocean regions in both hemispheres. Below 5 m depth, the data coverage is even more limited (Fig. 1B). At all depths, regions with reasonable spatial coverage of $\delta^{18}\text{O}_{\text{sw}}$ data contain limited temporal coverage. Typically, only a few years of regular measurements are available from the most highly sampled regions. For example, only 13% of locations contain at least 12 measurements spanning two years within a 2° latitude x 2° longitude grid box (Fig. 4A), and only 9% of the grid boxes contain data that cover at least 50% of the annual cycle (i.e., 6/12 months of the calendar year; Fig. 4B). While the coverage of seawater isotope data has been growing over the last decade, these measurements are still sparse in space and time, thus highlighting the need for globally coordinated sampling campaigns and archiving efforts.




Figure 4: Temporal distribution of near-surface (upper 5m) $\delta^{18}\text{O}_{\text{sw}}$ measurements in the database, aggregated in 2°x2° (lat x lon) grid boxes. (A) Total number of $\delta^{18}\text{O}_{\text{sw}}$ measurements in each grid box. (B) Fraction of calendar year with $\delta^{18}\text{O}_{\text{sw}}$ measurements.

Outside of the tropics and subtropics, the coverage of $\delta^{18}\text{O}$ data in the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database is more sparse, since only hidden datasets were collected from all latitudes and all depths across the global ocean, while datasets from public repositories were only incorporated into the database if the measurements were made in the upper 50 m between 35°N and 35°S. Future database development efforts will include incorporating additional hidden and public datasets.

3.2 Data-model comparisons of the seawater $\delta^{18}\text{O}$ data

To assess how the $\delta^{18}\text{O}_{\text{sw}}$ data in the database compares with isotope-enabled climate model simulations and other products, we compare the climatological annual cycle in $\delta^{18}\text{O}_{\text{sw}}$ at different island sites using four data products: two simulations of isotope-enabled General Circulation Models [the National Center for Atmospheric Research Community Earth System Model Last Millennium Ensemble (1000 years; Brady et al., 2019) and the NASA Goddard Institute for Space Studies E2-R last millennium simulation (ensemble member E4rhLMgTck; 255 years; Colose et al., 2016), a regional ocean model of the Pacific



called isoROMS (44 years; Stevenson et al., 2018), and a gridded dataset of global monthly mean $\delta^{18}\text{O}_{\text{sw}}$ based on data
290 assimilation with the MITgcm (Breitkreuz et al., 2018). The Breitkreuz dataset is based on a 400-year quasi-equilibrated
simulation of a water isotope-enabled global ocean general circulation model constrained by global monthly $\delta^{18}\text{O}_{\text{sw}}$ data
collected from 1950 to 2011 and climatological salinity and temperature data collected from 1951 to 1980.

The characteristics of $\delta^{18}\text{O}_{\text{sw}}$ variability at the four selected sites in the tropical Pacific and Atlantic Oceans vary widely across
295 the different data products and the $\delta^{18}\text{O}_{\text{sw}}$ observations from the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database, with large differences
in both the amplitude and phase of the annual cycle of $\delta^{18}\text{O}_{\text{sw}}$ (Fig. 5). These differences could be due to deficiencies in the
models (associated with model resolution, subgrid-scale parameterizations, and treatment of atmospheric exchange or ocean
mixing processes), and/or uncertainties in the observational data given the low temporal resolution of the $\delta^{18}\text{O}_{\text{sw}}$ measurements.
Clearly, more observational data is needed to determine the source of the discrepancies, pointing to the need for more
300 coordinated and sustained seawater isotope sampling programs. Seawater isotope sampling at multinational observing systems
that are already in place, such as the Tropical Pacific Observing System (TPOS), Bermuda Atlantic Time-series Study (BATS),
GO-SHIP, and GEOTRACES, could expand and complement existing observational programs. For example, incorporating
new sampling devices such as long-term osmotically pumped fluid samplers (Jannasch et al., 2004; Khare et al., 2021) could
provide a relatively straightforward way to add $\delta^{18}\text{O}_{\text{sw}}$ measurements to existing programs. The development of sustained
305 seawater isotope measurements at a network of observational hotspots around the global ocean would provide powerful new
constraints on hydrologic changes in the modern ocean, generating data that could be used to test theoretical predictions, assess
climate model performance and skill, and calibrate paleoclimate proxies for improved paleoclimate reconstruction.




Figure 5: Monthly climatology of $\delta^{18}\text{O}_{\text{sw}}$ at four island locations: Kiritimati Atoll in the Central Pacific Ocean, Palau in the western Pacific Ocean, Puerto Rico in the Caribbean Sea, and Dry Tortugas in the Gulf of Mexico. Five data sources are shown: observed $\delta^{18}\text{O}_{\text{sw}}$ (from this database; black dashed line), simulated $\delta^{18}\text{O}_{\text{sw}}$ from iCESM, iGISS, and isoROMS, and $\delta^{18}\text{O}_{\text{sw}}$ from the reanalysis product of Breitkreuz et al., 2018 (purple; monthly climatology constrained by observed monthly $\delta^{18}\text{O}_{\text{sw}}$ data collected from 1950 to 2011 and climatological salinity and temperature data collected from 1951 to 1980). The three Earth system models are the National Center for Atmospheric Research Community Earth System Model Last Millennium Ensemble (1,000 years; red) (Brady et al., 2019), the NASA Goddard Institute for Space Studies E2-R last millennium simulation (ensemble member E4rhLMgTck; 255 years; yellow) (Colose et al., 2016), and the isoROMs Pacific Ocean simulation (44 years; blue) (Stevenson et al., 2018).



4 Usage notes

4.1 General applications

320 While the primary motivation of the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database was for coral paleoclimate research, this database was designed to be useful to researchers from a wide range of disciplines, including paleoceanography and paleoclimatology, oceanography, marine biology, Earth science, and climatology. For example, of relevance to oceanography, the seawater $\delta^{18}\text{O}$ data can be coupled with the paired salinity data in the database to explore the relationship between these two parameters and investigate hydrological changes of the surface ocean (e.g., Conroy et al., 2017; Durack et al., 2012; LeGrande and Schmidt, 325 2006; Wagner and Slowey, 2011). Because this relationship varies with latitude and may vary with time (Conroy et al., 2017; LeGrande and Schmidt, 2011; Thompson et al., 2022), the seawater $\delta^{18}\text{O}$ database could be used to more comprehensively assess how this relationship varies in space and time.

As for applications in paleoclimatology, the pairing of seawater oxygen isotope data with salinity data can provide transfer 330 equations for reconstructing past salinity variations (e.g., Gagan et al., 1998; Kilbourne et al., 2004; McCulloch et al., 1994; Ren et al., 2003). For example: coral Sr/Ca, a temperature proxy, paired with coral $\delta^{18}\text{O}$, a proxy for both temperature and $\delta^{18}\text{O}_{\text{sw}}$, can be used to remove the temperature component from the coral $\delta^{18}\text{O}$ signal. The derived $\delta^{18}\text{O}_{\text{sw}}$ can then be converted to salinity using the local $\delta^{18}\text{O}_{\text{sw}}$ to salinity transfer equation (e.g., Kilbourne et al., 2004). The same method can be applied to foraminifera Mg/Ca and $\delta^{18}\text{O}$ records to reconstruct salinity variations and can also potentially be applied to bivalves, coralline 335 algae, ostracods, and otoliths (e.g., Light et al., 2018; Schmidt and Lynch-Stieglitz, 2011; Stott et al., 2004; Trofimova et al., 2020; Warner et al., 2022). Many studies have used this paired approach to reconstruct $\delta^{18}\text{O}_{\text{sw}}$ variations for a wide range of time scales (e.g., Brocas et al., 2019; Felis et al., 2009; Giry et al., 2013; Gorman et al., 2012; Hereid et al., 2013; Knebel et al., 2024; Wu et al., 2013); however, few studies have been able to validate their reconstructions with observed $\delta^{18}\text{O}_{\text{sw}}$ records that span more than one year (Conroy et al., 2017; O'Connor et al., 2021). Instead, most studies use reanalysis products such 340 as Simple Ocean Data Assimilation (SODA) (Carton et al., 2000, 2018), or satellite-derived sea surface salinity (SSS) products (e.g., NASA Aquarius, NASA SMOS) (Boutin et al., 2021) for validating the reconstructions (e.g., Cahyarini et al., 2008; Harbott et al., 2023; Hetzinger et al., 2006).

Additionally, the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database can be used in proxy-system model development, paleo-data 345 assimilation, and comparison studies between proxy reconstructions and climate model output (Dee et al., 2023; Evans et al., 2013; Reed et al., 2022; Sanchez et al., 2021; Smerdon, 2012; Stevenson et al., 2018; Thompson et al., 2011). Proxy-derived $\delta^{18}\text{O}_{\text{sw}}$ data can be directly compared with simulations from isotope-enabled models as part of the validation process and to understand oxygen isotope fractionation processes within the hydrological cycle (Dee et al., 2015; Stevenson et al., 2015, 2023). Furthermore, the proxy-derived salinity reconstructions can be compared with reanalysis and other salinity data 350 products, such as SODA, as a separate validation step (e.g., Cahyarini et al., 2008). Finally, the $\delta^{18}\text{O}_{\text{sw}}$ database offers the



opportunity for improved proxy system models with rigorous uncertainty quantification of proxy-derived estimates of salinity. With such estimates, long reconstructions of salinity would provide valuable insights into the low frequency variability of the hydrological cycle over the data sparse tropical oceans. This $\delta^{18}\text{O}_{\text{sw}}$ database is the most comprehensive to date and will be updated as new datasets are published to support ongoing research (see Section 6).

355 **5 Code/Data availability**

5.1 Accessing the database

The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database follows the FAIR data principles (Wilkinson et al., 2016) that strive to make scholarly data findable, accessible, interoperable, and reusable. The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database uses the Comma Separated Values (*.CSV) file format, a machine-readable format for archiving and describing seawater isotope data. Access
360 to the database has been granted for reviewers and editors during the review phase. The data are also available upon request for members of the public that wish to participate in the review process by emailing the corresponding author. Once the review period is complete, the database will be archived on the NOAA NCEI World Data Service for Paleoclimatology (study page: <https://www.ncei.noaa.gov/access/paleo-search/study/34575>) and issued a permanent DOI. A mirror copy of the database will also be hosted at [Waterisotopes.org](https://www.waterisotopes.org).

365 **5.2 Code availability**

Example scripts to help users filter and search the database are available on the CoralHydro2k Seawater Database GitHub page (https://github.com/CoralHydro2k/ch2kSeawater_Database).

5.3 Underlying data sources

The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database includes records (0–50 mbsl, 35°N to 35°S) from ten international databases,
370 including the NASA GISS Global Seawater Oxygen-18 database (Table 3). Literature searches were also conducted to find hidden seawater $\delta^{18}\text{O}$ data (from all depths and latitudes) published only in tables and supplemental data files of published papers, theses, and dissertations. Data was also sourced from author contributions sent directly to this project or the EarthChem community (earthchem.org/communities/seawater-oxygen-isotopes). Researchers should adhere to the data use policies for the underlying data sources (see Table 3 and Appendices A1–A4).

375

Table 3. Databases included

Database	URL (https://)	Notes	References
----------	----------------	-------	------------



British Oceanographic Data Centre	https://www.bodc.ac.uk/	See A1 for data usage policies	Acknowledge the source of the information by including any attribution statement, see A4.
CISE LOCEAN Seawater Isotope (SEANOE)	www.seanoe.org/data/00600/71186/	Public domain, see URL for data usage details.	(waterisotopes-CISE-LOCEAN 2024), (Reverdin et al. 2022)
GEOTRACES	geotraces.webdv.awi.de/	See A2 for data usage policies	(GEOTRACES Intermediate Data Product Group, 2023)
Global Ocean Data Analysis Project (GLODAPv2.2022)	www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.nodc:0257247	See A3 for data usage policies	(Key et al., 2023; Olsen et al., 2016)
NASA GISS Global Seawater Oxygen-18 (GISS)	data.giss.nasa.gov/o18data/	Public domain	(Bigg and Rohling, 2000; Schmidt, 1999; Schmidt et al., 1999)
NOAA National Centers for Environmental Information (NCEI)	www.ncei.noaa.gov/	Public domain	Cite original publication, online resource, dataset and publication DOIs (where available), and date accessed
NOAA NCEI Paleoclimatology	www.ncei.noaa.gov/products/paleoclimatology	Public domain	Cite original publication, online resource, dataset and publication DOIs (where available), and date accessed
NOAA NCEI World Ocean	www.ncei.noaa.gov/products/world-ocean-database	Public domain	(Boyer et al., 2018)
PANGAEA	www.pangaea.de/	Terms of use https://www.pangaea.de/about/terms.php	(Felden et al. 2023)
Waterisotopes.org (WIDB)	https://wateriso.utah.edu/waterisotopes/	See A4 for data usage policies	http://waterisotopes.org



6 Database submission of new datasets and versioning scheme

The CoralHydro2k Seawater $\delta^{18}\text{O}$ project will accept data submissions for updates to the database. All seawater $\delta^{18}\text{O}$ observations are welcomed regardless of location or water depth. To facilitate this process, a Seawater Oxygen Isotopes Community was developed within the EarthChem Library, an open-access repository for geochemical datasets (earthchem.org/communities/seawater-oxygen-isotopes), where researchers can submit their seawater isotope data and obtain a dataset DOI. The Seawater Oxygen Isotopes Community contains a template that can be downloaded to help researchers submit their data (scroll to the bottom of the webpage above and click on “Download Template”). This template is aligned with the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database to facilitate future updates to the database. The template has a README tab in the Microsoft Excel file with details on the template and an example. We hope that the creation of this site helps researchers publish their seawater isotope datasets, thus minimizing the number of “hidden” datasets.

The initial release of the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database will be Version 1.0.0 for this publication. With new submissions, the database will grow as new datasets are added. Database users who find errors in the database can use the “Report an issue” option in the GitHub site. Datasets submitted to the Seawater Oxygen Isotopes Community within the EarthChem Library (earthchem.org/communities/seawater-oxygen-isotopes) can be updated through that site.

As the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database is updated, it will be versioned following the scheme used by other PAGES data collection projects (Ahmed et al., 2013; Emile-Geay et al., 2017; Kaufman et al., 2020; McKay and Kaufman, 2014; Walter et al., 2023). The version number has three counters in the following form: $C_1.C_2.C_3$, where C_1 , C_2 , and C_3 are incrementing integers. When C_1 increases, C_2 and C_3 reset to zero. When C_2 increases, C_3 resets to zero. C_1 represents the number of publications describing the database. C_2 increments each time the set of records in the database changes (addition or removal of a dataset). C_3 increments when the data or metadata within the dataset changes, but the set of records remains the same. Upon updates, extensions, or corrections to the database, rather than issuing errata to this publication, changes will be included in subsequent versions of the database and updated and described through the online data repository.

7 Citation

This CoralHydro2k Seawater $\delta^{18}\text{O}$ Database descriptor publication should be cited when the database is used in whole or in part, including its metadata fields, for subsequent studies. We encourage end users of this database to also cite the original publications and/or data sources of the underlying primary data (Table 3). To facilitate this process, citation information for every data point is included in the metadata, including a full citation and DOI of the original publication, as well as a dataset citation and DOI for the original public archive of the data. Researchers should also adhere to the data use policies for the underlying data sources (see Appendices A1–A4).



8 Conclusions and anticipated applications of the Seawater $\delta^{18}\text{O}$ Database

Observational seawater $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data can place powerful constraints on the global water cycle, providing valuable information on the exchange of water between the ocean, atmosphere, and cryosphere, as well as on ocean-mixing processes. As such, these data provide an additional degree of freedom for understanding the complex hydrologic system, beyond what standard oceanographic variables like temperature and salinity can offer. They also provide a "common currency" that links paleoclimate reconstructions, modern climate observations, and isotope-enabled model simulations, allowing hydrologic processes to be evaluated on a wide range of time and spatial scales. Given the broad value of this data, and the growing number of seawater $\delta^{18}\text{O}$ and $\delta^2\text{H}$ datasets that have been generated since 2011, the CoralHydro2k Seawater $\delta^{18}\text{O}$ Database was developed to improve the accessibility of seawater isotope data for the Earth Science research community. This new, machine-readable, and metadata-rich database contains over 18,600 observational seawater $\delta^{18}\text{O}$ data points, paired with seawater $\delta^2\text{H}$ and salinity data and extensive metadata that makes the database suitable for a myriad of research applications. The metadata template also provides a set of best practices for reporting seawater isotope data in future studies.

The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database and its extensive metadata can provide insight into the multiple processes that impact seawater $\delta^{18}\text{O}$ and $\delta^2\text{H}$. Furthermore, the database can be used to better constrain the relationship between $\delta^{18}\text{O}_{\text{sw}}$ and salinity in the global ocean, and assess how this relationship varies in space and time. The database also provides updated seawater $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data critical for the calibration and validation of paleoclimate reconstructions using $\delta^{18}\text{O}$ and $\delta^2\text{H}$ to reconstruct past ocean temperature and salinity variations. For example, recent paleoclimate data assimilation efforts would greatly benefit from a spatial network of observational $\delta^{18}\text{O}_{\text{sw}}$ data for training the proxy system models that underlie those efforts. This database could also be used to construct a new gridded dataset of $\delta^{18}\text{O}_{\text{sw}}$ to update that of (LeGrande and Schmidt 2006), which has been widely used for providing climate model boundary conditions and to assess model performance and skill in resolving key features of the hydrologic cycle. In this way, the PAGES CoralHydro2k Seawater $\delta^{18}\text{O}$ Database can be used in a wide variety of applications to bolster our understanding of the modern climate system, while also providing new insights into past and future climate variability and change.

9 Author contribution

AA, KD, AM, TF, SL, SS, and ED designed the database, AA, AM, RP, SL, KD, JH, CM, SS, and AV entered data and/or metadata into the database, AA, AM, SL, RP, JH, CM, and ED performed quality control on the database, AA and KD prepared the manuscript, with contributions from all co-authors, RP developed the example Python code, and JH developed the Github site for the database.

10 Competing interests

The authors declare that they have no conflict of interest.



440 11 Acknowledgements

We gratefully acknowledge the many researchers, funding agencies, and international project teams responsible for the collection, quality control, and publication of the seawater isotope data, including GLODAP and GEOTRACES. We thank the PAGES CoralHydro2k team for their encouragement and effort in building this database, especially the helpful comments and suggestions from Amy Wagner and Hali Kilbourne. We also thank Gilles Reverdin for useful discussions about the database.

445 Many thanks to Erika Ornouski for her work in finding hidden $\delta^{18}\text{O}_{\text{sw}}$ data files in the early stages of the project. We are grateful to Kerstin Lehnert and the EarthChem team at Lamont Doherty Earth Observatory and Carrie Morrill and Edward Gille and Bruce Bauer at NOAA/WDS Paleoclimatology for providing opportunities to host the new database. We also recognize the efforts of Gavin Schmidt, Eelco Rohling, Grant Bigg, and Allegra LeGrande in building and maintaining the first seawater $\delta^{18}\text{O}$ database, as well as Gabriel Bowen at waterisotopes.org and Gilles Reverdin at LOCEAN-IPSL

450 (<https://www.locean.ipsl.fr/?lang=fr>) for their data collection and compiling efforts in continuing to make such data publicly available.

The GEOTRACES 2021 Intermediate Data Product version 2 (IDP2021v2) represents an international collaboration and is endorsed by the Scientific Committee on Oceanic Research (SCOR). The many researchers and funding agencies responsible

455 for the collection of data and quality control are thanked for their contributions to the IDP2021v2. This manuscript also contains data supplied by the Natural Environment Research Council.

PAGES provided funding support for this project through the PAGES Data Stewardship Scholarship (DSS_104 and DSS_114 awarded to A.R.A.). Additional support for this research came from the National Science Foundation awards OCE-1903640 to

460 A.R.A., OCE-2303245 to S.A.M., OCE-2303565 to S.C.S., and NSF-2102931 to K.L.D. and the Department of the Interior South Central Climate Adaptation Science Center Cooperative Agreement G19AC00086, Louisiana Board of Regents LEQSF(2021-22)-ENH-DE-05 to K.L.D. J.A.H and T.F. acknowledge Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 469906366 (T.F.) – SPP 2299/Project number 441832482.

465 References

Ahmed, M., Anchukaitis, K. J., Asrat, A., Borgaonkar, H. P., Braida, M., Buckley, B. M., Büntgen, U., Chase, B. M., Christie, D. A., Cook, E. R., Curran, M. A. J., Diaz, H. F., Esper, J., Fan, Z.-X., Gaire, N. P., Ge, Q., Gergis, J., González-Rouco, J. F., Goosse, H., Grab, S. W., Graham, N., Graham, R., Grosjean, M., Hanhijärvi, S. T., Kaufman, D. S., Kiefer, T., Kimura, K., Korhola, A. A., Krusic, P. J., Lara, A., Lézine, A.-M., Ljungqvist, F. C., Lorrey, A. M., Luterbacher, J., Masson-Delmotte, V.,

470 McCarroll, D., McConnell, J. R., McKay, N. P., Morales, M. S., Moy, A. D., Mulvaney, R., Mundo, I. A., Nakatsuka, T., Nash, D. J., Neukom, R., Nicholson, S. E., Oerter, H., Palmer, J. G., Phipps, S. J., Prieto, M. R., Rivera, A., Sano, M., Severi, M., Shanahan, T. M., Shao, X., Shi, F., Sigl, M., Smerdon, J. E., Solomina, O. N., Steig, E. J., Stenni, B., Thamban, M., Trouet, V., Turney, C. S. M., Umer, M., van Ommen, T., Verschuren, D., Viau, A. E., Villalba, R., Vinther, B. M., von Gunten, L.,



- 475 Wagner, S., Wahl, E. R., Wanner, H., Werner, J. P., White, J. W. C., Yasue, K., Zorita, E., and PAGES 2k Consortium: Continental-scale temperature variability during the past two millennia, *Nature Geoscience*, 6, 339–346, <https://doi.org/10.1038/ngeo1797>, 2013.
- Akhoudas, C. H., Sallée, J.-B., Haumann, F. A., Meredith, M. P., Garabato, A. N., Reverdin, G., Jullion, L., Aloisi, G., Benetti, M., Leng, M. J., and Arrowsmith, C.: Ventilation of the abyss in the Atlantic sector of the Southern Ocean, *Scientific Reports*, 11, 6760, <https://doi.org/10.1038/s41598-021-86043-2>, 2021.
- 480 Atwood, A. R., Moore, A. L., Long, S., Pauly, R., DeLong, K., Wagner, A., and Hargreaves, J. A.: The CoralHydro2k Seawater $\delta^{18}\text{O}$ Database, *PAGES Magazine*, 32, 59, 2024.
- Benetti, M., Reverdin, G., Pierre, C., Khatiwala, S., Tournadre, B., Olafsdottir, S., and Naamar, A.: Variability of sea ice melt and meteoric water input in the surface Labrador Current off Newfoundland, *Journal of Geophysical Research: Oceans*, 121, 2841–2855, <https://doi.org/10.1002/2015JC011302>, 2016.
- 485 Benway, H. M. and Mix, A. C.: Oxygen isotopes, upper-ocean salinity, and precipitation sources in the eastern tropical Pacific, *Earth and Planetary Science Letters*, 224, 493–507, <https://doi.org/10.1016/j.epsl.2004.05.014>, 2004.
- Biddle, L. C., Loose, B., and Heywood, K. J.: Upper Ocean Distribution of Glacial Meltwater in the Amundsen Sea, Antarctica, *Journal of Geophysical Research: Oceans*, 124, 6854–6870, <https://doi.org/10.1029/2019JC015133>, 2019.
- 490 Bigg, G. R. and Rohling, E. J.: An oxygen isotope data set for marine waters, *Journal of Geophysical Research: Oceans*, 105, 8527–8535, <https://doi.org/10.1029/2000JC900005>, 2000.
- Blossey, P. N., Kuang, Z., and Roms, D. M.: Isotopic composition of water in the tropical tropopause layer in cloud-resolving simulations of an idealized tropical circulation, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014554>, 2010.
- 495 Bong, H., Cauquoin, A., Okazaki, A., Chang, E.-C., Werner, M., Wei, Z., Yeo, N., and Yoshimura, K.: Process-Based Intercomparison of Water Isotope-Enabled Models and Reanalysis Nudging Effects, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD038719, <https://doi.org/10.1029/2023JD038719>, 2024.
- Bony, S., Risi, C., and Vimeux, F.: Influence of convective processes on the isotopic composition ($\delta^{18}\text{O}$ and δD) of precipitation and water vapor in the tropics: 1. Radiative-convective equilibrium and Tropical Ocean–Global Atmosphere–Coupled Ocean–Atmosphere Response Experiment (TOGA-COARE) simulations, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009942>, 2008.
- 500 Boutin, J., Reul, N., Koehler, J., Martin, A., Catany, R., Guimbard, S., Rouffi, F., Vergely, J. L., Arias, M., Chakroun, M., Corato, G., Estella-Perez, V., Hasson, A., Josey, S., Khvorostyanov, D., Kolodziejczyk, N., Mignot, J., Olivier, L., Reverdin, G., Stammer, D., Supply, A., Thouvenin-Masson, C., Turiel, A., Vialard, J., Cipollini, P., Donlon, C., Sabia, R., and Mecklenburg, S.: Satellite-Based Sea Surface Salinity Designed for Ocean and Climate Studies, *Journal of Geophysical Research: Oceans*, 126, e2021JC017676, <https://doi.org/10.1029/2021JC017676>, 2021.
- Boyer, T. P., Baranova, O. K., Coleman, C., Garcia, H. E., Grodsky, A., Locarnini, R. A., Mishonov, A. V., Paver, C. R., Reagan, J. R., Seidov, D., Smolyar, I. V., Weathers, K. W., and Zweng, M. M.: World Ocean Database 2018, NOAA Atlas NESDIS, 2018.
- 510 Brady, E., Stevenson, S., Bailey, D., Liu, Z., Noone, D., Nusbaumer, J., Otto-Bliesner, B. L., Tabor, C., Tomas, R., Wong, T., Zhang, J., and Zhu, J.: The Connected Isotopic Water Cycle in the Community Earth System Model Version 1, *Journal of Advances in Modeling Earth Systems*, 11, 2547–2566, <https://doi.org/10.1029/2019MS001663>, 2019.



- Breitkreuz, C., Paul, A., Kurahashi-Nakamura, T., Losch, M., and Schulz, M.: A Dynamical Reconstruction of the Global Monthly Mean Oxygen Isotopic Composition of Seawater, *Journal of Geophysical Research: Oceans*, 123, 7206–7219, <https://doi.org/10.1029/2018JC014300>, 2018.
- 515 Brocas, W. M., Felis, T., and Mudelsee, M.: Tropical Atlantic Cooling and Freshening in the Middle of the Last Interglacial From Coral Proxy Records, *Geophysical Research Letters*, 46, 8289–8299, <https://doi.org/10.1029/2019GL083094>, 2019.
- Cahyarini, S. Y., Pfeiffer, M., Timm, O., Dullo, W.-C., and Schönberg, D. G.: Reconstructing seawater $\delta^{18}\text{O}$ from paired coral $\delta^{18}\text{O}$ and Sr/Ca ratios: Methods, error analysis and problems, with examples from Tahiti (French Polynesia) and Timor (Indonesia), *Geochimica et Cosmochimica Acta*, 72, 2841–2853, <https://doi.org/10.1016/j.gca.2008.04.005>, 2008.
- 520 Carton, J. A., Chepurin, G., Cao, X., and Giese, B.: A Simple Ocean Data Assimilation Analysis of the Global Upper Ocean 1950–95. Part I: Methodology, *Journal of Physical Oceanography*, 30, 294–309, [https://doi.org/10.1175/1520-0485\(2000\)030<0294:ASODAA>2.0.CO;2](https://doi.org/10.1175/1520-0485(2000)030<0294:ASODAA>2.0.CO;2), 2000.
- Carton, J. A., Chepurin, G. A., and Chen, L.: SODA3: A New Ocean Climate Reanalysis, *Journal of Climate*, 31, 6967–6983, <https://doi.org/10.1175/JCLI-D-18-0149.1>, 2018.
- 525 Cauquoin, A., Werner, M., and Lohmann, G.: Water isotopes – climate relationships for the mid-Holocene and preindustrial period simulated with an isotope-enabled version of MPI-ESM, *Climate of the Past*, 15, 1913–1937, <https://doi.org/10.5194/cp-15-1913-2019>, 2019.
- Chamberlain, K. J., Lehnert, K. A., McIntosh, I. M., Morgan, D. J., and Worner, G.: Time to change the data culture in geochemistry, *Nature Reviews Earth & Environment*, 2, 737–739, <https://doi.org/10.1038/s43017-021-00237-w>, 2021.
- 530 Colose, C. M., LeGrande, A. N., and Vuille, M.: Hemispherically asymmetric volcanic forcing of tropical hydroclimate during the last millennium, *Earth System Dynamics*, 7, 681–696, <https://doi.org/10.5194/esd-7-681-2016>, 2016.
- Conroy, J. L., Thompson, D. M., Cobb, K. M., Noone, D., Rea, S., and Legerande, A. N.: Spatiotemporal variability in the $\delta^{18}\text{O}$ -salinity relationship of seawater across the tropical Pacific Ocean, *Paleoceanography*, 32, 484–497, <https://doi.org/10.1002/2016PA003073>, 2017.
- 535 Craig, H. and Gordon, L. I.: Stable Isotopes in Oceanographic Studies and Palaeotemperatures. Deuterium and oxygen-18 variations in the ocean and the marine atmosphere, *Laboratorio di Geologia Nucleare*, 9–130, 1965.
- Dansgaard, W.: The O^{18} -abundance in fresh water, *Geochimica et Cosmochimica Acta*, 6, 241–260, [https://doi.org/10.1016/0016-7037\(54\)90003-4](https://doi.org/10.1016/0016-7037(54)90003-4), 1954.
- Dee, S., Noone, D., Buenning, N., Emile-Geay, J., and Zhou, Y.: SPEEDY-IER: A fast atmospheric GCM with water isotope physics, *Journal of Geophysical Research: Atmospheres*, 120, 73–91, <https://doi.org/10.1002/2014JD022194>, 2015.
- 540 Dee, S., Bailey, A., Conroy, J. L., Atwood, A., Stevenson, S., Nusbaumer, J., and Noone, D.: Water isotopes, climate variability, and the hydrological cycle: recent advances and new frontiers, *Environmental Research: Climate*, 2, 022002, <https://doi.org/10.1088/2752-5295/accbe1>, 2023.
- DeLong, K., Atwood, A., Moore, M., and Sanchez, S.: Clues from the Sea Paint a Picture of Earth’s Water Cycle, *Eos*, 103, 2022.
- 545



- Diekmann, C. J., Schneider, M., Ertl, B., Hase, F., García, O., Khosrawi, F., Sepúlveda, E., Knippertz, P., and Braesicke, P.: The global and multi-annual MUSICA IASI $\{\text{H}_2\text{O}, \delta\text{D}\}$ pair dataset, *Earth System Science Data*, 13, 5273–5292, <https://doi.org/10.5194/essd-13-5273-2021>, 2021.
- 550 Doubleday, Z. A., Martino, J. C., and Trueman, C.: Harnessing universal chemical markers to trace the provenance of marine animals, *Ecological Indicators*, 144, 109481, <https://doi.org/10.1016/j.ecolind.2022.109481>, 2022.
- Durack, P. J., Wijffels, S. E., and Matear, R. J.: Ocean Salinities Reveal Strong Global Water Cycle Intensification During 1950 to 2000, *Science*, 336, 455–458, <https://doi.org/10.1126/science.1212222>, 2012.
- Eglinton, T. I. and Eglinton, G.: Molecular proxies for paleoclimatology, *Earth and Planetary Science Letters*, 275, 1–16, <https://doi.org/10.1016/j.epsl.2008.07.012>, 2008.
- 555 Emile-Geay, J., McKay, N. P., Kaufman, D. S., von Gunten, L., Wang, J., Anchukaitis, K. J., Abram, N. J., Addison, J. A., Curran, M. A. J., Evans, M. N., Henley, B. J., Hao, Z., Martrat, B., McGregor, H. V., Neukom, R., Pederson, G. T., Stenni, B., Thirumalai, K., Werner, J. P., Xu, C., Divine, D. V., Dixon, B. C., Gergis, J., Mundo, I. A., Nakatsuka, T., Phipps, S. J., Routson, C. C., Steig, E. J., Tierney, J. E., Tyler, J. J., Allen, K. J., Bertler, N. A. N., Björklund, J., Chase, B. M., Chen, M.-T., Cook, E., de Jong, R., DeLong, K. L., Dixon, D. A., Ekaykin, A. A., Ersek, V., Filipsson, H. L., Francus, P., Freund, M.
- 560 B., Frezzotti, M., Gaire, N. P., Gajewski, K., Ge, Q., Goosse, H., Gornostaeva, A., Grosjean, M., Horiuchi, K., Hormes, A., Husum, K., Isaksson, E., Kandasamy, S., Kawamura, K., Kilbourne, K. H., Koç, N., Leduc, G., Linderholm, H. W., Lorrey, A. M., Mikhalenko, V., Mortyn, P. G., Motoyama, H., Moy, A. D., Mulvaney, R., Munz, P. M., Nash, D. J., Oerter, H., Opel, T., Orsi, A. J., Ovchinnikov, D. V., Porter, T. J., Roop, H. A., Saenger, C., Sano, M., Sauchyn, D., Saunders, K. M., Seidenkrantz, M.-S., Severi, M., Shao, X., Sicre, M.-A., Sigl, M., Sinclair, K., St. George, S., St. Jacques, J.-M., Thamban,
- 565 M., Kuwar Thapa, U., Thomas, E. R., Turney, C., Uemura, R., Viau, A. E., Vladimirova, D. O., Wahl, E. R., White, J. W. C., Yu, Z., Zinke, J., and PAGES2k Consortium: A global multiproxy database for temperature reconstructions of the Common Era, *Scientific Data*, 4, 170088, <https://doi.org/10.1038/sdata.2017.88>, 2017.
- Evans, M. N., Tolwinski-Ward, S. E., Thompson, D. M., and Anchukaitis, K. J.: Applications of proxy system modeling in high resolution paleoclimatology, *Quaternary Science Reviews*, 76, 16–28, <https://doi.org/10.1016/j.quascirev.2013.05.024>,
 570 2013.
- Felden, J., Möller, L., Schindler, U., Huber, R., Schumacher, S., Koppe, R., Diepenbroek, M., and Glöckner, F. O.: PANGAEA - Data Publisher for Earth & Environmental Science, *Scientific Data*, 10, 347, <https://doi.org/10.1038/s41597-023-02269-x>, 2023.
- Felis, T., Suzuki, A., Kuhnert, H., Dima, M., Lohmann, G., and Kawahata, H.: Subtropical coral reveals abrupt early-twentieth-century freshening in the western North Pacific Ocean, *Geology*, 37, 527–530, <https://doi.org/10.1130/G25581A.1>, 2009.
- 575 Field, R. D., Kim, D., LeGrande, A. N., Worden, J., Kelley, M., and Schmidt, G. A.: Evaluating climate model performance in the tropics with retrievals of water isotopic composition from Aura TES, *Geophysical Research Letters*, 41, 6030–6036, <https://doi.org/10.1002/2014GL060572>, 2014.
- Finkenbiner, C. E., Li, B., Spencer, L., Butler, Z., Haagsma, M., Fiorella, R. P., Allen, S. T., Anderegg, W., Still, C. J., Noone, D., Bowen, G. J., and Good, S. P.: The NEON Daily Isotopic Composition of Environmental Exchanges Dataset, *Scientific Data*, 9, 353, <https://doi.org/10.1038/s41597-022-01412-4>, 2022.
- 580 Fiorella, R. P., Siler, N., Nusbaumer, J., and Noone, D. C.: Enhancing Understanding of the Hydrological Cycle via Pairing of Process-Oriented and Isotope Ratio Tracers, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002648, <https://doi.org/10.1029/2021MS002648>, 2021.



- 585 Frew, R. D., Dennis, P. F., Heywood, K. J., Meredith, M. P., and Boswell, S. M.: The oxygen isotope composition of water masses in the northern North Atlantic, *Deep Sea Research Part I: Oceanographic Research Papers*, 47, 2265–2286, [https://doi.org/10.1016/S0967-0637\(00\)00023-6](https://doi.org/10.1016/S0967-0637(00)00023-6), 2000.
- Gagan, M. K., Ayliffe, L. K., Hopley, D., Cali, J. A., Mortimer, G. E., Chappell, J., McCulloch, M. T., and Head, M. J.: Temperature and Surface-Ocean Water Balance of the Mid-Holocene Tropical Western Pacific, *Science*, 279, 1014–1018, 590 <https://doi.org/10.1126/science.279.5353.1014>, 1998.
- Galewsky, J., Steen-Larsen, H. C., Field, R. D., Worden, J., Risi, C., and Schneider, M.: Stable isotopes in atmospheric water vapor and applications to the hydrologic cycle, *Reviews of Geophysics*, 54, 809–865, <https://doi.org/10.1002/2015RG000512>, 2016.
- Gat, J. R.: OXYGEN AND HYDROGEN ISOTOPES IN THE HYDROLOGIC CYCLE, *Annual Review of Earth and Planetary Sciences*, 24, 225–262, <https://doi.org/10.1146/annurev.earth.24.1.225>, 1996. 595
- GEOTRACES Intermediate Data Product Group: The GEOTRACES Intermediate Data Product 2021 version 2 (IDP2021v2)., <https://doi.org/10.5285/ff46f034-f47c-05f9-e053-6c86abc0dc7e>, 2023.
- Giry, C., Felis, T., Kölling, M., Wei, W., Lohmann, G., and Scheffers, S.: Controls of Caribbean surface hydrology during the mid- to late Holocene: insights from monthly resolved coral records, *Climate of the Past*, 9, 841–858, 600 <https://doi.org/10.5194/cp-9-841-2013>, 2013.
- Gorman, M. K., Quinn, T. M., Taylor, F. W., Partin, J. W., Cabioch, G., Austin Jr., J. A., Pelletier, B., Ballu, V., Maes, C., and Saustrop, S.: A coral-based reconstruction of sea surface salinity at Sabine Bank, Vanuatu from 1842 to 2007 CE, *Paleoceanography*, 27, <https://doi.org/10.1029/2012PA002302>, 2012.
- Gupta, P., Noone, D., Galewsky, J., Sweeney, C., and Vaughn, B. H.: Demonstration of high-precision continuous measurements of water vapor isotopologues in laboratory and remote field deployments using wavelength-scanned cavity ring-down spectroscopy (WS-CRDS) technology, *Rapid Communications in Mass Spectrometry*, 23, 2534–2542, 605 <https://doi.org/10.1002/rcm.4100>, 2009.
- Harbott, M., Wu, H. C., Kuhnert, H., Jimenez, C., González-Díaz, P., and Rixen, T.: A Warming Southern Gulf of Mexico: Reconstruction of Anthropogenic Environmental Changes From a *Siderastrea siderea* Coral on the Northern Coast of Cuba, *Paleoceanography and Paleoclimatology*, 38, e2023PA004717, <https://doi.org/10.1029/2023PA004717>, 2023. 610
- Hargreaves, J., DeLong, K., Felis, T., Abram, N., Cobb, K., and Sayani, H.: Tropical ocean hydroclimate and temperature from coral archives, *PAGES Magazine*, 28, 2020.
- Henze, D., Noone, D., and Toohey, D.: Aircraft measurements of water vapor heavy isotope ratios in the marine boundary layer and lower troposphere during ORACLES, *Earth System Science Data*, 14, 1811–1829, <https://doi.org/10.5194/essd-14-1811-2022>, 2022. 615
- Hereid, K. A., Quinn, T. M., and Okumura, Y. M.: Assessing spatial variability in El Niño–Southern Oscillation event detection skill using coral geochemistry, *Paleoceanography*, 28, 14–23, <https://doi.org/10.1029/2012PA002352>, 2013.
- Hetzinger, S., Pfeiffer, M., Dullo, W.-C., Ruprecht, E., and Garbe-Schönberg, D.: Sr/Ca and $\delta^{18}\text{O}$ in a fast-growing *Diploria strigosa* coral: Evaluation of a new climate archive for the tropical Atlantic, *Geochemistry, Geophysics, Geosystems*, 7, 620 <https://doi.org/10.1029/2006GC001347>, 2006.



Imbrie, J., Hays, J. D., Martinson, D. G., McIntyre, A., Mix, A. C., Morley, J. J., Pisias, N. G., Prell, W. L., and Shackleton, N. J.: The Orbital Theory of Pleistocene Climate: Support from a Revised Chronology, of the Marine $\delta^{18}\text{O}$ Record, in: Berger, A. Ed., *Milankovitch and Climate, Part 1*, Springer, New York, 269–305, 1984.

625 Jacobs, S. S., Fairbanks, R. G., Horibe, Y., and Jacobs, S. S.: Origin and evolution of water masses near the Antarctic continental margin: Evidence from $\text{H}_2^{18}\text{O}/\text{H}_2^{16}\text{O}$ ratios in seawater, vol. 43, American Geophysical Union, Washington, D. C., 59–85, <https://doi.org/10.1029/ar043p0059>, 1985.

Jannasch, H. W., Wheat, C. G., Plant, J. N., Kastner, M., and Stakes, D. S.: Continuous chemical monitoring with osmotically pumped water samplers: OsmoSampler design and applications, *Limnology and Oceanography: Methods*, 2, 102–113, <https://doi.org/10.4319/lom.2004.2.102>, 2004.

630 Kaufman, D., McKay, N., Routson, C., Erb, M., Dätwyler, C., Sommer, P. S., Heiri, O., and Davis, B.: Holocene global mean surface temperature, a multi-method reconstruction approach, *Scientific Data*, 7, 201, <https://doi.org/10.1038/s41597-020-0530-7>, 2020.

635 Key, R. M., Olsen, A., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterstrom, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., and Suzuki, T.: Global Ocean Data Analysis Project, Version 2 (GLODAPv2), ORNL/CDIAC-162, NDP-093. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, https://doi.org/10.3334/CDIAC/OTG.NDP093_GLODAPv2, 2023.

640 Khare, A., Hughes, H. P., Kilbourne, K. H., and Schijf, J.: An ICP-AES method for routine high-precision measurement of seawater Sr/Ca ratios to validate coral paleothermometry calibrations, *Limnology and Oceanography: Methods*, 19, 416–430, <https://doi.org/10.1002/lom3.10434>, 2021.

Kilbourne, K. H., Quinn, T. M., Taylor, F. W., Delcroix, T., and Gouriou, Y.: El Niño–Southern Oscillation–related salinity variations recorded in the skeletal geochemistry of a *Porites* coral from Espiritu Santo, Vanuatu, *Paleoceanography*, 19, <https://doi.org/10.1029/2004PA001033>, 2004.

645 Knebel, O., Felis, T., Asami, R., Deschamps, P., Kölling, M., and Scholz, D.: Last Deglacial Environmental Change in the Tropical South Pacific From Tahiti Corals, *Paleoceanography and Paleoclimatology*, 39, e2022PA004585, <https://doi.org/10.1029/2022PA004585>, 2024.

650 Konecky, B. L., McKay, N. P., Churakova (Sidorova), O. V., Comas-Bru, L., Dassié, E. P., DeLong, K. L., Falster, G. M., Fischer, M. J., Jones, M. D., Jonkers, L., Kaufman, D. S., Leduc, G., Managave, S. R., Martrat, B., Opel, T., Orsi, A. J., Partin, J. W., Sayani, H. R., Thomas, E. K., Thompson, D. M., Tyler, J. J., Abram, N. J., Atwood, A. R., Cartapanis, O., Conroy, J. L., Curran, M. A., Dee, S. G., Deininger, M., Divine, D. V., Kern, Z., Porter, T. J., Stevenson, S. L., von Gunten, L., and Members, I. P.: The Iso2k database: a global compilation of paleo- $\delta^{18}\text{O}$ and $\delta^2\text{H}$ records to aid understanding of Common Era climate, *Earth System Science Data*, 12, 2261–2288, <https://doi.org/10.5194/essd-12-2261-2020>, 2020.

655 Konecky, B. L., McKay, N. P., Falster, G. M., Stevenson, S. L., Fischer, M. J., Atwood, A. R., Thompson, D. M., Jones, M. D., Tyler, J. J., DeLong, K. L., Martrat, B., Thomas, E. K., Conroy, J. L., Dee, S. G., Jonkers, L., Churakova (Sidorova), O. V., Kern, Z., Opel, T., Porter, T. J., Sayani, H. R., Skrzypek, G., Abram, N. J., Braun, K., Carré, M., Cartapanis, O., Comas-Bru, L., Curran, M. A., Dassié, E. P., Deininger, M., Divine, D. V., Incarbona, A., Kaufman, D. S., Kaushal, N., Kläbe, R. M., Kolus, H. R., Leduc, G., Managave, S. R., Mortyn, P. G., Moy, A. D., Orsi, A. J., Partin, J. W., Roop, H. A., Sicre, M.-A., von Gunten, L., Yoshimura, K., and Iso2k Project Members: Globally coherent water cycle response to temperature change during the past two millennia, *Nature Geoscience*, 16, 997–1004, <https://doi.org/10.1038/s41561-023-01291-3>, 2023.



- Kurita, N., Noone, D., Risi, C., Schmidt, G. A., Yamada, H., and Yoneyama, K.: Intraseasonal isotopic variation associated with the Madden-Julian Oscillation, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2010JD015209>, 2011.
- 665 Lee, J. and Fung, I.: “Amount effect” of water isotopes and quantitative analysis of post-condensation processes, *HYDROLOGICAL PROCESSES*, 22, 1–8, <https://doi.org/10.1002/hyp.6637>, 2008.
- LeGrande, A. N. and Schmidt, G. A.: Global gridded data set of the oxygen isotopic composition in seawater, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026011>, 2006.
- LeGrande, A. N. and Schmidt, G. A.: Water isotopologues as a quantitative paleosalinity proxy, *Paleoceanography*, 26, <https://doi.org/10.1029/2010PA002043>, 2011.
- 670 Light, T., Williams, B., Halfar, J., Hou, A., Zajacz, Z., Tsay, A., and Adey, W.: Advancing Mg/Ca Analysis of Coralline Algae as a Climate Proxy by Assessing LA-ICP-OES Sampling and Coupled Mg/Ca- $\delta^{18}\text{O}$ Analysis, *Geochemistry, Geophysics, Geosystems*, 19, 2876–2894, <https://doi.org/10.1029/2018GC007504>, 2018.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, <https://doi.org/10.1029/2004PA001071>, 2005.
- 675 McCulloch, M. T., Gagan, M. K., Mortimer, G. E., Chivas, A. R., and Isdale, P. J.: A high-resolution Sr/Ca and $\delta^{18}\text{O}$ coral record from the Great Barrier Reef, Australia, and the 1982–1983 El Niño, *Geochimica et Cosmochimica Acta*, 58, 2747–2754, [https://doi.org/10.1016/0016-7037\(94\)90142-2](https://doi.org/10.1016/0016-7037(94)90142-2), 1994.
- McKay, N. P. and Kaufman, D. S.: An extended Arctic proxy temperature database for the past 2,000 years, *Scientific Data*, 1, 140026, <https://doi.org/10.1038/sdata.2014.26>, 2014.
- 680 Meredith, M. P., Grose, K. E., McDonagh, E. L., Heywood, K. J., Frew, R. D., and Dennis, P. F.: Distribution of oxygen isotopes in the water masses of Drake Passage and the South Atlantic, *Journal of Geophysical Research: Oceans*, 104, 20949–20962, <https://doi.org/10.1029/98JC02544>, 1999.
- Noone, D. and Simmonds, I.: Associations between $\delta^{18}\text{O}$ of Water and Climate Parameters in a Simulation of Atmospheric Circulation for 1979–95, *Journal of Climate*, 15, 3150–3169, [https://doi.org/10.1175/1520-0442\(2002\)015<3150:ABOOWA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<3150:ABOOWA>2.0.CO;2), 2002.
- 685 Nusbaumer, J., Wong, T. E., Bardeen, C., and Noone, D.: Evaluating hydrological processes in the Community Atmosphere Model Version 5 (CAM5) using stable isotope ratios of water, *Journal of Advances in Modeling Earth Systems*, 9, 949–977, <https://doi.org/10.1002/2016MS000839>, 2017.
- O’Connor, G. K., Cobb, K. M., Sayani, H. R., Atwood, A. R., Grothe, P. R., Stevenson, S., Baum, J. K., Chen, T., Claar, D. C., Hitt, N. T., Lynch-Stieglitz, J., Mortlock, R. A., Schmidt, G. A., and Walter, R.: Coral Oxygen Isotopic Records Capture the 2015/2016 El Niño Event in the Central Equatorial Pacific, *Geophysical Research Letters*, 48, e2021GL094036, <https://doi.org/10.1029/2021GL094036>, 2021.
- 695 Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) – an internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 2016.



- PAGES 2k Network Coordinators: Understanding the climate of the past 2000 years: Phase 3 of the PAGES 2k Network, *PAGES Magazine*, 25, 110–110, 2017.
- PAGES Hydro2k Consortium: Comparing proxy and model estimates of hydroclimate variability and change over the Common Era, *Climate of the Past*, 13, 1851–1900, <https://doi.org/10.5194/cp-13-1851-2017>, 2017.
- Reed, E. V., Thompson, D. M., and Anchukaitis, K. J.: Coral-Based Sea Surface Salinity Reconstructions and the Role of Observational Uncertainties in Inferred Variability and Trends, *Paleoceanography and Paleoclimatology*, 37, e2021PA004371, <https://doi.org/10.1029/2021PA004371>, 2022.
- Ren, L., Linsley, B. K., Wellington, G. M., Schrag, D. P., and Hoegh-guldberg, O.: Deconvolving the $\delta^{18}\text{O}$ seawater component from subseasonal coral $\delta^{18}\text{O}$ and Sr/Ca at Rarotonga in the southwestern subtropical Pacific for the period 1726 to 1997, *Geochimica et Cosmochimica Acta*, 67, 1609–1621, [https://doi.org/10.1016/S0016-7037\(02\)00917-1](https://doi.org/10.1016/S0016-7037(02)00917-1), 2003.
- Reverdin, G., Waelbroeck, C., Pierre, C., Akhondas, C., Aloisi, G., Benetti, M., Bourlès, B., Danielsen, M., Demange, J., Diverres, D., Gascard, J.-C., Houssais, M.-N., Le Goff, H., Lherminier, P., Lo Monaco, C., Mercier, H., Metzl, N., Morisset, S., Naamar, A., Reynaud, T., Sallée, J.-B., Thierry, V., Hartman, S. E., Mawji, E. W., Olafsdottir, S., Kanzow, T., Velo, A., Voelker, A., Yashayaev, I., Haumann, F. A., Leng, M. J., Arrowsmith, C., and Meredith, M.: The CISE-LOCEAN seawater isotopic database (1998–2021), *Earth System Science Data*, 14, 2721–2735, <https://doi.org/10.5194/essd-14-2721-2022>, 2022.
- Risi, C., Bony, S., Vimeux, F., and Jouzel, J.: Water-stable isotopes in the LMDZ4 general circulation model: Model evaluation for present-day and past climates and applications to climatic interpretations of tropical isotopic records, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013255>, 2010.
- Risi, C., Muller, C., and Blossey, P.: What Controls the Water Vapor Isotopic Composition Near the Surface of Tropical Oceans? Results From an Analytical Model Constrained by Large-Eddy Simulations, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002106, <https://doi.org/10.1029/2020MS002106>, 2020.
- Risi, C., Muller, C., and Blossey, P.: Rain Evaporation, Snow Melt, and Entrainment at the Heart of Water Vapor Isotopic Variations in the Tropical Troposphere, According to Large-Eddy Simulations and a Two-Column Model, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002381, <https://doi.org/10.1029/2020MS002381>, 2021.
- Sanchez, S. C., Hakim, G. J., and Saenger, C. P.: Climate Model Teleconnection Patterns Govern the Niño-3.4 Response to Early Nineteenth-Century Volcanism in Coral-Based Data Assimilation Reconstructions, *Journal of Climate*, 34, 1863–1880, <https://doi.org/10.1175/JCLI-D-20-0549.1>, 2021.
- Schmidt, G. A.: Forward modeling of carbonate proxy data from planktonic foraminifera using oxygen isotope tracers in a global ocean model, *Paleoceanography*, 14, 482–497, <https://doi.org/10.1029/1999PA900025>, 1999.
- Schmidt, G. A., Bigg, G. R., and Rohling, E. J.: Global Seawater Oxygen-18 Database - v1.22, 1999.
- Schmidt, G. A., LeGrande, A. N., and Hoffmann, G.: Water isotope expressions of intrinsic and forced variability in a coupled ocean-atmosphere model, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007781>, 2007.
- Schmidt, M. W. and Lynch-Stieglitz, J.: Florida Straits deglacial temperature and salinity change: Implications for tropical hydrologic cycle variability during the Younger Dryas, *Paleoceanography*, 26, <https://doi.org/10.1029/2011PA002157>, 2011.
- Schneider, M., Ertl, B., Diekmann, C. J., Khosrawi, F., Weber, A., Hase, F., Höpfner, M., García, O. E., Sepúlveda, E., and Kinnison, D.: Design and description of the MUSICA IASI full retrieval product, *Earth System Science Data*, 14, 709–742, <https://doi.org/10.5194/essd-14-709-2022>, 2022.



- 735 Smerdon, J. E.: Climate models as a test bed for climate reconstruction methods: pseudoproxy experiments, *WIREs Climate Change*, 3, 63–77, <https://doi.org/10.1002/wcc.149>, 2012.
- Stevenson, S., Powell, B. S., Merrifield, M. A., Cobb, K. M., Nusbaumer, J., and Noone, D.: Characterizing seawater oxygen isotopic variability in a regional ocean modeling framework: Implications for coral proxy records, *Paleoceanography*, 30, 1573–1593, <https://doi.org/10.1002/2015PA002824>, 2015.
- 740 Stevenson, S., Powell, B., Cobb, K. M., Nusbaumer, J., Merrifield, M., and Noone, D.: Twentieth Century Seawater $\delta^{18}\text{O}$ Dynamics and Implications for Coral-Based Climate Reconstruction, *Paleoceanography and Paleoclimatology*, 33, 606–625, <https://doi.org/10.1029/2017PA003304>, 2018.
- Stevenson, S., Cobb, K. M., Merrifield, M., Powell, B., Sanchez, S., Nusbaumer, J., O'Connor, G., and Atwood, A.: Contrasting Central Equatorial Pacific Oxygen Isotopic Signatures of the 2014/2015 and 2015/2016 El Niño Events, *Geophysical Research Letters*, 50, e2023GL104454, <https://doi.org/10.1029/2023GL104454>, 2023.
- 745 Stott, L., Cannariato, K., Thunell, R., Haug, G. H., Koutavas, A., and Lund, S.: Decline of surface temperature and salinity in the western tropical Pacific Ocean in the Holocene epoch, *Nature*, 431, 56–59, <https://doi.org/10.1038/nature02903>, 2004.
- Strain, P. M. and Tan, F. C.: Seasonal evolution of oxygen isotope-salinity relationships in high-latitude surface waters, *Journal of Geophysical Research: Oceans*, 98, 14589–14598, <https://doi.org/10.1029/93JC01182>, 1993.
- 750 Tada, M., Yoshimura, K., and Toride, K.: Improving weather forecasting by assimilation of water vapor isotopes, *Scientific Reports*, 11, 18067, <https://doi.org/10.1038/s41598-021-97476-0>, 2021.
- Tardif, R., Hakim, G. J., Perkins, W. A., Horlick, K. A., Erb, M. P., Emile-Geay, J., Anderson, D. M., Steig, E. J., and Noone, D.: Last Millennium Reanalysis with an expanded proxy database and seasonal proxy modeling, *Climate of the Past*, 15, 1251–1273, <https://doi.org/10.5194/cp-15-1251-2019>, 2019.
- 755 Thompson, D. M., Ault, T. R., Evans, M. N., Cole, J. E., and Emile-Geay, J.: Comparison of observed and simulated tropical climate trends using a forward model of coral $\delta^{18}\text{O}$, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL048224>, 2011.
- Thompson, D. M., Conroy, J. L., Konecky, B. L., Stevenson, S., DeLong, K. L., McKay, N., Reed, E., Jonkers, L., and Carré, M.: Identifying Hydro-Sensitive Coral $\delta^{18}\text{O}$ Records for Improved High-Resolution Temperature and Salinity Reconstructions, *Geophysical Research Letters*, 49, e2021GL096153, <https://doi.org/10.1029/2021GL096153>, 2022.
- 760 Tierney, J. E., Abram, N. J., Anchukaitis, K. J., Evans, M. N., Giry, C., Kilbourne, K. H., Saenger, C. P., Wu, H. C., and Zinke, J.: Tropical sea surface temperatures for the past four centuries reconstructed from coral archives, *Paleoceanography*, 30, 226–252, <https://doi.org/10.1002/2014PA002717>, 2015.
- 765 Trofimova, T., Alexandroff, S. J., Mette, M. J., Tray, E., Butler, P. G., Campana, S. E., Harper, E. M., Johnson, A. L. A., Morrongiello, J. R., Peharda, M., Schöne, B. R., Andersson, C., Andrus, C. F. T., Black, B. A., Burchell, M., Carroll, M. L., DeLong, K. L., Gillanders, B. M., Grønkjær, P., Killam, D., Prendergast, A. L., Reynolds, D. J., Scourse, J. D., Shirai, K., Thébault, J., Trueman, C., and de Winter, N.: Fundamental questions and applications of sclerochronology: Community-defined research priorities, *Estuarine, Coastal and Shelf Science*, 245, 106977, <https://doi.org/10.1016/j.ecss.2020.106977>, 2020.
- 770 Wagner, A. J. and Slowey, N. C.: Oxygen isotopes in seawater from the Texas-Louisiana Shelf, *Bulletin of marine science*, 87, 1–12, <https://doi.org/10.5343/bms.2010.1004>, 2011.



- Walter, R. M., Sayani, H. R., Felis, T., Cobb, K. M., Abram, N. J., Arzey, A. K., Atwood, A., Brenner, L. D., Dassie, E. P., DeLong, K. L., Ellis, B., Fischer, M. J., Goodkin, N. F., Hargreaves, J. A., Kilbourne, K. H., Krawczyk, H. A., McKay, N. P., Murty, S. A., Ramos, R. D., Reed, E. V., Samata, D., Sanchez, S. C., Zinke, J., and PAGES CoralHydro2k Project Members: NOAA/WDS Paleoclimatology - CoralHydro2k Database (Common Era coral $\delta^{18}\text{O}$ and Sr/Ca data compilation), NOAA National Centers for Environmental Information, 2022.
- Walter, R. M., Sayani, H. R., Felis, T., Cobb, K. M., Abram, N. J., Arzey, A. K., Atwood, A. R., Brenner, L. D., Dassié, É. P., DeLong, K. L., Ellis, B., Emile-Geay, J., Fischer, M. J., Goodkin, N. F., Hargreaves, J. A., Kilbourne, K. H., Krawczyk, H., McKay, N. P., Moore, A. L., Murty, S. A., Ong, M. R., Ramos, R. D., Reed, E. V., Samanta, D., Sanchez, S. C., Zinke, J., and Members, the P. C. P.: The CoralHydro2k database: a global, actively curated compilation of coral $\delta^{18}\text{O}$ and Sr/Ca proxy records of tropical ocean hydrology and temperature for the Common Era, *Earth System Science Data*, 15, 2081–2116, <https://doi.org/10.5194/essd-15-2081-2023>, 2023.
- Warner, J. P., DeLong, K. L., Chicoine, D., Thirumalai, K., and Andrus, C. F. T.: Investigating the influence of temperature and seawater $\delta^{18}\text{O}$ on *Donax obesulus* (Reeve, 1854) shell $\delta^{18}\text{O}$, *Chemical Geology*, 588, 120638, <https://doi.org/10.1016/j.chemgeo.2021.120638>, 2022.
- waterisotopes-CISE-LOCEAN: Water isotopes of sea water analyzed since 1998 at LOCEAN, SEANO, 2024.
- Wei, Z., Lee, X., and Patton, E. G.: ISOLESC: A Coupled Isotope-LSM-LES-Cloud Modeling System to Investigate the Water Budget in the Atmospheric Boundary Layer, *Journal of Advances in Modeling Earth Systems*, 10, 2589–2617, <https://doi.org/10.1029/2018MS001381>, 2018.
- Werner, M., Langebroek, P. M., Carlsen, T., Herold, M., and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: Toward high-resolution isotope modeling on a global scale, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD015681>, 2011.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 160018, <https://doi.org/10.1038/sdata.2016.18>, 2016.
- Wong, A. P. S., Wijffels, S. E., Riser, S. C., Pouliquen, S., Hosoda, S., Roemmich, D., Gilson, J., Johnson, G. C., Martini, K., Murphy, D. J., Scanderbeg, M., Bhaskar, T. V. S. U., Buck, J. J. H., Merceur, F., Carval, T., Maze, G., Cabanes, C., André, X., Poffa, N., Yashayaev, I., Barker, P. M., Guinehut, S., Belbéoch, M., Ignaszewski, M., Baringer, M. O., Schmid, C., Lyman, J. M., McTaggart, K. E., Purkey, S. G., Zilberman, N., Alkire, M. B., Swift, D., Owens, W. B., Jayne, S. R., Hersch, C., Robbins, P., West-Mack, D., Bahr, F., Yoshida, S., Sutton, P. J. H., Cancouët, R., Coatanoan, C., Dobbler, D., Juan, A. G., Gourrion, J., Kolodziejczyk, N., Bernard, V., Bourlès, B., Claustre, H., D'Ortenzio, F., Le Reste, S., Le Traon, P.-Y., Rannou, J.-P., Saout-Grit, C., Speich, S., Thierry, V., Verbrugge, N., Angel-Benavides, I. M., Klein, B., Notarstefano, G., Poulain, P.-M., Vélez-Belchí, P., Suga, T., Ando, K., Iwasaka, N., Kobayashi, T., Masuda, S., Oka, E., Sato, K., Nakamura, T., Sato, K., Takatsuki, Y., Yoshida, T., Cowley, R., Lovell, J. L., Oke, P. R., van Wijk, E. M., Carse, F., Donnelly, M., Gould, W. J., Gowers, K., King, B. A., Loch, S. G., Mowat, M., Turton, J., Rama Rao, E. P., Ravichandran, M., Freeland, H. J., Gaboury, I., Gilbert, D., Greenan, B. J. W., Ouellet, M., Ross, T., Tran, A., Dong, M., Liu, Z., Xu, J., Kang, K., Jo, H., et al.: Argo Data 1999–2019: Two Million Temperature-Salinity Profiles and Subsurface Velocity Observations From a Global Array of Profiling Floats, *Frontiers in Marine Science*, Volume 7-2020, <https://doi.org/10.3389/fmars.2020.00700>, 2020.



- 815 Worden, J. R., Kulawik, S. S., Fu, D., Payne, V. H., Lipton, A. E., Polonsky, I., He, Y., Cady-Pereira, K., Moncet, J.-L.,
Herman, R. L., Irion, F. W., and Bowman, K. W.: Characterization and evaluation of AIRS-based estimates of the deuterium
content of water vapor, *Atmospheric Measurement Techniques*, 12, 2331–2339, <https://doi.org/10.5194/amt-12-2331-2019>,
2019.
- Wu, H. C., Linsley, B. K., Dassié, E. P., Schiraldi, B., and deMenocal, P. B.: Oceanographic variability in the South Pacific
Convergence Zone region over the last 210 years from multi-site coral Sr/Ca records, *Geochemistry, Geophysics, Geosystems*,
14, 1435–1453, <https://doi.org/10.1029/2012GC004293>, 2013.
- 820 Yoshimura, K., Kanamitsu, M., Noone, D., and Oki, T.: Historical isotope simulation using Reanalysis atmospheric data,
Journal of Geophysical Research: Atmospheres, 113, <https://doi.org/10.1029/2008JD010074>, 2008.