

History of anthropogenic Phosphorus inputs (HaPi) to the terrestrial biosphere from 1860 to 2020

Zihao Bian^{1,2*}, Hao Shi^{3*}, Rui Li⁴, Fei Lun⁵, Francesco N. Tubiello⁶, Nathaniel D. Mueller^{7,8}, Shiyu You³, Rong Hao^{1,2}, Jiageng Ma³, Longhui Li^{1,2}, Changchun Huang^{1,2}, Bing He⁹, Yuanzhi Yao⁴, Hanqin Tian^{10,11*}

¹State Key Laboratory of Climate System Prediction and Risk Management, Nanjing Normal University, Nanjing 210023, China.

²School of Geography, Nanjing Normal University, Nanjing 210023, China

³State Key Laboratory for Ecological Security of Regions and Cities, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

⁴Key Laboratory of Geographic Information Science of the Ministry of Education, School of Geographic Sciences, East China Normal University, Shanghai 200241, China

⁵ College of Land Science and Technology, China Agricultural University, Beijing 100193, China

⁶Statistics Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome 00153, Italy

⁷Department of Ecosystem Science and Sustainability, Colorado State University, Fort Collins, CO 80523, USA

⁸Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO 80523, USA

⁹School of Computer Science, Chengdu University of Information Technology, Chengdu 610225, China

¹⁰Center for Earth System Science and Global Sustainability, Schiller Institute for Integrated Science and Society, Boston College, Chestnut Hill, MA 02467, USA

¹¹Department of Earth and Environmental Sciences, Boston College, Chestnut Hill, MA 02467, USA

**Corresponding authors:*

Zihao Bian (zhbian@njnu.edu.cn)

Hao Shi (haoshi@rcees.ac.cn)

Hanqin Tian (hanqin.tian@bc.edu)

Abstract

Nitrogen and Phosphorus (P) are essential nutrients for sustaining life on Earth and have been increasingly applied in global agriculture to meet the growing demand for food production. Quantifying the spatial and temporal dynamics of nutrient inputs to the terrestrial biosphere is crucial for analyzing nutrient flows in crop-livestock systems, managing nutrient resources sustainably, and mitigating nutrient-related environmental impacts. Here, built upon our previous work mapping global nitrogen inputs (History of anthropogenic Nitrogen inputs, HaNi), this study presents the History of anthropogenic P inputs (HaPi) dataset, a comprehensive quantification of human-driven P fluxes to terrestrial ecosystems. HaPi covers the period from 1860 to 2020 and has a spatial resolution of 5 arc-minutes (about 10 km at the equator) with an annual time-step. This harmonized dataset integrates seven components, including P fertilizer application on croplands and pastures, manure P application on croplands and pastures, manure P deposition on pastures and rangelands, and atmospheric P deposition. The results reveal that the global total P input increased from 3.8 Tg yr⁻¹ in the 1860s to 41.0 Tg yr⁻¹ in the 2010s, with mineral fertilizer and livestock manure contributing equally to the increase. Regional patterns have shifted significantly over the study period, with China, South Asia, and Brazil surpassing Europe and the USA as the regions with the highest P inputs in recent decades. Furthermore, mineral fertilizers dominate P inputs in most industrialized countries in the Northern Hemisphere, whereas manure P remains the primary source in many countries of the Southern Hemisphere. The HaPi dataset improves P mass budget calculations and provides essential forcing data for empirical or mechanistic models, supporting critical research in agricultural nutrient management, water quality control, and assessments of the coupled human-Earth system. The dataset is available at <https://doi.org/10.6084/m9.figshare.29930279> (Bian et al., 2026).

1 Introduction

The global nitrogen and phosphorus (P) cycle has been unprecedentedly disturbed in the Anthropocene, posing significant risks to planetary boundaries and societal sustainability (Steffen et al., 2015). Driven by rising food demand, mining of phosphate rocks for mineral P fertilizers and livestock feed has significantly increased (Cordell et al., 2009). Anthropogenic activities have substantially intensified global P flows, contributing to half of global agricultural soil P uptake in recent decades (Demay et al., 2023) and tripling the P mobilization in the land-water continuum (Yuan et al., 2018). However, agriculture's growing demand for mineral P fertilizers raises critical concerns regarding the depletion of non-renewable phosphate rock reserves. Moreover, the unevenly distributed phosphate rock resources and potential geopolitical conflicts threaten the resilience of agricultural systems (Barbieri et al., 2022; Elser and Bennett, 2011). To mitigate these challenges, a better understanding of the spatial and temporal distribution of P supply and demand would support improved assessment of future P requirements for crop and livestock production (Sattari et al., 2012).

The historical and current management of anthropogenic P in croplands and livestock production has resulted in serious environmental issues. In particular, the global P flux from terrestrial to aquatic ecosystems has been amplified by inefficient agricultural applications, which accelerated P losses to rivers, lakes, and oceans via water transport. Generally, P is the major limiting nutrient for phytoplankton growth in freshwaters, since its easy adsorption to particles reduces its biological availability (Conley et al., 2009). However, worldwide mineral P fertilizer applications and related runoff have resulted in elevated levels of bioavailable P in freshwaters, causing widespread eutrophication and ecological damage to freshwater systems (Carpenter, 2005; Fink et al., 2018). Beyond contemporary P inputs, legacy P accumulated in soils from historical fertilizer and manure applications continues to leach into aquatic environments. This reduces the efficacy of long-term conservation efforts aimed at improving water quality (Stackpoole et al., 2019). Spatial-temporal patterns of agricultural legacy P and

associated water pollution are fundamentally driven by historical fertilizer and manure applications. Consequently, understanding the quantity, sources, and trends of P pollution is useful information needed for advancing effective water quality management strategies.

To address the challenges of food security and environmental issues related to anthropogenic P use, accurately quantifying global anthropogenic P inputs since the pre-industrial era is critical for evaluating historical trajectories of soil P fertility, P use efficiency (PUE), and P pollution. Many efforts have been carried out to develop global P input datasets (Lu and Tian, 2017; Lun et al., 2018; Ringeval et al., 2024). The Food and Agriculture Organization (FAO) and the International Fertilizer Association (IFA) provide widely adopted country-level mineral P fertilizer data in agricultural land (FAO, 2024). This data has been used to evaluate P budgets and surpluses at the country level. For example, Zou et al. (2022) used FAO P fertilizer data to evaluate national P budget and PUE. However, FAO's manure data only covers national nitrogen excretion, necessitating estimation of manure P via N:P ratios. For spatially explicit modeling, several gridded datasets have emerged. Mueller et al. (2012) generated crop-specific P fertilizer maps at a global scale circa 2000. Bouwman et al. (2013) developed gridded P fertilizer and livestock manure data by combining FAO fertilizer use, animal production data, and multiple ancillary datasets. This P input dataset has been applied in the IMAGE-Global Nutrient Model to assess global soil P pools, crop uptake, and riverine P fluxes (Beusen et al., 2016). More recently, gridded data of P application rates for 173 crops, at 5 km resolution and the year 2020, have been published (Nguyen et al., 2024). Furthermore, Lu and Tian (2017) developed an annual global P fertilizer dataset from 1961 to 2013 at a resolution of $0.5^{\circ} \times 0.5^{\circ}$, based on the IFA national P fertilizer statistics. Bouwman's dataset and Lu's P fertilizer data have been used to develop the global dataset on P in agricultural soil, GPASOIL-v0 (Ringeval et al., 2017) and GPASOIL-v1 (Ringeval et al., 2024), respectively. GPASOIL-v1 also provides manure P application rates on cropland and pasture based on a global constant N:P ratio and spatially explicit manure N data developed by Xu et al. (2018) and Zhang et al.

(2017). Nevertheless, inconsistencies exist in temporal coverage, spatial resolution, spatial allocation algorithms, and the baseline land use data among different P input datasets, which may propagate uncertainties in assessments of soil P fertility, legacy P accumulation and its environmental impacts. To resolve these issues, we propose to reconstruct a harmonized History of Anthropogenic Phosphorus Inputs (HaPi) dataset, which will integrate available FAO statistics, historical land use, grid-level manure N:P ratios, and atmospheric P deposition within a consistent spatiotemporal framework.

In our recent study, we have developed the History of anthropogenic Nitrogen inputs (HaNi) dataset, which consists of N fertilizer, manure N, and atmospheric N inputs on cropland, pasture, and rangeland (i.e., grassland) from 1860 to 2020 (Tian et al., 2022). Since its publication, HaNi has been widely adopted to estimate global N₂O emission (Tian et al., 2024), NH₃ concentrations (Ma et al., 2025) and emissions, and N loading (Dai et al., 2023). Complementary to the HaNi data, the HaPi dataset provides P fertilizer/manure application to cropland, P fertilizer/manure application to pasture, manure P deposition on pasture/rangeland, and atmospheric P deposition. The HaPi dataset advances over previous datasets in five aspects: (1) the impact of dynamic crop rotation is considered when allocating P fertilizer on cropland; (2) the annual spatial-explicit manure N:P ratio data is developed to generate manure P data based on manure N inputs; (3) all P inputs are at an annual step from 1860 to 2020 and a high spatial resolution of 5 arc-min; (4) beyond agricultural land, P inputs to all terrestrial ecosystems are estimated; and (5) consistent baseline land use maps are used when allocating fertilizer and manure P to cropland, pasture, and rangeland. The HaPi dataset is anticipated to improve P mass budget calculation and serve as forcing data for empirical or mechanistic models, thereby supporting diverse research on soil fertility, water pollution, P resource sustainability, and food security, etc.

2 Methods

2.1 Fertilizer application on cropland and pasture

The fertilizer P application maps are developed by allocating country-level inventory data into grid cells according to land use maps, crop rotation maps, and crop-specific fertilizer application data (Table 1). Annual national P fertilizer inputs on agricultural land during 1961-2020 were obtained from FAOSTAT (2024). We first separated total P fertilizer use into P application on cropland and pasture by assuming fractions of P used for cropland and pasture are the same as those of N fertilizer, following the method used in Zou et al. (2022). The annual fractions of fertilizer use allocated to cropland and pasture for major countries from 1961 to 2019 were adopted from Einarsson et al. (2021) and Lassaletta et al. (2014). Then, the country-level P fertilizer data were spatially distributed with cropland and pasture maps following the workflow in Figure 1.

To determine the P fertilizer application on cropland, we considered 17 dominant crop types (wheat, maize, rice, barley, millet, sorghum, soybean, sunflower, potato, cassava, sugarcane, sugar beet, oil palm, rapeseed, groundnut, cotton, and rye) and crop-specific harvested areas to generate crop-area-weighted average of P fertilizer application in each grid during 1961-2020, which is calculated as follows:

$$\overline{F_{crop,y,g}} = \frac{\sum_i (F_{crop,i,2000} \times AH_{i,y})}{\sum_i AH_{i,y}} \quad (1)$$

where $\overline{F_{crop,y,g}}$ is the crop-area-weighted average of P fertilizer application rate in a cropland grid g (g P m^{-2} cropland yr^{-1}) in year y , $F_{crop,i,y}$ and $AH_{i,y}$ are P fertilizer application rate (g P m^{-2}) and harvested area (m^2), respectively, for crop type i in year y . $F_{crop,i,2000}$ used a fixed value derived from the global crop-specific P fertilizer application data circa 2000 (at a resolution of 5-arcmin) developed by Mueller et al. (2012). This dataset was used as the spatial baseline for distributing fertilizer inputs across cropland. The crop distribution maps for individual years were derived by

allocating FAO statistics for crop-specific harvested area into grids using the Dissever algorithm (Malone et al., 2012), which is a general spatial downscaling method using fine-resolution covariate data. Specifically, the covariate data included climate variables (temperature and precipitation) (Fick and Hijmans, 2017), soil information (FAO and IIASA, 2023), and topography. In each year, crop-specific distribution from SPAM2010 (Yu et al., 2020) was used as initial inputs for starting an iterative process to obtain an optimal machine learning model within a region in each year. Though this climate-based disaggregation strategy for statistical analysis inevitably brings uncertainties, it has been usually adopted in generating crop distribution maps (e.g., Li et al., 2025; Yu et al., 2020). In practice, to minimize these uncertainties, the machine learning-based distribution maps are adjusted to meet multiple constraints. For example, the sum of areas of all crops in a grid should not exceed the actual arable land area (see our cropland data generation using HYDE3.3 and LUHv2 in text below), and the area of a specific farming system should not exceed the suitable area within the grid (adopted from SPAM2010).

Annual national P fertilizer consumption from FAOSTAT was subsequently used to scale the baseline application rates, allowing fertilizer inputs to vary over time:

$$R_{crop,y,j} = \frac{FAO_{crop,y,j}}{\sum_{g=1}^{g=n \text{ in country } j} (\overline{F_{crop,y,g}} \times AC_{y,j,g})} \quad (2)$$

$$Pfer_{crop,y,j,g} = \overline{F_{crop,y,g}} \times R_{crop,y,j} \quad (3)$$

where $R_{crop,y,j}$ represents the regulation ratio (unitless) for the year y and country j , $FAO_{crop,y,j}$ is the national total P fertilizer usage (g P yr^{-1}) on cropland, $AC_{y,j,g}$ is the cropland area (m^2) for the year y and grid g of the country. The historical land use data (cropland, pasture, rangeland) covers 1860-2020 and has an annual time-step and a 5 arcmin spatial resolution, developed by reconciling the HYDE 3.3 (<https://landuse.sites.uu.nl/datasets/>) and LUHv2 datasets (Hurtt et al., 2020; Klein Goldewijk et al., 2017; Tian et al., 2022). For the period prior to 1949, HYDE 3.3 provides land use maps only at decadal intervals (i.e., one map every ten years). To

generate continuous annual data, we applied a temporal downscaling approach: the LUHv2 data for each year within a given decade were used to proportionally interpolate the HYDE 3.3 map from the beginning of that decade. This method preserves the decadal baseline provided by HYDE 3.3 while incorporating the interannual variability captured by LUHv2. The P fertilizer in cropland grid g of the country j in year y , $Pfer_{crop}$ ($\text{g P m}^{-2} \text{ yr}^{-1}$), was then calculated as the product of $F_{crop,y,g}$ and $R_{crop,y,j}$.

The earliest year of P fertilizer statistics that the FAO database provides is 1961. Prior to the widespread adoption of phosphate rock as a fertilizer in the 1940s, guano—accumulated bird droppings over millennia—and human excreta were utilized as fertilizers for food crops. The historical P fertilizer data during 1860–1960 were generated based on data reported by Cordell et al. (2009), which provide global total P fertilizer usage estimates back to 1800. Annual relative change rates of global P fertilizer were first calculated based on the estimates from Cordell et al. (2009). These change rates were then applied to the spatially explicit P fertilizer distribution in the reference year of 1961 to reconstruct annual global P fertilizer maps for the period 1860–1960. This approach assumes that temporal dynamics of P fertilizer data before 1961 were primarily driven by global-scale trends.




Figure 1. Workflow of developing data for P fertilizer application on cropland. The

blue box represents the annual data during 1961-2020, and the orange box represents the static variable.

Table 1. Main data sources utilized in HaPi development

Data	Data Source	Dataset	Reference
Products			
Fertilizer	FAOSTAT	Annual country-level P fertilizer to land from 1961 to 2020; Crop-specific harvested area	FAOSTAT (2024)
	EARTHSTAT	Fertilizer rates for major crops	Mueller et al. (2012)
	SPAM2010	Crop distribution maps	Yu et al. (2020)
	Hyde3.2/LUHv2	Cropland, Pasture, and rangeland area from 1860 to 2019	Hurt et al. (2020); Klein Goldewijk et al. (2017)
	Cordell et al. (2009)	Global total P fertilizer usage from 1860 to 1960	Cordell et al. (2009)
Manure	HaNi	Manure Nitrogen inputs from 1860 to 2020	Tian et al. (2022)
	GLW3	Livestock distribution maps	Gilbert et al. (2018)
	FAOSTAT	Annual country-level livestock statistics from 1961 to 2020	FAOSTAT (2024)
	Lun et al. (2018)	Animal-specific manure P:N ratios	Lun et al. (2018)
Atmospheric deposition	HEMCO	Natural P emissions (mineral dust, sea-salt aerosols, and volcanic eruptions)	Meng et al. (2021); Weng et al. (2020); Carn (2019)
	EDGAR	Anthropogenic P emissions	Crippa et al. (2023)

GFED v4	P emissions from wildfire	Randerson et al. (2015)
ERA5	Climate data (temperature, precipitation, wind speed, etc.)	Muñoz-Sabater et al. (2021)
MERRA2	Aerosol Optical Thickness	Randles et al. (2017)
GIMMS LAI4g	Leaf area index	Cao et al. (2023)

2.2 Manure application on cropland/pasture and deposition on pasture/rangeland

The grid-level manure P inputs, including manure application on cropland, manure application on pasture, manure deposition on pasture, and manure deposition on rangeland, were generated by multiplying the corresponding manure N inputs from HaNi (Tian et al., 2022) and manure P:N ratios (Figure 2):

$$Pman_{y,g} = Nman_{y,g} \times RPN_{y,g} \quad (4)$$

where $Pman_{y,g}$ and $Nman_{y,g}$ indicate the P and N manure application or deposition on grid g in year y , respectively. The HaNi dataset offers global manure input data on cropland, pasture, and rangeland during 1860-2020 at a resolution of 5 arcmin. $RPN_{y,g}$ represents the manure P:N ratio in grid g .

$$RPN_{y,g} = \frac{\sum_t APN_{y,t} \times Anum_{y,g,t}}{\sum_t Anum_{y,g,t}} \quad (5)$$

$$Anum_{y,g,t} = \begin{cases} AGLWnum_{g,t} \\ GLMnum_{g,t} \times \frac{LFAO_{y,t,j}}{\sum_{g=1}^{g=n \text{ in country } j} GLMnum_{g,t}} \end{cases} \quad (6)$$

The annual grid-level manure P:N ratios $RPN_{y,g}$ were estimated as the animal-number weighted sum of these animal-specific P:N ratios within grids. We obtained manure P:N ratio $APN_{y,t}$ for each animal species t from Lun et al. (2018). Livestock distribution data $Anum_{y,g,t}$ were derived by fusing the annual gridded livestock of the world (AGLW) dataset (Du et al., 2025) and a newly downscaled product. The

AGLW dataset covers 1961-2021 at a 5 km resolution and was developed based on annual FAOSTAT census information using a machine learning method. But, in some areas (particularly the former Soviet Union region), livestock information is missing. To fill the gaps, we used the national FAO livestock statistics and the Global Livestock of the World 3 (GLW3) database (Gilbert et al., 2018) to generate a long time-series gridded livestock distribution data. Specifically, GLW3 provides global livestock populations $GLMnum_{g,t}$ at a 5-arcmin resolution for cattle, buffaloes, horses, sheep, goats, pigs, chickens, and ducks, but the data are available for a single reference year. Therefore, the annual country-level livestock population data $LFAO_{y,t,j}$ from FAOSTAT were employed to extend the GLW3 dataset into a time series of livestock distribution maps spanning from 1961 to 2020 following the method used in equations 2-3. Based on the fused product of livestock distribution, a spatially explicit manure P:N ratio dataset covering the period 1961–2020 was constructed and subsequently used to estimate manure P inputs from corresponding N inputs. The final nutrient ratio dataset is highly consistent both spatially and temporally (Figure. S1). Notably, although temporal changes in livestock composition are considered when estimating manure P:N ratios, potential changes in species-specific manure nutrient ratios associated with evolving feed composition and livestock management practices are not explicitly represented, which may introduce biases in manure P estimates.




Figure 2. Workflow of developing manure P input data. The blue box represents the annual data during 1961-2020, and the orange box represents the static variable.

2.3 Atmospheric P deposition

Atmospheric P deposition originates from both natural and anthropogenic sources and was estimated by combining a P emission inventory, an atmospheric chemical model, and an algorithm (Figure 3 and Table 1) (You et al., 2026). Natural P emissions primarily comprise dust, sea salt, Primary Biological Aerosol Particles (PBAPs), volcanoes, and wildfires. In this study, emissions from dust, sea salt, and volcanoes were derived from the GEOS-Chem HEMCO data (Meng et al., 2021; Weng et al., 2020; Carn, 2019; Carn et al., 2015; Ge et al., 2016). PBAPs emissions were estimated by utilizing specific humidity data and LAI data according to the method used in Myriokefalitakis et al. (2016). Wildfire emission data were sourced from the GFED4 dataset (Randerson et al., 2015). Meanwhile, anthropogenic emissions were estimated using monthly sector-specific data from the EDGAR database (Crippa et al., 2023). By integrating all the emission sources, we developed a global atmospheric P emission inventory, which was subsequently used as input for the GEOS-Chem model (Feng et al., 2021; Li et al., 2024) to simulate global atmospheric P deposition from 2001 to

2019. GEOS-Chem is a global three-dimensional atmospheric chemical model driven by meteorological inputs provided by the Goddard Earth Observing System (GEOS) of the NASA Global Modeling and Assimilation Office. It has been widely applied to address various atmospheric composition issues. The modeled P deposition by GEOS-Chem was at a resolution of $2.5^{\circ} \times 2^{\circ}$.

To obtain the atmospheric P deposition at a high resolution, the diserver algorithm (Malone et al., 2012; Roudier et al., 2017), which is a mass-conserving spatial downscaling method, was applied to the GEOS-Chem outputs. First, the coarse resolution P deposition data was resampled to $0.1^{\circ} \times 0.1^{\circ}$; Second, the diserver algorithm was used to iteratively train an XGBoost model for eight different regions, including Africa, Asia excluding China and India, China, Europe, India, North America, Oceania, and South America, to retrieve resampled P deposition and fine-resolution environmental variables. These environmental variables included ERA5 climate factors (wind components, temperature, surface net solar radiation, surface pressure, and monthly total precipitation) (Muñoz-Sabater et al., 2021), LAI (Cao et al., 2023), Aerosol Optical Thickness (AOT, Randles et al., 2017), and P emission data. The iterative model training process was terminated when the improvement in performance between successive generations fell below a predefined threshold. Model performance was assessed using the Mean Absolute Error (MAE; unit: g ha^{-1}), calculated for each model's target region across all terrestrial grid cells using the test dataset. The convergence threshold for MAE was initially set to 0.001 g ha^{-1} . Using this approach, we generated global atmospheric P deposition estimates at a spatial resolution of 0.1° for the period 2000–2019. To ensure consistency with the HaPi dataset, the 0.1° data were subsequently resampled to a 5 arc-min resolution and temporally extended back to 1900 using annual P deposition change rates derived from Ringeval et al. (2024). P deposition for the period 1860–1899 was assumed to remain constant and equal to the early twentieth-century levels.




Figure 3. Workflow of developing atmospheric P input data. Where PBAPs represent Primary Biological Aerosol Particles. Prec refers to precipitation. T2m indicates the 2-meter air temperature, while T2m max and T2m min refer to the maximum and minimum air temperatures at 2-meter height. U10 and V10 denote the eastward and northward wind speeds at 10-meter height, respectively. AOT stands for Aerosol Optical Thickness. SP represents Surface Pressure, and SSRD refers to Surface Solar Radiation Downwards.

3 Results

3.1 Global total P inputs

Global anthropogenic P inputs have undergone dramatic changes since the Industrial Revolution, increasing from 3.8 Tg P yr⁻¹ in the 1860s to 41.0 Tg P yr⁻¹ in the 2010s, a nearly 11-fold increase (Figure 4). The most rapid acceleration occurred between 1945 and 1989, coinciding with post-war agricultural intensification. The amplified usage of mineral fertilizer and livestock manure P contributed nearly equally, over the study period, i.e., 52% and 46% of the increase in TP inputs to the terrestrial biosphere, respectively. The composition of P inputs shifted significantly over this period (Table

2). Prior to the 1980s, livestock manure accounted for over half of total P (TP) inputs. However, by the 2010s, mineral fertilizer P in TP inputs surpassed manure P for the first time. Atmospheric P deposition, while relatively stable, declined from 50% to only 7% of TP inputs from the 1860s to the 2010s, due to the marked increase in anthropogenic inputs. The intensified application of P fertilizer on cropland was the primary driver behind the increased TP inputs after 1945.

Regionally, the TP inputs initially increased in Europe and eastern USA between the 1860s and the 1910s, primarily due to enhanced manure inputs (Figures 5-6). Subsequently, P input hotspots emerged in Europe, driven by the rising use of P fertilizer. Europe (5.2 Tg P yr^{-1}) and the USA (2.8 Tg P yr^{-1}) were the top two regions with the highest P inputs in the 1960s. After the 1980s, TP inputs in Europe declined quickly, while leveling off in the USA. By the 2010s, new hotspots emerged in eastern China, northern India, southern Brazil, and eastern Africa, sourcing from the widespread application of mineral P fertilizer and the expansion of livestock production in these regions. Notably, TP inputs in China began to decrease around 2010, whereas inputs in South Asia and Brazil maintained growth trends. Livestock manure remains the dominant P source in most regions of the South Hemisphere (Africa, South America, Oceania), while P fertilizer plays a more important role in industrialized regions of the North Hemisphere (Asia, Europe, North America). In addition to Europe, TP inputs in Russia, Korea, and Japan (KAJ), and Central Asia (CAS) also declined substantially, largely due to the reduction in P fertilizer usage. Manure P inputs continued to increase in Africa, South/Southeast Asia, and Central America (CAM), but have shifted to a decreasing trend in Oceania, Europe, and Russia in recent decades.




Figure 4. Annual changes of anthropogenic phosphorus inputs to terrestrial ecosystems from 1860 to 2020.

Table 2. Decadal mean P inputs to the terrestrial biosphere (Tg P yr⁻¹)

Decade	P _{fer}	P _{fer}	P _{man}	P _{man}	P _{man}	P _{man}	P _{dep}	Total
	Crop	Pasture	Crop	App	Dep	Dep		
				Pasture	Pasture	Range		
1860s	0.1	0.0	0.5	0.2	0.7	0.3	1.9	3.8
1910s	0.6	0.0	1.3	0.4	1.8	1.1	1.9	7.1
1960s	6.2	0.4	2.8	0.7	5.1	3.5	2.0	20.6
1970s	10.5	0.9	3.3	0.8	5.9	3.9	2.2	27.6
1980s	13.9	1.1	3.8	0.9	6.7	4.2	2.4	32.9
1990s	13.2	1.0	3.9	0.8	7.3	4.5	2.4	33.2
2000s	14.7	1.3	4.0	0.8	7.9	4.8	2.5	35.9
2010s	17.9	1.5	4.4	0.8	8.8	5.0	2.7	41.0

Note: The following abbreviations are used in the table: P_{fer} – P fertilizer, P_{man} – manure P, P_{dep} – Atmospheric P deposition, App – Application, and Dep – Deposition.




Figure 5. Spatial patterns in total Phosphorus input in the (a) 1860s, (b) 1910s, (c) 1960s, and (d) 2010s. The labels in the inset pie charts represent the percentage of each component: Fc – P fertilizer applied to cropland, Fp – P fertilizer applied to pasture, Mc – Manure P application on cropland, Map – Manure P application on pasture, Mdp – Manure P deposition on pasture, Mr – Manure P deposition on rangeland, Ad – Atmospheric P deposition.




Figure 6. Annual variations of P inputs in 18 regions during 1860-2020. The 18 regions are the Canada (CAN), USA (USA), Europe (EU), Central Asia (CAS), Russia (RUS), Korea and Japan (KAJ), China (CHN), South Asia (SAS), Southeast Asia (SEAS), Oceania (OCE), Middle East (MIDE), South Africa (SAF), Equatorial Africa (EQAF), North Africa (NAF), South West Asia (SWSA), Brazil (BRA), North South America (NSA), and Central America (CAM).

Oceania (OCE), Middle East (MIDE), Southern Africa (SAF), Equatorial Africa (EQAF), Northern Africa (NAF), southwestern South America (SWSA), Brazil (BRA), northern South America (NSA), and Central America (CAM).

3.2 P fertilizer inputs on cropland and pasture

The global total P fertilizer inputs surged from 6.6 Tg P yr⁻¹ in the 1960s to 19.4 Tg P yr⁻¹ in the 2010s, with cropland receiving over 90% of these inputs (Figure 7). Annual P fertilizer applied to croplands increased rapidly at a rate of 0.4 Tg yr⁻² during 1961–1989, then declined sharply until 1995 before resuming an upward trajectory. Despite this global growth, significant regional disparities emerged (Figures 6 and 8). Europe and the USA were the dominant regions for early P fertilizer consumption, accounting for 38% and 24% of global usage, respectively, in the 1960s. By the 2010s, China (30%) and South Asia (20%) had become the predominant consumers of P fertilizers, driven by intensive agricultural expansion. Regional shifts in P fertilizer inputs were pronounced. In the USA, P fertilizer application to cropland decreased from 2.0 Tg P yr⁻¹ in the 1970s to 1.5 Tg P yr⁻¹ in the 2000s. In Europe, P fertilizer usage peaked in the 1980s but decreased by 66% in the 2010s. The increase in P fertilizer usage in China and South Asia contributed to 44% and 29% of the global total increase in P fertilizer application on cropland from the 1960s to the 2010s, respectively. However, China's usage plateaued in the 2010s, while South Asia's continued to increase rapidly. In the 2020s, hotspot regions with high P fertilizer application rates (>3 g P m⁻² yr⁻¹) were mainly distributed in eastern China, northern India, and southern Brazil. The P fertilizer application rates on cropland were consistently low across most areas in Africa over the whole study period, accounting for only 4% of global consumption in the recent decade.

Global P fertilizer application on pasture increased from 0.3 to 1.6 Tg P yr⁻¹ during 1961-2020. Before the 1980s, European countries were the predominant consumers of P fertilizer on pasture (Figure S2). Thereafter, the USA and India significantly increased

their usage, accounting for 23% and 22% of total P fertilizer application on pasture, respectively, in the 2010s. P fertilizer application rate per pasture land was high in India, Japan, and southern Canada, but remained low in most other countries in recent decades.




Figure 7. Annual variations and trends in P fertilizer inputs during 1860–2020.




Figure 8. Global patterns of P fertilizer application on cropland in the 1960s, 1980s, 2000s, and 2010s.

3.3 Manure P inputs on cropland, pasture, and rangeland

Livestock manure served as the primary P source for agricultural soils historically, with global manure P inputs on cropland, pasture, and rangeland increasing from 1.7 to 18.9

Tg P yr⁻¹ during the period from the 1860s to the 2010s (Figure 9). Pasture received half of total manure P inputs, while cropland and rangeland each shared a similar proportion of the remaining half. Over 1961–2020, manure P deposition on pasture kept increasing at a rate of 0.07 Tg P yr⁻², while manure P application on pasture was relatively stable. Meanwhile, annual manure inputs on cropland and rangeland both increased, but at relatively slow rates than those on pasture. In the context of rising global manure P inputs, developing countries have demonstrated stronger growth rates than developed countries in recent decades. China has surpassed Europe as the region receiving the largest manure P inputs since the 2000s, as manure P inputs in Europe have declined since the 1980s. Over the past four decades, manure usage in North Africa (NAF) and Equatorial Africa (EQAF) has grown at the fastest rate, increasing by 96% and 138%, respectively. In the 2010s, China, South Asia, Brazil, and North Africa, as the top four regions, received 13%, 11%, 11%, and 10% of the global total manure P inputs, respectively.

Application rates of manure P per unit area of cropland increased significantly in Asia, Europe, and North and South America since 1860 (Figure 10). Manure P application on cropland initially increased in western Europe in the 1910s, with subsequent intensified application occurring in eastern Asia and northern South America by the 2010s. Manure application and deposition rates on pasture were extremely high in South and Southeast Asia over the last century (Figures 11, S3, and S4). The proportion of manure deposition in total manure inputs to pasture gradually increased from 79% to 92% from the 1860s to the 2010s. Prominent manure P deposition on rangeland was observed in South and Southeast Asia, with new hotspots developing in Central Africa, eastern South America, northern China, and Europe in the 2010s (Figure 12).




Figure 9. Annual variations and trends in manure P inputs from 1860 to 2020




Figure 10. Global patterns of manure P application on cropland in the 1860s, 1910s, 1960s, and 2010s.




Figure 11. Global patterns of manure P application and deposition on pasture in the 1860s, 1910s, 1960s, and 2010s.




Figure 12. Global patterns of manure P deposition on rangeland in the 1860s, 1910s, 1960s, and 2010s.

3.4 Atmospheric P deposition

Atmospheric P deposition is the major P source for natural ecosystems, such as forests and shrubs, but plays a less important role in cropland. Although anthropogenic activities have led to the increase of atmospheric P deposition from 2.0 Tg P yr⁻¹ in the 1960s to 2.7 Tg P yr⁻¹ in the 2010s, the proportion of atmospheric P deposition in total P inputs decreased dramatically to only 7% by the 2010s. China was the region with the largest atmospheric P deposition (21% of the global total) in the 2010s, followed by North Africa (15%), the USA (11%), and Europe (9%) (Figure 13). Atmospheric P deposition in northern Africa was primarily derived from natural sources and remained relatively stable during the study period. Influenced by anthropogenic activities, atmospheric P deposition in China has continually increased since the 1960s, while that in Europe has shifted from an increase to a decrease in the 1980s. Atmospheric P deposition was relatively low across most regions, with the exception of several notable hotspots in northern Africa, eastern China, central Europe, and eastern USA.




Figure 13. Global patterns of atmospheric P deposition in the 1960s, 1980s, 2000s, and 2010s.

4 Discussion

4.1 Comparison with previous studies

The HaPi dataset provides spatially explicit, annual estimates of anthropogenic P inputs to the terrestrial biosphere. This gridded P input dataset reveals subnational heterogeneity, enabling the detection of localized hotspots of high P input that national averages may obscure. When aggregated to the global scale, HaPi aligns well with existing studies, particularly for cropland fertilizer P inputs, which have been extensively evaluated (Table 3). Nearly all these studies rely on the country-level fertilizer inventory data provided by FAO or IFA, which give consistent global totals. Additional validation against independent national statistics from China and the U.S. shows that HaPi captures comparable magnitudes and similar spatial-temporal patterns of P fertilizer use (Figure S5), although with slightly lower estimates relative to both datasets. These results provide further support for the robustness of national-scale temporal dynamics in HaPi.

In contrast, the HaPi estimates of manure P applied to cropland are consistent with those reported by Zou et al. (2022) but are lower than other estimates, while manure P inputs in grasslands (pasture and rangeland) are higher than previous estimates. Despite these differences, total manure P inputs in HaPi remain consistent with other studies. The primary source of discrepancy arises from differences in the allocation of manure inputs between cropland and grassland and is partly driven by inconsistencies in cropland definitions, especially regarding the inclusion of temporary pastures. In previous studies, manure P inputs were calculated by multiplying manure N inputs provided by FAO and P:N ratio in livestock manure products. According to the latest FAOSTAT dataset, around 75% of total manure N inputs are deposited on grassland, and the remaining 25% is applied to cropland (around 21%) and grassland (4%) soils. The majority of manure is directly deposited on grasslands through grazing animal excretion, whereas a smaller proportion is collected and subsequently applied to pastures and croplands as a managed nutrient input. HaPi estimates 23% of manure P was applied on cropland, closely matching FAOSTAT's current methodology. In contrast, Lun et al. (2018) and Sattari et al. (2016) estimated 36% and 38% of total manure P inputs to cropland.

Compared to previous studies, the HaPi dataset offers the most comprehensive coverage of anthropogenic P input fluxes, featured by a high spatial resolution and extended historical coverage. These features support the analysis of legacy P accumulation and depletion, and simultaneously provide consistent forcing data for land surface and biogeochemical models. This harmonized P input dataset enhances our ability to assess regional nutrient trends, environmental risks, and management needs at multiple spatial scales.

Table 3. Comparison of P input data with other global estimates

Global P inputs	Previous studies (Tg P Yr ⁻¹)	Year	This study (Tg P Yr ⁻¹)	Reference
Fertilizer input on cropland	16.4	2002-2010	17.6	Lun et al. (2018)
	13.8	2000	13.2	Bouwman et al. (2009)
	0.4-15.8	1900-2010	0.3-17.6	Zhang et al. (2017b)
	4.6-17.5	1961-2013	4.5-18.0	Lu and Tian (2017)
	4.5-21.0	1961-2020	4.5-19.7	Ludemann et al. (2024)
	16.8	2008-2012	16.7	Zou et al. (2022)
Fertilizer input on pasture	0.4	2002-2010	1.5	Lun et al. (2018)
Manure input on cropland	1.6-7.2	1900-2010	1.0-4.3	Zhang et al. (2017b)
	7.1	2002-2010	4.1	Lun et al. (2018)
	6.3	2005	4.0	Sattari et al. (2016)
	4.3-6.6	1961-2020	2.7-4.5	Ludemann et al. (2024)
	4.7	2008-2012	4.2	Zou et al. (2022)

Manure input on	12.7	2002-2010	13.7	Lun et al. (2018)
grassland (pasture and rangeland)	10.2	2005	13.7	Sattari et al. (2016)
Total manure input	17.1	2000	16.8	Bouwman et al. (2013)
	16.5	2005	17.7	Sattari et al. (2016)
	19.8	2002-2010	17.8	Lun et al. (2018)

494

495 4.2 The implication of changes in manure and fertilizer P inputs

496 Globally, mineral P fertilizer has surpassed manure P as the largest P input to cropland,
497 while livestock manure remains the dominant P source for pasture and rangeland.
498 Overall, livestock manure is a more significant source of P for terrestrial ecosystems
499 compared to mineral fertilizer. In recent decades, livestock manure contributed over
500 half of the P sources in most regions, especially in Africa where around 90% of total P
501 inputs were derived from manure. As P fertilizer application began to decrease in
502 European countries, manure P application on cropland played an increasingly important
503 role in food production. Grassland, including pasture and rangeland, received over 70%
504 of global manure P inputs and over 40% of total P inputs; therefore, it is critical to take
505 account of grassland when investigating global P balance and P-related environmental
506 issues. For instance, it is estimated that P loading to the Gulf of Mexico originated
507 primarily from manure on pasture and rangeland (37%), followed by corn and soybeans
508 (25%) within the Mississippi River Basin (Alexander et al., 2008). Despite high manure
509 P input, manure excretion is still partly an internal P cycling in the grassland-livestock
510 system since it derives from soil P uptake by grass. Furthermore, global manure P input
511 cannot compensate for the grazing P output in grassland systems because around 23%
512 of manure was transferred from grassland to cropland systems. Due to the limited
513 application of fertilizer to grasslands, global grasslands are facing the challenge of a
514 negative P budget (Sattari et al., 2016). Given the projected increase in livestock

production to meet future demand for meat and milk, effectively managing P flows in the whole crop-livestock production systems is critical for the sustainable human P cycle (Bouwman et al., 2013).

Socio-economic development clearly drives the long-term changes in total anthropogenic P inputs to agricultural lands (Figure S6). The rapid increase in P fertilizer application on cropland in the latter half of the 20th century stimulated crop production but ultimately resulted in a decrease in the P use efficiency. After about 60 years of growth, global total P fertilizer use has leveled off in the 2010s. Europe, previously the largest consumer of P fertilizer, began reducing its usage in the 1980s. China, currently the largest consumer, has also shown a decrease in P fertilizer consumption since 2013. Although P fertilizer application decreases, crop yield and P uptake may not decline correspondingly. The large P surpluses due to previous P application in European countries have built up soil residual P pool which can continually supply crop production (Sattari et al., 2012). Similarly, the reduced usage of P fertilizer in China may also result from the supply of soil legacy P resources accumulated over the last decades. The increased crop yield and decreased P fertilizer usage indicated an enhancement in P use efficiency in these industrialized countries. Meanwhile, many developing countries, such as India and Brazil, continue to experience elevated P fertilizer usage and an amplified P surplus on croplands (Zhang et al., 2017b). Utilizing residual soil P can be a key strategy to reduce reliance on imported mineral P fertilizer and improve the sustainability of agriculture in these countries.

4.3 Limitations in Data Development

Despite numerous improvements in the HaPi dataset over previous P input datasets, several limitations remain in its development. First, the fertilizer input dataset was developed using the spatial distribution of crop-specific fertilizer application rates around the year 2000 as a baseline map. Although fertilizer application rates vary over time through scaling with national statistics, the fixed spatial pattern of fertilizer use

would disregard subnational variability in fertilizer application rates for each crop. Furthermore, the climate-based disaggregation strategy for FAOSTAT crop harvest area needs further improvement. A key challenge is the lack of globally consistent, dynamic maps (e.g., yearly suitable area for specific farming systems) for constraining spatial allocation of statistical data. To improve the crop-specific harvest area estimates, more efforts are required for obtaining fine-grained statistics (national or provincial level statistics data is used in this study) of varied crops. Another plausible solution is to fuse satellite-retrieved crop distribution products with these statistical disaggregation maps. But the major limits of this pathway are the discontinuous temporal coverage and lack of global availability for many crop types in satellite products.

Second, we used historical cropland, pasture, and rangeland data from HYDE/LUHv2 to spatialize country-level P fertilizer use amounts, but HYDE/LUHv2 data have been shown to exhibit inconsistent spatial and temporal patterns of land use relative to satellite-derived land use at the regional scale. For example, the HYDE dataset overestimates the cropland area in India, which can lead to the underestimation of the P fertilizer application rate on cropland. In this study, we assumed that the ratios of P fertilizer application on pasture relative to cropland were the same as those for N fertilizer, which may not accurately reflect the actual allocation of P fertilizer usage on pasture. Since country-level data for P fertilizer are only available from 1961 onwards, we assumed that the change rates of global fertilizer inputs before 1961 followed the annual global trends reported by Cordell et al. (2009). Although this approach provides a temporally continuous long-term dataset, it cannot explicitly represent historical changes in crop composition, fertilizer accessibility, and agricultural management practices, and it ignores regional variations in the changes of P fertilizer usage across different countries before 1961. Consequently, uncertainties are substantially higher in the pre-1961 period, and the resulting maps should be interpreted primarily as broad-scale reconstructions rather than precise representations of historical subnational fertilizer distributions.

Third, we calculated manure P inputs based on manure N inputs and P:N ratio with spatiotemporal heterogeneity, but the gridded manure P data were not constrained by survey data, as country-level manure P inputs data were not available from the Soil Nutrient Budget database in FAOSTAT. Although temporal changes in livestock composition are considered when estimating manure N:P ratios, potential changes in species-specific manure nutrient ratios associated with evolving feed composition and livestock management practices are not explicitly represented, which may introduce biases in manure P estimates. Aside from mineral fertilizer, livestock manure, and atmospheric deposition, other anthropogenic P sources, including guano, livestock bones, and human excreta, were historically used to enhance soil fertility. These P sources were difficult to quantify at a global level without reliable data sources and were therefore not included or accurately quantified in the HaPi dataset.

4.4 Uncertainty

The uncertainties in the HaPi dataset mainly arise from the input datasets and methodological assumptions used in developing the global P input estimates. Quantifying the overall uncertainties in the HaPi dataset is challenging due to the heterogeneity of underlying data sources and the scarcity of independent datasets for robust validation. Four major sources of uncertainty were identified as contributing to the overall uncertainty in the HaPi dataset. First, the FAO survey data, which serve as the primary constraint for national total P inputs, represents a critical source of uncertainty in estimating total P inputs to the terrestrial biosphere. Based on expert judgment, a generic uncertainty of approximately 20% was assigned to the FAO national estimates. Consequently, the spatially explicit P input maps derived from these national data inherently carry at least the same level of uncertainty, with a minimum estimated uncertainty of 20%. Second, uncertainties arise from the land use maps used to spatialize P inputs. Uncertainties in land cover classification or temporal interpolation in HYDE/LUHv2 data may propagate into the spatial allocation of P fertilizer and manure applications, thereby affecting local input rates and hotspot

identification. Third, uncertainties stem from empirical assumptions and static parameters used during dataset construction, including the assumption of partitioning ratios for fertilizer use between cropland and pasture, the use of time-invariant crop-specific fertilizer and manure application patterns, the fixed manure N:P ratios for different livestock species, and the application of globally uniform change rates for reconstructing pre-1961 data. Finally, uncertainties in atmospheric P deposition simulated by the GEOS-Chem model are primarily associated with model structure and parameterization. Although the HaPi dataset primarily focuses on anthropogenic P inputs, it additionally includes both natural and anthropogenic sources. Collectively, these sources contribute to uncertainties in both the magnitude and spatial distribution of anthropogenic P inputs. Given the limitations and uncertainties in the HaPi dataset, it is important to collect or conduct surveys of crop-specific P fertilizer and manure use at subnational scales and update global land use data to reflect more precise regional patterns of global fertilizer and manure P inputs.

Data availability

The History of Anthropogenic P Inputs (HaPi) dataset is available at <https://doi.org/10.6084/m9.figshare.29930279> (Bian et al., 2026).

Competing interests

At least one of the (co-)authors is a member of the editorial board of Earth System Science Data. The peer review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Author contributions

Z.B., H.S., and H.T. designed this work and developed the datasets. R.L., S.Y., B.H., and J.M. estimated atmospheric P deposition. F.N.T. provided the FAO dataset. N.D.M.

provided the crop-specific P fertilizer application datasets. L.F., H.R., L.L., H.C., and Y.Y. contributed to the development of methodology. All authors contributed to the writing of the manuscript.

Acknowledgements

Z.B. acknowledges funding support from the National Natural Science Foundation of China (42401015; U24A20577) and Jiangsu Provincial Department of Science and Technology (BK20240599). H.S. received support from the Joint CAS-MPG Research Project (HZXM20225001MI) and the National Key Research and Development Program of China (Grant No. 2023YFF1303700). H.T. acknowledges the funding support from U.S. National Science Foundation (Grant No. 1903722) and USDA CBG (Grant No. TENX12899). FAOSTAT data are produced under FAO regular budget, with inputs from national experts in member countries. The views expressed in this work are the authors' only and do not reflect FAO positions on the subject matter.

References

- Alexander, R. B., Smith, R. A., Schwarz, G. E., Boyer, E. W., Nolan, J. V., and Brakebill, J. W.: Differences in Phosphorus and Nitrogen Delivery to The Gulf of Mexico from the Mississippi River Basin, *Environ. Sci. Technol.*, 42, 822–830, <https://doi.org/10.1021/es0716103>, 2008.
- Barbieri, P., MacDonald, G. K., Bernard de Raymond, A., and Nesme, T.: Food system resilience to phosphorus shortages on a telecoupled planet, *Nat. Sustain.*, 5, 114 – 122, <https://doi.org/10.1038/s41893-021-00816-1>, 2022.
- Beusen, A. H. W., Bouwman, A. F., Van Beek, L. P. H., Mogollón, J. M., and Middelburg, J. J.: Global riverine N and P transport to ocean increased during the 20th century despite increased retention along the aquatic continuum, *Biogeosciences*, 13, 2441–2451, <https://doi.org/10.5194/bg-13-2441-2016>, 2016.
- Bian, Z; Shi, H; Tian, H.: HaPi: A Historical dataset of Anthropogenic Phosphorus Inputs to the terrestrial biosphere (1860-2020), <https://doi.org/10.6084/m9.figshare.29930279>, 2026
- Bouwman, A. F., Beusen, A. H., and Billen, G.: Human alteration of the global nitrogen and phosphorus soil balances for the period 1970–2050, *Glob. Biogeochem. Cycles*, 23, 2009.

656 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P.,
657 Willems, J., Rufino, M. C., and Stehfest, E.: Exploring global changes in nitrogen and phosphorus
658 cycles in agriculture induced by livestock production over the 1900-2050 period, *Proc. Natl. Acad.*
659 *Sci.*, 110, 20882–20887, <https://doi.org/10.1073/pnas.1012878108>, 2013.

660 Cao, S., Li, M., Zhu, Z., Wang, Z., Zha, J., Zhao, W., Duanmu, Z., Chen, J., Zheng, Y., Chen, Y.,
661 Myneni, R. B., and Piao, S.: Spatiotemporally consistent global dataset of the GIMMS leaf area
662 index (GIMMS LAI4g) from 1982 to 2020, *Earth Syst. Sci. Data*, 15, 4877 – 4899,
663 <https://doi.org/10.5194/essd-15-4877-2023>, 2023.

664 Carn, S.: Multi-Satellite Volcanic Sulfur Dioxide L4 Long-Term Global Database V3,
665 <https://doi.org/10.5067/MEASURES/SO2/DATA404>, 2019.

666 Carn, S. A., Yang, K., Prata, A. J., and Krotkov, N. A.: Extending the long-term record of volcanic
667 SO₂ emissions with the Ozone Mapping and Profiler Suite nadir mapper, *Geophys. Res. Lett.*, 42,
668 925–932, <https://doi.org/10.1002/2014GL062437>, 2015.

669 Carpenter, S. R.: Eutrophication of aquatic ecosystems: Bistability and soil phosphorus, *Proc. Natl.*
670 *Acad. Sci.*, 102, 10002–10005, <https://doi.org/10.1073/pnas.0503959102>, 2005.

671 Conley, D. J., Paerl, H. W., Howarth, R. W., Boesch, D. F., Seitzinger, S. P., Havens, K. E., Lancelot,
672 C., and Likens, G. E.: Controlling Eutrophication: Nitrogen and Phosphorus, *Science*, 323, 1014–
673 1015, <https://doi.org/10.1126/science.1167755>, 2009.

674 Cordell, D., Drangert, J.-O., and White, S.: The story of phosphorus: Global food security and food
675 for thought, *Glob. Environ. Change*, 19, 292–305, <https://doi.org/10.1016/j.gloenvcha.2008.10.009>,
676 2009.

677 Crippa, M., Guizzardi, D., Schaaf, E., Monforti-Ferrario, F., Quadrelli, R., Risquez Martin, A., Rossi,
678 S., Vignati, E., Muntean, M., Brandao De Melo, J., Oom, D., Pagani, F., Banja, M., Taghavi-
679 Moharamli, P., Köykkä, J., Grassi, G., Branco, A., and San-Miguel, J.: GHG emissions of all world
680 countries: 2023, Publications Office of the European Union, 2023.

681 Dai, M., Zhao, Y., Chai, F., Chen, M., Chen, N., Chen, Y., Cheng, D., Gan, J., Guan, D., Hong, Y.,
682 Huang, J., Lee, Y., Leung, K. M. Y., Lim, P. E., Lin, S., Lin, X., Liu, X., Liu, Z., Luo, Y.-W., Meng,
683 F., Sangmanee, C., Shen, Y., Uthaiapan, K., Talaat, W. I. A. W., Wan, X. S., Wang, C., Wang, D.,
684 Wang, G., Wang, S., Wang, Y., Wang, Y., Wang, Z., Wang, Z., Xu, Y., Yang, J.-Y. T., Yang, Y.,
685 Yasuhara, M., Yu, D., Yu, J., Yu, L., Zhang, Z., and Zhang, Z.: Persistent eutrophication and hypoxia
686 in the coastal ocean, *Camb. Prisms Coast. Futur.*, 1, e19, <https://doi.org/10.1017/cft.2023.7>, 2023.

687 Demay, J., Ringeval, B., Pellerin, S., and Nesme, T.: Half of global agricultural soil phosphorus
688 fertility derived from anthropogenic sources, *Nat. Geosci.*, 16, 69 – 74,
689 <https://doi.org/10.1038/s41561-022-01092-0>, 2023.

690 Du, Z., Yu, L., Zhao, Y., Li, X., Liu, X., Li, X., Hao, P., Chen, Z., Guo, Z., You, L., Ma, X., and
691 Wang, H.: Annual global grided livestock mapping from 1961 to 2021, *Earth Syst. Sci. Data*, 17,

692 5543–5556, <https://doi.org/10.5194/essd-17-5543-2025>, 2025.

693 Einarsson, R., Sanz-Cobena, A., Aguilera, E., Billen, G., Garnier, J., van Grinsven, H. J. M., and
694 Lassaletta, L.: Crop production and nitrogen use in European cropland and grassland 1961–2019,
695 *Sci. Data*, 8, 288, <https://doi.org/10.1038/s41597-021-01061-z>, 2021.

696 Elser, J. and Bennett, E.: A broken biogeochemical cycle, *Nature*, 478, 29 – 31,
697 <https://doi.org/10.1038/478029a>, 2011.

698 FAO: Cropland nutrient balance – Global, regional and country trends, 1961–2022, FAO ;, Rome,
699 2024.

700 FAO and IIASA: Harmonized World Soil Database Version 2.0, Rome Laxenburg, 2023.

701 Feng, X., Lin, H., Fu, T.-M., Sulprizio, M. P., Zhuang, J., Jacob, D. J., Tian, H., Ma, Y., Zhang, L.,
702 Wang, X., Chen, Q., and Han, Z.: WRF-GC (v2.0): online two-way coupling of WRF (v3.9.1.1) and
703 GEOS-Chem (v12.7.2) for modeling regional atmospheric chemistry–meteorology interactions,
704 *Geosci. Model Dev.*, 14, 3741–3768, <https://doi.org/10.5194/gmd-14-3741-2021>, 2021.

705 Fick, S. E. and Hijmans, R. J.: WorldClim 2: new 1-km spatial resolution climate surfaces for global
706 land areas, *Int. J. Climatol.*, 37, 4302–4315, <https://doi.org/10.1002/joc.5086>, 2017.

707 Fink, G., Alcamo, J., Flörke, M., and Reder, K.: Phosphorus Loadings to the World's Largest Lakes:
708 Sources and Trends, *Glob. Biogeochem. Cycles*, 32, 617 – 634,
709 <https://doi.org/10.1002/2017GB005858>, 2018.

710 Ge, C., Wang, J., Carn, S., Yang, K., Ginoux, P., and Krotkov, N.: Satellite-based global volcanic
711 SO₂ emissions and sulfate direct radiative forcing during 2005–2012, *J. Geophys. Res. Atmos.*, 121,
712 3446–3464, <https://doi.org/10.1002/2015JD023134>, 2016.

713 Gilbert, M., Nicolas, G., Cinardi, G., Van Boeckel, T. P., Vanwambeke, S. O., Wint, G. R. W., and
714 Robinson, T. P.: Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens
715 and ducks in 2010, *Sci. Data*, 5, 180227, <https://doi.org/10.1038/sdata.2018.227>, 2018.

716 Hurtt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk,
717 J., Fujimori, S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F.,
718 Jungclaus, J., Kaplan, J. O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz,
719 O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello,
720 F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of global land use change and management
721 for the period 850 – 2100 (LUH2) for CMIP6, *Geosci. Model Dev.*, 13, 5425 – 5464,
722 <https://doi.org/10.5194/gmd-13-5425-2020>, 2020.

723 Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates
724 for the Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927 – 953,
725 <https://doi.org/https://doi.org/10.5194/essd-9-927-2017>, 2017.

726 Lassaletta, L., Billen, G., Grizzetti, B., Anglade, J., and Garnier, J.: 50 year trends in nitrogen use

727 efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland,
 728 *Environ. Res. Lett.*, 9, 105011, <https://doi.org/10.1088/1748-9326/9/10/105011>, 2014.

729 Li, R., Gao, Y., Zhang, L., Shen, Y., Xu, T., Sun, W., and Wang, G.: Global estimates of ambient
 730 reactive nitrogen components during 2000–2100 based on the multi-stage model, *Atmos. Chem.*
 731 *Phys.*, 24, 7623–7636, <https://doi.org/10.5194/acp-24-7623-2024>, 2024.

732 Li, X., Yu, L., Du, Z., and Liu, X.: Crop Statistic to Annual Map: Tracking spatiotemporal dynamics
 733 of crop-specific areas through machine learning and statistics disaggregating, *Sci. Data*, 12, 1249,
 734 <https://doi.org/10.1038/s41597-025-05572-x>, 2025.

735 Lu, C. C. and Tian, H.: Global nitrogen and phosphorus fertilizer use for agriculture production in
 736 the past half century: shifted hot spots and nutrient imbalance, *Earth Syst. Sci. Data*, 9, 181, 2017.

737 Ludemann, C. I., Wanner, N., Chivenge, P., Dobermann, A., Einarsson, R., Grassini, P., Gruere, A.,
 738 Jackson, K., Lassaletta, L., Maggi, F., Obli-Laryea, G., van Ittersum, M. K., Vishwakarma, S., Zhang,
 739 X., and Tubiello, F. N.: A global FAOSTAT reference database of cropland nutrient budgets and
 740 nutrient use efficiency (1961–2020): nitrogen, phosphorus and potassium, *Earth Syst. Sci. Data*, 16,
 741 525–541, <https://doi.org/10.5194/essd-16-525-2024>, 2024.

742 Lun, F., Liu, J., Ciais, P., Nesme, T., Chang, J., Wang, R., Goll, D., Sardans, J., Peñuelas, J., and
 743 Obersteiner, M.: Global and regional phosphorus budgets in agricultural systems and their
 744 implications for phosphorus-use efficiency, *Earth Syst. Sci. Data*, 10, 1 – 18,
 745 <https://doi.org/10.5194/essd-10-1-2018>, 2018.

746 Ma, J., Shi, H., Zhu, Y., Li, R., Wang, S., Lu, N., Yao, Y., Bian, Z., and Huang, K.: The Evolution
 747 of Global Surface Ammonia Concentrations during 2001–2019: Magnitudes, Patterns, and Drivers,
 748 *Environ. Sci. Technol.*, 59, 5066–5079, <https://doi.org/10.1021/acs.est.4c14020>, 2025.

749 Malone, B. P., McBratney, A. B., Minasny, B., and Wheeler, I.: A general method for downscaling
 750 earth resource information, *Comput. Geosci.*, 41, 119 – 125,
 751 <https://doi.org/10.1016/j.cageo.2011.08.021>, 2012.

752 Meng, J., Martin, R. V., Ginoux, P., Hammer, M., Sulprizio, M. P., Ridley, D. A., and van Donkelaar,
 753 A.: Grid-independent high-resolution dust emissions (v1.0) for chemical transport models:
 754 application to GEOS-Chem (12.5.0), *Geosci. Model Dev.*, 14, 4249 – 4260,
 755 <https://doi.org/10.5194/gmd-14-4249-2021>, 2021.

756 Mueller, N. D., Gerber, J. S., Johnston, M., Ray, D. K., Ramankutty, N., and Foley, J. A.: Closing
 757 yield gaps through nutrient and water management, *Nature*, 490, 254 – 257,
 758 <https://doi.org/10.1038/nature11420>, 2012.

759 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta,
 760 S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fern
 761 ández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global
 762 reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349 – 4383,
 763 <https://doi.org/10.5194/essd-13-4349-2021>, 2021.

764 Myriokefalitakis, S., Nenes, A., Baker, A. R., Mihalopoulos, N., and Kanakidou, M.: Bioavailable
 765 atmospheric phosphorous supply to the global ocean: a 3-D global modeling study, *Biogeosciences*,
 766 13, 6519–6543, <https://doi.org/10.5194/bg-13-6519-2016>, 2016.

767 Nguyen, T. H., Tang, F. H. M., Conchedda, G., Casse, L., Obli-Laryea, G., Tubiello, F. N., and
 768 Maggi, F.: NPKGRIDS: a global georeferenced dataset of N, P₂O₅, and K₂O fertilizer application
 769 rates for 173 crops, *Sci. Data*, 11, 1179, <https://doi.org/10.1038/s41597-024-04030-4>, 2024.

770 Randerson, J. T., Van Der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire
 771 Emissions Database, Version 4.1 (GFEDv4), ORNL DAAC,
 772 <https://doi.org/10.3334/ORNLDAAAC/1293>, 2015.

773 Randles, C. A., Silva, A. M. da, Buchard, V., Colarco, P. R., Darmenov, A., Govindaraju, R., Smirnov,
 774 A., Holben, B., Ferrare, R., Hair, J., Shinozuka, Y., and Flynn, C. J.: The MERRA-2 Aerosol
 775 Reanalysis, 1980 Onward. Part I: System Description and Data Assimilation Evaluation,
 776 <https://doi.org/10.1175/JCLI-D-16-0609.1>, 2017.

777 Ringeval, B., Augusto, L., Monod, H., van Apeldoorn, D., Bouwman, L., Yang, X., Achat, D. L.,
 778 Chini, L. P., Van Oost, K., Guenet, B., Wang, R., Decharme, B., Nesme, T., and Pellerin, S.:
 779 Phosphorus in agricultural soils: drivers of its distribution at the global scale, *Glob. Change Biol.*,
 780 23, 3418–3432, <https://doi.org/10.1111/gcb.13618>, 2017.

781 Ringeval, B., Demay, J., Goll, D. S., He, X., Wang, Y.-P., Hou, E., Matej, S., Erb, K.-H., Wang, R.,
 782 Augusto, L., Lun, F., Nesme, T., Borrelli, P., Helfenstein, J., McDowell, R. W., Pletnyakov, P., and
 783 Pellerin, S.: A global dataset on phosphorus in agricultural soils, *Sci. Data*, 11, 17,
 784 <https://doi.org/10.1038/s41597-023-02751-6>, 2024.

785 Roudier, P., Malone, B. P., Hedley, C. B., Minasny, B., and McBratney, A. B.: Comparison of
 786 regression methods for spatial downscaling of soil organic carbon stocks maps, *Comput. Electron.*
 787 *Agric.*, 142, 91–100, <https://doi.org/10.1016/j.compag.2017.08.021>, 2017.

788 Sattari, S. Z., Bouwman, A. F., Giller, K. E., and van Ittersum, M. K.: Residual soil phosphorus as
 789 the missing piece in the global phosphorus crisis puzzle, *Proc. Natl. Acad. Sci.*, 109, 6348–6353,
 790 <https://doi.org/10.1073/pnas.1113675109>, 2012.

791 Sattari, S. Z., Bouwman, A. F., Martínez Rodríguez, R., Beusen, A. H. W., and van Ittersum, M. K.:
 792 Negative global phosphorus budgets challenge sustainable intensification of grasslands, *Nat.*
 793 *Commun.*, 7, 10696, <https://doi.org/10.1038/ncomms10696>, 2016.

794 Stackpoole, S. M., Stets, E. G., and Sprague, L. A.: Variable impacts of contemporary versus legacy
 795 agricultural phosphorus on US river water quality, *Proc. Natl. Acad. Sci.*, 116, 20562–20567,
 796 <https://doi.org/10.1073/pnas.1903226116>, 2019.

797 Steffen, W., Richardson, K., Rockstrom, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R.,
 798 Carpenter, S. R., de Vries, W., de Wit, C. A., Folke, C., Gerten, D., Heinke, J., Mace, G. M., Persson,
 799 L. M., Ramanathan, V., Reyers, B., and Sorlin, S.: Planetary boundaries: Guiding human

800 development on a changing planet, *Science*, 347, 1259855 – 1259855,
801 <https://doi.org/10.1126/science.1259855>, 2015.

802 Tian, H., Bian, Z., Shi, H., Qin, X., Pan, N., Lu, C., Pan, S., Tubiello, F. N., Chang, J., Conchedda,
803 G., Liu, J., Mueller, N., Nishina, K., Xu, R., Yang, J., You, L., and Zhang, B.: History of
804 anthropogenic Nitrogen inputs (HaNi) to the terrestrial biosphere: a 5 arcmin resolution
805 annual dataset from 1860 to 2019, *Earth Syst. Sci. Data*, 14, 4551 – 4568,
806 <https://doi.org/10.5194/essd-14-4551-2022>, 2022.

807 Tian, H., Pan, N., Thompson, R. L., Canadell, J. G., Suntharalingam, P., Regnier, P., Davidson, E.
808 A., Prather, M., Ciais, P., Muntean, M., Pan, S., Winiwarter, W., Zaehle, S., Zhou, F., Jackson, R. B.,
809 Bange, H. W., Berthet, S., Bian, Z., Bianchi, D., Bouwman, A. F., Buitenhuis, E. T., Dutton, G., Hu,
810 M., Ito, A., Jain, A. K., Jeltsch-Thömmes, A., Joos, F., Kou-Giesbrecht, S., Krummel, P. B., Lan, X.,
811 Landolfi, A., Lauerwald, R., Li, Y., Lu, C., Maavara, T., Manizza, M., Millet, D. B., Mühle, J., Patra,
812 P. K., Peters, G. P., Qin, X., Raymond, P., Resplandy, L., Rosentreter, J. A., Shi, H., Sun, Q., Tonina,
813 D., Tubiello, F. N., van der Werf, G. R., Vuichard, N., Wang, J., Wells, K. C., Western, L. M., Wilson,
814 C., Yang, J., Yao, Y., You, Y., and Zhu, Q.: Global nitrous oxide budget (1980–2020), *Earth Syst.*
815 *Sci. Data*, 16, 2543–2604, <https://doi.org/10.5194/essd-16-2543-2024>, 2024.

816 Weng, H., Lin, J., Martin, R., Millet, D. B., Jaeglé, L., Ridley, D., Keller, C., Li, C., Du, M., and
817 Meng, J.: Global high-resolution emissions of soil NO_x, sea salt aerosols, and biogenic volatile
818 organic compounds, *Sci. Data*, 7, 148, <https://doi.org/10.1038/s41597-020-0488-5>, 2020.

819 Xu, R., Pan, S. F., Chen, J., Chen, G. S., Yang, J., Dangal, S. R. S., Shepard, J. P., and Tian, H. Q.:
820 Half - century ammonia emissions from agricultural systems in Southern Asia: Magnitude,
821 spatiotemporal patterns, and implications for human health, *GeoHealth*, 2, 40–53, 2018.

822 You, S., Ma, J., Zheng, H., Liu, X., Tian, H., Wu, Y., Zaehle, S., Wang, Y., Shi, H., Li, R., and Shi,
823 H.: Human Activities Dominate Global Trends in Terrestrial Phosphorus Deposition, *Environ. Sci.*
824 *Technol.*, 60, 10850–10860, <https://doi.org/10.1021/acs.est.5c16208>, 2026.

825 Yu, Q., You, L., Wood-Sichra, U., Ru, Y., Joglekar, A. K. B., Fritz, S., Xiong, W., Lu, M., Wu, W.,
826 and Yang, P.: A cultivated planet in 2010 – Part 2: The global gridded agricultural-production maps,
827 *Earth Syst. Sci. Data*, 12, 3545–3572, <https://doi.org/10.5194/essd-12-3545-2020>, 2020.

828 Yuan, Z., Jiang, S., Sheng, H., Liu, X., Hua, H., Liu, X., and Zhang, Y.: Human Perturbation of the
829 Global Phosphorus Cycle: Changes and Consequences, *Environ. Sci. Technol.*, 52, 2438–2450,
830 <https://doi.org/10.1021/acs.est.7b03910>, 2018.

831 Zhang, B., Tian, H., Lu, C., Dangal, S. R. S., Yang, J., and Pan, S.: Global manure nitrogen
832 production and application in cropland during 1860–2014: a 5 arcmin gridded global dataset for
833 Earth system modeling, *Earth Syst. Sci. Data*, 9, 667–678, <https://doi.org/10.5194/essd-9-667-2017>,
834 2017a.

835 Zhang, J., Beusen, A. H. W., Van Apeldoorn, D. F., Mogollón, J. M., Yu, C., and Bouwman, A. F.:
836 Spatiotemporal dynamics of soil phosphorus and crop uptake in global cropland during the 20th

837 century, *Biogeosciences*, 14, 2055–2068, <https://doi.org/10.5194/bg-14-2055-2017>, 2017b.

838 Zou, T., Zhang, X., and Davidson, E. A.: Global trends of cropland phosphorus use and sustainability
839 challenges, *Nature*, 611, 81–87, <https://doi.org/10.1038/s41586-022-05220-z>, 2022.

840