



1 **Title: A revisiting of early 18th century environmental data to**
2 **identify Gulf of Lion properties before the industrial era**

3 Short title: Gulf of Lion environmental properties in early '700.

4

5 Marina Locritani¹, Sara Garvani^{1,2}, Giuseppe Manzella³, Giancarlo Tamburello¹, Antonio Guarnieri⁴

6 ¹Istituto Nazionale di Geofisica e Vulcanologia, Roma, 00143, Italy

7 ²Università di Genova, Genova, 16126, Italy

8 ³OceanHis, Torino, 10153, Italy

9 ⁴Istituto Nazionale di Geofisica e Vulcanologia, Roma, 00143, Italy, now at DG MARE - European Commission,
10 Bruxelles, 1049, Belgium

11

12 *Correspondence to:* Marina Locritani (marina.locritani@ingv.it)

13 **Keywords:** historical oceanography, Marsili, Marsigli, Histoire Physique de la Mer, bathymetry, water specific gravity, density and
14 salinity, temperature, sea level, Gulf of Lion, Mediterranean Sea

15 **Abstract**

16 The work "*Histoire Physique de la mer*", authored by Luigi Ferdinando Marsili (or Marsigli) and published in 1725, is
17 one of the earliest texts detailing observations of the physical, biological, and bathymetric characteristics of the sea,
18 mainly concentrating on the Gulf of Lion in southern France's Mediterranean area. Nonetheless, understanding Marsili's
19 findings is difficult due to the application of non-standard measurement units and the imprecision of georeferencing data.
20 The MACMAP project (A Multidisciplinary Analysis of Climate Change Indicators in the Mediterranean and Polar
21 Regions), which is funded by the Istituto Nazionale di Geofisica e Vulcanologia (INGV), has involved a thorough
22 recalibration of Marsili's observations. This project focused on transforming water weight measurements obtained from
23 different locations in the Gulf of Lion from June 1806 to January 1807 into water density values. The sampling sites were
24 digitized, bathymetric profiles were reconstructed, and tide amplitudes were examined. The main objective is to make
25 this historical data available to compare with current measurements.

26 **1. Introduction**

27 The study of oceanography took a methodological turn with the 'recommendations' made by Murray and Hooke in 1666,
28 which emphasized the need for 'diligent' observations from 'masters of ships, pilots, and other fit individuals during their
29 sea voyages.' These 'recommendations' outlined various measures, along with the methodologies and technologies that
30 should be employed. A significant enhancement to the concept of 'diligent observations' at sea was introduced by Luigi
31 Ferdinando Marsili, a member of the Society who stressed the importance of a robust sampling strategy to analyze the
32 'physical nature of the sea'. Marsili's contributions to oceanography were largely recognized again during the 300th
33 anniversary of his birth, thanks primarily to Olson and Olson's article, "Luigi Ferdinando Marsigli, the lost father of
34 oceanography" (1958), as well as Deacon's book (1971).

35 Marsili led a dynamic life as a traveler, soldier, and scientist (see Appendix B for more details). His most renowned work,
36 *Osservazioni Interne al Bosforo Tracio* (1685), outlines the two-layer circulation model that later elucidated the flow into
37 and out of the Gibraltar Strait. Also significant is *Histoire Physique de la Mer* (1725), an extensively illustrated
38 publication offering invaluable historical and scientific insights on the Gulf of Lion (the digital version of this volume
39 can be accessed here: https://www.google.it/books/edition/Histoire_physique_de_la_mer/RpsSxRY4bRcC).

40 Marsili was affiliated with the Paris Académie Royale des Sciences and the Royal Society of London. He greatly admired
41 Robert Boyle but was critical of Boyle's work "De fundo Maris," which he addressed in the first chapter of the *Histoire*,
42 where he asserted the necessity of verifying observations without dismissing sailors' theories. To fully grasp the
43 innovative concepts in the *Histoire*, one must understand the author's discussions with other Académie des Sciences de
44 Paris members. Initially, Marsili believed that stony formations like coral and Madrepores were not true plants (contrary
45 to the views of Joseph Pitton de Tournefort and others), but rather 'drips' created by the sea's viscous material (Archives
46 Départementales de l'Hérault, Registre de la Société royale des sciences de Montpellier, Hérault D. 116; *Memoires de la*
47 *Société Royale in Histoire de la Société royale des sciences établie à Montpellier, 1778*), as noted by McConnell (1990).
48 Through his Observations, Marsili ultimately realized his assumption was mistaken. However, it's crucial to acknowledge
49 his investigative approach, grounded in the Baconian framework of science, which posits that understanding nature begins



50 with collecting and methodically examining facts. To assess coral growth and explore those 'drips' or 'coral milk,' he
51 conducted physical measurements of various sea properties, including sea level, currents, and density.

52 Marsili's goals extended beyond simply determining whether corals were flowers or something different. He sought to
53 create a comprehensive natural history of the sea, its seabed, wind effects, and fish characteristics. His ambition was to
54 compile a significant treatise on the Earth's organic structure. A key component of this research involved studying lakes
55 and sea structures, underpinned by the belief in a morphological and lithological connection between mountains and the
56 seabed, all vital for his theoretical development. 'My intention [...] is to show the organic structure of the Earth' (first
57 chapter of the *Histoire*, dedicated to the marine basin). Marsili explored the Languedoc mountains, but while this research
58 remained unpublished, his investigations of the sea first culminated in the *Brieve Ristretto* (1711), an essay printed in
59 Venice, and later in the *Histoire* of 1725.

60 This is not the first occasion to analyze Marsili's measurements compared to contemporary data. This comparison supports
61 the accuracy and meticulousness of the Bolognese general's data collection methods. The seawater weight data gathered
62 by Marsili in the Constantinople Channel, as detailed in *Osservazioni intorno al Bosforo Tracio*, were examined by
63 Soffientino and Pilson (2005). Their analysis indicated that Marsili's findings were inconsistent with current data because
64 they were treated as salinity data before being adjusted to align with the figures in Zupko's work (Zupko, 1981). In
65 contrast, Pinardi et al. (2018) interpreted the Bosphorus seawater weight data as density data, finding it consistent with
66 modern measurements.

67
68 This paper compares physical data— seawater weight, bathymetric profiles, and sea level variations—found in *Histoire*
69 *Physique de la Mer* to current measurements. This research is part of the MACMAP project (A Multidisciplinary Analysis
70 of Climate Change Indicators in the Mediterranean and Polar Regions), which receives funding from the Istituto Nazionale
71 di Geofisica e Vulcanologia (INGV).

72 Data on seawater weight from the *Histoire Physique de la Mer* were transformed into density values using details from
73 the book's preface (Marsili, 1725). A specific, unnumbered page in the preface states: "*J'ai pesé les Corps solides avec*
74 *la Balance ordinaire, mais très-exacte, me servant de la livre composée de douze onces, l'once de huit dragmes, & la de*
75 *la dragme de soixante grains; & les fluides avec l'Areometre de verre, de la forme, grandeur, & poids que l'on trouvera*
76 *décrits.*" The locations of the sampling stations were determined using maps and georeferenced within a geographic
77 system, facilitating a comparison between Marsili's data and contemporary measurements from the exact locations.
78 Marsili's water samples represent one of the earliest examples of modern field sampling. He employs a methodical
79 approach that aligns with the reproducibility principle of measurements, enabling comparisons between seventeenth-
80 century collected data and current datasets (Pinardi et al., 2018).

81 The bathymetric profiles illustrated in Table III of *Histoire* have been digitized and contrasted with the transects presented
82 in Table II, which have been georeferenced for this study. The findings have been analyzed alongside "modern" data.
83 Variations in sea level listed in Tables X and XI of *Histoire* have also been examined and compared with "modern" data.

84 2. Material and methods

85 2.1. The *Histoire Physique de la Mer*

86 Luigi Ferdinando Marsili's "*Historie physique de la mer*," published in 1725, details the physical, biological, and
87 bathymetric attributes of the Provençal coastal region up to the shelf break. Across all his works, from "*Opus Danubialis*"
88 to his "*Histoire Physique de la mer*," Marsili embraced a humble yet ambitious "Baconian" approach to science. During
89 his tenure as a military engineer, he undertook comprehensive scientific studies that equipped him with a systematic
90 method, which became particularly beneficial after settling on France's southern coast in 1706. While engaged in maritime
91 research, he authored a treatise on the seabed and its waters, dedicating a significant portion to corals and other lithophytes
92 erroneously identified as plants.

93 The *Histoire* was initially printed across 173 pages, featuring 40 plates depicting animals, minerals, fossils, and plants,
94 presented as 12 fold-out plates containing large-scale data, maps of the Gulf of Lion, and coastal profiles. Luigi
95 Ferdinando Marsili employed a strong methodology for obtaining quantitative, well-organized in situ seawater density
96 measurements, including precise time and location specifications. Marsili's primary instruments for data collection
97 consisted of a thermometer, an areometer to assess water weight, a sampler for surface and depth water collection, a
98 "depth gauge," and a metered pole for evaluating sea level variations.



99

2.1.1. Bathymetry and sea level

100 In the first chapter, *Du Bassin* illustrates the bathymetry using graphs that detail the continental shelf's edge. Marsili
101 measured the Gulf of Lion's depth at 14 points, which he represented on a map. These findings allowed him to outline
102 both the seabed and coastal profiles. This volume contains the initial map of the Gulf of Lion, showcasing the bathymetric
103 slope that separates the continental shelf from the abyssal plain (Table I, page 3, titled *Carte du Golfe del Lion entre le*
104 *Cap Sisie en Provence et le Cap de Quiers en Roussillon*). Another map depicts the coastline from Cap Canaille to the
105 Croisette and the nearby islands within Cassis' territory in Provence (Carte Particuliere de la Coste, Table II, page 4).
106 Bathymetric profiles can be found in Table III, page 4 (Profils ou Coupes du Bassin de la Mer), Table IV, page 4 (*Profils*
107 *ou Coupes du Bassin de la Mer sur la Coste de Provence*), and Table V, page 7 (Porte Miou). Neither map features a
108 Coordinate Reference System and they are based on earlier maps. In the volume's preface, Marsili credits the prior
109 research that informed his reconstruction of the Gulf of Lion's bathymetry, including De Basville's study of the Languedoc
110 coast and De Chazelles' map of Provence and Roussillon, who was also an Engineer des Galères and a member of the
111 Paris Academy of Sciences and a hydrography professor in Marseilles.

112 Marsili determined the depth using a traditional method involving a weight on a graduated rope. He heavily relied on
113 measurements provided by fishermen, who would say 'the abyss has no bottom' (l'Abime n'a point de fond) when
114 depths were beyond their measuring lines. Marsili considered this saying inaccurate, as he aligned with the prevailing
115 thought of his time that sea bathymetry reflected terrestrial altimetry.

116 To gauge changes in sea level near a coastal section and the Cassis Sea basin, Marsili employed a metered pole. From
117 January 4 to April 9, 1707, he recorded the water's elevations and depressions at various times. He details his study of
118 sea level variation conducted at Cassis port in the third chapter of *Histoire*.
119

120

2.1.2. Physical characteristics

121 The *Histoire* details temperature and salinity, featuring illustrations of the instruments used and data collection tables.
122 Information on temperature can be found in Table VI, on page 16, titled "*des Experiences fautes avec le Thermometre*
123 *dans la mer à différentes profondeurs*." The sea water temperature was measured using a Florence thermometer, as
124 suggested by Cotte (1774). Marsili likely utilized a three-hook wine-spirit thermometer, similar to those detailed by
125 Camuffo (2020). Measurements were taken at the sea surface and various depths by securing the thermometer to a rope
126 weighted appropriately. Marsili's thermometer was attached to a wooden board, featuring a double scale that is numbered
127 in reverse: 1-55 and 55-1. The thermometer consisted of worked glass supported by three iron wires attached to the board.
128 Two horizontal hooks were standard, located above and below, while an unusual third vertical hook at the top prevented
129 the thermometer from sliding down (Cotte 1774, Camuffo, 2020). Cotte was unable to convert the temperature readings
130 from Marsili's scale to Reaumur units, and sadly, Marsili's thermometer was lost in a maritime conflict in 1707 with an
131 enemy brig or pirate vessel. Consequently, the thermometer's scale remains unknown today, and we lack information
132 about the duration for which the instrument was submerged, despite Marsili's meticulous record of the time immersion.

133 In Marsili's time, the concept of water density was understood in terms of sea gravity. As noted by Manzella et al. (2021),
134 measurements of gravity and salinity were conducted using a known-sized vial with a narrow neck or a graduated glass
135 tube. Gravity was assessed by weighing the water, while salinity was calculated based on the weight of the residue left
136 after evaporation.

137 The water weight was gauged using the hydrostatic ampoule (also known as the hydrostatic carafe or areometer, see
138 Locritani & Garvani, 2024 and Appendix B), which consists of a sealed glass sphere with a tapering neck (Montanari,
139 1696). Viviani, a student of Galileo, described the measurement process: the ampoules needed to include enough lead
140 flakes so they would float in the liquid being assessed. Additional known-weight rings were placed on the neck until the
141 ampoule was submerged. The weight of these rings that caused the ampoule to sink equaled the weight of the measured
142 water. For further details, see Pinardi et al. (2018). Sea water samples were collected using a wooden container sealed
143 with a valve. Marsili first weighed the sea water onboard the vessel and then used a balance later in a lab. The reference
144 water was sourced from a well near Marsili's lab in Cassis. Marsili gathered water samples between Cap Canaille and Cap
145 Croisette, which he subsequently analyzed in his lab using a thermometer and the hydrostatic ampoule or balance. The
146 hydrostatic method involved measuring an object's weight by submerging it in a liquid of known weight. The second
147 chapter includes tables and figures that present data on the weight of salty and fresh waters, a description of the areometer
148 or hydrostatic ampoule, and the salt concentration in the water samples.



149 **2.1.3. Vegetation**

150 The fourth chapter of the *Histoire*, titled *De la vegetation des plantes*, is the largest section. It features 40 copper-engraved illustrations
 151 by Matthys Pool (1676-1740), depicting animals, minerals, fossils, and plants, all numbered with corresponding references in the text.
 152 This section includes many plates that portray corals, which Marsili classified as "plants." Each plate indicates the location where the
 153 corals were discovered and describes how they were gathered by fishermen, along with the local names or those assigned by Marsili,
 154 as Linnaeus' nomenclature had not yet been adopted.

155 **3. Data conversion**

156 **3.1. Bathymetry**

157 The initial step in extracting historically significant information from historical-geographical maps involves defining a
 158 set of immutable and identifiable ground control points (GCPs) for georeferencing these maps in QGIS and then digitizing
 159 the sampling points. The scale of the Marsili maps is expressed in "toises" or "teses", which correspond to 1.2 brasses
 160 marine (fathoms), an ancient unit of length roughly equal to 0.32 feet or 1.95 meters, and according to Angelo Martini's
 161 metrology manual, 1 meter is approximately 0.001 mile marine (Martini, 1881). The direction and length of the transect
 162 propagation is measured in miles, nearly equivalent to nautical miles; specifically, 1 marine mile equals 1851 meters, as
 163 noted by Martini (1881, p. 466). In the initial phase, distances from the Marsili maps were obtained using Adobe Illustrator
 164 2022 and compared with those in contemporary maps (Google Satellite in QGIS 3.22). The mean, standard deviation, and
 165 root mean square for distances between pairs of GCPs from both the Marsili and Google Satellite maps were calculated
 166 in QGIS 3.22. Following recommendations by Bitterer (2006) and Hvizdák (2023), a statistically adequate sample
 167 requires GCPs greater than 30, which poses challenges with historical maps. To align the ancient and modern maps more
 168 closely, a restricted area near Marseille (from Toulon to Port de Bouc) was selected, resulting in 8 GCPs being recorded
 169 and an additional 4 GCPs zoomed in on the Cassis area. The second map was entirely georeferenced using these 8 GCPs.
 170 The coordinates of the GCPs are detailed in Table 1. Subsequently, the maps were georeferenced in QGIS 3.22 applying
 171 "Thin Plate Spline (TPS)" interpolation and the "Nearest Neighbour" sampling method.

Cassis form Pl II pag 4		
GCP	Latitude [°N]	Longitude [°E]
1	43.184196°	5.563402°
2	43.207667°	5.368603°
3	43.212021°	5.539110°
4	43.211178°	5.337976°
5	43.175865°	5.382420°
6	43.203079°	5.511857°
7	43.204568°	5.426200°
8	43.203280°	5.452809°
Marseille from Pl I pag 3		
GCP	Latitude [°N]	Longitude [°E]
1	43.278345°	4.890243°
2	43.282043°	5.346703°
3	43.279782°	5.325148°
4	43.393690°	4.985789°
5	43.045905°	5.859005°
6	43.213436°	5.337643°
7	43.341584°	5.265206°
8	42.880143°	5.308412°
Cassis from from Pl I pag 3		
GCP	Latitude [°N]	Longitude [°E]
1	43.214906°	5.336222°
2	43.160508°	5.607515°
3	43.173813°	5.398702°
4	43.209136°	5.539286°

172
 173 **Table 1: GCP used to georeferenced the Marsili maps (Gulf of Lion map - Pl. I, page 3 in the Marseille coastal area and the**
 174 **Cassis coastal area and Cassis map - Pl. II, page 4) in QGIS 3.2.**

175 Additionally, the EMODnet_satellite_coastline_MSL has been superimposed onto the georeferenced historical
 176 maps. Upon completing this georeferentiation, we could extrapolate the coordinates of the small urns and the
 177 positions of the transects.

178 The bathymetric profiles were digitized using WebPlotDigitizer (Rohatgi, 2017). The profile length in nautical
 179 miles (nmi) is indicated in the legend at the upper-right corner of the original figure. This facilitated the
 180 conversion of the horizontal distance from pixels to nautical miles. The maps display lines representing
 181 transects along which Marsili conducted bathymetric measurements. Descriptions in the *Histoire* tables
 182 provide starting points, total transect lengths, and depth measurements in fathoms in a few locations (*brasses*;



183 1 ftm is 1.624197 m, Martini, 1881); this data was used to convert the vertical distance from pixels to meters.
184 Some profiles lacked depth information; therefore, they were digitized in adimensional units (zero for sea level
185 and -1 for the profile's maximum depth). To compare with current bathymetry, the spatial coordinates of the
186 georeferenced transects from the maps were utilized to extrapolate bathymetric values from EmodNET
187 (<https://emodnet.ec.europa.eu/geoviewer/>) and Gebco 2023 (<https://download.gebco.net/>) data in QGIS 3.22.
188

189 3.2. Sea Level in Cassis

190 The Marsili sea level measurements were conducted not in a well but likely in a protected coastal area using a 68 pouces
191 rod, referred to as an inch rod by Marsili (1 French inch is roughly 2.7 cm, according to Martini, 1881). A timing issue
192 exists; the measurements lacked precise timestamps (such as sunrise, noon, sunset around 9 pm, and midnight). For the
193 analysis, measurements from sunrise to sunset were assumed to be recorded at roughly 6:45 am, noon, 6:45 pm, 9 pm,
194 and midnight. It is also assumed that each time point included two measurements (maximum and minimum) to mitigate
195 wave effects through averaging.

196 A data series was created using Marsili's data from January 5 to April 10, 1707. The minimum and maximum values were
197 averaged at each time interval, and then the mean was subtracted. The resultant series exhibited significant noise, leading
198 to the application of multiple filters to identify key signals. A primary low-pass filter was introduced to smooth the time
199 series. Notably, periodicities of about 5-7 days and 13-14 days were observed, which may relate to atmospheric influences
200 (Esposito and Manzella, 1982).

201 3.3. Density

202 Water weight measurements were recorded in ounces, drachmas, and grains. Maps marked the water sampling sites using
203 the amphora symbol. To convert the weight to Kg/m^3 , it was assumed that 1 pound corresponds to 12 ounces, 1 ounce to
204 8 drachmas, and 1 drachma to 60 grains (Marsili, 1725). The transformation formula used to convert weight to density is
205 $\rho = 1000 * P / Pr$, where ρ represents the density of water in Kg/m^3 ; 1000 is the density of distilled water in Kg/m^3 ; P is the
206 weight of seawater in grains as measured by Marsili; and Pr is the weight of reference water in grains from Marsili's
207 measurements. The minimum potential error in the measurements was assessed by considering values below 1000 Kg/m^3
208 for freshwater (such as rivers, fountains, and wells). This data has been compared with the water density measurements
209 from SeaDataNet covering the same sites from 1990 to 2018.

210 4. Results and Discussion

211 4.1. Bathymetry: historical map analysis

212 Marsili's maps include scales that indicate distances of 10000, 5000, 1000, 500, or 100 "teses". Due to the numerous
213 variations, it was necessary to evaluate the consistency within each map and then compute the distances. Table 1 lists 20
214 Ground Control Points (GCPs) selected for distance calculations from the Marsili maps, showing the relative distances
215 measured between pairs of GCPs in both maps, along with the differences in lengths (errors). Table 2 presents the Root
216 Mean Square Error (RMSE), as well as the means and standard deviations. The RMSE for the Gulf du Lion map is 6087
217 m, while for the Cassis map, it is 548 m. The mean distances are 2912 ± 6057 m for the Gulf du Lion map and 324 ± 472
218 m for the Cassis map. The errors are uneven, evidenced by the standard deviations exceeding the mean values. Improved
219 results are achieved by remapping the coastline and bathymetric lines for more localized areas (refer to Figure A1 in
220 Appendix A). This adjustment reduces the error, resulting in a comparison with EMODnet coastlines and bathymetric
221 lines showing an error of 631.38 ± 559.72 m for the Gulf of Lion and 163.83 ± 157.33 m for the Cassis area.



Site name	Latitude start [°N]	Longitude start [°E]	Latitude end [°N]	Longitude end [°E]	Lenght in Marsili map [m]	Lenght in Google Satellite [m]	Error beetween new - old
Map of Cassis coastal area							
Ile Riou	43,179382°	5,371768°	43,173686°	5,398688°	2006,10	2274,00	267,90
Ile maire	43,212390°	5,328571°	43,209194°	5,340005°	715,52	996,71	281,19
Cap Croisette-Cassis	43,214517°	5,327929°	43,214658°	5,537063°	14872,38	16333,00	1460,62
Calanque de Port Pin	43,197710°	5,509915°	43,203919°	5,510848°	667,24	695,00	27,76
Cap Croisette	43,214517°	5,327929°	43,214993°	5,336232°	570,66	863,72	293,06
Ile Calseraigne	43,190199°	5,381306°	43,186181°	5,391561°	978,91	948,00	-30,91
Ile Jarre	43,201178°	5,355180°	43,193853°	5,371627°	1400,32	1565,00	164,68
Port Pin	43,204062°	5,513849°	43,211172°	5,521307°	930,62	997,00	66,38
Map of Gulf of Lion							
Ile Riou	43,179382°	5,371768°	43,173686°	5,398688°	2075,86	2274,00	198,14
Ile maire	43,212390°	5,328571°	43,209194°	5,340005°	1579,92	996,71	-583,21
Cap Croisette - Cassis	43,214517°	5,327929°	43,214658°	5,537063°	16953,95	16333,00	-620,95
Cap Croisette - Cap Creus	42,319361°	3,322315°	43,214993°	5,336232°	175533,48	192526,00	16992,52
Ile Pommegue	43,276142°	5,310985°	43,262178°	5,287012°	3621,07	2501,00	-1120,07
Ile Ratonneau	43,286226°	5,323944°	43,280135°	5,291584°	4781,82	2717,00	-2064,82
Ile Planier	43,197204°	5,228365°	43,198999°	5,231435°	1242,18	317,75	-924,43
Cap Creus - Brescon	42,319361°	3,322315°	43,263251°	3,501627°	105101,15	105896,00	794,85


222

223 **Table 2: Comparison between the lengths measured in Marsili maps (Gulf of Lion map - Pl. I, page 3 and Cassis map - Pl. II,**
 224 **page 4) and modern map (Google Satellite in QGIS 3.2). The first column listed the names of the 16 sites (the first 8 in Cassis**
 225 **map - Pl. II, page 4 and the second eight in the Gulf du Lion map - Pl. I, page 3). The second and fifth columns indicate the**
 226 **starting and ending points coordinates, the sixth and seventh columns indicate the lengths measured in Marsili and Google**
 227 **Satellite maps, and the last column reports the relative difference between lengths.**

228

4.2. Bathymetry: Profiles


229 Figure 1 illustrates the transects design from Marsili's maps found in Histoire (Table I) and reconstructed as a bathymetric
 230 profile in Table III of Histoire. Figure 2 showcases a comparative analysis of the depth profiles recorded by Marsili,
 231 converted into meters, alongside those obtained from the EMODnet dataset. One significant observation is that Marsili
 232 underestimates depths greater than 100 meters. This discrepancy likely stems from the limitations of the measurement
 233 techniques available during his era, which impeded accurate depth readings. Additionally, our attempts to georeference
 234 Marsili's maps unveiled more challenges; the spatial alignment of his depth profiles does not correctly match the
 235 geographical features reflected in the EMODnet data. For example, in Marsili's profile F6, a straight line intersection with
 236 both Les Iles and Planier islands reveals an inconsistency, as such a cross-sectional view would not realistically occur in
 237 mapping scenarios like those of EMODnet. Marsili's depiction of seafloor features notably shifts towards the coastline.
 238 This issue is especially apparent in profiles E5 and G7, as shown in Figures 2a-c.



239

240 **Figure 1: The map shows the points (P4, P5, P6) in which anfores have been drawn in the Marsili maps on *Histoire* (Table I**
241 **and II). The anfores represent the sampling points for water weight measurements. Moreover, the figure shows the transects**
242 **(A1, B2, C3, D4, E5, F6, G7, H8, I9, L10, M11, N12) drawn in the Marsili maps on *Histoire* (Table I). The map has been**
243 **extrapolated to Google Satellite in QGIS 3.2.**

244 Despite these differences, Marsili has accurately pinpointed certain geographical features. For example, he clearly
245 outlined the continental shelf in profiles E5 and F6 (see Figures 2a-b) and recognized a canyon in profile H8 (Figure 2d).
246 These correspondences add credibility to Marsili's work, indicating that although his methods might have led to some
247 inaccuracies, he effectively captured essential elements of the seafloor topology that are still pertinent today.



248

249 **Figure 2: Comparison between matching EMODnet depth profiles and a) profile E5, b) profile F6, c) profile G7 and d) profile**
250 **H8 extracted from the figures shown in page 3 and page 4 of Marsili's work.**

251

4.3. Sea Level in Cassis

252 Applying a bandpass filter to the dataset helps isolate the primary tidal components (M2 and K1) during specific times,
253 such as the first two weeks of March, revealing oscillations consistent with the M2 tides. The resulting amplitude measures
254 around 10-12 cm, surpassing the 6.8 cm determined by Woppelmann et al. (2016) in Marseille, which is close to Cassis.
255 While the values obtained are credible, a more refined methodology and thorough analysis are necessary for accurately
256 extracting tidal components from the Marsili data.

257

4.4. The water weight measurements

258 Seawater density is fundamentally tied to the technologies and methods employed in its calculation. One of the earliest
259 instruments to gauge seawater's 'gravity' or weight was the explorator qualitum. Following this, specific gravity was
260 evaluated, which is the weight of seawater relative to an equal volume of freshwater. A significant issue arose from the
261 choice of sample water. Hooke (1635 - 1703) conducted specific gravity measurements using Thames water collected at
262 low tide in Greenwich (Derham, 1726), while Marsili opted for well water. Phipps became the first to utilize distilled
263 water during his Voyage toward the North Pole in 1774. Manzella and Novellino (2022) provide descriptions of the
264 methodologies and technologies for measuring seawater density.

265 Using a 'non-standard' reference water can introduce errors that must be considered in data analysis. To assess the quality
266 of the Marsili data, a comparison was made with measurements from recent years. Historical data from the Mediterranean
267 Sea, spanning from 1806 to 2022, was selected based on proximity to the Marsili measurement points. The TEOS-10
268 framework was employed to calculate the physical properties of seawater using measurements taken between 1990 and
269 2018.

270 The historical archive spans the Mediterranean Sea from 1864 to 2022. Initially, the archive was gathered by the former
271 Italian National Committee for Nuclear Energy (CNEN) through the Center National pour l'Exploitation des Océans
272 (Cnexo), which later merged with the Institut scientifique et technique des pêches maritimes (ISTPM) to form the current
273 L'Institut français de recherche pour l'exploitation de la mer (Ifremer). CNEN subsequently provided the initial archive
274 to the Mediterranean Oceanographic Data Base (MODB) project in the early 1990s. After three years, the MODB data



275 were incorporated into the Mediterranean Data Archeology and Rescue (MEDAR/MedAtlas) project, which also
 276 integrated additional data. This collective dataset then fed into the SeaDataNet system, where more information was
 277 added. Besides the SeaDataNet entries, the archive includes data from NESDIS/NOAA and public repositories like
 278 SEANOE and PANGAEA, alongside contributions based on personal knowledge. It should be noted that the archive does
 279 not fully cover all positions for the Marsili observations; where there was considerable variability, minimum and
 280 maximum values are provided.

281 Table 3 presents water density converted from Histoire tables. Data were gathered at various locations in the Gulf of Lion
 282 between June 18, 1706, and January 18, 1707 (illustrated in Figure 1 and Figure 3 with green points). The minimum
 283 possible measurement error has been calculated, yielding results of $\pm 1.23 \text{ Kg/m}^3$.


284

Latitude [°N]	Longitude[°E]	Site	Data [month day year]	Sea surface water density [Km/m3]	SeaDataNet density (from 1990 to 2018) [Kg/m3]
43.277825°	5.312381°	Marseille islands	June 18th 1706	1028,26087	1014 - 1028,7
43.395628°	4.985985°	Port the Bouc	June 13rd 1706	1026,09	
43.248569°	4.844061°	Opposite the mouth of the grand Rosne 5 miles offshore	July 14th 1706	1021,01	
43.371753°	4.600481°	from Chemin du Rosne to Cette in linea retta	July 14th 1706	1024,64	
43.388558°	3.690230°	Port of Cette	July 15th 1706	1028,26	
43.313049°	4.655259°	At the mouth of the little Rosne into the sea	October 26th 1706	1009,06	1014
43.211923°	5.533302°	At the mouth of the port of Cassis	December 4th 1706	1027,54	1027 - 1028,8
43.199698°	5.448772°	To the great Chandelle	December 7th 1706	1028,99	
43.211923°	5.533302°	At the mouth of the Port of Cassis during the storm		1028,62	
43.195922°	5.500447°	Castello Vieux	January 14th 1707	1027,54	1027 - 1028,8
43.184370°	5.551525°	To Cassidagne	January 18th 1707	1030,43	
43.204045°	5.513703°	At Port Miou where the river water flows into the sea	January 18th 1707	1015,94	

285

286 **Table 3: The table shows the sampling points coordinates (the anfores in Marsili maps) in which Marsili took water samples to**
 287 **measure water weight. Each coordinate is associated with a site and a date indicated by Marsili in Pl. VII pag. 23. The following**
 288 **columns show the measure of water weight converted in water surface density by Marsili data and the average surface water**
 289 **density extrapolated by SeaDataNET data (from 1990 to 2018).**

290 Table 3 indicates that Marsili's measurements align well with the latest data. The density value of 1030.43 calculated on
 291 January 18, 1707, is relatively high yet still falls within the range of the estimated minimum error.



292

293 **Figure 3: The map shows the points (P1, P2, P3, P7, P8, P9, P10, P11, P12) in which anfores have been drawn in the Marsili**
294 **maps on *Histoire* (Table I and II). The anfores represent the sampling points for water weight measurements.**

295

296 5. Data Availability

297 The data are available at:

298 <http://oceanbo.bo.ingv.it/erddap/search/index.html?page=1&itemsPerPage=1000&searchFor=cassid> (Locritani et al.,
299 2024a).

300 <http://oceanbo.bo.ingv.it/erddap/search/index.html?page=1&itemsPerPage=1000&searchFor=water+density+marsili>
301 (Locritani M. & Garvani S., 2024).

302 <http://oceanbo.bo.ingv.it/erddap/search/index.html?page=1&itemsPerPage=1000&searchFor=marsili> (Locritani et al.,
303 20224b).

304 6. Conclusion

305 This study presents a comprehensive analysis of eighteenth-century maritime observations in the Gulf of Lion, focusing
306 on data derived from the significant historical text by Marsili. Collected between 1706 and 1707, this dataset represents
307 one of the earliest systematic efforts to document the oceanographic characteristics of the region.

308 While Marsili's data collection methods do not adhere to contemporary standards due to the instrumentation limitations
309 and the accuracy of units available in his time, we have taken steps to evaluate the relevance of his findings. We aim to
310 contextualize the results within modern frameworks by applying error margins to the various measurements. For instance,
311 several parameters, including the weight of surface seawater, were in reasonable agreement with mean values observed
312 over the last two centuries, suggesting that Marsili's observations hold some validity despite methodological limitations.

313 Conversely, data related to tidal patterns requires more rigorous methodologies and analyses. The precision of Marsili's
314 maps was variable, with minimum errors estimated at around 160 meters, indicating potential discrepancies in geographic
315 representation. These inaccuracies highlight the challenges faced by early oceanographers, yet they also underscore the
316 pioneering nature of Marsili's work.



317 Despite these limitations, Marsili's contributions remain invaluable. His work not only provided the first systematic
318 description of the Gulf of Lion's continental shelf and abyssal regions but also laid the groundwork for future
319 oceanographic studies in the area. By bridging the historical insights and contemporary analysis, this study reaffirms the
320 significance of early oceanographic research and its relevance to our understanding of marine environments today.
321 Through this exploration, we honour Marsili's legacy while acknowledging the evolution of oceanographic methodologies
322 over the centuries.

323

324 **Author contribution:** ML conceptualized and administrated the research, validated the georeferenced maps and wrote
325 the original draft preparation, SG investigated the historical part of the project and wrote original draft preparation, GM
326 validated sea level data and supervised the entire study and wrote review and editing of the text, GT validated the
327 bathymetric data and wrote original draft preparation, AG providing funding acquisition as MACMAP project
328 coordinator.

329 **Competing interests:** Author GM is a member of the editorial board of the journal.

330 **Publisher's note:** Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and
331 institutional affiliations.

332 **Acknowledgements:** The authors are deeply indebted to the owner of the volume *Histoire physique de la mer* (Marsili,
333 1725) allowed to study on the original book. The authors are grateful to Grammarly and CatGPT who helped correct the
334 grammar of the entire paper and rephrase the sentences in the conclusion section.

335 **Financial support:** This research has been developed in the framework of the MACMAP (A Multidisciplinary Analysis
336 of Climate change indicators in the Mediterranean And Polar regions) project funded by Istituto Nazionale di Geofisica
337 e Vulcanologia (Environment Department), ([https://progetti.ingv.it/index.php/it/progetti-
338 dipartimentali/ambiente/macmap](https://progetti.ingv.it/index.php/it/progetti-dipartimentali/ambiente/macmap)).

339

340 References

- 341 Archives Départementales de l'Hérault, Registre de la Société royale des sciences de Montpellier, Hérault D. 116
- 342 Bitterer, L. *Vyrovňovací Počet*; Žilinská Univerzita v Žiline: Žilina, Slovakia, 280 p, 2006.
- 343 Boyle R., *Observations And Experiments About The Saltiness Of The Sea*, In tracts consisting of observations about the
344 saltiness of the sea: an account of a statical hygroscope and its uses, London, R. Davis, pp. 1-40, 1674.
- 345 Camuffo, D., Key problems in early wine-spirit thermometers and the “true Réaumur” thermometer. *Climatic Change*.
346 163. 1-20. 10.1007/s10584-020-02910-3, 2020.
- 347 Carpine Lancre J., McConnell A., Le comte L.F. Marsigli et la Société royale des sciences de Montpellier, «Actes du
348 110e Congrès national des sociétés savantes. Montpellier 1985», Paris, C.T.H.S, I, pp. 33-44, IT\ICCU\UBO\4690917,
349 1985.
- 350 Cotte, L., *Traité de Météorologie*, contenant 1. l'Histoire des Observations Meteorologiques. 2. Un Traite des Meteores.
351 3. l'Histoire et la description du Barometre, du Thermometre, et des autres Instrumens meteorologiques. 4. Les Tables des
352 Observations meteorolo, De l'imprimerie Royale, A Paris, 1774, IT\ICCU\NAPE\01448.
- 353 Deacon, M., *Scientists and the Sea, 1650-1900: a study of marine science*, London-New York, Academic Press,
354 IT\ICCU\UBO\0294370, ISBN1859283527, 1971.
- 355 Derham, F.R.S., *Philosophical Experiments and Observation of the Late Eminent Dr. Robert Hook S.R.S. And Geom.*
356 *Prof. Gresh and Other Eminent Virtuoso's in His Time*. Royal Society, 403 p, IT\ICCU\UBOE\109572 , 1726.



- 357 Esposito, A., Manzella G., Current Circulation in the Ligurian Sea, Editor(s): Jacques C.J. Nihoul, Elsevier Oceanography
358 Series, Elsevier, Volume 34, 187-203 pp., ISSN 0422-9894, ISBN 9780444420770, <https://doi.org/10.1016/S0422->
359 [9894\(08\)71245-5](https://doi.org/10.1016/S0422-9894(08)71245-5), 1982.
- 360 Hvizdák, L., Tometzová D., Iannaccone B., Šoltésová M., Domaracká L., and Kyšef'a K., Old Mine Map Georeferencing:
361 Case of Marsigli's 1696 Map of the Smolník Mines, ISPRS International Journal of Geo-Information 12, no. 8: 345.
362 <https://doi.org/10.3390/ijgi12080345>, 2023.
- 363 Locritani M., Garvani S., Manzella G., Sea Level data reported in Histoire Physique de la mer (1725) of L.F. Marsili
364 and transformed to calculate the M2 tidal amplitude. Istituto Nazionale di Geofisica e Vulcanologia (INGV), ERDDAP,
365 <https://doi.org/10.13127/histoiremarsili/sealevel>, [data set], 2024a.
- 366 Locritani, M. and Garvani S., Water density data extrapolated by water weight measurements reported in Histoire
367 Physique de la mer (1725) of L.F. Marsili. Istituto Nazionale di Geofisica e Vulcanologia (INGV), ERDDAP,
368 <https://progetti.ingv.it/it/progetti-dipartimentali/ambiente/macmap#water-density-l-f-marsili>, [data set], 2024.
- 369 Locritani M., Garvani S., Tamburello G., Fratianni C. (2024). Bathymetric measurements reported in Histoire Physique
370 de la mer (1725) of L.F. Marsili. Istituto Nazionale di Geofisica e Vulcanologia (INGV), ERDDAP,
371 <https://progetti.ingv.it/it/progetti-dipartimentali/ambiente/macmap#datasets>, [data set], 2024b.
- 372 Directions for Observations and Experiments to Be Made by Masters of Ships, Pilots, and Other Fit Persons in Their Sea-
373 Voyages., Philosophical Transactions (1665-1678), vol. 2, pp. 433–48. JSTOR, <http://www.jstor.org/stable/101197>.
374 Accessed 14 Oct. 2024, 1666.
- 375 Lettera postuma di Geminiano Montanari, Istruzione scritta gia' da tempo all'illustrissimo et eccellentissimo sig. Pietro
376 Grimani dal Signor Geminiano Montanari, primo professore delle matematiche ... sopra il modo di conoscere la
377 differenza del peso de' fluidi fra loro. Pubblicata in La Galeria di Minerva overo Notizie Universali, Girolamo Albrizzi,
378 390 p., 1696.
- 379 Manzella, G., Novellino A., Ocean Science Data, Collection, Managements, Networking and Services, Elsevier,
380 Radarweg 29, PO Box 211, 1000 AE Amsterdam, Netherlands, 2022.
- 381 Marsili L. F., Osservazioni intorno al Bosforo Tracio overo Canale di Costantinopoli, rappresentate in lettera alla sacra
382 real maestà di Cristina regina di Svezia, Roma, Nicolò Angelo Tinassi, IT\ICCU\UFIE\001269, 1681.
- 383 Marsili L. F., Brieve ristretto del saggio fisico intorno alla storia del mare scritta alla Regia Accademia delle Scienze di
384 Parigi. Ora esposto in una lettera all'eccellentiss. signor Cristino Martinelli, Venezia, presso Andrea Poletti,
385 IT\ICCU\RMSE\000883, 1711.
- 386 Marsili L. F., Histoire physique de la mer. Ouvrage enrichi de figures dessinées d'après le naturel. Par Louis Ferdinand
387 comte de Marsilli, membre de l'Academie royale des sciences de Paris, A Amsterdam aux de'pens de la Compagnie,
388 IT\ICCU\PUVE\000546, 1725.
- 389 Martini, A., Manuale di metrologia, Torino, Lotscher, IT\ICCU\SBL\0052845, 1881.
- 390 McConnell A., The flowers of coral. Some unpublished conflicts from Montpellier and Paris during the early 18th century,
391 «History and Philosophy of Life Science», XII, 51-66 pp., IT\ICCU\UBO\1798875, 1990.
- 392 Memoires de la Société Royale in Histoire de la Société royale des sciences établie à Montpellier avec les mémoires de
393 mathématiques et de physique, tirés des registres de cette société, tome second, 456 p., IT\ICCU\TOO\0185632, 1778.
- 394 Olson, F. C. W., Olson M. A., Luigi Ferdinando Marsigli, The lost father of oceanography, «Quarterly Journal of the
395 Florida Academy of Sciences», Vol. 21, No. 3, 227-234 pp., JSTOR, <http://www.jstor.org/stable/24315236>. Accessed 4
396 Nov. 2024, September 1958.
- 397 Phipps, C. J., A Voyage Towards the North Pole Undertaken by His Majesty's Command 1773, Published by London
398 Printed by W. Bowyer and J. Nichols, for J. Nourse, Bookseller to His Majes, IT\ICCU\LIGE\009473, 1774.



399 Pinardi, N., Özsoy E., Latif M.A, Moroni F., Grandi A., Manzella G., De Strobel F., and Lyubartsev V., Measuring the
400 Sea: Marsili's Oceanographic Cruise (1679–80) and the Roots of Oceanography, *J. Phys. Oceanogr.*, 48, 845–860, doi:
401 10.1175/JPO-D-17-0168.1, 2018.

402 Rohatgi, A., WebPlotDigitizer 4.0, <https://automeris.io/WebPlotDigitizer/>, 2017.

403 Soffientino, B., Pilson M.E.Q., The Bosphorus Strait: A special place in the history of oceanography, «*Oceanography*»,
404 Vol.18,No.2, 16–23 pp., <https://doi.org/10.5670/oceanog.2005.38>, 2005.

405 Zupko, R. E., Italian Weights and Measures from the Middle Ages to the Nineteenth Century, American Philosophical
406 Society, Memoir 145, Philadelphia, PA, 179 p., IT\ICCU\PUV\0197644, 1981.

407

408 Appendix A



409

410 **Figure A1: Comparison of three coastlines in the area near Cassis. The dotted green one is the coastline extrapolated from**
411 **EMODnet data, the purple line represents the digitised coastline derived from the historical map of the Gulf of Lion. In**
412 **contrast, the white line depicts the coastline extrapolated from the historical map of Cassis. The map has been extrapolated to**
413 **Google Satellite in QGIS 3.2.**

414 Appendix B

415 **Marsili biography and the scientific contest of reference**

416 Luigi Ferdinando Marsigli (Bologna, 1658-1730) was a natural scientist and soldier whose remarkable and idiosyncratic
417 accomplishments in both fields warrant recognition. During his youth, Marsigli accompanied his father and other family
418 members on various journeys, which allowed him to become acquainted with numerous distinguished scholars, including
419 the renowned librarian Antonio Magliabechi (1633-1714), the medical philosopher Tomaso Cornelio (1614-1684), and
420 Giovanni Borelli (1608-1679). These formative experiences fostered in Marsigli a profound and enduring interest in the



421 marvels of the natural world. Marsigli pursued extensive studies, albeit with irregular intervals. At the University of
422 Bologna, he studied under the anatomist Marcello Malpighi (1628-1694), the botanist Lelio Trionfetti (1647-1722), and
423 the mathematician and astronomer Geminiano Montanari (1632-1687), with whom he maintained a regular
424 correspondence. As a result of the influence of his teachers, Marsigli was able to embrace the contemporary principles,
425 methodologies and approaches of the Royal Society of London. In particular, Marsigli was profoundly influenced by the
426 instructions on experimentation and instrumentation that he received from his mentor, Geminiano Montanari. In 1665,
427 Montanari established the Accademia della Traccia or dei Filosofi in Bologna. This institution combined mathematical
428 and experimental approaches and emphasised the instrumentation, discoveries, and experiments conducted by the Royal
429 Society. Marsigli's expertise was also significantly shaped by one of the founders of the Royal Society. Robert Boyle
430 (1627-1691) operationalised Marsigli's procedure by translating the intentions expressed by Boyle in his treatises (*Tracts*
431 *consisting of Observations about the Saltness of the Sea*, London, 1674), this involved adapting the procedure to the
432 specific context.

433 Marsigli's first book of 1681, *Osservazioni intorno al Bosforo Tracio*, dedicated to Queen Christina of Sweden, reported
434 observations and experiments he had carried out during and shortly after his year-long visit to Istanbul in 1679 and 1680.
435 While this work is recognised by oceanographers as the first description of a simple laboratory-scale hydrological
436 experiment to show how differences in the density of two water masses lead to two-layer currents, another treatise by
437 Marsigli consecrated him as the lost father of oceanography (Olson, 1958): *Histoire physique de la mer*, published in 1725.

438 This remarkable treatise was written during the time Marsigli spent in France. After a few years of training in chemistry in
439 Paris (1704-1706), Marsigli expressed his desire to study the organic structure of the Earth. He first travelled to
440 Montpellier, where he was accepted as a member of the newly formed 'Societe royale', modelled on the Paris Academy
441 of Science (Carpine-Lancre and McConnell, 1985), during which time he contributed by providing a list of valuable
442 correspondents and suggesting essential books and journals for the Society to acquire. He illustrated the members of his
443 research with watercolour illustrations, including a location map, a sectional drawing of the seabed showing where the
444 coral was dredged and the apparatus used. At first, Marsigli thought it was a mineral concretion like those he had seen in
445 petrified springs. The true nature of coral - mineral, plant or animal - was debated from antiquity until the late 18th
446 century. After Montpellier, Marsigli moved on to Cassis, where he regularly joined the coral fishermen, taking
447 temperature readings, making soundings and collecting water samples. Marsigli observed the tides, studied waves and
448 currents and documented the colours of the water. In his laboratory on land, he measured the gravity of seawater samples,
449 analysed corals and used a microscope to study their structure and function.

450 In a letter to Abbé M. Bignon, he outlined his research intentions: "I have begun research on the history of the sea, where
451 I hope to treat the nature of the water of the sea and its diverse movements; of the differences of the bottoms of the sea,
452 which seem to me to be related to the structure of the mountains, of the effect of winds on this water, of the nature of fish
453 developed through analysis of the vegetation growing on the bottom of the sea."

454 This productive period, spanning a mere two years, was interrupted by Marsigli's second call to military service, this time
455 by Pope Clement XI. In 1708, Marsigli briefly departed for the Adriatic coast to command the Papal States' troops, seizing
456 the opportunity to conduct limited oceanographic studies in those waters. Upon his return to Cassis, he commenced work
457 on his treatise, publishing a brief extract in 1711: *Brieve ristretto del Saggio fisico*. In 1715, he was once more summoned
458 to serve the Pope. This was to be his final deployment. Upon the cessation of hostilities, he relinquished his military
459 obligations and subsequently dedicated the remainder of his life to the pursuit of his studies. His extensive collection of
460 scientific materials was donated to his hometown of Bologna on 13 March 1714, which marked the establishment of the
461 Bologna Institute of Science and Art. In 1715, he was elected to the Paris Academy of Sciences. He subsequently travelled
462 to England, where he formed a friendship with Newton and Halley. In 1722, he was elected a Fellow of the Royal Society.
463 The results of his two-year research at Cassis were published in 1725 in the form of his monumental work, *Histoire*
464 *Physique de la Mer*, which he dedicated to the Academy of Sciences in Paris. The work was eventually published in
465 Amsterdam in 1725.

466 Areometer or hydrostatic ampulla Note:

467 The instrument employed for the measurement of weight in both surface and deep water is designated as a hydrostatic
468 ampulla, hydrostatic bulb, or areometer. The description of an instrument called a "hydrostatic ampulla," constructed
469 following Montanari's technical instructions and closely resembling the apparatus utilized by Marsigli in the Sea of
470 Provence, can be found in a posthumous publication by Montanari in the cultural journal "La Galleria di Minerva" in
471 1704 (Montanari, 1696). The water weight instrument, which consisted of a small, long-necked bottle weighted with
472 sufficient lead shot to enable it to float upright, was used by Marsigli in the Bosphorus. This instrument was purchased by
473 Montanari himself (Soffientino & Pilon, 2005). It is noteworthy that Marsigli himself stated regarding the use of the
474 hydrostatic bulb that: "*l'altra parte che compone la natura di queste acque è il sapore salso, universale a tutti i mari che*
475 *più, e meno sono di esso abbondanti, come l'esperienza me l'ha mostrato mediante il peso rilevato con l'Ampolla*



476 *Idrostatica, secondo gli insegnamenti datimi dal Sig. Dott. Montanari, celebre matematico, e mio riverito Maestro, che*
477 *fin da primi anni cominciò a dimostrarmi i principi di simili studi, e tra gli altri, il fondamento dell'uso di tale Istromento,*
478 *e la perfezione alla quale col suo nobile ingegno l'ha ridotto; e ne conservo una lettera in cui se ne parla diffusamente”*
479 (Marsili, 1681, pag. 71). In the *Histoire Physique de la mer*, the areometer is illustrated in Plate VII, accompanied by its
480 weight in air (1 ounce, 3 drachms and 10 grains) and the lead rings with the corresponding weights. The ampoule is used
481 to measure density, which was also referred to as 'gravity'. When Marsili measures a weight, he is measuring gravity.
482 From the concept of generic gravity, we move on to that of specific gravity, whereby the measurement is related to a
483 sample. This is a concept that emerged from the French Revolution onwards and is now used for water. However, Marsili
484 uses rainwater instead.

485 **Additional references**

486 Adamson, G.C.D., Private diaries as information sources in climate research, *Wiley Interdiscip. Rev. Clim. Change*, 6,
487 599-611 pp., [10.1002/wcc.365](https://doi.org/10.1002/wcc.365), 2015.

488 Adamson, G. C. D., Hannaford, M. J., & Rohland, E. J., Re-thinking the present: the role of a historical focus in climate
489 change adaptation research. *Global Environmental Change*, 48, 195-205 pp., 2018.

490 Allan, R., Endfield, G., Damodaran, V., Adamson, G., Hannaford, M., Carroll, F., ... & Bliuc, A. M., Toward, integrated
491 historical climate research: the example of Atmospheric Circulation Reconstructions over the Earth. *Wiley*
492 *Interdisciplinary Reviews: Climate Change*, 7(2), 164-174, [10.1002/wcc.408](https://doi.org/10.1002/wcc.408), 2016.

493 Camuffo, D., Sturaro, G., Sixty-cm Submersion of Venice Discovered Thanks to Canaletto's Paintings., *Climatic Change*
494 58, 333–343 pp., <https://doi.org/10.1023/A:1023902120717>, 2003.

495 Camuffo, D., Sturaro, G., Use of proxy-documentary and instrumental data to assess the risk factors leading to sea
496 flooding in Venice, *Global and Planetary Change*, Volume 40, Issues 1–2, 93-103 pp., ISSN 0921-8181,
497 [https://doi.org/10.1016/S0921-8181\(03\)00100-0](https://doi.org/10.1016/S0921-8181(03)00100-0), 2004.

498 Camuffo, D., The Treatise on Waters by Cornaro (1560) and a quantitative assessment of the historical sea surges “Acqua
499 Alta” in Venice. *Climatic Change* 176, 18, <https://doi.org/10.1007/s10584-023-03492-6>, 2023.

500 Della Valle, A., Camuffo, D., Becherini, F. et al. Recovering, correcting, and reconstructing precipitation data affected
501 by gaps and irregular readings: The Padua series from 1812 to 1864. *Climatic Change* 176, 9,
502 <https://doi.org/10.1007/s10584-023-03485-5>, 2023.

503 Folland, C. K., & Parker, D. E., Correction of instrumental biases in historical sea surface temperature data. *Quarterly*
504 *Journal of the Royal Meteorological Society*, 121(522), 319-367 pp., 1995.

505 MacEachren, A. M., The evolution of thematic cartography/a research methodology and historical review. *The Canadian*
506 *Cartographer*, 16(1), 17-33 pp., 1979.

507 Pfister, C., Brázdil, R., Glaser, R., Barriendos, M., Camuffo, D., Deutsch, M., & Rodrigo, F. S., Documentary evidence
508 on climate in sixteenth-century Europe. *Climatic change*, 43, 55-110 pp., 1999.

509 Wilkinson, C., Woodruff, S. D., Brohan, P., Claesson, S., Freeman, E., Koek, F., & Wheeler, D., Recovery of logbooks
510 and international marine data: the RECLAIM project. *International Journal of Climatology*, 31(7), 968-979 pp., 2011.

511