

Global Thermocline Vertical Velocities: a Novel Observation Based Estimate

Diego Cortés-Morales^{1,2}, Alban Lazar², Diana Ruiz Pino², and Juliette Mignot²

¹Mediterranean Institute of Advances Studies (IMEDEA), Mallorca, Spain

²Laboratoire d'Océanographie et du Climat: Expérimentations et Approches Numériques (LOCEAN-IPSL), Paris, France

Correspondence: Diego Cortés-Morales (dcortes@imedea.csic-uib.es)

Abstract.

Vertical velocities at large scales are crucial for understanding ocean dynamics, influencing large-scale circulation and associated biochemical processes, yet their rationale is poorly understood, and their three-dimensional ~~distribution is almost unknown~~ mean distribution and temporal variability is mainly known by models. This paper introduces OLIV3 (Observation-based LInear Vorticity Vertical Velocities), a novel observation-based estimation product of vertical velocities over the global thermocline. This product relies on the geostrophic linear vorticity balance (LVB) applied to ARMOR3D observation-based meridional velocities with ERA5 Ekman pumping vertical velocity as surface boundary condition. It covers the ~~entire water column~~ water column over 71 isopycnal levels, with $1/4^\circ$ horizontal resolution at annual frequency during the 1993 – 2019 period, ~~available on both depth and isopycnal levels~~. Since the geostrophic LVB-derived vertical velocities only capture the geostrophic component of the vertical velocity, their performance is tested using an OGCM numerical model data against the total vertical flow. In the ~~upper ocean~~ thermocline, the LVB accurately reproduces the ~~annual~~ interannual variability and captures the climatology of the large-scale total vertical flow (~~for horizontal~~ scales larger than 5° horizontal resolution) with errors below 50% across the major ocean gyres. Focusing on surface-thermocline exchanges, one of the most common applications that needs vertical velocities, OGCM results indicates that baroclinic geostrophic vertical velocities are largely more accurate than the classic Ekman pumping proxy at estimating the interannual variability of the total vertical flow in the ocean interior. OLIV3 capability to estimate ~~vertical velocities in different ocean circulation regimes~~ real ocean vertical velocities is assessed against three reference datasets: two ~~reanalysis-based and one~~ reanalyses and the observation-based product, OMEGA3D. A strong spatial and temporal correlation is evidenced between OLIV3 and reanalysis datasets, in contrast to the ~~observation-based product~~ OMEGA3D, demonstrating even higher correlation than within themselves and ~~proving that while the geostrophic components of the reanalyses are highly correlated, the ageostrophic part is not~~ supporting the dominance of the geostrophic component of interannual variability of vertical movements. OLIV3 also reconstructs a baroclinic vertical velocity field, consistent with the basin oceanographic concept of Sverdrup balance theory. ~~Regarding one of the most common applications of vertical velocities, the transfers between the surface and thermocline, results from an OGCM simulation show that the baroclinic geostrophic vertical velocity is a better estimator of the temporal variability of the vertical flow in the ocean interior than Ekman pumping, and it is essential to consider the variability of the horizontal~~ By building on theoretical advances made since the introduction of Sverdrup and Ekman transport theories, OLIV3 provides a simplified

yet physically consistent estimate of large-scale vertical transport. The OLIV3 dataset developed in this study is available on 50 vertical levels (~~;~~ Cortés-Morales and Lazar, 2025a) and 71 isopycnal levels (at <https://doi.org/10.5281/zenodo.16962780> ; Cortés-Morales and Lazar, 2025b) (Cortés-Morales and Lazar, 2025).

30 1 Introduction

Ocean vertical motion is fundamental to understanding ocean dynamics. These motions serve as critical mechanisms for the exchange of properties between the ocean surface and interior, as well as within the ocean interior. The vertical exchanges encompass essential components such as heat, salinity, CO₂, oxygen, nutrients (silicates, nitrates), and contribute to shape the large-scale thermocline circulation and the Earth's climate regulation (Leach, 1987; Fischer et al., 1989; Klein and Lapeyre, 2009; Mahadevan, 2016; DeVries et al., 2017; Jacox et al., 2018). ~~At finer scales, mesoscale and submesoscale subduction processes contribute to the ventilation of the thermocline, altering the water mass properties in the ocean's interior, and impacting biogeochemical cycles (Freilich and Mahadevan, 2021).~~ Upwelling motions are ~~also~~ key for sustaining high primary production by supplying nutrients to the euphotic zone, thereby regulating the biological carbon pump (Freilich and Mahadevan, 2019; Uchida et al., 2019; Yang et al., 2021). They also constitute one of the key physical processes of the most productive fisheries regions globally (Pauly and Christensen, 1995). Moreover, the extreme oligotrophy observed in the anticyclonic gyres is attributed to persistent downwelling vertical velocities (Falkowski et al., 1991).

Although the importance of large-scale vertical velocities (w) in ocean dynamics has long been recognised, direct measurements of these motions remain a formidable challenge. This difficulty stems from the extremely weak intensity of the vertical velocity field relative to large-scale horizontal flows. Near the surface, typical magnitudes are on the order of $10^{-5} \text{ m s day}^{-1}$, decreasing to $10^{-6} \text{ m s } 0.1 \text{ m day}^{-1}$ within the thermocline and reaching $10^{-7} 0.01 - 10^{-8} \text{ m s } 0.001 \text{ m day}^{-1}$ at deeper levels in the ocean interior (e.g. Stommel and Arons, 1959; Schott and Stommel, 1978). As a result, the basin-scale ocean's vertical flow remains one of the open questions of physical oceanography. Vertical velocities span nearly four orders of magnitude across spatial and temporal scales. Observations from Lagrangian neutrally buoyant floats, ADCPs (Acoustic Doppler Current Profilers), and the Sentinel V ADCP have captured fine-scale vertical velocities with amplitudes ranging from 10^{-2} to 10^{-4} m s^{-1} , or about 1000 to 10 m day⁻¹ (e.g. Bower et al., 1989; Song et al., 1995; Rossby, 2016; D'Asaro et al., 2018; Comby et al., 2022). Such observations are limited to small regions (covering often only a few degrees or less) and energetic features that do not reflect the magnitude of large-scale circulation. Therefore, a combination of observational data and mathematical tools is required to estimate the vertical flow.

Traditional approaches for estimating vertical velocities across different ocean regions were derived from tracer fluxes (e.g. Stommel and Arons, 1959; Robinson and Stommel, 1959; Wyrтки et al., 1961; Munk, 1966; Wunsch, 1984) or the application of the continuity equation to ~~current measurements (e.g. Stommel and Arons, 1959; Robinson and Stommel, 1959 ; Wyrтки et al., 1961; Munk, 1966;~~ horizontal current measurements obtained from hydrographic station data (e.g. Stommel and Schott, 1977; Schott and Stommel, 1978; Wyrтки, 1981; Roemmich, 1983; ~~Wunsch, 1984;~~) and mooring measurements (e.g. Halpern and Freitag, 1987; Halpern et al., 1989; Weingartner and Weisberg, 1991; Helber and Weisberg, 2001). These

60 early methods provided insight into the small vertical velocities' order of magnitude and upwelling/downwelling patterns of the vertical motions. Vertical velocities have also been inferred from the divergence of horizontal velocity in numerical models (e.g. Madec et al., 2019), ~~although it~~. This methodology remains impractical for global observation-based applications because of the sparse distribution of direct current measurements. ~~An exception~~ Exceptions to such application to observations ~~is~~ are Freeland (2013), which used in situ Argo float observations to estimate w at a single depth in a limited domain of several degrees, assuming zero vertical flow at the surface, and Colin De Verdière and Ollitrault (2016) and Colin De Verdière et al. (2019), which computed vertical velocities from the horizontal divergence of geostrophic velocities inferred from Argo-derived three-dimensional thermohaline fields and float displacements at their parking depth. In the last decade, alternative approaches used isopycnal displacements (Giglio et al., 2013; Christensen et al., 2023), ~~the use of~~ mooring data combined with the momentum and density balances (Sevellec et al., 2015), and biogeochemical tracers (Garcia-Jove et al., 70 2022). The theoretical frameworks have also expanded to include methods based on the Bernoulli function to infer w (Tailleux et al., 2023).

A widely used approach for diagnosing vertical velocities is the quasi-geostrophic (QG) omega equation developed initially for the atmosphere by Hoskins et al. (1978). It links the vertical flow to various processes, including the thermal wind imbalance trend, deformation, kinematic deformation, turbulent buoyancy, and turbulent momentum. This diagnostic equation 75 (solvable from a single snapshot) has been used extensively in regional studies (e.g. Tintore et al., 1991; Pollard and Regier, 1992; Rudnick, 1996; Allen et al., 2001; Buongiorno Nardelli et al., 2001; Gomis et al., 2001; Naveira Garabato et al., 2001; Rodriguez et al., 2001). However, solving the omega equation requires high-resolution 3D fields and well-defined lateral and vertical boundary conditions. The equation was then updated to account for additional physical processes while maintaining the original Hoskins framework (~~e.g.~~ Giordani et al., 2006). It was used to assess mesoscale structures in the Atlantic (Ruiz et al., 2014), Southern (Buongiorno Nardelli et al., 2018) and global Oceans-oceans (Buongiorno Nardelli, 2020) ~~and front and submesoscale dynamics (Ruiz et al., 2019; Freilich and Mahadevan, 2021; Tzortzis et al., 2021).~~ OMEGA3D (Buongiorno Nardelli et al., 2018; Buongiorno Nardelli, 2020) is the only existing global vertical velocity product based on the omega equation, integrating in situ and satellite-derived fields. ~~While it~~ Although OMEGA3D provides a valuable benchmark, the absence of observation-based ground truth for w underscores the need for alternative approaches and new products to estimate vertical 85 velocities ~~at~~ on global circulation scales.

In contrast to the complexity of the omega equation, the linear vorticity balance (LVB; $\beta v = f \partial w / \partial z$) offers a simple diagnostic tool for estimating the geostrophic vertical flow on the β -plane. When vertically integrated from the surface to a level of no motion, it yields the classic Sverdrup balance, a foundational concept in wind-driven circulation theory (Sverdrup et al., 1947). The recent study of Cortés-Morales and Lazar (2024) (CM24 in the rest of the paper) has shown in a reference 90 OGCM simulation in the North Atlantic the capability of the gesotrophic-geostrophic LVB-derived vertical velocities to capture accurately the large-scale interannual variability of the vertical velocities, as well as a significant percentage of their time-mean structure, particularly within the thermocline, but also in the intermediate and deep oceans. The LVB's hypothesis breaks down in certain regions. In particular, the approximation is no longer valid near the equator, within the mixed layer, and along western boundary currents where nonlinear processes are no longer negligible in the vorticity balance.

95 ~~Despite these inherent limitations, this~~ This work extends the approach developed for the North Atlantic ~~by Cortés-Morales and Lazar (20~~
in CM24 to reconstruct observation-based geostrophic vertical velocities globally. It thereby delivers the Observation-based
 Lnear Vorticity Vertical Velocities (OLIV3). The work is structured around the following key objectives: (i) Implementation
 of the LVB framework using geostrophic meridional velocities from ARMOR3D and Ekman pumping from ERA5 as the sur-
 face boundary condition (Section 3.1). (ii) Validation and assessment of limitations of the OLIV3 product through an OGCM
 100 simulation, treated as a perfect model reference (Section 3.2). (iii) Evaluation of the robustness of OLIV3-derived thermocline
 vertical velocities in reproducing the known large-scale characteristics of the global vertical circulation with existing observa-
 tion and model-based estimates (Section 3.3). (iv) Possible physical reasons are proposed to explain why this simplification
 may or may not be valid in certain regions of the ocean (Section 3.4). (v) Comparison with Ekman pumping estimates to
 identify regions where LVB simplifications can be valid and necessary for describing the ocean interior (Section 3.5).

105 2 Methodology and Data

2.1 Revisiting Ekman Pumping Theory for and Reconstructing Geostrophic Vertical Velocities

~~The LVB provides a suitable and well-known foundation for describing the ocean interior flow, as thoroughly described in~~
~~Cortés-Morales and Lazar (2024). In geostrophic form, it is expressed as:~~ Assuming incompressibility of the total three-dimensional
flow implies:

$$110 \quad \nabla \cdot \mathbf{u} = 0, \quad (1)$$

where the velocity vector is written as $\mathbf{u} = (u, v, w) = (\mathbf{u}_h, w)$ where \mathbf{u}_h denotes the horizontal velocity. The total velocity
can be written as the sum of the geostrophic (\mathbf{u}_g) and Ekman components (\mathbf{u}_{Ek}):

$$\nabla \cdot \mathbf{u}_g + \nabla \cdot \mathbf{u}_{Ek} = 0, \quad (2)$$

Because the Ekman circulation vanishes below the base of the Ekman layer, the vertical integration of Eq. 2 over the water
 115 column can be separated into the Ekman layer, extending from the surface ($z = 0$) to its base ($z = D_{Ek}$), and into the ocean
interior, extending from the Ekman layer base to a given depth z where we neglect ageostrophic contributions:

$$\underbrace{\int_{D_{Ek}}^0 \nabla \cdot \mathbf{u}_g dz' + \int_{D_{Ek}}^0 \nabla \cdot \mathbf{u}_{Ek} dz'}_{\text{Ekman Layer}} + \underbrace{\int_z^{D_{Ek}} \nabla \cdot \mathbf{u}_g dz'}_{\text{Ocean Interior}} = 0, \quad (3)$$

The horizontal geostrophic current is calculated from the pressure gradient as

$$\underbrace{u_g = -\frac{1}{\rho_0 f} \frac{\partial p}{\partial y}; v_g = \frac{1}{\rho_0 f} \frac{\partial p}{\partial x}}_{\text{Geostrophic Current}} \quad (4)$$

120 Therefore, in a β -plane, the divergence of the horizontal geostrophic flow can be expressed as

$$\underline{\beta v_g \nabla_h \mathbf{u}_{h,g}} = \underline{f \partial_z w_g} - \underline{\frac{\beta v_g}{f}}. \quad (5)$$

125 assuming $f \neq 0$ (Pedlosky, 1996), where f is the Coriolis parameter and β is the meridional gradient of f , and v_g and w_g are the geostrophic meridional and vertical components of the velocity, respectively. Assuming a geostrophic flow on a β -plane, this formulation describes a local mass balance between meridional divergent flow and an opposing vertical convergence. Therefore, this formulation allows the estimation of the vertical velocity field in the ocean interior (Charney, 1955; Cortés-Morales and Lazar, 2024) without requiring the classical assumption of a linear variation of f across the domain.

The contribution of the Ekman layer in Eq. 3 can be expanded as

$$\underline{\int_{D_{ek}}^0 -\frac{\beta v_g}{f} dz'} + \underline{\int_{D_{ek}}^0 \frac{\partial w_g}{\partial z} dz'} + \underline{\int_{D_{ek}}^0 \nabla \cdot \mathbf{u}_{h,Ek} dz'} + \underline{\int_{D_{ek}}^0 \frac{\partial w_{Ek}}{\partial z} dz'}. \quad (6)$$

Evaluating analytically the vertical integrals of the vertical velocity gradients gives

$$130 \underline{\int_{D_{ek}}^0 -\frac{\beta v_g}{f} dz' + w_g(z=0) - w_g(z=D_{Ek})} + \underline{\int_{D_{ek}}^0 \nabla \cdot \mathbf{u}_{h,Ek} dz' + w_{Ek}(z=0) - w_{Ek}(z=D_{Ek})}. \quad (7)$$

We assume that $w_{Ek}(z=D_{Ek})=0$, as all components of the three-dimensional Ekman flow are dissipated at D_{Ek} . Furthermore, the Ekman theory relates the Ekman transport,

$$\underline{\mathbf{U}_{h,Ek} = \int_{D_{Ek}}^0 \mathbf{u}_{h,Ek} dz'}. \quad (8)$$

to the wind stress through

$$135 \underline{\nabla \cdot \mathbf{U}_{h,Ek} = \nabla \times \left(\frac{\boldsymbol{\tau}}{\rho_0 f} \right)}, \quad (9)$$

where $\mathbf{U}_{h,Ek}$ denotes the horizontal Ekman transport in the Ekman layer, ρ_0 represents the water density at the surface, and $\boldsymbol{\tau}$ is the wind stress.

At the sea surface, we impose zero total vertical flow (w_{tot}),

$$\underline{w_{tot}(z=0) = 0}. \quad (10)$$

140 This condition is exact for the stationary flow of the time-mean state, but it is only an approximation for time-varying flows. In the latter case, this approximation neglects the time derivative of sea surface height, $d(\eta)/dt$, where η represents the sea surface height relative to the mean state reference surface level. The total vertical velocity is decomposed into geostrophic and ageostrophic components ($w_{tot} = w_g + w_{ag}$), where the ageostrophic contribution is assumed, at first order, to be produced by surface wind stress and friction (Ekman three-dimensional flow). Under these assumptions, the surface condition implies:

$$145 \quad w_g(z=0) = -w_{gEk}(z_{ref}=0) - \int_z^{z_{ref}} \frac{\beta v_g}{f} dz'. \quad (11)$$

The geostrophic form of the LVB and the corresponding vertical velocity w_g can be derived from the divergence of the geostrophic flow on a β -plane (Pedlosky, 1996). Thus, this w can be considered as the geostrophic component of the total vertical flow. One common approach to setting the reference condition assumes a level of no motion, where all velocity components are zero. While some studies support this assumption (e.g. Koelling et al., 2020), a degree of uncertainty remains under real oceanic conditions. Alternatively, satellite observations, with their extensive spatial and temporal coverage, provide an adequate boundary condition. Therefore, it is proposed to use Ekman pumping at the ocean surface (w_{Ek}) as boundary condition at the surface. Substituting the Ekman transport (Eq. 9) into Eq. 7 and imposing the surface condition of zero total vertical velocity, the contribution of the Ekman layer in Eq. 3 yields

$$155 \quad w_{Ek} = \frac{1}{\rho_0} \nabla \times \left(\frac{\tau^{surf}}{f} \right)$$

$$\int_{D_{ek}}^0 -\frac{\beta v_g}{f} dz' - w_g(z = D_{Ek}) + \nabla \times \left(\frac{\tau}{\rho_0 f} \right). \quad (12)$$

Where ρ represents the water density at surface (1025 kg m^{-3}), and τ^{surf} the wind stress at the oceanic surface. In the ocean interior, only the geostrophic currents is not negligible, such that the contribution of ocean interior to the Eq. 3 can be derived

$$\int_z^{D_{ek}} -\frac{\beta v_g}{f} dz' + \int_z^{D_{ek}} \frac{\partial w_g}{\partial z} dz'. \quad (13)$$

160 Evaluating the vertical integral of the contribution of ocean interior to the Eq. 3 leads to

$$\int_z^{D_{ek}} -\frac{\beta v_g}{f} dz' + w_g(z = D_{Ek}) - w_g(z). \quad (14)$$

Combining the Ekman layer (Eq. 12) and ocean interior (Eq. 14) contributions, the terms corresponding to $w_g(D_{Ek})$ cancel each other, and the expression for the geostrophic vertical velocity in the ocean interior is obtained

$$w_g(z) = \nabla \times \left(\frac{\tau}{\rho_0 f} \right) - \int_z^0 \frac{\beta v_g}{f} dz'. \quad (15)$$

165 In this scenario, the horizontal geostrophic flow is divergent and generates a net geostrophic vertical velocity in the ocean interior. Equation 15 corresponds to the indefinite vertical integral of the LVB applied to the geostrophic flow. While the LVB is an approximation of the vorticity equation valid for large-scale flows, it provides a suitable and well-known foundation for describing the ocean interior flow, as thoroughly described in CM24. As a result of the approximations considered to derive the continuity equation (Eq. 2), discrepancies between the geostrophic vertical velocity and the total vertical velocity are expected,
170 particularly in regions where ageostrophic and nonlinear processes are non-negligible in the ocean interior.

Equation 15 further shows that, although w_g can only be explicitly diagnosed below the Ekman layer, due to the fact that we have defined z below D_{Ek} in Eq. 3, the magnitude of the vertical flow at the base of the Ekman layer differs from its surface value due to the vertically integrated divergence of the horizontal geostrophic flow within the Ekman layer. At the sea surface or if the geostrophic flow divergence is negligible, the geostrophic vertical velocity is equal to the Ekman pumping. However,
175 CM24 reported a non-negligible contribution of the β -plane geostrophic divergence for an OGCM with the Ekman layer. This effort is consistent with that of Jacox et al. (2018), which albeit different, similarly emphasises the need to revisit classical assumption of Ekman theory. Consequently, in this study, Ekman pumping (w_{Ek}) is adopted as surface boundary condition for the computation of geostrophic vertical velocities.

2.2 Deriving Observation-Based Geostrophic Vertical Velocities (OLIV3)

180 ~~While Cortés-Morales and Lazar (2024) aimed to illustrate the potential of the LVB framework as a main estimator of the North Atlantic thermocline dynamics, the primary objective of this study is to introduce a new observation-based product of geostrophic velocities derived from the geostrophic LVB.~~ To estimate the global ocean vertical velocities from the divergence of the geostrophic flow, meridional geostrophic velocities from ARMOR3D are used in combination with Ekman pumping vertical velocities derived from ERA-ERA5 wind stress as boundary condition at the ocean surface. A summary of the main
185 characteristics of the input datasets, including physical variables, cover period, temporal frequency, horizontal and vertical resolution is provided in Table 1.

ARMOR3D dataset (Guinehut et al., 2012; Mulet et al., 2012) integrates satellite-derived and in situ observations. Its derivation first involves the construction of synthetic temperature and salinity fields from altimetric surface level anomaly (SLA; AVISO+, 2015), and sea surface temperature and salinity (SST and SSS) (Reynolds et al., 2007; Droghei et al., 2018) via linear regression method and covariances calculated from historical in situ observations (EN3 dataset (Ingleby and Huddleston, 2007) and Argo floats(?)). The synthetic and observed temperature and salinity profiles are then merged using optimal interpolation (Bretherton et al., 1976). Finally, geostrophic velocities are computed using the thermal wind equation, referenced to

the surface geostrophic velocities estimated from the altimetric absolute dynamic topography. The mixed layer depth (MLD) is obtained from the minimum of temperature and density threshold equivalent to a 0.2°C decrease. ARMOR3D is available at <https://doi.org/10.48670/moi-00052>.

Ekman pumping vertical velocities are computed from monthly wind stress data provided by ERA5 (Hersbach et al., 2023), the fifth-generation reanalysis of the European Centre for Medium-Range Weather Forecasts (ECMWF). The ERA5 fields are provided at 0.25° horizontal resolution and can be downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store ([DOI: 10.24381/cds.f17050d7](https://doi.org/10.24381/cds.f17050d7)).

The resulting Observation-based Linear Vorticity Vertical Velocities (OLIV3) product consists of geostrophic vertical velocities derived from ARMOR3D meridional velocities and surface Ekman pumping from ERA5 wind stress, using [the integrated geostrophic LVB \(Eq. ??\) Eq. 15 over the native ARMOR3D vertical levels and then interpolated over isopycnal levels defined by the neutral density \(Jackett et al., 2006\) of ARMOR3D thermohaline field](#). The product spans the 1993 - 2019 period, with a horizontal resolution of 0.25° and [available in two vertical grids: 50 vertical levels and 71 isopycnal levels](#). [Both versions are](#) Figure 1 represents the OLIV3 time-mean (1993-2019) vertical velocity at various sigma levels across the tachocline, understood as the upper ocean layer defined by a strong vertical shear in the velocity field (CM24). The product is quality-flagged based on the [time-mean](#) relative error and interannual correlation coefficient between w_g and w_{tot} in the OGCM perfect model test. [The datasets are available on 50 vertical levels \(Cortés-Morales and Lazar, 2025a\) and 71 isopycnal levels \(and available at](#) <https://doi.org/10.5281/zenodo.16962780> ; [Cortés-Morales and Lazar, 2025b](#) (Cortés-Morales and Lazar, 2025). A low-resolution version, used in the intercomparison test of this study (Section 3.3), is available upon request.

2.3 Existing Estimates of Vertical Velocities

To evaluate OLIV3 performance, we compare the vertical velocities with two reanalyses (GLORYS12v1, ECCOV4r4), and an observation-based product (OMEGA3D). To facilitate comparison, the main attributes of the validation products sources are summarised in Table 1.

Following [Cortés-Morales and Lazar, 2024](#) CM24, the reference OGCM simulation for the assessment of validity of the methodology is the Nucleus for European Modelling of the Ocean (NEMO) OGCM OCCITENS run from the OCCIPUT project (Penduff et al., 2014; Bessieres et al., 2017, Madec et al., 2019). This simulation is forced by DFS5.2 forcing set, using ERA-Interim and ERA40 reanalyses (Dussin et al., 2016). Neutral density field and the isopycnal surfaces were computed for this study using thermohaline and sea surface height fields applied to Jackett et al. (2006) formulation. The MLD provided by the NEMO OCCITENS simulation is computed using a density criterion of 0.01 kg m^{-3} of density change from the surface following the procedure defined by de Boyer et al. (2004). Outputs from the NEMO OGCM OCCITENS simulation are available upon request (thierry.penduff@cns.fr). Geostrophic velocities are derived from the model pressure field (calculated using sea surface height and the hydrostatic equation) using the geostrophic equation via the codes available at <https://github.com/meom-group/CDFTOOLS>.

The GLObal Ocean ReanALYSIS and Simulations (GLORYS; Verezemskaya et al., 2021, Lellouche et al., 2021) assimilates via Kalman filter along-track altimeter SLA, satellite SST, sea ice concentration, and in situ temperature and salinity profiles,




Figure 1. OLIV3 time-mean (1993-2019) vertical velocity at various sigma levels across the tachocline. The field has been smoothed with a 5° running mean. Translucent black shading represents regions within the maximum mixed layer over the study period. The black contour lines represent the depth of the isopycnal surface in meters.

using NEMO as the model component (Lellouche et al., 2018). The MLD is defined following the same methodology as in the NEMOS-NEMO OGCM simulation. This dataset is hereafter referred to as GLORYS12v1 and accessed at https://tds.mercator-ocean.fr/thredds/glorys12v1/glorys12v1_pgn_monthlymeans.html.

230 The Estimating the Circulation and Climate of the Ocean (ECCO) in its fourth release, version 4 (Forget et al., 2015; Fukumori et al., 2018) employs a 4D-VAR assimilation scheme, integrating satellite altimetry, in situ temperature and salinity profiles from Argo, satellite sea surface salinity and temperature, and ocean bottom pressure, together with the MIT General

Circulation Model (Adcroft et al., 2004). The mixed layer depth is defined following the procedure developed by Kara et al. (2000, 2003), which [finds](#) [find](#) that the optimal estimated of turbulent mixing penetration is obtained with a mixed layer depth definition of $\Delta T = 0.8^\circ$ C. This reanalysis hereafter referred to as ECCOv4r4 is available at <https://www.ecco-group.org/products-ECCO-V4r4.htm>.

Finally, OMEGA3D is an observation-based global estimate of vertical velocities derived from the QG omega equation (Buongiorno Nardelli et al., 2018; Buongiorno Nardelli, 2020). It is based on ARMOR3D thermohaline field and geostrophic velocities, and ERA-Interim (Dee et al., 2011) surface air-sea fluxes. OMEGA3D is available at https://doi.org/10.25423/cmcc/multiobs_glo_phy_w_rep_015_007.

Table 1. Summary of datasets used for OLIV3 computation and validation.

| Dataset | Type | Resolution (hor./vert.) | Period | Frequency | Variables Used |
|---------------|--------------|-------------------------|--------------|-----------|------------------------------|
| ARMOR3D | Obs-based | 0.25° / 50 levels | 1993–2019 | weekly | v_g, T, S, MLD |
| ERA5 | Reanalysis | 0.25° / surface | 1979–present | monthly | Wind stress |
| NEMO-OCCITENS | Model (OGCM) | 0.25° / 75 levels | 1960–2015 | monthly | $w_t, w_g, T, S, \text{MLD}$ |
| GLORYS12v1 | Reanalysis | 1/12° / 50 levels | 1993–2019 | monthly | w_t, T, S, MLD |
| ECCOv4r4 | Reanalysis | 1° / 50 levels | 1992–2017 | monthly | w_t, T, S, MLD |
| OMEGA3D | Obs-based | 0.25° / 75 levels | 1993–2019 | weekly | w (QG) |
| OLIV3 | Obs-based | 0.25° / 50 levels | 1993–2019 | yearly | w_g |

2.4 Validation Methodology

Validating OLIV3 with observational data is problematic due to the lack of a ground truth for large-scale vertical velocities. Here, the performance of OLIV3 relies on the consistency across existing w estimates: GLORYS12v1, ECCOv4r4 and OMEGA3D. The intercomparison is conducted on a common spatiotemporal resolution: annual means at 5° horizontal resolution and isopycnal levels. This choice reduces vertical grid and thermohaline structure differences. It also allows to focus on large-scale dynamics, which are better resolved by the LVB framework ([Cortés-Morales and Lazar, 2024CM24](#)). The isopycnal levels are defined by the neutral density [of each dataset](#) (Jackett et al., 2006) [of each dataset](#). For the OLIV3 and OMEGA3D datasets, the thermohaline field used to interpolate w onto isopycnal levels is ARMOR3D, since the velocity field was constructed using it (Buongiorno Nardelli, 2020). Diagnostics are computed over the overlapping 23 year period (1993 - 2015) and include [the](#) time-mean [horizontal pattern](#), [the](#) time-mean [vertical gradient](#), [diapycnal gradient](#), [the interannual](#) variance and correlation coefficient (R). Regions within the equator band (5° S/N) are excluded as the geostrophic equation cannot be solved at these latitudes.

3 Results and discussion

3.1 Observation-based Linear Vorticity Vertical Velocities (OLIV3)

255 Twenty ~~five~~ seven year mean geostrophic vertical velocities (w_g) stemming from the OLIV3 product at σ_{26} isopycnal surface are presented in Fig. 2a. This isopycnal level was chosen to assess the vertical velocity estimates across most of the extension of the global subtropical gyres, while maintaining a focus on thermocline dynamics, where the LVB framework performs best (~~Cortés-Morales and Lazar, 2024~~see CM24 for the North Atlantic Ocean). The geostrophic vertical velocity field (OLIV3) within the tachocline, ~~understood as the upper ocean layer defined by a strong vertical shear in the velocity~~
260 ~~field (Cortés-Morales and Lazar, 2024)~~, reproduces the well-known wind-driven circulation features represented by Ekman pumping (Fig. 2b), generally with upwelling at tropical latitudes and downwelling at the subtropics. This emphasises the role of atmospheric forcing wind driven divergence as the primary driver of vertical flow within the upper ocean (e.g. Huang and Russell, 1994; Liang et al., 2017). The Pacific and Atlantic eastern tropical upwelling systems that continue along the eastern coast up to subtropical latitudes are associated with maximum positive values near the coast around $2 \times 10^{-6} \text{ m s}^{-1}$ 0.2 m day⁻¹
265 (Aristegui et al., 2009). The anticyclonic circulation of subtropical systems is characterised by negative w (downwelling) with maximum values found along western boundaries. This pattern differs from the agreement among reanalyses about upwelling in most of the western boundary current systems, in particular the Gulf Stream and the Kuroshio Current ~~as demonstrated by Liao et al. (2022)~~(Liao et al., 2022). In the Gulf Stream case, this poor agreement with reanalyses ~~in the literature~~ aligns with the lack of confidence in the LVB as an estimator of the vertical flow, as ~~demonstrated by Cortés-Morales and Lazar (2024)~~we
270 already demonstrated at the Atlantic Ocean. Outside the Northern Hemisphere subtropical band, some upwelling occurs over the extension of the Gulf Stream and Kuroshio systems, which can be explained by Ekman pumping (Qiu and Huang, 1995).

To analyse the temporal and vertical variability of the vertical velocity estimates, Fig. 3 shows regionally averaged w_g as a function of isopycnal level and year for three distinct regions in the North Atlantic Ocean: the eastern tropical, subtropical and subpolar gyres (purple regions in Fig. 2a). The bottom of the layer was selected as the level where the estimates' climatology changes sign. Figure 3 evidences that the direction of geostrophic vertical velocity estimates maintains their sign throughout most of the layer's thickness. Before reaching this depth, the annual velocities reduce in amplitude, supporting the baroclinic nature of vertical flow ~~in the tachocline found in Cortés-Morales and Lazar (2024)~~found in the North Atlantic tachocline (CM24). This behaviour is consistent with the requirement of a level of no motion at depth to satisfy the Sverdrup balance (Thomas et al., 2014).

280 The velocity sign remains unchanged over time in the three regions (Fig. 3), but temporal variability is evident across them. Weaker upwelling events in North Atlantic Tropical Gyre (e.g. 1995, 2001, 2005 and 2010 in top panel in Fig. 3) seem associated with the negative North Atlantic Oscillation (NAO) phases shown by Pinto et al. (2012) and Roch et al. (2024), while positive NAO phases correspond to stronger upwelling. This suggests that the variability of the eastern tropical gyre is modulated by the phase of the NAO. Interestingly, the magnitude of the oceanic response does not reflect the NAO intensity presented in the references above. Focusing on the subtropical gyre (central panel in Fig. 3), maximum downwelling events do not correlate with the NAO index as clearly as the tropical upwelling does. ~~Studies such as Pinto et al. (2012) suggest~~




Figure 2. (a) OLIV3 time-mean (1993-2019) vertical velocity at σ_{26} (1993-2018)-26. The field has been smoothed with a 5° running mean. Translucent black shading represents regions within the maximum mixed layer over the study period. The black contour lines represent the depth of the isopycnal surface in meters. Purple rectangles indicate the average regions for Fig. 3. (b) Time-mean Ekman pumping from ERA5. The field has been smoothed with a 5° running mean. White hatching masks the regions above σ_{26} for comparison purposes.

that the atmospheric changes modify both the intensity of the downwelling and the location of the subduction maximum (Pinto et al., 2012). Therefore, the out-of-phase relationship between subtropical vertical flow variability and NAO index may suggest that the region selected for this study this region does not fully reproduce the gyre interannual variability. However, other components can influence the variability of the region as the NAO is not the sole contributor to atmospheric forcing variability (Zhao and Johns, 2014). In the subpolar gyre (bottom panel in Fig. 3), oscillations in upwelling amplitude exhibit a periodicity of approximately 5 years, with particularly strong positive velocity events during 2002-2003, 2007 and 2009. These features align with observed changes in the subpolar gyre from satellite altimetry (Foukal et al., 2017) and volume transport estimates of the East Greenland Current (Daniault et al., 2011). These findings suggest that OLIV3, while capturing the baro-

290




Figure 3. Regionally averaged vertical velocity estimates from OLIV3 as a function of isopycnal surface and year for three subregions in the North Atlantic Ocean (NATL): (top) Tropical Gyre (30-16°W, 8-16°N), (middle) Subtropical Gyre (50-30° W, 15-30°N), and (bottom) Subpolar Gyre (55-30°W, 50-60°N). For each region, the first isopycnal level corresponds to the shallowest level that does not intersect with the sea surface. Contours of vertical velocities have been included for readability (m day^{-1}).

295 clinic nature of the vertical velocity field in the tachocline, is also capable of transmitting some of the surface interannual signal into the ocean interior.

3.2 Linear Vorticity Balance Framework in a Perfect Model Test

The vertical flow computed by applying the Eq. [??-15](#) to the ARMOR3D dataset represents the geostrophic component only. Therefore, before comparing OLIV3 with estimates of the total vertical flow, the limitations of this formulation are evaluated using an OGCM simulation, considered as a "perfect model test". This global-scale analysis extends the regional assessment conducted by [Cortés-Morales and Lazar \(2024\) CM24](#) for the North Atlantic Ocean. [The To evaluate the ability of the geostrophic component to represent the spatiotemporal variability of the total vertical flow, we examine three diagnostics: the absolute relative error between the time-mean geostrophic vertical velocity \(\$w_g\$ \) and the total vertical velocity \(\$w_{tot}\$ \), their interannual correlation coefficient at \$\sigma_{26}\$, as well as the relative error in the vertical-time-mean diapycnal gradient between the bottom of the mixed layer \(MLD\) and \$\sigma_{27}\$ computed using OGCM over the period 1993 - 2015 are presented in \(Fig. 4-The vertical\). The time-mean diapycnal gradient of vertical velocities between the base of the maximum mixed layer and \$\sigma_{27}\$ has been normalised by the density interval distance \(in meters\) between these isopycnal levels, as follows:](#)

$$\frac{\partial \Delta_z w}{\partial z} = \frac{\frac{|w_{\sigma_{27}}| - |w_{\sigma_{MLD}}|}{\sigma_{27} - \sigma_{MLD}}}{\frac{|z_{\sigma_{27}} - z_{\sigma_{MLD}}|}}. \quad (16)$$

Negative values indicate a decrease in magnitude with depth, while positive values indicate an increase. The velocity fields were smoothed with a $5^\circ \times 5^\circ$ running mean to retain large-scale structures that LVB can describe ([Cortés-Morales and Lazar, 2024 CM24](#)).

As shown in [Cortés-Morales and Lazar \(2024\) CM24](#) for the North Atlantic, w_g succeeds in estimating w_{tot} over most parts of the basins. Fig. 4a shows that values are accurate, with a relative error below 50%, across most of the global tropical and subtropical gyres (yellow, orange and red regions), as well as the eastern part of the subpolar Pacific gyre and the arctic Beaufort gyre. This result shows that the geostrophic vertical velocity field generally reproduces the spatial structure and amplitude of the thermocline vertical flow within the major gyres in the model simulation, suggesting that a similar behaviour may occur in the real ocean.

Relative errors exceed 50% in several regions. These include the intergyres bands, where vertical velocities change sign, as well as larger regions of intense current systems, such as the Gulf Stream and its northeastward extension, the Brazil-Malvinas Current, the Kuroshio Current, the Eastern Australia Current and the Algalhas Current (see locations in Barceló-Llull et al., 2025). High relative errors are also found in regions dominated by strong zonal flows, like the deep tropics, such as the Equatorial Currents and Countercurrents, but also open ocean poleward extensions of western boundary currents like the North Atlantic Drift (NAD), and the Antarctic Circumpolar Current (ACC), more visible on deeper isopycnals (not shown). They are characterised by an intense zonal component, which geostrophic part, unlike the meridional component, does not generate divergence and geostrophic vertical velocity ([Cortés-Morales and Lazar, 2024 CM24](#)). Most eastern boundaries, particularly eastern boundary upwelling systems (e.g. California or Benguela), as well as the northern Indian basins, suffer from high errors in the estimation of their time-mean vertical velocities. Many of these discrepancies typically arise in regions where the LVB (Eq. [??](#)) no longer holds (hatching in Fig. 4a). In such areas, nonlinear processes, friction and lateral diffusion become

essential to close the time-mean vorticity budget (e.g. Sonnewald et al., 2019; Waldman and Giordani, 2023; Khatri et al.,
330 2023).

It is interesting to note that in certain regions, the large differences are not consistent with the local validity of the LVB at the current isopycnal level. For example, in the North Atlantic intergyre region, a large relative difference between w_g and w_{tot} appears as a narrow band around 15°N, despite the LVB terms showing relative agreement within 10% (i.e., outside hatching). In contrast, within the tropical gyre (centred around 10°N), the relative error between w_g and w_{tot} is found to fall below
335 50%, yet the LVB is not valid on σ_{26} there (hatching areas). Indeed, as ~~discussed in Cortés-Morales and Lazar (2024), the~~ [for the North Atlantic application of LVB \(CM24\)](#) the geostrophic vertical velocity at a given depth is computed via vertical integration of the meridional transport above that level. Therefore, local deviations from the LVB at a given depth, implying that the export of water is not fully conserved, do not necessarily lead to large errors in the integrated geostrophic velocity. This is more pronounced at deeper levels where the amplitude of the meridional transport is reduced due to the baroclinic structure
340 of the tachocline flow.

Considering now the reconstruction of the interannual variability, the LVB method appears to be strikingly accurate (Fig. 4b). Correlation between w_g and w_{tot} exceeds 0.9, ~~and a variance explained at~~ [indicating that the geostrophic component explains](#) more than 80% [of the total vertical velocity variance](#), across most of the tropical, subtropical global ocean and northern subpolar gyres (only Pacific subpolar gyre visible), and the Beaufort polar gyre. These good results extend those previously reported
345 for the North Atlantic ~~in Cortés-Morales and Lazar (2024)~~. Weaker, or even negative, correlation values are found within the major western boundary current systems of the Pacific, Atlantic and Indian Oceans, where nonlinear dynamics are stronger. Similarly large differences also characterise Atlantic and Pacific deep tropics, the open ocean poleward extensions of western boundary currents, and the entire ACC, visible southeast of Africa. These regions are all dominated by flows with strong zonal components relative to the meridional flow, likely generating mostly ageostrophic vertical velocities that cannot be captured by
350 the LVB framework. Remarkably, very high correlations persist even in regions where the time-mean geostrophic component fails to replicate the total vertical velocity pathways (Fig. 4a), as well as in regions where the LVB does not hold, such as the mixed layer (translucent black surfaces in Fig. 4a).

Figure 4c illustrates the ability of time-mean w_g to represent the vertical structure of the velocity field shown in Fig. 4a, by displaying the relative error of a proxy of the ~~vertical~~ [time-mean diapycnal](#) gradient (Eq. 16) [between \$w_g\$ and \$w_{tot}\$](#) . Note
355 that the ~~vertical~~ [diapycnal](#) gradient of the [time-mean](#) total vertical velocity is almost everywhere positive (non-dotted areas), ~~corresponding to the downward decrease~~ [indicating a decrease in the magnitude of the vertical velocity toward the base of](#) ~~current intensity towards the bottom of the thermocline, suggesting that the velocity field is baroclinic and~~. [This structure is](#) [consistent with a baroclinic velocity field](#), generating a tachocline, ~~as~~ [underlined in the North Atlantic by Cortés-Morales and Lazar \(2024\)](#) ~~-This vertical~~ [\(CM24\)](#). [This diapycnal](#) gradient appears to be well represented by w_g , where the five main subtropical gyres
360 are characterised by relative error smaller than 20%. ~~A downward increase~~ [The magnitude of \$w_{tot}\$ intensity grows with depth](#) (dotted regions) ~~is found in~~ [within](#) western boundary current systems such as the Gulf Stream, Kuroshio and Brazil-Malvinas Currents, the North Pacific and North Atlantic deep tropics, and the intergyre regions between the major tropical and subtropical gyres. Nonetheless, in the Pacific and Atlantic tropical gyres, the errors in the ~~vertical~~ [diapycnal](#) gradient of the total




Figure 4. Assessment of OGCM geostrophic vertical velocity estimate against the OGCM total vertical velocity. (a) Absolute relative error between means of geostrophic vertical velocity and total vertical velocity at σ_{26} isopycnal surface. Translucent black shading represents regions within the maximum mixed layer over the study period and hatching delimits the areas where the LVB does not hold (relative error > 10%). (b) Correlation coefficient between the annual geostrophic vertical velocity and total vertical velocity at σ_{26} isopycnal surface. Black contours indicate correlation coefficients of 0.7 (dashed) and 0.9 (solid). (c) Relative difference in vertical diapycnal gradient of total and geostrophic time-mean vertical velocity ($\Delta_z w_{tot}$ and $\Delta_z w_g$ respectively) between the mixed layer base and σ_{27} , computed as $(w_{\sigma_{27}} - w_{\sigma_{MLD}}) / (\sigma_{27} - \sigma_{MLD}) \Delta_z w = (|w_{\sigma_{27}}| - |w_{\sigma_{MLD}}|) / (|z_{\sigma_{27}} - z_{\sigma_{MLD}}|)$ normalised by the magnitude of the diapycnal gradient of the total vertical velocity. Only the regions with a MLD shallower than σ_{27} are considered for this analysis. Dotted (no dotted) areas indicate regions where the total vertical velocity display a positive (negative) vertical diapycnal gradient, meaning increasing (decreasing) magnitude with depth.

and geostrophic vertical flow are larger than in the subtropical gyres. The spatial distribution of these errors is similar to the
365 pattern of relative errors above 10% between w_g and w_{tot} (Fig. 4a). Therefore, these results suggest the limitations of the LVB
framework in reproducing the time-mean value at a given depth and the vertical structure of the total vertical flow at tropical
regions and western boundary current systems.

These findings evidence the relevance of the geostrophic LVB framework for capturing and explaining the dynamics of the
large-scale vertical motion in an OGCM simulation. Most notably, the high and widespread ~~interannual synchrony~~ synchrony
370 at annual frequencies between w_{tot} and w_g suggests that the geostrophic component strongly dominates the interannual variability,
and therefore likely in the real ocean as well. While nonlinear processes influence the mean amplitude of the vertical
flow, they play a smaller role in modulating this variability. Thus, extending previous results from the North Atlantic Ocean to
global scales, it shows that geostrophic vertical velocity provides a reliable estimate of the total vertical flow for studying the
climatological flow structure within the interior of the major gyres and the interannual variability of vertical motion throughout
375 much of the tropical and subtropical oceans, parts of northern subpolar gyres and polar gyres. This supports the relevance of
applying the same reconstruction methodology to observation-based data. In the rest of the paper, we assess ~~our~~ the named
OLIV3, observation-based estimate of the global thermocline vertical velocities, ~~named OLIV3~~.

3.3 Assessment of OLIV3 Relative to Existing Vertical Velocity Estimates Over the Global Thermocline

Due to the lack of direct observations for large-scale vertical velocities, it is necessary to evaluate the performance of OLIV3
380 relative to commonly used products. In particular, acknowledging the multidimensional nature of the vertical velocity field,
a comprehensive evaluation must address the mean three-dimensional structure, as well as the temporal signals relevant for
future studies.

3.3.1 Large-scale ~~climatological vertical flow features~~ Climatological Vertical Flow Features

The ability of OLIV3 to represent the large-scale climatological upwelling ~~and~~ and downwelling structures is evaluated by com-
385 paring the 23-year mean at σ_{26} against three reference datasets: GLORYS12v1 and ECCOv4r4 reanalyses, and OMEGA3D
observation-based product. The large-scale key wind-driven features captured by OLIV3 in Fig. 5 are consistent at first order
with those in the reference datasets, with vertical velocity amplitude falling within a common range of values from ~~0.1 to 10 x~~
~~10⁻⁶ m s⁻¹ to 1 m day⁻¹~~ 0.01 to 1 m day⁻¹.

In the tropics, OLIV3, ECCOv4r4 and GLORYS12v1 (Fig. 5a, c and d respectively) exhibit spatially variable upwelling
390 across all the oceanic basins, including along the Pacific and Atlantic eastern boundaries (Elmoussaoui et al., 2005, Faye et
al., 2015). OMEGA3D (Fig. 5b) captures a broader and smoother field with maximum upwelling at the centre of the basins.
These OMEGA3D upwelling patterns were already reported in Buongiorno Nardelli (2020). Notably, the deep tropical band
(5-10°N/S) show low agreement between reanalyses and OLIV3, matching the regions where the LVB errors in the OGCM
exceed 10% (black hatching in Fig. 5a). Some regions exhibit amplitude discrepancies among the three datasets (e.g. North
395 Pacific tropical band), and some others even display opposite sign (e.g. Indian Ocean).

In subtropical latitudes, ~~despite~~ OLIV3 and OMEGA3D ~~tend to overestimate downwelling relative to the reanalyses. Despite this overestimation, OLIV3 estimates align more closely with reanalyses than OMEGA3D. Some reproduce the large-scale direction of the vertical flow, some~~ regional differences emerge ~~for~~ (Fig. 5e-i). For example, in the North Pacific, ~~where~~ OLIV3 and GLORYS12v1 show a maximum downwelling centred near 30°N, 140°W, while OMEGA3D and ECCOv4r4 display strong downwelling across the entire 30°N band. In the North Atlantic, the downwelling maximum in OLIV3 and GLORYS12v1 is found in the southeastern part of the gyre. Moreover, OMEGA3D captures this maximum closer to the western boundary current extension and ECCOv4r4 centres it at 30°N. In the South Atlantic and South Indian Oceans, all datasets reproduce maximum downwelling near the intersection of σ_{26} with the bottom of the mixed layer. In the South Pacific, only OMEGA3D presents a maximum near the intersection between the isopycnal level and the bottom of the mixed layer, GLORYS12v1 produces weaker amplitudes, and OLIV3 and ECCOv4r4 locate a maximum downwelling at around 30°S, 90°W.

Western boundary currents also reveal discrepancies. In the Gulf Stream and Kuroshio current extensions, OLIV3 reproduces upwelling patterns consistent with Ekman pumping (Qiu and Huang, 1995), OMEGA3D and reanalyses, ~~while capturing and captures~~ strong downwelling at 30°N along the continental section of both currents. At this latitude, in the Gulf Stream case, OMEGA3D reveals some downwelling, while ECCOv4r4 displays positive vertical velocities, in agreement with other model results (GODAS and SODA, among others) shown by Liao et al. (2022). GLORYS12v1 reproduces upwelling and downwelling on both sides of the current. In the Kuroshio Current, ECCOv4r4, GLORYS12v1 and OMEGA3D feature mainly upward flow. The Brazil Current is associated with upwelling flow in all datasets, although OLIV3 tends to overestimate its amplitude.

One may argue that a likely source of discrepancies across the isopycnal level between OLIV3 and the reanalyses is that OLIV3 reconstructs only the geostrophic component of the vertical velocity, whereas the reanalyses estimate the ~~full-total~~ vertical velocity field. However, the comparison between OGCM's w_g and w_{tot} ~~in~~ (Fig. 4a) illustrates better agreement in spatial patterns and intensity than when comparing OLIV3 with the reanalyses ~~in~~ (Fig. 5a, c and d). For instance, in the subtropical gyres, where the relative differences between OGCM's w_g and w_{tot} are typically below 10%, the ~~relative~~ differences between OLIV3 and any of the reanalyses often exceed this threshold ~~for~~ (Fig. 5f and g). This suggests that much of the observed differences derive from the observation-based input fields (ARMOR3D meridional velocities and ERA5 wind stress) rather than the geostrophic component reconstructed, which mostly dominates the total flow in the subtropics and upper tropics (Fig. 5). ~~Particularly, western boundary current systems correspond to regions with large errors in the geostrophic LVB-derived vertical velocities (hatching in Fig. 5a). In these regions, additional terms of the vorticity equation, such as the bottom pressure torque, close the vorticity budget (e.g. Hughes and De Cuevas, 2001; Gula et al., 2015; Schoonover et al., 2016).~~

OLIV3 demonstrates a reasonable ability to qualitatively capture the time-mean vertical velocity structure of the major ocean gyres at σ_{26} . Even in regions where the LVB assumption is no longer valid, such as the deep tropics, western boundary currents and the subpolar Pacific, OLIV3 estimates often remain within the uncertainty range defined by the intercomparison datasets.




Figure 5. Time-mean vertical velocity fields at σ_{26} and $5^\circ \times 5^\circ$ spatial resolution from: (a) OLIV3, (b) OMEGA3D, (c) GLORYS12v1 reanalysis, and (d) ECCOV4r4 reanalysis. White hatching represents regions within the maximum mixed layer defined by the thermohaline field corresponding to each velocity. (e-j) Differences between modules of the various datasets. In panel (a), black hatching indicates the areas where the LVB is not satisfied in the OGCM (relative error $>10\%$).

3.3.2 Vertical-Diapycnal Structure of Time-mean Vertical Flow

The presence of vertical shear in the vertical velocity field is fundamental for establishing the Sverdrup balance and defining the volumes of water influenced by these dynamics. In classical Sverdrup theory, vertical velocities in the deep ocean are assumed to be very weak, allowing the vertically integrated meridional transport to be related to the wind stress curl. Global estimates of the vertical velocity offer insights into the fundamental physics of the ocean interior circulation. When the LVB

holds, geostrophic vertical velocities in the ocean interior can be interpreted as the residue of the evacuation by meridional transport of the vertical mass flow input from the layer above. If the geostrophic vertical velocity at a given depth is effectively negligible, the divergence of the horizontal flow fully compensates the wind driven divergence above this level, implying that the Sverdrup balance adequately describes the ocean dynamics down to that depth. In this context, the vertical profile of the vertical velocity field adds information about the flow evacuation ratio under Sverdrup’s framework. As demonstrated ~~in Cortés-Morales and Lazar (2024)~~ for the North Atlantic Ocean in CM24, this assumption holds reasonably well in subtropical basins but breaks down at high latitudes. This finding is consistent with recent studies that have directly evaluated Sverdrup balance (e.g. Thomas et al., 2014). Despite its relevance, studies focusing on the Sverdrup balance rarely address the vertical structure of the vertical velocity field.

To further evaluate the ability of OLIV3 to reproduce the vertical structure of the vertical flow, the diapycnal gradient of the absolute value of time-mean vertical velocity (Eq. 16) for multiple estimates of w is computed (Fig. 6). Most datasets (OLIV3, ECCOv4r4 and GLORYS12v1) feature a reduction in downwelling amplitude with depth across the subtropical gyres, consistent with a baroclinic structure (panels a, c and d in Fig. 6). The largest negative gradients reach values ~~around $-10 \times$ above $-1.4 \times 10^{-7} \text{ m s}^{-4} \text{ m day}^{-1} / \text{kg m}^{-3}$ in the Atlantic and Indian subtropical gyres, and near $-5 \times 10^{-7} \text{ m s}^{-1} / \text{kg m}^{-3}$ in the Pacific Ocean~~ m^{-1} in the global subtropical gyres. Compared with the rest of estimates, OMEGA3D ~~showcase (Fig. 6b) displays~~ a more barotropic profile (Fig. 6b, e, h and i). OLIV3 (Fig. 6a) generally captures the reanalyses’ downwelling weakening with depth in the subtropics ~~, but (Fig. 6f and g). Nevertheless,~~ it displays positive vertical diapycnal gradients in the eastern tropical gyres, indicating increasing magnitude with depth, which contrasts with the other datasets. This difference may arise from a shallower thermocline at tropical latitudes compared to the subtropics (Salmon, 1982), which causes σ_{27} to lie below the bottom threshold of the thermocline. When the gradient is recalculated using $\sigma_{26.5}$, as a lower limit, OLIV3 captures a negative gradient in both Pacific and Atlantic tropical gyres (Fig. A1 in Appendix A). As ~~shown~~ for the North Atlantic ~~Ocean~~ application of LVB ~~in Cortés-Morales and Lazar, 2024 (CM24)~~, tropical gyre’s vertical velocities decrease rapidly in the thermocline, remaining an order of magnitude smaller than those at the top of the thermocline. This implies that the lower bounds of the gradient do not substantially bias the vertical structure in a model simulation, as seen in the small relative gradient error between OGCM w_{tot} and w_g in the Pacific and Atlantic tropical gyres ~~in (Fig. 4c)~~. However, there are large uncertainties for observation-based datasets like OLIV3 compared to the reanalyses below the thermocline. OLIV3 shows growing vertical velocities with depth in regions where the time-mean vertical velocities at a given depth also differ, such as the western boundaries and the deep tropics. This suggests that when OLIV3 fails to capture the correct amplitude of the vertical flow, it also fails to reproduce the local vertical structure.

The comparison with existing estimates demonstrates that OLIV3 reproduces the structure in the subtropics and upper tropics (particularly above $\sigma_{26.5}$), capturing both the amplitude and vertical structure of the vertical flow, indicating that the geostrophic component is primary contributor to the observation-based vertical flow and that Ekman pumping vertical velocity is a suitable boundary condition, consistent with the results of the perfect model test, ~~shown in Fig. 4 and Cortés-Morales and Lazar (2024).~~




Figure 6. [Vertical-Diapycnal](#) gradient of time-mean vertical velocity between the base of the maximum mixed layer depth and σ_{27} (Eq. 16) at $5^\circ \times 5^\circ$ spatial resolution, shown for: (a) OLIV3, (b) OMEGA3D, (c) GLORYS12v1, and (d) ECCOV4r4. Negative values indicate a decrease in vertical velocity magnitude with depth, while positive values depict increasing magnitude with depth. [\(e-j\) Differences between modules in the diapycnal gradient of time-mean vertical velocity for the various datasets.](#)

3.3.3 Vertical Velocity Time Variability

The perfect model test (Fig. 4) emphasises the high accuracy in terms of temporal variability of the total vertical velocity by the geostrophic component. To further assess this accuracy, we evaluate the annual variance and the correlation coefficient (R) across the various intercomparison datasets.

The key role of mesoscale activity (Wunsch, 2007) can be detected in the annual variance of vertical velocity at an isopycnal level within the tachocline (Fig. 7). In OLIV3, GLORYS12v1 and ECCOv4r4 (Fig. 7a, c and d), the highest variance values are found along western boundary current regions and in the lower-tropical band, while the subtropical gyre interior below the mixed layer generally displays lower variance. This pattern is consistent with known regions of high mesoscale eddy activity
475 [at the ocean surface](#), such as the western boundary current systems and the deep tropics (e.g. Wunsch, 2007; Barceló-Llull et al., 2025), that is transported into the ocean interior. OMEGA3D (Fig. 6b) deviates from this behaviour, showing a poleward variance increase near the intersection of the isopycnal level with the ocean surface ~~-(Fig. 8a, d and e)~~. ECCOv4r4 (Fig. 7d) maintains a similar variance spatial distribution compared with OLIV3 and GLORYS12v1 but with a weaker variance gradient between the subtropical gyre centres and the western boundary current regions, due to lower maximum values ~~-(Fig. 8c and f)~~.
480 Variance values within the tropical Indian basin, as well as the western tropical Pacific and Atlantic basins, exhibit considerable uncertainty across datasets. Nevertheless, OLIV3 reconstructs a field with variance comparable to that in GLORYS12v1 ([Fig. 8b](#)), even in regions where LVB does not hold. This supports the ability of the geostrophic component to capture the temporal variability of vertical motion at first order, as evidenced by Fig. 4.

In addition to the variance, assessing the ability of OLIV3 to represent interannual variability of w is evaluated through
485 the correlation coefficient between dataset pairs at low resolution at the representative σ_{26} (Fig. 9) and as a function of latitude at three different sigma surfaces (Fig. 10). Across most dataset pairs, the highest correlation values are found in the centres of the subtropical gyres, while the lowest occur in the tropical band and western boundary currents. The comparison of the two reanalyses (Fig. 9a and Fig. 10) ~~showcases~~ [shows](#) an overall relatively low correlation over large fractions of the global thermocline, with maxima within subtropical gyres and parts of the deep tropics, but with latitudinal median values
490 remaining below 0.5 almost everywhere. The comparison of these two reference datasets provides reference values quantifying the uncertainties inherent to the estimation of the vertical velocity component of the flow.

~~It appears that~~ OLIV3 exhibit significant large correlations ($R > 0.6$) with the reanalyses across large portions of the global subtropics, with values exceeding 0.8 in the Pacific and Atlantic Oceans (Fig. 9c and 9e). In tropical regions where the perfect model test indicates weak correlation between w_g and w_{tot} (hatching in Fig. 9a), R typically falls below 0.4. Notably, reanalyses
495 generally show a lower correlation with each other than with OLIV3 in most of the global subtropical band (Fig. 9a vs. 9c, 9e). ~~As shown in Fig. 4, the~~ [The](#) geostrophic component dominates the interannual variability of the vertical flow at these depths ~~-(Fig. 4)~~. Therefore, the reduced inter-reanalysis correlation is likely evidence of the lack of synchronisation in the nonlinear components of vertical flow in assimilated products, while the geostrophic component variability, captured by OLIV3, remains highly correlated. The magnitude and structure of the variability reproduced by the OLIV3 fall within the
500 [envelope range](#) of variability spanned by very commonly used reanalysis-based estimates of w . Although OMEGA3D reaches significant correlation values (R up to 0.7) with the intercomparison reanalyses and OLIV3 in some areas within the open ocean, particularly within the Pacific and Atlantic subtropical gyres (Fig. 9b, d and f), its overall performance is poorer and more spatially limited compared to the results for OLIV3.

The general better performance of OLIV3 compared to OMEGA3D in capturing the temporal variability is further illustrated
505 in Fig. 10, which displays the median correlation coefficient value as a function of latitude at three isopycnal levels. Latitudinal




Figure 7. Annual variance (logarithmic scale) of vertical velocity at σ_{26} and $5^\circ \times 5^\circ$ spatial resolution for: (a) OLIV3, (b) OMEGA3D, (c) GLORYS12v1, and (d) ECCOv4r4. In panel (a), black-gray hatching represents regions where correlation coefficient between OGCM w_{tot} and OGCM w_g is smaller than 0.5 (Fig. 4b).

median correlation values between OLIV3 and ECCOv4r4 in the Northern Hemisphere subtropical band ($20\text{--}40^\circ\text{N}$) are reduced from around 0.6 at $\sigma_{25.5}$ to 0.4 at σ_{27} . Although correlation coefficients between datasets tend to weaken with depth, OLIV3 (solid lines) consistently exhibits higher correlations with the model and reanalyses than OMEGA3D (dashed lines) throughout the entire thermocline except for high latitudes in the Northern Hemisphere.

w variance differences at $\sigma 26 \text{ kg m}^{-3}$




Figure 8. Difference between annual variances of vertical velocities at $\sigma 26$ and $5^\circ \times 5^\circ$ spatial resolution for the various datasets in Fig. 7.

510 3.4 Unveiling Methodological Differences Among the Existing Estimates

Several fundamental methodological differences among OLIV3, OMEGA3D and the two reanalyses may account for the discrepancies observed in the climatological horizontal and baroclinic structure, as well as the temporal evolution of the vertical movements. These include the reconstruction methodology, the components of vertical velocity reconstructed, the atmospheric forcing, the three-dimensional horizontal velocity inputs, and the spatiotemporal resolution of the datasets.

515 Atmospheric forcing does not appear to be the primary source of discrepancies, as all datasets employ variants of ERA atmospheric reanalysis products to force the oceanic surface. Similarly, both OLIV3 and OMEGA3D are based on the ARMOR3D geostrophic velocity field, while reanalyses compute vertical velocities directly from the total assimilated horizontal velocity field. Despite their common input, OLIV3 and OMEGA3D exhibit large differences. In contrast, OLIV3 aligns more closely with the reanalyses despite their distinct origin in either observation-based or assimilated horizontal velocity fields. However,
 520 ~~as discussed in Cortés-Morales and Lazar (2024)~~, the ageostrophic component of the horizontal velocity field is negligible in most of the tropical and subtropical tachocline, as discussed in CM24. Again, while these differences may induce some discrepancies, they are not dominant.

w correlation coefficient at $\sigma 26 \text{ kg m}^{-3}$




Figure 9. Correlation coefficient (R) between the vertical estimates from OLIV3, GLORYS12v1, ECCOv4r4, and OMEGA3D at $\sigma 26$ and $5^\circ \times 5^\circ$ resolution. Dotted squares indicate correlations significant at the 95% confidence level based on the Student t-test. Black hatching represents regions where correlation coefficient between OGCM w_{tot} and OGCM w_g is smaller than 0.5 (Fig. 4b).

Differences in native spatial and temporal resolution may play a significant role, even when data are averaged to a common resolution. Certain phenomena may be retained across scales and not entirely removed (Yeager, 2015). For example, GLORYS12v1 has a resolution of $1/12^\circ$, capturing smaller mesoscale features compared to ECCOv4r4 at 1° resolution. Higher spatial resolutions allow GLORYS12v1 to preserve events at seasonal and sub-seasonal timescales, which could not be maintained in coarser native resolutions. Previous studies on reanalysis intercomparison (e.g. Balmaseda et al., 2015; Carton et al., 2019) have shown that such differences due to resolution (in particular in temperature and salinity) are more pronounced in the tropics. In particular, Carton et al. (2018) shows how the coarser resolution of ECCO leads to very distinct results from other eddy-permitting datasets. The spatial resolutions in the [study-mentioned-mentioned study](#) range from 0.25° to 1° . Therefore, the uncertainty observed here between GLORYS12v1 and ECCOv4r4 probably reproduces and magnifies the uncertainty observed in the cited studies.

OLIV3 and reanalyses estimate w through vertical integration of horizontal velocity fields, because the governing equations only contain the vertical derivative of w . In contrast, OMEGA3D employs the omega equation, which [incorporates the](#) [, although it also requires vertical integration, explicitly includes](#) second-order vertical derivatives and horizontal derivatives




Figure 10. Median correlation coefficient (R) as a function of latitude for $\sigma_{25.5}$, σ_{26} , σ_{27} . The analysis includes only regions where the correlation between OGCM w_g and w_{tot} exceeds 0.5. Solid, dashed and dotted lines correspond to OLIV3, OMEGA3D intercomparisons, and reanalyses intercomparison respectively.

of w . Consequently, Dirichlet (vertical velocities are set to zero) and Neumann (partial derivatives of vertical velocity are set to zero) conditions are imposed as boundary conditions (Buongiorno Nardelli et al., 2018; Buongiorno Nardelli, 2020). The inclusion of the horizontal and second-order derivatives of w in the OMEGA3D framework may explain the discrepancies with the other datasets. However, a comprehensive examination of the sources of these differences would require a separate in-depth investigation, given the complex physics, constraints and assumptions underlying the omega equation.

3.5 Near Surface Interannual Variability of Vertical Flow: Improvement Relative to Ekman Pumping

Beyond the importance of providing an estimate of the vertical profile of thermocline vertical velocities, one might wonder how [near-surface-ocean interior](#) w_g compares with Ekman pumping (w_{Ek}) [from Eq. 9](#), the most commonly used observation-based




Figure 11. (a) Map of correlation coefficients for OGCM w at σ_{26} with Ekman pumping (1993-2015). (b) Differences between Fig. 4b and Fig. 11a. (c) Map of correlation coefficients for OGCM w and w_g at 100 m depth. (d) Map of correlation coefficients for OGCM w at 100 m depth with Ekman pumping. Same isolines as in Fig. 4b for panels a, c and d. For panel c the black contour lines represent the zero. The fields have been smoothed with a 5° running mean.

reference product for vertical transfers between thermocline and nitracline, and the surface waters. Indeed, this wind-based
545 computation is frequently employed to validate against observations, or calculate water mass fluxes (e.g. Marshall et al., 1993; Lazar et al., 2002), as well as transport of biogeochemical tracers (Oschlies, 2002) and marine ecosystem parameters, from fish (Parrish et al., 1981) up to whales (Croll et al., 2005). In these types of studies, Ekman pumping is generally considered as a vertical velocity proxy at a variety of levels depending on questions, time-scales and community habits. This level ranges from the bottom of the Ekman layer to that of the winter mixed layer (Williams et al., 2006), an isopycnal surface close to σ_{26} or a
550 fixed depth of a few tens of meters up to 200 meters (Palter et al., 2013). Here, ~~we chose to present results in both the interannual synchrony of the Ekman pumping at the ocean surface and both the geostrophic and total vertical velocities is examined in isopycnal and depth frames, on σ_{26} and at 100 m depth. Again, the computation is conducted with the reference OGCM simulation, considered to be a dynamically coherent estimate of the real ocean.~~ Figure 11 coordinates (Fig. 11) to highlight the assess of the geostrophic current divergence on the variability of vertical velocities in the ocean interior. Specifically, Fig. 11a displays the correlation between w_{Ek} at ocean surface and w_{tot} on σ_{26} and the difference with the correlation, while Fig. 11b represents the correlation difference between w_g and w_{tot} shown Fig. 4b. Panels c and d in Fig. 11 illustrate the the Figures 11c and 11d illustrate, respectively, the correlation between w_g and w_{tot} and at 100m depth, and the correlation between between w_{Ek} at ocean surface and w_{tot} at 100m depth, respectively. This comparison also illustrates the extent to which Ekman pumping can serve as a reliable proxy for describing the temporal variability of the total vertical flow within
560 the ocean interior. Again, the computation is conducted with the reference OGCM simulation, considered to be a dynamically coherent estimate of the real ocean.

Ekman pumping estimates well ($R > 0.7$) up to very well ($R > 0.9$) the variability of total vertical velocities along this isopycnal surface over a relatively small proportion of the thermocline, limited to certain parts of the subtropical gyres and the tropical Atlantic Ocean (Fig. 11a). However, comparison with the correlation between w_{tot} and w_g (Fig. 4b), quantified in
565 Fig. 11b, shows that w_g is accurate over much larger areas of the globe. In other words, w_{Ek} is only more accurate than w_g in regions where both estimates are poor ($R < 0.7$), such as western boundary currents and intense open ocean currents, such as the NAD and ACC with a strong zonal component. A marked decrease in the correlation coefficient is observed towards the western boundaries, where the isopycnal surfaces deepen, within the western boundary current systems, as well as in the deep tropics. When comparing this pattern to the correlation between w_g and w_{tot} shown in Fig. 4b, it is remarkable that the areas
570 with high correlation are even more extensive in all basins.

We extended the comparison to other vertical levels, in particular to depths of 50 (not shown) and 100 m (Fig. 11c and d), and reached the same conclusion. Overall, these results show that the Ekman pumping ~~model~~ alone is insufficient to account for interannual variability in vertical ~~flux-flow~~ in most regions of the globe. However, the inclusion of geostrophic meridional transport divergence in Ekman pumping, i.e. w_g , brings about a significant improvement. This highlights the advancement of
575 geostrophic vertical velocities as an estimator of total vertical fluxflow variability. Moreover, vertical movement in most eastern boundary upwelling systems, such as Benguela or Peru, shows weak correlations with Ekman pumping. This is consistent with the importance of remote forcing from trapped waves on the coast in these regions (Polo et al., 2008; Illig et al., 2014; Bachèlery et al., 2016).

4 Conclusions

580 ~~A-This study introduces a~~ novel observation-based global dataset of geostrophic vertical velocities (w) within the thermocline (Observation-based Linear Vorticity Vertical Velocities; OLIV3) is presented and validated against other existing estimates. ~~It is derived from the depth-integrated geostrophic~~ By applying the linear vorticity balance (LVB) ~~applied to in situ and satellite geostrophic meridional velocity fields and satellite surface wind stress.~~ to ARMOR3D meridional velocities and ERA5 Ekman pumping. OLIV3 provides global ~~ocean estimates of geostrophic vertical velocities at horizontal~~ a physically consistent
585 and observation-based framework to resolve large-scale vertical transport at a 0.25° resolution, 71 vertical levels horizontal resolution and annual frequency, covering the period 1993 - 2019. The ~~product estimates, to the first order, significant parts of the main well-known features of the large-scale tropical and subtropical global thermoelne vertical circulation at effective horizontal scales larger than 5°, in agreement with previous estimates. Most importantly, it also allows a reconstruction of the interannual variability of most of the ocean thermocline, including eastern basin boundaries, to the notable exception of western boundaries, the North Atlantic Drift (NAD) and the Antarctic Circumpolar Current (ACC).~~ dataset is available publicly
590 at <https://doi.org/10.5281/zenodo.16962780>.

The feasibility of the LVB framework for reconstructing vertical flow is demonstrated by the good agreement between the OGCM geostrophic vertical velocity estimates and the model's total velocity output. The spatial distribution and vertical shear of the temporal means of total velocity are well-reproduced by the geostrophic estimates, except in in regions dominated by

595 ~~nonlinear dynamics such as~~ the deep tropics and along western ~~boundaries across all basins, where nonlinear components~~
~~of w have a substantial contribution~~boundary currents. Furthermore, despite some mean biases across the globe, the model's
geostrophic vertical velocity field effectively captures the interannual variability of vertical flow, with correlations exceeding
0.9 across most of the tropical, subtropical, and extratropical global oceans, excluding western boundary currents and also rel-
atively intense zonal currents of the deep tropics, the NAD, and the ACC. This analysis confirms the dominance of geostrophic
600 meridional transport in driving the interannual variability, while nonlinear components primarily influence the time-mean am-
plitude.

~~A The~~ comparison of OLIV3 against ~~three independent datasets, including~~ two reanalyses (ECCOv4r4 and GLORYS12v1)
~~and an observation-based product (OMEGA3D), over the 1993-2015 period,~~ demonstrates that OLIV3 reproduces the large-
scale horizontal patterns and baroclinic vertical structure of the climatological tachocline circulation ~~It over the 1993-2015~~
605 period. OLIV3 also captures the interannual variability in most of the open-ocean tropical and subtropical regions when com-
pared to the reanalyses. The poorest performance of OLIV3 in the various metrics analysed is found across the same regions
where LVB failed to reconstruct vertical flow in the OGCM perfect model test: western boundary currents, the zonal tropical
currents, the NAD, and the ACC, ~~aligning with the OGCM preliminary test~~. Remarkably, OLIV3 often correlates as well or
better with each reanalysis than the reanalyses do with each other, suggesting that the geostrophic signal variability is co-
herently captured, while nonlinear components suffer from a high degree of uncertainty in ~~present~~ estimates of global ocean
610 thermocline vertical velocities.

~~With the introduction of OLIV3, two observation-based three-dimensional estimates of the vertical velocity field are now~~
~~available: OLIV3 derived from the LVB, and OMEGA3D based on the quasi-geostrophic omega equation.~~ The intercomparison
with OMEGA3D, the only other existing observation-based product, evidences the systematic improvement offered by OLIV3.
615 ~~In particular,~~ OMEGA3D reproduces a relatively barotropic structure, which contrasts with the vertical shear observed in other
products and the baroclinic ocean required to sustain the Sverdrup balance. Additionally, OMEGA3D exhibits an overall lower
interannual synchrony with reanalyses. ~~We suppose that the discrepancies using the~~ These discrepancies likely arise from
the complexity of the omega equation compared to the LVB ~~arise from the large number of terms in the former~~, including
higher-order vertical and horizontal derivatives of w , ~~many of which that~~ require boundary conditions extremely difficult to
620 compute.

~~An additional comparison within the OGCM reveals that geostrophic vertical velocities (w_g) and Ekman pumping (w_{EK})~~
OLIV3 has shown to be a useful tool for investigating interannual variability across the thermocline in polar, subtropical,
and tropical gyres, with the exception of their western boundary currents, as well as most of the eastern boundary upwelling
systems, offshore of the continental plateau. We strongly encourage its use in biogeochemical and biological studies focused on
625 the vertical structure and exchange of ocean biogeochemical tracers, and on their impact on marine ecosystems, at interannual
scales over large regions of the global ocean thermocline.

The wind-driven divergence at the surface and the vertical flow in the ocean interior are strongly correlated across large
portions of the global tropical and subtropical gyres. ~~Nonetheless, this coupling weakens toward the western boundaries and~~
~~the deep tropics, indicating that meridional transport dynamics, from which the geostrophic component of the vertical flow is~~

630 ~~derived, increasingly dominate the variability of the vertical flow. Furthermore, the analysis demonstrates that w_g , as supported~~
~~by the comparison between the OGCM estimation of the total vertical velocities in the ocean interior and Ekman pumping~~
~~(w_{Ek}). However, geostrophic vertical velocities offers a systematically better or equal accuracy than Ekman pumping in capturing~~
~~the interannual variability of the total vertical flow than w_{Ek} , except in regions of relatively intense zonal~~
635 ~~confidence. These results emphasise the better performance of geostrophic vertical velocities as an estimator of the vertical~~
~~flow variability in the ocean interior compared to the Ekman pumping alone. We propose to extend this result these findings~~
~~to the real ocean and consider, suggesting that OLIV3 should be more accurate provides a more accurate velocity field than~~
~~Ekman pumping for estimating mass, and physical (temperature and salinity) and biogeochemical (nutrients, oxygen, CO₂)~~
~~tracer fluxes between the surface and the pycnocline and nutrient line layers, nutricline layers. Observation-based knowledge~~
640 ~~of these flows is fundamental not only for a better understanding of the ocean's role in ongoing climate change but also for~~
~~assessing the extent of oligotrophy, oxygen minimum zone, and acidification in the anthropogenic ocean.~~

~~OLIV3 is a new and useful tool for investigating interannual variability in polar, subtropical, and tropical gyres, with the~~
~~exception of their western boundary currents, as well as most parts of the eastern boundary upwelling systems, offshore of the~~
~~continental plateau. We strongly encourage its use in biogeochemical and biological studies focused on the vertical exchange~~
645 ~~of ocean biogeochemical tracers and their impact on marine ecosystems.~~

~~In the future, OLIV3 could be improved to take into account seasonal and monthly scales and represent the specific processes~~
~~of coastal areas while maintaining the simplicity of the depth-integrated formalism. Overcoming the current limitations in~~
~~OLIV3 (spatiotemporal scales coarser than 5° and one year) would require incorporating full total meridional velocities and ad-~~
~~ditional terms from the vorticity equation, such as the horizontal advection of relative vorticity. This improvement could extend~~
650 ~~OLIV3's applicability to finer spatial and temporal scales, including seasonal and monthly variability, and to coastal regions~~
~~while maintaining the simplicity of the depth-integrated formalism. Detailed works focusing on regions strongly affected by~~
~~climate change and extreme events will be conducted in future studies using the upcoming improved version of OLIV3.~~

Code and data availability. The OLIV3 dataset developed in this study is available at <https://doi.org/10.5281/zenodo.16962780> (Cortés-Morales and Lazar, 2025). A low-resolution version, used in the intercomparison test of this study, is available upon request. Codes in MatLab
655 and Python to compute geostrophic velocities (OLIV3 and OGCM), apply the linear vorticity balance and calculate the intercomparison
metrics is available at the following repository: https://github.com/dcortales/compute_wglvb.




Figure A1. Vertical Diapycnal gradient of time-mean vertical velocity between the base of the maximum mixed layer depth and $\sigma_{26.5}$ (Eq. 16) at $5^\circ \times 5^\circ$ spatial resolution, shown for: (a) *OLIV3*, (b) *OMEGA3D*, (c) *GLORYS12v1*, and (d) *ECCOV4r4*. Negative values indicate a decrease in vertical velocity magnitude with depth, while positive values depict increasing magnitude with depth.

Appendix A: Supplementary figures

Author contributions. D. C.-M and A. L. conceptualized and designed the study. DCM processed the data, produced the figures and first draft of the manuscript, together with the associated data products. All authors have reviewed the manuscript. All authors have read and agreed to the published version of the manuscript.

660

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. This work was supported by CNES, LEFE-MErcator and the European Space Agency 4DMED-SEA project (ESA Contract No. 4000141547/23/I-DT). We deeply thank the anonymous reviewers for their useful fundamental and detailed suggestions that allowed the improvement of the manuscript.

665 References

- Adcroft, A., Hill C., Campin, J.-M., Marshall, J., and Heimbach, P.: Overview of the formulation and numerics of the mit gcm. In: Proceedings of the ECMWF seminar series on Numerical Methods, Recent developments in numerical methods for atmosphere and ocean modelling, pp. 139–149, 2004
- Allen, J., Smeed, D., Nurser, A., Zhang, J., and Rixen, M.: Diagnosis of vertical velocities with the qg omega equation: An examination of the errors due to sampling strategy. *Deep Sea Research Part I: Oceanographic Research Papers* 48(2):315–346, [https://doi.org/10.1016/S0967-0637\(00\)00035-2](https://doi.org/10.1016/S0967-0637(00)00035-2), 2001
- 670 Aristegui, J., Barton, E. D., Alvarez-Salgado, X. A., Santos, A. M. P., Figueiras, F. G., and Kifani Kifani, S., Hernandez-Leon, S., Mason, E., Machu, E., and Demarcq, H.: Sub-regional ecosystem variability in the canary current upwelling. *Progress in Oceanography* 83(1-4):33–48, <https://doi.org/10.1016/j.pocean.2009.07.031>, 2009
- 675 AVISO+: SSALTO/DUACS User Handbook, CLS-DOS-NT-06- 034, Issue 4.4, SALP-MU-P-EA-21065-CLS, 2015
- Bachèlery, M. L., Illig, S., and Dadou, I.: Interannual variability in the South-East Atlantic Ocean, focusing on the Benguela Upwelling System: Remote versus local forcing. *Journal of Geophysical Research: Oceans*, 121(1), 284-310, <https://doi.org/10.1002/2015JC011168>, 2016
- Balmaseda, M. A., Hernandez, F., Storto, A., Palmer, M., Alves, O., Shi, L., Smith, G. C., and Toyoda, T., Valdivieso M., Barnier B., et al.: The ocean reanalyses intercomparison project (ora-ip). *Journal of Operational Oceanography* 8(sup1):s80–s97, <https://doi.org/10.1080/1755876X.2015.1022329>, 2015
- 680 Barceló-Llull, B., Rosselló, P., Combes, V., Sánchez-Román, A., Pujol, M. I., and Pascual, A.: Kuroshio Extension and Gulf Stream dominate the Eddy Kinetic Energy intensification observed in the global ocean. *Scientific Reports*, 15(1), 21754, <https://doi.org/10.1038/s41598-025-06149-9>, 2025
- 685 Bessieres, L., Leroux, S., Brankart, J.-M., Molines, J.-M., Moine, M.-P., Bouttier, P.-A., Penduff, T., Terray, L., Barnier, B., and Serazin, G.: Development of a probabilistic ocean modelling system based on nemo 3.5: application at eddying resolution. *Geoscientific Model Development* 10(3):1091–1106, <https://doi.org/10.5194/gmd-10-1091-2017>, 2017
- Bower, A. S. and Rossby, T.: Evidence of cross-frontal exchange processes in the gulf stream based on isopycnal rafof float data. *Journal of Physical Oceanography* 19(9):1177–1190, [https://doi.org/10.1175/1520-0485\(1989\)019<1177:EOCFEP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1989)019<1177:EOCFEP>2.0.CO;2), 1989
- 690 Bretherton, F. P., Davis, R. E., and Fandry, C.: A technique for objective analysis and design of oceanographic experiments applied to mode-73. In: *Deep Sea Research and Oceanographic Abstracts*, Elsevier, vol 23, pp. 559–582, [https://doi.org/10.1016/0011-7471\(76\)90001-2](https://doi.org/10.1016/0011-7471(76)90001-2), 1976
- Buongiorno Nardelli, B., Santoleri, R., and Sparnocchia, S.: Small mesoscale features at a meandering upper-ocean front in the western ionian sea (mediterranean sea): Vertical motion and potential vorticity analysis. *Journal of physical oceanography* 31(8):2227–2250, [https://doi.org/10.1175/1520-0485\(2001\)031<2227:SMFAAM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<2227:SMFAAM>2.0.CO;2), 2001
- 695 Buongiorno Nardelli B., Mulet S., and Iudicone D.: Three-dimensional ageostrophic motion and water mass subduction in the southern ocean. *Journal of Geophysical Research: Oceans* 123(2):1533–1562, <https://doi.org/10.1002/2017JC013316>, 2018
- Buongiorno Nardelli, B.: A multi-year time series of observation-based 3d horizontal and vertical quasi-geostrophic global ocean currents. *Earth System Science Data* 12(3):1711–1723, <https://doi.org/10.5194/essd-12-1711-2020>, 2020
- 700 Carton, J. A., Chepurin, G. A., and Chen, L.: Soda3: A new ocean climate reanalysis. *Journal of Climate* 31(17):6967–6983, <https://doi.org/10.1175/JCLI-D-18-0149.1>, 2018

- Carton, J. A., Penny, S. G., and Kalnay, E.: Temperature and salinity variability in the soda3, ecco4r3, and oras5 ocean reanalyses, 1993–2015. *Journal of Climate* 32(8):2277–2293, <https://doi.org/10.1175/JCLI-D-18-0605.1>, 2019
- Charney J.: ~~The generation of oceanic currents by wind. *Journal of Marine Research* https://elischolar.library.yale.edu/journal_of_marine_research/858, 1955-~~
- 705
- Christensen, K. M., Gray, A., and Riser, S.: Global estimates of mesoscale vertical velocity near 1000 m from argo observations. Authorea Preprints <https://10.22541/essoar.168500356.69644187/v1>, 2023
- [Colin de Verdière, A., and Ollitrault, M.: A direct determination of the World Ocean barotropic circulation. *Journal of Physical Oceanography*, 46\(1\), 255-273, https://doi.org/10.1175/JPO-D-15-0046.1, 2016](https://doi.org/10.1175/JPO-D-15-0046.1)
- 710 [Colin de Verdière, A., Meunier, T., and Ollitrault, M.: Meridional overturning and heat transport from Argo floats displacements and the planetary geostrophic method \(PGM\): Application to the subpolar North Atlantic. *Journal of Geophysical Research: Oceans*, 124\(8\), 6270-6285, https://doi.org/10.1029/2018JC014565, 2019](https://doi.org/10.1029/2018JC014565)
- Comby, C., Barrillon, S., Fuda, J.-L., Doglioli, A. M., Tzortzis, R., Gregori, G., Thyssen, M., and Petrenko, A.: Measuring vertical velocities with adcps in low-energy ocean. *Journal of Atmospheric and Oceanic Technology* 39(11):1669–1684, <https://doi.org/10.1175/JTECH-D-21-0180.1>, 2022
- 715
- Cortés-Morales, D., and Lazar, A.: Vertical Velocity 3D Estimates from the Linear Vorticity Balance in the North Atlantic Ocean. *Journal of Physical Oceanography*, 54(10), 2185-2203, <https://doi.org/10.1175/JPO-D-24-0011.1>, 2024
- ~~Cortés-Morales, D., and Lazar, A.: Global Observation-based LInear Vorticity Vertical Velocities (OLIV3) over isopycnal levels [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.16962780>, 2025a-2025~~
- 720 ~~Cortés-Morales, D., and Lazar, A.: Global Observation-based LInear Vorticity Vertical Velocities (OLIV3) over vertical levels Data set. Zenodo. <https://doi.org/10.5281/zenodo.16981061>, 2025b-~~
- Croll, D. A., Marinovic, B., Benson, S., Chavez, F. P., Black, N., Ternullo, R., and Tershly, B. R.: From wind to whales: trophic links in a coastal upwelling system. *Marine Ecology Progress Series*, 289, 117-130, <https://doi.org/10.3354/meps289117>, 2005
- Daniault, N., Mercier, H., and Lherminier, P.: The 1992–2009 transport variability of the East Greenland-Irminger Current at 60 N. *Geophysical Research Letters*, 38(7), 2011
- 725
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *Journal of Geophysical Research: Oceans*, 109(C12), <https://doi.org/10.1029/2004JC002378>, 2004
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The era-interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the royal meteorological society* 137(656):553–597, <https://doi.org/10.1002/qj.828>, 2011
- 730
- DeVries, T., Holzer, M., and Primeau, F.: Recent increase in oceanic carbon uptake driven by weaker upper-ocean overturning. *Nature* 542(7640):215–218, <https://doi.org/10.1038/nature21068>, 2017
- Droghei, R., Buongiorno Nardelli, B., and Santoleri, R.: A new global sea surface salinity and density dataset from multivariate observations (1993–2016). *Frontiers in Marine Science* 5:84, <https://doi.org/10.3389/fmars.2018.00084>, 2018
- 735
- Dussin, R., Barnier, B., Brodeau, L., and Molines, J. M.: Drakkar forcing set dfs5. MyOcean Report https://FR-330-715-368-00032-IFR_DRAKKAR_DFSS5.2, 2016

- D'Asaro, E. A., Shcherbina, A. Y., Klymak, J. M., Molemaker, J., Novelli, G., Guigand, C. M., Haza, A. C., Haus, B. K., Ryan, E. H., Jacobs, G. A., et al.: Ocean convergence and the dispersion of flotsam. *Proceedings of the National Academy of Sciences* 115(6):1162–1167, 745 <https://doi.org/10.1073/pnas.1718453115>, 2018
- Elmoussaoui, A., Arhan, M., and Treguier, A.-M.: Model-inferred upper ocean circulation in the eastern tropics of the north atlantic. *Deep Sea Research Part I: Oceanographic Research Papers* 52(7):1093–1120, <https://doi.org/10.1016/j.dsr.2005.01.010>, 2005
- Falkowski, P. G., Ziemann, D., Kolber, Z., and Bienfang, P. K.: Role of eddy pumping in enhancing primary production in the ocean. *Nature*, 352(6330), 55-58, <https://doi.org/10.1038/352055a0>, 1991
- 745 Faye, S., Lazar, A., Sow, B. A., and Gaye, A. T.: A model study of the seasonality of sea surface temperature and circulation in the Atlantic North-eastern Tropical Upwelling System. *Frontiers in Physics*, 3, 76, <https://doi.org/10.3389/fphy.2015.00076>, 2015
- Fischer, J., Leach, H., and Woods, J. D.: A synoptic map of isopycnic potential vorticity in the seasonal thermocline. *Journal of physical oceanography*, 19(4), 519-531, [https://doi.org/10.1175/1520-0485\(1989\)019<0519:ASMOIP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1989)019<0519:ASMOIP>2.0.CO;2), 1989
- ~~Forget~~
- 750 ~~Forget, G., Campin, J.-M., Heimbach, P., Hill, C., Ponte, R., and Wunsch, C.: Ecco version 4: An integrated framework for non-linear inverse modeling and global ocean state estimation. *Geoscientific Model Development* 8(10):3071–3104, <https://doi.org/10.5194/gmd-8-3071-2015>, 2015~~
- Foukal, N. P., and Lozier, M. S.: Assessing variability in the size and strength of the North Atlantic subpolar gyre. *Journal of Geophysical Research: Oceans*, 122(8), 6295-6308, 2017
- 755 Freeland, H. J.: Vertical velocity estimates in the north pacific using argo floats. *Deep Sea Research Part II: Topical Studies in Oceanography* 85:75–80, <https://doi.org/10.1016/j.dsr2.2012.07.019>, 2013
- Freilich, M. and Mahadevan, A.: Decomposition of vertical velocity for nutrient transport in the upper ocean. *Journal of Physical Oceanography* 49(6):1561–1575, <https://doi.org/10.1175/JPO-D-19-0002.1>, 2019
- 760 ~~Freilich M. and Mahadevan A.: Coherent pathways for subduction from the surface mixed layer at ocean fronts. *Journal of Geophysical Research: Oceans* 126(5), <https://doi.org/10.1029/2020JC017042>, 2021~~
- Fukumori, I., Heimbach, P., Ponte R. M., and Wunsch C.: A dynamically consistent, multivariable ocean climatology. *Bulletin of the American Meteorological Society* 99(10):2107–2128, <https://doi.org/10.1175/BAMS-D-17-0213.1>, 2018
- ~~Garabato A. C. N., Leach H., Allen J. T., Pollard R. T., and Strass V. H.: Mesoscale subduction at the antarctic polar front driven by baroclinic instability. *Journal of Physical Oceanography* 31(8):2087–2107, [https://doi.org/10.1175/1520-0485\(2001\)031<2087:MSATAP>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<2087:MSATAP>2.0.CO;2), 2001~~
- 765 Garcia-Jove, M., Mourre, B., Zarokanellos, N. D., Lermusiaux, P. F., Rudnick, D. L., and Tintore, J.: Frontal dynamics in the alboran sea: 2. processes for vertical velocities development. *Journal of Geophysical Research: Oceans* 127(3), <https://doi.org/10.1029/2021JC017428>, 2022
- 770 Giglio, D., Roemmich, D., and Cornuelle, B.: Understanding the annual cycle in global steric height. *Geophysical research letters* 40(16):4349–4354, <https://doi.org/10.1002/grl.50774>, 2013
- Giordani, H., Prieur, L., and Caniaux, G.: Advanced insights into sources of vertical velocity in the ocean. *Ocean Dynamics* 56(5):513–524, <https://doi.org/10.1007/s10236-005-0050-1>, 2006

- Gomis, D., Ruiz, S., and Pedder, M. A.: Diagnostic analysis of the 3d ageostrophic circulation from a multivariate spatial interpolation of ctd and adcp data. *Deep Sea Research Part I: Oceanographic Research Papers* 48(1):269–295, [https://doi.org/10.1016/S0967-0637\(00\)00060-1](https://doi.org/10.1016/S0967-0637(00)00060-1), 2001
- Guinehut, S., Dhomp, A.-L., Larnicol, G., and Le Traon, P.-Y.: High resolution 3-d temperature and salinity fields derived from in situ and satellite observations. *Ocean Science* 8(5):845–857, <https://doi.org/10.5194/os-8-845-2012>, 2012
- Gula, J., Molemaker, M. J., and McWilliams, J. C.: Gulf stream dynamics along the southeastern us seaboard. *Journal of Physical Oceanography* 45(3):690–715, <https://doi.org/10.1175/JPO-D-14-0154.1>, 2015
- Halpern, D. and Freitag, H. P.: Vertical motion in the upper ocean of the equatorial eastern pacific. *Oceanologica Acta, Special Issue* <https://archimer.ifremer.fr/doc/00267/37842/>, 1987
- Halpern, D., Knox, R. A., Luther, D. S., and Philander, S. G. H.: Estimates of equatorial upwelling between 140 and 110 w during 1984. *Journal of Geophysical Research: Oceans* 94(C6):8018–8020, <https://doi.org/10.1029/JC094iC06p08018>, 1989
- 785 [Helber, R. W., and Weisberg, R. H.: Equatorial upwelling in the western Pacific warm pool. *Journal of Geophysical Research: Oceans*, 106\(C5\), 8989-9003, <https://doi.org/10.1029/2000JC000401>, 2001](https://doi.org/10.1029/2000JC000401)
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 monthly averaged data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.f17050d7 (Accessed on 01/07/2024), 2023
- 790 Hoskins, B. J., Draghici, I., and Davies, H. C.: A new look at the ω -equation. *Quarterly Journal of the Royal Meteorological Society*, 104(439), 31-38, <https://doi.org/10.1002/qj.49710443903>, 1978
- Huang, R. X. and Russell, S.: Ventilation of the subtropical north pacific. *Journal of Physical Oceanography* 24(12):2589–2605, [https://doi.org/10.1175/1520-0485\(1994\)024<2589:VOTSNP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<2589:VOTSNP>2.0.CO;2), 1994
- Hughes, C. W. and De Cuevas, B. A.: Why western boundary currents in realistic oceans are inviscid: A link between form stress and bottom pressure torques. *Journal of Physical Oceanography* 31(10):2871–2885, of the subtropical north pacific. *Journal of Physical Oceanography* 24(12):2589–2605, [https://doi.org/10.1175/1520-0485\(2001\)031<2871:WWBCIR>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<2871:WWBCIR>2.0.CO;2), 2001
- 795 Illig, S., Dewitte, B., Goubanova, K., Cambon, G., Boucharel, J., Monetti, F., Romero, C., Purca, S. and Flores, R.: Forcing mechanisms of intraseasonal SST variability off central Peru in 2000–2008. *Journal of Geophysical Research: Oceans*, 119(6), 3548-3573, <https://doi.org/10.1002/2013JC009779>, 2014
- 800 Ingleby, B. and Huddleston, M.: Quality control of ocean temperature and salinity profiles historical and real-time data. *Journal of Marine Systems* 65(1-4):158–175, <https://doi.org/10.1016/j.jmarsys.2005.11S019>, 2007
- Jackett, D. R., McDougall, T. J., Feistel, R., Wright, D. G., and Griffies, S. M.: Algorithms for density, potential temperature, conservative temperature, and the freezing temperature of seawater. *Journal of Atmospheric and Oceanic Technology* 23(12):1709–1728, <https://doi.org/10.1175/JTECH1946.1>, 2006
- 805 Jacox, M. G., Edwards, C. A., Hazen, E. L., and Bograd, S. J.: Coastal upwelling revisited: Ekman, Bakun, and improved upwelling indices for the U.S. West Coast. *Journal of Geophysical Research: Oceans*, 123, 7332–7350, <https://doi.org/10.1029/2018JC014187>, 2018
- ~~Jean-Michel L., Eric G., Romain B.-B., Gilles G., Angélique M., Marie D., Clément B., Mathieu H., Olivier L. G., Charly R., et al.: The copernicus global 1/12 oceanic and sea ice glorys12 reanalysis. *Frontiers in Earth Science* 9:698,876, <https://doi.org/10.3389/feart.2021.698876>, 2021~~
- 810 Kara, A. B., Rochford, P. A., and Hurlburt, H. E.: An optimal definition for ocean mixed layer depth. *Journal of Geophysical Research: Oceans*, 105(C7), 16803-16821, <https://doi.org/10.1029/2000JC900072>, 2000

- Kara, A. B., Rochford, P. A., and Hurlburt, H. E.: Mixed layer depth variability over the global ocean. *Journal of Geophysical Research: Oceans*, 108(C3), <https://doi.org/10.1029/2000JC000736>, 2003
- 815 Khatri, H., Griffies, S. M., Storer, B. A., Buzzicotti, M., Aluie, H., Sonnewald, M., Dussin, R., and Shao, A. E.: A scale-dependent analysis of the barotropic vorticity budget in a global ocean simulation. *Journal of Advances in Modeling Earth Systems* <https://doi.org/10.22541/essoar.168394747.71837050/v1>, 2023
- Klein, P. and Lapeyre, G.: The oceanic vertical pump induced by mesoscale and submesoscale turbulence. *Annual review of marine science* 1:351–375, <https://doi.org/10.1146/annurev.marine.010908.163704>, 2009
- ~~Koelling, J., Send, U., and Lankhorst, M.: Decadal strengthening of interior flow of north atlantic deep water observed by grace satellites. *Journal of Geophysical Research: Oceans* 125(11), <https://doi.org/10.1029/2020JC016217>, 2020~~
- 820 Lazar, A., Inui, T., Malanotte-Rizzoli, P., Busalacchi, A. J., Wang, L., and Murtugudde, R.: Seasonality of the ventilation of the tropical Atlantic thermocline in an ocean general circulation model. *Journal of Geophysical Research: Oceans*, 107(C8), 18-1, <https://doi.org/10.1029/2000JC000667>, 2002
- Leach, H.: The diagnosis of synoptic-scale vertical motion in the seasonal thermocline. *Deep Sea Research Part A. Oceanographic Research Papers*, 34(12), 2005-2017, [https://doi.org/10.1016/0198-0149\(87\)90095-1](https://doi.org/10.1016/0198-0149(87)90095-1), 1987.
- 825 Lellouche, J.-M., Greiner, E., Le Galloudec, O., Garric, G., Regnier, C., Drevillon, M., Benkiran, M., Testut, C.-E., Bourdalle-Badie, R., Gasparin, F., et al.: Recent updates to the copernicus marine service global ocean monitoring and forecasting real-time 1/12 high-resolution system. *Ocean Science* 14(5):1093–1126, <https://doi.org/10.5194/os-14-1093-2018>, 2018
- [Lellouche, J.-M., Greiner, E., Bourdalle-Badie, R., Garric, G., Melet, A., Drevillon, M., Bricaud, C., Hamon, M., Le Galloudec, O., Regnier, C., et al.: The copernicus global 1/12 oceanic and sea ice glorys12 reanalysis. *Frontiers in Earth Science* 9:698,876, <https://doi.org/10.3389/feart.2021.698876>, 2021](https://doi.org/10.3389/feart.2021.698876)
- 830 Liang, X., Spall, M., and Wunsch, C.: Global ocean vertical velocity from a dynamically consistent ocean state estimate. *Journal of Geophysical Research: Oceans*, 122(10), 8208-8224, <https://doi.org/10.1002/2017JC012985>, 2017
- Liao, F., Liang, X., Li, Y., and Spall, M.: Hidden upwelling systems associated with major western boundary currents. *Journal of Geophysical Research: Oceans*, 127(3), e2021JC017649, <https://doi.org/10.1029/2021JC017649>, 2022
- 835 Madec, G., Bourdalle-Badie, R., Chanut, J., Clementi, E., Coward, A., Ethe, C., Iovino, D., Lea, D., Levy, C., Lovato, T., et al.: Nemo ocean engine, scientific notes of climate modelling center, 2019
- Mahadevan, A.: The impact of submesoscale physics on primary productivity of plankton. *Annual review of marine science* 8:161–184, <https://doi.org/10.1146/annurev-marine-010814-015912>, 2016
- 840 Marshall, J. C., Williams, R. G., and Nurser, A. G.: Inferring the subduction rate and period over the North Atlantic. *Journal of Physical Oceanography*, 23(7), 1315-1329, [https://doi.org/10.1175/1520-0485\(1993\)023<1315:ITSRAP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<1315:ITSRAP>2.0.CO;2), 1993
- Mulet, S., Rio, M.-H., Mignot, A., Guinehut, S., and Morrow, R.: A new estimate of the global 3d geostrophic ocean circulation based on satellite data and in situ measurements. *Deep Sea Research Part II: Tropical Studies in Oceanography* 77:70–81, <https://doi.org/10.1016/j.dsr2.2012.04.012>, 2012
- 845 Munk, W. H.: Abyssal recipes. In: *Deep sea research and oceanographic abstracts*, Elsevier, vol 13, pp. 707–730, [https://doi.org/10.1016/0011-7471\(66\)90602-4](https://doi.org/10.1016/0011-7471(66)90602-4), 1966
- ~~Ohlraut, M. and Rannou, J.-P.: Andro: An argo-based deep displacement dataset~~

- Naveira Garabato, A. C., Leach, H., Allen, J. T., Pollard, R. T., and Strass, V. H.: Mesoscale subduction at the antarctic polar front driven by baroclinic instability. *Journal of Atmospheric and Oceanic Technology* 30(4):759 *Physical Oceanography* 31(8):2087–7882107, <https://doi.org/10.1175/JTECH-D-12-00073.1>, 2013-1520-0485(2001)031<2087:MSATAP>2.0.CO;2, 2001
- 850 Oschlies, A.: Nutrient supply to the surface waters of the North Atlantic: A model study. *Journal of Geophysical Research: Oceans*, 107(C5), 14-1, <https://doi.org/10.1029/2000JC000275>, 2002
- Palter, J.B., Marinov, I., Sarmiento, J.L., and Gruber, N.: Large-Scale, Persistent Nutrient Fronts of the World Ocean: Impacts on Biogeochemistry. In: Belkin, I.M. (eds) *Chemical Oceanography of Frontal Zones. The Handbook of Environmental Chemistry*, vol 116. Springer, Berlin, Heidelberg, https://doi.org/10.1007/698_2013_241, 2013
- 855 Parrish, R. H., Nelson, C. S., and Bakun, A.: Transport mechanisms and reproductive success of fishes in the California Current. *Biological Oceanography*, 1(2), 175-203, <https://doi.org/10.1080/01965581.1981.10749438>, 1981
- Pauly, D., and Christensen, V.: Primary production required to sustain global fisheries. *Nature*, 374(6519), 255-257, <https://doi.org/10.1038/374255a0>, 1995
- 860 Pedlosky, J.: *Ocean circulation theory*. Springer Science and Business Media, 1996
- Penduff, T., Barnier, B., Terray, L., Bessieres, L., Serazin, G., Gregorio, S., Brankart, J., Moine, M., Molines, J., and Brasseur, P.: Ensembles of eddy ocean simulations for climate. *CLIVAR Exchanges, Special Issue on High Resolution Ocean Climate Modelling* 19, 2014
- Pinto, J. G., and Raible, C. C.: Past and recent changes in the North Atlantic oscillation. *Wiley Interdisciplinary Reviews: Climate Change*, 3(1), 79-90, 2012
- 865 Pollard, R. T. and Regier, L. A.: Vorticity and vertical circulation at an ocean front. *Journal of Physical Oceanography* 22(6):609–625, [https://doi.org/10.1175/1520-0485\(1992\)022<0609:VAVCAA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1992)022<0609:VAVCAA>2.0.CO;2), 1992
- Polo, I., Lazar, A., Rodriguez-Fonseca, B., and Arnault, S.: Oceanic Kelvin waves and tropical Atlantic intraseasonal variability: 1. Kelvin wave characterization. *Journal of Geophysical Research: Oceans*, 113(C7), <https://doi.org/10.1029/2007JC004495>, 2008
- Qiu, B. and Huang, R.X.: Ventilation of the North Atlantic and North Pacific: subduction versus obduction. *Journal of Physical Oceanography*, 870 25(10), pp.2374-2390, 1995
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily high-resolution blended analyses for sea surface temperature. *Journal of climate* 20(22):5473–5496, <https://doi.org/10.1175/2007JCLI1824.1>, 2007
- Robinson, A. and Stommel, H.: The oceanic thermocline and the associated thermohaline circulation 1. *Tellus* 11(3):295–308, <https://doi.org/10.1111/j.2153-3490.1959.tb00035.x>, 1959
- 875 Roch, M., Brandt, P., Schmidtko, S., and Tuchen, F. P.: Impact of the North Atlantic Oscillation on the decadal variability of the upper subtropical-tropical Atlantic Ocean. *Journal of Geophysical Research: Oceans*, 129(4), 2024
- Rodriguez, J., Tintore, J., Allen, J. T., Blanco, J. M., Gomis, D., Reul, A., Ruiz, J., Rodriguez, V., Echevarria, F., and Jimenez-Gomez, F.: Mesoscale vertical motion and the size structure of phytoplankton in the ocean. *Nature* 410(6826):360–363, <https://doi.org/10.1038/35066560>, 2001
- 880 Roemmich, D.: The balance of geostrophic and ekman transports in the tropical atlantic ocean. *Journal of Physical Oceanography* 13(8):1534–1539, [https://doi.org/10.1175/1520-0485\(1983\)013<1534:TBOGAE>2.0.CO;2](https://doi.org/10.1175/1520-0485(1983)013<1534:TBOGAE>2.0.CO;2), 1983
- Rossby, T.: Visualizing and quantifying oceanic motion. *Annual review of marine science* 8:35–57, <https://doi.org/10.1146/annurev-marine-122414-033849>, 2016
- Rudnick, D. L.: Intensive surveys of the azores front: 2. inferring the geostrophic and vertical velocity fields. *Journal of Geophysical Research: Oceans* 101(C7):16,291–16,303, <https://doi.org/10.1029/96JC01144>, 1996
- 885

- Ruiz, S., Pelegri, J. L., Emelianov, M., Pascual, A., and Mason, E.: Geostrophic and ageostrophic circulation of a shallow anticyclonic eddy off cape bojador. *Journal of Geophysical Research: Oceans* 119(2):1257–1270, <https://doi.org/10.1002/2013JC009169>, 2014
- ~~Ruiz S., Claret M., Pascual A., Olita A., Troupin C., Capet A., Tovar-Sanchez A., Allen J., Poulain P.-M., Tintore J., et al.: Effects of oceanic mesoscale and submesoscale frontal processes on the vertical transport of phytoplankton. *Journal of Geophysical Research: Oceans* 124(8):5999–6014, <https://doi.org/10.1029/2019JC015034>, 2019~~
- Salmon, R.: The shape of the main thermocline. *Journal of Physical Oceanography*, 12(12), 1458-1479, [https://doi.org/10.1175/1520-0485\(1982\)012<1458:TSOTMT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1982)012<1458:TSOTMT>2.0.CO;2), 1982
- Schoonover, J., Dewar, W., Wienders, N., Gula, J., McWilliams, J. C., Molemaker, M. J., Bates, S. C., Danabasoglu, G., and Yeager, S.: North atlantic barotropic vorticity balances in numerical models. *Journal of Physical Oceanography* 46(1):289–303, <https://doi.org/10.1175/JPO-D-15-0133.1>, 2016
- Schott, F. A. and Stommel, H.: Beta spirals and absolute velocities in different oceans. *Deep Sea Research* 25(11):961–1010, [https://doi.org/10.1016/0146-6291\(78\)90583-0](https://doi.org/10.1016/0146-6291(78)90583-0), 1978
- Sevellec, F., ~~Garabato A Naveira Garabato, A. C. N.~~, Brearley, J., and Sheen, K.: Vertical flow in the southern ocean estimated from individual moorings. *Journal of Physical Oceanography* 45(9):2209–2220, <https://doi.org/10.1175/JPO-D-14-0065.1>, 2015
- Sonneveld, M., Wunsch, C., and Heimbach, P.: Unsupervised learning reveals geography of global ocean dynamical regions. *Earth and Space Science* 6(5):784–794, <https://doi.org/10.1029/2018EA000519>, 2019
- Song, T., Rossby, T., and Carter, E.: Lagrangian studies of fluid exchange between the gulf stream and surrounding waters. *Journal of physical oceanography* 25(1):46–63, [https://doi.org/10.1175/1520-0485\(1995\)025<0046:LSOFEB>2.0.CO;2](https://doi.org/10.1175/1520-0485(1995)025<0046:LSOFEB>2.0.CO;2), 1995
- Stommel, H. and Arons, A.: On the abyssal circulation of the world ocean. stationary planetary flow patterns on a sphere. *Deep Sea Research* (1953) 6:140–154, [https://doi.org/10.1016/0146-6313\(59\)90065-6](https://doi.org/10.1016/0146-6313(59)90065-6), 1959
- Stommel, H. and Schott, F.: The beta spiral and the determination of the absolute velocity field from hydrographic station data. *Deep Sea Research* 24(3):325–329, [https://doi.org/10.1016/0146-6291\(77\)93000-4](https://doi.org/10.1016/0146-6291(77)93000-4), 1977
- Sverdrup, H. U.: Wind-driven currents in a baroclinic ocean; with application to the equatorial currents of the eastern pacific. *Proceedings of the National Academy of Sciences* 33(11):318–326, <https://doi.org/10.1073/pnas.33.11.318>, 1947
- Tailleux, R.: On the determination of the 3d velocity field in terms of conserved variables in a compressible ocean. *Fluids* 8(3):94, <https://doi.org/10.3390/fluids8030094>, 2023
- Tintore, J., Gomis, D., Alonso, S., Parrilla, G.: Mesoscale dynamics and vertical motion in the alboran sea. *Journal of Physical Oceanography* 21(6):811–823, [https://doi.org/10.1175/1520-0485\(1991\)021<0811:MDAVMI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1991)021<0811:MDAVMI>2.0.CO;2), 1991
- Thomas, M. D., De Boer, A. M., Johnson, H. L., and Stevens, D. P.: Spatial and temporal scales of sverdrup balance. *Journal of Physical Oceanography* 44(10):2644–2660, <https://doi.org/10.1175/JPO-D-13-0192.1>, 2014
- ~~Tzortzis R., Doglioli A. M., Barrillon S., Petrenko A. A., d’Ovidio F., Izard L., Thyssen M., Pascual A., Barcelo-Llull B., Cyr F., et al.: Impact of moderately energetic fine-scale dynamics on the phytoplankton community structure in the western mediterranean sea. *Biogeosciences* 18(24):6455–6477, <https://doi.org/10.5194/bg-18-6455-2021>, 2021~~
- Uchida, T., Balwada, D., Abernathey, R., McKinley, G., Smith, S., ~~Levy and Levy~~, M.: The contribution of submesoscale over mesoscale eddy iron transport in the open southern ocean. *Journal of Advances in Modeling Earth Systems* 11(12):3934–3958, <https://doi.org/10.1029/2019MS001805>, 2019

- Verezemskaya, P., Barnier, B., Gulev, S. K., Gladyshev, S., Molines, J.-M., Gladyshev, V., Lellouche, J.-M., [Gavrikov and Gavrikov, A.](#): Assessing eddying (1/12) ocean reanalysis glorys12 using the 14-yr instrumental record from 59.5 n section in the atlantic. *Journal of Geophysical Research: Oceans* 126(6), <https://doi.org/10.1029/2020JC016317>, 2021
- 925 Waldman, R. and Giordani, H.: Ocean barotropic vorticity balances: theory and application to numerical models. *Journal of Advances in Modeling Earth Systems* 15(4), <https://doi.org/10.1029/2022MS003276>, 2023
- Weingartner, T. J. and Weisberg, R. H.: On the annual cycle of equatorial upwelling in the central Atlantic ocean. *Journal of Physical Oceanography* 21(1):68–82, [https://doi.org/10.1175/1520-0485\(1991\)021<0068:OTACOE>2.0.CO;2](https://doi.org/10.1175/1520-0485(1991)021<0068:OTACOE>2.0.CO;2), 1991
- Williams, R.G., Roussenov, V. and Follows, M.J.: Nutrient streams and their induction into the mixed layer. *Global Biogeochemical Cycles*, 930 20(1), <https://doi.org/10.1029/2005GB002586>, 2006
- Wunsch, C.: An estimate of the upwelling rate in the equatorial atlantic based on the distribution of bomb radiocarbon and quasi-geostrophic dynamics. *Journal of Geophysical Research: Oceans* 89(C5):7971–7978, <https://doi.org/10.1029/JC089iC05p07971>, 1984
- Wunsch, C.: The past and future ocean circulation from a contemporary perspective. *Ocean Circulation: Mechanisms and Impacts—Past and Future Changes of Meridional Overturning*, 173, 53-74, <https://doi.org/10.1029/173GM06>, 2007
- 935 Wyrtki, K.: The thermohaline circulation in relation to the general circulation in the oceans. *Deep Sea Research (1953)* 8(1):39–64, [https://doi.org/10.1016/0146-6313\(61\)90014-4](https://doi.org/10.1016/0146-6313(61)90014-4), 1961
- Wyrtki, K.: An estimate of equatorial upwelling in the pacific. *Journal of Physical Oceanography* 11(9):1205–1214, [https://doi.org/10.1175/1520-0485\(1981\)011<1205:AEOEUI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1981)011<1205:AEOEUI>2.0.CO;2), 1981
- Yang, P., Jing, Z., Sun, B., Wu, L., Qiu, B., Chang, P., [Ramachandran and Ramachandran, S.](#): On the upper-ocean vertical eddy heat transport 940 in the kuroshio extension. part i: Variability and dynamics. *Journal of Physical Oceanography* 51(1):229–246, <https://doi.org/10.1175/JPO-D-20-0068.1>, 2021
- Yeager, S.: Topographic coupling of the atlantic overturning and gyre circulations. *Journal of Physical Oceanography* 45(5):1258–1284, <https://doi.org/10.1175/JPO-D-14-0100.1>, 2015
- Zhao, J., and Johns, W.: Wind-forced interannual variability of the Atlantic Meridional Overturning Circulation at 26.5 N. *Journal of Geophysical Research: Oceans*, 119(4), 2403-2419, <https://doi.org/10.1002/2013JC009407>, 2014 945