

# OLIGOTREND, a global database of multi-decadal chlorophyll-a and water quality timeseries time series for rivers, lakes and estuaries

Camille Minaudo<sup>1,2</sup>, Andras Abonyi<sup>3,4,5</sup>, Carles Alcaraz<sup>6</sup>, Jacob Diamond<sup>7</sup>, Nicholas J.K. Howden<sup>8,9</sup>, Michael Rode<sup>10,11</sup>, Estela Romero<sup>12,1</sup>, Vincent Thieu<sup>13</sup>, Fred Worrall<sup>14</sup>, Qian Zhang<sup>15</sup>, Xavier Benito<sup>6,1</sup>

<sup>1</sup>Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona (UB), Diagonal 643, 08028 Barcelona, Spain

<sup>2</sup>Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Diagonal 643, 08028 Barcelona, Spain

<sup>3</sup>MTA-ÖK Lendület Fluvial Ecology Research Group, Karolina Street 29, H-1113 Budapest, Hungary

<sup>4</sup>HUN-REN Centre for Ecological Research, Institute of Aquatic Ecology, Karolina Street 29, H-1113 Budapest, Hungary

<sup>5</sup>WasserCluster Lunz – Biologische Station GmbH, Dr. Carl Kupelwieser Promenade 5, A-3293 Lunz am See, Austria

<sup>6</sup>Marine and Continental Waters Program, Institute of Agrifood Research and Technology (IRTA), 43540 La Ràpita, Catalonia, Spain

<sup>7</sup>Intergovernmental Hydrological Programme, UNESCO, Paris, France, 7 Place de Fontenoy 75015

<sup>8</sup>School of Civil, Aerospace and Design Engineering, University of Bristol, Bristol, BS8 1TR, UK

<sup>9</sup>Cabot Institute, University of Bristol, Bristol, BS5 9LT, UK

<sup>10</sup>Department of Aquatic Ecosystem Analysis and Management, Helmholtz Centre for Environmental Research - UFZ, Magdeburg 39104, Germany

<sup>11</sup>Institute of Environmental Science and Geography, University of Potsdam, Potsdam 14476, Germany

<sup>12</sup>Global Ecology Unit, Centre for Ecological Research and Forestry Applications (CREAF), Campus UAB, Bellaterra, Spain

<sup>13</sup>Sorbonne Université, CNRS, EPHE, UMR 7619 METIS, 4 place Jussieu, Box 105, 75005, Paris, France

<sup>14</sup>Department of Earth Sciences, University of Durham, Durham, UK

<sup>15</sup>University of Maryland Center for Environmental Science. U.S. Environmental Protection Agency Chesapeake Bay Program, 1750 Forest Drive, Suite 130, Annapolis, MD 21401, United States

*Correspondence to:* Camille Minaudo ([camille.minaudo@ub.edu](mailto:camille.minaudo@ub.edu))

**Abstract.** Reversed eutrophication, called oligotrophication, has widely been documented globally over the last 30 years in rivers, lakes, and estuaries. However, the absence of a comprehensive and harmonized dataset has hindered a deeper understanding of its ecological consequences. To address this data gap, we developed the OLIGOTREND database, which contains multi-decadal timeseries time series of chlorophyll-a, nutrients (nitrogen and phosphorus), and related physicochemical parameters, totalling 4.3 million observations. These data originate from 1,894 unique monitoring locations across estuaries (n = 238), lakes (687), and rivers (969). Most time series covered the period 1986–2022 and comprised at least 15 years of chlorophyll-a observations. Each location is associated to catchment and hydroclimatic attributes. Trend and breakpoint analyses were applied to all timeseries time series. Chlorophyll-a showed temporally variable and ecosystem-specific responses to nutrient declines with an overall declining trend for 18% of the time series, contrasting greatly with a majority of declining trends for nutrient concentrations. We harmonized the database to ensure reproducibility, ease of access, and support future updates and contributions. Available at <https://doi.org/10.6073/pasta/a7ad060a4dbc4e7dfcb763a794506524> (Minaudo & Benito, 2024) the OLIGOTREND database supports collaborative efforts aimed at further advancing our understanding of

biogeochemical and biological mechanisms underlining oligotrophication, and ecological impacts of global long-term environmental change.

**Short summary.** Many waterbodies undergo nutrient decline globally, called oligotrophication, but a comprehensive dataset to understand ecosystem responses is lacking. The OLIGOTREND database comprises multi-decadal chlorophyll-*a* and nutrient ~~timeseries~~*time series* from rivers, lakes, and estuaries with 4.3 million observations from 1,894 unique measurement locations. The database provides empirical evidence for oligotrophication responses with a spatial and temporal coverage exceeding previous efforts.

## Introduction

Decades of freshwater and estuarine eutrophication in the 20th century spurred coordinated national efforts to reduce aquatic nutrient loads and subsequent algal blooms (Pinay et al., 2017). The most effective actions have included improved wastewater collection and treatment, better coordinated watershed management, and the regulation of phosphorus in detergents (Conley et al., 2009; Némery & Garnier, 2016). Evidence from rivers, lakes, and estuaries already suggests that such efforts can indeed reverse eutrophication at time scales ranging from months to years and decades, in a process termed oligotrophication or re-oligotrophication. However, our understanding of oligotrophication is still ~~\_not fully understood~~*incomplete* (Anneville et al., 2019; Hoyer et al., 2002; Ibáñez & Peñuelas, 2019), and the magnitude, direction, and timing of ecological responses to water quality improvements remain to be better detected and quantified. Declines in nutrients often coincide with a transition in primary producers in terms of quantity and community composition. The most reported change in inland and estuarine ecosystems is the systematic replacement of phytoplankton by submerged macrophytes (Ibáñez & Peñuelas, 2019). However, these shifts can follow nonlinear trajectories, typically explained by the occurrence of alternative stable states in lakes (Scheffer & Carpenter, 2003), rivers (Verdonschot et al., 2013), and estuaries (Duarte et al., 2009; Elliott & Quintino, 2007). Additional complexities in predicting primary producer shifts arise due to nutrient legacies in the landscape that can create lags in ecosystem response (Stackpoole et al., 2019; Van Meter et al., 2021), and the presence of dams and weirs that alter the spatiotemporal variability of nutrient mobilization and transport (Zeng et al., 2023). Indeed, a wide range of contrasting trends in nutrients and primary production (as indicated by chlorophyll-*a* [*chl**a*]) are possible (Greening & Janicki, 2006; Kronvang et al., 2005; Murphy et al., 2022), including natural causes such as forest growth (Nilsson et al., 2024). Due to the complexity of ecosystem responses to watershed nutrient reduction, a common predictive framework remains elusive, highlighting the need for ~~a~~*cross*-ecosystem analysis of oligotrophication trends.

Available water quality datasets, while plentiful, remain heterogeneous and often irregularly collected and reported, hindering their use in across-system studies. Moreover, oligotrophication has been primarily focused on local and regional-scale studies (e.g. Abonyi et al., 2018; Greening et al., 2014; Minaudo et al., 2021; Sabel et al., 2020) and isolated aquatic ecosystems. Thus, the spatial extent of oligotrophication trends remain poorly constrained, and we lack an understanding of the connectivity of

oligotrophication responses across the watershed to estuary continuum. Even the best available harmonized, large-scale water quality databases commonly exclude *chla* (e.g., GRQA, Virro et al., 2021), limiting their utility to evaluate oligotrophication. Likewise, some databases may cover large numbers of observations, but exclude parallel measurements of *chla* and nutrients, mainly phosphorus (Nilsson et al., 2024; Spaulding et al., 2024) or are temporally limited relative to oligotrophication timescales (Brehob et al., 2024). Therefore, there is a clear need for a centralized database of paired nutrient and primary producer observations at oligotrophication-relevant timescales across different ecosystems.

Here we present OLIGOTREND (Minaudo & Benito, 2024), a database of 4.3 million quality assessed public and open access observations of water quality variables and *chla* from rivers, lakes and reservoirs, estuaries and coastal bays, enabling the joint assessment of multi-decadal oligotrophication trends across spatial scales. We collected and harmonized multi-decadal [timeseriesime series](#) to facilitate its structure and reuse. The database also covers geo-spatial data, including catchment and waterbody attributes, climate variables, and a robust trend analysis of all water quality [timeseriesime series](#). Here we highlight some of the main findings from our first analyses of the database and describe possible research directions that OLIGOTREND holds the potential to answer.


## 2. Data and Methods

We followed a transparent and reproducible approach to produce the OLIGOTREND database, in line with best practices for Open Science in Ecology (Powers & Hampton, 2019). In particular, the entire data processing pipeline (Figure 1) was developed collaboratively in a version control GitLab repository (<https://gitlab.com/OLIGOTREND/wp1-unify>). Data are referenced according to their level (“L”) in the processing pipeline. [Timeseriesime series](#) extracted from various sources were defined as “L0a”, preserving the original data structure and formatting. [Timeseriesime series](#) were then harmonized (“L0b”), and a selection of variables of interest (see Section 2.1) at sampling sites with at least 15 years of *chla* data qualified for the data quality assessment and check (QA/QC, see Section 2.2) and to be matched with geospatial data (see Section 2.3). Harmonized and curated [timeseriesime series](#) together with catchment and waterbody attributes constitute “L1” data, i.e., analysis- and sharing-ready data. Any additional processing of L1 data, e.g. trend analyses, was considered as “L2” (see Section 2.4).

### 2.1. Data collection

In-situ *chla* concentrations and physicochemical parameters were extracted from open-source international, national, and regional water quality databases (Table 1). We first obtained data from queries to the Earth System Science Data portal (<https://www.earth-system-science-data.net/>), the Environmental Data Initiative repository (<https://edirepository.org/>), and the Scientific Data portal (<https://www-nature-com.sire.ub.edu/sdata/>). We then conducted a literature search on Web of Science (<https://www.webofscience.com/wos/>) and Scopus (<https://www.scopus.com/>) for further existing long-term *chla* and nutrient [timeseriesime series](#). To do so, we used the following search terms: “TITLE or ABSTRACT (oligotrophication,

reoligotrophication, chlorophyll, timeseries); and in TITLE or ABSTRACT (lake, river, estuary, coastal, estuarine)-; and in EVERYTHING (trend, long term, multi-decadal)”, and ~~W~~-directly extracted the datasets when public and accessible. we directly extracted the datasets and proceeded with data harmonization. The database architecture (Figure 1) allows researchers to easily complement it with additional ~~timeseries~~time series in the future. New additions to the database will be eased by a set of scripts available in a dedicated version control GitLab repository (<https://gitlab.com/OLIGOTREND/wp1-unify>) allowing to reproduce, update or add more timeseries from level L0a to higher data levels and products.



**Figure 1.** Data levels and procedure followed to produce the OLIGOTREND database, an ensemble of harmonized and curated timeseries time series of *chl a* and water quality paired with catchment and waterbody attributes. QA/QC stands for quality assessment and quality check.

We gathered data as raw measurements, i.e. unprocessed or non-aggregated [timeseries](#)[time series](#), and defined herein this data as level L0a. Extracted variables included chlorophyll-a (*chl*a), water temperature (*wtemp*), conductivity (*cond*), pH, dissolved oxygen as concentration (*o2*) and percentage of saturation (*o2sat*), dissolved inorganic nitrogen (*din*), nitrate (*no3*), nitrate + nitrite (*no23*), ammonium nitrogen (*nh4*), Kjeldahl nitrogen (*nkjehl*), total nitrogen (*tn*), orthophosphate or soluble reactive phosphorus (*po4*), total phosphorus (*tp*), dissolved organic carbon (*doc*), and total suspended solids (*tss*). The ecosystem types covered in this database included lakes and reservoirs, rivers, estuaries and coastal bays.

We primarily targeted databases identified with long periods of records without any filter on geographic location (Table 1). We discarded *chla* datasets obtained with remote sensing techniques, to ensure a strict comparability among observations. For stratifying deep lakes, we extracted values either for the euphotic layer, or from the upper 10 m if euphotic depth was unavailable, to avoid using data from light-limited conditions.

Table 1. Data sources of the OLIGOTREND database.

Source	Link to data (and date of extraction when appropriate)	Spatial coverage
naiades French water quality portal	<a href="https://naiades.eaufrance.fr/">https://naiades.eaufrance.fr/</a> (last accessed 07/05/2024)	French national territory
Naderian et al., 2024	<a href="https://doi.org/10.1016/j.resconrec.2023.107401">https://doi.org/10.1016/j.resconrec.2023.107401</a>	Global
Chesapeake Bay Program	<a href="https://www.chesapeakebay.net/what/downloads/cbp-water-quality-database-1984-present">https://www.chesapeakebay.net/what/downloads/cbp-water-quality-database-1984-present</a> (last accessed 30/01/2024)	Chesapeake Bay and watershed
LAGOS-NE	<a href="https://doi.org/10.1093/gigascience/gix101">https://doi.org/10.1093/gigascience/gix101</a>	North-East USA
UK Harmonized Monitoring Dataset	<a href="https://datamap.gov.wales/documents/2633">https://datamap.gov.wales/documents/2633</a> (last accessed 17/06/2024)	England and Wales
Lake PCI	<a href="https://doi.org/10.20383/102.0488">https://doi.org/10.20383/102.0488</a>	Temperate and cold northern lakes
Danish monitoring program	<a href="https://odaforalle.au.dk/login.aspx">https://odaforalle.au.dk/login.aspx</a> (last accessed 14/06/2024)	Denmark
Sacramento Bay Interagency monitoring	<a href="https://doi.org/10.6073/pasta/f58f8217c18f469e7fd565997a47813c">https://doi.org/10.6073/pasta/f58f8217c18f469e7fd565997a47813c</a>	Sacramento-San Joaquin Delta (USA)
Elbe monitoring program	<a href="https://www.fgg-elbe.de/fachinformationssystem.html">https://www.fgg-elbe.de/fachinformationssystem.html</a> (last accessed 12/12/2023)	Elbe River watershed and estuary (Germany)
Filazzola et al., 2020	<a href="https://doi.org/10.1038/s41597-020-00648-2">https://doi.org/10.1038/s41597-020-00648-2</a>	Global
USGS-NWIS Data Retrieval	<a href="https://doi.org/10.5066/P9X4L3GE">https://doi.org/10.5066/P9X4L3GE</a> (last accessed 19/12/2023)	USA
GEMStat	<a href="https://gemstat.org/">https://gemstat.org/</a> (last accessed 11/06/2024)	Global
LTER Florida Everglades	<a href="https://doi.org/10.6073/pasta/f45fbf88dcf1f78f0d74c1dbdaaa8c7d">https://doi.org/10.6073/pasta/f45fbf88dcf1f78f0d74c1dbdaaa8c7d</a>	Florida Everglades (USA)
Danube River public program (HUN-REN CER, IAE)	<a href="https://doi.org/10.1111/fwb.13084">https://doi.org/10.1111/fwb.13084</a>	Middle section of the Danube River (N-Budapest, Hungary)
Victoria State Government	<a href="http://www.data.water.vic.gov.au/">http://www.data.water.vic.gov.au/</a> (last accessed 17/05/2024)	Victoria State (Australia)
Commission pour la Protection des Eaux du Léman (CIPEL)	<a href="https://www.cipel.org/en/">https://www.cipel.org/en/</a> (last accessed 03/02/2023)	Lake Geneva, France-Switzerland
Ebro River monitoring program	<a href="https://doi.org/10.1016/j.scitotenv.2011.11.059">https://doi.org/10.1016/j.scitotenv.2011.11.059</a>	Ebro River at Tortosa (Spain)
Romero et al., 2013	<a href="https://doi.org/10.1007/s10533-012-9778-0">https://doi.org/10.1007/s10533-012-9778-0</a>	Southwestern Europe

2.2. Data harmonization and quality control

First, L0a *timeseriesime series* were individually reformatted into standard units and data matrix headers, forming an ensemble of *timeseriesime series* defined here as level L0b. Nutrient concentrations were expressed as mg L<sup>-1</sup> except *chl*a, which remained in µg L<sup>-1</sup>. *Timeseriesime series* were named with a unique identifier (*uniquID*) per site corresponding to the concatenation of the following data separated by underscores: “ecosystem type”, “basin”, “station ID”, e.g., “river\_loire\_04000100”. Basin names were derived from site geographic coordinates and the corresponding watershed according to the FAO dataset (Food and Agriculture Organization of the United Nations & FAO Land and Water Division, 2024). Ecosystem type was either “estuary”, “lake” or “river”, corresponding to estuary or coastal bay, lake or reservoir, and river, respectively. The “station ID” was the one provided by the original data source. For each sampling site, the geographic

coordinates found in the original metadata ~~was-were~~ used to create a point shapefile labelled with the station unique identifier (*uniqueID*) as explained above. Stations with no geographic coordinates were discarded from the database.

Data quality was assessed and checked for all L0b ~~timeseriesime series~~ from sampling stations presenting at least 15 years of *chl*a data. The resulting dataset comprises the OLIGOTREND L1 data level (Figure 1). ~~We did not remove any data in response to data curation (QA/QC) to allow users to design their own quality check procedure. To offer the possibility to OLIGOTREND users to design their own quality check procedure, we did not remove any data in response to data curation (QA/QC).~~ Instead,

we flagged potentially anomalous or suspicious observations. Valid observations were indicated with flag = 0. Quality control identified missing values (flag = 1), possible outliers (flag = 2), and abnormally repetitive values (flag = 3). Observations were considered as outliers when the corresponding values exceeded 3 times the interquartile range defined by site. Observations were considered abnormally repetitive when, at a given site and for a given variable, the corresponding value appeared more than 5 % of the time in the ~~timeseriesime series~~, not necessarily consecutively. Obvious mistakes in the units found in the original datasets at level L0b were identified ~~and corrected~~ by plotting the density of distribution of observed concentrations and scatter plots by pairs of variables (e.g., *chl*a vs *tp*, *tp* vs *po4*, ...etc.) throughout the database.

### 2.3. Link with watershed and ecosystem properties

We linked inland sampling stations with the global HydroATLAS database (Lehner et al., 2022; Linke et al., 2019). The HydroATLAS has three distinct datasets: BasinATLAS, RiverATLAS, and LakeATLAS which represent sub-basin delineations (polygons), the river network (lines), and lake shorelines (polygons), respectively. ~~Although we proceeded with the spatial join between HydroATLAS and OLIGOTREND stations, we acknowledge there may be a potential temporal mismatch between HydroATLAS properties and OLIGOTREND temporal coverage. Yet, we considered this spatial join would succeed at demonstrating the great variability of watershed and ecosystem properties encountered in the OLIGOTREND database.~~

First, we linked all OLIGOTREND sampling stations to the BasinATLAS by spatial selection of polygons of sub-basins (Pfafstetter level 12, i.e., the highest hierarchical sub-basin level in the BasinATLAS), overlapping with the point shapefile of L1 OLIGOTREND stations. A selection of watershed properties related to their physiography, climate, land cover, hydrology and anthropogenic pressures were extracted and linked to each station present in the database at the L1 level and intersecting with one of the BasinATLAS sub-basins. Similarly, the intersection of LakeATLAS lake polygons with L1 stations provided an ensemble of lake characteristics for 61% of the lake stations (418 out of 687). ~~Finally, OLIGOTREND L1 river stations were linked to the RiverATLAS database by identifying the three nearest river segments using the function joinbynearest() in QGIS 3.26.2. For each possible station-segment match, the distance between the station and each segment was calculated, and the quality of the spatial join was assessed using a flagging system: if the distance to the nearest segment exceeded 500 m, a flag (flag = 1) was raised, indicating that the distance might be too large for the join to be considered valid. If the distance to the second or third nearest segment was less than 10% greater than the distance to the nearest segment, a flag (flag = 2) was raised indicating that several river segments could potentially be selected. In that case, if these segments were associated with~~

multiple sub-basins (HYBAS L12 in HydroATLAS documentation), a flag value of 2.1 was set. If these segments were linked to multiple drainage basins (MAIN\_RIV in HydroRIVERS), a flag value of 2.2 was set. All other associations identified during the spatial join were considered as valid, and flag value was set to flag = 0. Only stations with flag = 0 were considered reliable. Overall, out of 924 river stations, 90% were considered as valid. We found that 6.1% of stations were more than 500 m away from the closest HydroRIVERS segment, and 3.9% shown possible multiple associations (flag ≥ 2), sometimes with different sub-basins (1.3%, flag = 2.1) or drainage basins (0.3%, flag = 2.2). We acknowledge that there is some uncertainty in the spatial join between OLIGOTREND river stations and HydroRIVERS given the spatial resolution of the HydroSHEDS (15 arc-second). This uncertainty could be reduced by using a river network derived from a higher-resolution Digital Elevation Model. Finally, OLIGOTREND L1 river stations were linked to the RiverATLAS database by finding the nearest river segment with a maximum distance of 200 m using the function *joinbynearest()* in QGIS. The identified proportion of uncertain We acknowledge an important uncertainty for this steps could be attributable given to the the spatial resolution of the HydroSHEDS (15 arc-second). However, we found a unique corresponding river stretch for 87% of river stations (844 out of 969).

Stations with unmatched basin, lake or river segment from the HydroATLAS database were not removed from the OLIGOTREND database, but we did not account for them in the statistics and description of watershed attributes.

#### 2.4. Timeseriesime series metrics and trend analysis

We described the OLIGOTREND timeseriesime series based on multiple metrics. These included the number of observations by each variable, the extent of the period of record, as well as the median, average and standard deviation of all valid values over the entire timeseriesime series.

As a first step into the trend analysis, we quantified the proportion of timeseriesime series showing lower annual averages in the second half of the timeseriesime series compared to the first one. We chose annual averages over growing season averages to increase robustness in the metric because sampling frequency was sometimes unequally distributed seasonally. This further simplified the question of how to identify the growing season among sites across latitudes. We considered that a lower average value in the 2nd half of the timeseriesime series indicated decline, regardless of the level of trend-complexity found in the timeseriesime series.

A breakpoint and segmented regression analysis was performed using the R package *segmented* (Fasola et al., 2018). Whenever the Davies test (Davies, 1987) did not identify any non-constant linear regressions in time series, we conducted a Mann-Kendall trend analysis on annual averages with the R package *trend* (Pohlert, 2023). When the Mann-Kendall test detected a monotonic trend ( $p < 0.01$ ), we calculated a Sen's slope over the complete dataset. Whenever the Davies test identified non-constant linear regressions, we fitted a segmented regression to the data with two joined segments, and the position of the temporal breakpoint and the corresponding interval estimation were identified. The Sen's slope was then quantified for both sides of the given breakpoint. For each segment, there were three possible trend types: declining, no trend, rising, noted as “-”, “0” and “+”, respectively. The combination of two joined segments or a single segment only when no breakpoint was



detected provided a total of 12 possible trend types: "-", "--", "+-", "0-", "-0", "0", "00", "+0", "-+", "0+", "+", "++". We acknowledge a segmented regression with one breakpoint unlikely captures all the variety in trend patterns, but it may provide a comprehensive first assessment for non-linear and non-monotonic temporal patterns, robust enough to provide a first overview on multi-decadal temporal trajectories. Outputs from the trend analysis and above-described statistical descriptors constitute level L2 data.

### 3. Database characteristics

#### 3.1. Timeseriesime series characteristics

We collected L0 data from 3,718 sampling stations, producing a total of 41,979 timeseriesime series. Among these, 1,894 stations had at least chla for over 15 years and were selected for quality check and harmonization at level L1 (Figure 1). Following quality check, the OLIGOTREND database includes 4.3 million observations. Across all variables and timeseriesime series, 83,807 observations (1.7 % of total observations) were flagged as outliers, and 691,000 (13.7 % of total observations) as repetitive observations. The highest proportion of abnormally repetitive observations were found for nh4 and tp (34 % and 21 % of the observations, respectively, Table 2), likely related to detection and/or quantification limits above the actual concentrations. For chla, 13 % of the observations were flagged as repetitive (9.9%) or extreme outliers (3.4%). We only included the valid data points for all subsequent analysis and timeseriesime series descriptions. Most L1 timeseriesime series were multi-decadal with a median timeseriesime series length of 33 years (Table 2).

The majority of chla timeseriesime series included 5 observations per year (Table 2); only 16 % of timeseriesime series were based on monthly sampling. We counted that 95% of chla timeseriesime series exceeded 15 years, and 75%, 43% and 11% covered 20, 30 and 40 years, respectively. The longest chla timeseriesime series covering more than 45 years originated from the LakePCI dataset (10 lake chla timeseriesime series located in Sweden), the UK Harmonized Monitoring Program (42 rivers in England and Wales), and the Sacramento Bay Interagency monitoring (13 stations in estuarine area).

**Table 2. Overview of L1a data and percentage of data points flagged as invalid for each of the main variables. Ranges are presented as “median (10<sup>th</sup> percentile– 90<sup>th</sup> percentile)”. The percentage of flagged observations (last column) correspond to possible outliers and abnormally repetitive values.**

Variable	Number of timeseriesime series	Timeseriesime series length [yr]	Number of individual years covered	Number of observations	Frequency [observations/yr-1]	% of flagged observations
chla	1885	29 (16-41)	22 (15-36)	158 (58-463)	5 (3-14)	13.3
cond	783	36 (20-43)	31 (18-42)	270 (168-527)	8 (5-13)	1.1
din	207	34 (15-35)	35 (16-36)	429 (176-588)	12 (11-17)	1.7
doc	157	23 (14-35)	22 (15-35)	267 (147-550)	11 (7-21)	2.8
nh4	916	33 (16-43)	26 (15-42)	139 (54-344)	4 (2-10)	38.1
nkjel	654	30 (15-43)	23 (12-35)	104 (31-221)	3 (1-6)	57.8


<i>no23</i>	176	22 (16-43)	20 (11-34)	188 (36-480)	7 (2-14)	18.9
<i>no3</i>	1008	34 (19-43)	30 (17-42)	245 (138-453)	8 (4-12)	4.8
<i>o2</i>	1005	35 (21-42)	33 (18-42)	302 (179-567)	10 (5-15)	0.8
<i>o2sat</i>	997	35 (21-42)	33 (18-42)	299 (182-557)	10 (5-15)	1.5
<i>ph</i>	1028	34 (17-42)	28 (16-38)	130 (64-377)	4 (2-11)	45.5
<i>po4</i>	1014	34 (19-43)	29 (17-42)	218 (87-422)	7 (3-11)	20.5
<i>tn</i>	434	32 (17-37)	24 (16-36)	262 (50-574)	10 (2-16)	1.3
<i>tp</i>	1451	32 (16-39)	26 (15-36)	167 (43-474)	6 (2-14)	23
<i>tss</i>	1027	34 (20-42)	33 (18-42)	237 (123-500)	7 (4-14)	15.8
<i>wtemp</i>	1155	35 (19-42)	33 (18-42)	305 (182-573)	10 (6-15)	0.7

Timeseriesime series duration and mean observation frequency for all other variables was generally similar to the *chla* timeseriesime series. The median period of record was 32 years for both *tp* and *tn*. Median sampling frequency was 6 and 10 observations per year for *tp* and *tn*, respectively. A small proportion (2% and 1.8%, respectively) of *tp* and *tn* timeseriesime series were shorter than 15 years. For *tp*, 84%, 57% and 9% of the timeseriesime series were longer than 20, 30 and 40 years, respectively. For *tn*, 83%, 61% and 5% of the timeseriesime series were longer than 20, 30 and 40 years, respectively. There were 444 stations with joint *chla*, N and P observations for over 15 years. Among these, 220 corresponded to river stations, 169 to estuary stations, and 55 to lake stations.

Across all timeseries, the median temporal coverage was 1986 to 2022 (Table 3 and Figure 2). Yet, OLIGOTREND featured early and long *chla* time series with 19 of them starting before 1970 and an average of 50 year-long timeseries, most of them found in the Lake PCI dataset. Across all variables, the 2000s and 2010s are the decades with the highest coverage. The 2020s were not as covered as the 2010s were, likely indicating that databases are not systematically updated with the most recent observations.

Formatted: Font: Not Italic

Formatted: Font: Italic



**Figure 2. Temporal coverage of OLIGOTREND timeseries for all environmental variables included.** The y-axis “count” quantifies how many times the OLIGOTREND time series with valid observations in a series of valid observations cover a given year for each variable, from between 1960 and 2024. Only 35 time series started before 1960; 20 concerned *tss* and only one *chla*. Vertical red lines indicate median starting and ending years across the pooled data-set, i.e. the periods with the highest number of observations globally. Only 35 time series started before 1960, and 20 of them were *tss* observations and only one was *chla*. Red vertical lines indicate for each variable the median starting and ending period years across all the time series available for a given variable of records across ecosystems, i.e. the period with a majority of the most observations across all stations.

### 3.2. Spatial coverage

The OLIGOTREND L1 database contains 13,992 ~~timeseries~~ *time series* originating from 1,894 sampling stations spanning across 5 continents (Table 1, Figure 32). There are 238, 687, and 969 stations located in estuaries or coastal bays, lakes or

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: 9 pt, Italic

**Formatted:** Font: 9 pt, Italic

**Formatted:** Font: (Default) Times New Roman, 10 pt, Font color: Auto

reservoirs, and rivers, respectively (Table 3). The 3 largest data sources are the French national water quality monitoring (775 stations), a global database of water quality measurements in lakes (Naderian et al., 2024 — 378 stations), and the United States' Chesapeake Bay Program (199 stations).


Geographically, the L1 dataset includes stations from 33 different large watersheds (Figure 32 and see Table S1 for detailed list of these watersheds). The 5 most represented large watersheds are the Seine (France, 320 stations), the United States North Atlantic Coast (266 stations), the Mississippi-Missouri basin (231 stations), the French West Coast (183 stations), and England and Wales (163 stations). In total, 7 large watersheds contain more than 100 stations. Data from the Chesapeake Bay (United States North Atlantic Coast watershed) and the Elbe River watershed are particularly remarkable in terms of data contribution, covering hundreds of stations along the main rivers, encompassing both freshwater and estuarine zones.

260

**Table 3. Characteristics of the *timeseriesime series* constituting the OLIGOTREND database, organized by data source (see Table 1). See Table S1 in the Supplementary material for similar statistics organized by basins. For the length of *timeseriesime series*, number of observations per *timeseriesime series*, and *chla* sampling frequencies, we provide the median value, and 10th and 90th percentiles are indicated in brackets.**

Source	Median period of record	n stations (in estuary – lake – river)	Length [years]	n <sub>obs</sub> per <i>timeseriesime series</i>	Average <i>chla</i> sampling frequency [n/year]	Total number of observations
naiades French water quality portal	1988-2023	774	34	201	4	2,118,792
		(24 - 1 - 749)	(16-42)	(71-416)	(2-6)	
Naderian2024	1986-2011	378	25	120	6	106,480
		(0 - 378 - 0)	(17-35)	(37-260)	(3-11)	
Chesapeake Bay program	1985-2019	199	34	408	12	822,961
		(157 - 0 - 42)	(19-35)	(193-588)	(10-17)	
LAGOS-NE	1985-2010	140	24	85	5	56,616
		(0 - 140 - 0)	(18-32)	(35-248)	(2-12)	
UK Harmonized Monitoring Dataset	1978-2012	133	35	299	10	168,474
		(0 - 0 - 133)	(20-44)	(177-547)	(6-15)	
Lake PCI	1988-2018	95	23	246	11	93,580
		(0 - 95 - 0)	(15-49)	(116-1174)	(5-21)	
Danish monitoring program	1983-2020	56	33	165	6	75,608
		(0 - 56 - 0)	(21-42)	(33-481)	(2-15)	
Sacramento Bay Interagency monitoring	1975-2021	46	42	297	13	50,126
		(46 - 0 - 0)	(18-46)	(109-592)	(7-18)	
Elbe monitoring program	1985-2016	25	31	581	15	701,431
		(2 - 0 - 23)	(22-38)	(145-8490)	(4-20)	
Filazzola et al., 2020	2001-2018	13	17	123	3	7,852
		(0 - 13 - 0)	(16-28)	(32-387)	(1-12)	
USGS-NWIS Data Retrieval	1991-2021	10	30	682	22	7,337
		(0 - 0 - 10)	(21-31)	(512-1093)	(17-35)	
GEMStat	1980-2016	9	26	398	11	12,737
		(0 - 3 - 6)	(16-41)	(158-645)	(9-24)	
LTER Florida Everglades	1991-2008	9	17	207	11	25,027
		(9 - 0 - 0)	(16-33)	(188-366)	(10-12)	
Danube River public program (HUN-REN CER, IAE)	1979-2012	2	33	1100	32	13,032
		(0 - 0 - 2)	(33-33)	(1010-1127)	(32-32)	
Victoria State Government	1990-2024	2	34	782	39	17,536
		(0 - 0 - 2)	(26-34)	(329-1685)	(36-41)	

Commission pour la Protection des Eaux du Léman (CIPEL)	1980-2018	1 (0 - 1 - 0)	38	815 (815-815)	12	8,150
Ebro River monitoring program	1980-2004	1 (0 - 0 - 1)	24 (15-24)	284 (133-323)	18 (18-18)	2,039
Romero et al., 2013	1982-2016	1 (0 - 0 - 1)	34 (29-34)	304 (176-362)	4 (4-4)	1,684
TOTAL	1986-2022	1,894	33 (17-42)	220 (71-507)	5 (3-14)	4,281,312



**Figure 32.** a) Map highlighting the 1894 sampling stations included in the OLIGOTREND database at level L1, categorized by ecosystem types. b) close-up on the Eastern side of the US, and c) on Europe showcasing most data points from France, UK and Denmark.

The OLIGOTREND database covers 1,229 sub-basins from the HydroATLAS database, distributing over 257 spatially independent large watersheds with no hydrological connections. OLIGOTREND covers a wide range of eco-physiographic contexts (Table 4). It covers medium to large watersheds (10th to 90th percentiles were 142 to 11,416 km<sup>2</sup>), primarily lowlands.

Stations extend to four climate zones, from extremely cold and mesic to hot and dry. Share among land-use types also covers a wide range, from 100% forest or natural grassland areas to heavily impacted urban areas and croplands. Some of the stations are located in nearly pristine areas, but most of them are in highly populous areas.

Similarly, lakes and rivers represented by the OLIGOTREND database cover a wide range of morphometry, from shallow (e.g., Hickling Broad lake, England, average water column depth ~0.7 m) to deep and large lakes (e.g., Lake Geneva, France-Switzerland, average depth ~155 m), and from headwater streams (e.g., the Evel river in French Brittany draining a basin of 5 km<sup>2</sup>) to large rivers (e.g., Mississippi, Danube, Rhine, Loire, Seine, Ebro, Susquehanna Rivers).

**Table 4. Basin characteristics covered by the OLIGOTREND database based on the HydroATLAS (level 12), the HydroLAKES and HydroRIVERS databases. Column “Range” indicates median values; and percentiles 10 and 90 are shown in brackets.**

Category	Variable	Description	Aggregation	Range	Units
Physiography	up_area	Watershed area	Upstream sub-basin	573.8 (142-11,416)	km <sup>2</sup>
	ele_mt_sav	Elevation	Sub-basin	125 (28-417)	m a.s.l.
	slp_dg_uav	Terrain slope	Upstream sub-basin	25 (10-71)	degrees
Climate	tmp_dc_syr	Air temperature average	Sub-basin	10.1 (6.3-12.5)	degrees Celsius
	pre_mm_sy	Precipitation average	Sub-basin	755 (625-1,106.2)	mm
	clz_cl_smj	Climate zone <sup>(*)</sup>	Sub-basin	10 (7-11)	class
Land cover	for_pc_use	Forest cover extent	Upstream sub-basin	15 (0-90)	%
	crp_pc_use	Cropland cover extent	Upstream sub-basin	33 (4-64)	%
	pst_pc_use	Pasture cover extent	Upstream sub-basin	10 (1-36)	%
Hydrology	dis_m3_pyr	Natural discharge	Sub-basin	7.7 (1.5-131)	m <sup>3</sup> /s
	run_mm_sy	Land surface runoff	Sub-basin	376 (204-776)	mm
	lka_pc_use	Limnicity	Upstream sub-basin	2 (0-60)	%
	dor_pc_pva	Degree of regulation	Upstream sub-basin	0 (0-176)	%
Anthropogenic	pop_ct_usu	Population	Upstream sub-basin	38 (2.5-874)	inhab. (x1000)
	ppd_pk_ua	Population density	Upstream sub-basin	53.7 (11-294)	inhab./km <sup>2</sup>
	urb_pc_use	Urban cover extent	Upstream sub-basin	2 (0-15)	%
Lake characteristics	Lake_area	Lake area	Lake body	1.1 (0.2-25)	km <sup>2</sup>
	Depth_avg	Average lake depth	Lake body	5 (2.9-14.7)	m
	Res_time	Residence time	Lake body	289 (33-1394)	days
River characteristics	upland_skm	Watershed area	Upstream river segment	<del>583-629</del> (6165-12725) 13,249	km <sup>2</sup>
	dis_av_cms	Average interannual discharge	River segment pourpoint	<del>78.37</del> (0.758-142) 143	m <sup>3</sup> /s
	ord_stra	Strahler order	River segment	3 (2-5)	d.l.

<sup>\*</sup>: Climate zone classes encompass the following classes: Extremely cold and mesic, Cool temperate, Warm temperate and Hot and dry.

3.3. OLIGOTREND timeseriesime series ranges and relationships

For most variables, long-term averages are clustered by ecosystem type (Figure 43). The lowest *chl*a concentrations were found in rivers ( $7.8 \pm 10.7 \text{ ug L}^{-1}$ ) followed by estuaries ( $11.8 \pm 9.9 \text{ ug L}^{-1}$ ) and then lakes ( $18.0 \pm 25.3 \text{ ug L}^{-1}$ ). This greatly contrasted with most P, N, and oxygen timeseriesime series: for instance, *tp* and *tn* distributions showed the highest ranges in rivers ( $0.13 \pm 0.11 \text{ mg P L}^{-1}$  and  $3.1 \pm 1.8 \text{ mg N L}^{-1}$ ), and the lowest in lakes ( $0.06 \pm 0.13 \text{ mg P L}^{-1}$  and  $1.9 \pm 0.9 \text{ mg N L}^{-1}$ ). For DOC, most timeseriesime series remained within a similar range of values regardless of ecosystem type, except for four lakes located in the North-East US (global lake database; Naderian et al., 2024). The highest conductivity values appeared in estuaries, much higher than in rivers or lakes. There were only 9 lakes with conductivity timeseriesime series, explaining the density distribution peaks for this ecosystem type. The warmest waters were also found in estuaries.





Figure 43. Distribution of inter-annual average concentrations of all the OLIGOTREND timeseriesime series. Number of timeseriesime series for each variable are indicated in brackets for each variable.

Across the entire database, *chl*a annual averages showed moderate to strong correlation with *tp* and *tn* (Figure 54). *Chl*a was strongly and positively correlated with *tp* (Pearson,  $r = 0.39$ ) across all ecosystem types. The positive correlation was the strongest for lakes ( $r = 0.82$ ), moderate for rivers ( $r = 0.37$ ), and estuaries showed the weakest relationship ( $r = 0.29$ ). *Chl*a was positively correlated with *tn* (Pearson,  $r = 0.40$ ), which was the highest in lakes ( $r = 0.75$ ), moderate in rivers ( $r = 0.49$ )



and lowest in estuaries ( $r = 0.30$ ). Variables  $tp$  and  $tn$  were positively correlated across all ecosystem types (Pearson,  $r = 0.59$ ), with the strongest correlation found in lakes ( $r = 0.74$ ), slightly lower in rivers ( $r = 0.63$ ), and the weakest one in estuaries ( $r = 0.35$ ). There was a clear cluster outlier for these variables in estuaries, characterized by low  $chla$  and  $tp$  but rather high  $tn$ . These observations corresponded exclusively to the Florida Coastal Everglades.




**Figure 54.** Relationships between  $chl-a$  and  $tp$  (a),  $chl-a$  and  $tn$  (b), and  $tp$  and  $tn$  (c). Each dot represents the annual mean for a given [timeseries](#). Dark dots for estuary stations highlight the observations in the Florida Coastal Everglades which clearly stand out from all other estuarine observations. Pearson correlations are all statistically significant ( $p$ -value  $< 2e-16$ ) and corresponding coefficients ( $r$ ) are indicated in each panel.

### 3.4. Trends in the OLIGOTREND database

Comparing the mean value of annual averages between the second and the first halves of [timeseries](#) proved to be a simple but effective way to overview temporal behaviour of [timeseries](#) in the database. Across all variables and ecosystem types, 60% of [timeseries](#) showed a lower average value in the second half. 63% of  $chl-a$  [timeseries](#) showed lower values in the second half (Figure 65). For N and P nutrient [timeseries](#), 78% to 87% showed an average concentration lower in the second half (it was 85%, 87%, 78%, 85%, 86% for  $tp$ ,  $po4$ ,  $tn$ ,  $din$ ,  $nh4$ , respectively). An exception was found for  $no3$  with only 45% [timeseries](#) with a lower concentration in the second half of the [timeseries](#). Interestingly, we found that the majority (74%) of  $tss$  [timeseries](#) had a lower concentration in

the second half, whereas *o2*, *o2sat*, *pH*, and *cond* showed no clear differences in the second half of the *timeseriesime series* with 49%, 43%, 42%, and 42%. For *wtemp*, there was a clear indication of a warming trend with 64% of *timeseriesime series* with higher averages in the second half of the *timeseriesime series*.



**Figure 65.** Distribution of ratio between 2nd half *timeseriesime series* averages over 1st half averages. Values significantly below 1 likely indicate declining trends, regardless of the complexity of the temporal trajectory.

The breakpoint and trend analysis (Figure 76) revealed 15% of *chla timeseriesime series* were best represented with a segmented trend component while 62% had no trend detected, 18% presented a monotonic declining trend, 5% a monotonic rising trend (predominantly found in estuaries, see Figure 87). The predominant segmented trend types were “00” (32%), “0-” (21%), “-0” (19%) and “+” (7%).

For *tp* and *po4*, 29-31% of the *timeseriesime series* had a breakpoint with a segmented trend, 26-32% had no trend detected, while 35-42% presented a declining monotonic trend and 1-2% were rising. For *tp timeseriesime series*, 72% of segmented trends had a declining trend type, while it was 65% for *po4 timeseriesime series*. Compared to rivers and estuaries, a lower proportion of declining *tp* trends were observed in lake *timeseriesime series*.

For N species, *timeseriesime series* were dominated by the no-trend type (38-61%) and significant trends were contrasted: *tn*, *din* and *nh4* showed a large number of declining trends (36-42%) and a small proportion of rising trends (less than 2%), while *no3* and *no23* were characterized by a larger proportion of rising trends (7% for *no23* and 17% for *no3*) and segmented trends (14% for *no23* and 25% for *no3*). For *no3*, 57% of segmented trends had a declining trend type on the most recent part of the time series as 34% were “0-” and 23% were “+”. Other variables were characterized by 50-60% of no-trend time series.

Interestingly, among the detected trends, *tss* showed a significant proportion of declining trend types, while *o2*, *o2sat*, *ph* and *wtemp* showed a predominance of rising trends.





Figure 76. Overview of trend significance and trend types identified in the OLIGOTREND database. Blue stripes are indicative of declining trends, grey stripes of no-trend, and red stripes of rising trends. Empty stripes indicate variables or ecosystems where the number of time series available was lower than 30. Refer to Section 2.4 for a detailed explanation of trend symbols indicated in the legend.




**Figure 87.** Overview of all *chla* annual *timeseries* normalized by interannual averages (thin lines), organised by trend types (panels) and ecosystem type (colour). Thick grey lines are smoothed curves of all *timeseries* within a given panel, only displayed to guide the reader. Refer to Section 2.4 for a detailed explanation of trend symbols indicated on top of each panel.

For *chla*, Sen's slopes in estuaries were smaller in magnitude compared to lakes and rivers, regardless of the trend type (Figure 98a). Lakes exhibited a median Sen's slope of  $-0.7 \mu\text{g L}^{-1} \text{ year}^{-1}$ ; it was  $-0.4 \mu\text{g L}^{-1} \text{ year}^{-1}$  in rivers and  $-0.3 \mu\text{g L}^{-1} \text{ year}^{-1}$  in estuaries. The fastest declines (below  $-4 \mu\text{g L}^{-1} \text{ year}^{-1}$ ) were found in the Sacramento Bay in California, the River Loire (France), and several shallow lakes in the Mississippi-Missouri basin, the Denmark Germany Coast, and England and Wales. The largest positive *chla* trends were found in rivers, with a median slope of  $0.79 \mu\text{g L}^{-1} \text{ year}^{-1}$ , compared to  $0.13$  and  $0.23 \mu\text{g L}^{-1} \text{ year}^{-1}$  in estuaries and lakes, respectively. The fastest rises (above  $4 \mu\text{g L}^{-1} \text{ year}^{-1}$ ) were found in the River Loire (France). For *tp*, the fastest rises and declines were observed in river ecosystems (Figure 98b) with median slopes of  $4.0 \times 10^{-3}$  and  $-4.7 \times 10^{-3} \text{ mgP L}^{-1} \text{ year}^{-1}$ , respectively, one order of magnitude greater than the slopes observed in lakes and estuaries. The fastest declines (below  $-0.1 \text{ mgP L}^{-1} \text{ year}^{-1}$ ) were observed in the Rhône and Seine Rivers (France).

For *tn*, although the fastest declines were observed in estuary stations (Florida Coastal Everglades) down to  $-0.4 \text{ mgN L}^{-1} \text{ year}^{-1}$ , the median value for declining slopes was overall faster in rivers with median slopes of  $-0.14 \text{ mgN L}^{-1} \text{ year}^{-1}$  (Figure 98c). It was  $-6 \times 10^{-3} \text{ mgN L}^{-1} \text{ year}^{-1}$  in estuaries and  $-7 \times 10^{-3} \text{ mgN L}^{-1} \text{ year}^{-1}$  in lakes. Only 11 stations showed rising *tn* trends (Figure 76), and among them, 3 were in the Chesapeake Bay (US North Atlantic Coast) which contrasted with the 145 other

estuarine stations in this basin which either showed declining trends (n=89) or no trends (n=56). Note that only 7 lacustrine stations showed rising *tn* and in rivers, and none of the *tn* [timeseries](#) showed a rising pattern.



**Figure 98.** Overview of all Sen's slopes calculated for *chl a* (a), *tp* (b) and *tn* (c) whether they are showing a declining (negative values) or a rising trend (positive values). Medians by ecosystem type are indicated with a plain circle, 10th and 90th percentiles correspond to the segment ends. The numbers of [timeseries](#) found for each variable, ecosystem and trend type are indicated at the bottom or the top of each segment. See Fig. S1 in the Supplementary material for a similar figure for all variables included in OLIGOTREND.

We identified 444 stations with joint *chl a*, P and N data over 15 years and more than 6 observations per year. Among these, 100 (or 23%) *chl a* [timeseries](#) showed a linear declining trend, 251 (or 57%) had no trend, and 37 (or 8%) were rising. Declining *chl a* [timeseries](#) were also linked to declining trends in N and P (Figure 109a). Nearly half of the *chl a* [timeseries](#) with no trend had corresponding no-trend or declining patterns in nutrient [timeseries](#) (Figure 109b). Rising *chl a* [timeseries](#) predominantly corresponded to no-trend or declining patterns in nutrient [timeseries](#). Only 18% of the rising *chl a* [timeseries](#) also had significant rising trends in N or P.




Figure 109. Relative share of trend types found for nitrogen and phosphorus concentrations related to *chl a* timeseries with declining trends (a), no trends (b), and rising trends (c). This analysis is based on 444 stations having parallel measurements of *chl a*, and N (*din* and/or *no3* and/or *tn*) and P (*po4* and/or *tp*) for at least 15 years. Empty rows correspond to variables with less than 30 timeseries.

#### 4. Potential implications of OLIGOTREND for future research

The OLIGOTREND database has the potential to answer some important questions in large-scale aquatic ecology, biogeochemistry, and global change studies. Below, we highlight the most important findings of the database and discuss potential implications for future research beyond disciplinary boundaries.

##### 4.1. Unravelling the ambiguous links between *chl a* and nutrient levels for lakes, rivers, and estuaries

The development of primary producers is far more complex than a single relationship with nutrient availability, especially if one also considers the differences among ecosystem types. Hydraulic flushing, turbulence, exposition to solar radiation, temperature (e.g. Reynolds, 2006), and light climate (Hilt et al., 2011) are crucial environmental variables in lotic systems. Water residence time, internal loading (Jeppesen et al., 2005; Krishna et al., 2021), stratification regime, and underwater light climate are other crucial factors controlling lentic ecosystems (Donis et al., 2021). Such differences are also reflected in the OLIGOTREND database. For instance, on one hand rivers had the highest P and N concentrations, followed by estuaries and lakes, and on the other hand, the highest *chl a* concentrations were found in lakes followed by estuaries and then rivers (Figure 43). Further, only 18% of the *chl a* timeseries showed a linear declining trend which contrasted greatly with a dominating decreasing trend for most nutrient concentrations (Figures 65, 76 and 109). Moreover, although lake timeseries showed the highest correlation between *chl a* and nutrients (Figure 4), they were also the ones with the highest proportion of non-significant trends (Figure 76). In this context, we argue that the OLIGOTREND database provides a unique opportunity and foundation to further investigate the ambiguous links existing between *chl a* and nutrient levels over many contrasted water bodies located in basins with different environmental and climatic conditions.

#### 4.2. Is oligotrophication specific to aquatic ecosystem types?

The OLIGOTREND database evidenced different responses of the individual ecosystem types to nutrients declines (Figures 76, 87 and 109). For instance, compared to estuaries and lakes, rivers showed the highest proportion of declining *chla* (Figure 76). The inherent specificities of different ecosystems could partly explain why oligotrophication seems to be ecosystem-specific: i) the successful P reduction in many rivers worldwide (e.g., Le Moal et al., 2019) has led to more frequent P limitation for phytoplankton (Elser et al., 2007), although N or Si may also be limiting primary production (Paerl et al., 2016); ii) in lakes, longer water residence time, and internal nutrient loading can either delay (Jeppesen et al., 2005) or amplify (i.e., through algal blooms; e.g., Krishna et al., 2021) the ecological response following nutrient declines; iii) temporal shifts in phytoplankton assemblages towards taxa better adapted to low P levels, or taxa that are barely controlled by zooplankton grazing (e.g. filamentous cyanobacteria; Selmečzy et al., 2019) can often represent overlooked effects explaining rising or weak trends in primary producers despite nutrient decline over time (Anneville et al., 2019); iv) in estuaries, the dynamic of primary producers is also largely affected by marine waters, where coastal phytoplankton, sensitive to N (Elser et al., 2007), or N and P availability meets freshwater phytoplankton primarily sensitive to P (Kemp et al., 2005). Future analysis of OLIGOTREND timeseriesime series together with catchment and waterbody attributes could improve our understanding of how aquatic ecosystems respond to nutrient trends in a wide variety of aquatic ecosystems.

#### 4.3. Abrupt and gradual changes in long-term water quality timeseriesime series

The OLIGOTREND database could be explored to further evidence the extent of gradual changes or abrupt regime shifts in water quality timeseriesime series. In fact, some of the waterbodies represented in OLIGOTREND are known for shifting their primary producer's structure and function following oligotrophication. This is the case of the Loire (France) and the Ebro Rivers (Spain), which are known for their long-term gradual regime shifts from phytoplankton to macrophytes in response to phosphorus decline (Diamond et al., 2021; Ibáñez et al., 2012; Minaudo et al., 2015, 2021). Similarly, phytoplankton of the middle Danube now more frequently contains benthic taxa, predominantly diatoms, potentially indicating a long-term regime shift from pelagic to benthic production in recent decades (Abonyi et al., 2018). Moreover, oligotrophication can result in a shift from heterotrophic conditions to dominantly autotrophic processes with lower pollution, like it was observed for the Elbe River (Wachholz et al. 2024). OLIGOTREND timeseriesime series could be further analysed to detect possible temporal changes in variance (as a possible early-warning signal, Dakos et al., 2015), seasonal patterns, and relationships between *chla*, nutrients and ecosystem metabolism. This could enhance our understanding of crucial factors underlying regime shifts in river ecosystems, which are comparatively less well known than in lakes (Gilaranz et al., 2022).

In OLIGOTREND, we highlighted a significant number of no-trend or rising *chla* timeseriesime series despite declining nutrient levels (Figure 109c). This could be related to climatic effects and long-term changes of ecosystem structure, such as in the Chesapeake Bay (Harding et al., 2019). Future analysis of the OLIGOTREND will provide an invaluable source of data to disentangle the effects of climate change and watershed biogeochemistry on multi-decadal *chla* and nutrient trends.

**Commented [MR1]:** Wachholz, A., Jawitz, J.W., Borchardt, D. (2024): From Iron Curtain to green belt: shift from heterotrophic to autotrophic nitrogen retention in the Elbe River over 35 years of passive restoration *Biogeosciences* **21** (15), 3537 - 3550 <https://doi.org/10.5194/bg-21-3537-2024>

#### 430 4.4. Combining OLIGOTREND with large-scale datasets to foster interdisciplinary aquatic data science

The OLIGOTREND database can help boost water quality research if it is combined with other large-scale or long-term ecological datasets. For instance, it is known that shifting baselines because of temporal changes in different, covarying environmental factors can preclude the return of primary producer to pre-eutrophication conditions (Carstensen et al., 2011; Duarte et al., 2009). As global change intensifies, leading to novel ecosystems (Hobbs et al., 2009), the temporal extension of  
435 most available water quality datasets limits a correct estimation of pre-eutrophication baselines. Only a fraction of the OLIGOTREND database covers *chl*a and/or nutrients during the eutrophication phase, which renders pre-oligotrophication reference conditions impossible to discern; and hence, makes it difficult to validate nutrient remediation actions (Pinay et al., 2017). In this context, combining paleolimnological observations with water quality monitoring data could have a potential not fully implemented at large spatial scales and across different aquatic ecosystem types (Bennion et al., 2015; Bhattacharya  
440 et al., 2022; Dong et al., 2012).

Recent research has shown that nutrient concentrations link to nutrient loads (point and nonpoint sources) at the catchment scale (Ehrhardt et al., 2021; Jarvie et al., 2012; Murphy et al., 2022). Yet, only a few studies have established a mechanistic link between nutrient inputs management and the development of the phytoplankton biomass. Data-based approaches that jointly analyse decreasing nutrient loadings over multiple decades and sites with corresponding measurements of *chl*a and  
445 nutrients can help better characterize how successful catchment management and environmental measures can be to reverse eutrophication. OLIGOTREND holds the potential to approach oligotrophication longitudinally at the basin scale, where long-term trajectories can be assessed from small streams, rivers, lakes/reservoirs towards estuaries/coastal ecosystems along with their hydrologically connected time series.

Remote sensing could further supplement crucial water quality information organised in OLIGOTREND. Remote sensing can  
450 provide [timeseriesime series](#) data on water quality for inland and coastal aquatic ecosystems, which, if combined with in-situ measurements, can increase *chl*a data coverage both spatially and temporally (Ross et al., 2019; Spaulding et al., 2024). Moreover, regional and Earth System numerical models will improve further if calibrated or validated by in situ observations (Casquin et al., 2024; Liu et al., 2024). The OLIGOTREND database readily represents a centralized and harmonized dataset open for calibration and validation by remotely sensed water quality data, and available for training and validating regional  
455 and large-scale numerical models.

Finally, there is a growing interest in large-scale observations that integrate new and existing databases to answer key questions in aquatic ecology (Barquín et al., 2015). Long-term observations of community data (e.g. via LTER and eLTER, GBIF, Biofresh) may include key functional groups of aquatic food webs, such as phytoplankton, zooplankton, macroinvertebrates (Welti et al., 2024), and fish (Comte et al., 2021). For a selection of sites, *chl*a trends can be further analysed jointly with long-  
460 term community data to investigate the role that community composition and biodiversity may play in responding to long-term environmental change (Jochimsen et al., 2013). Some of the OLIGOTREND [timeseriesime series](#) are linked to lotic



community data (i.e., phytoplankton), which have been seldom explored compared to lakes when testing the biodiversity effect on ecosystem functioning and services (Filstrup et al., 2019; Ptacnik et al., 2008).

## 5. Code and data availability

All the data are openly available along with the R scripts used for data processing from raw measurements at L0a level to higher data processing levels. All R scripts produced to extract, harmonize and process the OLIGOTREND data were stored and organized in a dedicated GitLab repository (<https://gitlab.com/OLIGOTREND/wp1-unify>). Data at levels L1 and L2 (Figure 1) were deposited in an Environmental Data Initiative Data Package accessible on the EDI data portal (<https://doi.org/10.6073/pasta/a7ad060a4dbc4e7dfcb763a794506524>, Minaudo & Benito, 2024). Original links to data sources of L0a data are provided in Table 1 and in the EDI Data Package. Additionally, we also provide [in the GitLab repository](#) all the GIS files emerging from the data extraction step, including shapefiles of L0 and L1 stations, the corresponding basins, lakes and rivers characteristics resulting from the spatial join between OLIGOTREND stations and the HydroATLAS.

## 6. Conclusions

The OLIGOTREND database provides invaluable information in aquatic ecology and Earth system science. We evidenced oligotrophication at large temporal and spatial scales and unveiled the complexity of the chlorophyll-*a* response following oligotrophication and the relationships between chlorophyll-*a* and nutrients in inland and transitional waters covering a wide range of climatic and environmental conditions. While the database is not exhaustive, its flexible structure and reproducible processing pipeline facilitate the inclusion of additional datasets in the future. We also see a strong need to continuously update the database due to the accelerating climate change and the resulting impacts on the loading and processing of nutrients and the associated ecological implications (van Vliet et al., 2023). Finally, OLIGOTREND will support collaborative efforts aimed at advancing our understanding of the complex biogeochemical and biological mechanisms driving oligotrophication and the broader ecological impacts of global environmental change.

## Author contributions

CM and XB both secured the funding for this study and contributed equally for the Conceptualization, Methodology, Data curation, Formal analysis, Investigation. They wrote together the original draft. All other authors provided datasets and participated in the revisions of the original draft.

### Funding sources

This study was funded by the Iberian Association of Ecology (SIBECOL) through the 2022 early-career advanced grant to CM and XB. Both CM and XB have received funding from the postdoctoral fellowships programme Beatriu de Pinós, funded by the Secretary of Universities and Research (Government of Catalonia) and by the Horizon 2020 programme of research and innovation of the European Union under the Marie Skłodowska-Curie grant agreement No 801370. AA was supported by the National Research, Development and Innovation Office, Hungary (FK 142485 project) and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. ~~ER is thankful for the funding provided by the Severo Ochoa Excellence Program (CEX2018-000828-S, Spanish Research Agency) and the Spanish MCIN project KALORET (PID2021-128778OA-I00).~~ ER acknowledges the support of the Grant “Severo Ochoa Centres of Excellence” (CEX2018-000828-S) funded by MCIN/AEI/10.13039/501100011033, and the project KALORET (PID2021-128778OA-I00) funded by MCIN/AEI/10.13039/501100011033 and by “ERDF A way of making Europe”.

### References

- Abonyi, A., Ács, É., Hidas, A., Grigorszky, I., Várbíró, G., Borics, G., & Kiss, K. T. (2018). Functional diversity of phytoplankton highlights long-term gradual regime shift in the middle section of the Danube River due to global warming, human impacts and oligotrophication. *Freshwater Biology*, 63(5), 456–472. <https://doi.org/10.1111/fwb.13084>
- Anneville, O., Chang, C., Dur, G., Souissi, S., Rimet, F., & Hsieh, C. (2019). The paradox of re-oligotrophication: the role of bottom-up versus top-down controls on the phytoplankton community. *Oikos*, 128(11), 1666–1677. <https://doi.org/10.1111/oik.06399>
- Barquín, J., Benda, L. E., Villa, F., Brown, L. E., Bonada, N., Vieites, D. R., Battin, T. J., Olden, J. D., Hughes, S. J., Gray, C., & Woodward, G. (2015). Coupling virtual watersheds with ecosystem services assessment: a 21st century platform to support river research and management. *Wiley Interdisciplinary Reviews: Water*, 2(6), 609–621. <https://doi.org/10.1002/WAT2.1106>
- Bennion, H., Simpson, G. L., & Goldsmith, B. J. (2015). Assessing degradation and recovery pathways in lakes impacted by eutrophication using the sediment record. *Frontiers in Ecology and Evolution*, 3(AUG), 148732. <https://doi.org/10.3389/FEVO.2015.00094/BIBTEX>
- Bhattacharya, R., Lin, S. G. M., & Basu, N. B. (2022). Windows into the past: lake sediment phosphorus trajectories act as integrated archives of watershed disturbance legacies over centennial scales. *Environmental Research Letters*, 17(3), 34005. <https://doi.org/10.1088/1748-9326/ac4cf3>
- Brehob, M. M., Pennino, M. J., Handler, A. M., Compton, J. E., Lee, S. S., & Sabo, R. D. (2024). Estimates of Lake Nitrogen, Phosphorus, and Chlorophyll- a Concentrations to Characterize Harmful Algal Bloom Risk Across the United States. *Earth's Future*, 12(8), e2024EF004493. <https://doi.org/10.1029/2024EF004493>

- Carstensen, J., Sánchez-Camacho, M., Duarte, C. M., Krause-Jensen, D., & Marbà, N. (2011). Connecting the Dots: Responses  
520 of Coastal Ecosystems to Changing Nutrient Concentrations. *Environmental Science & Technology*, 45(21), 9122.  
<https://doi.org/10.1021/es202351y>
- Casquin, A., Silvestre, M., & Thieu, V. (2024). *nuts-STEauRY dataset: hydrochemical and catchment characteristics dataset  
for large sample studies of Carbon, Nitrogen, Phosphorus and Silicon in french watercourses*.  
<https://doi.org/10.5281/ZENODO.10830852>
- 525 Comte, L., Carvajal-Quintero, J., Tedesco, P. A., Giam, X., Brose, U., Erős, T., Filipe, A. F., Fortin, M., Irving, K., Jacquet,  
C., Larsen, S., Sharma, S., Ruhi, A., Becker, F. G., Casatti, L., Castaldelli, G., Dala-Corte, R. B., Davenport, S. R.,  
Franssen, N. R., ... Olden, J. D. (2021). RivFishTIME: A global database of fish time-series to study global change  
ecology in riverine systems. *Global Ecology and Biogeography*, 30(1), 38–50. <https://doi.org/10.1111/geb.13210>
- Conley, D. J., Paerl, H. W., Howarth, R. W., Boesch, D. F., Seitzinger, S. P., Havens, K. E., Lancelot, C., & Likens, G. E.  
530 (2009). Controlling Eutrophication: Nitrogen and Phosphorus. *Science*, 323(5917), 1014–1015.  
<https://doi.org/10.1126/science.1167755>
- Dakos, V., Carpenter, S. R., van Nes, E. H., & Scheffer, M. (2015). Resilience indicators: prospects and limitations for early  
warnings of regime shifts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1659), 1–10.  
<https://doi.org/10.1098/RSTB.2013.0263>
- 535 Davies, R. B. (1987). Hypothesis testing when a nuisance parameter is present only under the alternative. *Biometrika*, 74(1),  
33–43. <https://doi.org/10.1093/biomet/74.1.33>
- Diamond, J. S., Moatar, F., Cohen, M. J., Poirel, A., Martinet, C., Maire, A., & Pinay, G. (2021). Metabolic regime shifts and  
ecosystem state changes are decoupled in a large river. *Limnology and Oceanography*, n/a(n/a).  
<https://doi.org/https://doi.org/10.1002/lno.11789>
- 540 Dong, X., Bennion, H., Maberly, S. C., Sayer, C. D., Simpson, G. L., & Battarbee, R. W. (2012). Nutrients exert a stronger  
control than climate on recent diatom communities in Esthwaite Water: evidence from monitoring and  
palaeolimnological records. *Freshwater Biology*, 57(10), 2044–2056. <https://doi.org/10.1111/J.1365-2427.2011.02670.X>
- Donis, D., Mantzouki, E., McGinnis, D. F., Vachon, D., Gallego, I., Grossart, H., Senerpont Domis, L. N., Teurlincx, S.,  
545 Seelen, L., Lüring, M., Verstijnen, Y., Maliaka, V., Fonvielle, J., Visser, P. M., Verspagen, J., Herk, M., Antoniou, M.  
G., Tsiarta, N., McCarthy, V., ... Ibelings, B. W. (2021). Stratification strength and light climate explain variation in  
chlorophyll <scp> a </scp> at the continental scale in a European multilake survey in a heatwave summer. *Limnology  
and Oceanography*, 1–20. <https://doi.org/10.1002/lno.11963>
- Duarte, C. M., Conley, D. J., Carstensen, J., & Sánchez-Camacho, M. (2009). Return to Neverland: Shifting Baselines Affect  
550 Eutrophication Restoration Targets. *Estuaries and Coasts*, 32(1), 29–36. <https://doi.org/10.1007/s12237-008-9111-2>

- Ehrhardt, S., Ebeling, P., Dupas, R., Kumar, R., Fleckenstein, J. H., & Musolff, A. (2021). Nitrate Transport and Retention in Western European Catchments Are Shaped by Hydroclimate and Subsurface Properties. *Water Resources Research*, 57(10), e2020WR029469. <https://doi.org/10.1029/2020WR029469>
- 555 Elliott, M., & Quintino, V. (2007). The Estuarine Quality Paradox, Environmental Homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. *Marine Pollution Bulletin*, 54(6), 640–645. <https://doi.org/10.1016/J.MARPOLBUL.2007.02.003>
- Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., Ngai, J. T., Seabloom, E. W., Shurin, J. B., & Smith, J. E. (2007). Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters*, 10(12), 1135–1142. <https://doi.org/10.1111/j.1461-0248.2007.01113.x>
- 560 Fasola, S., Muggeo, V. M. R., & Küchenhoff, H. (2018). A heuristic, iterative algorithm for change-point detection in abrupt change models. *Computational Statistics*, 33(2), 997–1015. <https://doi.org/10.1007/S00180-017-0740-4/FIGURES/7>
- Filazzola, A., Mahdiyan, O., Shuvo, A., Ewins, C., Moslenko, L., Sadid, T., Blaggrave, K., Imrit, M. A., Gray, D. K., Quinlan, R., O'Reilly, C. M., & Sharma, S. (2020). A database of chlorophyll and water chemistry in freshwater lakes. *Scientific Data*, 7(1), 310. <https://doi.org/10.1038/s41597-020-00648-2>
- 565 Filstrup, C. T., King, K. B. S., & McCullough, I. M. (2019). Evenness effects mask richness effects on ecosystem functioning at macro-scales in lakes. *Ecology Letters*, 22(12), 2120–2129. <https://doi.org/10.1111/ele.13407>
- Food and Agriculture Organization of the United Nations, & FAO Land and Water Division. (2024, October 1). *Major hydrological basins of the world*. <https://data.apps.fao.org/catalog/iso/7707086d-af3c-41cc-8aa5-323d8609b2d1>
- 570 Gilarranz, L. J., Narwani, A., Odermatt, D., Siber, R., Dakos D Edited, V., & Hastings, A. (2022). *Regime shifts, trends, and variability of lake productivity at a global scale*. <https://doi.org/10.1073/pnas>
- Greening, H., & Janicki, A. (2006). Toward reversal of eutrophic conditions in a subtropical estuary: Water quality and seagrass response to nitrogen loading reductions in Tampa Bay, Florida, USA. *Environmental Management*, 38(2), 163–178. <https://doi.org/10.1007/S00267-005-0079-4/TABLES/2>
- 575 Greening, H., Janicki, A., Sherwood, E. T., Pribble, R., & Johansson, J. O. R. (2014). Ecosystem responses to long-term nutrient management in an urban estuary: Tampa Bay, Florida, USA. *Estuarine, Coastal and Shelf Science*, 151, A1–A16. <https://doi.org/10.1016/J.ECSS.2014.10.003>
- Harding, L. W., Mallonee, M. E., Perry, E. S., Miller, W. D., Adolf, J. E., Gallegos, C. L., & Paerl, H. W. (2019). Long-term trends, current status, and transitions of water quality in Chesapeake Bay. *Scientific Reports*, 9(1), 6709. <https://doi.org/10.1038/s41598-019-43036-6>
- 580 Hilt, S., Köhler, J., Kozerski, H.-P., van Nes, E. H., & Scheffer, M. (2011). Abrupt regime shifts in space and time along rivers and connected lake systems. *Oikos*, 120(5), 766–775. <https://doi.org/10.1111/j.1600-0706.2010.18553.x>
- Hobbs, R. J., Higgs, E., & Harris, J. A. (2009). Novel ecosystems: implications for conservation and restoration. *Trends in Ecology & Evolution*, 24(11), 599–605. <https://doi.org/10.1016/j.tree.2009.05.012>

- 585 Hoyer, M. V., Frazer, T. K., Notestein, S. K., & Canfield Jr., D. E. (2002). Nutrient, chlorophyll, and water clarity relationships in Florida's nearshore coastal waters with comparisons to freshwater lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 59(6), 1024–1031. <https://doi.org/10.1139/f02-077>
- Ibáñez, C., Alcaraz, C., Caiola, N., Rovira, A., Trobajo, R., Alonso, M., Duran, C., Jiménez, P. J., Munné, A., & Prat, N. (2012). Regime shift from phytoplankton to macrophyte dominance in a large river: Top-down versus bottom-up effects. *Science of the Total Environment*, 416, 314–322. <https://doi.org/10.1016/j.scitotenv.2011.11.059>
- 590 Ibáñez, C., & Peñuelas, J. (2019). Changing nutrients, changing rivers. *Science*, 365(6454), 637–638. <https://doi.org/10.1126/science.aay2723>
- Jarvie, H. P., Sharpley, A. N., Scott, J. T., Haggard, B. E., Bowes, M. J., & Massey, L. B. (2012). Within-River Phosphorus Retention: Accounting for a Missing Piece in the Watershed Phosphorus Puzzle. *Environmental Science & Technology*, 46(24), 13284–13292. <https://doi.org/10.1021/es303562y>
- 595 Jeppesen, E., Søndergaard, M., Jensen, J. P., Havens, K. E., Anneville, O., Carvalho, L., Coveney, M. F., Deneke, R., Dokulil, M. T., Foy, B., Gerdeaux, D., Hampton, S. E., Hilt, S., Kangur, K., Köhler, J., Lammens, E. H. h. r., Lauridsen, T. L., Manca, M., Miracle, M. R., ... Winder, M. (2005). Lake responses to reduced nutrient loading – an analysis of contemporary long-term data from 35 case studies. *Freshwater Biology*, 50(10), 1747–1771. <https://doi.org/10.1111/j.1365-2427.2005.01415.x>
- 600 Jochimsen, M. C., Kümmerlin, R., & Straile, D. (2013). Compensatory dynamics and the stability of phytoplankton biomass during four decades of eutrophication and oligotrophication. *Ecology Letters*, 16(1), 81–89. <https://doi.org/10.1111/ele.12018>
- Krishna, S., Ulloa, H. N., Kerimoglu, O., Minaudo, C., Anneville, O., & Wüest, A. (2021). Model-based data analysis of the effect of winter mixing on primary production in a lake under reoligotrophication. *Ecological Modelling*, 440, 109401. <https://doi.org/10.1016/j.ecolmodel.2020.109401>
- 605 Kronvang, B., Jeppesen, E., Conley, D. J., Søndergaard, M., Larsen, S. E., Ovesen, N. B., & Carstensen, J. (2005). Nutrient pressures and ecological responses to nutrient loading reductions in Danish streams, lakes and coastal waters. *Journal of Hydrology*, 304(1), 274–288. <https://doi.org/10.1016/j.jhydrol.2004.07.035>
- 610 Lehner, B., Messenger, M. L., Korver, M. C., & Linke, S. (2022). Global hydro-environmental lake characteristics at high spatial resolution. *Scientific Data*, 9(1), 1–19. <https://doi.org/10.1038/s41597-022-01425-z>
- Le Moal, M., Gascuel-Oudou, C., Ménesguen, A., Souchon, Y., Étrillard, C., Levain, A., Moatar, F., Pannard, A., Souchu, P., Lefebvre, A., & Pinay, G. (2019). Eutrophication: A new wine in an old bottle? *Science of The Total Environment*, 651, 1–11. <https://doi.org/10.1016/j.scitotenv.2018.09.139>
- 615 Linke, S., Lehner, B., Ouellet Dallaire, C., Ariwi, J., Grill, G., Anand, M., Beames, P., Burchard-Levine, V., Maxwell, S., Moidu, H., Tan, F., & Thieme, M. (2019). Global hydro-environmental sub-basin and river reach characteristics at high spatial resolution. *Scientific Data*, 6(1), 283. <https://doi.org/10.1038/s41597-019-0300-6>

620 Liu, M., Raymond, P. A., Lauerwald, R., Zhang, Q., Trapp-Müller, G., Davis, K. L., Moosdorf, N., Xiao, C., Middelburg, J.  
 J., Bouwman, A. F., Beusen, A. H. W., Peng, C., Lacroix, F., Tian, H., Wang, J., Li, M., Zhu, Q., Cohen, S., van Hoek,  
 W. J., ... Regnier, P. (2024). Global riverine land-to-ocean carbon export constrained by observations and multi-model  
 assessment. *Nature Geoscience*, 17(9), 896–904. <https://doi.org/10.1038/s41561-024-01524-z>  
 Minaudo, C., Abonyi, A., Leitão, M., Lançon, A. M., Floury, M., Descy, J.-P., & Moatar, F. (2021). Long-term impacts of  
 nutrient control, climate change, and invasive clams on phytoplankton and cyanobacteria biomass in a large temperate  
 river. *Science of The Total Environment*, 756, 144074. <https://doi.org/10.1016/j.scitotenv.2020.144074>  
 625 Minaudo, C., & Benito, X. (2024, January 30). *OLIGOTREND, a global database of multi-decadal timeseriesime series of*  
*chlorophyll-a and nutrient concentrations in inland and transitional waters, 1986-2023 ver 3*. Environmental Data  
 Initiative. <https://doi.org/10.6073/pasta/a7ad060a4dbc4e7dfcb763a794506524>  
 Minaudo, C., Meybeck, M., Moatar, F., Gassama, N., & Curie, F. (2015). Eutrophication mitigation in rivers: 30 years of  
 trends in spatial and seasonal patterns of biogeochemistry of the Loire River (1980–2012). *Biogeosciences*, 12(8), 2549–  
 630 2563. <https://doi.org/10.5194/bg-12-2549-2015>  
 Murphy, R. R., Keisman, J., Harcum, J., Karrh, R. R., Lane, M., Perry, E. S., & Zhang, Q. (2022). Nutrient Improvements in  
 Chesapeake Bay: Direct Effect of Load Reductions and Implications for Coastal Management. *Environmental Science*  
*& Technology*, 56(1), 260–270. <https://doi.org/10.1021/acs.est.1c05388>  
 Naderian, D., Noori, R., Heggy, E., Bateni, S. M., Bhattarai, R., Nohegar, A., & Sharma, S. (2024). A water quality database  
 635 for global lakes. *Resources, Conservation and Recycling*, 202, 107401. <https://doi.org/10.1016/j.resconrec.2023.107401>  
 Némery, J., & Garnier, J. (2016). The fate of phosphorus. *Nature Geoscience*, 9(5), 343–344. <https://doi.org/10.1038/ngeo2702>  
 Nilsson, J. L., Camiolo, S., Huser, B., Agstam-Norlin, O., & Futter, M. (2024). Widespread and persistent oligotrophication  
 of northern rivers. *Science of The Total Environment*, 955, 177261. <https://doi.org/10.1016/j.scitotenv.2024.177261>  
 Paerl, H. W., Scott, J. T., McCarthy, M. J., Newell, S. E., Gardner, W. S., Havens, K. E., Hoffman, D. K., Wilhelm, S. W., &  
 640 Wurtsbaugh, W. A. (2016). It Takes Two to Tango: When and Where Dual Nutrient (N & P) Reductions Are  
 Needed to Protect Lakes and Downstream Ecosystems. *Environmental Science & Technology*, 50(20), 10805–10813.  
<https://doi.org/10.1021/acs.est.6b02575>  
 Pinay, G., Gascuel, C., Ménesguen, A., Souchon, Y., & Le Moal, M. (2017). *Eutrophication: manifestations, causes,*  
*consequences and predictability*.  
 645 Pohlert, T. (2023). *trend: Non-Parametric Trend Tests and Change-Point Detection*. [https://CRAN.R-](https://CRAN.R-project.org/package=trend)  
[project.org/package=trend](https://CRAN.R-project.org/package=trend)  
 Powers, S. M., & Hampton, S. E. (2019). Open science, reproducibility, and transparency in ecology. *Ecological Applications*,  
 29(1), e01822. <https://doi.org/10.1002/eap.1822>  
 Ptacnik, R., Solimini, A. G., Andersen, T., Tamminen, T., Brettum, P., Lepistö, L., Willén, E., & Rekolainen, S. (2008).  
 650 Diversity predicts stability and resource use efficiency in natural phytoplankton communities. *Proceedings of the*  
*National Academy of Sciences*, 105(13), 5134–5138. <https://doi.org/10.1073/pnas.0708328105>

Formatted: Spanish (Spain)

- Reynolds, C. S. (2006). *The ecology of phytoplankton* (M. Usher, D. Saunders, R. Peet, & A. Dobson, Eds.; Cambridge). Cambridge University Press.
- Romero, E., Garnier, J., Lassaletta, L., Billen, G., Le Gendre, R., Riou, P., & Cugier, P. (2013). Large-scale patterns of river inputs in southwestern Europe: Seasonal and interannual variations and potential eutrophication effects at the coastal zone. *Biogeochemistry*, 113(1–3), 481–505. <https://doi.org/10.1007/S10533-012-9778-0>~~FIGURES/6~~
- Ross, M. R. V., Topp, S. N., Appling, A. P., Yang, X., Kuhn, C., Butman, D., Simard, M., & Pavelsky, T. (2019). AquaSat: a dataset to enable remote sensing of water quality for inland waters. *Water Resources Research*, 1–14. <https://doi.org/10.1029/2019wr024883>
- Sabel, M., Eckmann, R., Jeppesen, E., Rösch, R., & Straile, D. (2020). Long-term changes in littoral fish community structure and resilience of total catch to re-oligotrophication in a large, peri-alpine European lake. *Freshwater Biology*, 65(8), 1325–1336. <https://doi.org/10.1111/fwb.13501>
- Scheffer, M., & Carpenter, S. R. (2003). Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends in Ecology & Evolution*, 18(12), 648–656. <https://doi.org/10.1016/J.TREE.2003.09.002>
- Selmeczy, G. B., Abonyi, A., Krienitz, L., Kasprzak, P., Casper, P., Telcs, A., Somogyvári, Z., & Padisák, J. (2019). Old sins have long shadows: climate change weakens efficiency of trophic coupling of phyto- and zooplankton in a deep oligo-mesotrophic lowland lake (Stechlin, Germany)—a causality analysis. *Hydrobiologia*, 831(1), 101–117. <https://doi.org/10.1007/s10750-018-3793-7>
- Spaulding, S. A., Platt, L. R. C., Murphy, J. C., Covert, A., & Harvey, J. W. (2024). Chlorophyll a in lakes and streams of the United States (2005–2022). *Scientific Data*, 11(1), 611. <https://doi.org/10.1038/s41597-024-03453-3>
- Stackpoole, S. M., Stets, E. G., & Sprague, L. A. (2019). Variable impacts of contemporary versus legacy agricultural phosphorus on US river water quality. *Proceedings of the National Academy of Sciences*, 201903226. <https://doi.org/10.1073/pnas.1903226116>
- Van Meter, K. J., McLeod, M. M., Liu, J., Tenkouano, G. T., Hall, R. I., Van Cappellen, P., & Basu, N. B. (2021). Beyond the Mass Balance: Watershed Phosphorus Legacies and the Evolution of the Current Water Quality Policy Challenge. *Water Resources Research*, 57(10). <https://doi.org/10.1029/2020WR029316>
- van Vliet, M. T. H., Thorslund, J., Stokal, M., Hofstra, N., Flörke, M., Ehalt Macedo, H., Nkwasa, A., Tang, T., Kaushal, S. S., Kumar, R., van Griensven, A., Bouwman, L., & Mosley, L. M. (2023). Global river water quality under climate change and hydroclimatic extremes. *Nature Reviews Earth & Environment* 2023 4:10, 4(10), 687–702. <https://doi.org/10.1038/S43017-023-00472-3>
- Verdonschot, P. F. M., Spears, B. M., Feld, C. K., Brucet, S., Keizer-Vlek, H., Borja, A., Elliott, M., Kernan, M., & Johnson, R. K. (2013). A comparative review of recovery processes in rivers, lakes, estuarine and coastal waters. *Hydrobiologia*, 704(1), 453–474. <https://doi.org/10.1007/s10750-012-1294-7>
- Virro, H., Amatulli, G., Kmoch, A., Shen, L., & Uemaa, E. (2021). GRQA: Global River Water Quality Archive. *Earth System Science Data*, 13(12), 5483–5507. <https://doi.org/10.5194/essd-13-5483-2021>

Wachholz, A., Jawitz, J.W., & Borchardt, D. (2024). From Iron Curtain to green belt: shift from heterotrophic to autotrophic nitrogen retention in the Elbe River over 35 years of passive restoration. *Biogeosciences* 21 (15), 3537 - 3550 <https://doi.org/10.5194/bg-21-3537-2024>

Formatted: Font: Italic

Formatted: Font: Italic

Welti, E. A. R., Bowler, D. E., Sinclair, J. S., Altermatt, F., Álvarez-Cabria, M., Amatulli, G., Angeler, D. G., Archambaud, G., Arrate Jorrín, I., Aspin, T., Azpiroz, I., Baker, N. J., Bañares, I., Barquín Ortiz, J., Bodin, C. L., Bonacina, L., Bonada, N., Bottarin, R., Cañedo-Argüelles, M., ... Haase, P. (2024). Time series of freshwater macroinvertebrate abundances and site characteristics of European streams and rivers. *Scientific Data*, 11(1), 601. <https://doi.org/10.1038/s41597-024-03445-3>

Zeng, L., Ji, J., Xu, S., Cao, Y., & Chen, X. (2023). Decoupling of nitrogen, phosphorus and biogenic silica in floodplain sediments in response to land use change and hydrological alterations. *Journal of Hydrology*, 623, 129833. <https://doi.org/10.1016/j.jhydrol.2023.129833>