

Title: ABCFlux v2: Arctic–boreal CO₂ and CH₄ monthly flux observations and ancillary information across terrestrial and freshwater ecosystems

Journal: Earth System Science Data

Authors: Anna-Maria Virkkala¹, Isabel Wargowsky¹, Judith Vogt², Simran Madaan², Tiffany Windholz¹, Kyle A. Arndt¹, Gerard Rocher-Ros⁵⁵, Mathias Göckede², David Olefeldt⁸⁰, Edward A. G. Schuur¹⁵, David Bastviken⁸², Richard O'Keefe¹, McKenzie A. Kuhn^{3,1}, Brendan M. Rogers¹, Jennifer D. Watts¹, Kelcy Kent¹, Kristoffer Aalstad⁷⁵, Kelly Aho³⁵, Joonatan Ala-Könni¹¹⁹, Haley Alcock⁶⁸, Inge Althuizen¹⁵², Christopher D. Arp¹⁸⁵, Jun Asanuma¹⁸¹, Katrin Attermeyer^{65, 184}, Mika Aurela⁹⁷, Sivakiruthika Balathandayuthabani^{82, 171}, Alan Barr¹⁰⁶, Maialen Barret¹³⁷, Ochirbat Batkhishig¹²⁸, Christina Biasi^{56, 61}, Mats P. Björkman^{34, 107}, Andrew Black⁹⁴, Elena Blanc-Betes¹²³, Pascal Bodmer⁵¹, Julia Boike^{7, 115}, Abdullah Bolek², Frédéric Bouchard^{17, 28}, Ingeborg Busmann⁸³, Lea Cabrol^{6, 142}, Eleonora Canfora²³, Sean Carey¹⁶³, Karel Castro-Morales¹⁰², Namyi Chae¹³¹, Andreas Christen⁸⁹, Torben R. Christensen^{58, 186}, Casper T. Christiansen³⁸, Housen Chu²², Graham Clark⁴², Francois Clayer¹⁵⁴, Patrick Crill⁷², Christopher Cunada¹⁰⁸, Scott J. Davidson²⁶, Joshua F. Dean¹⁶⁷, Sigrid Dengel¹³⁸, Matteo Detto¹¹⁴, Catherine Dieleman¹⁶⁵, Florent Domine¹⁷⁰, Egor Dyukarev^{132, 188}, Colin Edgar¹²⁴, Bo Elberling⁷⁴, Craig A. Emmerton³¹, Eugenie Euskirchen¹²⁴, Grant Falvo¹⁵, Thomas Friborg⁷⁴, Michelle Garneau¹⁰⁴, Mariasilvia Giamberini¹²⁹, Mikhail V. Glagolev^{95, 126}, Miquel A. Gonzalez-Meler³², Gustaf Granath⁵⁴, Jón Guðmundsson⁹³, Konsta Happonen⁹⁹, Yoshinobu Harazono¹³⁴, Lorna Harris¹¹⁸, Josh Hashemi⁷, Nicholas Hasson¹⁷⁶, Janna Heerah⁷⁸, Liam Heffernan⁴⁷, Manuel Helbig^{78, 105}, Warren Helgason²⁰, Michal Heliasz⁴¹, Greg Henry³, Geert Hensgens^{8, 47}, Tetsuya Hiyama¹²², Macall Hock^{37, 178}, David Holl¹³³, Beth Holmes⁵⁰, Jutta Holst⁴¹, Thomas Holst⁴¹, Gabriel Hould-Gosselin⁶⁸, Elyn Humphreys⁶⁶, Jacqueline Hung¹, Jussi Huotari^{91, 140}, Hiroki Ikawa¹⁴³, Danil V. Ilyasov¹⁷², Mamoru Ishikawa⁹², Go Iwahana^{9, 134}, Hiroki Iwata^{62, 121}, Marcin Antoni Jackowicz-Korczynski⁵⁷, Joachim Jansen¹¹⁹, Järvi Järveoja⁶⁴, Vincent E.J. Jassey¹⁷⁴, Rasmus Jensen⁵⁸, Katharina Jentsch⁷, Robert G. Jaspersen¹⁷³, Carl-Fredrik Johannesson^{74, 153}, Cheristy P. Jones⁴⁵, Anders Jonsson⁵⁵, Ji Young Jung¹³⁶, Sari Juutinen⁹⁸, Evan Kane²⁴, Jan Karlsson⁵⁵, Sergey Karsanaev¹²⁰, Kuno Kasak⁷¹, Julia Kelly⁴¹, Kasha Kempton⁸⁰, Marcus Klaus⁶⁴, George W. Kling⁵³, Natascha Kljun⁴¹, Jacqueline Knutson¹⁵⁴, Hideki Kobayashi¹³⁵, John Kochendorfer¹⁵¹, Kukka-Maaria Kohonen^{63, 119}, Pasi Kolari¹⁷⁹, Mika Korkiakoski⁹⁷, Aino Korrensalo^{61, 150}, Pirkko Kortelainen⁹⁶, Egle Koster¹⁶⁶, Kajar Koster⁶⁰, Ayumi Kotani¹¹⁰, Praveena Krishnan¹⁵¹, Juliya Kurbatova⁴, Lars Kutzbach¹³³, Min Jung Kwon^{48, 136}, Ethan D. Kyzivat^{11, 44}, Jessica Lagroix¹⁷⁷, Theodore Langhorst¹⁸⁰, Elena Lapshina¹⁷², Tuula Larmola¹⁴⁹, Klaus S. Larsen⁷⁴, Isabelle Laurion^{18, 17}, Justin Ledman¹⁵, Hanna Lee³⁶, A. Joshua Leffler⁷⁷, Lance Lesack⁸⁴, Anders Lindroth⁴¹, David Lipson³⁷, Annalea Lohila^{97, 119}, Efrén López-Blanco^{57, 59}, Vincent L. St. Louis³¹, Erik Lundin⁵, Miska Luoto⁷³, Takashi Machimura¹¹¹, Marta Magnani^{130, 144, 146}, Avni Malhotra¹², Marja Maljanen⁶¹, Ivan Mammarella¹¹⁹, Elisa Männistö¹⁵⁶, Luca Beletti Marchesini¹⁰⁰, Phil Marsh¹⁸⁷, Pertti J. Martikainen⁶¹, Maija E. Marushchak⁶¹, Mikhail Mastepanov^{57, 155}, Alex Mavrovic^{13, 17, 175}, Trofim Maximov¹²⁰, Christina Minions¹, Marco Montemayor¹, Tomoaki Morishita¹⁰¹, Patrick Murphy¹, Daniel F. Nadeau⁴⁰, Erin Nicholls⁴⁹, Mats B. Nilsson⁶⁴, Anastasia Niyazova¹⁷², Jenni Nordén¹⁵³, Koffi Dodji Noumonvi⁶⁴, Hannu Nykanen⁶¹, Walter Oechel³⁷, Anne Ojala¹⁴⁹, Tomohiro Okadera¹⁴⁵, Sujun Pal⁹⁰, Alexey V. Panov¹⁸³, Tim Papakyriakou¹⁹, Dario Papale¹⁴⁷, Sang-Jong Park¹³⁶, Frans-Jan W. Parmentier¹⁴,

Gilberto Pastorello¹⁶⁸, Mike Peacock^{29, 67}, Matthias Peichl⁶⁴, Roman Petrov¹²⁰, Kyra St. Pierre⁴³,
 Norbert Pirk⁷⁵, Jessica Plein³⁷, Vilmantas Preskienis^{18, 17, 27, 113}, Anatoly Prokushkin¹⁸³, Jukka
 45 Pumpanen⁶¹, Hilary A. Rains¹⁵⁷, Niklas Rakos⁵, Aleksi Räsänen¹⁰³, Helena Rautakoski⁹⁸, Riikka
 Rinnan³⁸, Janne Rinne¹⁴⁹, Adrian Rocha³³, Nigel Roulet¹⁴¹, Alexandre Roy^{17, 175}, Anna
 Rutgersson⁴⁶, Aleksandr F. Sabrekov¹⁷², Torsten Sachs^{105, 127}, Erik Sahlée⁴⁶, Alejandro Salazar⁹³,
 Henrique Oliveira Sawakuchi⁸², Christopher Schulze^{68, 80}, Roger Seco¹²⁵, Armando Sepulveda-
 Jauregui^{79, 139}, Svetlana Serikova⁵⁵, Abbey Serrone^{21, 34}, Hanna M. Silvennoinen¹⁵³, Sofie
 50 Sjogersten¹⁶², June Skeeter^{3, 148}, Jo Snöäl⁷⁰, Sebastian Sobek^{54, 76}, Oliver Sonnentag⁶⁸, Emily H.
 Stanley¹⁶, Maria Strack¹⁸², Lena Strom⁴¹, Patrick Sullivan⁸⁸, Ryan Sullivan⁹⁰, Anna Sytiuk¹⁷⁴,
 Torbern Tagesson⁴¹, Pierre Taillardat¹⁰, Julie Talbot⁶⁸, Suzanne E. Tank³¹, Mario Tenuta⁸¹, Irina
 Terenteva⁶⁹, Frederic Thalasso³⁹, Antoine Thibault⁴⁰, Halldor Thorgeirsson¹¹⁷, Fenix Garcia
 Tigreros¹¹², Margaret Torn²², Amy Townsend-Small¹⁶⁴, Claire Treat^{7, 158}, Alain Tremblay¹¹⁶, Carlo
 55 Trotta²³, Eeva-Stiina Tuittila¹⁵⁶, Merritt Turetsky⁵², Masahito Ueyama¹⁰⁹, Muhammad Umair⁶⁸, Aki
 Vähä¹¹⁹, Lona van Delden⁷, Maarten van Hardenbroek²⁵, Andrej Varlagin⁴, Ruth K. Varner^{45, 72},
 Elena Veretennikova^{132, 169}, Timo Vesala¹⁷⁹, Tarmo Virtanen⁸⁷, Carolina Voigt^{7, 48, 68}, Jorien E.
 Vonk⁴⁷, Robert Wagner³⁷, Katey Walter Anthony¹⁸⁵, Qinxue Wang¹⁴⁵, Masataka Watanabe¹⁶¹,
 Hailey Webb^{52, 160}, Jeffrey M. Welker^{30, 86}, Andreas Westergaard-Nielsen⁷⁴, Sebastian
 60 Westermann¹⁴, Jeffrey R. White⁸⁵, Christian Wille¹⁰⁵, Scott N. Williamson¹⁵⁹, Scott Zolkos¹,
 Donatella Zona³⁷, Susan M. Natali¹

* shared first-authorship

¹Woodwell Climate Research Center, Falmouth, MA, USA

²Max Planck Institute for Biogeochemistry, Jena, Germany

65 ³Department of Geography, University of British Columbia, Vancouver, Canada

⁴A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

⁵Abisko Scientific Research Station, Swedish Polar Research Secretariat, Sweden

⁶Aix-Marseille University, University Toulon, CNRS, IRD, Mediterranean Institute of Oceanography (M.I.O.) Marseille, France

70 ⁷Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

⁸Arctic Geology, University Centre in Svalbard, Svalbard, Norway

⁹Arctic Research Center, Hokkaido University, Japan

¹⁰Asian School of the Environment, Nanyang Technological University, 639798 Singapore

¹¹Biodiversity Research Institute, Portland, ME, USA

75 ¹²Biological Sciences Division, Pacific Northwest National Laboratory, USA

¹³Cegep de Sherbrooke, Canada

¹⁴Center for Biogeochemistry of the Anthropocene, Department of Geosciences, University of Oslo, Oslo, Norway

¹⁵Center for Ecosystem Science and Society, Northern Arizona University, USA

80 ¹⁶Center for Limnology, University of Wisconsin-Madison, Madison, WI 53706, USA

¹⁷Centre d'études nordiques (CEN), Université Laval, Québec, Canada

¹⁸Centre Eau Terre Environnement, Institut national de la recherche scientifique, Canada

¹⁹Centre for Earth Observation Science, University of Manitoba, MB, Canada

²⁰Chemical and Biological Engineering, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

85 ²¹Civil and Environmental Engineering, University of Washington, USA

²²Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, USA

²³CMCC Foundation - Euro-Mediterranean Center on Climate Change, Italy

- ²⁴College of Forest Resources and Environmental Science, Michigan Technological University, USA
- 90 ²⁵Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, Netherlands
- ²⁶Département des sciences biologiques, Université du Québec à Montréal, Montréal, Canada
- ²⁷Département des sciences fondamentales, Université du Québec à Chicoutimi, Saguenay, Canada
- ²⁸Department of Applied Geomatics, Université de Sherbrooke, Sherbrooke, Canada
- 95 ²⁹Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, Sweden
- ³⁰Department of Biological Sciences, University of Alaska Anchorage, Anchorage, AK, USA
- ³¹Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada
- ³²Department of Biological Sciences, University of Illinois at Chicago, Chicago, IL 60607, USA
- ³³Department of Biological Sciences, University of Notre Dame, USA
- 100 ³⁴Department of Biology and Environmental Sciences, University of Gothenburg, Box 463, SE-40530 Gothenburg, Sweden
- ³⁵Department of Biology, Boston University, USA
- ³⁶Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway
- ³⁷Department of Biology, SDSU, San Diego, CA, USA
- 105 ³⁸Department of Biology, University of Copenhagen, Denmark
- ³⁹Department of Biotechnology and Bioengineering Department, Center for Research and Advanced Studies (Cinvestav), Mexico City, Mexico
- ⁴⁰Department of Civil and Water Engineering, Université Laval, QC, Canada
- 110 ⁴¹Department of Earth and Environmental Sciences, Lund University, Sölvegatan 12, 223 62 Lund, Sweden
- ⁴²Department of Earth and Environmental Sciences, St. Francis Xavier University, Antigonish, Canada
- ⁴³Department of Earth and Environmental Sciences, University of Ottawa, Ottawa ON Canada, K1N 6N5, Canada
- ⁴⁴Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA
- 115 ⁴⁵Department of Earth Sciences and Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, NH
- ⁴⁶Department of Earth Sciences, Uppsala University, Sweden
- ⁴⁷Department of Earth Sciences, Vrije Universiteit, Amsterdam, Netherlands
- ⁴⁸Department of Earth System Sciences, University of Hamburg, Hamburg, Germany
- 120 ⁴⁹Department of Earth, Energy and Environment, University of Calgary, Canada
- ⁵⁰Department of Earth, Ocean, and Atmospheric Science, Florida State University, USA
- ⁵¹Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, USA
- ⁵²Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Boulder, CO USA
- ⁵³Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA
- 125 ⁵⁴Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden
- ⁵⁵Department of Ecology, Environment and Geoscience, Umeå University, Umeå, Sweden
- ⁵⁶Department of Ecology, University of Innsbruck, Innsbruck, Austria
- ⁵⁷Department of Ecoscience and Arctic Research Centre, Aarhus University, Roskilde, Denmark
- ⁵⁸Department of Ecoscience, Aarhus University, Denmark
- 130 ⁵⁹Department of Environment and Minerals, Greenland Institute of Natural Resources, Nuuk, Greenland
- ⁶⁰Department of Environmental and Biological Sciences, University of Eastern Finland, Joensuu, Finland
- ⁶¹Department of Environmental and Biological Sciences, University of Eastern Finland, Kuopio, Finland
- ⁶²Department of Environmental Science, Shinshu University, Matsumoto, Japan
- 135 ⁶³Department of Environmental Systems Science, Institute for Agricultural Sciences, ETH Zurich, Zurich, Switzerland

- ⁶⁴Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå, Sweden
- ⁶⁵Department of Functional and Evolutionary Ecology, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria
- 140 ⁶⁶Department of Geography & Environmental Studies, Carleton University, Ottawa, Canada
- ⁶⁷Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool, UK
- ⁶⁸Department of Geography, Université de Montréal, QC, Canada
- ⁶⁹Department of Geography, University of Calgary, Calgary, AB T2N 1N4, Canada
- 145 ⁷⁰Department of Geography, University of Exeter, United Kingdom
- ⁷¹Department of Geography, University of Tartu, Estonia
- ⁷²Department of Geological Sciences and Bolin Centre for Climate Research, Stockholm University, Sweden
- ⁷³Department of Geosciences and Geography, University of Helsinki, Finland
- 150 ⁷⁴Department of Geosciences and Natural Resource Management, University of Copenhagen, Denmark
- ⁷⁵Department of Geosciences, University of Oslo, Oslo, Norway
- ⁷⁶Department of Limnology, Uppsala University, Sweden
- ⁷⁷Department of Natural Resource Management, South Dakota State University, Brookings, SD 57007, USA
- 155 ⁷⁸Department of Physics and Atmospheric Science, Dalhousie University, Halifax, NS, Canada
- ⁷⁹Department of Plankton and Microbial Ecology, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Stechlin, Germany
- ⁸⁰Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada
- ⁸¹Department of Soil Science, University of Manitoba, Winnipeg, Manitoba, Canada
- 160 ⁸²Department of Thematic Studies-Environmental Change, Linköping University, Linköping, Sweden
- ⁸³Department Shelf Sea System Ecology, Alfred Wegener Institute, Helgoland, Germany
- ⁸⁴Departments of Geography and Biological Sciences, Simon Fraser University, Canada
- ⁸⁵Earth and Atmospheric Sciences, Environmental Sciences, Indiana University Bloomington, USA
- ⁸⁶Ecology and Genetics Research Unit, University of Oulu, Oulu, Finland
- 165 ⁸⁷Ecosystems and Environment Research Programme, University of Helsinki, Finland
- ⁸⁸Environment and Natural Resources Institute, University of Alaska Anchorage, Anchorage, AK 99508, USA
- ⁸⁹Environmental Meteorology, Faculty of Environment and Natural Resources, University of Freiburg, Germany
- 170 ⁹⁰Environmental Science Division, Argonne National Laboratory, Lemont, IL, USA
- ⁹¹Faculty of Biological and Environmental Sciences, Ecosystems and Environment Research Programme, Lammi Biological Station, University of Helsinki, Lammi FI-16900, Finland
- ⁹²Faculty of Earth Environmental Science, Hokkaido University, Japan
- ⁹³Faculty of Environmental and Forest Sciences, Agricultural University of Iceland, Iceland
- 175 ⁹⁴Faculty of Land and Food Systems, University of British Columbia, Vancouver, British Columbia, Canada
- ⁹⁵Faculty of Soil Science, Lomonosov Moscow State University, building 12, Leninskie Gory, GSP-1, 119991, Moscow, Russia
- ⁹⁶Finnish Environment Institute, Finland
- 180 ⁹⁷Finnish Meteorological Institute, Climate System Research, Finland
- ⁹⁸Finnish Meteorological Institute, Finland
- ⁹⁹Finnish Youth Research Society, Finland
- ¹⁰⁰Forest Ecology Unit, Research and Innovation Centre, Edmund Mach Foundation, Italy

- 185 ¹⁰¹Forestry and Forest Products Research Institute, Tohoku Research Center, Japan
¹⁰²Friedrich Schiller University Jena, Jena, Germany
¹⁰³Geography Research Unit, University of Oulu, Oulu, Finland
¹⁰⁴Geotop, Université du Québec à Montréal, Montreal, QC, Canada
¹⁰⁵GFZ Helmholtz Centre for Geosciences, Potsdam, Germany
¹⁰⁶Global Institute for Water Security, University of Saskatchewan, Saskatoon, Saskatchewan, Canada
- 190 ¹⁰⁷Gothenburg Global Biodiversity Centre, Box 463, SE-405 30 Gothenburg, Sweden
¹⁰⁸Government of the Northwest Territories, Canada
¹⁰⁹Graduate School of Agriculture, Osaka Metropolitan University, Sakai 599-8531, Japan
¹¹⁰Graduate School of Bioagricultural Sciences, Nagoya University, Japan
¹¹¹Graduate School of Engineering, The University of Osaka, Japan
- 195 ¹¹²Graduate School of Oceanography, University of Rhode Island, Narragansett, RI, USA
¹¹³Groupe de recherche interuniversitaire en limnologie (GRIL), Quebec, Canada
¹¹⁴High Meadows Environmental Institute, Princeton University, USA
¹¹⁵Humboldt University of Berlin, Germany
¹¹⁶Hydro-Quebec, Quebec, Canada
- 200 ¹¹⁷Icelandic Climate Council, Iceland
¹¹⁸Independent Researcher
¹¹⁹Institute for Atmospheric and Earth System Research, Faculty of Science, University of Helsinki, Finland
¹²⁰Institute for Biological Problems of Cryolithozone of the Siberian Branch of the RAS - Division of
- 205 Federal Research Centre "The Yakut Scientific Centre of the Siberian Branch of the Russian Academy of Sciences", Russia
¹²¹Institute for Mountain Science, Shinshu University, Matsumoto, Japan
¹²²Institute for Space-Earth Environmental Research, Nagoya University, Japan
¹²³Institute for Sustainability, Energy, and Environment, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
- 210 ¹²⁴Institute of Arctic Biology, University of Alaska Fairbanks, USA
¹²⁵Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona, Spain
¹²⁶Institute of Forest Science, Russian Academy of Sciences, 21 Sovetskaya st., Uspenskoe, Moscow region 143030, Russia
- 215 ¹²⁷Institute of Geoecology, Technische Universität Braunschweig, Braunschweig, Germany
¹²⁸Institute of Geography and Geoecology, Mongolian Academy of Sciences, Ulaanbaatar, Mongolia
¹²⁹Institute of Geosciences and Earth Resources, National Research Council (CNR-IGG), 56124, Pisa, Italy
¹³⁰Institute of Geosciences and Earth Resources, National Research Council (CNR-IGG), Turin, Italy
- 220 ¹³¹Institute of Life Science and Natural Resources, Korea University, Seoul, Republic of Korea
¹³²Institute of Monitoring of Climatic and Ecological Systems, Siberian Branch of the Russian Academy of Sciences, Russia
¹³³Institute of Soil Science, Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany
- 225 ¹³⁴International Arctic Research Center, University of Alaska Fairbanks, USA
¹³⁵Japan Agency for Marine-Earth Science and Technology, Japan
¹³⁶Korea Polar Research Institute, Incheon, South Korea
¹³⁷Laboratoire d'Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, Toulouse, France
- 230 ¹³⁸Lawrence Berkeley National Laboratory, USA
¹³⁹Limnological Institute, Department of Biology, University of Konstanz, Konstanz, Germany

- 235 ¹⁴⁰Masinotek Oy, Ensimmäinen Savu 2, Vantaa FI-01510, Finland
¹⁴¹McGill University, Canada
¹⁴²Millennium Institute Biodiversity of Antarctic and Subantarctic Ecosystems (BASE), Santiago 7800003, Chile
¹⁴³National Agriculture and Food Research Organization, Japan
¹⁴⁴National Biodiversity Future Center (NBFC), Palermo, Italy
¹⁴⁵National Institute for Environmental Studies, Tsukuba, 305-8506, Japan
¹⁴⁶National Institute of Nuclear Physics (INFN), Turin, Italy
240 ¹⁴⁷National Research Council (CNR) - IRET, Monterotondo Scalo, Italy
¹⁴⁸Natural Resources Canada, Canada
¹⁴⁹Natural Resources Institute Finland, Helsinki, Finland
¹⁵⁰Natural Resources Institute Finland, Natural Resources Unit, Joensuu, Finland
¹⁵¹NOAA Air Resources Laboratory, Atmospheric Turbulence and Diffusion Division, Oak Ridge, TN
245 37830, USA
¹⁵²NORCE Norwegian Research Centre and Bjerknes Centre for Climate Research, Bergen, Norway
¹⁵³Norwegian Institute for Nature Research (NINA), Oslo, Norway
¹⁵⁴Norwegian Institute for Water Research (NIVA), Oslo, Norway
¹⁵⁵Oulanka Research Station, University of Oulu, Kuusamo, Finland
250 ¹⁵⁶Peatland and Soil Ecology Research Group, School of Forest Sciences, University of Eastern Finland, Joensuu, Finland
¹⁵⁷PG Environmental, Fairfax, Virginia, USA
¹⁵⁸Pioneer Center Land-CRAFT, Aarhus University, Denmark
¹⁵⁹Polar Knowledge Canada, Canadian High Arctic Research Station, Cambridge Bay, Nunavut, Canada
255 ¹⁶⁰Renewable and Sustainable Energy Institute, University of Colorado Boulder, Boulder, CO USA
¹⁶¹Research and Development Initiative, Chuo University, Tokyo 112-8551, Japan
¹⁶²School of Biosciences, University of Nottingham, Nottingham, United Kingdom
¹⁶³School of Earth, Environment and Society, McMaster University
¹⁶⁴School of Environment and Sustainability, University of Cincinnati, USA
260 ¹⁶⁵School of Environmental Sciences, University of Guelph, Canada
¹⁶⁶School of Forest Sciences, University of Eastern Finland, Joensuu, Finland
¹⁶⁷School of Geographical Sciences, University of Bristol, Bristol, United Kingdom
¹⁶⁸Scientific Data Division, Lawrence Berkeley National Laboratory, USA
¹⁶⁹Siberian State Medical University, Russia
265 ¹⁷⁰Takuvik Joint International Laboratory, Université Laval (Canada) and CNRS-INSU (France), Québec City, Canada
¹⁷¹Tamil Nadu Agricultural University, Coimbatore, India
¹⁷²UNESCO Department "Environmental Dynamics and Global Climate Changes", Yugra State University, Khanty-Mansiysk, Russia
270 ¹⁷³United States Forest Service, University of Alaska Anchorage, USA
¹⁷⁴Université de Toulouse, Toulouse INP, CNRS, IRD, CRBE, Toulouse, France
¹⁷⁵Université du Québec à Trois-Rivières, Research Centre for Watershed-Aquatic ecosystem interactions (RIVE) , Canada
¹⁷⁶University of Alaska Fairbanks, Fairbanks, Alaska, USA
275 ¹⁷⁷University of Alberta, Canada
¹⁷⁸University of California Davis, USA
¹⁷⁹University of Helsinki, Finland
¹⁸⁰University of Massachusetts Amherst, USA
¹⁸¹University of Tsukuba, Tsukuba, Japan

280 ¹⁸²University of Waterloo, Canada
¹⁸³V. N. Sukachev Institute of forest, Siberian Branch of the Russian Academy of Sciences, Russia
¹⁸⁴WasserCluster Lunz, Biologische Station, Dr. Carl Kupelwieser Promenade 5, 3293 Lunz am See,
Austria
¹⁸⁵Water and Environmental Research Center, University of Alaska Fairbanks, Fairbanks, AK, USA
285 ¹⁸⁶Water, energy and environmental engineering research unit, Oulu University, Finland
¹⁸⁷Wilfred Laurier University, Canada
¹⁸⁸Yugra State University, Russia

Correspondence to: Anna-Maria Virkkala (avirkkala@woodwellclimate.org)

290

Abstract

Measurements of surface-atmosphere carbon dioxide (CO₂) and methane (CH₄) fluxes have been relatively sparse across the Arctic tundra and boreal biomes, causing significant
295 uncertainties in carbon budget estimates from the region. While the availability of Arctic-boreal carbon flux data has increased substantially over the past decade, the data have remained spread across different repositories, scientific articles, and unpublished sources, making it difficult to leverage. Here we present a new dataset of monthly Arctic-boreal carbon fluxes (ABCFlux v2) across terrestrial (wetlands and uplands) and freshwater (lakes and rivers)
300 ecosystems compiled from previous syntheses including the Arctic-boreal CO₂ flux database (ABCFlux v1), the Boreal-Arctic Wetland and Lake Methane Dataset (BAWLD-CH₄), and the Global River Methane Database (GRiMeDB). In addition, we consider data from general-purpose (e.g., Zenodo) and flux network repositories, literature, and site principal investigators. The dataset includes surface-atmosphere CO₂ fluxes of gross primary production (GPP),
305 ecosystem respiration (Reco), and net ecosystem exchange (NEE), alongside CH₄ fluxes. For aquatic ecosystems, we split CH₄ fluxes into diffusive and ebullitive flux pathways, and included potential emissions from transient storage in the water column ("storage fluxes"), alongside CO₂ and CH₄ concentrations dissolved in the surface water. Fluxes are measured through a variety of methods including chamber and eddy covariance techniques alongside bubble traps, ice-surveys, and concentration-based turbulence-driven modelling in aquatic ecosystems. The
310 monthly flux data are reported together with supporting methodological and environmental metadata. The resulting ABCFlux v2 has 23,847 flux site-months, 8,182 concentration site-months, and 199 seasonal observations from 1,024 sites, and includes 56,139 reported fluxes (i.e. sum of GPP, Reco, NEE, and CH₄ fluxes) from the years 1984 to 2024. The majority of
315 monthly observations occurred after 1999. Wetlands had the highest number of site-month observations (8,758), followed by boreal forest (6,981), lotic ecosystems (6,275), lentic ecosystems (3,799) and upland tundra (3,308). Measurements of CO₂ dominated the dataset across most ecosystem types (25,222) except for lentic ecosystems, where CH₄ flux site-months (3,098) were more frequent than CO₂ flux site-months (2,915). Overall, ABCFlux v2
320 includes 160% more site-months for terrestrial CO₂ flux data compared to ABCFlux v1. Integrating and updating BAWLD-CH₄ flux data from growing season averages to monthly fluxes resulted in 5,671 site-months of chamber CH₄ data compared to 762 site-years. This

collaborative initiative, involving contributions from over 260 researchers, provides a comprehensive overview of the current state of the Arctic-boreal carbon flux network and its data, and serves as an important step in reducing uncertainties in Arctic-boreal carbon budgets and in enhancing our understanding of climate feedbacks. The data can be accessed at ORNL DAAC at <https://doi.org/10.3334/ORNLDAAC/2448>.

1. Introduction

The Arctic-boreal region has historically been sparsely measured for carbon dioxide (CO₂) and methane (CH₄) fluxes (Baldocchi et al., 2018; Pallandt et al., 2022). This data sparsity, together with rapid warming, changes in hydrology, permafrost thaw, and other environmental shifts (Biskaborn et al., 2019; O'Neill et al., 2023; Rantanen et al., 2022; Webb et al., 2022), has created significant uncertainties in Arctic-boreal carbon budget estimates (Hugelius et al., 2024; Treat et al., 2024), hindering our capability to understand the fate of its large soil organic carbon stocks (Hugelius et al., 2014; Schuur et al., 2008, 2022). However, over the past decade, the availability of Arctic-boreal carbon flux data has increased substantially (Vogt et al., 2025), giving rise to new flux synthesis datasets (Kuhn et al., 2021; Virkkala et al., 2022) that have been widely used to improve our process-understanding (Kuhn et al., 2021), model intercomparisons (Tao et al., 2021; Treat et al., 2024), site-level trend assessments (See et al., 2024), and Arctic-boreal carbon budgets (Kuhn et al., 2025; Ramage et al., 2024; Virkkala et al., 2025; Vonk et al., 2025; Yuan et al., 2024).

While significant progress has been made in Arctic-boreal carbon flux datasets, most existing syntheses do not include recently published flux data from 2020 onward, a period marked by rapid warming (Minobe et al., 2025) and increased disturbances - such as fires (Euskirchen et al., 2024; Kelly et al., 2024; Korkiakoski et al., 2023), thermokarst (Jorgenson et al., 2025), and vegetation shifts (Frost et al., 2025). Moreover, global flux repositories often fail to include Arctic-boreal-specific variables, such as permafrost, high-latitude vegetation types, or lake origin (e.g. glacial or thermokarst lake). Additionally, CO₂ and CH₄ fluxes from freshwater and terrestrial ecosystems have typically been studied and synthesized separately, hindering a holistic understanding of the Arctic-boreal carbon cycle, which can also lead to double counting of carbon fluxes (Casas-Ruiz et al., 2023; Kyzivat and Smith, 2023; Thornton et al., 2016). Finally, data remain scattered across repositories, scientific publications, and unpublished sources, making it difficult to understand how comprehensive and representative the current network of Arctic-boreal flux measurements is.

To address these research gaps, we compiled a dataset of Arctic-boreal CO₂ and CH₄ fluxes in terrestrial and freshwater ecosystems (ABCFlux v2) from flux repositories, data syntheses, literature, and data contributors, which are presented here. We built upon recent syntheses (Golub et al., 2023; Kuhn et al., 2021; Song et al., 2024; Stanley et al., 2022; Virkkala et al., 2022) and earlier Arctic-boreal terrestrial and freshwater CO₂ and CH₄ flux datasets (Belshe et al., 2013; McGuire et al., 2012; Natali et al., 2019; Olefeldt et al., 2013; Treat et al., 2018; Wik et al., 2016b), including ABCFlux v1 (Virkkala et al., 2022). The structure of ABCFlux v2 follows

v1, which synthesized monthly terrestrial CO₂ fluxes. Compared to v1, ABCFlux v2 includes not only updated terrestrial CO₂ fluxes, but also terrestrial CH₄ fluxes. In addition, we expanded the dataset to freshwater ecosystems, including lentic (lakes, ponds, reservoirs, pools) and lotic (rivers and streams) waterbodies, and synthesized carbon fluxes and surface concentrations of dissolved CO₂ and CH₄. We also added several new variables to ABCFlux v2 from the BAWLD-CH₄ and GRiMeDB database to include variables specific to freshwaters (see section 3 for details) (Kuhn et al., 2021; Stanley et al., 2023).

ABCFlux v2 comprises several measurement techniques that provide different measurement frequencies across multiple ecosystem scales. Eddy covariance is a common method for measuring temporal dynamics in terrestrial carbon fluxes on ecosystem scales and quantifying year-round net carbon emissions. In many cases, eddy covariance data are actively shared, processed and curated in global (Pastorello et al., 2020) and regional (Heiskanen et al., 2022; Novick et al., 2018; Ueyama et al., 2025) flux networks. However, not all Arctic-boreal sites are part of these networks, and overall, the Arctic-boreal region has a particularly low coverage of eddy covariance towers (Pallandt et al., 2022), especially for lakes (Eugster et al. 2022, Golub et al. 2023). Furthermore, eddy covariance, which aggregates fluxes over ecosystem scales (hundreds of meters), often cannot resolve issues regarding local-scale spatial heterogeneity in emission and uptake driven by small-scale variation in vegetation, hydrology, soil microclimate (Chen et al., 2012; Virkkala et al., 2024). Moreover, CH₄ flux estimates derived from eddy covariance generally do not distinguish between the multiple CH₄ emission pathways (diffusion, ebullition, plant-mediated transport; but see (Ueyama et al., 2023)), which are important to understanding processes controlling the total CH₄ fluxes (Bastviken et al., 2004; Kyzivat et al., 2022). Thus, relying solely on eddy covariance towers is insufficient for a comprehensive understanding of Arctic-boreal carbon fluxes and the ability to predict current and future emissions more accurately, emphasizing the value of other kinds of flux measurements including small-scale, ground-based techniques.

Small-scale, ground flux techniques most often consist of static or automated chamber measurements in terrestrial and freshwater ecosystems and concentration-based turbulence-driven modeling approaches in freshwaters (Kuhn et al., 2021; Stanley et al., 2023; Virkkala et al., 2021). Chamber techniques can assess fluxes across small footprints (ca. 0.3 to 1 m²), allowing for detailed assessments of environmental controls on fluxes (Kuhn et al., 2021). Flux gradient approaches, wherein gas samples are taken from the air and throughout the soil or snow profile to estimate net flux, have also been used in some terrestrial sites (Pirk et al., 2016). In freshwaters, diffusive fluxes can also be estimated from measurements of gas concentrations dissolved in the surface water and using turbulence-driven modelling approaches based on gas transfer velocities (Klaus and Vachon, 2020; Vachon and Prairie, 2013). Ebullitive fluxes can be derived from concentration bursts during chamber measurements (Bastviken et al., 2004), but are most commonly assessed using bubble traps, which can be coupled with ice-bubble surveys to reduce spatial uncertainties (Huttunen et al., 2001; Walter Anthony and Anthony, 2013; Wik et al., 2013). For freshwaters, ground-based measurement techniques are also used to capture storage fluxes. Storage fluxes refer to the sudden diffusive efflux triggered by lake turnover in spring and fall, or by gas that accumulates under the frozen layer of a lake and is emitted to the

405 atmosphere when the ice melts in the spring (Jammet et al., 2015). Storage fluxes are measured
 from the difference between measured waterbody content of dissolved gas before and after the
 turnover or ice-off (Karlsson et al., 2013). While these micro-scale approaches advance the
 understanding of local processes and spatial variability in carbon fluxes, they are also
 410 accompanied by uncertainties due to potentially limited spatial and temporal representativeness,
 the disturbance that collars, floating bubble traps, and chambers can cause on the ground or
 water surface (Welles et al., 2001), oxidation in the water column prior to ice out (Pajala et al.,
 2023), the wide range of available methods to determine gas transfer velocities (Hall and
 Ulseth, 2020; Klaus and Vachon, 2020; Raymond et al., 2012), and the temporal
 representativeness of the manual sampling campaigns (Golub et al., 2023; McGuire et al., 2012;
 415 Wik et al., 2016). Automated chambers and continuous concentration measurements in
 terrestrial and aquatic ecosystems, provide more temporally representative sampling relative to
 those from more limited manual sampling campaigns, but do not solve for potential artifacts
 derived from ground disturbances. Overall, combining all fluxes measured with these different
 techniques (Table 1) is an important benefit of ABCFlux v2 compared to other efforts focused on
 420 a single flux measurement technique or gas species.

In this community-driven effort, we integrated surface-atmosphere CO₂ and CH₄ fluxes into a
 single, unified Arctic-boreal-specific dataset. Below, we provide a description of the dataset and
 a summary of the flux network, and synthesize flux magnitudes across key land cover types,
 spanning both terrestrial and freshwater ecosystems.

425 **Table 1.** A summary of the measurement techniques, carbon flux and concentration
 observations, and key ecosystems included in ABCFlux v2. Terrestrial classes include dry and
 moist tundra, bogs, fens, marshes, tundra wetlands, permafrost bogs, and boreal forest
 ecosystems. Freshwater classes include lentic and lotic waterbodies. Storage flux refers to the
 transient accumulative release of gasses during ice-out and water column mixing events. We do
 430 not include plant mediated aquatic fluxes of CH₄ due to a lack of data available in this region.
 Percentages represent the percent of site-months from the respective ecosystem (i.e. % of
 terrestrial site-months or % of freshwater site-months). Percentages for the aquatic ecosystems
 do not add up to 100 % due to overlaps, for example, where both diffusion fluxes and dissolved
 concentrations were measured.

Measurement technique	Terrestrial	Freshwater
Eddy covariance	Eddy covariance (CO ₂ , CH ₄ flux) (64.4%)	Eddy covariance (CO ₂ , CH ₄ flux) (4.6 %)
Non-eddy covariance	Manual or automated chamber (CO ₂ , CH ₄ flux) (33.9%)	Diffusion based on chambers and turbulence-driven modeling derived from concentrations (CO ₂ , CH ₄ flux) (42.8 %)
	Flux gradient approach (CO ₂ flux) (1.7%)	Ebullition based on bubble traps or chambers (CO ₂ , CH ₄ flux) (4.7 %)

		Storage flux: Water column survey (CO ₂ , CH ₄ flux) (3.6 %) <i>Note: Total CH₄ flux = CH₄ diffusion + CH₄ ebullition</i>
Dissolved gas concentration		Dissolved concentration at the water surface (CO ₂ , CH ₄) (86.4 %)

435

2. Data compilation and search

ABCFlux v2 focuses on the Arctic tundra and boreal biomes, as characterized in Dinerstein et al. (2017; Figure 1). It also includes some hemiboreal sites located within 500 km south of the boreal biome boundary, when data were available through public repositories or provided by data contributors (~2.5% of sites). These sites were included because hemiboreal ecosystems share key characteristics with boreal systems and may provide insight into potential trajectories of boreal ecosystems under changing climate (Berner and Goetz, 2022). The dataset compiles in situ measured CO₂ and CH₄ fluxes aggregated to monthly time periods (unit: g C m⁻² per month, i.e. g CO₂-C m⁻² per month for CO₂ fluxes and g CH₄-C m⁻² per month for CH₄ fluxes) from terrestrial and freshwater ecosystems, including boreal forests, wetlands, tundra, lentic and lotic waterbodies. In forested ecosystems, chamber CO₂ measurements were excluded as they typically do not represent the whole ecosystem fluxes (i.e. chamber measurements exclude trees). However, for CH₄, we accepted measurements of understory CH₄ fluxes as the CH₄ fluxes from trees are expected to be minimal (see Sect. 8.3 for challenges associated with this assumption). For freshwater ecosystems, we also included monthly average CO₂ and CH₄ concentrations dissolved in surface waters because this information helps to understand flux dynamics and can also be used to estimate fluxes (e.g. (Holgerson and Raymond, 2016)). We excluded flux data from experimental manipulation sites, with the exception for control sites within manipulations experiments. We included data from managed forests but excluded croplands, as croplands typically undergo intensive annual management (e.g., tillage, fertilization, and harvesting). In contrast, managed forests are generally managed on decadal timescales, which allows them to retain some functional characteristics of natural ecosystems, particularly in later stages of regrowth.

We used a monthly aggregation interval as it is a common and standard temporal frequency across many site-level, synthesis, and modeling studies, remote sensing products, and process models. However, there were some seasonally-aggregated data from previous syntheses and studies that we were not able to incorporate in a monthly format; these were kept in the dataset in seasonal format alongside the start and end date of the measurement period. Seasonal fluxes make up only <1% of the dataset. Monthly fluxes were primarily found derived by multiplying daily means (g C m⁻² d⁻¹) by the number of days in each month to calculate monthly

cumulative fluxes ($\text{g C m}^{-2} \text{ month}^{-1}$), although methods varied based on available data and temporal resolution (see section 5.1).

The data compilation steps are detailed in a flow chart (Fig. 2). We compiled and harmonized data from syntheses, global and regional flux repositories and general data repositories (Table 2), publications, and direct submissions from data contributors. In cases where data for the same sites and periods were available from multiple sources, we prioritized user-contributed data over data extracted from repositories, syntheses, and publications. This prioritization was chosen due to the benefits associated with the expertise of data contributors with data processing at their site (e.g., gap-filling), and the inclusion of ancillary data. There are no duplicate observations in the main dataset, i.e. only one flux estimate per flux type is given for each site-month, even though there may be multiple different data sources.





Fig. 1. Numbers of site-months for terrestrial CO₂ and CH₄ flux sites, and aquatic CO₂ and CH₄ flux sites across the Arctic-boreal region. The number of months represented by circles refers to

total, not necessarily consecutive months. See Supplementary Fig. 8 for zoomed-in maps for the densely measured areas.

Fig. 2. Flowchart representing the main data compilation steps to produce ABCFlux v2. Note that repositories are listed based on the direct source of data acquisition; for example,



terrestrial NEON fluxes are included via the AmeriFlux repository, whereas aquatic gas concentrations were sourced directly from NEON.

2.1 Recent data syntheses

We incorporated data from various data syntheses into our dataset (Table 2). We integrated terrestrial CO₂ flux data included in ABCFluxv1 into our v2 dataset with some modifications to the original dataset. Notably, GPP values in v1 were reported as negative, whereas in v2 they are presented as positive to align with the convention used throughout this synthesis. We removed soil respiration data from forest floors as our focus here was on whole-ecosystem CO₂ fluxes. Some monthly chamber fluxes within the v1 dataset were spatial replicates (same coordinates and land cover) and for the purposes of v2, we aggregated these by taking a mean flux. We updated the soil moisture classification for several sites as well as several site names

from v1 to be consistent with site names used in v2. Additionally, some data from v1 were replaced by more recent versions of the data found in flux repositories.

The terrestrial and aquatic CH₄ chamber flux data from the BAWLD-CH₄ dataset (Kuhn et al., 2021) were originally presented as average daily fluxes over the growing season (wetlands and uplands) and open-water season (lentic ecosystems) for each site, with the last year of data collection being 2019 (Table 2). For sites where data at monthly resolution were available in publications, we extracted monthly flux and supporting environmental data. If monthly flux or other ancillary data were not extractable from the literature, we reached out to the lead author for data contribution.

Data in the Global River Methane Database (GRiMeDB, Table 2; (Stanley et al., 2023)) were aggregated to monthly resolution in ABCFlux v2. Fluxes in the global CO₂ lake and reservoir synthesis (Golub et al., 2023) were presented in half-hourly timesteps (Table 2), but were aggregated to monthly resolution. Dissolved gas concentrations in rivers from Liu et al. (2022) were converted to monthly averages. Overall, a significant portion of data (30%) incorporated in ABCFlux v2 originated from these terrestrial and aquatic data syntheses.

Table 2. Global and regional flux syntheses and repositories used in the data compilation of terrestrial and/or aquatic data. Information about the ecosystem, the name, reference or web page, flux method, gas species, temporal resolution, spatial extent and period of data coverage of the respective synthesis or repository are given.

Ecosphere	Type	Ecosystem	Name of repository or synthesis	Reference or web page (last access: August 4, 2025)	Flux method	Gas	Temporal resolution	Spatial extent	Data coverage
Both	Synthesis	Wetland, Lentic	BAWLD-CH ₄	Kuhn et al. (2021)	Mostly chamber	CH ₄	Seasonal	Arctic-boreal	1984-2020
Terrestrial	Synthesis	Terrestrial	ABCFlux v1	(Virkkala et al., 2022)	Eddy covariance, chamber	CO ₂	Monthly	Arctic-boreal	1989-2020
Terrestrial	Repository	Terrestrial	FLUXNET2015	(Pastorello et al., 2020); https://fluxnet.org/data/fluxnet2015-dataset/	Eddy covariance	CO ₂	Half-hourly to yearly	Global	1994-2014
Terrestrial	Repository	Terrestrial	FLUXNET-CH ₄	(Delwiche et al., 2021); https://fluxnet.org/data/fluxnet-ch4-community-product/	Eddy covariance	CH ₄	Half-hourly to daily	Global	2006-2018
Terrestrial	Repository	Terrestrial	Ameriflux	(Chu et al., 2023);	Eddy covariance	CO ₂ , CH ₄	Half-hourly to yearly	Americas	1994-2021

				https://ameriflux.lbl.gov/					
Terrestrial	Repository	Terrestrial	ICOS	(Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre, 2020); https://www.icos-cp.eu/	Eddy covariance	CO2, CH4	Half-hourly to yearly	Europe	1996-2023
Terrestrial	Repository	Terrestrial	JapanFlux2024	Ueyama et al. (2025)	Eddy covariance	CO2, CH4	Half-hourly to yearly	Japan and East Asia	1990-2023
Terrestrial	Repository	Terrestrial	European Fluxes Database Cluster	(Valentini, 2003); https://www.europe-fluxdata.eu/	Eddy covariance	CO2	Daily to monthly	Europe	1996-2008
Terrestrial	Repository	Terrestrial	Arctic Data Center	https://arcticdata.io/	Eddy covariance, chamber	CO2, CH4	Varies	Global	Varies
Terrestrial	Repository	Terrestrial	Zenodo	https://zenodo.org/	Chamber	CO2, CH4	Varies	Global	Varies
Terrestrial	Repository	Terrestrial	Next Generation Ecosystem Experiments	https://ngee-arctic.ornl.gov/	Chamber	CO2, CH4	Varies	USA	Varies
Terrestrial	Repository	Terrestrial	EMERGE-DB	https://emerge-db.asc.ohio-state.edu/	Chamber	CO2, CH4	Varies	Sweden	Varies
Aquatic	Synthesis	Lentic		Golub et al. (2023)	Eddy covariance	CO2	Half-hourly	Global	2005-2015
Aquatic	Synthesis	Lotic	GRiMeDB	Stanley et al. (2023)	Chamber, concentration	CO2, CH4	Daily to seasonal	Global	1973-2021
Aquatic	Synthesis	Lotic		Liu et al. (2022)	Concentration	CO2	Daily to seasonal	Global	
Aquatic	Repository	Lentic, lotic	NEON	https://www.neonscience.org/	Concentration	CO2, CH4	Half-hourly	US	2016-2022
Aquatic	Repository	Lentic	Bolin	https://bolin.su.se/data	Chamber, concentration, ebullition	CO2, CH4	Varies	Global	Varies
Aquatic	Repository	Lentic, lotic	PANGAEA	https://doi.org/10.1594/PANGAEA.919986	Concentration	CH4	Varies	Global	Varies
Aquatic	Repository	Lentic, lotic	SITES	https://www.fieldsites.se/	Chamber, concentration	CO2	Varies	Global	Varies

515 2.2 Data repositories

We obtained a majority of terrestrial eddy covariance data from flux network repositories listed in Table 2. Because aquatic ecosystem eddy covariance data were sparsely available through these flux networks, we relied on user-contributed data and those published in Golub et al., 2023. The repository data were downloaded between August 2023 and January 2024.

520 The terrestrial data that we downloaded from these repositories came in a variety of formats, though we gave preference to CO₂ flux data processed with the ONEFlux pipeline (Pastorello et al., 2020) and CH₄ flux data processed as part of the FLUXNET-CH₄ community product (Delwiche et al., 2021) when available since these data were gap-filled and quality-checked (i.e., FLUXNET2015, ICOS, and some of the Ameriflux datasets). This decision was made
525 because of the strengths associated with the consistent data processing, quality-control, and recent updates. The ONEFlux pipeline produces datasets aggregated to different time resolutions (half-hourly, daily, monthly, and yearly) along with fluxes processed with various partitioning methods and friction velocity (USTAR) criteria, to remove data under low turbulence conditions (i.e., turbulence filters to correct biases). However, the ONEFlux pipeline does not
530 perform footprint partitioning or fetch screening based on wind direction. Therefore, we assumed that any such filtering (e.g., for wind direction or land cover representativeness) was performed by the data provider of the site prior to ONEFlux processing, or that the reported fluxes represent the entire tower footprint. We used the datasets pre-aggregated to monthly mean fluxes ($\text{g C m}^{-2} \text{ d}^{-1}$) and opted for fluxes processed with a constant USTAR threshold and
535 flux partitioning according to Reichstein et al. (2005) when available (for a justification, see Virkkala et al., 2022). To calculate gap-fill percentages for each month, we used the half-hourly datasets produced by the ONEFlux pipeline and the quality flag associated with each flux (QC= 0 measured value).

For terrestrial eddy covariance data not processed with ONEFlux, we prioritized gap-filled data,
540 though where it was not available, we accepted data that were not gap-filled (10% of all the monthly eddy covariance data). The level of pre-processing (i.e. USTAR filtering, storage correction, etc) of this non-gap-filled data varied by data source but all data were quality checked prior to monthly aggregation. We justified this approach to increase the amount of data in this data-sparse region, and carefully assessed that the aggregated monthly fluxes were
545 within a realistic range (within minimum and maximum monthly fluxes in similar environments).

We also searched terrestrial and aquatic data through several data repositories not focused solely on fluxes (see Table 2). To identify datasets of interest we used the same search words in the repositories as in the literature search (see Section 2.3). If the datasets identified from this search had associated publications, we reviewed the publications and extracted relevant
550 information including that describing the environmental conditions of the study site.

2.3 Literature search

We conducted an exhaustive *Web of Science* search with search words (“carbon flux” or “carbon dioxide flux” or “methane flux” or “CH₄ flux” or “CO₂ flux” or “NEE” or “net ecosystem

exchange”) and (“arctic” or “tundra” or “boreal”) up to December, 2023. For aquatic data, this search was extended up to August 2024. Based on the literature search, we added data from three additional publications for terrestrial ecosystems focusing on chambers and 105 for aquatic ecosystems beyond those already included in BAWLD-CH₄, GRiMeDB, Liu et al., 2022 and Golub et al., 2023. The total number of new papers identified for terrestrial ecosystems was low because the recent BAWLD-CH₄ and ABCFluxv1 datasets already encompass studies up to 2018–2020, and more recent data were sourced from repositories or submitted directly by data contributors.

2.4 Community-contributed data

In order to capture data that are not easily extractable from literature nor found in repositories, we contacted ~180 researchers that were identified based on earlier reviews and syntheses, and our expert knowledge. This call for data began in April 2023, with final data submissions due in June 2024, though most contributions were given prior to January 2024. Out of the ~180 researchers contacted, we received 98 terrestrial and 43 aquatic datasets. These user-contributed flux data constitute 42% of observations in the overall dataset (47% of terrestrial, 32% of aquatic, 39% of CO₂, and 51% of CH₄). In addition to flux data, submissions from data contributors were more likely to include detailed site descriptions and data for ancillary variables that were not often available in repositories or papers. Submitted data from 23 new sites were previously unpublished (i.e., not published in scientific papers; 4.1% of site-months for terrestrial and 2.7% for aquatic) but had been processed using standard processing protocols or similar tools used at the site before.

3. Data columns

3.1 Summary of data columns

The ABCFlux v2 dataset is organized such that each row represents a unique combination of site and month. The data is grouped by site and arranged chronologically. To facilitate ease of use, the dataset is provided as two separate files based on ecosystem type. The aquatic file (including lentic and lotic systems) contains 103 variables while the terrestrial file (including boreal forest, wetlands, and tundra) contains 93 variables. All columns included in ABCFlux v2 are listed in Table 3, together with the percentage of data in each column, which has been subset by applicable data types (e.g. soil temperature applies only to terrestrial data). Across both files, there are a total of 141 columns in ABCFlux v2 including 15 that contain flux data and 12 pertaining to measurement technique details (e.g. flux method details, partition method) and data quality (e.g. gap-fill percentage, number of chamber measurement days).

For eddy covariance measurements in terrestrial ecosystems, net ecosystem exchange of CO₂ (NEE) can be partitioned into gross primary productivity (GPP) and ecosystem respiration (Reco). In this dataset we include NEE as well as GPP and Reco, when available but we did not perform any data processing or flux partitioning, aside from unit conversions. Throughout this dataset, we report GPP and Reco as positive. NEE and CH₄ fluxes are reported with respect to the atmosphere, where positive values are a net source to the atmosphere from the ecosystem,

and negative values are a net sink from the atmosphere to the ecosystem. For aquatic ecosystems, only the overall (non-partitioned) CO₂ flux is given because of the multiple origins of the CO₂ (Battin et al., 2023). The partitioning of GPP from Reco in aquatic systems is usually done with other methods (i.e. as commonly done with oxygen mass balances; (Staeher et al., 2010), but are not included in this dataset. For freshwater CH₄ fluxes derived with methods other than eddy covariance, we differentiate between diffusive and ebullitive emission pathways and assume the sum of diffusion and ebullition to yield the total CH₄ flux. We also compiled CO₂ and CH₄ storage fluxes (as done for CH₄ by Kuhn et al., 2021 and Wik et al. 2016). Partitioning plant-mediated carbon fluxes from freshwaters is rare and not considered here.

In addition to carbon fluxes and dissolved gas concentrations, we gathered information describing environmental conditions of the site as well as general site characteristics. There are 48 variables pertaining only to aquatic data (e.g. waterbody depth, water temperature) and 38 tailored to terrestrial data (e.g. vegetation information, soil temperature). Fifty three of the columns represent static variables (e.g., biome, land cover), whereas the remaining columns vary monthly (e.g., fluxes, soil temperatures), seasonally (seasonal flux if monthly was not available), or annually (active layer depth). New columns were added to v2 compared to v1 (Virkkala et al., 2022) to represent detailed descriptions of plant functional types (e.g., evergreen and deciduous shrub coverage), deep soil temperatures (<10 cm), and permafrost thaw presence or absence in the top two meters. We also added new columns pertaining to CH₄ fluxes that were not included in previous flux synthesis efforts, such as a categorical moisture class (wet-moist-dry) and BAWLD classes (Olefeldt et al., 2021).

To differentiate between terrestrial and freshwater ecosystem classes, we followed the BAWLD classification system (Olefeldt et al., 2021) and specified the classes in column “bawld_class”. BAWLD was specifically designed to separate key classes relevant for CH₄ cycling. There are 8 terrestrial classes and 10 freshwater classes listed in the variable descriptions in Table 3, and the classes are more thoroughly discussed in their respective papers and metadata documents (Kuhn et al., 2021; Olefeldt et al., 2021). The column “bawld_class” was designed to differentiate between plot-level (sub-meter) variability in land cover types. Eddy covariance sites that had highly heterogeneous footprint with multiple BAWLD landcover classes were assigned the dominant BAWLD class for the tower footprint and should be interpreted with caution given the different landscape classification scales. When assigning lotic classes, we followed the BAWLD river size distinction (large rivers have a Strahler order >5, small rivers have a Strahler order ≤5). We first used the description provided by data contributors to determine if a small river was organic-poor or organic-rich. If the data contributors did not provide a site description, we deferred to the organic carbon geospatial data by Hugelius et al. (2020). We selected a 20% organic soil coverage threshold for this split based on a comparison with a small dataset containing site-level classifications of organic-poor and organic-rich soils, combined with our expert knowledge and cross-checked with the gridded percent cover data for rivers from BAWLD. However, since most lotic sites were ultimately classified using geospatial rather than

site-level data, these classifications are inherently more uncertain. In some cases where classification was unclear, the site was left as 'Unknown'.

635 In addition to the BAWLD classes, and to acknowledge classes relevant to CO₂ fluxes, we used
a classification system from the European Space Agency (ESA) Climate Change Initiative (CCI)
land cover product (ESA CCI 2016) which has been used in earlier syntheses (Virkkala et al.,
2021). It differentiates between six different boreal forest/vegetation classes instead of the
broad Boreal forest class in BAWLD. There are in total 20 terrestrial classes and one general
aquatic class in ESA CCI, which are listed in the column descriptions in Table 3 and are more
640 thoroughly discussed in their respective metadata document (ESA CCI 2016). For the ESA CCI
class, we created two columns: "land_cover_plot" for the plot level and "land_cover_eco" for the
ecosystem level, to acknowledge the extent and scale of the land cover type associated with
each measurement. For example, a dry shrub-dominated plot at a palsamire received a shrub
class at the plot level, while the ecosystem-level class was water-logged, characterizing the
645 mostly wetland-dominated status of the ecosystem. For both the BAWLD and ESA CCI
columns, the categories were defined by data contributors or extracted from papers and
repositories and then unified by dataset developers through an expert assessment utilizing
additional columns (e.g., plant cover and vegetation description for BAWLD and ESA CCI
terrestrial classes, and lake size and sediment type for BAWLD aquatic classes). If no
650 information was available, the class was left as "NA".

Table 3. Variable names and their description, and the percentage of data present. Percentages were calculated based on the total relevant data for each variable. Subset refers to the data category the respective variable applies to: eddy covariance (EC), and non-eddy covariance (Non-EC) including all other measurement methods (chambers, concentration, etc.). Where the subset remains blank, the variable applies to the whole dataset.

655

Variable category	Subset	Variable name	Unit	Description	Percent data present
Metadata		site_name		Site name as specified in data source. E.g. Hyytiälä	100.0
		site_reference		A more specific name used in data source. E.g. the name of the chamber plot (e.g. shrub) or the abbreviation used in eddy covariance data repositories (e.g. FI-Hyy).	100
		data_contributor_or_author		Data contributor(s) or primary author(s) associated with data set or publication	100
		email		Primary author email(s)	72.9
		extraction_source		Data source where data were extracted and compiled.	100
		citation		Citation for the data source: journal article, data citation, and/or other source (online repository link etc.). If the user contributed unpublished data, a journal article citation describing already published data can be added here.	90.9
		country		Country of the study site	100
		latitude	decimal degrees	Latitude of study site. Note that spatial accuracy is subject to the limitations of the primary data sources and may vary across the dataset.	100
		longitude	decimal degrees	Longitude of study site. Note that spatial accuracy is subject to the limitations of the primary data sources and may vary across the dataset.	100
		year	YYYY	Year in which data were recorded	99.8
		month	MM	Measurement month	99.2
Fluxes	Terrestrial	nee	g C m ⁻² month ⁻¹	Monthly cumulative net ecosystem exchange (NEE) for the entire measurement interval in g C in CO ₂ . Negative flux represents a net sink to the ecosystem.	78.5

	Terrestrial	gpp	g C m-2 month-1	Monthly cumulative gross primary productivity (GPP) for the entire measurement interval in g C in CO ₂ . Note: GPP is presented as positive (uptake) values.	61.2
	Terrestrial	reco	g C m-2 month-1	Monthly cumulative Ecosystem respiration (Reco) for the entire measurement interval in g C in CO ₂	68.5
	Aquatic	co2_flux	g C m-2 month-1	Monthly cumulative CO ₂ flux for the entire measurement interval. Refers to aquatic fluxes only where fluxes are commonly not partitioned between photosynthesis and respiration. Negative flux represents a net sink.	36.6
		ch4_flux_total	g C m-2 month-1	Monthly cumulative total CH ₄ flux for the entire measurement interval. Negative flux represents a net sink to the ecosystem.	27.7
	Aquatic-Non-EC	co2_flux_ebullition	g C m-2 month-1	Monthly cumulative ebullitive CO ₂ flux for the entire measurement interval	0.9
	Aquatic-Non-EC	ch4_flux_ebullition	g C m-2 month-1	Monthly cumulative ebullitive CH ₄ flux for the entire measurement interval	12.5
	Aquatic-Non-EC	ch4_flux_diffusion	g C m-2 month-1	Monthly cumulative diffusive CH ₄ flux for the entire measurement interval	93.6
	Aquatic-Non-EC	ch4_flux_storage	g C m-2 month-1	Monthly cumulative storage CH ₄ flux for the entire measurement interval. Storage flux is a diffusion flux but can be measured separately and is separated here because of its non-continuous rapid burst emission nature.	9.7
	Aquatic-Non-EC	co2_flux_storage	g C m-2 month-1	Monthly cumulative storage CO ₂ flux for the entire measurement interval. Storage flux is a diffusion flux but can be measured separately and is separated here because of its non-continuous rapid burst emission nature.	0.1
	Terrestrial	nee_seasonal	g C m-2 season-1	Cumulative NEE flux in a seasonal format (e.g. June 3rd to August 25th). This column is filled only if monthly data are not available. Negative flux represents a net sink.	0.4

	Aquatic	co2_flux_seasonal	g C m-2 season-1	Cumulative CO2 flux from freshwater systems in a seasonal format (e.g. June 3rd to August 25th). This column is filled only if monthly data are not available. Negative flux represents a net sink.	0.5
		ch4_flux_total_seasonal	g C m-2 season-1	Cumulative CH4 total flux in a seasonal format (e.g. June 3rd to August 25th). This column is filled only if monthly data are not available.	0.9
	Aquatic-Non-EC	ch4_flux_ebullition_seasonal	g C m-2 season-1	Cumulative CH4 flux from ebullition in a seasonal format (e.g. June 3rd to August 25th). This column is filled only if monthly data are not available.	0.5
	Aquatic-Non-EC	ch4_flux_diffusion_seasonal	g C m-2 season-1	Cumulative CH4 flux from diffusion in a seasonal format (e.g. June 3rd to August 25th). This column is filled only if monthly data are not available.	0.1
Meteorological		tair	°C	Monthly mean air temperature	53.7
		tair_height	m	Height of the air temperature measurement	23.5
		precip	mm	Monthly cumulative precipitation mm	35.5
		snow_depth	cm	Mean snow depth during the measurement interval	8.2
		ppfd	μmol m-2 s-1	Mean photosynthetically active radiation during measurement interval (in Photosynthetic Photon Flux Density, PPFD)	29.1
	Terrestrial	tsoil_surface	°C	Monthly mean surface soil temperature in ca. 0-10 cm depth	77.5
	Terrestrial	tsoil_surface_depth	cm	Depth of the surface soil temperature measurement	58.5
	Terrestrial	tsoil_deep	°C	Monthly mean deeper soil temperature at > 10 cm depth	25.7
	Terrestrial	tsoil_deep_depth	cm	Depth of the deeper soil temperature measurement	29.0

	Terrestrial	soil_moisture	VWC %	Monthly mean surface soil moisture in ca. 0-10 cm depth	45.4
	Terrestrial	moisture_depth	cm	Depth for the surface soil moisture measurement	30.4
	Terrestrial	thaw_depth	cm	Mean thaw depth during the measurement interval. Positive values represent depth below the soil surface.	8.3
	Terrestrial	alt	cm	Active layer thickness (maximum thaw depth). Thickness changes annually. Positive values represent depth below the soil surface.	14.4
	Terrestrial	water_table_depth	cm	Mean water table depth during the measurement interval; positive is below the surface, negative is above (inundated)	17.4
Measure ment details		flux_method	EC, Chamber, Snow diffusion, Ebullition, Concentration, Concentration (no flux)	Broad categories for how flux values were measured, may list more than one	100
		flux_method_detail		Details related to how flux values were measured. Specifies between manual vs. automated chambers, snow diffusion, open vs. closed-path eddy covariance, water sampling, ebullition trap, ice sampling	96.5
		flux_method_description		Details related to measurement method, e.g. chamber size/volume and deployment time, tubing length, use of bubble shield or not, height of the tower and wind measurements.	69.5
		gap_fill		Approach used to gap-fill the data	85.7
	Terrestrial	partition_method		Method used to partition NEE into GPP and RECO.	64.9
	EC	tower_corrections		Details related to processing corrections employed, including time, duration, and thresholds for u^* and heat corrections	75.9

	Chamber	diurnal_coverage	Day, Day and Night	Indicator whether data was collected during the day or during day and night	91.2
		instrumentation		Description of instrumentation used (e.g. type of greenhouse gas analyzer)	49.2
	EC- CO ₂	gap_fill_perc_co2	%	% of eddy covariance or automated chamber CO ₂ (nee, co2_flux) data that was gap-filled in the measurement interval (relative to standard measurement time step)	73.5
	EC-CH ₄	gap_fill_perc_ch4	%	% of eddy covariance or automated chamber CH ₄ data that was gap-filled in the measurement interval (relative to standard measurement time step)	29.9
	Non-EC-CO ₂	chamber_nr_measurement_days_co2		Number of days with chamber measurement of CO ₂ within the measurement interval (month)	42.7
	Non-EC-CH ₄	chamber_nr_measurement_days_ch4		Number of days with chamber measurements of CH ₄ within the measurement interval (month)	50.8
	Aquatic Non-EC-CO ₂	ebullition_nr_measurement_days_co2		Number of days with ebullition measurements of CO ₂ within the measurement interval (month)	0.4
	Aquatic Non-EC-CH ₄	ebullition_nr_measurement_days_ch4		Number of days with ebullition measurements of CH ₄ within the measurement interval (month)	0.6
	Terrestrial	nee_seasonal_interval	MM/DD/YYYY-MM/DD/YYYY	The period for the seasonal estimate (e.g. 06/04/2015-08/25/2015)	0.4
	Aquatic	co2_flux_seasonal_interval	MM/DD/YYYY-MM/DD/YYYY	The period for the seasonal estimate (e.g. 06/04-08/25)	0.5
		ch4_flux_seasonal_interval	MM/DD/YYYY-MM/DD/YYYY	The period for the seasonal estimate (e.g. 06/04-08/25)	1.1
	CO ₂	expert_flag_co2	0,1,2,3,4	0: no known issues, 1: terrestrial fluxes outside the 1st and 99th percentiles, aquatic lentic fluxes outside the 99th percentile, 2: terrestrial non-eddy covariance growing season measurements with 3 or less measurement days in the month and no	73.5

				modeling used to gap-fill, 3: terrestrial eddy covariance data with 3 or more consecutive months of 100% gap-filling, 4: site does not represent typical environmental conditions of the Arctic Boreal Zone	
	Terrestrial-CO ₂	expert_flag_gpp	0,1	0: no known issues, 1: terrestrial fluxes outside the 1st and 99th percentiles	70.9
	Terrestrial-CO ₂	expert_flag_reco	0,1	0: no known issues, 1: terrestrial fluxes outside the 1st and 99th percentiles	79.4
	CH ₄	expert_flag_ch4	0,1,2,3,4	0: no known issues, 1: terrestrial fluxes greater than 30 g C m ⁻² month ⁻¹ , aquatic lentic fluxes outside the 99th percentiles, 2: terrestrial non-eddy covariance growing season measurements with 3 or less measurement days in the month and no modeling used to gap-fill, 3: terrestrial eddy covariance data with 3 or more consecutive months of 100% gap-filling, 4: site does not represent typical environmental conditions of the Arctic Boreal Zone	66.8
Site information		biome	Tundra, Boreal, Temperate	Biome of the study site	100
		land_cover_eco	40=Mosaic natural vegetation (>50%) / cropland (<50%); 60=Tree cover, broadleaved deciduous; 70=Tree cover, needleleaved evergreen; 80=Tree cover, needleleaved deciduous; 90=Tree cover, mixed leaf type;	Dominant land cover class for the site following expert assignment and using class names of the ESA CCI land cover product. http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf (section 9.1, global classification, page 81)	99.8

			100=Mosaic tree & shrub (>50%) / herbaceous (<50%); 110=Mosaic herbaceous (>50%) / tree & shrub (<50%); 120=Shrubland; 121=Shrubland evergreen; 122=Shrubland deciduous; 130=Grassland; 140=Lichens & mosses; 150=Sparse vegetation (<15%); 151=Sparse tree; 152=Sparse shrub; 153=Sparse herbaceous; 160=Tree cover, flooded fresh/brackish; 170=Tree cover, flooded saline; 180=Shrub/herbaceous flooded; 200=Bare areas; 210=Waterbodies		
		land_cover_plot	Same classes used as for "land_cover_eco" (see above)	Dominant land cover class for the site following expert assignment and using class names of the ESA CCI land cover product. http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf (section 9.1, global classification, page 81)	99.8
		bawld_class	Bog, Fen, Marsh, Permafrost Bog, Wet Tundra, Dry Tundra, Moist Tundra, Boreal Forest, Rocklands, Large Lake,	Dominant ecosystem class for the site following Boreal-Arctic Wetland-Lake Database (BAWLD) classes (Olefeldt et al. 2021). https://essd.copernicus.org/articles/13/51/27/2021/	100

			Midsize Glacial Lake, Small Glacial Lake, Midsize Peatland Lake, Small Peatland Lake, Midsize Yedoma Lake, Small Yedoma Lake, Large River, Small Organic-Rich River, Small Organic-Poor River		
		veg_detail		Detailed vegetation description from data source/contributor, incl. dominant species and a description of vegetation changes at the site. In an aquatic site, this should have information on both the terrestrial and aquatic plants.	57.5
		permafrost	Yes, No	Reported presence or absence of permafrost	68.2
		permafrost_thaw	Yes, No	Permafrost thaw present	32.0
		thaw_category	Active layer thickening with subsidence, Active layer thickening without subsidence, Active layer thickening with talik formation, Ice-wedge degradation, Thaw lake/pond expansion, Wetland collapse, Active layer detachment sliding, Retrogressive thaw slumping	Broad category for types of thaw	14.6
		thaw_extent	0-33%, 34-66%, 67-100%	Spatial extent of thaw within the site (i.e. tower footprint, chamber/sampling plot, drainage basin for aquatic ecosystems)	16.2

		thaw_severity	Low: Change to the ecosystem that is temporary or does not have substantial impact on ecosystem identity or function, Moderate: Change to the ecosystem that is temporary but has a substantial impact on ecosystem function, High: Change to the ecosystem that is lasting with consequences for ecosystem identity and function	Severity of dominant type of thaw following the classifications of ecosystem change presented in (Webb et al., 2025)	14.6
		thaw_dominant	Active layer detachment, Active layer thickening, Active layer thickening/Thaw Ponds, Gradual thaw, Palsa thaw, Retrogressive thaw slumps, River bank erosion, Subsidence, Thaw lake, Thermokarst, Thermokarst (pond), Thermokarst bog formation, and Upland thermokarst mounds with talik	Dominant type of thaw. If multiple categories were chosen in thaw_category, thaw_extent and thaw_severity primarily refer to the type of thaw listed here	14.2
		landform		Description of the geomorphological landforms associated with the site. E.g., polygonal features, palsas, cryoturbation, abrupt thaw features, drained lake basins.	14.6

		disturbance		Description of the recent disturbance history of the site or list "No" if there are no disturbances. Note that the disturbance might have been caused due to natural reasons (e.g. fire) and/or due to anthropogenic influences (e.g. drained peatland, harvested forest). If several disturbances have occurred during the recent decades, these can all be listed here but please list the last dominant disturbance first.	54.8
		disturbance_category	Altered hydrology, Animal herbivory, Artificial pond, Beavers, Drained lake, Drainage, Erosion, Extreme weather, Fire, Forestry, Human paths, Insect herbivory, Land use change, None, Other, Peat mining, Reservoir, Roads, Seismic lines, Thaw, Tidal effects, Wastewater	Broad categories for disturbances	77.4
		disturb_dominant	Animal herbivory, Artificial pond, Beavers, Drained lake, Drainage, Erosion, Erosion/Thaw, Extreme weather, Fire, Forestry, Human paths, Insect herbivory, Land use change, None, Other, Reservoir, Roads, Seismic lines, Thaw, Wastewater	Dominant disturbance of the site. If multiple categories are listed in disturbance category, disturb_extent and disturb_severity primarily refer to the disturbance listed here	20.8
		disturb_year	Numeric variable (year), 0 = annual (e.g., annual	Year of last dominant disturbance, 0-[year] indicates the disturbance is ongoing and began in the year given	22.3

			grazing, annual nutrient additions)		
		disturb_severity	Low: Change to the ecosystem that is temporary or does not have substantial impact on ecosystem identity or function, Moderate: Change to the ecosystem that is temporary but has a substantial impact on ecosystem function, High: Change to the ecosystem that is lasting with consequences for ecosystem identity and function	Relative severity of last dominant disturbance following the classifications of ecosystem change presented in Webb et al. 2025	19.7
		disturb_extent	0-33%, 34-66%, 67-100%	Spatial extent of the dominant disturbance within the site (i.e. tower footprint, chamber/sampling plot, drainage basin for aquatic ecosystems)	14.6
		site_activity	Active, Non-active	The current flux measurement activity status of the site (i.e., measurements conducted each year).	72.6
Terrestrial variables	Terrestrial	dec_shrub	Absent, Present, Dominant	Are deciduous shrubs absent, present, or dominant? Examples of deciduous shrubs: Betula nana, Salix sp, Vaccinium uliginosum, Vaccinium myrtillus, Rubus chamaemorus	46.3
	Terrestrial	ev_shrub	Absent, Present, Dominant	Are evergreen shrubs absent, present, or dominant? Examples of evergreen shrubs: Empetrum sp, Cassiope sp, Loiseleuria sp, Vaccinium vitis-idaea, Rhododendron sp, Phyllodoce caerulea, Dryas octopetala	46.7
	Terrestrial	sedge	Absent, Present, Dominant	Are sedges absent, present, or dominant	45.5

	Terrestrial	non_sedge_herbaceous	Absent, Present, Dominant	Are grasses, rushes and forbs absent, present, or dominant	43.2
	Terrestrial	ev_needle_tree	Absent, Present, Dominant	Are evergreen needleleaf trees absent, present, or dominant	51.1
	Terrestrial	dec_needle_tree	Absent, Present, Dominant	Are deciduous needleleaf trees absent, present, or dominant	37.5
	Terrestrial	dec_broad_tree	Absent, Present, Dominant	Are deciduous broadleaf trees absent, present, or dominant	40.4
	Terrestrial	sphagnum_cover	Absent, Present, Dominant	Are Sphagnum mosses absent, present, or dominant	43.6
	Terrestrial	other_moss_cover	Absent, Present, Dominant	Are other mosses (non-Sphagnum) mosses absent, present, or dominant	37.2
	Terrestrial	canopy_height	m	Height of the vegetation canopy	24.0
	Terrestrial	soil_moisture_class	Wet = At least sometimes inundated or water table close to surface, Dry = well-drained, Moist = in between wet and dry	General descriptor of site moisture	53.9
	Terrestrial-forest	forest_age	YYYY	Forest age since last disturbance (anthropogenic/natural)	11.3
	Terrestrial	soil_depth	cm	Soil organic layer depth	21.1
	Terrestrial	soil_ph		Surface soil pH in ca. 0-10 cm depth	11.3
	Terrestrial	soil_perc_c	%	Surface soil C % in ca. 0-10 cm depth	13.9
	Terrestrial	soil_perc_n	%	Surface soil N % in ca. 0-10 cm depth	7.6
	Terrestrial	c_stock	kg C m-2	Soil organic carbon stock, ideally for the entire soil profile.	22.1
	Terrestrial	stock_depth	cm	Soil depth used in the stock calculation.	21.3

	Terrestrial	soil_type_detail		Soil type description	22.9
	Terrestrial	lai	m-2 m-2	Leaf area index	14.5
	Terrestrial	ndvi		Normalized difference vegetation index	4
Aquatic variables	Aquatic	waterbody_type	Lentic, Lotic	Type of waterbody: lentic (standing water) or lotic (flowing water)	99.6
	Aquatic	aquatic_site_sampling_location	Edge, Center, Both	The locations of the measurements within the waterbody	34.8
	Aquatic	water_body_trophic_status	Oligotrophic, Mesotrophic, Eutrophic	Trophic state classification describing the productivity of a lentic waterbody	38.6
	Aquatic	water_area	m2	Area of lentic waterbody	31.9
	Aquatic	water_depth	m	Mean, maximum or point-level depth of a lentic or lotic waterbody	34.7
	Aquatic	water_depth_location	Mean, Sampling location, Maximum	Description representing the location of the water depth measurement	5.7
	Aquatic	water_ph		Mean pH at water surface during the measurement interval	61.8
	Aquatic	water_n	mg l-1	Mean total nitrogen at water surface during the measurement interval	41.9
	Aquatic	water_p	mg l-1	Mean total phosphorus at water surface during the measurement interval	35.6
	Aquatic	water_doc	mg l-1	Mean dissolved organic carbon content at water surface during the measurement interval	62.2
	Aquatic	water_iceon	MM/DD/YYYY	Ice-on date	10.4
	Aquatic	water_iceoff	MM/DD/YYYY	Ice-off date	10.9
	Aquatic	water_temperature	°C	Mean surface water temperature during the measurement interval	76.9
	Aquatic	benthic_veg	Yes, No	Is benthic vegetation occurring at the site?	7.2
	Aquatic	emergent_veg		Details related to the emergent vegetation	5.3

	Aquatic	sediment	Minerogenic, Organic, Peat, Yedoma, Unspecified	Sediment type	23.8
	Aquatic	water_chlorophyll	mg l ⁻¹	Mean concentration of chlorophyll at water surface during the measurement interval	2.5
	Aquatic	water_do	mg l ⁻¹	Mean dissolved oxygen at water surface during the measurement interval	29.5
	Aquatic	water_co2	μmol l ⁻¹	Mean dissolved CO2 concentration at water surface during the measurement interval	77.0
	Aquatic	water_ch4	μmol l ⁻¹	Mean dissolved CH4 concentration at water surface during the measurement interval	75.3
	Aquatic	water_turbidity	FNU (Formazin Nephelometric Unit)	Mean turbidity of the water during the measurement interval	9.3
	Aquatic	water_conductivity	μS/cm	Mean electrical conductivity at water surface during the measurement interval	45.3
	Aquatic	k600	cm h ⁻¹	Gas transfer velocity normalized to a Schmidt number of 600	16.0
	Aquatic	k600_equation		Equation used to determine gas transfer velocity (k600)	17.4
	Aquatic	k600_method		Method used to determine gas transfer velocity (k600)	25.9
	Aquatic	stream_discharge	m ³ s ⁻¹	Stream discharge of lotic waterbody	18.0
	Aquatic	stream_velocity	m s ⁻¹	Stream velocity of lotic waterbody	2.1
	Aquatic	water_mixing_regime	E.g. Monomictic, Dimictic, Polymictic	Mixing regime of lentic waterbody to indicate the frequency of mixing throughout the year (once = monomictic, twice = dimictic, multiple times = polymictic)	8.3
	Aquatic	strahler_order		Strahler order of lotic waterbody to define stream size	26.1
	Aquatic EC	fetch_screening	Yes, No, Unknown	Indicator whether fetch screening was applied	99.2

	Aquatic EC	fetch_detail		Details about fetch screening	92.1
	Aquatic	air_co2	ppmv	Mole fraction of CO2 in the air above water surface during the measurement interval	18.4
	Aquatic	air_ch4	ppbv	Mole fraction of CH4 in the air above water surface during the measurement interval	13.8
	Aquatic Non-EC	water_d13ch4	permil	Ratio of stable carbon isotopes of CH4 at water surface during the measurement interval	5.8
	Aquatic Non-EC	water_d13co2	permil	Ratio of stable carbon isotopes of CO2 at water surface during the measurement interval	2.9
	Aquatic	isotopic_analysis_detail		Details regarding the isotopic analysis, e.g. determined from dissolved gas or ebullition	1.3
Policies and notes		data_usage	Tier1 = data are open and free for scientific and educational purposes, Tier2 = data producers must have opportunities to collaborate and consult with data users, Other (please specify)	Instruction of data usage	98.4
		data_version		Version number for data extracted from repositories or version number given by the data contributor based on their version tracking.	24.4
		notes		Additional relevant information	38.3
		id		Unique identifier given to each individual monthly entry at each site	100

3.2 Definition of site

To differentiate between measurement locations, we used two attributes: `site_name` and `site_reference`. The column “`site_name`” (e.g., Stordalen Mire) is considered a more general description of a site whereas “`site_reference`” (e.g., Stordalen Mire_Palsa Site_Chamber) is a more specific description of a plot/sub-site within a broader site and indicates the method of measurement. The distinction between `site_name` and `site_reference` is most evident among chamber studies where measurements may have been made across different types of vegetation or landscape characteristics within a single site (i.e., several `site_references` corresponding to a single “`site_name`”). We assigned a unique `site_reference` to a site, as long as it had a distinct land cover class, coordinates, or unique related environmental data. For eddy covariance tower measurements, the distinction between “`site_name`” and “`site_reference`” is less significant. Eddy covariance towers from the major flux repositories (e.g., FLUXNET, Ameriflux, ICOS) often have a FluxID assigned which was reflected in the `site_reference` of the data (e.g., Stordalen_SE-St1_tower). For user-contributed tower data that did not have a FluxID, `site_reference` is often the `site_name` along with a name specified by the data contributor with the addition of “_tower”. In instances where footprint analysis was applied to split tower data in addition to the ecosystem-level fluxes (sites Ranskalankorpi, Iskoras, Stordalen), the “`site_reference`” column specifies which ecosystem the flux comes from (e.g., Iskoras_NO-Isk_palsa_tower and Iskoras_NO-Isk_pond_tower). ABCFlux v2 comprises 1,024 individual site names and 5,121 individual site references. In order to not exaggerate the number of sites in this synthesis, we refer to the number of unique “`site_name`” unless otherwise specified.

Flux data for lentic waterbodies were aggregated to the waterbody level by taking the unweighted average of all observational data (e.g., “Edge” and “Center”) for sites where multiple measurements were conducted. Therefore, spatial within-lake differences were not individually accounted for. However, the column “`aquatic_site_sampling_location`” provides information about the location within the waterbody where measurements were conducted, and differentiates between the edge and the center of the waterbody, or both if measurements were conducted across the waterbody which can have implications for total flux calculations. (Ray et al., 2023). It should be noted that the sampling location within the waterbody remained unknown (no location information was available) for 66% of the aquatic flux measurements, 22% were sampled from the center of the waterbody, 7% from both (center and edge), and 5% from the edge of the waterbody. If measurements were taken on different days, the number of days is specified in the various “`nr_measurement_days`” columns. Where measurements were taken within a single campaign along a large lotic waterbody over several kilometers, the river was divided into sections and flux data were aggregated for each section separately. This spatial aggregation was handled on a case-by-case basis and in close collaboration with data contributors. In instances where this spatial aggregation was applied, the “`site_name`” represents the river name (e.g., Teno) and “`site_reference`” reflects the river section (e.g., Teno_Karigasniemi_chamber, where Teno river is the name of the river, and Karigasniemi the measurement location).

4. Data quality and screening

700 We screened and cleaned data in ABCFlux v2 using expert judgement, informed by the gap-filled data percentage, quality flags and number of measurements days, if available. Our primary approach was to visually assess the time series of meteorological variables and fluxes for each site as well as the overall magnitudes in flux and supporting environmental data. In general, the quality control of all data was carried out in close cooperation with data providers.

705 4.1 Terrestrial fluxes

For repository data, we encountered 49 occurrences in eddy covariance site-level time series of CO₂ and CH₄ fluxes that had “flat lines” over several months of data (see Supplementary Fig. 1 panel A for an example), i.e., relatively constant flux values that did not vary by more than 3 g C m⁻² month⁻¹ over consecutive months. These “flat lines” often occurred at the beginning and end of time series and usually had a very high gap-fill percentage (a mean gap-fill percentage of 97%), indicating the value was based on very little measured data and was most often entirely gap-filled. We excluded flux data with “flat lines” if they were made up of three or more consecutive months with high gap-fill percentages (>75%).

715 Another issue that was identified during the quality check of the eddy covariance data submitted by data contributors, or extracted from data repositories, pertained to the winter months (Dec-Feb). During this period, NEE was occasionally found to be exactly zero with a gap-fill percentage of 100%. We excluded these zeroes from our dataset, as there should always be some variability due to measurement and data processing uncertainties, even during low-flux conditions. Moreover, previous studies (Kittler et al., 2017b; Natali et al., 2019; Watts et al., 2021) have shown that Arctic-boreal ecosystems can exhibit winter-season fluxes of significance, making the assumption of an exactly zero winter flux unlikely.

725 For 26% of CO₂ and 70% of CH₄ eddy covariance site-months (including data provided by data contributors and from flux repositories), there was no information about data quality and/or gap-filled data percentage. Consequently, we were not able to solely and systematically rely on these metadata for quality screening. Thus, we calculated the 1st and 99th percentiles for each combination of month, biome, and flux measurement method (EC and non-EC) and used them, together with the visual assessment of time series, to identify data that either strongly deviated from expected seasonal patterns or fell outside these percentile thresholds (Supplementary Table 2). We removed data if it was both outside of these percentiles and stood out visually (see Supplementary Fig. 1, panel C and D for examples). Primarily, we excluded months that were entirely gap-filled during winter and showed net CO₂ uptake beyond the 99th percentile. The visual inspection of time series also led to the removal of flux data from sites where one year in the dataset showed unrealistic patterns, such as potentially reversed signs (e.g., winter uptake and summer sources). These datasets were often downloaded from general data repositories like the Arctic Data Center and attempts to resolve the issues by contacting data contributors or

reviewing relevant publications from the site were unsuccessful. At some sites, flux time series followed a realistic seasonal pattern within the percentile thresholds (see Supplementary Table 2) despite some of the monthly fluxes being entirely gap-filled (often during a few months of the winter season, or due to, e.g., one year of missing data in a longer time series). Due to the limited amount of data in the Arctic-boreal region, these fluxes were kept in the dataset, and this is noted in the gap-filled data percentage column. In total, we kept 27 sites that included 90-100% gap-filled flux data during the peak winter months (Dec-Feb) across all the measured years because the seasonal dynamics and magnitudes matched those from other years of data from the same site or similar ecosystem types (see e.g. Supplementary Fig. 1, panel B). Though we did remove months where the data repository had winter months that were 100% gap-filled and data contributors provided only growing season data and advised against including repository data.

We examined terrestrial non eddy covariance (i.e. chamber and snow pack diffusion) using the same approach as eddy covariance, accessing the 1st and 99th percentiles along with the visual inspection of each site time series. These fluxes were not removed, as the limited temporal coverage of these measurement methods made it difficult to interpret seasonal patterns from time series graphs. Additionally, converting these often temporally limited observations into monthly cumulative fluxes can yield values with considerable uncertainty; however, we retained these data given the overall scarcity of measurements in this region and instead rely on the quality flags described in Section 5.3 to guide data users.

For the supporting environmental data, we removed data with unchanging values across three or more months as they likely represented a seasonal average and not monthly data. We also removed soil temperature observations that were above 40°C and below -40°C as these were the approximate temperature ranges seen in ABCFlux v1 and BAWLD-CH4 and values outside of this range were assumed to be errors. The largest amount of cleaning was done for the water table depth to ensure that the sign of the data was aligned with the variable description for ABCFlux v2 (i.e., positive is below the soil surface, negative is above).

4.2 Freshwater fluxes

The footprints of eddy covariance towers over waterbodies often include surrounding non-aquatic ecosystems. Therefore, a fetch screening is commonly applied by data contributors of sites with mixed footprints (e.g. Lake Villasjön in Sweden; (Jammet et al., 2017) to separate flux contributions from aquatic and adjacent terrestrial ecosystems. The simplest approach to remove non-aquatic flux contributions is to apply a wind-directional fetch screening that excludes half-hourly fluxes from wind directions associated with land surfaces during data processing (also done in Golub et al., 2023). Apart from the wind-directional screening, more sophisticated approaches have been used (e.g. Bayesian modeling (Pirk et al., 2024)). In ABCFlux v2, we indicated whether fetch screening was applied and for which wind directions. Regardless of the type of approach used to filter out non-aquatic carbon fluxes for eddy covariance data, the number of data gaps tends to be larger than for terrestrial towers. Because of this, and the fact that most gap-filling approaches are tailored to terrestrial ecosystems, gap-

780 filling for aquatic towers remains challenging. We included both gap-filled and non-gap-filled aquatic tower data and derived monthly cumulative fluxes based on the available data.

Within the process of quality screening, we also unified some variables such as the gas transfer velocity, which can be used to estimate diffusive fluxes based on dissolved gas concentrations and hydraulic properties. Various methods to derive gas transfer velocities can be used, which
785 differ slightly for lentic and lotic ecosystems (Hall and Ulseth, 2020; Klaus and Vachon, 2020; Raymond et al., 2012). The gas transfer velocity can be expressed as a magnitude independent of gas and temperature when normalized to a Schmidt number (Sc) of 600 (k_{600} , i.e. normalized gas transfer velocity) for freshwater at 20 °C. The Schmidt number is defined as the ratio between kinematic viscosity and mass diffusivity but is often empirically determined, and
790 quantifies the temperature-dependent molecular transport properties of each gas (Jähne et al., 1987). Where gas transfer velocities for a specific gas species (k_{gas}) were not normalized to a Schmidt number of 600, we converted them accordingly ($k_{600} = (600/Sc)^n * k_{gas}$; (Cole and Caraco, 1998)), where n is determined by windspeed (Guérin et al., 2007). Furthermore, dissolved CO_2 and CH_4 concentrations (or partial pressures) were converted from a range of
795 given units (ppm, ppb, μatm , mol/L, mmol/L, nmol/L, mmol/m³, mg/L) to $\mu mol/L$ following previous procedures from GRiMeDB (github code: <https://github.com/lukeloken/GlobalRiverMethane>).

5. Data usage notes

800 Despite extensive efforts in dataset cleaning, users of ABCFlux v2 should remain aware of certain considerations to avoid potential misinterpretation of the data.

5.1 Uncertainties related to gap-filling and flux partitioning

The approach used to gap-fill and estimate monthly cumulative fluxes varied within and across measurement methods. For eddy covariance data, the most common gap-filling technique was
805 Marginal Distribution Sampling (MDS), used in 57% of site-months, following the ONEFlux pipeline processing approach (Pastorello et al., 2020). However, we also incorporated fluxes that were gap-filled using other methods including neural networks and non-linear regression. For terrestrial and aquatic eddy covariance data, the cumulative monthly flux was most commonly obtained by multiplying the gap-filled monthly mean flux rate given as per day ($g\ C\ m^{-2}\ day^{-1}$) by the number of days in the month. In instances where gap-filled data at terrestrial sites
810 were not available, we multiplied the daily mean of the respective month by the days in that particular month, and indicated that no gap-filling was applied in the “gap_fill” column. With these non-gap-filled data, gaps covered 53% for NEE and 69% for CH_4 flux per month on average. For aquatic eddy covariance sites, half-hourly flux data adopted from Golub et al.
815 (2023) had been gap-filled following (Pastorello et al., 2020) and were aggregated to monthly cumulative fluxes for ABCFlux v2, but data contributors also shared non-gap-filled data which we then aggregated to monthly cumulative fluxes and indicated that in the “gap_fill” column. As mentioned above, additional gaps in flux data occur where fetch screening was applied to

heterogeneous tower footprints. For aquatic sites, the gap-fill percentage often reflects both general data gaps, such as those caused by sensor issues, maintenance, power outages, or poor turbulence conditions, and the additional exclusion of data due to fetch screening. Consequently, gap-fill percentages between terrestrial and aquatic sites in ABCFlux v2 cannot be directly compared. Gap-fill percentages for aquatic data averaged 68% for CO₂ and 65% for CH₄.

The methods used to partition CO₂ fluxes into GPP and Reco at terrestrial eddy covariance sites varied, with the most common approach (66% of site-months) being that of (Lasslop et al., 2010; Reichstein et al., 2005)), which has been widely applied in global synthesis and upscaling studies (e.g., (Nelson et al., 2024)). In addition to the Reichstein et. al (2005) method, this dataset also includes fluxes based on other partitioning methods such as (Lasslop et al., 2010; Reichstein et al., 2005)) and (Runkle et al., 2013). Potential limitations and differences among partitioning methods in the Arctic-boreal context have been extensively discussed in Virkkala et al. (2022) (under “Uncertainties in eddy covariance flux partitioning”). In particular, nighttime partitioning (Reichstein et al., 2005) can introduce uncertainty in high-latitude regions where low-light nighttime conditions are limited during summer. However, when comparing multiple gap-filling and partitioning methods across sites, we found that the variability in annual GPP and Reco estimates was small (Desai et al., 2008; Keenan et al., 2019), lending confidence to the partitioned GPP and Reco estimates derived from the diverse methods used in this dataset.

Chamber and other non-eddy covariance flux measurements, although generally more temporally sporadic than eddy covariance, were often converted to monthly cumulative fluxes using a similar method as eddy covariance (i.e., measurements averaged and multiplied by days) or, in some cases, gap-filled with light- and temperature-response models; details related to the approach can be found in the “gap_fill” column. Similarly, meteorological data were often collected only during these sporadic measurements, and are thus not based on continuous meteorological measurements within a month. 20% and 33% of terrestrial non-eddy covariance NEE and CH₄ flux measurements, respectively, 67% and 50% of lentic chamber CO₂ and CH₄ flux measurements, and 14% and 16% of lotic chamber CO₂ and CH₄ flux measurements were derived from one single measurement day (however note that the majority of lotic data do not have this information, 80% of CO₂ and 76% of CH₄ data). Furthermore, these measurements were often conducted during daytime only (69% of terrestrial chamber measurements, 55% of lentic, and 90% of lotic flux measurements). The sporadic nature and lower data coverage of the non-eddy covariance data leads to uncertainty in monthly flux and meteorological data. The bias toward daytime measurements may lead to an overestimation of net CO₂ sinks in vegetated ecosystems due to less photosynthesis at night (Lai et al., 2012; Järveoja et al., 2020) and an overestimation of CH₄ emissions in lentic ecosystems due to calmer winds and cooler temperatures at night (López-Blanco et al., 2017; Sieczko et al., 2020; Voigt et al., 2023). Similarly, in lotic systems, measurements were typically conducted during the day, even though nighttime emissions may exceed daytime release, potentially leading to an underestimation of monthly CO₂ emissions (Attermeyer et al., 2021; Gómez-Gener et al., 2021). However, such biases were not clearly evident in the data, even when comparing monthly fluxes derived using simple averages versus light- and temperature-response models (Supplementary Fig. 2 and 3).

Site-months based on daytime-only measurements, however, tended to show greater variability, with lower minimum and higher maximum fluxes compared to those including both day- and night-time data. The columns “diurnal_coverage”, “chamber_nr_measurement_days_co2”, “chamber_nr_measurement_days_ch4” and “ebullition_nr_measurement_days” help the data users understand the temporal representativeness of the data.

5.2 Fluxes and periods captured by the dataset

Gas flux and dissolved gas concentration measurements were more abundant during the growing season (May-August) compared to the non-growing season (September-April). The average total number of observations per month was 4,058 during growing season months vs 1,479 during non-growing season months, and 50% of terrestrial data and 28% of aquatic data were collected in the non-growing season despite two thirds of the year being non-growing season. Non-growing season monthly fluxes were often more heavily gap-filled than those during the growing season (54% in the growing season vs. 68% in the non-growing season for CO₂, 61% vs. 63% for CH₄), which further contributes to higher uncertainties. This is problematic as the non-growing season, and in particular spring and autumn season CO₂ and CH₄ emissions in both terrestrial and freshwater ecosystems, are important for the annual carbon balance (Arndt et al., 2023; Lyu et al., 2024). In freshwater ecosystems, the CO₂ and CH₄ emissions during the spring ice-out period (“storage flux”) are known to contribute significantly to annual fluxes (e.g., 11 to 59% in subarctic lakes (Jammet et al., 2015; Juutinen et al., 2009a; Karlsson et al., 2013, 2024; Prėskienis et al., 2021)), but measurements during the spring period are limited. Assuming a spring ice-out period in May-June for lentic ecosystems and March-April for lotic ecosystems (following Song et al., 2024), 28% of the lentic site-months and only 6% of the lotic site-months captured this period. Furthermore, information on the timing of the ice-on and ice-off was rarely given for non-growing season measurements at aquatic sites. Therefore, annual CO₂ and CH₄ fluxes from aquatic ecosystems may be underestimated due to the lack of data during these influential seasonal periods. However, the extent of this underestimate is uncertain given the unknown role of CH₄ oxidation and diel variation in CO₂ consumption in the water prior to emissions (Pajala et al., 2023; Rudberg et al., 2021). At the same time, simple averaging of the dominating daytime summer fluxes to spring and autumn seasons may instead generate substantial overestimates for CH₄, while simultaneously underestimating CO₂ emissions in productive lakes where emissions can occur outside the growing season (Natchimuthu et al., 2016; Rudberg et al., 2021).

A substantial part (23%) of freshwater carbon fluxes synthesized in ABCFlux v2 were gained from samples taken exclusively in the center of waterbodies, excluding edges. This may be problematic as the spatial variability of carbon fluxes across depth zones in lentic ecosystems can be large and plays a significant role when estimating ecosystem-level emissions, particularly for CH₄ fluxes (Kuhn et al., 2023). Studies have shown that CO₂ emissions might be overestimated when only considering measurements from the center of lakes (Loken et al., 2019), whereas CH₄ emissions might be underestimated in some lakes (Juutinen et al., 2003), but not others (Schmiedeskamp et al., 2021). It is also important to note that the location of eddy covariance towers in freshwater ecosystems can contribute to under- or over-estimating fluxes depending on the location of localized emission hotspots (e.g. thermokarst features).

Therefore, both under- and overestimations in aquatic cumulative ecosystem-level fluxes are possible, depending on the gas, sampling location, sampling frequency, and seasonal dynamics involved (Ray et al., 2023).

5.3 Quality flags based on expert assessment

To help data users assess data quality in the flux records, we included additional columns “expert_flag_co2”, “expert_flag_gpp”, “expert_flag_reco” and “expert_flag_ch4” (Table 4, Supplementary Fig. 4). These columns may aid in filtering out sites and/or observations that are atypical or highly uncertain and should be used with particular caution when scaling monthly fluxes to estimate budgets across larger domains. In other use cases, these observations may still be useful.

As a broad overview, we flagged CO₂ fluxes outside of the 1st and/or 99th percentiles (flag 1), uncertain fluxes due to sporadic non-eddy covariance measurements (flag 2), long periods of eddy covariance data that have been entirely gap-filled (flag 3), and sites that do not represent typical conditions across the Arctic-boreal region due to human induced changes (flag 4, Table 4). All other CO₂ fluxes were marked with a zero (i.e. representative, good-quality data).

In more detail, we used 1st and 99th percentiles for terrestrial CO₂ fluxes calculated separately for each combination of month, biome (tundra, boreal, temperate), and measurement method (EC, chamber, snow diffusion) to flag extremely high or low fluxes with 1. This procedure was applied independently to NEE (expert_flag_co2), GPP (expert_flag_gpp), and Reco (expert_flag_reco). This flag considered, for example, unusually low NEE values during peak winter months (i.e. fluxes from less than -25 g C m⁻² month⁻¹ between December and February), which represent unrealistically high winter net uptake values likely due to issues in eddy covariance data collection and gap-filling (Jentsch et al., 2021; Kittler et al., 2017). During the summer months (June-August), flag 1 captured eddy covariance NEE data below -140 g C m⁻² month⁻¹ and chamber NEE data below -425 g C m⁻² month⁻¹. For chamber data, these large negative NEE values were often based on single daytime measurements alone. Flag 2 marks terrestrial chamber CO₂ and CH₄ flux growing season (May-Aug) measurements with fewer than three observation days in a month, where monthly values were calculated using simple averages rather than models incorporating light and temperature. Such limited sampling and simplistic averaging can introduce bias by failing to capture environmental variability, especially during the growing season. The three-day threshold is based on the assumption that approximately one measurement per week is necessary to produce a more reliable cumulative estimate, as suggested by Virkkala et al. (2022). To account for long periods of missing and entirely gap-filled data, we introduced flag 3 which marks eddy covariance data with 3 or more consecutive months of 100% gap-filling. For flag 4, we flagged eddy covariance sites where there have been significant atypical changes to the landscape from humans such as sites where the organic soils were removed with a bulldozer (Euskirchen et al., 2017; Walter Anthony et al., 2024). While flag 1 was applied separately to NEE, GPP, and Reco, flags 2–4 were applied only to expert_flag_co2, as they pertain to measurement methods and site conditions rather than to the flux variables themselves.

For aquatic CO₂, fluxes from lentic ecosystems that exceeded the monthly 99th percentile (between 26.4 g C m⁻² month⁻¹ in January to 320.9 g C m⁻² month⁻¹ in June) were marked with flag 1. We refrained from flagging fluxes from lotic ecosystems since these may be affected, for example, by higher turbulence and larger resulting gas exchange compared to stiller conditions in lentic ecosystems.

For terrestrial CH₄ fluxes, we applied flag 1 to fluxes higher than 30 g C m⁻² month⁻¹ as such high values are well beyond the range of previously published estimates (Kuhn et al., 2021). Flags 2–4 were applied to terrestrial CH₄ fluxes following the same criteria used for CO₂ fluxes.

Similarly to terrestrial data, we flagged diffusive CH₄ fluxes for lentic ecosystems that were higher than 30 g C m⁻² month⁻¹ and were excluded from analyses in previous studies (such as BAWLD-CH₄) due to exceptionally high gas transfer velocities with flag 1. This flag was not applicable to aquatic eddy covariance sites since total CH₄ fluxes did not exceed the chosen threshold. Furthermore, we refrained from flagging high diffusive CH₄ fluxes for lotic ecosystems for the same reasons as mentioned above.

Flag 2 was not used for aquatic fluxes since 90% of the aquatic monthly cumulative CO₂ fluxes and 86% of CH₄ diffusion fluxes from non-eddy covariance methods were derived from 3 or less measurements per month. Furthermore, none of the aquatic eddy covariance sites showed 3 or more consecutive months of 100% gap-filled data (flag 3), and none of the aquatic sites experienced significant land use change which could be linked with extremely high fluxes (flag 4).

Table 4. Quality flags assigned to the terrestrial and aquatic CO₂ and CH₄ fluxes based on expert assessment (columns “expert_flag_co2”, “expert_flag_gpp”, “expert_flag_reco” and “expert_flag_ch4”). Note that the data that the respective flags apply to may change between terrestrial and aquatic, as well as lentic and lotic fluxes, and between flux methods. Further detail is given in Sect. 5.3.

CO ₂ Flag	Requirements	Applicable data
0	Terrestrial: NEE, GPP, Reco within the 1st and 99th percentiles Aquatic-lentic: CO ₂ fluxes within 99th percentile Aquatic-lotic: all CO ₂ fluxes	All terrestrial and aquatic
1	Terrestrial: NEE, GPP, Reco outside of 1st and 99th percentiles Aquatic-lentic: CO ₂ fluxes greater than 99th percentile	All terrestrial and aquatic (lentic)
2	Growing season NEE fluxes with 3 or less measurement days in a month AND no modeling used in gap-filling	May-August terrestrial non-eddy covariance

3	NEE fluxes with 3 or more consecutive months of 100% gap-filling	Terrestrial eddy covariance
4	Sites with significant land use changes from humans	Selected terrestrial sites
CH₄ Flag		
0	Less than 30 g C m ⁻² month ⁻¹	All terrestrial, aquatic: only lentic non-eddy covariance diffusion and eddy covariance total flux
1	Greater than 30 g C m ⁻² month ⁻¹	All terrestrial, Aquatic: only lentic non-eddy covariance diffusion
2	Growing season CH ₄ fluxes with 3 or less measurement days in a month AND no modeling used in gap-filling	May-August terrestrial non-eddy covariance
3	CH ₄ fluxes with 3 or more consecutive months of 100% gap-filling	Terrestrial eddy covariance
4	Sites with significant land use changes from humans	Selected terrestrial sites

6. Spatial and temporal distribution of the dataset

980 Throughout the following sections, we describe the spatial and temporal distribution of the dataset. We use the column “site_name” to identify unique sites, and group the flux measurement techniques following the three measurement categories presented in Table 1. The dissolved gas-category includes concentration data alone and is included in the site-month numbers presented Figures 3-6. The term “site-month” refers to monthly data (fluxes and concentrations) and excludes seasonal fluxes (199 observations in the overall dataset). The key ecosystem categories used in visualizations are described in Supplementary Text 1; the terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

6.1. Number of site-months, unique flux values and sites

990 The ABCFlux v2 dataset comprises 29,121 site-months, where each row represents a month with one to several unique gas fluxes (e.g. one site may include both a CH₄ and GPP flux for a given month) for a given site. In total, the dataset includes 56,139 unique flux values spanning CO₂ and CH₄ fluxes and their surface-atmosphere transport pathways. This “unique flux” value reflects the sum of all non-NA flux entries in the “nee”, “gpp”, “reco”, “co2_flux”, “diffusion”, “ebullition”, “total_ch4”, “storage”, and “seasonal” columns (see Supplementary Table 1). For 995 aquatic concentrations, there are 8,801 site-months and 15,668 unique aquatic concentration

measurements. Throughout the following sections, we focus on site-months rather than unique flux values. The number of site-months per site varied from 1 month to 330 site-months (1 to 330 site-months at terrestrial sites, 1 to 149 site-months at aquatic sites), with an average of 15 site-months (38 for terrestrial sites, 4 for aquatic sites); note that some sites had some sporadic months that were completely gap-filled in these estimates. We identified six large site clusters in the data (> 800 monthly terrestrial and aquatic observations within a 30 km² radius): Toolik Lake (USA), Abisko-Stordalen (Sweden), Hyytiälä-Siikanen (Finland), Degerö-Flakaliden (Sweden), Fairbanks (USA), and Utqiagvik (formerly Barrow; USA), see Supplementary Figure 11.

The dataset includes a total of 1,024 sites, comprising 337 terrestrial and 712 aquatic sites (Table 5). Most sites collected data primarily during the growing season (May-August), while 119 sites operate year-round, the majority of which (108) are eddy covariance sites in terrestrial ecosystems. ABCFlux v2 includes 16 terrestrial year-round CO₂ flux sites with >10 years of data, mainly located in the boreal biome (14 sites). The longest time series of CO₂ fluxes in ABCFlux v2 in the boreal biome were from Hyytiälä (27 years), Degerö (23 years), Fyodorovskoye (22 years), University of Alaska Fairbanks (21 years), and in the tundra biome sites Eight Mile Lake (14 years), Imnavait Creek Watershed Heath Tundra (14 years), and Imnavait Creek Watershed Tussock Tundra (14 years). There were 33 terrestrial sites with both year-round CO₂ and CH₄ flux data (at least for some of the years; see Supplementary Table 3). For CH₄, there were 34 terrestrial sites with year-round data with the longest time series at Trail Valley Creek (9 years), Cherskii reference (8 years), Cherskii (8 years), and Scotty Creek Landscape (8 years). For the aquatic dataset, CO₂ fluxes were measured year-round at 7 eddy covariance and non-eddy covariance sites (Bernard Lake, Eastmain Reservoir, Iskoras, Kuivajärvi, Romaine-2 reservoir, Vanajavesi, Simpevarp) and at three of them CH₄ fluxes were measured year-round (Bernard Lake, Iskoras, Romaine-2 reservoir). ABCFlux v2 also includes automatic chamber measurements from 23 sites. However, these sites have shorter time series and contribute 15% of monthly CO₂ chamber measurements and 6% of monthly CH₄ chamber measurements.

Table 5. Number of sites and site-months for the whole dataset separated by terrestrial (CO₂ and CH₄), aquatic (CO₂ and CH₄), CO₂ only and CH₄ for growing season (May-August) and non-growing season (September- April) months. The table also lists the number of year-round sites and year-round sites with 5+ years of data. Note that some sites are both terrestrial and aquatic. A year-round site was defined as having at least some data for all months (no 100% gap-filled months) for at least one full year in the time series; if no data on gap-filling percentage was provided then we assumed there was some data for all months. The sum of terrestrial and aquatic site counts exceeds the total number of sites because some sites have both terrestrial and aquatic measurements.

	Gas species	All		Growing season			Non-growing season		Year-round	
		Sites	Site-months	Sites	Site-months	Sites with 5+ years	Sites	Site-months	Sites	Sites with 5+ years
ABCFlux v2		1,024	29,062	947	16,409	122	564	12,610	119	52
Terrestrial		337	18,952	304	9,075	108	265	9,046	112	50
	CO ₂	256	16,257	239	7,367	102	221	9,007	108	50
	CH ₄	173	7,433	156	4,456	31	121	3,066	34	9
Aquatic		712	10,110	665	7,160	15	312	2,792	7	2
	CO ₂	628	8,791	606	6,282	11	292	2,522	7	2
	CH ₄	464	8,855	455	6,414	12	279	2,413	3	1
All CO ₂		875	25,101	829	13,649	107	502	11,529	115	52
All CH ₄		651	16,415	596	10,870	34	392	5,479	37	10

6.2 Regional coverage

Terrestrial and aquatic sites are widely distributed across the Arctic-boreal domain (Fig. 1). However, the distribution of sites in ABCFlux v2 varies significantly across regions and key ecosystems in terms of the number of sites, site-months, and the gas species measured (Table 5, Supplementary Fig. 6). In terms of site-months in the full dataset, Alaska showed the highest coverage (24.3% of the dataset) with Sweden (20.9%), Canada (19.2%), and Finland (17.4%) following closely behind, and finally Russia (11.2%), Norway (3%), and Greenland (2.2%) (Supplementary Table 4). Among just the terrestrial data, Alaska, Finland, and Canada account for a majority of the site-months (23.4%, 20.5%, and 20.3% respectively), with Sweden only making up 14.7% of terrestrial observations. In contrast, measurements from Sweden made up

the highest number (32.5%) of the aquatic site-months followed by Alaska (25.9%), Canada (17.6%), and Finland (11.5%). The majority of aquatic sites came from Finland (31.6%), and this dominance can be attributed to a few key aquatic studies that were incorporated in our dataset (Juutinen et al., 2009b; Kortelainen et al., 2006). The density of sites in these countries is also quite variable in terms of terrestrial and aquatic systems as well as the type of flux (Supplementary Fig. 5).

6.3 Temporal and seasonal coverage

Data in ABCFlux v2 span the years 1984 to 2024, with the majority of monthly observations occurring after 1999 (92.5% in total, 95.0% terrestrial, 87.4% aquatic, Fig. 3). Years with the largest amount of data are 2014-2019. The distribution of sites with more recent data is less comprehensive (2022 as an example; Supplementary Fig. 9). This does not imply these sites discontinued data collection, but rather that the data were not made available at the time of this synthesis. Regarding seasonal coverage, eddy covariance data are more evenly distributed across the year compared to chamber and other non-eddy covariance measurement methods (Supplementary Fig. 6). Terrestrial ecosystems show the highest data availability in July, with 1.5 to 3 times more data than other months, depending on the ecosystem (Fig. 4). In contrast, data coverage in aquatic ecosystems peaks in August, although overall coverage from June to August remains similar. Aquatic data show a sharp decline during spring and autumn, whereas terrestrial ecosystems experience a more gradual decrease outside the summer months. Among terrestrial ecosystems, wetlands exhibit the strongest seasonal bias, with substantially more observations in summer compared to other seasons.




Fig. 3. The number of CO₂ and CH₄ flux site-months across years. Numbers of observations are shown as stacked bars for each year. If a site-month measured both CO₂ and CH₄, it is counted as separate site-months in the figure (1 CO₂ site-month and 1 CH₄ site-month). The distribution of sites with more recent data is less comprehensive (2022 as an example). This does not imply these sites discontinued data collection, but rather that the data were not made available at the time of this synthesis




Fig. 4 The number of CO₂ and CH₄ monthly observations across months and key ecosystem types (A = terrestrial, B = aquatic). Aquatic concentration site-months are included in the figure. The terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

6.4 Land cover type and disturbance coverage

Wetlands (including bogs, fens, permafrost bogs, tundra wetlands, marshes) had the highest number of flux observations (8,758 site-months), driven by the abundance of CO₂ flux measurements (Fig. 5). Boreal forests were the second most measured ecosystem type (6,981 site-months), while tundra systems were the least studied (3,308 site-months). CO₂ flux site-months dominated the dataset across most ecosystem types (25,222 site-months), except for lentic ecosystems, where CH₄ flux measurements (3,098 site-months) were more frequent than CO₂ flux measurements (2,915 site-months).





Fig. 5. The number of CO₂ and CH₄ flux and concentration site-months per key ecosystem types. Aquatic concentration site-months are included in the figure. The terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

To better understand data coverage and calculate aerial extent across more detailed land and waterbody types, we used a combination of BAWLD and ESA CCI land cover data for terrestrial ecosystems (see Supplementary Text 1 for details) and a simplified version of BAWLD for aquatic ecosystems, focused on waterbody sizes. Overall, flux site-months were somewhat unevenly distributed across key land cover types relative to their areal extent (Fig. 6). Some classes were measured more in comparison to the area they cover (e.g., bogs, fens, permafrost bogs, wet tundra), while others showed more balanced representation (e.g., shrublands, graminoid ecosystems, evergreen needleleaf forests, mixed forests). In contrast, some classes were sparsely measured relative to their large areal extent (e.g., deciduous needleleaf forests, sparse vegetation and barren). For aquatic ecosystems, the most pronounced coverage biases were observed in the large lentic class, which was underrepresented relative to its total surface area, while small lotic ecosystems were disproportionately sampled compared to their small surface area extent. However, since flux data in our dataset are collected at the lake level, each site-month observation is weighted equally, regardless of lake size. This may not accurately reflect larger lakes, where a single observation can represent a much greater area. Similarly, this approach does not account for flux magnitudes or variability, which are variable across ecosystems. For example, lotic and wetland ecosystems often exhibit substantial CH₄ flux variability, necessitating more frequent observations for accurate representation compared to, for instance, barren ecosystems.



1115 **Fig. 6.** The coverage of monthly flux observations and areas of key terrestrial (A) and aquatic (B) ecosystem types from the ESA land cover model. Aquatic concentration site-months are included in the figure. Some observations were left unclassified due to the lack of descriptive data and are not shown in the figure. There are 61 site-months of data from marshes, resulting in their proportion appearing nearly zero in the figure.

1120 There are 272 sites in ABCFlux v2 that reported a disturbance, constituting 11,558 site-months. These disturbances were broadly classified into 21 categories: altered hydrology, animal herbivory, artificial pond, beavers, drainage, drained lake, erosion, extreme weather, fire, forestry, human paths, insect herbivory, land use change, peat mining, reservoir, seismic lines, thaw, tidal effects, wastewater, and other. Among these categories, the most common
1125 disturbance noted was “Thaw” (131 sites, 6,091 site-months), which can be attributed to the thaw category including various types of permafrost thaw. We further divide the thaw category into 8 categories: [active layer thickening with subsidence](#), [active layer thickening without subsidence](#), [active layer thickening with talik formation](#), [ice-wedge degradation](#), [thaw lake/pond expansion](#), [wetland collapse](#), [active layer detachment sliding](#), and [retrogressive thaw slumping](#). Aside from thaw, fire and forestry were the most frequent disturbances in terms
1130 of terrestrial sites (35 and 26 sites respectively). For aquatic sites, thaw was still the most common disturbance (71 sites) followed by wastewater, though the number of sites with this disturbance was much less (9 sites). The years that disturbances occurred varied from hundreds of years ago to more recent (2023) and on-going disturbances.

1135 6.5 Flux site data coverage comparison to earlier synthesis datasets

ABCFlux v2 includes 59% more sites and 160% more site-months for terrestrial CO₂ flux data compared to ABCFlux v1. ABCFlux v2 has substantially more sites with CO₂ data from the non-growing season (September-April; 213 vs. 141 in ABCFlux v1). ABCFlux v2 also has more recent data, with 2020-2024 making up 16% of the data; though the most recent years 2023-
1140 2024 account for 2.1% of site-months likely due to a delay in publication or processing of data.

While ABCFlux v2 has substantial overlap with the major international eddy covariance data repositories (FLUXNET2015, FLUXNET-CH₄, Ameriflux, and ICOS), ABCFlux v2 also incorporates a large number of additional sites and site-months from community-contributed
1145 data (i.e., data provided directly by site PIs or researchers), enhancing both the spatial coverage and temporal range of the dataset. FLUXNET-2015 covers 25% of the terrestrial eddy covariance CO₂ site-months included in ABCFlux v2. However, many of these observations were also directly provided by data contributors or removed during data cleaning (see Sect. 3.1 for more details), reducing the net contribution of FLUXNET2015 to 18% of the terrestrial eddy
1150 covariance CO₂ site-months. While other repositories such as Ameriflux (17% of terrestrial CO₂ site-months) and ICOS (10% of site-months) remain important data sources in the Arctic-boreal region, ultimately community-contributed data represent a substantial portion of the dataset (31% of site-months). This trend is even more pronounced in the terrestrial CH₄ eddy covariance data, where FLUXNET-CH₄ covers 28% of site-months in ABCFlux v2 but
1155 contributes only 24% to the final dataset, while community-contributed data account for 60% of the observations.

Supporting data coverage remained relatively similar between the v1 and v2 datasets. For instance, soil organic carbon stock data were available for 16% of site-months in v1 compared
1160 to 22% in v2, and soil moisture data for 35% in v1 versus 45% in v2. However, the v2 database shows improved coverage of disturbance information, increasing from 30% in v1 to 44% in v2.

BAWLD-CH₄ terrestrial CH₄ flux data included 555 site-years (daily average over the growing season) of chamber data from 214 sites. Integrating and updating terrestrial CH₄ chamber data to monthly format in ABCFlux v2 resulted in 4520 site-months from 126 sites (using BAWLD-CH₄ “site” definition for comparison). BAWLD-CH₄ aquatic flux data included 396 site-years (daily average over the open-water season) of diffusive fluxes and 168 site-years of ebullitive fluxes from 391 and 151 sites, respectively. Integrating and updating aquatic flux data to monthly format in ABCFlux v2 resulted in 939-site-months from 286 sites for diffusion and 212 site-months from 27 sites for ebullition. The apparent loss of ebullition sites is because many of the original BAWLD-CH₄ sites include ebullitive flux data only at the seasonal timeframe and it was not possible to partition fluxes into monthly intervals. Seasonal ebullitive data were included in the “ch4_flux_ebullition_seasonal” column instead.

Of the lotic data, about 85% of site-months were transferred from the global river methane database (GRiMeDB), with the remaining 15% of data largely being submitted by data contributors or extracted from recent publications. Nine eddy covariance towers within the Arctic-boreal domain were included in the global analysis of lake and reservoir CO₂ fluxes (Golub et al., 2023). In comparison, ABCFlux v2 contains CO₂ flux data from 15 eddy covariance towers covering aquatic ecosystems, one of which was deployed on a Finnish river. Otherwise, new tower sites were added from lakes in Canada and Scandinavia.

We also compared the number of sites in ABCFlux v2 with the recent ARctic greenhouse Gas Observation metadata version 1 (ARGO) (Vogt et al., 2025), where metadata across observational platforms in the Arctic-boreal region were collected. Our comparison focused on the number of sites rather than site-months because ARGO does not contain the flux data itself. While we cross-checked ARGO when preparing v2 and it now includes all ABCFlux v2 sites, our dataset contains approximately 40 fewer eddy covariance sites with carbon flux data from terrestrial and aquatic ecosystems. This discrepancy is due to data for those sites not being publicly available. In particular, ABCFlux v2 excludes recently established sites, whose data are not yet fully processed or shared, and older, short-term towers from the early 2000s included in publications for which data could not be accessed. ABCFlux v2 includes recent data since 2022 for only around half of the existing tower sites that are active according to ARGO, in most cases likely due to the delay in making data available. For non-eddy covariance sites, the site counts vary more strongly between ABCFlux v2 and ARGO due to differences in defining a ‘site’. When comparing unique site names between both datasets, ABCFlux v2 contains roughly three quarters of the sites from the observational platforms Chamber and Dissolved in ARGO. In both datasets, the number of aquatic sites is larger than for terrestrial sites, whereas lentic and lotic sites are split evenly.

7. Flux synthesis

Here we show a summary of monthly and annual CO₂ and CH₄ flux variability to provide a synthesis of flux magnitudes showing the data spread as well as highlight some uncertainties that the data user needs to be aware of. We used columns “nee” for terrestrial data and

“co2_flux” for aquatic data for the CO₂ balance estimate, and the “ch4_flux_total” column for the CH₄ flux for both ecosystems without differentiation between diffusion and ebullition. Monthly fluxes were summarized using the full dataset (i.e., not using any quality flags) and are reported in the main text for the key ecosystem classes as well as in the Supplementary material for the more detailed ecosystem classes (Supplementary Tables 5-10). We additionally summarize monthly concentrations of CH₄ and CO₂ for lentic and lotic ecosystems.

Total annual estimates were calculated by finding the monthly median fluxes per overarching class type (wetland, tundra, boreal forest, lentic lotic) across the entire dataset, and these were summed to obtain annual fluxes per each class. In this approach, median monthly fluxes from the entire year from terrestrial ecosystems were considered. For freshwater ecosystems, we summed emissions from May-October (6 months), and we assumed ice-emissions accounted for 17% of the total annual emissions (following Liu et al., 2022; Ramage et al., 2024). We calculated annual emissions using a subset of the data where all data with flags 1–4 were removed based on “expert_co2_flag” and “expert_ch4_flag” columns. The flags considered unrealistically high or low fluxes, uncertain fluxes due to sporadic chamber measurements, eddy covariance data with long periods of entirely gap-filled data, and sites that do not represent typical conditions across the Arctic-boreal region.

The monthly and annual CO₂ and CH₄ flux estimates from our dataset exhibit magnitudes and variability comparable to previous estimates across key ecosystems (Supplementary Figs. 12-14) and are briefly described below.

7.1 Average monthly fluxes and concentrations

Monthly NEE varied widely, ranging from -81.3 to 122.4 g C m⁻² month⁻¹ (2.5th–97.5th percentiles) and from -621.7 to 3850.0 g C m⁻² month⁻¹ (min–max) across the entire dataset (Fig. 7). The most extreme values were often driven by non-eddy covariance measurements with low data coverage per month. When subset by the five main ecosystems, monthly median NEE fluxes varied from -34.1 to 603.4 g C m⁻² month⁻¹ (Table 6). A few lotic sites in the boreal biome remained unfrozen in January and February and showed very high median fluxes (603.4 g C m⁻² month⁻¹); however it is generally assumed that most lotic, and also lentic, ecosystems are frozen and not actively releasing carbon this time of the year. Monthly median net emissions were highest in lotic ecosystems, followed by lentic ecosystems and October-April months in boreal ecosystems. Strongest net uptake values were found in boreal forests, followed by wetlands and tundra, whereas some aquatic net uptake up to -43.2 g C m⁻² month⁻¹ was observed as well (in total 24 lotic and 219 lentic site-months). Terrestrial ecosystems showed clear seasonal patterns in NEE, with uptake peaking in July (tundra and wetland) or June (boreal forests) and net emissions peaking in October-November (boreal forests and wetland) and September-October (tundra) (Table 6). Seasonal patterns in aquatic NEE were less clear, but net emissions were high throughout most of the year.

Monthly CH₄ flux varied from -104 to 6146 (2.5th–97.5th percentiles) and from -9272 to 145,389 (min–max) mg C m⁻² month⁻¹. The extreme values were primarily associated with aquatic non-eddy covariance measurements, where diffusion and ebullition fluxes were summed to estimate total CH₄ flux. Ebullition fluxes in lentic ecosystems accounted on average for 53% (±33% standard deviation, 308 site-months) of the total CH₄ flux, and for 44% (±40%, 16 site-months) in lotic ecosystems. Six monthly total CH₄ fluxes exceeded expected ranges due to exceptionally high ebullition fluxes (>30 g C m⁻² month⁻¹). Monthly median CH₄ fluxes across key ecosystem types varied from -10.48 to 3712 mg C m⁻² month⁻¹ (Table 7). Monthly median CH₄ emissions were highest in lotic ecosystems, but numbers of site-months were small. Lentic ecosystems and wetlands followed (Fig. 8). Boreal forests were consistent small net CH₄ sinks (median) throughout the July–November period, while tundra remained neutral (0.0 g C m⁻² month⁻¹) or exhibited small net emissions, particularly in spring, autumn, and winter. 14% of tundra (non-wetland) and 15% of boreal forest site-months represented net CH₄ sinks. Emissions from wetlands, lentic and lotic ecosystems showed clear seasonal patterns with emissions peaking in July–August.

Table 6. Monthly median NEE (terrestrial) or CO₂ flux (aquatic) in g C m⁻² month⁻¹ across key ecosystems, with 25th–75th quantiles in parentheses. The terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

Month	Boreal Forest	Tundra	Wetland	Lentic	Lotic
1	12.4 (6.5 to 19.8)	5.3 (1.5 to 9.7)	5.2 (2.5 to 8.4)	2.5 (0.6 to 7.3)	139.4 (59.7 to 386.3)
2	10.1 (4.7 to 15.8)	3.3 (0.3 to 8.4)	4.2 (1.7 to 6.6)	13.8 (2.7 to 22.8)	603.4 (547.8 to 659)
3	8.9 (3.1 to 15.6)	1.6 (0.3 to 6.1)	3.8 (1.3 to 7.2)	2.8 (1.1 to 10.4)	14.8 (6.0 to 57.2)
4	1.5 (-16.9 to 9.8)	1.6 (0.4 to 5.5)	2.4 (0.2 to 6.3)	6.9 (3.9 to 13.4)	50.6 (24.9 to 128.2)
5	-26.0 (-48.2 to -9.9)	4.0 (-0.3 to 9.5)	-2.1 (-11.5 to 3.2)	23.4 (6.9 to 52.8)	41.4 (17.4 to 90.2)
6	-48.0 (-71.9 to -26.1)	-13.8 (-33.3 to 2.8)	-15.3 (-29.8 to -3.1)	11.5 (4.8 to 26.1)	78.9 (35.5 to 195.9)
7	-34.1 (-59.8 to -14.7)	-26.6 (-53.3 to -4.4)	-32.2 (-55.6 to -13.2)	6.1 (2.0 to 16.6)	49.6 (17.9 to 161.3)

8	-11.8 (-33.6 to 7.1)	-11.9 (-27.5 to -1.3)	-17.5 (-33.5 to -8.0)	8.3 (2.3 to 19.9)	62.0 (24.1 to 167.6)
9	11.6 (-4.1 to 25.8)	12.2 (4.7 to 18.0)	1.7 (-5.4 to 10.7)	16.3 (5.2 to 36.8)	63.2 (25.5 to 164.6)
10	20.2 (12.3 to 33.8)	11.6 (7.5 to 17)	9.4 (5.6 to 15.4)	14.9 (5.6 to 25.8)	38.3 (15.0 to 75.3)
11	19.5 (12.1 to 29.7)	8.7 (4.4 to 13.7)	7.9 (5.0 to 11.8)	10.9 (7.4 to 14.8)	41.2 (14.7 to 122.3)
12	14.8 (7.6 to 24.4)	6.8 (2.5 to 11.6)	6.5 (3.7 to 10.1)	2.6 (-0.2 to 7.5)	14.8 (0 to 71.2)

1265

Table 7. Monthly median total CH₄ flux (mg C m⁻² month⁻¹) across key ecosystems, with 25th–75th quantiles in parentheses. The terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

Month	Boreal Forest	Tundra	Wetland	Lentic	Lotic
1	-0.54 (-8.79 to 65.63)	9.6 (9.6 to 80.87)	124.1 (42.5 to 344.4)	78.8 (12.3 to 179.5)	15.5 (10.8 to 18.3)
2	-0.3 (-9.1 to 83.41)	8.67 (8.67 to 45.62)	108.1 (34.6 to 232.8)	54.2 (6.7 to 151.2)	15 (8 to 22)
3	0.35 (-0.26 to 54.32)	0.59 (-0.32 to 3.23)	105.7 (23 to 248.6)	79.7 (7.1 to 177.9)	83 (71 to 179)
4	-0.21 (-14.17 to 3.22)	0.27 (-0.68 to 2.38)	108.2 (25.9 to 349.5)	100.7 (18.5 to 325.4)	80 (37 to 91)
5	0 (-11.75 to 168.75)	124.84 (20.33 to 278.64)	285.3 (52.3 to 869.6)	176.5 (42.9 to 933.1)	331 (89.5 to 3192.1)
6	-0.55 (-26.23 to 134.61)	4.34 (-3.61 to 176.38)	433.9 (64.4 to 1291.7)	392.3 (119.8 to 890)	1123.2 (840 to 2218)

7	-10.48 (-50.84 to 157.47)	-3.31 (-36.41 to 87.45)	929.1 (183.5 to 2503.7)	684.5 (218.5 to 1308)	3712 (335.5 to 17600.3)
8	-10.47 (-37.13 to 95.25)	0 (-12.03 to 373.81)	1156.1 (305.5 to 2445.2)	705.7 (329.9 to 1253)	3071 (1515.5 to 8802.1)
9	-4.55 (-34.72 to 187.88)	82.26 (8.11 to 705.62)	573.8 (125 to 1522.8)	393.5 (162.4 to 734.5)	1161 (513 to 1678)
10	-10.45 (-30.11 to 55.46)	21.4 (3.76 to 143.6)	437.4 (178.6 to 774)	200 (122.4 to 338.2)	264 (158 to 1071)
11	-7.55 (-30.28 to 61.89)	25.65 (-6.13 to 135.63)	264.6 (150.6 to 468.2)	180.7 (5.1 to 252.8)	133 (124 to 133)
12	-3.4 (-20.15 to 36.9)	6.84 (1.08 to 108.55)	196.7 (105 to 387)	194 (3.1 to 277.5)	72 (44.8 to 110.2)

1270




Fig. 7. Monthly NEE (terrestrial) and total CO₂ flux (freshwater) variability across key terrestrial and aquatic classes together with the number of site-months. Note that y axes for the lentic and lotic fluxes follow a pseudo-log scale and represent the CO₂ flux. The terrestrial tundra class characterizes non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

1275




Fig. 8. Monthly CH₄ fluxes across key terrestrial and aquatic classes together with the number of site-months. Note that y axes follow a pseudo-log scale. The tundra classes characterize non-wetland ecosystems in the tundra biome (i.e., dry and moist tundra).

Monthly dissolved CO₂ concentrations varied from 12.6 to 5588 (2.5th–97.5th percentiles) and from 0 to 5480 (min–max) μmol L⁻¹. Monthly median CO₂ concentrations were generally highest in lotic ecosystems and followed a seasonal pattern of lower concentrations during the summer months (Fig. 9). Monthly CO₂ concentrations for lentic systems were highest in spring, but these months also had the smallest sampling sizes. Monthly dissolved CH₄ concentrations varied from 0 to 13.1 (2.5th–97.5th percentiles) with a maximum concentration of 530 μmol L⁻¹. Lotic ecosystems trended towards CH₄ higher concentrations and there were no clear monthly patterns for either freshwater ecosystem.




Figure 9. Monthly dissolved CH₄ and CO₂ concentrations across freshwater classes together with the number of site-months. Note that y axes follow a pseudo-log scale.

7.2 Average annual fluxes

Our annual NEE estimates showed a clear sink-to-source transition from net CO₂ sinks in wetlands (-34.0 g C m⁻² yr) forests (-22.2 g C m⁻² yr), and tundra (-10.7 g C m⁻² yr) to CO₂ sources in lentic (142 g C m⁻² yr) and especially lotic ecosystems (1180 g C m⁻² yr). However, the interquartile range for all three terrestrial classes included positive NEE estimates (Table 8). Annual CH₄ fluxes demonstrated the largest median net CH₄ source for wetlands (5.6 g C m⁻² yr⁻¹) followed by lentic waterbodies (3.1 g C m⁻² yr⁻¹), lotic waterbodies (2.4 g C m⁻² yr⁻¹) and then tundra (0.6g C m⁻² yr⁻¹). Boreal ecosystems were near neutral (0.07 m⁻² yr⁻¹), with an interquartile range between -0.26 and 1.53 g C m⁻² yr⁻¹). It's important to note that these figures are not upscaled estimates and therefore may not accurately reflect the overall sink–source status of the Arctic–boreal region. For example, many freshwater systems exhibit strong CO₂ sink during the summer months (Fig. 8) and our approach using the median values for each ecosystem type likely masks this uptake signal and CH₄ uptake in upland ecosystems. Furthermore, we provide only a simplified estimate of the contribution of ice-out emissions to total annual estimates for freshwater ecosystems.

Table 8. Estimates of annual fluxes (g C m⁻² yr⁻¹) for each broadly defined ecosystem. Values represent the median annual emission estimate followed by first and third quartiles.

Ecosystem	Annual CH ₄ flux (g C m ⁻² yr ⁻¹)	Annual NEE flux (g C m ⁻² yr ⁻¹)
Wetland	5.6 (1.9, 11)	-34.0 (-111, 44.2)
Boreal Forest	0.071 (-0.26, 1.5)	-22.2 (-182, 117)
Tundra	0.62 (0.1, 3.6)	-10.7 (-96, 82.6)
Lentic	3.1 (0.46, 7.1)	142 (64.8, 245)
Lotic	2.4 (1.5, 4.0)	1180 (893, 2050)

8. Discussion

8.1 Future research directions

While a detailed analysis of the underlying mechanisms of flux patterns is beyond the scope of this data description paper, our data compilation offers several new perspectives and research opportunities. For example, ABCFlux v2 opens up for the opportunity to further investigate the detailed characterization of land cover types, waterbody classifications, and disturbance history and with that provides valuable context often overlooked in recent syntheses, which have typically employed coarse classifications (e.g., treating "boreal forest" as a single category, as seen in Virkkala et al., 2021 and Ramage et al., 2024 studies, or aquatic ecosystems split only across permafrost zones as in Song et al., 2024). With ABCFlux v2, it is possible to effectively detect temporal trends, including those in CH₄ fluxes. The inclusion of both CO₂ and CH₄ fluxes also allows for further analysis of their ratios under changing environmental conditions. Moreover, the monthly format provides a clearer understanding of seasonal dynamics, offering an improvement over earlier studies that primarily focused on growing season or annual cumulative fluxes (Kuhn et al., 2021; Ramage et al., 2024; Virkkala et al., 2021).

ABCFlux v2 further allows for an improved understanding of some under-studied flux dynamics. While summertime net CO₂ uptake was previously found at individual sites (Emmerton et al., 2016; Prėskienis et al., 2021), a broader-scale analysis of the underlying conditions and the extent of CO₂ uptake is worth undertaking. This points towards CO₂ and CH₄ sources playing a more complex role in regional carbon budgets than previously understood (Bogard et al., 2019; Tank et al., 2009). Lotic systems exhibit highly variable yet substantial per m² emission rates that, according to our synthesis, appear higher than those of any other ecosystem type. However, the degree to which these elevated emissions occur across entire river networks remains uncertain, as does the influence of local (e.g., steep or shallow sections with high gas transfer velocities and emissions vs. flatter or deeper sections with lower emissions; (Natchimuthu et al., 2017) and circumpolar (e.g., climate gradients) drivers of variability.

Importantly, the overall contribution of lotic systems to regional carbon budgets is constrained by their limited spatial extent—they occupy only about 0.5% of the Arctic–boreal domain (Olefeldt et al., 2021b). As a result, despite high per m² fluxes, total emissions from rivers may be smaller than those from more widespread ecosystems such as wetlands, which have lower per m² emissions but a much greater areal extent (Casas-Ruiz et al., 2023). Nevertheless, ABCFlux v2 offers a valuable opportunity to address these aquatic knowledge gaps.

Our dataset also shows distinctive patterns in the seasonal dynamics of CH₄ exchange across different ecosystems. Particularly noteworthy is the contrast between non-wetland tundra and boreal forest systems: tundra areas function as net CH₄ sinks in ABCFlux v2 for only a single month, while boreal forest ecosystems maintain CH₄ uptake throughout the summer period as well as autumn period (Fig. 8).

8.2 Remaining gaps

ABCFlux v2 demonstrates clear improvements in carbon flux data quantity over time but also shows that some ecosystems (e.g., deciduous needleleaf forests, barren and sparsely vegetated ecosystems, large lentic ecosystems) or periods (non-growing season, and especially the challenging spring ice-off period in aquatic ecosystems) still remain poorly captured. Moreover, the studied sites are heavily clustered in a few regions in Alaska and Fennoscandia, therefore leaving significant spatial gaps in coverage in spite of the relatively high absolute number of locations. In particular, despite Russia's large land mass (close to 60% of the domain), data from Russia only comprise 10.8% of the dataset in terms of site-months and 15.3% of sites in ABCFlux v2 (with similar representativeness among terrestrial and aquatic data), making this region a critical data gap. At local scales, flux data across lentic depth zones and throughout entire lotic networks are critically needed to better estimate spatially representative waterbody fluxes. Furthermore, most freshwater measurements were taken during daytime, leaving nighttime dynamics less constrained. To more reliably estimate the net ecosystem carbon balance, incorporating lateral fluxes into future studies would be a large benefit, especially in landscapes affected by permafrost thaw (Zolkos et al., 2022).

Additionally, some specific flux mechanisms and environmental controls remain undersampled. This is particularly true for ebullition, which only makes up 12% of aquatic CH₄ fluxes in the dataset but can account for up to 90% of total aquatic CH₄ emissions (Walter et al., 2010; Kuhn et al., 2021). Moreover, plant-mediated CH₄ emissions in aquatic systems and CH₄ fluxes from terrestrial trees are not explicitly partitioned in the dataset; instead, they are included as part of total fluxes measured primarily by eddy covariance. Improved partitioning of these flux pathways would enhance our understanding of the processes driving CH₄ dynamics and help to better constrain landscape-scale carbon budgets (Gauci et al., 2024; Iwata et al., 2018; Juutinen et al., n.d.; Kankaala et al., 2005; Kyzivat et al., 2022). Important site-level environmental data such as thaw depth, soil carbon stocks, river width, and comprehensive information on plant and microbial communities are partly or entirely missing, yet would provide valuable insights into the processes governing fluxes. Likewise, more detailed disturbance metrics and a stronger

integration of disturbed sites into flux monitoring networks are needed to better capture the impacts of these changes on carbon fluxes and budgets.

Although we excluded chamber measurements for CO₂ fluxes taken underneath the canopy of forests, as chambers do not capture the full ecosystem dynamics it is notable that these chamber measurements could be valuable end points and quality assessments when co-located with towers, providing a lower limit on respiration estimates. We did include below canopy chamber CH₄ flux measurements under the assumption that CH₄ fluxes from trees are negligible. However, recent evidence suggests that trees may play a globally significant role in CH₄ uptake, even considering that CH₄ uptake by trees decreases with increasing latitude and approaches zero in low mean annual temperatures (Gauci et al., 2024; Sundqvist et al., 2012). Furthermore, some studies suggest that boreal trees emit CH₄ from tree stems (Klaus et al., 2024; Machacova et al., 2023; Vainio et al., 2022), suggesting more tree-based flux measurements are needed and should be incorporated into future synthesis efforts. Furthermore, given the low data coverage of 2% for stable isotope CO₂ and CH₄ measurements among the aquatic data in ABCFlux v2, the lack of our understanding of emission pathways and sources becomes apparent. Therefore, increasing the number of observations across ecosystems would significantly improve the source attribution of emissions as well as our process understanding.

While long-term, year-round terrestrial CO₂ flux sites have become more common, the need remains to expand the network to more aquatic and under-represented terrestrial sites where year-round measurements of both CO₂ and CH₄ fluxes are collected simultaneously. A larger number of sites measuring these fluxes throughout the year would also inform and improve wintertime process understanding. Long-term CH₄ flux monitoring sites remain scarce, hindering our ability to detect temporal trends in CH₄ emissions—despite their potentially critical role in a changing climate (Turetsky et al., 2020). Thus, maintaining existing sites and setting up new permanent CH₄ flux sites is critical for accurate understanding on the changing carbon cycle in Arctic-boreal regions. The availability of recent, “real-time”, data remains challenging as often post-processing times can delay the release of the data. Improving the turn-around time and associated pipelines from data collection to availability will improve future synthesis and modeling efforts.

An additional notable data gap is the lack of information on nitrous oxide (N₂O), another significant GHG that is currently absent from the ABCFlux v2 dataset, which primarily focuses on CO₂ and CH₄. This omission reflects a broader data gap, despite growing evidence that N₂O emissions may become increasingly important for regional and global climate feedbacks as permafrost thaw accelerates (Voigt et al., 2017). As N₂O has a global warming potential over 300 times that of CO₂ over a 100-year timescale, even relatively small fluxes may significantly contribute to climate feedbacks. Incorporating N₂O measurements in future efforts would be critical to achieving a more comprehensive understanding of GHG dynamics in the Arctic-boreal domain.

8.3 The importance of regional networks

While contributing data to global repositories remains highly encouraged, regional syntheses like this provide valuable insights into network status and development needs across the Arctic-boreal domain. In our case, it has helped in establishing a collaborative community that bridges terrestrial and aquatic researchers while integrating CO₂ and CH₄ flux data in a standardized format, creating a foundation for collectively improving Arctic-boreal carbon cycle understanding. Looking ahead, we strongly advocate for maintaining a community-driven approach in future flux syntheses, similar to earlier ABCFlux initiatives (See et al., 2024; Virkkala et al., 2022, 2025) and those supported by, for example, the Permafrost Carbon Network (Schuur et al., 2015, 2022). Including data contributors as co-authors not only ensures proper recognition of their contributions but also draws on their expertise in data interpretation. Moreover, there is potential for efforts like ABCFlux to gradually evolve into more dynamic, continuously updated resources. While not yet realized, the aspiration to create a database capable of integrating recent flux measurements and associated metadata remains important for enabling iterative updates (e.g. <https://fluxnet.org/fluxnet-data-system/>) and the incorporation of new knowledge (see e.g., (Arctic Terrestrial Carbon Cycling, 2025)). Ultimately, such a collaborative and adaptive approach enhances both the technical quality and scientific context of carbon flux syntheses, helping to advance our understanding of high-latitude carbon dynamics in a rapidly changing environment.

9. Data use guidelines

ABCFlux v2 data is categorized into two usage tiers depending on the data source, as indicated in the "data_usage" column. Tier 1 data is open and free to use for scientific and educational purposes. In contrast, Tier 2 requires that data users give data producers the opportunity to collaborate and consult with them. However, it is recommended that all researchers reach out and collaborate with the dataset developers and relevant site data producers when using the dataset as a core component of their analysis. This encourages a more informed and context-rich use of the data.

If used, the dataset should be referenced by citing both this paper and the dataset (DOI: 10.3334/ORNLDAAAC/2448).

10. Data availability

The dataset associated with this publication can be found in ORNL DAAC at <https://doi.org/10.3334/ORNLDAAAC/2448>

11. Conclusions

ABCFlux v2 provides the most comprehensive dataset of surface-atmosphere Arctic-boreal ecosystem CO₂ and CH₄ fluxes to date. It is particularly useful for machine learning or regression modeling for drivers of fluxes, process model tuning, remote sensing-based

upscaling, and empirical studies aiming to understand carbon budgets and regional variability in flux magnitudes, as well as changes in fluxes through time.

1465 Author contributions

The ABCFlux v2 dataset was conceptualized and developed by a team led by AMV, IW, JV, and MAK, supervised by BR, MG, and SN, and further supported by KA, KK, GRR, JW, DO, EAGS, DB, and ROK. IW, JV, MAK, AMV, SM, and TW compiled the data. AMV drafted and coordinated the manuscript in close collaboration with IW, MAK, JV, SN, BR, MG, KA, KK, 1470 GRR, JW, DO, EAGS, DB and ROK. Other authors contributed data to the ABCFlux v2 dataset and reviewed the manuscript. Data contributors whose data were extracted from publications are not coauthors in this paper, unless new data were provided.

Acknowledgements

1475 This work was supported by funding catalysed by the TED Audacious Project (Permafrost Pathways) as well as funding through the Gordon and Betty Moore Foundation (grant no. 8414), the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 951288, Q-Arctic) and under the Horizon Europe Climate programme (grant agreement no. 101056921, GreenFeedBack), and the Natural 1480 Sciences and Engineering Research Council (NSERC) Discovery Grant (DGECR-2025-00337). We thank ICOS, Ameriflux, Fluxnet Canada and the Canadian Carbon Program (CCP), FLUXNET, and AsiaFlux networks for openly sharing data for our synthesis. We thank Christina Shintani for her work creating the maps shown in Figure 1 and Supplementary Figure 8.

1485 Adrian Rocha was funded by NSF# 2103539. Aino Korrensalo was funded by the Research Council of Finland, grant number 338980. Aki Vähä was funded by the Research Council of Finland (ICOS-FIRI and project no. 322432), ICOS-FI via University of Helsinki funding, and the EU Horizon Europe Framework Programme for Research and Innovation (GreenFeedback grant no. 101056921). Alan Barr acknowledges the support of our boreal flux sites from the 1490 Fluxnet Canada Research Network (FCRN), the Canadian Carbon Program (CCP), and the Changing Cold Regions Network (CCRN) research networks funded by the Natural Sciences and Engineering Research Council of Canada (NSERC). Aleksandr F. Sabrekov, Elena Lapshina, and Egor Dyukarev's work was part of the most important innovative project of national importance "Development of a system for ground-based and remote monitoring of 1495 carbon pools and greenhouse gas fluxes in the territory of the Russian Federation, ensuring the creation of recording data systems on the fluxes of climate-active substances and the carbon budget in forests and other terrestrial ecological systems" (registration no. 123030300031-6) and the IMCES SB RAS work was supported by the basic state project FWRG-2021-0001 (No121031300154-1). Aleksi Räsänen and Tarmo Virtanen was funded by the Research 1500 Council of Finland (grant number 308513). Alex Mavrovic and Alexandre Roy would like to warmly thank the Indigenous communities who have welcomed us and shared their territory with

us in Iqaluktuuttiaq and Inuvik. AM and AR's dataset was acquired thanks to the contributions of the Natural Sciences and Engineering Research Council of Canada (NSERC), the Fonds de recherche du Quebec - Nature et technologies (FRQNT) and Polar Knowledge Canada (POLAR). Alexey V. Panov was funded by the State Assignment FWES-2024-0023. Amy Townsend-Small was funded by NSF: AON-1107607. Anatoly Prokushkin was funded by the State Assignment FWES-2024-0023. Anders Lindroth acknowledges the European Union, Formas, Swedish Research Council. Andreas Christen acknowledges support by the Canada Foundation for Innovation – IF 2015 (grant no. 33600), NSERC Discovery Grants Program (grant no. RGPIN-2017-03958), NSERC Discovery Grants – Northern Research Supplement (grant no. RGPNS-503529), and NSERC Discovery Grants Program – Accelerator Supplement (grant no. RGPAS-507854). Andreas Westergaard-Nielsen was funded by Villum foundation (project no. 42069) and the Danish Ministry of Climate, Energy and Utilities. Andrew Black and Warren Helgason are grateful for the support of our boreal flux sites from the Fluxnet Canada Research Network (FCRN), the Canadian Carbon Program (CCP), and the Changing Cold Regions Network (CCRN) research networks funded by the Natural Sciences and Engineering Research Council of Canada (NSERC). Anne Ojala acknowledges University of Helsinki for providing access to field stations as well making it possible to use the facilities with the help of highly skilled staff. Armando Sepulveda-Jauregui, Maialen Barret, and Frederic Thalasso acknowledge European project ERANET-LAC "METHANOBASE" ELAC2014-DCC-0092-A. Avni Malhotra was funded by NSERC Discovery grant, NSF REU, NASA Terrestrial Ecology Program (project 80NSSC22K1253). Ayumi Kotani acknowledges JSPS KAKENHI 19H05668, ArCS (JPMXD1300000000), ArCSII (JPMXD1420318865). Bo Elberling acknowledges the Danish National Research Foundation, Center for Permafrost, CENPERM DNR100 (B.E). The work of Carl-Fredrik Johannesson, Jenni Nordén, Klaus Steenberg Larsen and Hanna Silvennoinen was in relation to the ForBioFunCtioN project, funded by the Research Council of Norway (grant number 303006 to JN). They are grateful to Esben Kirk Hansen and colleagues at Bymiljøetaten of the municipality of Oslo, and Holger Lange, PI at the ICOS-Norway site Hurdal, for granting us research permits and for providing help with practical issues concerning the field data collection. Carlo Trotta was funded by Horizon Europe (Copernicus Climate Change Service Evolution (CERISE)- project, 101082139). Carolina Voigt was funded by Canada Foundation for Innovation project Changing Arctic Network (CANet, grant no. 33661); the Canada Research Chairs Program (CRC-2018-00259 to Oliver Sonnentag); the European Research Council Starting Grant COLDSPOT (no. 101163177 to Carolina Voigt). Catherine Dieleman and Hailey Webb acknowledge the Bonanza Creek Long-Term Ecological Research Site. Cheristy P. Jones work was supported by the Department of Energy (DE-SC0023456), the EMERGE BII (NSF-DBI 2022070), the Swedish Polar Research Secretariat and SITES for the support of the work done at the Abisko Scientific Research Station (SRC 4.3-2021-00164), the FLUXNET Secondment Program, the Greenland Ecosystem Monitoring program (<https://g-e-m.dk/>) for work done at the Kobbefjord Research Station, the University of New Hampshire CEPS First Year PhD Fellowship and the CARPE NRT Fellowship (NSF #2125868). Christina Biasi was supported by the Austrian Science Fund (FWF project PERNO; no. 10.55776/M3335) and the Academy of Finland through multiple projects, including N-PERM (no. 341348), NOCA (no. 314630), and the Yedomia-N project (no. 287469). Christopher D. Arp received funding from NSF: AON-1107607. Claire Treat, Lona van Delden, and Josh Hashemi acknowledge ERC

StG FluxWIN 851181. Craig A. Emmerton acknowledges Natural Sciences and Engineering Research Council (NSERC; V.L.S.L. and C.A.E.). Danil V. Ilyasov and Anastasia Niyazova's work was supported by the Russian Science Foundation, project number 25-17-20042. Dario Papale thanks the support of the ITINERIS project (IR0000032) Funded by EU - Next

1550 Generation EU PNR. David Bastviken and associated data contributions were funded by the European Research Council (ERC H2020 grant 725546), the Swedish Research Councils VR (grant 2022-03841), and Formas (grant 2018-01794). Donatella Zona was funded by NSF award number 1932900, and 2149988. Edward A. G. Schuur was funded by: National Science

1555 Flagship, ICOS-Finland, Research Council of Finland through project and infrastructure funding (grant numbers 337064 and 330840). Efrén López-Blanco, Rasmus Jensen, Torben R. Christensen, and Mikhail Mastepanov's work was supported by the Greenland Ecosystem Monitoring (g-e-m.dk) funded by the Danish Environmental Protection Agency and the Danish Energy Agency. ELB, TRC and MM consider this study a contribution to GreenFeedBack

1560 (Greenhouse gas fluxes and earth system feedbacks) funded by the European Union's HORIZON research and innovation program under grant agreement No 101056921. Miquel A. Gonzalez-Meler and Elena Blanc-Betes acknowledge the Toolik Lake Field Station staff and the CH2MHill Polar Services for logistical support and their research was funded by the DOE, Terrestrial Ecosystem Science Program (DE-SC 0006607) while experimental site

1565 establishment and maintenance was funded by NSF Office of Polar Programs (Award numbers 9321730, 9617643, 0856728, 1504141). Elena Veretennikova acknowledges the Russian Academy of science in the framework of basic research - No121031300154-1. Eleonora Canfora was funded by Horizon Europe (Open-Earth-Monitor Cyberinfrastructure project, 101059548). Elisa Männistö acknowledges the ACCC Flagship, ICOS-Finland, Research

1570 Council of Finland through project and infrastructure funding (grant numbers 337064 and 330840). Emily H. Stanley acknowledges that the assembly of GRiMeDB was supported by the U.S. National Science Foundation (grant no. DEB-2025982, NTL LTER), the Canada Research Chair program, the Swedish Research Council (Vetenskapsradet) (grant nos. 2021-06667 and 2021-04058), and Svenska Forskningsradet Formas (grant no. 2019-01105). Erik Lundin

1575 acknowledges the Abisko Scientific Research Station & ICOS SE, funded by the Swedish Research Council (SRC) and the consortium partners since 2010. Erik Sahlée's and Anna Rutgersson's work was sponsored by the Swedish research council FORMAS project number: 2009-1692.. Eugenie Euskirchen was funded by NSF Grants DEB LTREB 1354370 and 2011257, DEB-0425328, DEB-0724514, and DEB-0830997 as well as US Geological Survey

1580 Climate R&D program. Fenix Garcia Tigreros was funded by NASA ABoVE Project 80NSSC19M0104 and the U.S. Geological Survey Biological Carbon Sequestration Program. Francois Clayer's work was supported by NIVA core funding (Research Council of Norway, contract #342628/L10) Global Change at Northern Latitudes and GreenSense projects, as well as the BIOGOV project (Research Council of Norway, project #323945). Frans-Jan W.

1585 Parmentier was supported by the research council of Norway (project nrs. 323945 & 352260). Frédéric Bouchard was funded by and acknowledges the Natural Sciences and Engineering Research Council (NSERC), the Polar Continental Shelf Program (PCSP), and the W. Garfield Weston Foundation. Geert Hensgens and Jorien Vonk were supported by the Netherlands Earth Science System Centre (NESSC), which was funded by the Dutch Ministry of Education,

1590 Culture and Science (OCW) (grant no. 024.002.001). George W. Kling's work was supported by grants from U.S. NSF AON 1936769, DEB 2224743. Gerard Rocher-Ros was funded by the European Union (ERC, ARIMETH, 101161308). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held

1595 responsible for them. Greg Henry acknowledges the Natural Sciences and Engineering Research Council of Canada, Polar Continental Shelf Project, Canada Foundation for Innovation. Hanna Lee acknowledges the Horizon Europe project (IM4CA: 101183460). Hannu Nykanen's work was funded by the European Union contract (ENV4-CT97-0583) - CONGAS project and TUNDRA (ENV4-CT97-0522), and the Arctica projects was funded by the Academy

1600 of Finland. Hannu Nykanen acknowledges Kevo Subarctic Research Institute at Utsjoki for logistic support during the field work. Helena Rautakoski was funded by the European Union - NextGenerationEU instrument and from the Research Council of Finland (grant no. 337552, 347794, 324259). Henrique Oliveira Sawakuchi acknowledges that this project has received funding from the European Research Council under the European Union's Horizon 2020

1605 research and innovation program (Grant Agreement No. 725546), the Knut and Alice Wallenberg Foundation (Grant 2016.0083), the Knut and Alice Wallenberg Foundation (Wallenberg Academy Fellowship) to DS, the Swedish Research Council (Grant 2016-04829), and FORMAS (the Swedish Research Council for Sustainable Development (Grant 2018-01794). Henrique Oliveira Sawakuchi thanks Shun Koizumi, Isolde Puts, and Hendricus

1610 Verheijen (Umea University) for help measuring oxygen and temperature profiles, and Erik Geibrink (Umea University) for his help setting up the experiment. Hideki Kobayashi was funded by JSPS JP25H00454, ArCS-3. Housen Chu acknowledges that AmeriFlux data were made available through the data portal (<https://ameriflux.lbl.gov>) and processing maintained by the AmeriFlux Management Project, supported by the U.S. Department of Energy Office of Science,

1615 Office of Biological and Environmental Research, under contract number DE-AC02-05CH11231. Isabelle Laurion acknowledges the Network of Centres of Excellence program ArcticNet, Fonds quebécois de la recherche sur la nature et les technologies (today: Fonds de recherche du Quebec - Nature et technologies, FRQNT), Natural Sciences and Engineering Research Council of Canada (CRSNG), Natural Resources Canada Polar Continental Shelf Program,

1620 International Polar Year, CREATE program EnviroNorth, W. Garfield Weston Foundation, Indian and Northern Affairs Canada (NSTP program of Canadian Gov, today: managed by Polar Knowledge Canada), Center for Northern Studies, Parks Canada. Ivan Mammarella acknowledges financial support from Research Council of Finland (NPERM project nr 341349, ICOS-FIRI), ICOS-FI via University of Helsinki funding, EU-INTERACT, the EU Horizon Europe

1625 +/- Framework Programme for Research and Innovation (GreenFeedback nr. 101056921 and LiweFor nr. 101079192). Jacqueline Knutson was supported by NIVA core funding (Research Council of Norway, contract #342628/L10) Global Change at Northern Latitudes and GreenSense projects, as well as the BIOGOV project (Research Council of Norway, project #323945). Järvi Järveoja acknowledges Formas (grant no. 2021-00611). Jeffrey M. Welker was

1630 funded by NSF Arctic System Science & Arctic Observing Network proposals 9321730, 9617643, 0856728, 1504141, 1836873 & JMW's UArctic Research Chairship. Ji Young Jung acknowledges RS-2025-24683148 (KOPRI-PN25010). Jo Snöäliv acknowledges Stockholm University and Abisko Scientific Research Station (ANS). Joachim Jansen acknowledges The

1635 Abisko Scientific Research Station (ANS) and the Swedish Polar Research Secretariat. John Kochendorfer's work was supported by NSF grant 1203583. Jukka Pumpanen was funded by the Research Council of Finland (project number 286685). Julia Kelly's research was funded by the Swedish Research Council FORMAS grant 2018-02700 and the Swedish Research Council FORMAS grant 2019-00836, the Crafoord foundation grant 20190763, Skogssallskapet Stina Werner Fond grant 2021-094, the Royal Physiographic Society of Lund and the Swedish
 1640 government through the Strategic Research Areas BECC (Biodiversity and Ecosystem Services in a Changing Climate) and MERGE (Modelling the Regional and Global Earth system). Julie Talbot and Daniel F. Nadeau's research was funded by the Natural Sciences and Engineering Research Council of Canada (NSERC) through Grant RDCPJ508080-16 entitled "Observation and modelling of net evaporation from a boreal hydroelectric complex (water footprint)". Juliya
 1645 Kurbatova's study was done with the support of the state assignment of IPEE RAS (Theme No. 1022033100172-2-1.6.19). June Skeeter would like to thank the staff at the Aurora Research Institute in Inuvik for providing logistical support (especially Edwin Amos, for the local expertise and invaluable assistance he provided in the field), Rick Ketter for helping to design and install the floating EC system, and the Polar Continental Shelf Program, Natural Resources Canada
 1650 for additional logistical support. June Skeeter also acknowledges support by the Canada Foundation for Innovation – IF 2015 (grant no. 33600), NSERC Discovery Grants Program (grant no. RGPIN-2017-03958), NSERC Discovery Grants – Northern Research Supplement (grant no. RGPNS-503529), and NSERC Discovery Grants Program – Accelerator Supplement (grant no. RGPAS-507854). Kajar Koster was supported by the Research Council of Finland
 1655 (Project Nos. 286685, 294600, 307222) and the Kone Foundation. Karel Castro-Morales' project, PROPERAQUA, was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project No. 396657413. Katey Walter Anthony was supported by NSF NNA 2022577 and a gift from Tito's Handmade Vodka. Katrin Attermeyer and Pascal Bodmer thank the EuroRun team. Kuno Kasak was supported by the Estonian Research Council grant
 1660 nr PSG714 and by the Estonian Ministry of Education and Research, Center of Excellence for Sustainable Land Use (FutureScapes, TK232). Kyra St. Pierre acknowledges the Danish National Research Foundation, Natural Sciences and Engineering Research Council, Canadian Initiative for Nordic Studies. Lance Lesack was funded by NSERC (Discovery Grant and Northern Research Supplement programs to Lesack), the Polar Continental Shelf Project
 1665 (helicopter support to Lesack), and from the Northern Scientific Training Program, Polar Knowledge Canada. Logistical, technical, and financial support has been provided by the Aurora Research Institute (Western Arctic Research Centre). Lance Lesack's published data remains the joint intellectual property of his publication co-authors and Simon Fraser University (SFU). SFU's Summit repository is being used to comply with NSERC's open access publication
 1670 and data policy. Lea Cabrol acknowledges Millennium Science Initiative Program ICN2021_002 (Millenium Institute BASE) from the Chilean National Agency of Research and Innovation, ANID European project ERANET-LAC "METHANOBASE" ELAC2014-DCC-0092-A. Macall Hock was funded by NSF award number 1932900. Maija E. Marushchak was financed by the EU 6th Framework Programme project CARBO-North (no. 036993) and Research Council of Finland
 1675 project Thaw-N (no. 349503). Manuel Helbig and Janna Heerah acknowledge support through the Natural Sciences and Engineering Research Council of Canada (NSERC) Grant ALLRP 561142-20. Marcin Antoni Jackowicz-Korczynski acknowledges the Greenland Ecosystem

Monitoring (GEM) program and the Integrated Carbon Observing System (ICOS) as funded by the Danish Ministry of Climate, Energy and Utilities. Margaret Torn's work on US-NGB and US-NGC was part of The Next-Generation Ecosystem Experiments (NGEE) Arctic project supported by the Office of Biological and Environmental Research in the Department of Energy Office of Science. Mariasilvia Giamberini and Marta Magnani acknowledge Italian National Research Council (CNR), CNR-DSSTA and CNR-ISP for the logistics in the Ny Ålesund research base. Masahito Ueyama acknowledges the Arctic Challenge for Sustainability II (ArCS II; JPMXD1420318865) and ArCS III (JPMXD1720251001). Mats P. Björkman acknowledges the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie (657627), FORMAS - Swedish Research Council for Sustainable Development (2016-01187 and 2022-00786), the Swedish research council (2021-04011), and the strategic research environment BECC - Biodiversity and Ecosystem services in a Changing Climate. Matthias Peichl, Mats B. Nilsson, Koffi Dodji Noumonvi, and Järvi Järveoja acknowledge funding from the Swedish Research Council (VR, grant no. 2018-03966 and 2019-04676), the Kempe Foundation (grant no. JCK-1712, JCSMK23-0221) and support from the Swedish Infrastructure for Ecosystem Science (SITES) and the Swedish Integrated Carbon Observation System (ICOS-Sweden). Mika Aurela, Mika Korkiakoski, and Annalea Lohila's work was financially supported by the Ministry of Transport and Communications through the Integrated Carbon Observing System (ICOS) research and the ACCC flagship funded by the Academy of Finland (grant no. 337552). Mikhail V. Glagolev's study was conducted under the state assignment of Lomonosov Moscow State University. Namyi Chae was supported by the NRF-RS-2025-24683148. Natascha Kijun's research was funded by the Swedish Research Council FORMAS grant 2018-02700 and the Swedish Research Council FORMAS grant 2019-00836, the Crafoord foundation grant 20190763, Skogssällskapet Stina Werner Fond grant 2021-094, the Royal Physiographic Society of Lund and the Swedish government through the Strategic Research Areas BECC (Biodiversity and Ecosystem Services in a Changing Climate) and MERGE (ModELLing the Regional and Global Earth system). Nigel Roulet acknowledges NSERC Discovery. Norbert Pirk was supported by the European Research Council (project #101116083), the Norwegian Research Council (project #323945), the Norwegian Environment Agency, and the strategic research initiative LATICE (#UIO/GEO103920). Oliver Sonnentag acknowledges infrastructure funding from the Canada Foundation for Innovation project Changing Arctic Network (CANet, grant no. 33661, awarded to P.M.), and further financial support through the Canada Research Chair (CRC-2018-00259) and NSERC Discovery Grants programs (DGPIN-2018-05743), ArcticNet, a Network of Centres of Excellence Canada (grant no. P216), the Canada First Research Excellence Fund's Global Water Futures program (Northern Water Futures), and the Polar Continental Shelf Program. Patrick Crill was funded by Vetenskapsrådet (Swedish Research Council), Stockholm University, USNSF, USDOE and assisted by Swedish Polar Research. Pierre Tailladat was funded by the Natural Sciences and Engineering Research Council of Canada and Hydro-Quebec to Michelle Garneau (RDCPJ 514218-17). Pirkko Kortelainen acknowledges the Finnish Environment Institute. Praveena Krishnan was supported by an NSF grant 1848694. Qinxue Wang acknowledges monitoring systems were supported by ERTDF (2E-1203), Ministry of the Environment Government of Japan. Riikka Rinnan acknowledges that Abisko Stordalen birch forest data collection was funded by Danish National Research Foundation (grant No. DNRF168), European Research

Council (grant No. 771012), and Independent Research Fund Denmark (grant No. 1026-00127B). Robert Wagner was funded by the NSF Office of Polar Programs (award number 1204263) with additional logistical support funded by the Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE), an Earth Ventures (EV-1) investigation, under contract with the National Aeronautics and Space Administration (NASA) and acknowledges this research was conducted on land owned by the Ukpeagvik Inupiat Corporation (UIC). Roger Seco acknowledges a Ramon y Cajal Grant (RYC2020-029216-I) funded by MICIU/AEI/10.13039/501100011033 and by "ESF Investing in your future", and project PID2021-122892NA-I00 funded by MICIU/AEI and by "ERDF A way of making Europe". Sang-Jong Park acknowledges RS-2025-24683148 (KOPRI-PN25010; S.-J.P.). Sari Juutinen was funded by the Research Council of Finland. Scott J. Davidson's work at Barrow-BEO, Barrow-BES, Atqasuk and Ivotuk sites was funded by the Office of Polar Programs of the National Science Foundation (NSF) awarded to Donatella Zona (award number 1204263) and Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE), an Earth Ventures (EV-1) investigation, under contract with the National Aeronautics and Space Administration, and a Royal Society International Exchange awarded to Donatella Zona (2013/R1), and a NERC Arctic Research Programme, CYCLOPS Grant (NE/K00025X/1) to Gareth K. Phoenix. This research was conducted on land owned by the Ukpeagvik Inupiat Corporation (UIC) and the Arctic Slope Regional Corporation (ASRC). For Scott J. Davidson's work at Poplar Fen, IPAD and Carmon Creek sites he would like to acknowledge that this research takes place within the boundaries of Treaty 8, traditional lands of the Dene and Cree, as well as the traditional lands of the Metis of northern Alberta. The University of Waterloo is located on the traditional territory of the Neutral, Anishnaabeg, and Haudenosaunee Peoples. The University of Waterloo is situated on the Haldimand Tract, land promised to Six Nations, which includes six miles on each side of the Grand River. Scott J. Davidson's research is part of the Boreal Ecosystem Recovery and Assessment (BERA) project (www.bera-project.org), and was supported by the Natural Sciences and Engineering Research Council of Canada Alliance Grant (ALLRP 548285 - 19) awarded to Maria Strack in conjunction with Alberta-Pacific Forest Industries Inc., Canadian Natural Resources Ltd., Cenovus Energy, ConocoPhillips Canada Resources Corp., Imperial Oil Resources Ltd., Canadian Forest Service's Northern Forestry Centre, and Alberta Biodiversity Monitoring Institute. Sean Carey thanks Dr. Gordon Drewitt for initial data processing, Dr. Michael Treberg for instrument maintenance and calibration and Tyler de Jong, David Barrett and Rosy Tutton for field assistance and acknowledges the continued support of the Water Resources Branch, Government of Yukon, for the operation of Wolf Creek Research Basin. Sean Carey was funded by the National Science and Engineering Council of Canada through Discovery Grants and the Global Water Futures Program. Sigrid Dengel acknowledges the work on US-NGB and US-NGC was part of The Next-Generation Ecosystem Experiments (NGEE) Arctic project supported by the Office of Biological and Environmental Research in the Department of Energy Office of Science. Sofie Sjogersten acknowledges the University of Nottingham Faculty of Science Pump-Prime Research Accelerator Fund and is grateful for the fieldwork support from Mattias Dalkvist, Veronica Escubar Ruiz and staff at the Abisko Scientific Research Station. Sujan Pal and Ryan Sullivan acknowledge observations from the Atmospheric Radiation Measurement (ARM) user facility are supported by the U.S. Department of Energy Office of Science user facility managed by the Biological and Environmental

Research Program. Work at Argonne National Laboratory was supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under contract DEAC0206CH11357. Suzanne E. Tank acknowledges financial support from the Natural Sciences and Engineering Research Council of Canada to SET, POLAR Knowledge Canada (Taiga Plains and Taiga Shield data) and Polar Continental Shelf Program (Peel Plateau data). Tetsuya Hiyama was funded by the Grants-in-Aid for Scientific Research (Grant Numbers JP19H05668 and JP25H00507) of the Japan Society for the Promotion of Science (JSPS). Tim Papakyriakou was funded by the Natural Sciences and Engineering Research Council of Canada to TNP. Timo Vesala acknowledges University of Helsinki for providing access to field stations as well, making it possible to use the facilities with the help of highly skilled staff. Trofim Maximov and Roman Petrov's work was carried out within the framework of the state assignment of the Ministry of Education and Science of Russia under the project 'Study of biogeochemical cycles and adaptive reactions of plants of boreal and arctic ecosystems of the north-east of Russia', state registration number AAAA-A21-121012190034-2. Tuula Larmola acknowledges the Research Council of Finland. Vincent E.J. Jassey and Anna Sytiuk were supported by the MIXOPEAT project (Grant No. ANR-17-CE01-0007 to VEJJ) and thank the Swedish Polar Research Secretariat and SITES for the support of the work done at the Abisko Scientific Research Station. SITES is supported by the Swedish Research Council's grant 4.3-2021- 00164. Vincent L. St. Louis acknowledges Natural Sciences and Engineering Research Council (NSERC; V.L.S.L. and C.A.E.). Walter Oechel was funded by NOAA/EPP Grant # NA16SEC4810008. Ruth Varner was funded National Science Foundation grant EAR-1063037 EF-1241037 and DBI-2022070, National Aeronautics and Space Administration grant NNX17AK10G, Department of Energy grants DE-SC0016440 and DE-SC0023456, and Swedish Research Council Vetenskapsrådet grant 2019-05764.

References

- Arndt, K. A., Hashemi, J., Natali, S. M., Schiferl, L. D., and Virkkala, A.-M.: Recent Advances and Challenges in Monitoring and Modeling Non-Growing Season Carbon Dioxide Fluxes from the Arctic Boreal Zone, *Current Climate Change Reports*, 9, 27–40, <https://doi.org/10.1007/s40641-023-00190-4>, 2023.
- Attermeyer, K., Casas-Ruiz, J. P., Fuss, T., Pastor, A., Cauvy-Fraunié, S., Sheath, D., Nydahl, A. C., Doretto, A., Portela, A. P., Doyle, B. C., Simov, N., Gutmann Roberts, C., Niedrist, G. H., Timoner, X., Evtimova, V., Barral-Fraga, L., Bašić, T., Audet, J., Deininger, A., Busst, G., Fenoglio, S., Catalán, N., de Eyto, E., Pilotto, F., Mor, J.-R., Monteiro, J., Fletcher, D., Noss, C., Colls, M., Nagler, M., Liu, L., Romero González-Quijano, C., Romero, F., Pansch, N., Ledesma, J. L. J., Pegg, J., Klaus, M., Freixa, A., Herrero Ortega, S., Mendoza-Lera, C., Bednařík, A., Fonvielle, J. A., Gilbert, P. J., Kenderov, L. A., Rulík, M., and Bodmer, P.: Carbon dioxide fluxes increase from day to night across European streams, *Commun. Earth Environ.*, 2, 1–8, <https://doi.org/10.1038/s43247-021-00192-w>, 2021.
- Baldocchi, D., Chu, H., and Reichstein, M.: Inter-annual variability of net and gross ecosystem carbon fluxes: A review, *Agric. For. Meteorol.*, 249, 520–533, <https://doi.org/10.1016/j.agrformet.2017.05.015>, 2018.

- 1810 Bastviken, D., Cole, J., Pace, M., and Tranvik, L.: Methane emissions from lakes: Dependence of lake characteristics, two regional assessments, and a global estimate: LAKE METHANE EMISSIONS, *Global Biogeochem. Cycles*, 18, <https://doi.org/10.1029/2004gb002238>, 2004.
- 1815 Battin, T. J., Lauerwald, R., Bernhardt, E. S., Bertuzzo, E., Gener, L. G., Hall, R. O., Jr, Hotchkiss, E. R., Maavara, T., Pavelsky, T. M., Ran, L., Raymond, P., Rosentreter, J. A., and Regnier, P.: River ecosystem metabolism and carbon biogeochemistry in a changing world, *Nature*, 613, 449–459, <https://doi.org/10.1038/s41586-022-05500-8>, 2023.
- Belshe, E. F., Schuur, E. A. G., and Bolker, B. M.: Tundra ecosystems observed to be CO₂ sources due to differential amplification of the carbon cycle, *Ecol. Lett.*, 16, 1307–1315, <https://doi.org/10.1111/ele.12164>, 2013.
- 1820 Berner, L. T. and Goetz, S. J.: Satellite observations document trends consistent with a boreal forest biome shift, *Glob. Chang. Biol.*, 28, 3275–3292, <https://doi.org/10.1111/gcb.16121>, 2022.
- 1825 Biskaborn, B. K., Smith, S. L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P., Romanovsky, V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R., Diekmann, B., Drozdov, D., Etzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T., Isaksen, K., Ishikawa, M., Johansson, M., Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P., Kröger, T., Lambiel, C., Lanckman, J.-P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M., Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q., Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale, *Nat. Commun.*, 10, 264, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.
- 1830 Bogard, M. J., Kuhn, C. D., Johnston, S. E., Striegl, R. G., Holtgrieve, G. W., Dornblaser, M. M., Spencer, R. G. M., Wickland, K. P., and Butman, D. E.: Negligible cycling of terrestrial carbon in many lakes of the arid circumpolar landscape, *Nat. Geosci.*, 12, 180–185, <https://doi.org/10.1038/s41561-019-0299-5>, 2019.
- 1835 Casas-Ruiz, J. P., Bodmer, P., Bona, K. A., Butman, D., Couturier, M., Emilson, E. J. S., Finlay, K., Genet, H., Hayes, D., Karlsson, J., Paré, D., Peng, C., Striegl, R., Webb, J., Wei, X., Ziegler, S. E., and Del Giorgio, P. A.: Integrating terrestrial and aquatic ecosystems to constrain estimates of land-atmosphere carbon exchange, *Nat. Commun.*, 14, 1571, <https://doi.org/10.1038/s41467-023-37232-2>, 2023.
- 1840 Chen, B., Coops, N. C., Fu, D., Margolis, H. A., Amiro, B. D., Black, T. A., Arain, M. A., Barr, A. G., Bourque, C. P.-A., Flanagan, L. B., Lafleur, P. M., McCaughey, J. H., and Wofsy, S. C.: Characterizing spatial representativeness of flux tower eddy-covariance measurements across the Canadian Carbon Program Network using remote sensing and footprint analysis, *Remote Sens. Environ.*, 124, 742–755, <https://doi.org/10.1016/j.rse.2012.06.007>, 2012.
- 1845 Chu, H., Christianson, D. S., Cheah, Y.-W., Pastorello, G., O'Brien, F., Geden, J., Ngo, S.-T., Hollowgrass, R., Leibowitz, K., Beekwilder, N. F., Sandesh, M., Dengel, S., Chan, S. W., Santos, A., Delwiche, K., Yi, K., Buechner, C., Baldocchi, D., Papale, D., Keenan, T. F., Biraud, S. C., Agarwal, D. A., and Torn, M. S.: AmeriFlux BASE data pipeline to support network growth and data sharing, *Sci. Data*, 10, 614, <https://doi.org/10.1038/s41597-023-02531-2>, 2023.
- 1850 Cole, J. J. and Caraco, N. F.: Atmospheric exchange of carbon dioxide in a low-wind oligotrophic lake measured by the addition of SF₆, *Limnol. Oceanogr.*, 43, 647–656, <https://doi.org/10.4319/lo.1998.43.4.0647>, 1998.

- 1855 Delwiche, K. B., Knox, S. H., Malhotra, A., Fluet-Chouinard, E., McNicol, G., Feron, S., Ouyang, Z., Papale, D., Trotta, C., Canfora, E., and Others: FLUXNET-CH4: A global, multi-ecosystem dataset and analysis of methane seasonality from freshwater wetlands (Appendix B and Figure 3), 2021.
- Desai, A. R., Richardson, A. D., Moffat, A. M., Kattge, J., Hollinger, D. Y., Barr, A., Falge, E., Noormets, A., Papale, D., Reichstein, M., and Stauch, V. J.: Cross-site evaluation of eddy covariance GPP and RE decomposition techniques, *Agric. For. Meteorol.*, 148, 821–838, <https://doi.org/10.1016/j.agrformet.2007.11.012>, 2008.
- 1860 Emmerton, C. A., L., S. L. V., Lehnherr, I., Graydon, J. A., Kirk, J. L., and Rondeau, K. J.: The importance of freshwater systems to the net atmospheric exchange of carbon dioxide and methane with a rapidly changing high Arctic watershed, *Biogeosciences*, 13, 5849–5863, 2016.
- ESA: Land Cover CCI Product User Guide Version 2.0, available at http://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf, last access: 16 June 2025.
- 1865 Euskirchen, E. S., Edgar, C. W., Sydonia Bret-Harte, M., Kade, A., Zimov, N., and Zimov, S.: Interannual and seasonal patterns of carbon dioxide, water, and energy fluxes from ecotonal and thermokarst-impacted ecosystems on carbon-rich permafrost soils in northeastern Siberia, *J. Geophys. Res. Biogeosci.*, 122, 2651–2668, <https://doi.org/10.1002/2017jg004070>, 2017.
- 1870 Euskirchen, E. S., Edgar, C. W., Kane, E. S., Waldrop, M. P., Neumann, R. B., Manies, K. L., Douglas, T. A., Dieleman, C., Jones, M. C., and Turetsky, M. R.: Persistent net release of carbon dioxide and methane from an Alaskan lowland boreal peatland complex, *Glob. Chang. Biol.*, 30, e17139, <https://doi.org/10.1111/gcb.17139>, 2024.
- 1875 Lai, D.Y.F., Roulet, N. T., Humphreys, E. R., Moore, T. R., and Dalva, M.: The effect of atmospheric turbulence and chamber deployment period on autochamber CO₂ and CH₄ flux measurements in an ombrotrophic peatland, *Biogeosciences*, 9, 3305–3322, <https://doi.org/10.5194/bg-9-3305-2012>, 2012.
- 1880 Frost, G. V., Bhatt, U. S., Macander, M. J., Berner, L. T., Walker, D. A., Raynolds, M. K., Magnússon, R. Í., Bartsch, A., Bjerke, J. W., Epstein, H. E., Forbes, B. C., Goetz, S. J., Hoy, E. E., Karlsen, S. R., Kumpula, T., Lantz, T. C., Lara, M. J., López-Blanco, E., Montesano, P. M., Neigh, C. S. R., Nitze, I., Orndahl, K. M., Park, T., Phoenix, G. K., Rocha, A. V., Rogers, B. M., Schaepman-Strub, G., Tømmervik, H., Verdonen, M., Veremeeva, A., Virkkala, A.-M., and Waigl, C. F.: The changing face of the Arctic: four decades of greening and implications for tundra ecosystems, *Front. Environ. Sci.*, 13, <https://doi.org/10.3389/fenvs.2025.1525574>, 2025.
- 1885 Gauci, V., Pangala, S. R., Shenkin, A., Barba, J., Bastviken, D., Figueiredo, V., Gomez, C., Enrich-Prast, A., Sayer, E., Stauffer, T., Welch, B., Elias, D., McNamara, N., Allen, M., and Malhi, Y.: Global atmospheric methane uptake by upland tree woody surfaces, *Nature*, 631, 796–800, <https://doi.org/10.1038/s41586-024-07592-w>, 2024.
- 1890 Golub, M., Koupaei-Abyazani, N., Vesala, T., Mammarella, I., Ojala, A., Bohrer, G., Weyhenmeyer, G., Blanken, P., Eugster, W., Koebisch, F., Chen, J., Czajkowski, K., Deshmukh, C., Guérin, F., Heiskanen, J., Humphreys, E., Jonsson, A., Karlsson, J., Kling, G., Lee, X., Liu, H., Lohila, A., Lundin, E., Morin, T., Podgrajsek, E., Provenzale, M., Rutgersson, A., Sachs, T., Sahlée, E., Serça, D., Shao, C., Spence, C., Strachan, I., Xiao, W., and Desai, A.: Diel, seasonal, and inter-annual variation in carbon dioxide effluxes from lakes and reservoirs,

- 1895 Environ. Res. Lett., 18, <https://doi.org/10.1088/1748-9326/acb834>, 2023.
- Gómez-Gener, L., Rocher-Ros, G., Battin, T., Cohen, M. J., Dalmagro, H. J., Dinsmore, K. J., Drake, T. W., Duvert, C., Enrich-Prast, A., Horgby, Å., Johnson, M. S., Kirk, L., Machado-Silva, F., Marzolf, N. S., McDowell, M. J., McDowell, W. H., Miettinen, H., Ojala, A. K., Peter, H., Pumpanen, J., Ran, L., Riveros-Iregui, D. A., Santos, I. R., Six, J., Stanley, E. H., Wallin, M. B., White, S. A., and Sponseller, R. A.: Global carbon dioxide efflux from rivers enhanced by high nocturnal emissions, *Nat. Geosci.*, 14, 289–294, <https://doi.org/10.1038/s41561-021-00722-3>, 2021.
- 1900 Guérin, F., Abril, G., Serça, D., Delon, C., Richard, S., Delmas, R., Tremblay, A., and Varfalvy, L.: Gas transfer velocities of CO₂ and CH₄ in a tropical reservoir and its river downstream, *J. Mar. Syst.*, 66, 161–172, <https://doi.org/10.1016/j.jmarsys.2006.03.019>, 2007.
- 1905 Hall, R. O., Jr and Ulseth, A. J.: Gas exchange in streams and rivers, *WIREs Water*, 7, <https://doi.org/10.1002/wat2.1391>, 2020.
- Heiskanen, J., Brümmer, C., Buchmann, N., Calfapietra, C., Chen, H., Gielen, B., Gkritzalis, T., Hammer, S., Hartman, S., Herbst, M., Janssens, I. A., Jordan, A., Juurola, E., Karstens, U., Kasurinen, V., Kruijt, B., Lankreijer, H., Levin, I., Linderson, M.-L., Loustau, D., Merbold, L., Myhre, C. L., Papale, D., Pavelka, M., Pilegaard, K., Ramonet, M., Rebmann, C., Rinne, J., Rivier, L., Saltikoff, E., Sanders, R., Steinbacher, M., Steinhoff, T., Watson, A., Vermeulen, A. T., Vesala, T., Vítková, G., and Kutsch, W.: The Integrated Carbon Observation System in Europe, *Bull. Am. Meteorol. Soc.*, 103, E855–E872, <https://doi.org/10.1175/bams-d-19-0364.1>, 2022.
- 1910
- 1915 Holgerson, M. A. and Raymond, P. A.: Large contribution to inland water CO₂ and CH₄ emissions from very small ponds, *Nat. Geosci.*, 9, 222–226, <https://doi.org/10.1038/ngeo2654>, 2016.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., and Others: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- 1920
- Hugelius, G., Ramage, J., Burke, E., Chatterjee, A., Smallman, T. L., Aalto, T., Bastos, A., Biasi, Canadell, J. G., Chandra, N., Chevallier, F., Ciais, P., Chang, J., Feng, L., Jones, M. W., Kleinen, T., Kuhn, M., Lauerwald, R., Liu, J., López-Blanco, E., Luijkx, I. T., Marushchak, M. E., Natali, S. M., Niwa, Y., Olefeldt, D., Palmer, P. I., Patra, P. K., Peters, W., Potter, S., Poulter, B., Rogers, B. M., Riley, W. J., Saunois, M., Schuur, E. A. G., Thompson, R. L., Treat, C., Tsuruta, A., Turetsky, M. R., Virkkala, A.-M., Voigt, C., Watts, J., Zhu, Q., and Zheng, B.: Permafrost region greenhouse gas budgets suggest a weak CO₂ sink and CH₄ and N₂O sources, but magnitudes differ between top-down and bottom-up methods, *Global Biogeochem. Cycles*, 38, <https://doi.org/10.1029/2023gb007969>, 2024.
- 1925
- 1930 Huttunen, J., Lappalainen, K., Saarijärvi, E., Vaisanen, T., and Martikainen, P.: A novel sediment gas sampler and a subsurface gas collector used for measurement of the ebullition of methane and carbon dioxide from a eutrophied lake, *Sci. Total Environ.*, 266, 153–158, [https://doi.org/10.1016/s0048-9697\(00\)00749-x](https://doi.org/10.1016/s0048-9697(00)00749-x), 2001.
- 1935
- Iwata, H., Hirata, R., Takahashi, Y., Miyabara, Y., Itoh, M., and Iizuka, K.: Partitioning eddy-covariance methane fluxes from a shallow lake into diffusive and ebullitive fluxes, *Boundary*

- Layer Meteorol., 169, 413–428, <https://doi.org/10.1007/s10546-018-0383-1>, 2018.
- 1940 Jähne, B., Münnich, K. O., Bössinger, R., Dutzi, A., Huber, W., and Libner, P.: On the parameters influencing air-water gas exchange, *J. Geophys. Res.*, 92, 1937–1949, <https://doi.org/10.1029/jc092ic02p01937>, 1987.
- 1945 Jammet, M., Crill, P., Dengel, S., and Friberg, T.: Large methane emissions from a subarctic lake during spring thaw: Mechanisms and landscape significance: LAKE METHANE EMISSIONS UPON SPRING THAW, *J. Geophys. Res. Biogeosci.*, 120, 2289–2305, <https://doi.org/10.1002/2015jg003137>, 2015.
- Jammet, M., Dengel, S., Kettner, E., Parmentier, F.-J. W., Wik, M., Crill, P., and Friberg, T.: Year-round CH₄ and CO₂ flux dynamics in two contrasting freshwater ecosystems of the subarctic, *Biogeosciences*, 14, 5189–5216, <https://doi.org/10.5194/bg-14-5189-2017>, 2017.
- 1950 Järveoja, J., Nilsson, M., Crill, P., and Peichl, M.: Bimodal diel pattern in peatland ecosystem respiration rebuts uniform temperature response, *Nat. Commun.*, 11, <https://doi.org/10.1038/s41467-020-18027-1>, 2020.
- Jentsch, K., Schulz, A., Pirk, N., Foken, T., Crewell, S., and Boike, J.: High levels of CO₂ exchange during synoptic-scale events introduce large uncertainty into the Arctic carbon budget, *Geophys. Res. Lett.*, 48, <https://doi.org/10.1029/2020gl092256>, 2021.
- 1955 Jorgenson, M. T., Douglas, T. A., Shur, Y. L., and Kanevskiy, M. Z.: Mapping the vulnerability of boreal permafrost in central Alaska in relation to thaw rate, ground ice, and thermokarst development, *J. Geophys. Res. Earth Surf.*, 130, <https://doi.org/10.1029/2024jf008030>, 2025.
- 1960 Juutinen, S., Rantakari, M., Kortelainen, P., Huttunen, J. T., Larmola, T., Alm, J., and Martikainen P.: Methane dynamics in different boreal lake types, *Biogeosciences*, 6, 209–223, 2009a.
- Juutinen, S., Rantakari, M., Kortelainen, P., Huttunen, J. T., Larmola, T., Alm, J., Silvola, J., and Martikainen, P. J.: Methane dynamics in different boreal lake types, *Biogeosciences*, 6, 209–223, <https://doi.org/10.5194/bg-6-209-2009>, 2009b.
- 1965 Juutinen, S., Alm, J., and Larmola, T.: Major implication of the littoral zone for methane release from boreal lakes, <https://doi.org/10.1029/2003GB002105>, 2003.
- Kankaala, P., Käki, T., Mäkelä, S., Ojala, A., Pajunen, H., and Arvola, L.: Methane efflux in relation to plant biomass and sediment characteristics in stands of three common emergent macrophytes in boreal mesoeutrophic lakes, *Glob. Chang. Biol.*, 11, 145–153, <https://doi.org/10.1111/j.1365-2486.2004.00888.x>, 2005.
- 1970 Karlsson, J., Giesler, R., Persson, J., and Lundin, E.: High emission of carbon dioxide and methane during ice thaw in high latitude lakes: CARBON EMISSION FROM LAKES IN SPRING, *Geophys. Res. Lett.*, 40, 1123–1127, <https://doi.org/10.1002/grl.50152>, 2013a.
- 1975 Karlsson, J., Verheijen, H. A., Seekell, D. A., Vachon, D., and Klaus, M.: Ice-melt period dominates annual carbon dioxide evasion from clear-water Arctic lakes, *Limnol. Oceanogr. Lett.*, 9, 112–118, <https://doi.org/10.1002/lol2.10369>, 2024.
- Keenan, T. F., Migliavacca, M., Papale, D., Baldocchi, D., Reichstein, M., Torn, M., and Wutzler,

- T.: Widespread inhibition of daytime ecosystem respiration, *Nat. Ecol. Evol.*, 3, 407–415, <https://doi.org/10.1038/s41559-019-0809-2>, 2019.
- 1980 Kelly, J., Kljun, N., Cai, Z., Doerr, S. H., D'Onofrio, C., Holst, T., Lehner, I., Lindroth, A., Thapa, S., Vestin, P., and Santin, C.: Wildfire impacts on the carbon budget of a managed Nordic boreal forest, *Agric. For. Meteorol.*, 351, 110016, <https://doi.org/10.1016/j.agrformet.2024.110016>, 2024.
- 1985 Kittler, F., Eugster, W., Foken, T., Heimann, M., Kolle, O., and Göckede, M.: High-quality eddy-covariance CO₂ budgets under cold climate conditions: Arctic Eddy-Covariance CO₂ Budgets, *J. Geophys. Res. Biogeosci.*, 122, 2064–2084, <https://doi.org/10.1002/2017jg003830>, 2017a.
- Kittler, F., Heimann, M., Kolle, O., Zimov, N., Zimov, S., and Göckede, M.: Long-Term Drainage Reduces CO₂ Uptake and CH₄ Emissions in a Siberian Permafrost Ecosystem, *Global Biogeochem. Cycles*, 31, 1704–1717, <https://doi.org/10.1002/2017gb005774>, 2017b.
- 1990 Klaus, M. and Vachon, D.: Challenges of predicting gas transfer velocity from wind measurements over global lakes, *Aquat. Sci.*, 82, <https://doi.org/10.1007/s00027-020-00729-9>, 2020.
- Klaus, M., Öquist, M., and Macháčová, K.: Tree stem-atmosphere greenhouse gas fluxes in a boreal riparian forest, *Sci. Total Environ.*, 954, 176243, <https://doi.org/10.1016/j.scitotenv.2024.176243>, 2024.
- 1995 Korkiakoski, M., Ojanen, P., Tuovinen, J.-P., Minkkinen, K., Nevalainen, O., Penttilä, T., Aurela, M., Laurila, T., and Lohila, A.: Partial cutting of a boreal nutrient-rich peatland forest causes radically less short-term on-site CO₂ emissions than clear-cutting, *Agric. For. Meteorol.*, 332, 109361, <https://doi.org/10.1016/j.agrformet.2023.109361>, 2023.
- 2000 Kortelainen, P., Rantakari, M., Huttunen, J. T., Mattsson, T., Alm, J., Juutinen, S., Larmola, T., Silvola, J., and Martikainen, P. J.: Sediment respiration and lake trophic state are important predictors of large CO₂ evasion from small boreal lakes: LARGE CO₂ EVASION FROM SMALL BOREAL LAKES, *Glob. Chang. Biol.*, 12, 1554–1567, <https://doi.org/10.1111/j.1365-2486.2006.01167.x>, 2006.
- 2005 Kuhn, M., Olefeldt, D., Arndt, K. A., Bastviken, D., Bruhwiler, L., Crill, P., DelSontro, T., Fluet-Chouinard, E., Grosse, G., Hovemyr, M., Hugelius, G., MacIntyre, S., Malhotra, A., McGuire, A. D., Oh, Y., Poulter, B., Treat, C. C., Turetsky, M. R., Varner, R. K., Walter Anthony, K. M., Watts, J. D., and Zhang, Z.: Current and future methane emissions from boreal-Arctic wetlands and lakes, *Nat. Clim. Chang.*, 15, 986–991, <https://doi.org/10.1038/s41558-025-02413-y>, 2025.
- 2010 Kuhn, M. A., Varner, R. K., Bastviken, D., Crill, P., MacIntyre, S., Turetsky, M., Walter Anthony, K., McGuire, A. D., and Olefeldt, D.: BAWLD-CH₄: a comprehensive dataset of methane fluxes from boreal and arctic ecosystems, *Earth System Science Data*, 13, 5151–5189, <https://doi.org/10.5194/essd-13-5151-2021>, 2021.
- 2015 Kuhn, M. A., Schmidt, M., Heffernan, L., Stührenberg, J., Knorr, K.-H., Estop-Aragonés, C., Broder, T., Gonzalez Moguel, R., Douglas, P. M. J., and Olefeldt, D.: High ebullitive, millennial-aged greenhouse gas emissions from thermokarst expansion of peatland lakes in boreal western Canada, *Limnol. Oceanogr.*, 68, 498–513, <https://doi.org/10.1002/lno.12288>, 2023.
- Kyzivat, E. D. and Smith, L. C.: A closer look at the effects of lake area, aquatic vegetation, and

double-counted wetlands on pan-Arctic lake methane emissions estimates, *Geophys. Res. Lett.*, 50, <https://doi.org/10.1029/2023gl104825>, 2023.

- 2020 Kyzivat, E. D., Smith, L. C., Garcia-Tigreros, F., Huang, C., Wang, C., Langhorst, T., Fayne, J. V., Harlan, M. E., Ishitsuka, Y., Feng, D., Dolan, W., Pitcher, L. H., Wickland, K. P., Dornblaser, M. M., Striegl, R. G., Pavelsky, T. M., Butman, D. E., and Gleason, C. J.: The importance of lake emergent aquatic vegetation for estimating arctic-boreal methane emissions, *J. Geophys. Res. Biogeosci.*, 127, <https://doi.org/10.1029/2021jg006635>, 2022.
- 2025 Lasslop, G., Reichstein, M., Papale, D., Richardson, A. D., Arneth, A., Barr, A., Stoy, P., and Wohlfahrt, G.: Separation of net ecosystem exchange into assimilation and respiration using a light response curve approach: critical issues and global evaluation: SEPARATION OF NEE INTO GPP AND RECO, *Glob. Chang. Biol.*, 16, 187–208, <https://doi.org/10.1111/j.1365-2486.2009.02041.x>, 2010.
- 2030 Liu, S., Kuhn, C., Amatulli, G., Aho, K., Butman, D. E., Allen, G. H., Lin, P., Pan, M., Yamazaki, D., Brinkerhoff, C., Gleason, C., Xia, X., and Raymond, P. A.: The importance of hydrology in routing terrestrial carbon to the atmosphere via global streams and rivers, *Proc. Natl. Acad. Sci. U. S. A.*, 119, e2106322119, <https://doi.org/10.1073/pnas.2106322119>, 2022.
- 2035 Loken, L. C., Crawford, J. T., Schramm, P. J., Stadler, P., Desai, A. R., and Stanley, E. H.: Large spatial and temporal variability of carbon dioxide and methane in a eutrophic lake, *J. Geophys. Res. Biogeosci.*, 124, 2248–2266, <https://doi.org/10.1029/2019jg005186>, 2019.
- 2040 López-Blanco, E., Lund, M., Williams, M., Tamstorf, M. P., Westergaard-Nielsen, A., Exbrayat, J.-F., Hansen, B. U., and Christensen, T. R.: Exchange of CO₂ in Arctic tundra: impacts of meteorological variations and biological disturbance, *Biogeosciences*, 14, 4467–4483, <https://doi.org/10.5194/bg-14-4467-2017>, 2017.
- Lyu, Z., Sommers, P., Schmidt, S. K., Magnani, M., Cimpoiasu, M., Kuras, O., Zhuang, Q., Oh, Y., De La Fuente, M., Cramm, M., and Bradley, J. A.: Seasonal dynamics of Arctic soils: Capturing year-round processes in measurements and soil biogeochemical models, *Earth Sci. Rev.*, 254, 104820, <https://doi.org/10.1016/j.earscirev.2024.104820>, 2024.
- 2045 Machacova, K., Warlo, H., Svobodová, K., Agyei, T., Uchytlová, T., Horáček, P., and Lang, F.: Methane emission from stems of European beech (*Fagus sylvatica*) offsets as much as half of methane oxidation in soil, *New Phytol.*, 238, 584–597, <https://doi.org/10.1111/nph.18726>, 2023.
- 2050 McGuire, A. D., Christensen, T. R., Hayes, D. J., Herault, A., Euskirchen, E., Yi, Y., Kimball, J. S., Koven, C., Lafleur, P., Miller, P. A., Oechel, W., Peylin, P., and Williams, M.: An assessment of the carbon balance of arctic tundra: comparisons among observations, process models, and atmospheric inversions, *Biogeosci. Discuss.*, 9, 4543, <https://doi.org/10.5194/bg-9-3185-2012>, 2012.
- 2055 Minobe, S., Behrens, E., Findell, K. L., Loeb, N. G., Meyssignac, B., and Sutton, R.: Global and regional drivers for exceptional climate extremes in 2023–2024: beyond the new normal, *Npj Clim. Atmos. Sci.*, 8, 138, <https://doi.org/10.1038/s41612-025-00996-z>, 2025.
- Arctic Terrestrial Carbon Cycling: <https://arctic.noaa.gov/report-card/report-card-2024/arctic-terrestrial-carbon-cycling/>, last access: 6 January 2025.
- Natali, S. M., Watts, J. D., Rogers, B. M., Potter, S., Ludwig, S. M., Selbmann, A.-K., Sullivan,

- 2060 P. F., Abbott, B. W., Arndt, K. A., Birch, L., Björkman, M. P., Bloom, A. A., Celis, G., Christensen, T. R., Christiansen, C. T., Commane, R., Cooper, E. J., Crill, P., Czimczik, C., Davydov, S., Du, J., Egan, J. E., Elberling, B., Euskirchen, E. S., Friborg, T., Genet, H., Göckede, M., Goodrich, J. P., Grogan, P., Helbig, M., Jafarov, E. E., Jastrow, J. D., Kalhori, A. A. M., Kim, Y., Kimball, J. S., Kutzbach, L., Lara, M. J., Larsen, K. S., Lee, B.-Y., Liu, Z., Lorant, M. M., Lund, M., Lupascu, M., Madani, N., Malhotra, A., Matamala, R., McFarland, J.,
- 2065 McGuire, A. D., Michelsen, A., Minions, C., Oechel, W. C., Olefeldt, D., Parmentier, F.-J. W., Pirk, N., Poulter, B., Quinton, W., Rezanezhad, F., Risk, D., Sachs, T., Schaefer, K., Schmidt, N. M., Schuur, E. A. G., Semenchuk, P. R., Shaver, G., Sonnentag, O., Starr, G., Treat, C. C., Waldrop, M. P., Wang, Y., Welker, J., Wille, C., Xu, X., Zhang, Z., Zhuang, Q., and Zona, D.: Large loss of CO₂ in winter observed across the northern permafrost region, *Nat. Clim. Chang.*, 9, 852–857, <https://doi.org/10.1038/s41558-019-0592-8>, 2019.
- 2070 Natchimuthu, S., Sundgren, I., Gålfalk, M., Klemetsson, L., Crill, P., Danielsson, Å., and Bastviken, D.: Spatio-temporal variability of lake CH₄ fluxes and its influence on annual whole lake emission estimates, *Limnol. Oceanogr.*, 61, S13–S26, <https://doi.org/10.1002/lno.10222>, 2016.
- 2075 Natchimuthu, S., Wallin, M. B., Klemetsson, L., and Bastviken, D.: Spatio-temporal patterns of stream methane and carbon dioxide emissions in a hemiboreal catchment in Southwest Sweden, *Sci. Rep.*, 7, 39729, <https://doi.org/10.1038/srep39729>, 2017.
- 2080 Nelson, J. A., Walther, S., Gans, F., Kraft, B., Weber, U., Novick, K., Buchmann, N., Migliavacca, M., Wohlfahrt, G., Šigut, L., Ibrom, A., Papale, D., Göckede, M., Duveiller, G., Knohl, A., Hörtnagl, L., Scott, R. L., Dušek, J., Zhang, W., Hamdi, Z. M., Reichstein, M., Aranda-Barranco, S., Ardö, J., Op de Beeck, M., Billesbach, D., Bowling, D., Bracho, R., Brümmer, C., Camps-Valls, G., Chen, S., Cleverly, J. R., Desai, A., Dong, G., El-Madany, T. S., Euskirchen, E. S., Feigenwinter, I., Galvagno, M., Gerosa, G. A., Gielen, B., Gode, I., Goslee, S., Gough, C. M., Heinesch, B., Ichii, K., Jackowicz-Korczynski, M. A., Klosterhalfen, A., Knox, S.,
- 2085 Kobayashi, H., Kohonen, K.-M., Korkiakoski, M., Mammarella, I., Gharun, M., Marzuoli, R., Matamala, R., Metzger, S., Montagnani, L., Nicolini, G., O'Halloran, T., Ourcival, J.-M., Peichl, M., Pendall, E., Ruiz Reverter, B., Roland, M., Sabbatini, S., Sachs, T., Schmidt, M., Schwalm, C. R., Shekhar, A., Silberstein, R., Silveira, M. L., Spano, D., Tagesson, T., Tramontana, G., Trotta, C., Turco, F., Vesala, T., Vincke, C., Vitale, D., Vivoni, E. R., Wang, Y., Woodgate, W.,
- 2090 Yepez, E. A., Zhang, J., Zona, D., and Jung, M.: X-BASE: the first terrestrial carbon and water flux products from an extended data-driven scaling framework, *FLUXCOM-X*, *Biogeosciences*, 21, 5079–5115, <https://doi.org/10.5194/bg-21-5079-2024>, 2024.
- 2095 Novick, K. A., Biederman, J. A., Desai, A. R., Litvak, M. E., Moore, D. J. P., Scott, R. L., and Torn, M. S.: The AmeriFlux network: A coalition of the willing, *Agric. For. Meteorol.*, 249, 444–456, <https://doi.org/10.1016/j.agrformet.2017.10.009>, 2018.
- Olefeldt, D., Turetsky, M. R., Crill, P. M., and McGuire, A. D.: Environmental and physical controls on northern terrestrial methane emissions across permafrost zones, *Glob. Chang. Biol.*, 19, 589–603, <https://doi.org/10.1111/gcb.12071>, 2013.
- 2100 Olefeldt, D., Hovemyr, M., Kuhn, M. K. A., Bastviken, D., Bohn, T. J., Connolly, J., Crill, P., Euskirchen, E. S., Finkelstein, S. A., Genet, H., Grosse, G., Harris, L. I., Heffernan, L., Helbig, M., Hugelius, G., Juutinen, S., Lara, M. J., Malhotra, A., Manies, K., McGuire, A. D., Natali, S. M., O'Donnell, J. A., Parmentier, F.-J. W., Räsänen, A., Schädel, C., Sonnentag, O., Strack, M., Tank, S. E., Treat, C., Varner, R. K., Virtanen, T., Warren, R. K., and Watts, J. D.: The Boreal-

- 2105 Arctic Wetland and Lake Dataset (BAWLD), Earth System Science Data, <https://doi.org/10.5194/essd-2021-140>Earth, 2021.
- O'Neill, H. B., Smith, S. L., Burn, C. R., Duchesne, C., and Zhang, Y.: Widespread permafrost degradation and thaw subsidence in northwest Canada, *J. Geophys. Res. Earth Surf.*, 128, <https://doi.org/10.1029/2023jf007262>, 2023.
- 2110 Pajala, G., Sawakuchi, H. O., Rudberg, D., Schenk, J., Sieczko, A., Gålfalk, M., Seekell, D., Sundgren, I., Thanh Duc, N., Karlsson, J., and Bastviken, D.: The effects of water column dissolved oxygen concentrations on lake methane emissions—results from a whole-lake oxygenation experiment, *J. Geophys. Res. Biogeosci.*, 128, <https://doi.org/10.1029/2022jg007185>, 2023.
- 2115 Pallandt, M. M., Kumar, J., Mauritz, M., Schuur, E. A. G., Virkkala, A.-M., Celis, G., Hoffman, F. M., and Göckede, M.: Representativeness assessment of the pan-Arctic eddy covariance site network and optimized future enhancements, *Biogeosciences*, 19, 559–583, <https://doi.org/10.5194/bg-19-559-2022>, 2022.
- 2120 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P.,
- 2125 Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De Ligne, A., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., et al.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Sci Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.
- 2130
- 2135 Pirk, N., Tamstorf, M. P., Lund, M., Mastepanov, M., Pedersen, S. H., Mylius, M. R., Parmentier, F.-J. W., Christiansen, H. H., and Christensen, T. R.: Snowpack fluxes of methane and carbon dioxide from high Arctic tundra, *J. Geophys. Res. Biogeosci.*, 121, 2886–2900, <https://doi.org/10.1002/2016jg003486>, 2016.
- Pirk, N., Aalstad, K., Mannerfelt, E. S., Clayer, F., de Wit, H., Christiansen, C. T., Althuizen, I., Lee, H., and Westermann, S.: Disaggregating the carbon exchange of degrading permafrost peatlands using Bayesian deep learning, *Geophys. Res. Lett.*, 51, <https://doi.org/10.1029/2024gl109283>, 2024.
- 2140
- Prėskienis, V., Laurion, I., Bouchard, F., Douglas, P. M. J., Billett, M. F., Fortier, D., and Xu, X.: Seasonal patterns in greenhouse gas emissions from lakes and ponds in a High Arctic polygonal landscape, *Limnol. Oceanogr.*, 66, <https://doi.org/10.1002/lno.11660>, 2021.
- 2145 Ramage, J., Kuhn, M., Virkkala, A.-M., Voigt, C., Marushchak, M. E., Bastos, A., Biasi, C., Canadell, J. G., Ciais, P., López-Blanco, E., Natali, S. M., Olefeldt, D., Potter, S., Poulter, B., Rogers, B. M., Schuur, E. A. G., Treat, C., Turetsky, M. R., Watts, J., and Hugelius, G.: The net

- GHG balance and budget of the permafrost region (2000–2020) from ecosystem flux upscaling, *Global Biogeochem. Cycles*, 38, e2023GB007953, <https://doi.org/10.1029/2023gb007953>, 2024.
- 2150 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3, 1–10, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- 2155 Raymond, P. A., Zappa, C. J., Butman, D., Bott, T. L., Potter, J., Mulholland, P., Laursen, A. E., McDowell, W. H., and Newbold, D.: Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers: Gas transfer velocity and hydraulic geometry, *Limnol. Oceanogr.*, 2, 41–53, <https://doi.org/10.1215/21573689-1597669>, 2012.
- 2160 Ray, N. E., Holgerson, M. A., Andersen, M. R., Bikše, J., Bortolotti, L. E., Futter, M., Kokorite, I., Law, A., McDonald, C., Mesman, J. P., Peacock, M., Richardson, D. C., Arsenault, J., Bansal, S., Cawley, K., Kuhn, M., Shahabnia, A. R., and Smufer, F.: Spatial and temporal variability in summertime dissolved carbon dioxide and methane in temperate ponds and shallow lakes, *Limnol. Oceanogr.*, 68, 1530–1545, <https://doi.org/10.1002/lno.12362>, 2023.
- 2165 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grunwald, T., Havrankova, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.-M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm, *Glob. Chang. Biol.*, 11, 1424–1439, <https://doi.org/10.1111/j.1365-2486.2005.001002.x>, 2005.
- 2170 Rudberg, D., Duc, N. T., Schenk, J., Sieczko, A. K., Pajala, G., Sawakuchi, H. O., Verheijen, H. A., Melack, J. M., MacIntyre, S., Karlsson, J., and Bastviken, D.: Diel variability of CO₂ emissions from northern lakes, *J. Geophys. Res. Biogeosci.*, 126, <https://doi.org/10.1029/2021jg006246>, 2021.
- 2175 Runkle, B. R., Sachs, T., Wille, C., Pfeiffer, E. M., and Kutzbach, L.: Bulk partitioning the growing season net ecosystem exchange of CO₂ in Siberian tundra reveals the seasonality of its carbon sequestration strength, *Biogeosciences*, 10, 1337–1349, 2013.
- 2180 Schmiedeskamp, M., Praetzel, L. S. E., Bastviken, D., and Knorr, K.-H.: Whole-lake methane emissions from two temperate shallow lakes with fluctuating water levels: Relevance of spatiotemporal patterns, *Limnol. Oceanogr.*, 66, 2455–2469, <https://doi.org/10.1002/lno.11764>, 2021.
- 2185 Schuur, E. A. G., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A., Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G., and Zimov, S. A.: Vulnerability of permafrost carbon to climate change: Implications for the global carbon cycle, *Bioscience*, 58, 701–714, <https://doi.org/10.1641/b580807>, 2008.
- 2190 Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.

- 2195 Schuur, E. A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven, C., Leshyk, V., Lawrence, D., Loranty, M. M., Mauritz, M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C., Strauss, J., Treat, C., and Turetsky, M.: Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic, *Annu. Rev. Environ. Resour.*, 47, 343–371, <https://doi.org/10.1146/annurev-environ-012220-011847>, 2022.
- 2200 See, C. R., Virkkala, A.-M., Natali, S. M., Rogers, B. M., Mauritz, M., Biasi, C., Bokhorst, S., Boike, J., Bret-Harte, M. S., Celis, G., Chae, N., Christensen, T. R., Murner (Connon), S. J., Dengel, S., Dolman, H., Edgar, C. W., Elberling, B., Emmerton, C. A., Euskirchen, E. S., Göckede, M., Grelle, A., Heffernan, L., Helbig, M., Holl, D., Humphreys, E., Iwata, H., Järveoja, J., Kobayashi, H., Kochendorfer, J., Kolari, P., Kotani, A., Kutzbach, L., Kwon, M. J., Lathrop, E. R., López-Blanco, E., Mammarella, I., Marushchak, M. E., Mastepanov, M., Matsuura, Y., Merbold, L., Meyer, G., Minions, C., Nilsson, M. B., Nojeim, J., Oberbauer, S. F., Olefeldt, D., Park, S.-J., Parmentier, F.-J. W., Peichl, M., Peter, D., Petrov, R., Poyatos, R., Prokushkin, A.
- 2205 S., Quinton, W., Rodenhizer, H., Sachs, T., Savage, K., Schulze, C., Sjögersten, S., Sonnentag, O., St. Louis, V. L., Torn, M. S., Tuittila, E.-S., Ueyama, M., Varlagin, A., Voigt, C., Watts, J. D., Zona, D., Zyryanov, V. I., and Schuur, E. A. G.: Decadal increases in carbon uptake offset by respiratory losses across northern permafrost ecosystems, *Nat. Clim. Chang.*, 14, 853–862, <https://doi.org/10.1038/s41558-024-02057-4>, 2024.
- 2210 Sieczko, A. K., Duc, N. T., Schenk, J., Pajala, G., Rudberg, D., Sawakuchi, H. O., and Bastviken, D.: Diel variability of methane emissions from lakes, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 21488–21494, <https://doi.org/10.1073/pnas.2006024117>, 2020.
- 2215 Song, C., Liu, S., Wang, G., Zhang, L., Rosentreter, J. A., Zhao, G., Sun, X., Yao, Y., Mu, C., Sun, S., Hu, Z., Lin, S., Sun, J., Li, Y., Wang, Y., Li, Y., Raymond, P. A., and Karlsson, J.: Inland water greenhouse gas emissions offset the terrestrial carbon sink in the northern cryosphere, *Sci. Adv.*, 10, eadp0024, <https://doi.org/10.1126/sciadv.adp0024>, 2024.
- 2220 Staehr, P. A., Bade, D., Van de Bogert, M. C., Koch, G. R., Williamson, C., Hanson, P., Cole, J. J., and Kratz, T.: Lake metabolism and the diel oxygen technique: State of the science: Guideline for lake metabolism studies, *Limnol. Oceanogr. Methods*, 8, 628–644, <https://doi.org/10.4319/lom.2010.8.0628>, 2010.
- Stanley, E. H., Loken, L. C., Casson, N. J., Oliver, S. K., Sponseller, R. A., Wallin, M. B., Zhang, L., and Rocher-Ros, G.: GRiMeDB: the Global River Methane Database of concentrations and fluxes, *Earth Syst. Sci. Data*, 15, 2879–2926, <https://doi.org/10.5194/essd-15-2879-2023>, 2023.
- 2225 Sundqvist, E., Crill, P., Mölder, M., Vestin, P., and Lindroth, A.: Atmospheric methane removal by boreal plants: METHANE REMOVAL BY BOREAL PLANTS, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012gl053592>, 2012.
- Tank, S. E., Lesack, L. F. W., and Hesslein, R. H.: Northern delta lakes as summertime CO₂ absorbers within the arctic landscape, *Ecosystems*, 12, 144–157, <https://doi.org/10.1007/s10021-008-9213-5>, 2009.
- 2230 Tao, J., Zhu, Q., Riley, W. J., and Neumann, R. B.: Warm-season net CO₂ uptake outweighs cold-season emissions over Alaskan North Slope tundra under current and RCP8.5 climate, *Environ. Res. Lett.*, 16, 055012, <https://doi.org/10.1088/1748-9326/abf6f5>, 2021.
- Thornton, B. F., Wik, M., and Crill, P. M.: Double-counting challenges the accuracy of high-

- 2235 latitude methane inventories: DOUBLE-COUNTING ARCTIC METHANE, *Geophys. Res. Lett.*, 43, 12,569–12,577, <https://doi.org/10.1002/2016gl071772>, 2016.
- Treat, C. C., Anthony Bloom, A., and Marushchak, M. E.: Nongrowing season methane emissions—a significant component of annual emissions across northern ecosystems, <https://doi.org/10.1111/gcb.14137>, 2018.
- 2240 Treat, C. C., Virkkala, A.-M., Burke, E., Bruhwiler, L., Chatterjee, A., Fisher, J. B., Hashemi, J., Parmentier, F.-J. W., Rogers, B. M., Westermann, S., Watts, J. D., Blanc-Betes, E., Fuchs, M., Kruse, S., Malhotra, A., Miner, K., Strauss, J., Armstrong, A., Epstein, H. E., Gay, B., Goeckede, M., Kalhori, A., Kou, D., Miller, C. E., Natali, S. M., Oh, Y., Shakil, S., Sonnentag, O., Varner, R. K., Zolkos, S., Schuur, E. A. G., and Hugelius, G.: Permafrost carbon: Progress on understanding stocks and fluxes across northern terrestrial ecosystems, *J. Geophys. Res. Biogeosci.*, 129, <https://doi.org/10.1029/2023jg007638>, 2024.
- 2245 Turetsky, M. R., Abbott, B. W., Jones, M. C., Anthony, K. W., Olefeldt, D., Schuur, E. A. G., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D. M., Gibson, C., Sannel, A. B. K., and David McGuire, A.: Carbon release through abrupt permafrost thaw, <https://doi.org/10.1038/s41561-019-0526-0>, 2020.
- 2250 Ueyama, M., Knox, S. H., Delwiche, K. B., Bansal, S., Riley, W. J., Baldocchi, D., Hirano, T., McNicol, G., Schafer, K., Windham-Myers, L., Poulter, B., Jackson, R. B., Chang, K.-Y., Chen, J., Chu, H., Desai, A. R., Gogo, S., Iwata, H., Kang, M., Mammarella, I., Peichl, M., Sonnentag, O., Tuittila, E.-S., Ryu, Y., Euskirchen, E. S., Göckede, M., Jacotot, A., Nilsson, M. B., and Sachs, T.: Modeled production, oxidation, and transport processes of wetland methane emissions in temperate, boreal, and Arctic regions, *Glob. Chang. Biol.*, 29, 2313–2334, <https://doi.org/10.1111/gcb.16594>, 2023.
- 2255 Ueyama, M., Takao, Y., Yazawa, H., Tanaka, M., Yabuki, H., Kumagai, T. O., and Ichii: The JapanFlux2024 dataset for eddy covariance observations covering Japan and East Asia from 1990 to 2023, *Earth System Science Data Discussions*, 2025, 1–47, 2025.
- 2260 Vachon, D. and Prairie, Y. T.: The ecosystem size and shape dependence of gas transfer velocity versus wind speed relationships in lakes, *Can. J. Fish. Aquat. Sci.*, 70, 1757–1764, <https://doi.org/10.1139/cjfas-2013-0241>, 2013.
- 2265 Vainio, E., Haikarainen, I. P., Machacova, K., Putkinen, A., Santalahti, M., Koskinen, M., Fritze, H., Tuomivirta, T., and Pihlatie, M.: Soil-tree-atmosphere CH₄ flux dynamics of boreal birch and spruce trees during spring leaf-out, *Plant Soil*, 478, 391–407, <https://doi.org/10.1007/s11104-022-05447-9>, 2022.
- 2270 Valentini, R.: EUROFLUX: An Integrated Network for Studying the Long-Term Responses of Biospheric Exchanges of Carbon, Water, and Energy of European Forests, in: *Fluxes of Carbon, Water and Energy of European Forests*, edited by: Valentini, R., Springer Berlin Heidelberg, Berlin, Heidelberg, 1–8, https://doi.org/10.1007/978-3-662-05171-9_1, 2003.
- 2275 Virkkala, A.-M., Aalto, J., Rogers, B. M., Tagesson, T., Treat, C. C., Natali, S. M., Watts, J. D., Potter, S., Lehtonen, A., Mauritz, M., Schuur, E. A. G., Kochendorfer, J., Zona, D., Oechel, W., Kobayashi, H., Humphreys, E., Goeckede, M., Iwata, H., Lafleur, P. M., Euskirchen, E. S., Bokhorst, S., Marushchak, M., Martikainen, P. J., Elberling, B., Voigt, C., Biasi, C., Sonnentag, O., Parmentier, F.-J. W., Ueyama, M., Celis, G., St Louis, V. L., Emmerton, C. A., Peichl, M., Chi, J., Järveoja, J., Nilsson, M. B., Oberbauer, S. F., Torn, M. S., Park, S.-J., Dolman, H.,

- Mammarella, I., Chae, N., Poyatos, R., López-Blanco, E., Christensen, T. R., Kwon, M. J., Sachs, T., Holl, D., and Luoto, M.: Statistical upscaling of ecosystem CO₂ fluxes across the terrestrial tundra and boreal domain: Regional patterns and uncertainties, *Glob. Chang. Biol.*, 27, 4040–4059, <https://doi.org/10.1111/gcb.15659>, 2021a.
- 2280 Virkkala, A. M., Natali, S. M., Rogers, B. M., Watts, J. D., Savage, K., Connon, S. J., and Zyryanov V: The ABCflux database: Arctic-Boreal CO₂ flux observations and ancillary information aggregated to monthly time steps across terrestrial ecosystems, *Earth System Science Data Discussions*, 2021, 1–54, 2021b.
- 2285 Virkkala, A.-M., Natali, S. M., Rogers, B. M., Watts, J. D., Savage, K., Connon, S. J., Mauritz, M., Schuur, E. A. G., Peter, D., Minions, C., Nojeim, J., Commane, R., Emmerton, C. A., Goeckede, M., Helbig, M., Holl, D., Iwata, H., Kobayashi, H., Kolari, P., López-Blanco, E., Marushchak, M. E., Mastepanov, M., Merbold, L., Parmentier, F.-J. W., Peichl, M., Sachs, T., Sonnentag, O., Ueyama, M., Voigt, C., Aurela, M., Boike, J., Celis, G., Chae, N., Christensen, T. R., Bret-Harte, M. S., Dengel, S., Dolman, H., Edgar, C. W., Elberling, B., Euskirchen, E., Grelle, A., Hatakka, J., Humphreys, E., Järveoja, J., Kotani, A., Kutzbach, L., Laurila, T., Lohila, A., Mammarella, I., Matsuura, Y., Meyer, G., Nilsson, M. B., Oberbauer, S. F., Park, S.-J., Petrov, R., Prokushkin, A. S., Schulze, C., St. Louis, V. L., Tuittila, E.-S., Tuovinen, J.-P., Quinton, W., Varlagin, A., Zona, D., and Zyryanov, V. I.: The ABCflux database: Arctic–boreal CO₂ flux observations and ancillary information aggregated to monthly time steps across terrestrial ecosystems, *Earth Syst. Sci. Data*, 14, 179–208, <https://doi.org/10.5194/essd-14-179-2022>, 2022.
- 2290 Virkkala, A.-M., Niittynen, P., Kemppinen, J., Marushchak, M. E., Voigt, C., Hensgens, G., Kerttula, J., Happonen, K., Tyystjärvi, V., Biasi, C., Hultman, J., Rinne, J., and Luoto, M.: High-resolution spatial patterns and drivers of terrestrial ecosystem carbon dioxide, methane, and nitrous oxide fluxes in the tundra, *Biogeosciences*, 21, 335–355, <https://doi.org/10.5194/bg-21-335-2024>, 2024.
- 2300 Virkkala, A.-M., Rogers, B. M., Watts, J. D., Arndt, K. A., Potter, S., Wargowsky, I., Schuur, E. A. G., See, C. R., Mauritz, M., Boike, J., Bret-Harte, M. S., Burke, E. J., Burrell, A., Chae, N., Chatterjee, A., Chevallier, F., Christensen, T. R., Commane, R., Dolman, H., Edgar, C. W., Elberling, B., Emmerton, C. A., Euskirchen, E. S., Feng, L., Goeckede, M., Grelle, A., Helbig, M., Holl, D., Järveoja, J., Karsanaev, S. V., Kobayashi, H., Kutzbach, L., Liu, J., Luijckx, I. T., López-Blanco, E., Lunneberg, K., Mammarella, I., Marushchak, M. E., Mastepanov, M., Matsuura, Y., Maximov, T. C., Merbold, L., Meyer, G., Nilsson, M. B., Niwa, Y., Oechel, W., Palmer, P. I., Park, S.-J., Parmentier, F.-J. W., Peichl, M., Peters, W., Petrov, R., Quinton, W., Rödenbeck, C., Sachs, T., Schulze, C., Sonnentag, O., St. Louis, V. L., Tuittila, E.-S., Ueyama, M., Varlagin, A., Zona, D., and Natali, S. M.: Wildfires offset the increasing but spatially heterogeneous Arctic–boreal CO₂ uptake, *Nat. Clim. Chang.*, 1–8, <https://doi.org/10.1038/s41558-024-02234-5>, 2025.
- 2305 Vogt, J., Pallandt, M. M. T. A., Basso, L. S., Bolek, A., Ivanova, K., Schlutow, M., Celis, G., Kuhn, M., Mauritz, M., Schuur, E. A. G., Arndt, K., Virkkala, A.-M., Wargowsky, I., and Goeckede, M.: ARGO: ARctic greenhouse Gas Observation metadata version 1, *Earth Syst. Sci. Data*, 17, 2553–2573, <https://doi.org/10.5194/essd-17-2553-2025>, 2025.
- 2310 Voigt, C., Marushchak, M. E., Lamprecht, R. E., Jackowicz-Korczyński, M., Lindgren, A., Mastepanov, M., Granlund, L., Christensen, T. R., Tahvanainen, T., Martikainen, P. J., and Biasi, C.: Increased nitrous oxide emissions from Arctic peatlands after permafrost thaw, *Proc.*
- 2320

- Natl. Acad. Sci. U.S.A., 114, 6238–6243, <https://doi.org/10.1073/pnas.1702902114>, 2017.
- 2325 Voigt, C., Virkkala, A.-M., Hould Gosselin, G., Bennett, K. A., Black, T. A., Detto, M., Chevrier-Dion, C., Guggenberger, G., Hashmi, W., Kohl, L., Kou, D., Marquis, C., Marsh, P., Marushchak, M. E., Nesic, Z., Nykänen, H., Saarela, T., Sauheitl, L., Walker, B., Weiss, N., Wilcox, E. J., and Sonnentag, O.: Arctic soil methane sink increases with drier conditions and higher ecosystem respiration, *Nat. Clim. Chang.*, 1–10, <https://doi.org/10.1038/s41558-023-01785-3>, 2023.
- 2330 Vonk, J. E., Fritz, M., Speetjens, N. J., Babin, M., Bartsch, A., Basso, L. S., Bröder, L., Göckede, M., Gustafsson, Ö., Hugelius, G., Irrgang, A. M., Juhls, B., Kuhn, M. A., Lantuit, H., Manizza, M., Martens, J., O'Regan, M., Suslova, A., Tank, S. E., Terhaar, J., and Zolkos, S.: The land–ocean Arctic carbon cycle, *Nat. Rev. Earth Environ.*, 6, 86–105, <https://doi.org/10.1038/s43017-024-00627-w>, 2025.
- 2335 Walter Anthony, K. M. and Anthony, P.: Constraining spatial variability of methane ebullition seeps in thermokarst lakes using point process models: THERMOKARST LAKE METHANE EBULLITION, *J. Geophys. Res. Biogeosci.*, 118, 1015–1034, <https://doi.org/10.1002/jgrg.20087>, 2013.
- 2340 Walter Anthony, K. M., Anthony, P., Hasson, N., Edgar, C., Sivan, O., Eliani-Russak, E., Bergman, O., Minsley, B. J., James, S. R., Pastick, N. J., Kholodov, A., Zimov, S., Euskirchen, E., Bret-Harte, M. S., Grosse, G., Langer, M., and Nitzbon, J.: Upland Yedoma taliks are an unpredicted source of atmospheric methane, *Nat. Commun.*, 15, 6056, <https://doi.org/10.1038/s41467-024-50346-5>, 2024.
- 2345 Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre: Warm Winter 2020 ecosystem eddy covariance flux product for 73 stations in FLUXNET-Archive format—release 2022-1, <https://doi.org/10.18160/2G60-ZHAK>, 2020.
- 2350 Watts, J. D., Natali, S. M., Minions, C., Risk, D., Arndt, K., Zona, D., Euskirchen, E. S., Rocha, A. V., Sonnentag, O., Helbig, M., Kalhori, A., Oechel, W., Ikawa, H., Ueyama, M., Suzuki, R., Kobayashi, H., Celis, G., Schuur, E. A. G., Humphreys, E., Kim, Y., Lee, B.-Y., Goetz, S., Madani, N., Schiferl, L. D., Commane, R., Kimball, J. S., Liu, Z., Torn, M. S., Potter, S., Wang, J. A., Jorgenson, M. T., Xiao, J., Li, X., and Edgar, C.: Soil respiration strongly offsets carbon uptake in Alaska and Northwest Canada, *Environ. Res. Lett.*, 16, 084051, <https://doi.org/10.1088/1748-9326/ac1222>, 2021.
- 2355 Webb, E. E., Liljedahl, A. K., Cordeiro, J. A., Loranty, M. M., Witharana, C., and Lichstein, J. W.: Permafrost thaw drives surface water decline across lake-rich regions of the Arctic, *Nat. Clim. Chang.*, 12, 841–846, <https://doi.org/10.1038/s41558-022-01455-w>, 2022.
- 2360 Webb, H., Fuchs, M., Abbott, B. W., Douglas, T. A., Elder, C. D., Ernakovich, J. G., Euskirchen, E. S., Göckede, M., Grosse, G., Hugelius, G., Jones, M. C., Koven, C., Kropp, H., Lathrop, E., Li, W., Loranty, M. M., Natali, S. M., Olefeldt, D., Schädel, C., Schuur, E. A. G., Sonnentag, O., Strauss, J., Virkkala, A.-M., and Turetsky, M. R.: A review of abrupt permafrost thaw: Definitions, usage, and a proposed conceptual framework, *Curr. Clim. Change Rep.*, 11, 7, <https://doi.org/10.1007/s40641-025-00204-3>, 2025.
- Welles, J. M., Demetriades-Shah, T. H., and McDermitt, D. K.: Considerations for measuring ground CO₂ effluxes with chambers, *Chem. Geol.*, 177, 3–13, [https://doi.org/10.1016/s0009-2541\(00\)00388-0](https://doi.org/10.1016/s0009-2541(00)00388-0), 2001.

- 2365 Wik, M., Crill, P. M., Varner, R. K., and Bastviken, D.: Multiyear measurements of ebullitive methane flux from three subarctic lakes: METHANE EBULLITION FROM SUBARCTIC LAKES, *J. Geophys. Res. Biogeosci.*, 118, 1307–1321, <https://doi.org/10.1002/jgrg.20103>, 2013.
- Wik, M., Thornton, B. F., Bastviken, D., Uhlbäck, J., and Crill, P. M.: Biased sampling of methane release from northern lakes: A problem for extrapolation, *Geophys. Res. Lett.*, 43, 1256–1262, <https://doi.org/10.1002/2015gl066501>, 2016a.
- 2370 Wik, M., Varner, R. K., Anthony, K. W., MacIntyre, S., and Bastviken, D.: Climate-sensitive northern lakes and ponds are critical components of methane release, *Nat. Geosci.*, 9, 99–105, <https://doi.org/10.1038/ngeo2578>, 2016b.
- Yuan, K., Li, F., McNicol, G., Chen, M., Hoyt, A., Knox, S., Riley, W. J., Jackson, R., and Zhu, Q.: Boreal-Arctic wetland methane emissions modulated by warming and vegetation activity, *Nat. Clim. Chang.*, 14, 282–288, <https://doi.org/10.1038/s41558-024-01933-3>, 2024.
- 2375 Zolkos, S., Tank, S. E., Kokelj, S. V., Striegl, R. G., Shakil, S., Voigt, C., Sonnentag, O., Quinton, W. L., Schuur, E. A. G., Zona, D., Lafleur, P. M., Sullivan, R. C., Ueyama, M., Billesbach, D., Cook, D., Humphreys, E. R., and Marsh, P.: Permafrost landscape history shapes fluvial chemistry, ecosystem carbon balance, and potential trajectories of future change, *Global Biogeochem. Cycles*, 36, <https://doi.org/10.1029/2022gb007403>, 2022.
- 2380