

# **CoralCache: a virtual coral core repository for transparent and reproducible annual growth rate analyses**

Thomas M. DeCarlo<sup>1\*</sup>, Oliwia Jasnos<sup>1</sup>, Avi Strange<sup>1</sup>, Andreas Andersson<sup>2</sup>, Angel T. Bautista VII<sup>3</sup>, Sierra  
5 Bloomer<sup>4</sup>, Isaiah W. Bolden<sup>5</sup>, Maartje Bosman<sup>6</sup>, Thomas C. Brachert<sup>7</sup>, Giulia B. Braz<sup>8</sup>, Gabriel O.  
Cardoso<sup>9</sup>, Juan P. Carricart-Ganivet<sup>10</sup>, Jessica E. Carilli<sup>11,†</sup>, Karl D. Castillo<sup>12</sup>, Leticia Cavole<sup>13</sup>, Sylvia  
Chan<sup>5</sup>, Xuefei Chen<sup>14</sup>, Ben Chomitz<sup>15,16</sup>, Thierry Correge<sup>17</sup>, Travis A. Courtney<sup>2,18</sup>, Mikayla Deigan<sup>19</sup>,  
Juan Pablo D’Olivo<sup>20</sup>, Robert Dunbar<sup>21</sup>, Ian C. Enochs<sup>15</sup>, Ludmilla Falsarella<sup>22</sup>, Thomas Felis<sup>23</sup>, Gabriela  
Gutierrez-Estrada<sup>10</sup>, Brighton Hedger<sup>24</sup>, Shijian Hu<sup>25</sup>, Seamus Jameson<sup>26</sup>, Stacy Jupiter<sup>27</sup>, Paul Kench<sup>28</sup>,  
10 Diego K. Kersting<sup>29</sup>, Ke Lin<sup>30</sup>, Yi-Wei Liu<sup>31</sup>, Carla A. B. Lorigados<sup>32</sup>, Derek P. Manzello<sup>33</sup>, Malcolm T.  
McCulloch<sup>34</sup>, Miguel Mies<sup>8</sup>, Rodrigo L. Moura<sup>35</sup>, Ferdinand Oberle<sup>26</sup>, Natan Pereira<sup>36</sup>, Nancy Prouty<sup>26</sup>,  
Riovie D. Ramos<sup>30</sup>, Haojia Ren<sup>37</sup>, Emma Ryan<sup>6</sup>, Diane M. Thompson<sup>19</sup>, Lauren T. Toth<sup>4</sup>, Marina J.  
Vergotti<sup>9</sup>, Jody M. Webster<sup>38</sup>, Jens Zinke<sup>39</sup>

<sup>1</sup>Department of Earth and Environmental Sciences, Tulane University, New Orleans, LA, USA

15 <sup>2</sup>Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

<sup>3</sup>Department of Science and Technology – Philippine Nuclear Research Institute (DOST-PNRI), Quezon City, NCR 1101,  
Philippines

<sup>4</sup>U.S. Geological Survey, St. Petersburg Coastal and Marine Science Center, St. Petersburg, FL, USA

<sup>5</sup>School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

20 <sup>6</sup>School of Environment, University of Auckland

<sup>7</sup>Institute for Earth System Science and Remote Sensing, Leipzig University, 04103 Leipzig, Germany

<sup>8</sup>Instituto Oceanográfico, Universidade de São Paulo, Praça do Oceanográfico, 191 – 05508-120 São Paulo, SP, Brazil

<sup>9</sup>Institute of Geological Sciences, Freie Universität Berlin, Berlin, Germany

- <sup>10</sup>Laboratorio de Esclerocronología de Corales Arrecifales, Unidad Académica de Sistemas Arrecifales, ICML, UNAM. Puerto  
25 Morelos, México
- <sup>11</sup>Australian Nuclear Science and Technology Organization, Sydney, Australia.
- <sup>12</sup>University of North Carolina at Chapel Hill, Department of Earth, Marine and Environmental Sciences, Chapel Hill, NC,  
USA
- <sup>13</sup>School of Arts, Sciences and Humanities, University of São Paulo, São Paulo, Brazil
- 30 <sup>14</sup>State Key Laboratory of Deep Earth Processes and Resources, Guangzhou Institute of Geochemistry, Chinese Academy of  
Sciences, Guangzhou 510640, China
- <sup>15</sup>Atlantic Oceanographic and Meteorological Laboratory, National Oceanic and Atmospheric Administration, Miami, FL,  
USA
- <sup>16</sup>Cooperative Institute for Marine and Atmospheric Studies, University of Miami, Miami, Florida, USA
- 35 <sup>17</sup>Université de Bordeaux (UMR EPOC – OASU CNRS 5805) Allée Geoffroy Saint-Hilaire, CS 50023 33615 Pessac, France
- <sup>18</sup>Department of Marine Sciences, University of Puerto Rico Mayagüez, Mayagüez, Puerto Rico
- <sup>19</sup>Department of Geosciences, University of Arizona, Tucson, AZ 85721
- <sup>20</sup>Unidad Académica de Sistemas Arrecifales, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma  
de México, Puerto Morelos, Mexico
- 40 <sup>21</sup>Oceans Department, Stanford University, Stanford CA 94305
- <sup>22</sup>Centro de Biologia Marinha, Universidade de São Paulo, São Sebastião, Brazil
- <sup>23</sup>MARUM – Center for Marine Environmental Sciences, University of Bremen, 28359 Bremen, Germany
- <sup>24</sup>NOAA Pacific Islands Fisheries Science Center, Honolulu, HI, USA
- <sup>25</sup>College of Oceanography, Hohai University, Nanjing, JS, China
- 45 <sup>26</sup>U.S. Geological Survey, Pacific Coastal and Marine Science Center, Santa Cruz, CA, USA
- <sup>27</sup>Global Marine Program, Wildlife Conservation Society, Bronx, NY, 10460 USA
- <sup>28</sup>Department of Geography, National University of Singapore
- <sup>29</sup>Instituto de Acuicultura Torre de la Sal (IATS, CSIC), 12595 Ribera de Cabanes, Spain.

<sup>30</sup>Earth Observatory of Singapore, Nanyang Technological University, 639798 Singapore

50 <sup>31</sup>Institute of Earth Sciences, Academia Sinica, Taipei, Taiwan

<sup>32</sup>Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo. Av. Prof. Orlando Marques de Paiva, 87 – 05508-270 São Paulo, SP, Brazil.

<sup>33</sup>Coral Reef Watch, US National Oceanic and Atmospheric Administration (NOAA), College Park, MD, USA

<sup>34</sup>The University of Western Australia, Oceans Institute, Western Australia 6009, Australia

55 <sup>35</sup>Universidade Federal do Rio de Janeiro, Brazil

<sup>36</sup>PGQA, Department of Exact and Earth Science, State University of Bahia, Salvador, Brazil

<sup>37</sup>Department of Geosciences, National Taiwan University, Taiwan

<sup>38</sup>Geocoastal Research Group, School of Geosciences, The University of Sydney, NSW 2006, Australia

<sup>39</sup>School of Geography, Geology and the Environment, University of Leicester, Leicester, United Kingdom

60 †Now at Naval Information Warfare Center Pacific, San Diego, CA, USA

\* *Correspondence to:* Thomas M. DeCarlo ([tdecarlo@tulane.edu](mailto:tdecarlo@tulane.edu))

**Abstract.** As science fields enter the Big Data revolution, open-access repositories are essential for addressing larger-scale questions than are possible for single researchers by making data findable, accessible, interoperable, and reusable (FAIR). Furthermore, transparent data and code are increasingly important for reproducible research, especially for data types that

65 inherently require subjective human interpretations. These ideas are applicable to coral sclerochronology, as the field has long been characterized by individual researchers collecting and analyzing coral skeletal cores from their study sites without substantial data sharing or archiving of the core images to meet FAIR principles. Here, we present CoralCache, a virtual coral core repository that not only archives image datasets (*i.e.*, digitized X-rays and computed tomography scans), but also observer interpretations of the density banding patterns. CoralCache is linked to a graphical user interface, CoralCT, which together

70 offer a way forward for coral growth rate analysis that is reproducible and collaborative. The data organization systems presented here could also be readily applied to related archives such as tree rings or bivalve shells.

## 1 Introduction

75 The density bands within coral skeletal cores are the “tree rings of the sea”. As massive- and columnar-morphology coral colonies build their skeletons over decades to centuries, they typically construct alternating bands of relatively high and low density on an annual basis (Knutson et al., 1972; Ma, 1934). Along with providing age-model constraints to researchers analyzing skeletal geochemistry to reconstruct environmental variations, these bands also reveal the annual growth history of the colony (Hudson, 1981; Ma, 1934). The growth rates themselves can be used as environmental proxies and indicators of  
80 the relative health of the colony in response to long-term trends of acidification and warming or to acute stressors such as pollution and bleaching episodes (Barkley et al., 2018; Barkley and Cohen, 2016; Baumann et al., 2019; Cantin et al., 2010; Cantin and Lough, 2014; Carilli et al., 2009; Carricart-Ganivet et al., 2012; Castillo et al., 2012; Cooper et al., 2008, 2012; Courtney et al., 2020; De’ath et al., 2009; DeCarlo et al., 2024; Felis et al., 2009; Lough and Barnes, 2000; Mallela et al., 2015; Manzello, 2010; Mollica et al., 2019; Rodgers et al., 2021; Saenger et al., 2009; Samperiz et al., 2025; Vergotti et al.,  
85 2025).

Despite the value of coral skeletal core archives, centralized or systematic data sharing and archiving practices are lacking. Whereas other core archives, such as deep-sea sediment or fossil reef cores, have large, centralized facilities for physically storing the cores (*e.g.*, International Ocean Discovery Program (IODP) Core Repositories), no such repository exists for coral cores. Rather, coral cores are physically stored by individual researchers, university archives, or in some cases, by  
90 government science agencies (Dassié et al., 2017; Oberle et al., 2026; Williams et al., 2017). Yet, the daunting effort and costs necessary to centralize physical core storage are not necessary for coral growth studies because X-ray or computed tomography (CT) images from coral cores (hereafter “core image data”) effectively capture the annual density banding patterns that can be interpreted for past growth rates. This makes digital data archives an effective, low-cost alternative to physical archives of cores for studies of coral growth variability. Core image data are also often used to screen cores for clear growth axes and  
95 early diagenesis prior to geochemical sampling (DeLong et al., 2013; Fouke et al., 2021). Further, these images provide critical information for age modelling, such as age-depth tie points, which contribute to age model refinement. Often these raw images and age-depth models are not publicly available, limiting meta-analyses among sites (Thompson, 2022). Many coral geochemical and growth rate datasets are archived at the National Centers for Environmental Information (NCEI) World Data


Service for Paleoclimatology (<https://www.ncei.noaa.gov/products/paleoclimatology>) and the World Data Center PANGAEA  
100 (<https://www.pangaea.de/>) (Felden et al., 2023). The paleoclimate community has also recently produced the PAGES (Past  
Global Changes) CoralHydro2k database hosted at NCEI, a global compilation of coral  $\delta^{18}\text{O}$  and Sr/Ca proxy records of  
tropical ocean hydrology and temperature for the Common Era (Atwood et al., 2026; Walter et al., 2023). However, without a  
comparable image archive, re-analysis and interpretation of these valuable datasets is limited to that obtainable from the  
original published accounts.

105 We present CoralCache, a virtual coral core repository with global coverage for archiving coral skeletal core image  
datasets. This repository fills a key gap by enabling researchers to transparently archive and share the raw core density image  
datasets and the visual analyses of those images, rather than simply the summarized growth rate data, which on their own  
cannot be traced or precisely reproduced. CoralCache is designed to meet the data-sharing principles of findability,  
accessibility, interoperability, and reusability (FAIR) (Wilkinson et al., 2016). Researchers can search the entire core list for  
110 suitable cores to address a particular question or that are from a certain region prior to expending time and resources to collect  
new samples. We describe here the standards and mechanisms by which CoralCache achieves these steps forward in archiving  
and transparency.

## 2 The CoralCache repository

115 The foundation of the CoralCache repository is storing CT scan and X-ray image datasets of coral cores, combined not only  
with output growth-rate data, but also with the observer interpretations of the banding patterns in each core. This is achieved  
through the repository being linked to a graphical user interface, called CoralCT (Fig. 1; DeCarlo et al., 2025b), that allows  
users to trace the annual bands in the images (the reader is directed to reference (DeCarlo et al., 2025b) for additional example  
images of coral skeletal cores and how users interact with them in the CoralCT application). CoralCT pulls image datasets  
120 from the CoralCache repository and automatically sends observer band traces back to the repository. These features are built  
into the CoralCT application; therefore, no extra steps are required for observers to share their annual density-banding  
interpretations. While there are options to limit public access to pre-publication datasets, the concept for the CoralCache  
repository is that, upon publication, authors will be able to readily share the raw core images, the output growth rate data, and

the exact interpretations they made in generating those growth rate data. This will allow other researchers to review, reproduce,  
125 or revisit growth rate analyses of prior works, and even to compile growth rate datasets of potentially hundreds or thousands  
of cores from across the world. The output data include annual skeletal bulk density, linear extension, and calcification rates,  
all with assigned calendar years based on the observer's interpretation of the date of the topmost band relative to the core  
collection date. Important features such as "stress bands" (Barkley and Cohen, 2016; Cantin and Lough, 2014; DeCarlo, 2020;  
DeCarlo et al., 2020; DeCarlo and Cohen, 2017; Mallela et al., 2015) can be annotated on publicly-visible snapshots and  
130 described in a notes section of the output files. The methods used to describe these parameters are published in previous works  
(DeCarlo et al., 2015; Duprey et al., 2012; Helmle et al., 2002; Lough, 2008), including the measurement procedures in the  
CoralCT application (Fig. 1; DeCarlo et al., 2025b). The present work focuses on archiving and sharing these datasets via  
CoralCache, which exists as an online data storage system accessible via secure file transfer protocol (SFTP).



135 **Figure 1. Default display of a partially-analyzed coral skeletal core in the CoralCT application. Sliders found on the right side of the screen are used to adjust the cores appearance to increase visual clarity of annual density bands. In the core image, clear alternations of high-density (bright) and low-density (dark) bands are visible, with a pair of high- and low-density bands together representing one year of growth. The yellow lines and digits indicate user-defined bands. On this core, the observer has identified bands as the darkest part of low-density bands. In the axial image toward the lower-right, the current slice location is displayed by the red box, and yellow dots indicate the distribution of tie-points for the present band (band 17 in this case).**

140 **Additional descriptions of the methods of the CoralCT application can be found in DeCarlo et al. (2025b).**

## 2.1 Metadata and data standards

145 Several different file types are associated with the CoralCache repository. Core image data can be either 3-dimensional (3-D) CT scans or 2-dimensional (2-D) X-ray images. CT scans must be in Digital Imaging and Communications in Medicine (DICOM) format (.dcm file extension), and the files must contain embedded within them at least the following metadata elements that are standard to .dcm files: PixelSpacing (in millimetres [mm]), either SliceLocation or ImagePositionPatient (in mm), and both RescaleSlope and RescaleIntercept (*i.e.*, to convert image data to the standard Hounsfield Unit/HU scale; 150 Hounsfield, 1973). The CT scans must be in a stack of images in which each image file is an axial slice (also known as a horizontal or transverse slice). If necessary, the CoralCT application has tools that allows users to (1) convert CT scan data from .tiff to .dcm, (2) rotate .dcm stacks if the individual files are not already in the axial plane, and (3) resize .dcm files if the file size of the dataset is larger than desired (*e.g.*, loading and response times in the application increase with file size). X-ray images must be in .tiff format, and they must include digital pixels per inch (dpi) as metadata. We developed a User Guide 155 that includes a description of file submission requirements that allows data contributors to ensure they understand their data and metadata prior to submission (DeCarlo, 2025).

All cores must contain basic sample collection metadata. These requirements include month and year of collection, latitude, longitude, depth, and coral genus. Additional metadata, described below, are preferred for optimal calibration of skeletal density. For CT scans, data submitters may have developed their own calibration between HU and skeletal density 160 (DeCarlo et al., 2015; Manzello et al., 2014), and if so, that can be added into the metadata linked to each core and used for the final data output from CoralCT. CoralCache also stores user-uploaded CT scans of density standards, which are then available for density calibration of specific cores. For example, if a user scanned three density standards alongside regular cores, they could submit the scans of those three standards and have them saved on CoralCache with a group name that links them to the appropriate cores. In CoralCT, the user can then derive the mean HU of each standard and create a calibration 165 equation against known density, and this equation can be applied to the metadata record of any cores for which that user is listed as a responsible person. Different standard sets and/or calibration equations can be applied to each core. While there is a default calibration equation (DeCarlo et al., 2015), conducting careful density calibrations with suitable standards is essential for accurate density and calcification rate data, especially if the CT scan is not on the regular HU scale (*i.e.*, air is -1000, water is 0, with maximum values of around 2000 for pure CaCO<sub>3</sub>) (Hounsfield, 1973). One way to validate density calibrations is to

170 weigh the physical core and calculate its density from the measured mass and CT-derived volume (an output from CoralCT), and then to compare this density value to the whole-core HU-derived density output (also a CoralCT output). Density calibration for X-rays is more difficult, as there is no standard scale, and the brightness of the images will depend on the settings of the X-ray machine as well as the thickness of the coral slab. Data submitters must develop their own calibration equations, and there are a variety of approaches for this, such as using wedges of aluminium or *Tridacna* spp. shells, pressed  
175 coral powders, or via gamma densitometry (Duprey et al., 2012). X-ray images should also either be corrected for the heel effect and the inverse square law (Duprey et al., 2012) prior to submission, or described in the “notes” section of the metadata that these corrections were not yet applied.

Besides core image data, the observer band maps are stored as .mat files. These are default files for the MATLAB language in which CoralCT is written, but they do not need to be opened directly by a user. It is important to note that CoralCT  
180 runs in MATLAB Runtime, which uses the MATLAB language but does not require the user to have MATLAB installed or any form of license. For CT scans, the .mat files have the same number of rows and columns as the CT scan they are linked to (*i.e.*, each DICOM image contains pixels in rows and columns), and the third dimension represents each band that has been identified. The data stored within this 3-D array are the vertical pixel indices that were clicked by the observer (pixels downward from the top of the core, beginning at index 1), with zeros filled in everywhere else. The .mat files also contain the  
185 following settings used by the observer when marking the bands (DeCarlo et al., 2025b): image brightness and contrast, the size of the filter used for smoothing bands (see DeCarlo et al., 2025), the projection type used (*e.g.*, minimum or mean), the slice thickness in mm, and the angle at which the core was rotated (if applicable). For X-rays, the .mat file likewise contains the band coordinates, but with rows representing the horizontal pixel index of the core image, columns representing the band numbers, and the value of entries representing the vertical pixel index from the top of the image.

190 Finally, output growth rate data are stored as comma separated values (.csv) files. These contain key metadata about the core, the username of the observer, any notes made during analysis, and any acknowledgements or citations linked to the core. The growth rate data are presented as annual linear extension (in centimeters per year [ $\text{cm yr}^{-1}$ ]), density (grams per cubic centimeter [ $\text{g cm}^{-3}$ ]), and calcification (grams per centimeter squared per year [ $\text{g cm}^{-2} \text{yr}^{-1}$ ]). CoralCT measures extension and density, and then calcification is calculated as the product of density and extension (Barnes and Lough, 1993). The calendar

195 years assigned to the annual data are based on counting backward from the observer's interpretation of the first calendar year measured. This is typically the year of collection or the year prior, depending on whether there is a complete or partial band formed at the top of the core (see D'Olivo et al., 2013). These .csv files are consistent in terms of the rows and columns where each piece of metadata and data are placed, enabling the development of automated computer codes for reading information from large sets of these files. Furthermore, the CoralCT application contains an option to export data in Linked Paleo Data  
200 (LiPD) format (McKay and Emile-Geay, 2016).

## 2.2 Data storage and access systems

The organization of the combined CoralCT–CoralCache system is more complex than a typical data repository for two reasons. First, it is an interactive system in which CoralCT allows users to not only access data in the CoralCache repository, but also  
205 continually add traceable interpretations (*i.e.*, band maps) to CoralCache. Therefore, continuous refinement of the banding pattern can be done across multiple sessions. This precludes a system where the files are read-only, and users are allowed to access the full repository directly. Rather, files must be both readable and writable to allow the continual addition and editing of users' band maps. Because at least some files must be writeable, users cannot have direct access to the entire data storage because band maps could be inadvertently edited, moved, or deleted. Second, the CoralCache repository holds datasets for  
210 which CoralCT analyses are incomplete or not yet published. Since most researchers will not want these data publicly accessible until after publication, there cannot be public access to the entire data storage.

Effective use of the CoralCache repository, while maintaining the data security constraints, is enabled through two mechanisms (Fig. 2). First, access to the main data storage is moderated by users logging in with unique usernames (see section below) and only reaching data through the CoralCT application, in which only certain actions are allowed. For example, users  
215 can download only publicly accessible datasets or those in which the core responsible person has granted access to specified users. Similarly, users can access and edit band maps linked to their username, but they can only view (not edit) other users' band maps. The CoralCT application includes buttons to download openly accessible datasets, including options for bulk downloading of all output growth-rate data in entire regions, or all growth-rate datasets linked to a particular core (*i.e.*, from multiple observers). However, these prescribed actions may limit the efficiency of some research applications, such as bulk

220 analysis of all growth-rate files or comparisons of band maps among observers. For this reason, the second mechanism is that there is a copy of the CoralCache repository that contains all publicly accessible files, including the core image data, output growth rate files, and the band-map files associated with the output files. We run a script (DeCarlo, 2025) separate from CoralCT that updates the CoralCache public copy every month by automatically transferring any new or updated files associated with cores that are tagged as publicly accessible from the main data storage. Since these are only copies of the  
225 original files and are not part of the interactive system linked to CoralCT, there is direct and unrestricted access to this data storage (DeCarlo et al., 2025a). With this approach, end-users are able to develop automated scripts that, for example, query the data storage for all files, or all files from a certain user, or all band map files from a particular region, etc. The two-tiered access approach effectively balances user convenience with essential data security. See Data Access section for information on accessing the data storage.

230




Figure 2: Schematic of the CoralCache system, a virtual repository designed to store and share data on coral skeletal cores, and its integration with the CoralCT application.

### 235 2.3 User registration and data submission options

The interactions with the main data storage require users to register in the CoralCT application. This approach is necessary to track which observers have identified bands in specific cores, enabling assessments of observer agreement scores (see below) or potentially weighting different observer contributions. No personal information is required other than an active email address to receive a temporary password. Users are then able to change their password directly in the CoralCT application. By

240 requiring active email addresses, we can subsequently notify all users about updates to the application or general progress notes related to the CoralCache repository. The registration system also enables the band maps (`.mat` files) to be linked to usernames, so that users can save their work and then re-open a core to continue their work at a later time or from a different

computer. Likewise, access to specific cores can be restricted to certain usernames by the core responsible person to protect the sharing of datasets prior to publication.

245

Data submission can proceed by either a flexible submission option or direct submission. In the flexible option, users may submit folders of data or zipped files for DICOMs, or any image file type for X-rays, along with a completed metadata template. Flexible data submissions go to the CoralCT data management team at Tulane University, who subsequently rename files to meet the application requirements and organize them on the CoralCache repository. As long as all of the essential image data and metadata are provided, the flexible submission option is relatively easy for data submitters because they do not need to follow strict file naming or organization requirements. Alternatively, data contributors may use direct submission, which requires DICOM data to be zipped and named in a certain way, X-rays to be submitted in .tiff format, and the metadata template to be completed in a precise way (see description in User Guide, DeCarlo, 2025). If done correctly, the files are directly added to CoralCache and are immediately available for analysis.

255

#### **2.4 Current status of the repository**

Since its inception in 2024, 130 users have registered in the CoralCT application and together have added 1,374 unique cores from across the tropics to the CoralCache repository (Fig. 3). The cores are mostly from living coral colonies, although some were collected from dead colonies with geochemical dating providing a top-of-core date. Others are reef-framework cores that contain fragments of skeletons from corals that built the reef over centuries to millennia. There are 37,108 total annual band traces that have been processed into output growth rate files from 592 measured cores (Fig. 4). Currently, 487 cores are fully publicly accessible, 328 are available for anyone to measure but with output data restricted to the core responsible person, and 559 cores are restricted to access only by the core responsible person and other assigned usernames.




Figure 3: Map showing the locations of all coral skeletal cores (pink dots) currently stored on the CoralCache repository.




Figure 4: Current status of the CoralCache repository. A) Total numbers of cores in the repository per region (black) and numbers of cores that have been processed for growth measurements (hatched). B) Total band measurements including multiple observers per core (black) and excluding multiple observers per core (*i.e.*, using the single longest record per core; hatched).

The directory of cores and associated metadata is viewable in the CoralCT application (Fig. 5). Even though cores may be completely restricted from access by anyone other than the responsible person, all cores are listed in the publicly viewable directory. This enables the community of researchers to know what cores have been collected from a region of interest, potentially helping to spur collaborations via sample discovery and/or avoiding duplicate research efforts and oversampling of Threatened or Endangered coral species. The core directory also lists how many observers have completed analysis of each core, allowing interested researchers to identify data gaps or potential collaborators in need of band interpretations for their cores.

	observers	confidence	citation	acknowledgement	region	sub region	genus	lat	lon
389-M99-E-013R-1	0	NaN		Please acknowledge the United States Geolo...	Hawaii	Island of H...	Porites	19.8352	-156.0924
TAN23_3	4	7.1546		Please acknowledge Juan D'Olivo	Quintana ...	Tanchacte	Pseudodi...	20.9277	-86.8330
TAN23_2	3	8.9255		Please acknowledge Juan D'Olivo	Quintana ...	Tanchacte	Pseudodi...	20.9277	-86.8330
TAN23_1	4	6.0944		Please acknowledge Juan D Olivo if using thi...	Quintana ...	Tanchacte	Pseudodi...	20.9277	-86.8330
IM1	0	NaN		Dr. Isaiah Bolden, Tabarrion G. Stoves, and S...	Jamaica	Portland	Orbicella	18.1779	-76.4120
IM2	0	NaN		Dr. Isaiah Bolden, Tabarrion G. Stoves, and S...	Jamaica	Portland	Orbicella	18.1779	-76.4120
IM3	1	8.4508		Dr. Isaiah Bolden, Tabarrion G. Stoves, and S...	Jamaica	Portland	Orbicella	18.1779	-76.4120
IM4	0	NaN		Dr. Isaiah Bolden, Tabarrion G. Stoves, and S...	Jamaica	Portland	Orbicella	18.1779	-76.4120
2022-P3-T1-CH61_1	1	8.1836		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1810	-169.6574
2022-P3-T1-CH61_2	1	5.5904		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1810	-169.6574
LY01	0	NaN		Please acknowledge Yi-Wei Liu	Taiwan	Lanyu	Porites	22.0000	121.5800
H01	2	7.0084	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3201	-157.6694
H02	3	8.6924	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3202	-157.6694
H03	1	9.0000	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3201	-157.6696
H04	1	9.0161	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3201	-157.6696
H05	1	5.8007	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3200	-157.6695
H06	1	NaN	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3200	-157.6695
H07	1	8.3480	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3200	-157.6695
H08	1	NaN	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3202	-157.6696
H09	1	5.1254	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3201	-157.6696
H10	3	6.9035	<a href="https://doi.org/10.1038/s43247-0...">https://doi.org/10.1038/s43247-0...</a>	Please acknowledge Tom DeCarlo if publishi...	Hawaii	Oahu	Porites	21.3201	-157.6696
2022-P3-T1-CH66_1	2	7.1509		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1809	-169.6573
2022-P3-T1-CH66_2	2	6.9512		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1809	-169.6573
2022-P2-T1-CH02	1	5.9923		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1843	-169.6649
2022-P2-T1-CH09	2	6.9447		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1849	-169.6661
2022-P2-T1-CH09_2	0	NaN		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1849	-169.6661
2022-P3-T1-CH64	2	5.2296		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1813	-169.6582
2022-P3-T1-CH65_1	1	8.7969		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1813	-169.6581
2022-P3-T1-CH65_2	1	0.9620		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1813	-169.6581
2022-P2-T1-CH06	1	7.6053		Please acknowledge the United States Geolo...	US Pacific...	Ofu	Porites	-14.1850	-169.6668
CO-02	0	NaN	<a href="https://doi.org/10.1038/s41598-0...">https://doi.org/10.1038/s41598-0...</a>	Gabriel O. Cardoso, Diego Kersting and Juan...	Antilles	Martinique	Siderastrea	14.4649	61.0180

280


Figure 5: An example section of the coral skeletal core directory currently available on the CoralCache repository showing basic collection metadata.

## 2.5 Inter-observer agreement scores

285 The automated saving of observer band maps by username enables calculation of inter-observer agreement scores, which could be used as quality control of output data given that un-restricted cores can be analyzed by any registered user. We describe here a method for how these scores are currently calculated from CoralCache data and displayed in CoralCT, but we note that other approaches could be developed from data in the public copy of the repository. Additionally, it is important to highlight that these are simply agreement scores, not necessarily a measure of accuracy. They are included for potential future meta-  
290 analyses including growth rate data from cores that have been interpreted by multiple observers, as data analysts may choose to weight or otherwise select individual observer data based on their agreement with other observers across all cores. Thus, beginning in CoralCT version 2.6.5, the output growth rate files contain the inter-observer agreement score (if one exists) for that observer.

295 In the first step of calculating inter-observer agreement scores, we search through all cores in CoralCache and only use data from cores that have at least three observers. Therefore, not all observers will necessarily have an inter-observer agreement score because they may not have interpreted bands on any cores with two or more additional observers. For each core, we calculate an among-observer variance score as follows. For each band in a core, we identify the area over which all observers traced that band. Some observers may have traced across the entire area of the core, while others may have identified a smaller  
300 region; therefore, we focus only on the overlapping horizontal extents (Fig. 6). We then calculate the standard error of the mean of vertical placement of the band for all the overlapping pixels in the horizontal extent. If the band placement by observers is randomly distributed about a mean position, then the standard error of the mean should decrease with more observers, although outlier placements could cause the standard error to increase. We then average the standard error among all bands in the core, and call this “among-observer variance” (*i.e.*, one value per core for a given set of observers).

305



**Figure 6: Illustration of how different observers identify varying horizontal extents of each band of a coral skeletal core in CoralCT, a graphical user interface. The left image shows a 3-D plot of the surfaces identified by four observers for a single band. The band is plotted in a different colour per observer, set to 50% transparency, and vertically exaggerated 2x to improve clarity. The right image shows a ‘top-down’ view with colours indicating the number of observers that traced the band across the horizontal extent of the CT scan. The lightest blue colour shows where all four observers overlapped, and this area would be used in the inter-observer agreement score calculations.**

The second step in calculating the inter-observer agreement score is to identify the set of observers that minimizes the among-observer variance. Because standard error is the standard deviation divided by the square root of the number of observers ( $n$ ), in this case, there will be a tendency for the optimal set of observers to favour retaining, rather than excluding, individuals. However, there is also the possibility that a small number of observers agree very closely even if they diverge from most other observers, and this set of the small number of observers could produce the smallest among-observer variance. For this reason, we set a requirement that the optimal set of observers must include at least half of the total observers. For cores with seven or fewer observers, we calculate among-observer variance for every possible combination of  $n/2$  or more. However, some cores have been interpreted by many observers, up to 26 observers in some cases, which equates to over 10 million possible combinations. For these cores (those with more than seven observers), we use a minimization function that searches

the parameter space to find the minimum value of among-observer variance. We developed this function by inputting a list of observers (*i.e.*, 1, 2, ... *n*) and allowing the search algorithm to adjust these numbers. We then round the adjusted numbers and  
325 retain the unique integers of observers. Effectively, this allows the search algorithm to shift the integers away from ‘poor’ observers and toward ‘good’ observers so that the unique set of integers contains only the ‘good’ observers. We use the simulated annealing function (`simulannealbnd`) in MATLAB to perform this search because it is effective at initially sampling a wide parameter space and then narrowing into the best solution with smaller adjustments. We allow 250 total function evaluations before reaching the optimal set of observers. We note that it is technically possible that there is a better set of  
330 observers beyond the sampled parameter space, but in general, the simulated annealing approach should converge on relatively good sets of observers with low among-observer variances.

The final step is to calculate the inter-observer agreement scores for each username. We calculate this as the proportion of possible cores (those with three or more observers) that an observer interpreted in which that observer’s band map was retained in the optimal set of observers. For example, if a user traced bands in 10 cores that met the requirement of  
335 three or more observers, and the user’s bands were included in the optimal set of observers for seven of those scores, their inter-observer agreement score would be 0.7. We note again that this score represents agreement, not necessarily accuracy, and that there is no set way to interpret or use this score. Rather, it is a potentially useful metric for developing statistical approaches to synthesizing multiple observers per core, and thus it is included in the CoralCT output files. Even if subsequent data analysts develop different approaches, a key step forward in the field is that there can be multiple observers per core, and  
340 it is possible to develop quantitative metrics of the skill of the observer(s), which has not previously been possible in sclerochronology or in related fields such as dendrochronology.

The inter-observer agreement scores are automatically updated once per month. We note that because these scores depend on interactions among all other users’ band maps, a particular user’s score could increase or decrease even without them making new band interpretations. The score printed in the output growth rate files is that user’s inter-observer agreement  
345 score at the time they processed that particular core, but subsequent analyses could re-calculate the inter-user agreement scores to suit a particular research question.

## 2.6 Observer confidence scores

At the time that observers choose to process growth rates from their completed band mapping, they have the option to assign a confidence (0-10) score to that core. These scores are saved in the output growth rate files. Similar to the inter-observer agreement scores, there is no prescribed method for how to use the confidence scores in subsequent analyses. However, data analysts could incorporate the confidence scores in various ways. For example, the contribution of each core to a climate reconstruction based on a network of cores could be weighted relative to the confidence scores for that core. Intuitively, higher confidence in interpreting bands by observers is likely to indicate more accurate data. Since observers may differ in their system for assigning relative confidence scores, data analyses incorporating these scores would likely benefit from standardizing the scores per observer. Full access to the public copy of the data storage is again key for enabling automated scripts for pulling these scores for all observers involved in any particular study. Incorporating observer confidence may further enhance data quality and reliability in large-scale syntheses.

## 2.7 Sustainability

A variety of approaches were used to ensure long-term sustainability of the repository. In earlier versions of the CoralCT application, only the flexible data submission option was available, and nearly all data currently in the repository were submitted in this way. While DICOM is a standardized file format, most data submissions required some organization steps, such as converting submitted CT scan .tiff files to .dcm, rotating the images onto the axial plane, or resizing files, in addition to zipping the files and creating correctly named folders on the data storage. The previous approach, in which the CoralCT data management team performed these organization tasks, was necessary because most data submitted thus far have come from works completed before the CoralCache repository was built in 2024. Data contributors mostly sent data upon request for archiving purposes, and many contributors were no longer active in coral core analysis, so minimizing time requirements for data contributors was crucial to achieving community buy-in. Future data submissions will likely more commonly be from users who plan to use CoralCT for analysis of their new cores, in which case it is reasonable for the data submitters to carefully read the User Guide to learn how to suitably organize their data for the direct submission option. Importantly, the direct

submission option requires no effort from the CoralCT maintenance team, so the repository can continue to grow and be used for new datasets regardless of continued personnel support.

Automated monitoring scripts have also been implemented to reduce time requirements for active management of the repository by the CoralCT team. The data storage is automatically queried for several tasks. If a user forgets their password, they may click a button in the CoralCT application, and a new password is created and emailed to the user. New user registrations are currently approved manually as a means to avoid programmed bot (*i.e.*, non-human) registrants, but the monitoring scripts have options for turning on automating approval for all or certain email addresses such as .gov, .ac, or .edu (DeCarlo, 2025). Additionally, the entire repository is backed up onto two separate external hard drives every three days. Finally, once per month, all open-access files are automatically sent from the main data storage to the public copy, and inter-observer agreement scores are updated. These monitoring scripts are provided in DeCarlo (2025), with only the main data storage login credentials redacted. Likewise, the code underlying CoralCT is available at <https://github.com/tdecarlo19/CoralCT>, and has nearly full functionality except that it is linked to a separate data storage with only example datasets.

The whole repository can run with minimal human effort for maintenance. Currently, the CoralCT team is needed to handle datasets submitted via the flexible option, confirm new user registrations without .edu email addresses, and ensure the monitoring scripts are running. Handling data submissions is by far the most time-consuming task because it can involve problem-solving, depending on the state of submitted files. However, if this support were to cease, the organization steps would simply shift onto data submitters without any loss of repository functioning. If needed, the remaining tasks of confirming new users and running automation scripts could readily be handed over to another team or organization. Beyond the publicly available scripts, only data storage login credentials are needed to run the system. As described above, refraining from public release of direct access to the main data storage is a necessary requirement of the interactive link between the CoralCT application and CoralCache repository, and for the option to keep pre-publication datasets in the repository.

### 3 Data availability

The data are available via the public copy of the CoralCache data storage. The current version of CoralCache is available at <https://doi.org/10.5281/zenodo.17208911> (DeCarlo et al., 2025a). Additionally, login information to access all of the data via a FTP client is available at <https://doi.org/10.5281/zenodo.17790935>.

#### 400 **4 Code availability**

The code for the CoralCT application and its interactions with the public copy of the CoralCache data storage is available at <https://github.com/tdecarlo19/CoralCT>. This code has been slightly redacted to remove features associated with usernames and access to the private, pre-publication component of the data storage. The desktop CoralCT application is available following DeCarlo et al. (2025b).

405

#### **5 Conclusions**

Historically, coral core image data have not been shared publicly or placed in a repository, and analyses of growth banding in those cores have largely been opaque and thus difficult to reproduce. While this was incompatible with the intent and arguably the rule of data sharing requirements of most funders and publishers, it was understandable given the lack of a repository for these types of data, especially the large file sizes associated with CT scans. CoralCT and CoralCache together remove these barriers to data archiving and transparency. The data organization and sharing system also meets the FAIR principles (Wilkinson et al., 2016) because the metadata in the core directory are consistent and the datasets are findable; post-publication data are readily accessible through the CoralCT application; output files are consistently formatted so that they are interoperable across software platforms; and the data are reusable because previously published datasets in CoralCache can be gathered into regional- or global-scale meta-analyses. With a graphical user interface and an extensive User Guide, the application and repository are easy to use. Furthermore, CoralCache enhances the robustness of coral growth analyses by enabling objective evaluation of inter-observer agreement and confidence in band interpretations. This additional layer of transparency strengthens the credibility of research findings and enables Big Data syntheses of growth variations in hundreds or thousands of cores. While we do not intend to prescribe a single method, the CoralCT application does offer ease-of-use for 2-D or 3-D growth rate measurements that are then automatically archived and transparent upon public release. Most importantly, even submitting core image data in CoralCache can fulfil a crucial obligation of data sharing for authors working

415

420

with coral cores, an obligation that is still unfortunately often bypassed by publishing only summarized data without the images necessary for reproducibility (e.g. Galochkina et al., 2026). Given the ability to protect pre-publication datasets, we argue that there is no downside to archiving in CoralCache, and that funders, publishers, and reviewers no longer need to accept claims  
425 for the lack of a suitable repository to meet data sharing requirements for coral core image datasets.

### **Author contributions**

T.M.D. conceived of the repository design with input from O.J. and A.S. All authors contributed data and critically evaluated  
430 the manuscript.

### **Competing interests**

The authors declare that they have no conflict of interest.

### **435 Acknowledgements**

This research was supported by the Louisiana Board of Regents Endowed Professorships subprogram and by NSF award OCE-2444864 to T.M.D. All substantial data contributors are included as authors, but we still wish to acknowledge all those who helped grow CoralCache, past and present. We also thank Dr. Bryan Black for inspiration. M.M. acknowledges the support from FAPESP grant #23/09180-3 for the production of coral skeleton cores. T.F. acknowledges Deutsche  
440 Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 469906366 (T.F.) – SPP 2299/Project number 441832482. D.K.K. acknowledges Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 401447620. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement of the U.S. Government. The scientific results and conclusions, as well as any views or opinions expressed herein, are those of the author(s) and do not necessarily reflect the views of NOAA or the Department of Commerce.

445

## References

- Atwood, A. R., Moore, A. L., DeLong, K. L., Long, S. E., Sanchez, S. C., Hargreaves, J. A., Morris, C. A., Pauly, R. E., Dassié, É. P., Felis, T., Voelker, A. H. L., Murty, S. A., and Cobb, K. M.: The PAGES CoralHydro2k Seawater  $\delta^{18}\text{O}$  Database: a FAIR-aligned compilation of seawater  $\delta^{18}\text{O}$  data to uncover “hidden” insights from the global ocean, *Earth System Science Data*, 18, 1921–1941, <https://doi.org/10.5194/essd-18-1921-2026>, 2026.
- 450
- Barkley, H., Cohen, A., Mollica, N., Brainard, R., Rivera, H., DeCarlo, T., Lohmann, G., Drenkard, E., Alpert, A., Young, C., Vargas-Angel, B., Lino, K., Oliver, T., Pietro, K., and Luu, V.: Repeat bleaching of a central Pacific coral reef over the past six decades (1960–2016), *Communications Biology*, 1, 177, 2018.
- Barkley, H. C. and Cohen, A. L.: Skeletal records of community-level bleaching in *Porites* corals from Palau, *Coral Reefs*, 35, 1407–1417, <https://doi.org/10.1007/s00338-016-1483-3>, 2016.
- 455
- Barnes, D. J. and Lough, J. M.: On the nature and causes of density banding in massive coral skeletons, *J. Exp. Mar. Biol. Ecol.*, 167, 91–108, 1993.
- Baumann, J. H., Ries, J. B., Rippe, J. P., Courtney, T. A., Aichelman, H. E., Westfield, I., and Castillo, K. D.: Nearshore coral growth declining on the Mesoamerican Barrier Reef System, *Global Change Biology*, 25, 3932–3945, <https://doi.org/10.1111/gcb.14784>, 2019.
- 460
- Cantin, N. E. and Lough, J. M.: Surviving Coral Bleaching Events: *Porites* Growth Anomalies on the Great Barrier Reef, *PLoS One*, 9, e88720, 2014.
- Cantin, N. E., Cohen, A. L., Karnauskas, K. B., Tarrant, A. M., and McCorkle, D. C.: Ocean warming slows coral growth in the central Red Sea, *Science*, 329, 322, <https://doi.org/DOI:10.1126/science.1190182>, 2010.
- 465
- Carilli, J. E., Norris, R. D., Black, B., Walsh, S. M., and McField, M.: Century-scale records of coral growth rates indicate that local stressors reduce coral thermal tolerance threshold, *Glob. Change Biol.*, 16, 1247–1257, <https://doi.org/10.1111/j.1365-2486.2009.02043.x>, 2009.
- Carricart-Ganivet, J. P., Cabanillas-Terán, N., Cruz-Ortega, I., and Blanchon, P.: Sensitivity of Calcification to Thermal Stress Varies among Genera of Massive Reef-Building Corals, *PLoS One*, 7, e32859, 2012.
- 470
- Castillo, K. D., Ries, J. B., Weiss, J. M., and Lima, F. P.: Decline of forereef corals in response to recent warming linked to history of thermal exposure, *Nature Climate Change*, 2, 756–760, <https://doi.org/10.1038/nclimate1577>, 2012.
- Cooper, T. F., De’ath, G., Fabricius, K. E., and Lough, J. M.: Declining coral calcification in massive *Porites* in two nearshore regions of the northern Great Barrier Reef, *Global Change Biology*, 14, 529–538, <https://doi.org/10.1111/j.1365-2486.2007.01520.x>, 2008.
- 475
- Cooper, T. F., O’Leary, R. A., and Lough, J. M.: Growth of Western Australian corals in the anthropocene., *Science (New York, N.Y.)*, 335, 593–6, <https://doi.org/10.1126/science.1214570>, 2012.
- Courtney, T. A., Kindeberg, T., and Andersson, A. J.: Coral calcification responses to the North Atlantic Oscillation and coral bleaching in Bermuda, *PLOS ONE*, 15, e0241854, <https://doi.org/10.1371/journal.pone.0241854>, 2020.
- 480
- Dassié, É., DeLong, K. L., Kilbourne, H., Williams, B., Abram, N. J., Brenner, L., Brahmi, C., Cobb, K. M., Corrège, T., Dissard, D., Emile-Geay, J., Evangelista, H., Evans, M. N., Farmer, J., Felis, T., Gagan, M., Gillikin, D. P., Goodkin, N., Khodri, M., Lavagnino, A., Lavigne, M., Lazareth, C. E., Linsley, B., Lough, J., Mcgregor, H. V., Nurhati, I., Ouellette, G.,

- Perrin, L., Raymo, M., Rosenheim, B., Sandstrom, M., Schöne, B., Sifeddine, A., Stevenson, S., Thompson, D., Waite, A., Wanamaker, A., and Wu, H. C.: Saving Our Marine Archives, *Eos, Transactions American Geophysical Union*, 98, <https://doi.org/10.1029/2017EO068159>, 2017.
- 485 De'ath, G., Lough, J. M., and Fabricius, K. E.: Declining coral calcification on the Great Barrier Reef, *Science*, 323, 116, <https://doi.org/DOI: 10.1126/science.1165283>, 2009.
- DeCarlo, T.: Server connection and monitoring files for CoralCache virtual coral core repository, , <https://doi.org/10.5281/zenodo.17790935>, 2025.
- 490 DeCarlo, T., Jasnos, O., Strange, A., Andersson, A., Bautista VII, A., Bloomer, S., Bolden, I., Bosman, M., Brachert, T., Braz, G., Cardoso, G., Carricart-Ganivet, J., Carilli, J., Castillo, K., Cavole, L., Chan, S., Chen, X., Chomitz, B., Correge, T., Courtney, T., Deigan, M., D'Olivo, J. P., Dunbar, R., Enochs, I., Falsarella, L., Felis, T., Gutiérrez-Estrada, G., Hedger, B., Hu, S., Jameson, S., Jupiter, S., Kench, P., Kersting, D., Lin, K., Liu, Y.-W., Lorigados, C., Manzello, D., McCulloch, M., Mies, M., Moura, R., Oberle, F., Pereira, N., Prouty, N., Ramos, R., Ren, H., Ryan, E., Thompson, D., Toth, L., Vergotti, M., Webster, J., and Zinke, J.: CoralCache coral core image dataset (1.0), <https://doi.org/10.5281/ZENODO.17208911>, 2025a.
- 495 DeCarlo, T. M.: The past century of coral bleaching in the Saudi Arabian central Red Sea, *PeerJ*, 8, e10200, <https://doi.org/10.7717/peerj.10200>, 2020.
- DeCarlo, T. M. and Cohen, A. L.: Dissepiments, density bands and signatures of thermal stress in *Porites* skeletons, *Coral Reefs*, 36, 749–761, <https://doi.org/10.1007/s00338-017-1566-9>, 2017.
- 500 DeCarlo, T. M., Cohen, A. L., Barkley, H. C., Cobban, Q., Young, C., Shamberger, K. E., Brainard, R. E., and Golbuu, Y.: Coral macrobioerosion is accelerated by ocean acidification and nutrients, *Geology*, 43, 7–10, <https://doi.org/10.1130/G36147.1>, 2015.
- DeCarlo, T. M., Gajdzik, L., Ellis, J., Coker, D. J., Roberts, M. B., Hammerman, N. M., Pandolfi, J. M., Monroe, A. A., and Berumen, M. L.: Nutrient-supplying ocean currents modulate coral bleaching susceptibility, *Science Advances*, 6, eabc5493, 2020.
- 505 DeCarlo, T. M., Cotton, J., Whelehan, A., Gramse, M., Berumen, M. L., Harrison, H. B., McCulloch, M. M., Whitaker, H. V., Falk, T., Groenvall, E., and Matthews, K.: Calcification trends in long-lived corals across the Indo-Pacific during the industrial era, *Commun Earth Environ*, 5, 1–9, <https://doi.org/10.1038/s43247-024-01904-8>, 2024.
- DeCarlo, T. M., Whelehan, A., Hedger, B., Perry, D., Pompel, M., Jasnos, O., and Strange, A.: CoralCT: A platform for transparent and collaborative analyses of growth parameters in coral skeletal cores, *Limnology and Oceanography: Methods*, 510 23, 97–116, <https://doi.org/10.1002/lom3.10661>, 2025b.
- DeLong, K. L., Quinn, T. M., Taylor, F. W., Shen, C., and Lin, K.: Improving coral-base paleoclimate reconstructions by replicating 350years of coral Sr/Ca variations, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 373, 6–24, <https://doi.org/10.1016/j.palaeo.2012.08.019>, 2013.
- 515 D'Olivo, J. P., McCulloch, M. T., and Judd, K.: Long-term records of coral calcification across the central Great Barrier Reef: assessing the impacts of river runoff and climate change, *Coral Reefs*, 32, 999–1012, <https://doi.org/10.1007/s00338-013-1071-8>, 2013.
- Duprey, N., Boucher, H., and Jiménez, C.: Digital correction of computed X-radiographs for coral densitometry, *Journal of Experimental Marine Biology and Ecology*, 438, 84–92, <https://doi.org/10.1016/j.jembe.2012.09.007>, 2012.

- 520 Felten, J., Möller, L., Schindler, U., Huber, R., Schumacher, S., Koppe, R., Diepenbroek, M., and Glöckner, F. O.: PANGAEA - Data Publisher for Earth & Environmental Science, *Sci Data*, 10, 347, <https://doi.org/10.1038/s41597-023-02269-x>, 2023.
- Felis, T., Suzuki, A., Kuhnert, H., Dima, M., Lohmann, G., and Kawahata, H.: Subtropical coral reveals abrupt early-twentieth-century freshening in the western North Pacific Ocean, *Geology*, 37, 527–530, <https://doi.org/10.1130/G25581A.1>, 2009.
- Fouke, K. W., Trop, J. M., and Sivaguru, M.: Changes in Coral Skeleton Growth Recorded by Density Band Stratigraphy, Crystalline Structure, and Hiatuses, *Front. Mar. Sci.*, 8, <https://doi.org/10.3389/fmars.2021.725122>, 2021.
- 525 Galochkina, M., Cohen, A. L., Oppo, D. W., and Ummenhofer, C. C.: Climate modes can be leveraged to forecast coral bleaching months in advance, *Commun Earth Environ*, <https://doi.org/10.1038/s43247-026-03438-7>, 2026.
- Helmle, K., Kohler, K., and Dodge, R.: Relative optical densitometry and the Coral X-radiograph Densitometry System: Coral XDS, International Society of Reef Studies European Meeting, 2002.
- 530 Hounsfield, G. N.: Computerized transverse axial scanning (tomography): Part 1. Description of system, *British Journal of Radiology*, 46, 1016–1022, <https://doi.org/10.1259/0007-1285-46-552-1016>, 1973.
- Hudson, J. H.: Growth rates in *Montastraea annularis*: a record of environmental change in Key Largo Coral Reef Marine Sanctuary, Florida, *Bulletin of Marine Science*, 31, 444–459, 1981.
- Knutson, D. W., Buddemeier, R. W., and Smith, S. V.: Coral chronometers: seasonal growth bands in reef corals., *Science*, 177, 270–2, <https://doi.org/10.1126/science.177.4045.270>, 1972.
- 535 Lough, J.: Coral calcification from skeletal records revisited, *Marine Ecology Progress Series*, 373, 257–264, <https://doi.org/10.3354/meps07398>, 2008.
- Lough, J. M. and Barnes, D. J.: Environmental controls on growth of the massive coral *Porites*, *J. Exp. Mar. Biol. Ecol.*, 245, 225–243, 2000.
- Ma, T. Y. H.: On the growth rate of reef corals and the sea water temperature in the Japanese Islands during the latest geological times, *Geology*, 16, 165–189, 1934.
- 540 Mallela, J., Hetzinger, S., and Halfar, J.: Thermal stress markers in *Colpophyllia natans* provide an archive of site-specific bleaching events, *Coral Reefs*, 35, 181–186, <https://doi.org/10.1007/s00338-015-1350-7>, 2015.
- Manzello, D. P.: Coral growth with thermal stress and ocean acidification: lessons from the eastern tropical Pacific, *Coral Reefs*, 29, 749–758, 2010.
- 545 Manzello, D. P., Enochs, I. C., Bruckner, A., Renaud, P. G., Kolodziej, G., Budd, D. A., Carlton, R., and Glynn, P. W.: Galápagos coral reef persistence after ENSO warming across an acidification gradient, *Geophysical Research Letters*, 41, 9001–9008, <https://doi.org/10.1002/2014GL062501>, 2014.
- McKay, N. P. and Emile-Geay, J.: Technical note: The Linked Paleo Data framework – a common tongue for paleoclimatology, *Climate of the Past*, 12, 1093–1100, <https://doi.org/10.5194/cp-12-1093-2016>, 2016.
- 550 Mollica, N. R., Cohen, A. L., Alpert, A. E., Barkley, H. C., Brainard, R. E., Carilli, J. E., DeCarlo, T. M., Drenkard, E. J., Lohmann, P., Mangubhai, S., Pietro, K. R., Rivera, H. E., Rotjan, R. D., Scott-Buechler, C., Solow, A. R., and Young, C. W.: Skeletal records of bleaching reveal different thermal thresholds of Pacific coral reef assemblages, *Coral Reefs*, 38, 743–757, <https://doi.org/10.1007/s00338-019-01803-x>, 2019.

- 555 Oberle, F. K. J., Toth, L. T., Prouty, N. G., Santos, B., Jacobs, J. A., Bloomer, S., Bagheri, K., Williams, B. N., Padgett, J. S., Stathakopoulos, A., and La Selle, S.: The USGS Rotating X-Ray Computed Tomography (RXCT) Coral-Core Archive: Scope, Access, and Standardization, *Journal of Marine Science and Engineering*, 14, 490, <https://doi.org/10.3390/jmse14050490>, 2026.
- 560 Rodgers, K. S., Richards Donà, A., Stender, Y. O., Tsang, A. O., Han, J. H. J., Weible, R. M., Prouty, N., Storlazzi, C., and Graham, A. T.: Rebounds, regresses, and recovery: A 15-year study of the coral reef community at Pila'a, Kaua'i after decades of natural and anthropogenic stress events, *Marine Pollution Bulletin*, 171, 112306, <https://doi.org/10.1016/J.MARPOLBUL.2021.112306>, 2021.
- Saenger, C., Cohen, A. L., Oppo, D. W., Halley, R. B., and Carilli, J. E.: Surface-temperature trends and variability in the low-latitude North Atlantic since 1552, *Nature Geoscience*, 2, 492–495, <https://doi.org/10.1038/ngeo552>, 2009.
- 565 Samperiz, A., Sosdian, S., Hendy, E., Johnson, K., John, E. H., Jupiter, S. D., and Albert, S.: Coastal seawater turbidity and thermal stress control growth of reef-building *Porites* spp. corals in Fiji, *Sci Rep*, 15, 17172, <https://doi.org/10.1038/s41598-025-02283-6>, 2025.
- Thompson, D. M.: Environmental records from coral skeletons: A decade of novel insights and innovation, *Wiley Interdisciplinary Reviews: Climate Change*, 13, e745, <https://doi.org/10.1002/WCC.745>, 2022.
- 570 Vergotti, M. J., D'Olivo, J. P., Brachert, T. C., Capdevila, P., Garrabou, J., Linares, C., Spreter, P. M., and Kersting, D. K.: Reconstruction of long-term sublethal effects of warming on a temperate coral in a climate change hotspot, *Journal of Animal Ecology*, 94, 125–138, <https://doi.org/10.1111/1365-2656.14225>, 2025.
- 575 Walter, R. M., Sayani, H. R., Felis, T., Cobb, K. M., Abram, N. J., Arzey, A. K., Atwood, A. R., Brenner, L. D., Dassíe, É. P., DeLong, K. L., Ellis, B., Emile-Geay, J., Fischer, M. J., Goodkin, N. F., Hargreaves, J. A., Kilbourne, K. H., Krawczyk, H., McKay, N. P., Moore, A. L., Murty, S. A., Ong, M. R., Ramos, R. D., Reed, E. V., Samanta, D., Sanchez, S. C., Zinke, J., and the PAGES CoralHydro2k Project Members: The CoralHydro2k database: a global, actively curated compilation of coral  $\delta^{18}\text{O}$  and Sr&thinsp;&thinsp;Ca proxy records of tropical ocean hydrology and temperature for the Common Era, *Earth System Science Data*, 15, 2081–2116, <https://doi.org/10.5194/essd-15-2081-2023>, 2023.
- 580 Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Sci Data*, 3, 160018, <https://doi.org/10.1038/sdata.2016.18>, 2016.
- 585 Williams, B. N., Schreppel, H. A., Reich, C. D., Smith, K. E. L., Tiling-Range, G., Stalk, C. A., Douglas, S. H., Dadisman, S. V., Flocks, J. G., Toth, L. T., and Stathakopoulos, A.: St. Petersburg Coastal and Marine Science Center's Geologic Core and Sample Database, <https://doi.org/10.5066/F7319TR3>, 2017.