

Rescue, Integration, and Analytical Application of historical data from eight pioneering geomagnetic observatories in China

Suqin Zhang¹, Changhua Fu¹, Jianjun Wang², Chuanhua Chen³, Guohao Zhu⁴, Qian Zhao⁵, Jun Chen⁴, Shaopeng He⁶, Bin Wang⁶, Pengkun Guo⁶, Na Deng⁷, Jinghui Lu⁸, Hongchi Yu⁹

¹Institute of Geophysics, China Earthquake Administration, Beijing, 100081, China

²Earthquake Administration of Gansu Province, Lanzhou, 730000, China

³Earthquake Administration of Shandong Province, Jinan, 250014, China

⁴Shanghai Earthquake Agency, Shanghai, 200062, China

⁵Earthquake Administration of Liaoning Province, Shenyang, 110034, China

⁶Hebei Earthquake Agency, Hebei Province, Shijiazhuang, 050022, China

⁷Earthquake Administration of Hubei Province, Wuhan, 430071, China

⁸Earthquake Administration of Guangdong Province, Guangzhou, 510070, China

⁹Earthquake Administration of Jilin Province, Guangzhou, 130117, China

Correspondence to: Changhua Fu (fuchanghua2004@163.com)

Abstract. Decades to centuries of continuous geomagnetic observation data have extensive scientific research and practical application value, especially in revealing the long-term variation of the geomagnetic field, which is irreplaceable. During the International Geophysical Year (1957-1958), China established geomagnetic observatories in Beijing, Lhasa, Lanzhou, Wuhan, Guangzhou, Changchun, and Urumqi, forming the initial structure of China's geomagnetic observation network together with the Shanghai Observatory. These observatories have continuously observed despite facing many challenges since their establishment, accumulating a large amount of valuable observational data, making significant contributions to the progress of geomagnetic scientific research and development. However, the scattered storage state of these historical data and the potential risk of damage pose a threat to the integrity and reliability of the data. This study conducted a rescue integration of the historical observational data from eight pioneering geomagnetic observatories in China, significantly improving data quality and facilitating long-term preservation and use of the data. This article introduces the basic conditions of eight observatories, including their locations, changes in location, observation environments, magnetic rooms, the magnetism of building materials, the layout of building facilities, measuring instruments, etc. These are the main prerequisites and foundations for ensuring the quality of observation data. Then, it introduces the integration and processing of historical data, including data collection, digitization, unification of formats, anomaly detection, and processing. Subsequently, the processed data were validated, including assessments of daily variations accuracy and long-term stability. The results show that the quality of the integrated historical data has been significantly improved. These datasets are of great value for improving historical geomagnetic field models, studying variable fields, main geomagnetic fields, and their long-term variations. Finally, we applied the data to the analysis and research of Sq and geomagnetic jerks, exploring the spatiotemporal variation characteristics of Sq and jerk in the China. Sq is mainly a daytime phenomenon, and its variation pattern in the middle and low latitude regions is mainly characterized by its dependence on latitude and local time. The geomagnetic jerk phenomenon exhibits significant regional differences and asynchronous occurrence times of jerks. Jerk events in 1969, 1979, 1991, 2003, and 2019 were observed at all observatories and had distinct jerk variation characteristics. Other jerks were only observed at some observatories or individual observatories. The maximum time difference for the occurrence of the same jerk event at different observatories was 2 years. This study aims to provide these precious datasets to the scientific community and the public so that these data can be integrated with data from other sources, thereby further exploring the spatiotemporal evolution and physical mechanisms of the geomagnetic field. The historical datasets of the eight geomagnetic observatories that have been integrated and quality controlled are available at <https://doi.org/10.5281/zenodo.15481895> (Zhang et al., 2025).

删除[susu]: rules

删除[susu]: <https://doi.org/10.5281/zenodo.14560950>

删除[susu]: 2024b

45 1 Introduction

Geomagnetism is an observational science. Geomagnetic observation data contains rich information about the near-Earth space. By studying this information, we can not only gain a deep understanding of the physical properties and dynamics within the Earth, but also investigate the structure of the magnetosphere and ionosphere, as well as the impact of solar activity on the human living environment, which holds significant importance. The application scope of geomagnetic observation data is extensive, covering fields such as aerospace, communication, mineral resource exploration, underground pipeline networks, power systems, oil drilling, space weather forecasting, and earthquake prediction (Reay et al., 2005; Marshall et al., 2011; Bolduc et al., 2002; Boteler et al., 1998; Liu et al., 2008, 2016; Liu et al., 2009), attracting widespread attention from scientists around the world. In particular, constructing historical geomagnetic models and researching the evolution of the Earth's magnetic field and its related dynamic processes largely depend on precise measurement data of long-term changes in the geomagnetic field provided by geomagnetic observatories (Clarke et al., 2009; Soloviev, et al., 2018; Gillet et al., 2013). Therefore, establishing and maintaining permanent geomagnetic observatories with a certain density and uniform distribution is crucial.

In 1834, German scientist Gauss established the world's first geomagnetic observatory to measure the three-component absolute values of the geomagnetic field—the Göttingen Geomagnetic Observatory. The establishment of the Göttingen Geomagnetic Observatory marked the beginning of systematic observation and research of geomagnetic field changes. To gain a more comprehensive understanding and mastery of the change patterns of the geomagnetic field, scientists realized the necessity of establishing more observatories worldwide. Currently, over 130 geomagnetic observatories globally have joined INTERMAGNET worldwide, providing continuous, quasi-real-time online data (<http://www.intermagnet.org>). These observatories are distributed around the world, forming a vast network for geomagnetic field observations.

In China, the first geomagnetic observatory was established in Beijing by Russian scientists in 1870 but ceased operations in 1882. In 1874, French missionaries established a geomagnetic observatory in Xujiahui, Shanghai, which was relocated to Lujiazui in 1908 and then to Sheshan in 1932, where it has continued to operate until now, becoming one of the longest-running geomagnetic observatories globally with nearly 150 years of continuous observation records. Additionally, the United Kingdom and Germany set up geomagnetic observatories in Hong Kong and Qingdao respectively. From the 1930s to the 1940s, Chinese scholars established observatories in places such as Zijin Mountain in Nanjing, Liangfeng in Guangxi, Guilin, and Beibei in Chongqing, but these observatories ceased operations before 1944.

During the International Geophysical Year, geomagnetic observatories were established in Beijing, Lhasa, Lanzhou, Wuhan, Guangzhou, Changchun, and Urumqi by China, together with the Shanghai Observatory, forming the eight pioneering geomagnetic observatories in China, collectively known as the “Old Eight observatories.” The layout of these observatories is roughly cross-shaped (Figure 1), forming the basic framework of geomagnetic observation network of China(Cheng,

删除[susu]: observation

1995). They undertake the critical task of monitoring the spatiotemporal changes of the geomagnetic field in China, laying the foundation for the construction and development of China's geomagnetic network, and making significant contributions to the geomagnetic scientific research and development in China. In 1984, the “Old Eight observatories” were awarded the "International Centennial Geomagnetic Observation Commemorative Medal" by the International Association of
80 Geomagnetism and Aeronomy (IAGA) in recognition of their outstanding contributions in the global field of geophysical observation. As the "Old Eight observatories" with a long history of observations, they have accumulated a large amount of valuable data, becoming an important foundation for research in geomagnetism. However, historical materials are scattered, with a large portion existing in paper form, which is extremely inconvenient to use. Moreover, due to the long storage period, complex and harsh preservation conditions, these precious materials are facing severe risks of damage. Over time, these
85 materials may be lost or damaged due to natural wear and tear, thus affecting their application in scientific research. Therefore, the rescue work of historical materials has always been a focus of attention in the global scientific community (Jonkers et al., 2003; Chulliat and Telali., 2007; Dawson et al., 2009; Dong et al., 2009; Korte et al., 2009; Reay et al., 2013; Morozova et al., 2014, 2021; Zhao et al., 2017; Sergeyeve et al., 2021; Zhang et al., 2022a) .

Thanks to the advancement of the national special project for the rescue of historical data in the field of earthquakes by the
90 China Earthquake Administration, we have carried out the rescue, organization, and analysis of these historical data First, we collected observational data that was stored in various locations and in multiple media formats. Then, we digitized the paper materials using digital photography and manual entry, and stored data from different periods and formats in a unified database for management. Subsequently, we conducted quality checks and processing on the data in the database, and validated the processed data. Finally, we applied and analyzed these data. Systematically organizing and summarizing these
95 data is of great significance for understanding the long-term evolution patterns of the geomagnetic field and and it also provides valuable data support for future related research.




Figure 1: Distribution map of the eight pioneering geomagnetic observatories in China

2 Data Acquisition

100 Ensuring data quality is crucial for the reliability of research outcomes and the effectiveness of their application. The quality of geomagnetic data is influenced by a variety of factors, including the regional geomagnetic background conditions of the observation site (such as the foundation, topography, etc.), Magnetic field gradient of the observation site, the magnetic properties of the building materials used, the layout of the magnetic room, the insulation of the observation room, the configuration and updating of observation instruments, measurement arrangements, and data processing, among others

105 (Jankowski and Sucksdorff 1996; CEA, 2004; Kudin, 2021; Linthe et al. 2013; Zhang et al. 2024a). This section will provide a detailed introduction to the geomagnetic observation conditions at various observatories, including observatory site selection and historical changes, magnetic environment of the observatories, building materials, magnetic room layout, insulation conditions of the observation room, and the configuration of observation instruments.

删除[susu]: study

2.1 Basic Information of observatories

110 Table 1 shows the location information of the eight pioneering geomagnetic observatories and their changes. These data, covering the latitude and longitude of each site, the start and end years of the observatory's operation, the distance between old and new sites, whether the observatories joined INTERMAGNET, are included in the yearbooks.

Table 1 Changes in the location of geomagnetic observatories

<u>IAGA CODE</u>	<u>Information</u>	<u>Location 1</u>	<u>Location 2</u>	<u>Location 3</u>	<u>In INTERMAGNET</u>
<u>CNH</u>	<u>LAT, LON</u>	<u>43.8°N, 125.3°E</u>	<u>44.0°N, 125.2°E</u>	<u>44.1°N, 125.0°E</u>	
	<u>Distance1</u>		<u>23.6km</u>	<u>/</u>	<u>/</u>
	<u>Distance2</u>	<u>/</u>		<u>19.5km</u>	<u>/</u>
	<u>Operating year</u>	<u>1957-1978</u>	<u>1979-2007</u>	<u>2008-present</u>	
<u>WMQ</u>	<u>LAT, LON</u>	<u>43.8°N, 87.7°E</u>	<u>44.4N, 86.9°E</u>	<u>/</u>	<u>/</u>
	<u>Distance1</u>		<u>92.4km</u>	<u>/</u>	<u>/</u>
	<u>Operating year</u>	<u>1978-2012</u>	<u>2013-present</u>	<u>/</u>	
<u>BJI</u>	<u>LAT, LON</u>	<u>40.0°N, 116.2°E</u>	<u>/</u>	<u>/</u>	<u>/</u>
	<u>Operating year</u>	<u>1957-present</u>	<u>/</u>	<u>/</u>	
<u>LZH</u>	<u>LAT, LON</u>	<u>36.1°N, 103.9°E</u>	<u>/</u>	<u>/</u>	<u>yes</u>
	<u>Operating year</u>	<u>1959-present</u>	<u>/</u>	<u>/</u>	
<u>SSH</u>	<u>LAT, LON</u>	<u>31.1°N, 121.2°E</u>	<u>/</u>	<u>/</u>	<u>/</u>
	<u>Operating year</u>	<u>1933-present</u>	<u>/</u>	<u>/</u>	<u>/</u>
<u>WHN</u>	<u>LAT, LON</u>	<u>30.5°N, 114.6°E</u>	<u>30.5°N, 114.5°E</u>	<u>30.9°N, 113.3°E</u>	
	<u>Distance1</u>		<u>9.6km</u>	<u>/</u>	<u>/</u>
	<u>Distance2</u>	<u>/</u>		<u>123.0km</u>	
	<u>Operating year</u>	<u>1959-1998</u>	<u>1999-2017</u>	<u>2018-present</u>	

LSA	LAT, LON	29.6°N, 91.0°E	/	/	/
	Operating year	1957-present	/	/	/
GZH	LAT, LON	23.1°N, 113.3°E	23.0°N, 112.5°E	/	
	Distance	82.6km		/	yes
	Operating year	1958-1995	2002-present	/	

115 2.1.1 Changchun observatory (IAGA code CNH)

The Changchun observatory, started construction in 1951 and is located in the Nanling area (43.8°N,125.3°E) in the southern suburbs of Changchun City, Jilin Province. The observatory was completed in 1954 and underwent instrument installation and trial observations from 1954 to 1956. Formal observation work began in 1957. The geology foundation of the observatory and surrounding area is composed of Quaternary alluvial layers, approximately 30 meters thick, consisting of loess and sub-clay. The magnetic field around the observatory is homogeneous, with a horizontal gradient less than 0.5 nT per meter. The absolute measurement house and variation recording room are integrated, with the absolute measurement house being a ground-level building and the recording room a semi-underground structure. It is built using weak magnetic materials such as limestone, wood, and copper. The range of the annual temperature in the recording room is 16°C, and the range of the daily temperature is 0.5°C.

125 Due to the rapid urbanization of Changchun City, the geomagnetic observations have inevitably been disturbed. Therefore, at the end of 1978, the Changchun observatory was relocated to Helong Town, Nong'an County, northwest of Changchun City (latitude 44.0°N, longitude 125.2°E). According to documented records (Sun, 2013), the site differences between the new and old observatories are D: -9.0', H: 26.6 nT, Z: -188.2 nT. The geology foundation of this site and surrounding area is Cretaceous strata, consisting of mudstone, shale, and fine sandstone. The magnetic field around the observatory is homogeneous, with a horizontal gradient less than 1 nT per meter. The absolute measurement house is a ground-level building, and the recording room is a semi-underground building. Both are constructed using weak-magnetic materials such as limestone, white bricks, pearl powder, copper, and wood. The annual temperature difference in the recording room is about 24°C, and the daily temperature difference is about 0.5°C.

删除[susu]: e observatory

135 Unfortunately, the construction of rail transit has once again caused interference to the geomagnetic observation. Consequently, the Changchun observatory was relocated to Sangan Township, Nong'an County (latitude 44.1°, longitude 125.0°) in 2007, and began normal observation on June 1st of the same year. The site differences between the new and old observatories are D: -4.56', H: -45.6 nT, Z: -145.8 nT (Sun, 2013). The site is built on a base of Cretaceous Songhua River Group thick sedimentary strata, consisting of non-magnetic and weakly magnetic mudstones, siltstones, and sandstones, with a horizontal gradient of the ground magnetic field less than 1nT per meter. Both the absolute measurement house and the relative recording room are constructed with weak magnetic materials. The measurement house is a ground-level building,

删除[susu]: observatory

while the recording room is a fully underground structure, with loess covering providing natural insulation. The annual temperature difference in the recording room is about 8°C, and the daily temperature difference is less than 0.3°C.

2.1.2 Beijing Geomagnetic Observatory (IAGA code BJI)

The Beijing Geomagnetic Observatory, started construction in 1952, is located south of Baijiatuan Village in the western suburbs of Beijing (latitude 40.0, longitude 116.2), and was fully completed in 1954. In 1955, the observatory began to install observation instruments and started trial observation. In 1957, the Beijing Geomagnetic Observatory officially undertook the task of observation. The geology foundation of the observatory and surrounding area is composed of diluvial gravel layers, with Ordovician limestone buried more than 54 meters below, and the upper part is covered with layers of sand, gravel, and pebbles. Within the observation area, the magnetic field gradient is less than 1nT per meter. Both the absolute measurement house and the relative recording room are constructed with weak magnetic materials. The measurement house is a wooden and stone structure on the ground floor. The recording room is a semi-underground wooden and stone structure, with an annual temperature difference of 17°C and a daily temperature difference of less than 0.3°C. However, since 1995, with the acceleration of urbanization, the observation environment of the Beijing Geomagnetic Observatory has gradually been affected by surrounding construction projects. After 2012, with the opening of new subway lines, the interference to the observation environment has become increasingly severe.

2.1.3 Guangzhou Observatory (IAGA code GZH)

The Guangzhou Observatory (IAGA code GZH) was started construction in 1954, located 10 kilometers southeast of Guangzhou in Shiliugang (latitude 23.1, longitude 113.3). The observatory was completed in 1955 and equipped with geomagnetic observation instruments the same year. In 1958, the Guangzhou Observatory officially began geomagnetic observations. The geology foundation of the observatory and surrounding area is composed of sandstone, with a homogeneous magnetic field distribution and a magnetic gradient of less than 1nT/meter. Both the absolute measurement house and the relative recording room are ground-level buildings, constructed with weak magnetic materials such as limestone and wood. The daily temperature variation in the recording room is less than 0.2°C, and the annual temperature variation is about 17°C.

However, due to the construction of the Guangzhou rail transit and the South China Expressway, it was inevitable that interference would occur with geomagnetic observations. Therefore, the Guangzhou Observatory decided to relocate. The new site was chosen in Shendang Village, Liantang Town, Gaoyao District, Zhaoqing City, approximately 120 kilometers southwest of Guangzhou (latitude 23.0, longitude 112.5), and the site selection and positioning work was completed between 1996 and 1997. Construction of the new observatory began in November 2000 and was completed in December 2001. The old observatory ceased observations by the end of 1995. Thus, there were no observation data available from 1996 to 2001. The bedrock at the site is composed of Carboniferous limestone. The magnetic field around the observatory is homogeneous, with a magnetic field gradient of less than 1nT per meter. Both the absolute measurement house and the relative recording room are

constructed with weak magnetic reinforced concrete structures. The absolute measurement house is a ground-level building, while the recording room is a fully underground structure. The annual temperature difference in the recording room is less than 4°C, and the daily temperature difference is less than 0.1°C.

2.1.4 Wuhan Geomagnetic Observatory (IAGA code WHN)

The Wuhan Geomagnetic Observatory (IAGA code WHN) is located in the southeastern suburbs of Wuhan city, Hubei Province, in the town of Baozixie, Wuchang County (latitude 30.5, longitude 114.6). The construction of the observatory began in the spring of 1957 and was completed by December of the same year. In 1958, the instruments were installed and calibrated, and in 1959, the observatory officially began its observations. The observatory is built on a hill with a thick overburden, with a surface layer of yellow clay, where the distribution of the magnetic field gradient is relatively homogeneous, with a gradient of less than 1nT per meter. Both the absolute measurement house and the recording room are constructed with weak magnetic materials as ground-level buildings. The annual temperature difference in the recording room is about 28°C, and the daily temperature difference is generally less than 0.5°C.

Later, affected by the rapid development of township enterprises, it was relocated in 1997 to Jiufeng Village within the territory of Baozixia Town (114.5°E, 30.5°N). In 1998, both Jiufeng site and Baoxie site conducted one-year geomagnetic observation work simultaneously. In 1999, the new site (Jiufeng site) officially began its observational. The site differences between the new and old observatories are D: 0.2', H: -13 nT, Z: -55 nT (Institute of Geophysics, China Earthquake Administration, 2005)The geology foundation of Jiufeng site, and surrounding area is is mainly composed of Upper Triassic quartz sandstone, with thin layers of marl interbedded within. The magnetic field around the observatory is homogeneous, with a magnetic gradient less than 1nT per meter. The absolute measurement house and recording room are both constructed with weak-magnetic materials;the measurement house is a ground structure made of wood and stone, while the recording room is a ground structure made of copper, stone, and plastic, covered with a three-meter-thick soil layer. The indoor daily temperature difference is less than 0.1°C, and the annual temperature difference is less than 9.0°C.

Due to the influence of the subway, it was relocated to Tangchi Town, Yingcheng City, Hubei Province (113.3°E, 30.9°N) in 2017. Due to the severe impact on the observation environment at Jiufeng site, parallel observations were not conducted at Jiufeng site and Yingcheng Site. The geology foundation of Yingcheng Site, and surrounding area is made of limestone with a maximum thickness of 7.4 meters. The magnetic field gradient is less than 1 nT per meter. Both the absolute measurement house and the relative recording room are ground buildings made of weak-magnetic wooden and stone structures. The daily temperature difference in the recording room is about 0.1°C, and the annual temperature difference is about 15°C.

2.1.5 Sheshan magnetic observatory (IAGA code SSH)

The Sheshan Geomagnetic Observatory (IAGA code SSH) and its predecessors has the longest continuous data series. The construction that started in 1874 was in Xujiahu, in Shanghai (1874-1908). Due to the construction of tram lines passing through Xujiahu in 1907, which inevitably interfered with geomagnetic recordings, the observatory was relocated to

205 Lukjapu, Kunshan (1908—1933) in 1908. Later, because of the construction of the Beijing-Shanghai Railway, it was moved to its current location in Songjiang District, Sheshan, Shanghai (since 1932). The SSH Geomagnetic Observatory is located about 20 kilometers southwest of Shanghai in the Xisheshan area of Songjiang District (31.1°N, 121.2°E). With nearly 150 years of history, it is one of the longest-operating geomagnetic observatories in China and the world. The geology foundation of the observatory and surrounding area is mainly composed of medium to acidic andesite from the Upper Jurassic to Lower Cretaceous periods. The magnetic field gradient is 2-3 nT per meter. Both the absolute measurement house and the relative recording room are constructed using materials with weak magnetic materials. The daily temperature difference in the recording room is less than 0.2°C, and the annual temperature difference is about 15°C.

210 **2.1.6 Lhasa Observatory (IAGA code LSA)**

The Lhasa Observatory (IAGA code LSA) is located on the west outskirts of Lhasa city (29.6 N, 91.0 E). Construction of the observatory began in 1956, it was completed in 1957, and started formal observations in July of the same year (Zhou, 2013). Due to instrument failure, the observatory suspended its observation work from 1974 to 1982, and resumed in 1983. The Lhasa Observatory was the first highland geophysical observatory established in the world and was, at the time, the highest altitude geomagnetic observatory globally, with an elevation of 3,655 meters. It is located in the central part of the alluvial basin of the Lhasa River Valley in the suburbs of the city. The magnetic field gradient at the observation site is about 1nT per meter, and is relatively homogeneous. Both the absolute measurement house and the recording room are non-magnetic ground buildings made of limestone, earth bricks, and wood structures. The daily temperature difference inside the recording room is less than 0.2°C, and the annual temperature difference is about 19°C.

删除[susu]: resuming

删除[susu]: The

220 **2.1.7 Lanzhou Observatory (IAGA code LZH)**

The Lanzhou Geomagnetic Observatory (IAGA code LZH) is located at Liujiaping, north of the Yellow River in Lanzhou City (36.1 N, 103.9 E). Construction began in 1956, was completed in 1957, geomagnetic instruments were installed in 1958, and trial observation were started. It officially commenced observation on January 1, 1959. The observatory is situated at Liujiaping, north of the Yellow River in Lanzhou City. The geology foundation of the observatory and surrounding area is mainly Quaternary loess layers. The ground magnetic field gradient is about 1nT per meter, and is relatively homogeneous. Both the absolute measurement house and the relative recording room use non-magnetic or weakly magnetic stones and earth bricks, as well as wooden structures for construction. The absolute measurement house is a ground-level building, while the recording room is an underground structure with a daily temperature difference of 0.3°C and an annual temperature variation of no more than 10°C.

230 **2.1.8 Urumqi Observatory (IAGA code WMQ)**

The Urumqi Observatory (IAGA code WMQ) was established in 1964 and officially began its observation in 1970. However, due to the long-term instability of the instruments, the first geomagnetic observation report was not officially published until

1978. The observatory is located in the eastern suburbs of Urumqi, Xinjiang Uygur Autonomous Region, 10 kilometers from the city center in Shuimogou (43.8 N, 87.7 E). The geology foundation of the observatory and surrounding area is consists of Triassic sandstone and mudstone, with a Quaternary loess layer approximately 4 meters thick covering it. The ground magnetic field gradient is less than 1nT per meter. Both the absolute measurement house and the recording room are ground buildings, constructed using weak magnetic soil bricks and a wooden stone structure. The annual temperature difference in the recording room is less than 8°C, and the daily temperature difference is less than 0.3°C.

Later, due to urban expansion and rapid socioeconomic development, the observation was significantly affected. Therefore, in 2012, the Urumqi Observatory was relocated to Yuanhucun Village, Hutubi County, Changji Hui Autonomous Prefecture, about 80 kilometers northwest of Urumqi City (44.4 N, 86.9 E). The old observatory ceased operations in December 2012, while the new observatory officially began its observation on January 1, 2013. In 2012, both Shuimogou site and Hutubi site conducted one-year geomagnetic observation work simultaneously. The site differences between the new and old observatories are D: -20', H: 420 nT, Z: -413 nT (Institute of Geophysics, China Earthquake Administration, 2012). The geology foundation of Hutubi site is covered with loess layers, and the observation field area has a homogeneous geomagnetic field with a ground magnetic gradient of less than 1nT per meter. The absolute measurement house and the recording room are both constructed with non-magnetic reinforced concrete and stone structures, with the absolute measurement house being a ground-level building and the recording room being a fully underground structure. The annual temperature difference in the recording room is less than 6°C, and the daily temperature difference is less than 0.1°C.

2.2 Instrument Configuration

All eight observatories involved in this study are equipped with the absolute measurement house and the relative recording rooms. The absolute observation room is a ground-level building, while the relative recording room can be either ground-level or underground. They are constructed using weak magnetic materials, providing an ideal environment for geomagnetic observations. Inside the absolute measurement house, the absolute measurement instruments are installed; in the relative recording room, the relative recording instruments are set up. The absolute measurement instruments is capable of measuring the absolute value of the geomagnetic field with high precision, but it cannot operate continuously; the relative recording instruments, on the other hand, can continuously record the changes in the geomagnetic field over long periods, although it cannot measure the absolute value of the geomagnetic field. Therefore, a combination of absolute observations and relative recordings is necessary to obtain long-term continuous absolute values of the geomagnetic field (Jankowsky and Sucksdorf 1996; Rasson 2007). In China, absolute measurement instruments are commonly used to determine the three independent elements of the geomagnetic field: magnetic declination (D), magnetic inclination (I), and total intensity (F). Absolute measurements are scheduled twice a week, usually on Mondays and Thursdays. The relative recording instruments are mainly used to measure the three elements of magnetic declination (D), horizontal component (H), and vertical component (Z).

Before 1975, the absolute observation instruments used on the eight observatories were produced by foreign manufacturers, including Askania, Schmidt, Smith, Cooke, Mating, QHM and other types of magnetometers, which were used to measure the

absolute values of the D, H, and I components. Relative record relied on the Type 57 and Type 72 magnetometers developed independently in China. In 1975, China successfully developed the CJ6 geomagnetic theodolite, after which this instrument was used to absolute measurement of magnetic declination and horizontal intensity. At the same time, they used proton precession magnetometers of the CZM, DTZ, and other types to measure the F total field. Entering the 21st century, the China Earthquake Administration initiated the "Seismic Precursor Station (Network) Technical Renovation" project, vigorously promoting the digitization observation of geomagnetic observatories, equipping them with GM-3 digital fluxgate magnetometers to continuously record changes in the geomagnetic field in the H, Z, and D components. In addition, they were equipped with CTM-DI fluxgate theodolites, combined with G856 proton precession magnetometers, to achieve absolute measurement of D, I, and F. Since 2007, the advancement of the "China Digital Seismological Observation Network" project has further updated and improved the observation instruments at the observatories, using MINGEO fluxgate theodolites and GSM-19T proton precession magnetometers for absolute measurement of D, I, and F, and equipped with GM4 fluxgate magnetometers for continuous relative recording of the D, H, and Z components. At the same time, the GSM-90F1 Overhauser magnetometer was introduced for continuous recording of the F total field.

删除[susu]: process

Table 2 provides a comprehensive listing of all instrument data adopted by eight geomagnetic observatories since 1933. This information includes the instrument models, types, countries and manufacturers of origin, measurement parameters, the year of first deployment at Chinese stations, recording methods, as well as which specific stations have utilized these particular instruments.

Table 2 Instrument data sheet of geomagnetic observatory (Department of Monitoring and Prediction, China Earthquake Administration, 2008)

Serial number	Observation method	Instrument name	Model	Generate country	Manufacturer	Measurement Components	Recording method	Starting year of observation	Observatories
1		Geomagnetic Inductor	Askania	West Germany	Askania Company	I		1957	BJI, WHN, LSA, LZH
2		Geomagnetic Theodolite	Schmidt	West Germany	Askania Company	D, H		1958	BJI, SSH, LSA, WMO,
3		Magnetometer	Smith	United Kingdom	Combridge Company	D, H		1907	GZH, SSH, CNH
4		Geomagnetic Theodolite	Cooke	United Kingdom	Messrs. Cooke, Troughton & Simms, Ltd.	D, H		1958	WHN
5	Absolute observation instrument	Geomagnetic Theodolite	Mating	East Germany	potsdam Mating U. Wiesenberg	D, H		1954	BJI, GZH, CNH
6		Horizontal intensity magnetometer	QHM	Denmark		H		The 1970s	WHN
7		Geomagnetic Theodolite	CJ6	China	Institute of Geophysics, Chinese Academy of Sciences	D, H		The 1980s	WHN, LSA, CNH, GZH
8		Proton Precession Magnetometer	CZM	China	Institute of Geophysics, Chinese Academy of Sciences	F		The 1970s	WHN, SSH, LSA, CNH

9		Proton Procession Magnetometer	DTZ	China	Institute of Geophysics, Chinese Academy of Sciences	F		The 1990s	CNH, BJI, GZH, LSA, WMQ
10		Geomagnetic Theodolite	CTM-DI	China	Institute of Geophysics, Chinese Academy of Sciences	D, I		The 1990s	CNH, BJI, WHN, LSA, LZH, WMQ
11		Proton Procession Magnetometer	G856	United States	EG&G Geometrics Company	F		The 1980s	CNH, WMQ, BJI, LZH, WHN, LSA, GZH
12		Declination/inclination magnetometer	DIM-100	United States	Earth Data Analysis	D, I		The 1990s	SSH
13		Geomagnetic Theodolite	MINGEO DIM	Hungary	MinGeo Ltd	D, I		2006	CNH, BJI, WHN, SSH, LSA, LZH, WMQ
14		Geomagnetic Theodolite	MAG-01H	United Kingdom	Barting Company	D, I		2000	GZH
15		Overhauser Magnetometer	GSM-19T	Canada	GEM	F	digital recording	2011	SSH, GZH, WHN, SSH, LZH, WMQ
16		Overhauser Magnetometer	GSM-90F1	Canada	GEM	F	digital recording	2007	WHN, WMQ
17		Recording Declinometer	Toepfer	Germany	Potsdam Toepfer commercial firm	D	photographic recording	The 1930s	SSH
18		Recording Magnetometer	Type 57	China	Institute of Geophysics, Chinese Academy of Sciences	D, H, Z	photographic recording	1957	GZH, WHN, LSA, LZH, WMQ
19		Recording Magnetometer	Type 72	China	Institute of Geophysics, Chinese Academy of Sciences	D, H, Z	photographic recording	The 1970s	CNH, BJI, WHN, SSH, LSA
20	Relative recording instrument	Fluxgate Magnetometer	GM-3	China	Institute of Geophysics, China Earthquake Administration	D, H, Z	digital recording	2001	BJI, GZH, WHN, LZH, WMQ
21		Fluxgate Magnetometer	GM4	China	Institute of Geophysics, China Earthquake Administration	D, H, Z	digital recording	2007	CNH, BJI, WHN, LSA, LZH, WMQ
22		Fluxgate Magnetometer	FGM-01	China	Institute of Geophysics, China Earthquake Administration	D, H, Z	digital recording	2013	GZH, WHN, LZH, WMQ
23		Fluxgate Magnetometer	FGE	Denmark	Danish Fluxgate Technologies	D, H, Z	digital recording	The early 21st century	SSH, WMQ
24		Vector Proton Procession Magnetometer	FHD-2B	China	Jiangsu Xinyi Jingwei Electronics Co., Ltd.	D, H, Z, F	digital recording	The early 21st century	CNH, BJI, WHN, LSA, LZH, WMQ

2.3 Data collection, organization, digitization, and standardization

290 Given that these valuable historical materials are distributed across different observatories or administrative departments, making it difficult to query and access data resources. Limited by the historical conditions at the time, storage methods vary, with some records on paper carriers, others stored on outdated floppy disks and other storage media, and still others stored on

modern hard drives. Due to the lack of unified standards and specifications, users find it difficult to obtain shared common data sources. Therefore, this study first aims to collect and organize these data from various scattered locations, including geomagnetic observatories and the archives of the Institute of Geophysics, China Earthquake Administration.

The data we collected prior to 1985 primarily originates from the observatory's paper-based annual reports. Continuous records from this period were achieved through photographic methods, and these records have already been published. However, due to the passage of time, we were unable to obtain the digital records from the time of publication. Therefore, We need to digitize the data from published numbers. For this portion of historical materials recorded on paper carriers, we have scanned and generated digital images stored on hard drives for long-term preservation. Since the paper materials have yellowed, and exhibit low legibility, past experience has shown that the accuracy of digital image recognition software often falls short of manual entry (Zhang et al., 2022b). Therefore, we adopted a fully manual input approach to ensure the digitization accuracy of this portion of data.

From 1985 to 2001, data was saved on floppy disks. For data stored on floppy disks, we needed to find compatible reading devices. Since floppy disk drives are rare on modern computers, we had to search everywhere and finally found a compatible drive at a second-hand electronics market, thus achieving the migration of data from the floppy disk to the hard drive.

Data from 2002 to 2006 was stored in Access databases in MDB format, while data from 2007 onwards has been saved in Oracle databases..For data stored on modern hard drives, although technically easier to handle, we also needed to conduct a thorough inspection to ensure that the data was not damaged or lost.

We converted the above collected data into a standardized format, and imported it into the database of Geomagnetic Network of China (GNC) for unified management and use. The data we integrated covers the absolute hourly mean values (AHMV) of the X (northward component), Y (eastward component) and Z components from the following observatories: CNH Observatory (1957-2022), BJI Observatory (1957-2022), LSA Observatory (1957-2022, but no observation data from 1974 to 1982 due to instrument failure), SSH Observatory (1933-2019, no observation from 2020-2022 due to renovation of the observation room), GZH Observatory (1958-2022, no observation data from 1996-2001 due to observatory relocation), WMQ Observatory (1978-2022), LZH Observatory (1959-2022), and WHN Observatory (1959-2022). The data after 2022 was not included because it was incomplete at the time of our analysis.

In order to verify the consistency of data from different sources, we compared the integrated data with that held at the World Data Center (WDC) for Geomagnetism in Edinburgh (<https://wdc.bgs.ac.uk/data.html>), as shown in Figure 2. The pale blue in the figure represents the integrated data, denoted as GNC, while the dark blue represents data from the WDC. The comparison reveals that, except for the data from the WMQ and LSA observatories, which show relatively consistent integrity, the data obtained from the WDC for the other observatories perform poorly in terms of integrity and continuity, with many discontinuous gaps.

删除[susu]: In addition, we also
删除[susu]: unified the storage format,
删除[susu]: all
删除[susu]: the data
删除[susu]: an Oracle
删除[susu]: , long-term preservation,
删除[susu]: collected




Figure 2: Consistency comparison of data from GNC and WDC

3 Quality Inspection and Processing

Quality inspection and preprocessing of observational data are critical steps for conducting data analysis and applications. This study first performs a preliminary examination of the time series data through visual inspection to identify and eliminate obvious errors due to measurement mistakes, data entry omissions, or equipment failures. Subsequently, the tolerance test (Zhang et al., 2022b; Korte et al., 2009) is used to further identify and remove outliers from the data. After completing these steps, the "Multi-scale Wavelet Analysis Step Detection and Signal Correction Algorithm(MWASDSC)" (Zhang et al., 2023) is applied for further detection and processing of the data, primarily aimed at detecting and correcting jump phenomena (Chulliat and Telali, 2007) in the data, which are often caused by replacing absolute instruments, changing standard observation pillars, or relocating observatories. The MWASDSC is as follows: (1) Low-pass filtering: Apply low-pass filtering to the original signal to eliminate high-frequency interference and obtain a smoother low-pass filtered signal; (2) Moving average processing: Perform moving average processing on the low-pass filtered signal based on different numbers of moving average points, resulting in a series of moving average signals. These signals reflect data characteristics at

different scales. (3) Constructing a dataset and wavelet transform: Based on the moving average signals obtained in (2),
340 construct a dataset and choose the largest scale moving average signal to build a ramp function as the wavelet function.
Perform a wavelet transform on it to obtain wavelet transform coefficients. At the same time, remove the largest scale
moving average signal from the dataset; (4) Step jump point detection: Use the position of the maximum wavelet transform
coefficient as the predetermined step jump point, and determine the optimal step jump point position and step height through
a search algorithm; (5) Enclose the area and repeat the search: Enclose the area that spans one ramp width before and after
345 the optimal step jump point, repeat step (4), and continue to search for the optimal step jump point position and step height
from the remaining area. (6) Constructing wavelet functions at different scales: Repeat steps (3) to (5), construct wavelet
functions at different scales in sequence, and use wavelet analysis to pick up step information at different scales, including
step jump point positions and step heights. When repeating step (3), look for the wavelet transform coefficients of the largest
scale moving average signal in the remaining scales moving average signals. (7) Statistics and correction: Statistically
350 analyze all step information found by the wavelet functions at all scales. If three or more scales can identify the same step, it
is preliminarily determined as a step that needs correction. Then, combine with the Kp index to investigate. If the Kp index is
less than 5, it is officially determined as a step that needs correction (Kp index greater than 5 is usually large magnetic
disturbance, which can also produce step effects). Based on the positions of the jump points and step heights of all confirmed
steps, correct the original signal in turn to obtain the corrected signal. (8) Iterative processing: Take the corrected signal in (7)
355 as the original signal for the next round of signal processing, repeat steps (1) to (7), and search for and correct the remaining
steps again.

This study successfully achieved quality inspection and processing of time series data through the aforementioned steps. The
preliminary review and tolerance test effectively eliminated obvious error values and spike anomalies from the data. The
MWASDSC can identify and correct the jump in the data. We estimated all jumps related to the observatory's relocation in
360 this way, which is clearly visible in Figure 3. Figure 3 shows the comparison of the AHMV before and after correction, with
the corrected data showing significant improvement in continuity and reliability.

The analysis of the corrected data shows that each component has a significant year-to-year variation, which over a
considerable period of time manifests as a monotonic increase or decrease. For the X component, there was a general trend
of monotonic increase before 1962, and a monotonic decrease for the subsequent 60 years. The SSH observatory, due to its
365 longer observation, can more clearly exhibit this characteristic of change. The Y-component time series from the WMQ, LZH,
and LSA observatories in the western region show a high degree of consistency in their trends. From the start of the
observation, these three observatories exhibited an overall upward trend in the Y-component, reaching their respective peaks
in 1997, 1995, and 1999, after which they transitioned into a downward trend until the present. At the same time, the other five
observatories have shown a general downward trend since the beginning of observations. Additionally, all eight observatories
370 displayed varying degrees of upward convexity in the Y-component between 1987 and 2008. As for the Z-component, the
CNH, BJI, LZH, and LSA observatories have shown a consistent year-on-year decline since observations began, reaching
minimum values around 1973, followed by an overall upward trend. The SSH observatory showed a slow upward trend before

删除[susu]: 2

1955, began a slow decline afterwards, reached its lowest point in 1973, and has been on an upward trend ever since. The remaining three observatories have roughly exhibited an upward trend since observations began.

375 Furthermore, in the time series curve of the AHMV, it is clear that the AHMV contain a rich variety of external field variation
contributions. Since the external field originates from the ionosphere and magnetosphere current systems outside the Earth, the
complexity and variability of these currents result in a diverse and complexly changing magnetic field morphology. It is due
to this origin that the spatial distribution of the varying magnetic field is extensive, with a global scale and long-distance
spatial correlation. This characteristic is particularly prominent in the X component, for example, during March 13th to 15th,
380 1989, all eight observatories recorded significant geomagnetic storm, and a noticeable decrease exhibited in the X
component, with different observatories exhibiting varying magnitude of change.

删除[susu]: components




Figure 3: Comparison of the AHMV before and after correction at the eight observatories

删除[susu]: 2

4 Validation of corrected data

385 Validating the corrected data is an important means to evaluate data quality. The content of validation includes the accuracy
of diurnal variation records and the stability of long-term changes (Curto and Marsal, 2007).

4.1 Validation of the accuracy of diurnal variation

Typically, the accuracy of daily variations is tested by comparing the consistency of daily variation data from neighboring observatories (Korte et al., 2007; Reda, 2011; Linthe, 2013). Since these eight observatories were among the earliest established in China, there was a lack of data from nearby observatories for comparison in the early days. In this study, we adopted a strategy of cross-validation using data from the eight observatories (Curto and Marsal, 2007). Specifically, We used Geomagnetic Data Processing Software (Zhang et al., 2016) to conduct consistency comparative analysis of the diurnal variation patterns of these eight observatories on daily and monthly timescales.

The comparison results show that the data from the eight observatories, after correction, exhibit a high degree of consistency in their daily variation patterns, with changes sometimes gentle and sometimes intense. Among these, one type of variation is gentle and periodic, cycling with a period of about 24 hours, with the magnetic field changes during the day being significantly greater than those at night. The other type of variation occurs occasionally, is turbulent and irregular, and disappears after a period of time (Xu, 2009). This phenomenon is mainly attributed to their origins from different current systems. The gentle and periodic variations originate from relatively stable current systems in the ionosphere (Pedatella et al., 2011; Yamazaki et al., 2016), while the intense and irregular variations originate from various transient current systems formed by solar particle flows in the magnetosphere and ionosphere (Campbell, 2003). In most cases, regular magnetic field changes are superimposed with some irregular changes. Taking the records from February 1986 as an example, as can be clearly observed from Figure 4, the daily records of each observatory include both regular periodic quiet daily variations and complex disturbed variations, especially during the magnetic storm from February 7th to 9th (As shown between the two vertical dashed lines in Fig. 3), where all observatories' three components showed a consistent intense disturbed response.

删除[susu]: 3

删除[susu]: observatories




Figure 4: Time series curve of AHMV of X, Y and Z components from eight observatories in February 1986

删除[susu]: 3

4.2 Validation of the stability of long-term changes

Comparing the processed data with the prediction data from mature and reliable model can serve as an important method for evaluating the long-term stability of data (Kudin et al., 2021), and can also be used to test the model's adaptability to ground observation data (Finlay et al., 2016). The comparison model selected for this study is the International Geomagnetic Reference Field (IGRF) model. The IGRF model is a mathematical model of the internal geomagnetic field widely recognized by the international academic community, constructed based on candidate models developed by multiple research teams. It is mainly describing describes the core or main field as detectable at Earth's surface. The IGRF model has also been widely accepted and applied in practical applications. The latest version of the IGRF model can be obtained from the website (<http://www.ngdc.noaa.gov/IAGA/vmod/>, last access: 20 May 2025, V-MOD Working Group, 2024) and supports online direct calculation of the seven elements of the geomagnetic field (X, Y, Z, F, D, I and H) and their secular variations at specified coordinates and dates. The update cycle of the IGRF model is every five years, and the coefficients of the 14th generation IGRF model were finally determined by IAGA working group in November 2024.

删除[susu]: the long-wavelength components of the main magnetic field.

To exclude interference from external fields, this study selected mean data from the midnight period from eight observatories to calculate the annual mean (He et al., 2019). The midnight mean data was obtained by calculating the arithmetic average of

the AHMV from 00:00 to 03:00 local time at each observatory. During this period, the observatories are located on the Earth's night side, where the geomagnetic field is relatively stable, effectively avoiding interference from the ionospheric current system and some of the magnetospheric current system. In addition, the midnight period also avoids interference caused by a large amount of human activity. Furthermore, to eliminate the influence of the lithospheric magnetic field, we subtracted their respective initial year's annual mean from the annual mean of each observatory and the IGRF model every year (Figure 5).

删除[susu]: 4




Figure 5: Comparison of annual mean of the observatories with annual mean calculated by the IGRF14 model

删除[susu]: 4

As observed, it is evident that the three components from observatories are highly consistent with the long-term variation trends shown by IGRF model. This indicates that the corrected data can accurately reflect the long-term characteristics of the geomagnetic field in the China. Specifically, both the X-component showed an upward trend before 1962, which then transitioned to a monotonic downward trend; for the Y-component, the WMQ, LZH, and LSA observatories in the western region exhibited a high degree of synchronicity in their trends. Before the late 1990s, the overall trend was upward, followed by a shift to a downward trend until the present. The other five observatories generally showed a downward trend; regarding

the Z-component, the CNH, BJI, LZH, and LSA observatories showed a slight annual decrease before 1973, after which the overall trend became upward; the SSH observatory showed a slow increase before 1955, followed by a decrease until reaching a minimum in 1973, after which it has been on an upward trend. The other three observatories have consistently shown an upward trend. Additionally, for the BJI observatory, the deviation between the observed and model for the X and Y components after 2000, and for the Z component after 2012, has shown an increasing trend. It is speculated that this phenomenon may be due to the increasing interference in the observation environment.

To further quantitatively analyze the differences between the observation and model, this study used formula (1) to calculate the differences between the observatory values and model values obtained above, as well as the standard deviation of the differences. The comparison of the differences is shown in Figure 6, and the statistical results of the standard deviation of the differences are shown in Table 3.

$$\Delta B = B_{OBS} - B_{IGRF}$$

$$S = \sqrt{\frac{\sum_{i=1}^n (\Delta B_i - \overline{\Delta B_i})^2}{n}} \quad (1)$$

In the formula, B_{OBS} represents the annual mean of the observatory, B_{IGRF} represents the annual mean of the IGRF model, ΔB represents the difference between the annual mean of the observatory and the value of the IGRF model, $\overline{\Delta B_i}$ represents the mean of the differences, S represents the standard deviation of the differences, n represents the total number of years, and i represents the i -th year.

删除[susu]: 5

删除[susu]: 2




Figure 6: Difference between observatory values and IGRF model values

删除[susu]: 5

Although the long-term trend of the observatories are consistent with the IGRF model values, it can be seen from Figure 6 that there are certain differences between them. The difference in the Z component of the LSA observatory is the largest, with the maximum difference exceeding 500 nT and the average difference being about 280 nT. Following that, the differences in the X and Z components of the BJI have significantly increased since 2000, with the maximum difference approaching 200 nT, which may be related to the gradual deterioration of the surrounding observation environment (Zhang, 2018). The differences at other observatories are basically all below 100 nT, or even smaller. The error statistics in Table 3 also show that, apart from the slightly larger errors in the X and Z components of LSA and BJI, the root mean square errors of the X, Y, and Z components at the other observatories are all within 50 nT. The preliminary analysis reveals that the reasons of the differences are primarily concentrated in two aspects: First, the remaining contributions from external magnetic fields—although we only selected data for night times, geomagnetic field activity was not taken into account, particularly since geomagnetic storms can still be substantial. On the other hand, due to the uneven distribution of observatories, missing data from these eight observatories, and data quality issues, errors were introduced during the construction of the IGRF model. Additionally, the five-year piecewise linear model adopted in the IGRF model is a simplification method, which has certain limitations in accurately describing continuous secular variations.

删除[susu]: 5

删除[susu]: 2

Table 3 Statistical results of standard deviations for the differences between observatories values and IGRF model values

	CNH	WMQ	BJI	LZH	SSH	WHN	LSA	GZH
X(nT)	24	22	64	22	18	22	59	31
Y(nT)	19	14	17	23	23	12	12	43
Z(nT)	27	13	57	32	30	45	143	34

By analyzing the differences between them, the overall accuracy of the IGRF model in China can also be assessed. Although there are certain differences, considering that the global estimation accuracy range of the IGRF model is between 5 and 300 nT (Lowes, 2000; Wang, 2003; Nie et al., 2017; Lowes, 2022), the model is capable of accurately depicting the spatial distribution of the main magnetic field in China. In scientific research and practical applications of the geomagnetic field, it is crucial to know the accuracy of the model in specific areas.

5 Analysis Application

The application scope of geomagnetic observation data is very extensive, as mentioned in the preface. In this section, we will introduce its applications in the study of Sq variations and geomagnetic jerks.

5.1 Analysis of the Characteristics of Sq Variation in the China

In the calm variations, the solar quiet daily variation (Sq) has attracted attention due to its theoretical significance and practical value. Sq mainly originates from the ionospheric dynamo current system (Vichare et al., 2017) and is the most significant variation component in the mid-low latitude regions during the quiet period of the geomagnetic field. This section will briefly analyze the characteristics of Sq variation in China using the corrected hourly mean data.

The data from the five calmest days of the geomagnetic field in February 1986 were selected, and the average Sq for that month was obtained through time series superposition. The international geomagnetic quiet day data comes from the World Data Center (<https://wdc.kugi.kyoto-u.ac.jp/>). As shown in Figure 7, the horizontal axis represents local time. From the figure, it can be seen that the variations of the three components of Sq at various geomagnetic observatories show the characteristic of large daytime variation and small nighttime variation. This is because the Sq current system has a counterclockwise daytime current vortex in the mid-low latitude part of the Northern Hemisphere (the center position of the maximum current intensity), and the center of this current vortex is located at approximately latitude 30°, while the current intensity at night is very weak (Zhao et al., 2014). It can also be seen from the figure that the X component reaches a significant extreme value before noon, the Y component shows a reverse extreme value around local noon, and the Z component has a characteristic of a minimum value appearing around noon. These characteristics are closely related to the current vortex center of the Sq current system being around local time 12 o'clock. In addition, the variation of Sq in the mid-

删除[susu]: Preliminary analysis suggests that the reasons for the differences mainly involve two aspects: first, factors such as the uneven distribution of observatories, the degree of participation of observation data, data quality, measurement errors, boundary effects, truncation errors, and extension errors, etc., introduce errors of different magnitudes during the IGRF modeling process (Lowes, 2000; Nie et al., 2017). Second, although this study has tried to exclude the interference of external fields and the lithospheric magnetic field as much as possible, the annual mean of the observatories may still contain other components besides the long-wavelength component of the main magnetic field that the IGRF model mainly reflects

删除[susu]: 2

删除[susu]: 6

low latitude regions is mainly influenced by two factors: latitude and local time (Xu, 2009; Feldstein and Zaitzev, 1968), which is also clearly reflected in Figure 7. Sq shows a significant dependence on local time. Comparing observatories with the same latitude but different longitudes, such as CNH (43.8 N, 125.3 E) and WMQ (43.8 N, 87.7 E), it can be seen that their Sq variation patterns are very similar, almost independent of longitude, showing characteristics of smooth changes with local time. The latitude variation characteristics of Sq are mainly manifested as the phase of the X component approximately reverses at latitude 30 degrees; the Y and Z components do not show a reversal phenomenon, because the Y and Z components usually show variations in the equatorial region (Campbell, 2003; Amory-Mazaudier, 2009; Yamazaki and Maute, 2017). In addition, the changes in the X and Z components show approximate symmetry around local noon, while the Y component shows asymmetry. These variation characteristics are consistent with the temporal and spatial variation patterns of Sq in the mid-low latitude regions during the winter solstice month of the Lloyd seasons.

We also analyzed the amplitude characteristics of Sq. The bar chart in the figure shows the daily variation amplitude of the three components at each observatories. The daily variation amplitude of the Y component is the largest at the GZH observatory with the lowest latitude. The daily variation amplitude of the Z component shows that for observatories with similar longitudes, the lower the latitude, the larger the daily variation amplitude. The variation amplitude of the X component does not show a clear regularity.

删除[susu]: 6

510




Figure 7: Sq in February 1986. The upper panel: the Sq variations in the X, Y, and Z components recorded at 8 observatories. The lower panel: the daily variation amplitudes of the X, Y, and Z components at each observatory.

删除[susu]: 6

5.2 Geomagnetic jerk analysis

515 Geomagnetic jerks, which refer to rapid changes in the geomagnetic field occurring over short periods, are characterized by abrupt changes in the secular variations of geomagnetic field elements, showing a "V " or "Λ " shape. The academic community generally agrees that the origin of geomagnetic jerks can be traced back to the Earth's core, and these sudden changes are usually related to the flow changes in the Earth's core (Malin, 1982; Alexandrescu, 1995; Manda, 2010; Chulliat, 2010; Qamili, 2013; Ou et al., 2016). Although geomagnetic jerks have received extensive attention and research in 520 the past forty years (Chulliat, 2010; Brown, 2013; Feng et al., 2019; Kang, 2020; Bai, 2023), there are still many unsolved mysteries in the study of their origin mechanisms and change characteristics. The purpose of this section is to analyze and discuss the geomagnetic jerks using the acquired long-term observation data in this study.

The most direct method for detecting geomagnetic jerks is to identify V-shaped or Λ-shaped changes in the secular variation (SV) of a specific component of the geomagnetic field, after excluding the external field components (Soloviev, 2017). This 525 part of the study still selects annual mean values calculated from AHMV based on the midnight period, effectively avoiding interference from external magnetic fields (variations in the ionosphere and magnetosphere) and human activities. The general view is that the X and Z components are more susceptible to the effects of external fields, while the Y component is least affected by external fields, making the jerk's manifestation most significant on the Y component (Manda et al., 2010; Duka, 2012; Kang et al., 2020). Based on this, this study focuses on analyzing jerks using the Y component.

530 Next, we obtain SV value by calculating the first-order difference of the annual mean of Y component (Manda et al., 2000; Nahayo et al., 2018; Kang et al., 2020). By taking the difference, we can eliminate the constant crustal magnetic field component (Bloxxham and Jackson, 1992; Feng et al., 2018). The SV is calculated as:

$$SV_Y = (Y(year) - Y(year - 1)) / 1 \quad (2)$$

In the formula, SV_Y represents the SV value of the Y component, $Y(year)$ represents the current annual mean, and 535 $Y(year - 1)$ represents the annual mean of the previous year.

In Figure 8, the black dots represent the SV values of the Y component at each observatory. The blank areas in the data are due to the lack of observation data during the corresponding time periods. Among them, the observation environment at the BJI observatory has been severely disturbed since 2012, so the data from 2012 and onwards were not included in the analysis. The GZH observatory was relocated and reconstructed due to environmental interference between 1996 and 2001, resulting 540 in no observation data during that period.

删除[susu]: 7




Figure 8: The SV of the annual Y series from the eight observatories

To quantitatively characterize the Jerk features, this study calculated the SA values before and after each jerk and the occurrence time of the jerk. The estimation of SA values is based on piecewise linear fitting of the time series, using the method proposed by Torta et al. (2015) and Kotzé (2011). The specific steps are as follows: First, to minimize variations related to the solar cycle 11, a 11-year moving average is applied to the first-order differences of the annual means (Alexandrescu, 1997; Korte, 2009). By observing the trend and turning points of the moving average curve, the starting and ending points of each time window are determined, and then piecewise linear fitting is performed. Figure 8 well depicts this process. The intersection time of two adjacent fitted linear segments in the figure represents the moment of jerk occurrence. The slope of the fitted straight line represents the SA value (Mandea et al., 2000; Pinheiro et al., 2011; Brown, 2013; Bai et al., 2023). The intensity of the Jerk can be quantified by its amplitude, which is the difference in SA values before and after each jerk, A , with its fitting formula as.

删除[susu]: 7

删除[susu]: 7

$$\begin{aligned}SV_Y(t_i) &= a_1 t_i + b_1, \quad t_i \leq t_0 \\SV_Y(t_i) &= a_2 t_i + b_2, \quad t_i \geq t_0 \\A &= a_2 - a_1\end{aligned}\tag{3}$$

555 In the formula, $SV_Y(t_i)$ represents the SV value of Y component in t_i year , a_2 indicates the SA value after the jerk, and a_1 is the SA value before the jerk.

As shown in Figure 8, several jerks appeared in the SV of the Y components at every observatory. The most notable events occurred in 1969, 1979, 1991, and 2003 (Mandea and Olsen, 2007), which were clearly reflected in the SV at each observatory. The 2003 jerk occurred distinctly at all observatories, further confirming Tozzi's view that this event was also global (Tozzi et al., 2009). The Jerk in 2007 (Kotzé, 2011) and 2013 (Torta et al., 2015; Chulliat et al., 2015; Finlay et al., 2016) were recorded by five observatories, with three western observatories not recording the event in 2007. The Jerk event in 1999 was only recorded by two observatories, CNH and SSH, and its morphological characteristics were not very clear, which may indeed be a local jerk (Pinheiro et al., 2011). These phenomena further illustrate that some jerk events have global extension characteristics, while some jerks are local events (Chulliat et al., 2010). In 2019, a geomagnetic jerk event was recorded by several observatories. This supports Duan's (2020) prediction of a possible new geomagnetic jerk between 2020 and 2021, which was forecasted by Duan et al. based on the relationship between LOD variations and geomagnetic jerks. This phenomenon deserves further attention.

The 1949 Jerk event (Alexandrescu et al., 1996) is clearly recorded in the Y component of SSH SV, with very clear characteristics of the Jerk. However, since this jerk event occurred relatively early when China only had the SSH observatory, we cannot determine whether this jerk event was prevalent across the entire territory of China or was merely a unique phenomenon confined to that particular observatory. Additionally, events with similar jerk characteristics were recorded by the CNH observatory in 1996, and by the SSH observatory in 1942, 1958, and 1965. Among these, the 1942 event (Mandea et al., 2000; Duka et al., 2012), and the 1958 event were only detected in certain observatories of the world (Lutephy, 2018). These jerks may be real or could be related to the length of geomagnetic records, data quality, etc., especially for data from the early initial of research, which requires more early data for confirmation.

Table 4 shows the occurrence times and amplitude A of each jerk event in the Y component of various geomagnetic observatories. Positive amplitude indicates that the secular variation exhibits a V-shaped feature, while negative amplitude suggests that the secular variation exhibits a Λ-shaped feature. The slashes (/) denote unavailable data, and “ - ” is used where data existed but no jerk was detected during the corresponding time period. The first column of the table lists the widely recognized occurrence times of geomagnetic jerks.

Table 4: The occurrence times and Amplitude of Jerks

Jerk event	Year/Amplitude	CNH	WMQ	BJI	LZH	SSH	WHN	LSA	GZH
1942	Time (year)	/	/	/	/	1942	/	/	/

删除[susu]: 7

删除[susu]: Furthermore, it is surprisingly that the latest jerk event in 2019 was recorded by various observatories, which further confirms the prediction by Duan (2020) that a new geomagnetic rapid change may occur between 2020–2021. Duan et al.'s research predicts, based on the correspondence between LOD changes and geomagnetic jerks, that a new geomagnetic jerk may occur between 2020 and 2021.This phenomenon deserves further attention.

删除[susu]: However, due to its early occurrence, there was only one observatory in China at that time, lacking data from other observatories to corroborate, thus it cannot be determined whether it was a national event or an isolated regional event.

删除[susu]: 3

删除[susu]: 3

	Amplitude (nT)	/	/	/	/	-4.6	/	/	/
1950	Time (year)	/	/	/	/	1950	/	/	/
	Amplitude (nT)	/	/	/	/	7.6	/	/	/
1958	Time (year)	/	/	/	/	1958	/	/	/
	Amplitude (nT)	/	/	/	/	-3.7	/	/	/
1965	Time (year)	/	/	/	/	1965	/	/	/
	Amplitude (nT)	/	/	/	/	2.5	/	/	/
1969	Time (year)	1971	/	1970	1971	1971	1971	1969	1971
	Amplitude (nT)	-3.9	/	-4.3	-4.8	-4.8	-4.7	-4.3	-4.4
1979	Time (year)	1980	/	1978	1979	1978	1979	/	1979
	Amplitude (nT)	2.8	/	3.3	5.7	4.2	5.9	/	6.6
1991	Time (year)	1992	1991	1991	1991	1993	1992	1991	1992
	Amplitude (nT)	-3.6	-6.4	-4.0	-8.8	-3.8	-6.6	-8.3	-7.0
1996	Time (year)	1996	-	-	-	-	-	-	/
	Amplitude (nT)	4.0	-	-	-	-	-	-	/
1999	Time (year)	1999	-	-	-	1999	-	-	/
	Amplitude (nT)	-4.2	-	-	-	-1.6	-	-	/
2003	Time (year)	2003	2003	2003	2004	2004	2003	2004	2004
	Amplitude (nT)	4.0	6.9	3.9	9.8	5.1	8.6	8.1	5.4
2007	Time (year)	2007	-	2007	-	2008	2008	-	2009
	Amplitude (nT)	-2.1	-	-4.0	-	-4.0	-7.3	-	-4.2
2013	Time (year)	2013	2013	-	2013	2013	-	2013	-
	Amplitude (nT)	-4.3	-7.7	-	-12.4	4.7	-	-10.6	-
2019	Time (year)	2019	2019	-	2019	2019	2019	2019	2019
	Amplitude (nT)	13.2	6.9	-	14.1	-13.9	8.4	13.0	12.3

As shown in Table 4, the geomagnetic jerks exhibit significant regional differences and asynchronous occurrence times (Torta et al., 2015). For the same jerk event, there are considerable differences in the amplitude recorded by different observatories. Overall, the jerk amplitudes recorded by observatories in the central and western observatories are greater than those in the eastern observatories, except for the jerk event in 2019.

For different jerk events, the amplitude at the same observatory also varies significantly, ranging from about 2nT yr⁻² to several tens of nT yr⁻². Notably, for jerks with more pronounced morphological features, the absolute value of their amplitude exceeds 4nT yr⁻² (Soloviev, 2017). The occurrence times of the jerks do not show any clear pattern in the eight

删除[susu]: 3

590 observatories studied; the time difference for the same jerk recorded at different observatories can reach up to 2 years. This result is consistent with the conclusions of other scholars, who noted that the time discrepancy of the same event at different observatories can reach 2 to 3 years (Mandea et al., 2010; Brown et al., 2013; Morozova et al., 2014; Soloviev, 2017). The intervals between jerks are often 4 years, 6 years, 8 years, and 12 years, which seems to be correlated with the 6-year and 8.6-year oscillation periods of Earth's rotation diurnal length variations (Ma, 2004; Holme and De Viron, 2013; Lutephy, 2018; Duan and Huang, 2020; Gvishiani and Soloviev, 2020).

600 **6 Data availability**

An integrated and quality-controlled historical datasets of eight pioneering geomagnetic observatories in China are available at: <https://doi.org/10.5281/zenodo.14560950>, 2024 (Zhang et al., 2024b). The data are provided in Txt format, including the observed AHMV files of the three components (X, Y and Z) at the eight observatories, and also the metadata files about the datasets.

7 Conclusion

Geomagnetic observatories play a crucial role in monitoring the spatiotemporal variations of the geomagnetic field, and their accumulated continuous observation data are particularly important for understanding the long-term evolution patterns of the geomagnetic field. Long time-series geomagnetic observation data have wide applications and significant value for scientific research and practical use. The rescue of geomagnetic historical data is a focal point of global scientific concern. This study conducted a salvage organization of the historical observation data from eight geomagnetic observatories (CNH, WMQ, BJI, LZH, SSH, WHN, LSA, and GZH) from the official start of observations to 2022. This included collecting disperse stored observational materials, digitizing paper materials, unified database management, anomaly detection and processing, quality assessment, and application analysis. After the salvage, the historical data is easier to preserve and use over the long term, with significantly improved data quality. These datasets are of great value for improving historical geomagnetic field models (Korte, 2009), studying changing magnetic fields (including diurnal variations and magnetic storm events, etc.), the main geomagnetic field, and its secular variations. However, it should be noted that after 2012, the data from the BJI observatory should be handled with caution due to severe interference in the observation.

In Chapter 5 of this study, we first applied the data to explore the changing characteristics of Sq in China. The analysis results indicate that the main features of Sq include significant changes during the day, weak changes at night, and the changing patterns also show dependence on latitude and local time. In addition, we also focus on the changing characteristics of jerks in the China. The research results revealed several landmark jerks in 1969, 1979, 1991, and 2003 that commonly appeared at eight observatories, thus verifying the global characteristics of these events. In contrast, events in 1999, 2007, and 2013 exhibited local characteristics. For events in 1942 and 1958, since they were only recorded at one observatory, more long-term

620 sequence data are needed for in-depth analysis and confirmation. Encouragingly, the jerk in 2019 was recorded at all observatories. For those jerks with significant morphological features, their amplitude absolute values all exceeded 4nT yr⁻². These findings further confirm the regional differences in geomagnetic jerks and the asynchronicity of the occurrence times of jerks, which is of significant scientific importance for understanding changes in the geomagnetic field.

Although this study reveals the spatiotemporal characteristics of Sq and jerks in China through analysis, given that the data
625 samples used may lack sufficient representativeness and the analytical methods may have certain limitations, these findings may not fully and accurately reflect the actual situation in China. Therefore, subsequent research will collect and integrate more high-quality data from observatories for more detailed and in-depth analysis. At the same time, we will focus on improving the accuracy of the detection algorithm and delve into the physical mechanisms of geomagnetic jerks.

This study aims to provide this valuable dataset to the scientific community and the public, so that these data can be
630 integrated with data from other sources to further explore the spatiotemporal evolution of the geomagnetic field and its physical mechanisms. It also hopes to inspire more scientific workers to engage in the protection of historical materials, with the aim of preserving these precious resources before they are destroyed or lost, and to further realize their scientific value.

Author contributions. SZ conducted data integration, quality inspection,, and analyzed application examples. SZ prepared the manuscript. CF was responsible for identifying and correcting data anomalies, created charts, and revised the manuscript.
635 CC, GZ, JC, QZ, BW, ND, JL, SH, PG and HY is responsible for scanning and digitizing paper materials, as well as standardizing data in various formats. CC designed a set of Excel templates. JW developed the data quality comparison software. JW and CC established database management system

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. We thank CNH, BJI, SSH, WMQ, LZH, LSA,GZH, WHN, and the reference room of the Institute of
640 Geophysics, China Earthquake Administration for providing the valuable data resources. We sincerely thank all the technical personnel and scientists working on the geomagnetic observatories. We also thank World Data Center for Geomagnetism, Kyoto, for providing online International Geomagnetic Quiet Day.

Financial support. This work was funded by the National Key R&D Program of China (grant no. 2023YFC3007404), the National Natural Science of Foundation of China (grant no. 42374092).

- Alexandrescu, M., Gibert, D., Hulot, G., Le Mouél, J.-L., and Saracco, G.: Detection of geomagnetic jerks using wavelet analysis, *J. Geophys. Res.*, 100, 12557–12572, <https://doi.org/10.1029/95JB00314>, 1995.
- Alexandrescu, M., Gibert, D., Hulot, G., Le Mouél, J.-L., and Saracco, G.: Worldwide wavelet analysis of geomagnetic jerks, *J. Geophys. Res.*, 101, 21975–21994, <https://doi.org/10.1029/96JB01648>, 1996.
- 650 Alexandrescu, M., Courtillot, V., and Le Mouél, J.-L.: High-resolution secular variation of the geomagnetic field in western Europe over the last 4 centuries: Comparison and integration of historical data from Paris and London, *J. Geophys. Res.*, 102, 20245–20258, <https://doi.org/10.1029/97JB01423>, 1997.
- Amory-Mazaudier, C.: Electric current systems in the earth's environment. *Niger. J. Space Res.*, 8, 178-255, 2009.
- Bai, C. H., Kang, G. F., Gao, G. M., Wen, L. M., and Sun, Y. P.: Response of geomagnetic jerk features to the geomagnetic secular acceleration in the China region, *Chinese Journal of Geophysics*, 66, 3747-3760, : 10.6038/cjg2022Q0270, 2023 (in Chinese).
- 655 Bloxham, J. and Jackson, A. Time-dependent mapping of the magnetic field at the core-mantle boundary *Journal of Geophysical Research* 97, 19537-19563, :10.1029/92JB01591.issn: 0148-0227, 1992.
- Bolduc, L., Langlois, P., Boteler, D., and Pirjola, R.: A study of geoelectromagnetic disturbances in quebec. II. Detailed analysis of a large event, *IEEE T. Power Deliver.*, 15, 272–278, 2002.
- 660 Boteler, D. H., Pirjola, R. J., and Nevanlinna, H.: The effects of geomagnetic disturbances on electrical systems at the Earth's surface, *Adv. Space Res.*, 22, 17–27, 1998
- Brown, W. J., Mound, J. E., and Livermore, P.W.: Jerks abound: an analysis of geomagnetic observatory data from 1957 to 2008, *Physics of The Earth and Planetary Interiors*, 223, 62–76, doi:10.1016/j.pepi.2013.06.001, 2013.
- 665 Campbell, W. H.: Introduction to Geomagnetic Fields. Cambridge University Press, pp. 62–102. Book DOI: <http://dx.doi.org/10.1017/CBO9781139165136>, 2003.
- Cheng, A. L.: Modernization technology album of geomagnetic stations and networks, *Recent developments in world Seismological*, 6, 1-40., <https://doi.org/10.1002/2015GL064067>, 1995 (in Chinese).
- China Earthquake Administration (CEA), Specification for the construction of seismic station Geomagnetic station, DB/T 9–2004, Beijing: China Earthquake Publishing House, 2004 (in Chinese).
- 670 Chulliat, A. and Telali, K.: World monthly means database project, *Publications of the Institute of Geophysics, Polish Academy of Sciences*, C-99, 268-274, 2007.
- Chulliat, A., Thébaud, E., and Hulot, G.: Core field acceleration pulse as a common cause of the 2003 and 2007 geomagnetic jerks, *Geophysical Research Letters*, 37, L07301, doi: 10.1029/2009GL042019, 2010.
- 675 Chulliat, A., Alken, P., and Maus, S.: Fast equatorial waves propagating at the top of the Earth's core, *Geophys Res Lett*, 42, 3321–3329, <https://doi.org/10.1002/2015GL064067>, 2015.

- Clarke, E., Flower, S., Humphries, T., McIntosh, R., McTaggart, F., McIntyre, B., Owenson, N., Henderson, K., Mann, E., MacKenzie, K., Piper, S., Wilson, L., and Gillanders, R.: The digitization of observatory magnetograms, poster presented at: 11th IAGA Scientific Assembly, Sopron, Hungary, 23–30 August 2009, <https://www.osti.gov/etdeweb/biblio/21389614> (last access: 3 January 2025), 2009.
- Curto, J. J. and Marsal, S.: Quality control of Ebro magnetic observatory using momentary values, *Earth Planets Space*, 59, 1187–1196, 2007.
- Dawson, E., Reay, S., Macmillan, S., Flower, S., and Shanahan, T.: Quality control procedures at the World Data Centre for Geomagnetism (Edinburgh), IAGA 11th Scientific Assembly, Sopron, Hungary, 23–30 August 2009, https://www.researchgate.net/publication/264590163_Quality_control_procedures_at_the_World_Data_Centre_for_Geomagnetism_Edinburgh (last access: 29 December 2024), 2009.
- Dong, X. H., Li, X. J., Zhang, G. Q., Shi, J., and Liu, C.: The study of digital identification of magnetogram, *Seismological and Geomagnetic Observation and Research*, 30, 49–55, 2009 (in Chinese).
- [Department of Monitoring and Prediction, China Earthquake Administration: Electromagnetic observation instruments, in: Comprehensive Collection of Seismic Monitoring Instruments, 191–251, 2008 \(in Chinese\).](#)
- Duan, P. and Huang, C.: Intradecadal variations in length of day and their correspondence with geomagnetic jerks, *Nature Communications*, 11, 2273, DOI:10.1038/s41467-020-16109-8, 2020.
- Duka, B., De Santis, A., Manda, M., Isac, A., and Qamili, E.: Geomagnetic jerks characterization via spectral analysis, *Solid Earth*, 3, 131–148, <https://doi.org/10.5194/se-3-131-2012>, 2012.
- Feldstein, Y. I. and Zaitzev, A. N.: Quiet and disturbed solar-daily variations of magnetic field at high latitudes during the IGY, *Tellus*, 20, 338–366, DOI: 10.3402/tellusa.v20i2.10014, 1968.
- Feng, Y., Holme, R., Cox, G.A., and Jiang, Y.: The geomagnetic jerk of 2003.5-characterisation with regional observatory secular variation data, *Physics of the Earth and Planetary Interiors*, 278, 47–58, DOI:10.1016/J.PEPI.2018.03.005, 2018.
- Feng, Y., Jiang, Y. S., Gu, J. L., Xu, F., Jiang, Y., and Liu S.: Geomagnetic jerk extraction based on the covariance matrix. *Appl. Geophys.* 16, 153–159. <https://doi.org/10.1007/s11770-019-0761-6>, 2019.
- Finlay, C.C., Olsen, N., Kotsiaros, S., Gillet, N., and Tøffner-Clausen, L.: Recent geomagnetic secular variation from Swarm and ground observatories as estimated in the CHAOS-6 geomagnetic field model. *Earth Planet Sp* 68, <https://doi.org/10.1186/s40623-016-0486-1>, 2016.
- Gillet, N., Jault, D., Finlay, C. C., and Olsen, N.: Stochastic modeling of the Earth's magnetic field: Inversion for covariances over the observatory era. *Geochemistry, Geophysics, Geosystems*, 14(4):766–786. DOI:10.1002/ggge.20041, 2013.
- Gvishiani, A., and Soloviev, A.: *Observations, Modeling and Systems Analysis in Geomagnetic Data Interpretation*, ISBN 978-3-030-58969-1 (eBook), <https://doi.org/10.1007/978-3-030-58969-1>, 2020.
- He, Y. F., Zhao, X. D., Zhang, S. Q., Yang, D. M., and Li, Q.: Geomagnetic jerks based on the midnight mean of the geomagnetic field from geomagnetic networks of China, *Acta Seismologica Sinica*, 41, 512–523, <https://doi.org/10.11939/jass.20190009>, 2019 (in Chinese).

Institute of Geophysics, China Earthquake Administration, Geomagnetic Observation Report of Wuhan Geomagnetic Observatory, 2005 (in Chinese).

Institute of Geophysics, China Earthquake Administration, Geomagnetic Observation Report of Urumqi Geomagnetic Observatory, 2012 (in Chinese).

- 715 Holme, R. and De Viron, O.: Characterization and implications of intradecadal variations in length of day, *Nature*, 499, 202–204, <https://doi.org/10.1038/nature12282>, 2013.
- Jankowski, J. and Sucksdorff, S.: Guide for magnetic measurements and observatory practice, The Polish National Space Research Press, Warszawa, Poland, 235pp, ISBN: 0-9650686-2-5, 1996.
- Jonkers, A. R. T., Jackson, A., and Murray, A.: Four centuries of geomagnetic data from historical records, *Rev. Geophys.*, 41, 1006, doi:10.1029/2002RG000115, 2003.
- 720 Kang, G. F., Gao, G. M., Wen, L. M., and Bai, C. H.: The 2014 geomagnetic jerk observed by geomagnetic observatories in China, *Chinese Journal of Geophysics*, 63, 4144–4153, doi: 10.6038/cjg2020N0337, 2020 (in Chinese).
- Korte, M., Manda, M., Olsen, N.: Worldwide observatory hourly mean values 1995 to 2003: an investigation of their quality, General Assembly European Geoscience Union, Vienna, Austria, 2007, EGU2007-A-02799, https://gfzpublic.gfz-potsdam.de/pubman/item/item_235551, 2007.
- 725 Korte, M., Manda, M., and Matzka, J.: A historical declination curve for Munich from different data sources, *Physics of The Earth and Planetary Interiors*, 177, 161–172. DOI:10.1016/j.pepi.2009.08.005, 2009.
- Kotzé, P. B.: Signature of the 2007 geomagnetic jerk at the Hermanus magnetic observatory, *South African Journal of Geology*, 114, 207–210, <https://doi.org/10.2113/gssajg.114.2.207>, 2011.
- 730 Kudin, D.V., Soloviev, A. A., Sidorov, R.V., Starostenko, V. I., Sumaruk, Y.P., and Legostayeva, O. V.: Advanced Production of Quasi-Definitive Magnetic Observatory Data of the INTERMAGNET Standard. *Geomagn. Aeron.* 61, 54–67, <https://doi.org/10.1134/S0016793221010096>, 2021.
- Linthe, H. J., Reda, J., Isac, A., Matzka, J., and Turbitt, C.: Observatory data quality control-the instrument to ensure valuable research, in: *Proceedings of the XVth IAGA Workshop on Geomagnetic Observatory Instruments, Data Acquisition and Processing: extended abstract volume*, edited by: Hejda, P., San Fernando, Spain, March 2013, 173–177, <https://core.ac.uk/download/pdf/20319986.pdf> (last access: 18 November 2022), 2013.
- 735 Liu, C. M., Liu, L. G., and Pirjola, R.: Geomagnetically induced currents in the high voltage power grid in China, *IEEE T. Power Deliver.*, 24, 2368–2374, 2009(in Chinese).
- Liu, L. G., Liu, C. M., Zhang, B., Wang, Z. Z., Xiao, X. N., and Han, L. Z.: Strong magnetic storm's influence on China's Guangdong power grid, *Chinese Journal of Geophysics*, 51, 976–981, <https://doi.org/10.3321/j.issn:0001-5733.2008.04.004>, 2008 (in Chinese).
- 740 Liu, L. G., Ge, X. N., Wang, K. R., Zong, W., and Liu C. M.: Observation studies of encroachment by geomagnetic storms on high-speed railways and oil-and-gas pipelines in China, *Sci. Sin. Tech.*, 46, 268–275, <https://doi.org/10.1360/N092015-00279>, 2016 (in Chinese).

- 745 Lowes, F. J.: IGRF Health Warning, Errors, and Limitations, <https://www.ncei.noaa.gov/products/international-geomagnetic-reference-field/health-warning>, 2022. (last access: 29 December 2024).
- Lowes, F.: An estimate of the errors of the IGRF/DGRF fields 1945–2000, *Earth Planets Space*, 52, 1207–1211, 2000.
- Lutephy, M.: Interplanetary external driven quasidynamo as the origin of geomagnetic jerks correlated with length of day and gravity anomaly, *Contributions to Geophysics and Geodesy*, 48, 23–74, <https://doi.org/10.2478/congeo-2018-000>,
750 2018.
- Ma, L. H., Han, Y. B., and Yin, Z. Q.: Progress on variable earth rotation rate and geophysical phenomena, *Progress In Geophysics*, 19, 968–974, DOI:10.3969/j.issn.1004-2903.2004.04.043, 2004 (in Chinese).
- Malin, S.R.C. and Hodder, B.M.: Was the 1970 geomagnetic jerk of internal or external origin? *Nature* 296, 726–728, 1982.
- Mandea, M., Bellanger, E., Le, Mouél, J.L.: A geomagnetic jerk for the end of the 20th century? *Earth and Planetary Science Letters*, 183, 369–373, 2000.
755
- Mandea, M. and Olsen, N.: Investigation of a secular variation impulse using satellite data: The 2003 geomagnetic jerk, *Earth Planet. Sc. Lett.*, 255, 94–105, 2007.
- Mandea, M., Holme, R., Pais, A., Pinheiro, K., Jackson, A., and Verbanac, G.: Geomagnetic jerks: rapid core field variations and core dynamics, *Space Science Reviews*, 155, 147–175, 2010.
- 760 Marshall, R. A., Smith, E. A., Francis, M. J., Waters, C. L., and Sciffer, M. D.: A preliminary risk assessment of the Australian region power network to space weather, *Space Weather*, 9, S10004, :10.1029/2011SW000685, 2011.
- Morozova, A. L., Ribeiro, P., and Pais, M. A.: Correction of artificial jumps in the historical geomagnetic measurements of Coimbra Observatory, Portugal, *Ann. Geophys.*, 32, 19–40, 2014.
- Morozova, A. L., Ribeiro, P., and Pais, M. A.: Homogenization of the historical series from the Coimbra Magnetic
765 Observatory, Portugal, *Earth Syst. Sci. Data*, 13, 809–825, <https://doi.org/10.5194/essd-13-809-2021>, 2021.
- Nahayo, E., Kotzé, P.B., Korte, M., and Webb, S.: A harmonic spline magnetic main field model for Southern Africa combining ground and satellite data to describe the evolution of the South Atlantic Anomaly in this region between 2005 and 2010. *Earth Planets Space*, 70: 30, <https://doi.org/10.1186/s40623-018-0796-6>, 2018.
- Nie, L. J., Qiu, Y. D., Shen, W. B., Zhang, S. Q., and Zhang, B. B.: Accuracy Evaluation and Applicability of IGRF12 and
770 WMM2015 Model in Chinese Mainland, *Geomatics and Information Science of Wuhan University*, 42, 1229–1235, DOI: 10.13203/j.whugis20160143, 2017 (in Chinese).
- Ou, J. M., Du, A. M., and Xu, W. Y.: Investigation of the SA evolution by using the CHAOS-4 model over 1997–2013, *Sci. China Earth Sci.* 59, 1041–1050, <https://doi.org/10.1007/s11430-016-5265-0>, 2016.
- Pedatella, N. M., Forbes, J. M., Maute, A., Richmond, A. D., Fang, T.-W., Larson, K. M., and Millward, G.: Longitudinal
775 variations in the F region ionosphere and the topside ionosphere-plasmasphere: Observations and model simulations, *Journal of Geophysical Research*, 116, A12309, :10.1029/2011JA016600, 2011.

Pinheiro, K. J., Jackson, A., and Finlay, C. C.: Measurements and uncertainties of the occurrence time of the 1969, 1978, 1991, and 1999 geomagnetic jerks, *Geochemistry, Geophysics, Geosystems*, 12:Q10015, : 10.1029/2011GC003706, 2011.

780 Qamili, E., De Santis, A., Isac, A., Manda, M., Duka, B., and Simonyan, A.: Geomagnetic jerks as chaotic fluctuations of the Earth's magnetic field, *Geochem. Geophys. Geosyst.* 14, 839–850, :10.1029/2012GC004398, 2013.

Rasson, J. J.: Observatories, instrumentation, in: *Encyclopedia of geomagnetism and paleomagnetism*, edited by: Gubbins, D. and Herrero-Bervera, E., Springer, New York, pp 711–713, DOI:10.1007/978-1-4020-4423-6_225, 2007.

785 Reay, S. J., Allen, W., Baillie, O., Bowe, J., Clarke, E., Lesur, V., and Macmillan, S.: Space weather effects on drilling accuracy in the North Sea, *Ann. Geophys.*, 23, 3081–3088, doi:10.5194/angeo-23-3081-2005, 2005.

Reay, S. J., Clarke, E., Dawson, E., and Macmillan, S.: Operations of the World Data Centre for Geomagnetism, Edinburgh, *Data Science Journal*, 12, WDS47–WDS51, 2013.

Reda, J., Fouassier, D., Isac, A., Linthe, H. J., Matzka, J., and Turbitt, C. W.: Improvements in Geomagnetic Observatory Data Quality, in: *Geomagnetic Observations and Models*, vol. 5, edited by: Manda, M. and Korte, M., Springer
790 Science + Business Media, 127–148, eBook ISBN 978-90-481-9858-0, 2011.

Sergeyeva, N., Gvishiani, A., Soloviev, A., Zabarinskaya, L., Krylova, T., Nisilevich, M., and Krasnoperov, R.: Historical *K* index data collection of Soviet magnetic observatories, 1957–1992, *Earth Syst. Sci. Data*, 13, 1987–1999, <https://doi.org/10.5194/essd-13-1987-2021>, 2021.

795 Soloviev, A., Chulliat, A., and Bogoutdinov, S.: Detection of secular acceleration pulses from magnetic observatory data, *Physics Earth Planet Interiors*, 270, 128–142. <https://doi.org/10.1016/j.pepi.2017.07.005>, 2017.

Soloviev, A., Lesur, V., and Kudin, D.: On the feasibility of routine baseline improvement in processing of geomagnetic observatory data, *Earth, Planets and Space*, 70: 16. DOI:10.1186/s40623-018-0786-8, 2018.

[Sun, X. L.: Analysis of long-term variation characteristics of geomagnetic field in Changchun, Journal of Disaster Prevention and Reduction, 29, 44-48, 2013\(in Chinese\).](#)

800 Torta, J. M., Pavón-Carrasco, F. J., Marsal, S., Finlay, C. C.: Evidence for a new geomagnetic jerk in 2014, *Geophys Res Lett*, 42, 7933–7940, <https://doi.org/10.1002/2015GL065501>, 2015.

Tozzi, R., De Michelis, P., and Meloni, A.: Geomagnetic jerks in the polar regions, *Geophysical Research Letters*, 36, L15304, DOI:10.1029/2009GL039359, 2009.

[V-MOD Working Group, IGRF 14th generation, http://www.ngdc.noaa.gov/IAGA/vmod/, last access: 20 May 2025.](http://www.ngdc.noaa.gov/IAGA/vmod/)

805 Vichare, G., Rawat, R., Jadhav, M., and Sinha, A.K.: Seasonal variation of the Sq focus position during 2006–2010, *Adv. Space Res*, 59, 542–556. <https://doi.org/10.1016/j.asr.2016.10.009>, 2017.

Wang, D. W.: Analysis of the International Geomagnetic Reference Field Error in the China Continent, *Chinese Journal Geophysics*, 46, 63-73, 2003(in Chinese).

810 Xu, W. Y.: *Physics of Electromagnetic Phenomena of the Earth*, Hefei: University of Science and Technology of China press, 558 pages, ISBN 978-7312022562, 2009 (in Chinese).

Yamazaki, Y., Kathrin, H., and Anderson, W. J.: Day-to-day variability of midlatitude ionospheric currents due to magnetospheric and lower atmospheric forcing, *J. Geophys. Res.: Space Phys.*, 121, 7067–7086, <https://doi.org/10.1002/2016JA022817>, 2016.

Yamazaki, Y. and Maute, A.: Sq and EEJ—A review on the daily variation of the geomagnetic field caused by ionospheric
815 dynamo currents, *Space Science Reviews*, 206, 299–405, 2017.

Zhang, S, Fu, C, Zhao, X, Zhang, X, He, Y, Li, Q, Chen, J, Wang, J and Zhao, Q. 2024. Strategies in the Quality Assurance of Geomagnetic Observation Data in China. *Data Science Journal*, 23: 9, pp. 1–11. <https://doi.org/10.5334/dsj-2024-009>

Zhang, S. Q., Fu, C. H., He, Y. F., Yang, D. M., Li, Q., Zhao, X. D., and Wang, J. J.: Quality Control of Observation Data by
820 the Geomagnetic Network of China, *Data Science Journal*, 15, p. 15, <https://doi.org/10.5334/dsj-2016-015>, 2016.

Zhang, S. Q., Zhu, G. H., Wang, J. J., Chen, C. H., He, S. P., and Guo, P. K.: Quality-controlled geomagnetic hourly mean values datasets of Sheshan observatory from 1933 to 2019, Zenodo [data set], <https://doi.org/10.5281/zenodo.7005471>, 2022a.

Zhang, S., Fu, C., Wang, J., Zhu, G., Chen, C., He, S., Guo, P., and Chang, G.: Rescue and quality control of historical
825 geomagnetic measurement at Sheshan observatory, China, *Earth Syst. Sci. Data*, 14, 5195–5212, <https://doi.org/10.5194/essd-14-5195-2022>, 2022b.

Zhang, S. Q., Fu, C. H., Wang, J. J., Zhao, Q., Hu, X. J., Zhang, X. X., Guo, P. K., He, S. P., and Chang, G. P.: A method for automatic identification and processing of step-like interference information in geomagnetic data. China: CN117687104 B, 2023,

830 Zhang, S. Q., Fu, C. H., Zhao, X. D., Zhang, X. X., He, Y. F., Li, Q., Chen, J., Wang, J. J., and Zhao, Q.: Strategies in the Quality Assurance of Geomagnetic Observation Data in China, *Data Science Journal*, 23: 9, pp. 1–11. DOI: <https://doi.org/10.5334/dsj-2024-009>, 2024a.

[Zhang, S., Zhu, G., Chen, J., Deng, . na ., Lu, . jinghui ., Yu, . hongchi ., Chen, . chuahua ., & Wang, . bin . \(2025\). An integrated and quality-controlled historical datasets of eight pioneering geomagnetic observatories in China \[Data set\]. Zenodo. https://doi.org/10.5281/zenodo.15481895,](https://doi.org/10.5281/zenodo.15481895)

Zhao, X. D., Yang, D. M., He, Y. Fei., Yu, P. Q., Liu, X. C., Zhang, S. Q., Luo, K. Q., Hu, X. J.: The study of Sq equivalent current during the solar cycle, *Chinese J. Geophys*, 57, 3777–3788, DOI: 10.6038/cjg20141131, 2014 (in Chinese).

Zhang, X. L., Chang, G. P., and Zhang, M. D.: Interferences on the geomagnetism observation system in Beijing National Earth Observatory, *Seismological and Geomagnetic Observation and Research*, 39, 108–113. DOI: 10.3969/j.issn.1003-
840 3246.2018.06.016, 2018.

Zhao, X. K., Wu, B. Y., and N., B. Q.: The geomagnetic dataset of Beijing Ming Tombs station (1991–2001), *China Scientific Data*, 2, 1–9, <https://www.sciengine.com/csD/doi/10.11922/csdata.170.2016.0074> (last access: 29 December 2024), 2017 (in Chinese).

删除[susu]: Zhang, S. Q., Zhu, G. H., Chen, J., Deng, N., Lu, J. H., Yu, H. C., Chen, C. H., and Wang, B.: An integrated and quality-controlled historical datasets of eight pioneering geomagnetic observatories in China [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.14560950>, 2024b

Zhou, J. P.: Reminiscences of the Lhasa Geomagnetic Observatory, Recent developments in world Seismological, 1, 34-40,
845 DOI:10.3969/j.issn.0235-4975.2013.01.009, 2013 (in Chinese).