

BEACH: Barbados and Eastern Atlantic Combined High-altitude dropsonde datasets

Helene M. Gloeckner^{1,2}, Theresa Mieslinger¹, Nina Robbins-Blanch^{2,3}, Geet George⁴, Lukas Kluft¹, Tobias Kölling¹, Sandrine Bony⁵, Julia Windmiller¹, and Bjorn Stevens¹

¹Max Planck Institute for Meteorology, Bundesstraße 53, 20146 Hamburg, Germany

²International Max Planck Research School on Earth System Modelling, Hamburg, Germany

³Meteorologisches Institut, Universität Hamburg, Bundesstraße 55, 20146 Hamburg, Germany

⁴Faculty of Civil Engineering and Geosciences, TU Delft, Stevinweg 1, 2628 CN Delft, Netherlands

⁵LMD / IPSL, CNRS, Sorbonne University, 4 place Jussieu, Paris F75252 Cedex 05, France

Correspondence: Helene M. Gloeckner (helene.gloeckner@mpimet.mpg.de)

Abstract. As part of the ORCESTRAs field campaign in August and September 2024, 1191 dropsondes were deployed over the Eastern and Western Atlantic ITCZ from the HALO aircraft coordinated by the PERCUSION and MAESTRO subcampaigns. Here, we describe the hierarchy and processing of the resulting Barbados and Eastern Atlantic Combined High-altitude (BEACH) dropsonde datasets. The Level 0 dataset contains measured meteorological variables, such as relative humidity (RH),
5 temperature (T), pressure (p), eastward (u), and northward (v) wind data as output by the AVAPS system. The corresponding ASPEN quality-controlled data is called Level 1. Level 2 adds further measurement-specific quality control flags. Level 3 builds the core of BEACH including all quality controlled dropsonde profiles interpolated to a common 10 m altitude grid and concatenated into a single dataset. We further derive mesoscale vorticity, divergence, and vertical velocities from 87 circular flight patterns in Level 4 using the regression method. These area-averaged variables will guide our understanding of mesoscale
10 processes acting within the ITCZ, one of the main goals of ORCESTRAs. All data levels are openly available on IPFS, while the processing code is made public on GitHub.

1 Introduction

The Organized Convection and EarthCARE Studies over the Tropical Atlantic (ORCESTRAs, Stevens et al., 2026) field campaign was designed to quantify drivers of mesoscale convective organisation in the tropics with a particular focus on the
15 structure and variability of the Atlantic Inter-tropical Convergence Zone (ITCZ). More than 1000 dropsondes were launched as part of PERCUSION¹, and in support of the MAESTRO campaign, two of the sub-campaigns of ORCESTRAs. The soundings were conducted in August and September, 2024, and later processed to derive area-averaged estimates of horizontal divergence and vertical velocity on the mesoscale (~ 200 km, 1h; close to the meso- β scale as per Orlanski (1975)). The datasets described here provide the first comprehensive mesoscale vertical velocity estimates derived from airborne dropsonde measurements
20 within the Atlantic ITCZ.

¹campaign and platform specific acronyms are specified in the appendix

Reliable profiles of the area averaged vertical wind ~~speed~~velocity, $W(z)$ are crucial to determine the magnitude and sign of vertical moist static energy advection in the tropics, which in turn helps to understand the interactions of cumulus convection with large scale circulations (Back and Bretherton, 2006), and patterns of tropical rainfall (Bernardez and Back, 2024). Directly measuring W , however, remains challenging due to its small magnitude compared to the horizontal wind components. While reanalysis data provides estimate of W , without independent measurements it is hard to know how well these estimates are constrained by data, even in cases when additional data from field campaigns are assimilated (Huaman et al., 2022).

Efforts to derive W ~~more~~-directly from observations have a long history. Already eighty years ago Panofsky (1946) proposed integrating area-averaged divergence of the horizontal wind velocity (\mathcal{D}) upward from the surface to compute W . An approach Bellamy (1949) developed further to graphically acquire divergence from three dislocated measurements which defined a triangle. From Gauss' theorem, the area averaged divergence is equal to the line integral of the normal wind around the perimeter of a polygon, whose vertices can be defined by point measurements from sondes. Ceselski and Sapp (1975) adopted this approach to derive \mathcal{D} from operational soundings over the Northern American continent. Yanai (1961) applied these methods to sounding measurements to provide the first estimates of W in the tropics. The utility of this approach was demonstrated in subsequent analyses of the tropical atmosphere (Reed and Recker, 1971; Yanai et al., 1973), during GATE, and in a great many field studies thereafter, as sounding arrays increasingly became incorporated in the design of field campaigns.

Panofsky and Bellamy's ideas were recast by Lenschow et al. (1999), who applied them to aircraft data. Lenschow and collaborators used airborne gust-probe measurements of the horizontal wind to estimate \mathcal{D} at the top of the boundary layer from straight and level legs arranged to close a polygon. They argued that circular flight patterns would be preferable, as they not only minimize the perimeter to area, but also avoid sharp turns required to transition between polygon edges, during which measurements are not useful. Lenschow et al. (2007) demonstrated the circle method to calculate \mathcal{D} , and showed that \mathcal{D} could equivalently be computed from spatial derivatives estimated from best fit linear-regression of the measured wind field to spatial distance.

Bony and Stevens (2019) combined the past approaches by using an aircraft to deploy dropsondes to construct a sounding array. By flying multiple circles with a diameter of approximately 200km following the mean wind they could provide independent estimates, and hence quantify the error. This allowed them to demonstrate that about twelve sondes were sufficient to derive a reliable mesoscale divergence profile and 6 to 8 sondes are tolerable to evaluate the structure of the calculated vertical velocity profile. Bony and Stevens (2019) also demonstrated that their measurements were amenable to the regression method. Using the ICON model to perform large eddy simulation for the observed conditions, they further demonstrated that the temporal decorrelation of the divergence is given by the advective timescale, and hence much less than the time required to fly a single circle.

As compared to the use of winds measured just at flight level, the use of dropsondes has the advantage of sounding arrays, in that they provide vertical profiles of \mathcal{D} , and hence W . Hence these methods were incorporated into the experimental design of EUREC⁴A (Bony et al., 2017) and other campaigns (Pincus et al., 2021) in the winter trades, as well as for HALO-(AC)³ in the Arctic (Wendisch et al., 2024). During the 2019 OTREC field campaign (Fuchs-Stone et al., 2020; López Carrillo and Raymond, 2011), the regression methods were generalized to a variational approach by which \mathcal{D} and W were estimated from

dropsonde data (Vömel et al., 2020) spread over a large area augmented by winds estimated from airborne doppler radar measurement following Mapes and Houze Jr (1995).

Methods to estimate W continue to evolve. Pujol and Bony (2024), for instance, have developed qualitatively new approaches to estimating W , by tracking humidity gradients in time using satellite measured radiances. Their method, however, requires an area devoid of cloud, making its application within the ITCZ problematic. Hence to expand our understanding of convective regimes, PERCUSION incorporated circular flight patterns to drop sondes in and around the Atlantic ITCZ and thereby quantify W . These measurements resulted in the Barbados and Eastern Atlantic Combined High-altitude (BEACH) datasets described in this paper. A name chosen in part because it extends the musical themes of other named elements within ORCESTRA through reference to Amy Beach, the first female US American composer to publish a symphony.

In what follows we present BEACH and the choices made in its construction. Section 2 outlines the dropsonde measurements during PERCUSION. Section 3 describes the methodological and technical details for the data processing, which is adapted from the JOANNE processing described in George et al. (2021). Section 4 gives a brief overview of the thermodynamic and dynamic structure of the tropical atmosphere as measured by the BEACH dropsondes.

2 Measurements

During PERCUSION 1191 sondes were dropped from the German research aircraft HALO. Data of 715 sondes, launched between August 27 and September 28, was assimilated into the IFS analysis. After quality control and other processing steps described in Section 3, 1115 sondes were used in the BEACH Level 3 gridded product. Most of these were grouped in 89 circles to form the BEACH Level-4 product.

Detailed sonde statistics for each flight are provided in Table 1, and the circles that were flown in coordination with the MAESTRO subcampaign are listed in Table B1. Flight tracks were designed with two major objectives: (1) to fly along the EarthCARE track coincident with an EarthCARE overpass to calibrate the satellite measurements and validate the retrievals, and (2) to provide estimates of the mesoscale vertical motions in and around the ITCZ (see Section 3.4).

The circles were designed to take approximately one hour at 14km altitude to complete, which resulted in a circle diameter of roughly 260km, which varied slightly based on flight altitude and hence speed. Additional circles with a smaller diameter of ≈ 140 km were flown at lower altitudes in approximately 40min to coordinate with MAESTRO measurements by the SAFIRE ATR-42 research aircraft near the Cape Verde island Sal, leading to a larger variation in circle diameter in the East.

A typical flight in the East included four circles: one near the center of the ITCZ, two at the edges and one in coordination with the SAFIRE ATR-42. During flight planning the ITCZ was identified as the region where total column water vapor values exceeded 48mm or where surface wind direction changed (Praturi and Stevens, 2026). However, especially in the West, the ITCZ was often not well defined (Stevens et al., 2026), as regions of elevated water vapor could extend over a wide range of latitudes. Even with a clearly defined ITCZ, restrictions from air-traffic control sometimes did not allow the orientation of the circles along the EarthCARE orbit and across the ITCZ. As a result, in the West the flight plans focused on distributing circles




Figure 1. Location of dropsonde launches. Sondes that were launched in ATR coordinated circles are marked in yellow, sondes in regular circles in teal, and sondes that were dropped close to the R/V Meteor in navy regardless their affiliation to a circle. Other sondes are marked in light blue. Sondes can be part of multiple groups (i.e. the sondes in coordination with the ATR were usually also part of a circle and hence appear twice in the legend).

within and across the ITCZ, with less regard to the orientation of the circles, except to maintain a similar inter-circle distance as in the East, and remain anchored to EarthCARE’s overpass.

90 Figure 1 shows all sondes with measurements that passed the basic quality control colored depending on whether they belong to a standard one-hour circle (teal), a smaller ATR circle (yellow), or were dropped in coordination with measurements by the research vessel R/V Meteor (navy). In total, 90 circles that were planned to have dropsondes were flown of which 87 have enough sonde measurements to derive vertical motion on the mesoscale (Bony and Stevens, 2019, see Section 3.4). The indicated regions East, North and West are the same used in Stevens et al. (2026) and are used in Section 4 to divide the data.

95 Additional sondes were sometimes dropped at the point of the EarthCARE overpass, and at the Southern or Northern most point of the overpass to validate other instruments, and for instrument calibration on board of HALO. On flight HALO-20240919a² many sondes were dropped along the flight path to provide a basis for testing a variety of sampling strategies. On this flight alone 73 sondes were launched, one on average every 7–8min.

2.1 Instruments and Sensors

100 The dropsondes used during PERCUSION are of the type RD41 developed by NCAR (Hock and Franklin, 1999; Aberson et al., 2023) and manufactured by Vaisala. Each sonde consists of a PTU unit measuring pressure, temperature, and relative humidity (RH) at 2 Hz sampling frequency. A GPS unit provides information on the dropsonde position at 4 Hz, and wind components are derived from the horizontal displacement of the sonde on its way to the surface. The dropsonde system during

²Flight IDs are based on the platform shortname and the ISO-8601 date alphabetically demarcated to allow more than one flight per day

PERCUSION could receive data from 8 sondes simultaneously, but usually no more than 6 sondes were in the air at once. After
105 the drop, different sensors need different equilibration times until the measurements are valid. The Aspen default equilibration
times, that were used in the processing, are listed in Vömel and Goodstein (2020) and more details on single sensors, their
resolution, and performance are given in George et al. (2021) and their Table 1.

The dropsonde sensors are the same as those included in radiosondes of type RS-41 launched from Barbados, and the
~~Meteor-RV~~ METEOR as part of ORCESTRAS. These radiosondes in addition to the radiosondes launched at INMG populate
110 the RAPSODI datasets (Winkler et al., 2026). Common variables and a uniform grid that is shared among RAPSODI's Level
2 and BEACH's Level 3 facilitate a combined analysis, even though the processing described in section 3 differs significantly
between the datasets due to different raw data formats and dataset requirements.

2.2 Problems during operation

After HALO-(AC)³ (Ehrlich et al., 2024), the last HALO campaign with extensive dropsonde operations before PERCUSION,
115 the antenna for the dropsonde receiver on HALO was moved from a central position on the bottom part of the fuselage behind
the wings of the aircraft to the port-side wing. As a consequence, a longer cable and an amplifier were installed to connect the
antenna with the AVAPS system. The connection between sondes in the air and the AVAPS system seemed interrupted during
flight maneuvers with high roll angles possibly due to the shift in antenna position. In addition, the network connection from
the AVAPS system to the dropsonde computer was unstable until 2024-09-14 which led to data loss from some sondes in the
120 air on HALO-20240827a and HALO-20240914a. Overall those problems did not lead to significantly worse quality control
drop outs than experienced during EUREC⁴A (Section 3.2).

In some instances, air traffic control restricted drops during flight operations resulting in circles with fewer sondes. In some of
these instances, parts of the circle could be reflight a second time, or additional sondes could be dropped to cover a wider area
(e.g. see flight reports for HALO-20240907a or HALO-20240926a). In one instance, on flight HALO-20240821a, clearance to
125 drop sondes was revoked during an entire circle, and it therefore remains without sondes.

In three cases, two measurements have the same serial id in the raw data file headers. This can happen if a sonde is initialized
twice without a drop in between, if the power-pin of a sonde is removed and re-plugged within a few milliseconds because this
leads to a factory reset of the sonde, or if two sondes are initialized to send data on the same frequency and the frequency is
changed at a later stage for one of those sondes. In case of a factory reset, the sonde forgets its calibration and the serial id
130 000007500 is assigned to it. To handle the above mentioned specific cases, BEACH uses a hash derived from the serial id and
launch time to uniquely identify each sonde.

3 Data Processing and Data Products

The datasets in BEACH (see Figure 2) are organized in five levels: Level 0 contains the raw data as recorded by the AVAPS
system; Level 1 contains ASPEN-processed netCDF files; Level 2 includes further customized quality controlled Level 1

Table 1. PERCUSION dropsonde statistics showing the number of sondes per flight and processing level.

flight ID	date	flight time (UTC)	Level 0	Level 1	Level 2	Level 3	Level 4	circles
HALO-20240809b	2024-08-09	09:41:13-15:57:28	3	3	3	3	0	0
HALO-20240811a	2024-08-11	11:59:34-20:35:57	54	49	49	48	46	4
HALO-20240813a	2024-08-13	14:15:39-23:18:05	50	49	49	49	48	4
HALO-20240816a	2024-08-16	11:35:40-20:03:22	52	49	49	49	48	4
HALO-20240818a	2024-08-18	10:04:39-19:03:57	40	38	38	38	38	3
HALO-20240821a	2024-08-21	12:23:35-19:52:41	42	38	36	36	36	3
HALO-20240822a	2024-08-22	11:23:08-19:40:12	55	53	53	53	52	5
HALO-20240825a	2024-08-25	09:14:49-18:58:42	52	47	47	47	47	4
HALO-20240827a	2024-08-27	09:59:43-19:08:18	55	51	49	44	42	4
HALO-20240829a	2024-08-29	12:20:37-20:30:09	52	51	51	51	49	4
HALO-20240831a	2024-08-31	08:51:01-17:41:37	51	50	50	50	49	4
HALO-20240903a	2024-09-03	11:32:07-20:24:25	50	46	46	46	46	4
HALO-20240906a	2024-09-06	10:36:20-17:56:59	12	12	12	12	12	1
HALO-20240907a	2024-09-07	12:49:52-20:40:32	41	41	41	41	41	3
HALO-20240909a	2024-09-09	11:40:40-20:46:33	36	36	36	36	36	3
HALO-20240912a	2024-09-12	11:29:50-20:05:06	51	45	45	45	45	4
HALO-20240914a	2024-09-14	11:29:18-20:04:07	54	45	45	45	45	4
HALO-20240916a	2024-09-16	11:36:59-20:56:44	57	55	55	55	39	3
HALO-20240919a	2024-09-19	11:01:16-19:56:21	76	73	73	73	68	5
HALO-20240921a	2024-09-21	11:22:33-20:07:37	64	61	61	61	61	5
HALO-20240923a	2024-09-23	11:13:40-20:06:12	62	59	59	59	58	5
HALO-20240924a	2024-09-24	15:37:24-21:46:15	49	48	48	48	30	3
HALO-20240926a	2024-09-26	11:42:36-20:23:42	64	60	60	60	58	5
HALO-20240928a	2024-09-28	10:47:02-20:02:36	69	66	66	66	64	5
Total			1191	1125	1121	1115	1058	89

135 data in zarr format; Level 3 consists of the Level 2 data interpolated onto a common altitude grid, as well as additional derived physical variables; Level 4 associates Level 3 data with circles and provides circle products, e.g., the mesoscale vertical motion.

All processing steps generating the BEACH datasets are openly available and embedded in the Python package `pydropsonde`. The BEACH datasets were created with `pydropsonde` version 0.5.5 which evolved out of the processing done for the JOANNE dataset (George et al., 2021). The basic structure of the data levels remains, while some parts of the processing
140 have been improved and expanded, as will be described in this section.




Figure 2. Schematic overview of the data processing from the AVAPS raw data to the Level 4 circle products.

3.1 Level 1 processing : ASPEN quality control

The Level 0 (Gloeckner et al., 2026a) or raw data generated by the AVAPS system is described by George et al. (2021, chapter 2.3.1 and Table 4). pydropsonde uses the raw data files “D-files” and the metadata files “A-files”. As a first step, the metadata of each sonde is checked for a detected launch, which occurs if the sonde ~~parachute opens properly~~ launch detect pin has been
 145 activated. The respective ”A-file” includes a line stating ”Launch Obs Done?” and possible flag values ”0” - *False* and ”1” - *True*. In case ~~if the parachute~~ does not open, or the opening is not detected, the sonde does not switch to a high power mode for transmitting data and the connection to the receiving unit is lost after falling a few hundred meters. Such profiles are of little value and discarded from any further analysis.

ASPEN is a software package developed by NCAR that is used for analysis and quality control (QC) of dropsonde data. For
 150 the BEACH processing, ASPEN version 4.0.4 was used. For each sonde that detected its launch, the ASPEN software (Martin and Suhr, 2021) is run on the raw data (D-file) using a container-based approach. A docker image containing the command-line functionality of ASPEN is utilized within the processing pipeline with the default editsonde configuration. The ASPEN processing includes several quality control steps, such as removal of the equilibration period, smoothing, and outlier checks as described in the ASPEN Manual (AspenDocs 1.0) and Dropsonde Data Quality Report (Vömel and Goodstein, 2020, based on
 155 the NRD41 sondes, which are built differently but contain the same sensors).

We encountered several special cases due to connection or manufacturing issues that needed individual treatment: Eleven sondes were missing metadata information due to an empty A-file. Since the processing with ASPEN is independent of the A-files, it was applied regardless. A flag stating a successful launch-detect is set to ‘None’ within the pydropsonde processing




Figure 3. Normalized sonde counts for the quality measures. Grey lines denote the thresholds for a sonde to pass a QC (i.e. a sonde is flagged if the fraction of missing values exceeds 0.2, if the number of near-surface measurements is lower than 50, or if the profile does not extent above 8000m). QC values for v are equal to those from u . QC values for p are indistinguishable from QC values for ta for the purpose of the plot and omitted for readability.

in those cases signaling that the status of launch-detect is unknown. Since all of those sondes have other problems as well, neither of them appears in Level 3 (see Table E1). Five of those sondes have a Level 2 file, but should be handled with care since their altitude coordinates are unreliable. They can be identified by a NaT `launch_time`. In addition, based on the metadata, 17 sondes were falsely configured by the manufacturer to be of type NRD41, often called *minisonde*, instead of the actual sonde type RD41. They have been processed by ASPEN with the respective *minisonde* configuration, because ASPEN does not allow a processing with the [default](#) RD41 configuration for those sondes. While this does not influence the variables used for the BEACH datasets of Level 2-3 and above, it does impact the estimated vertical wind component included in the Level 1 and 2 datasets of those sondes. The affected sondes are listed in supplement Table E2 [so that they can be manually reprocessed if needed](#).

In total, 1125 sondes reached Level 1 (Gloeckner et al., 2026b) with individual numbers per flight listed in Table 1. We call the untouched ASPEN netCDF output files Level 1 data and store it in form of single datasets per sonde.

3.2 Level 2 processing: Additional quality control

3.2.1 Quality Control Tests

The Level 2 (Gloeckner et al., 2026c) processing applies additional quality control (QC) tests to provide a basis for a combined analysis of all profiles. It includes four variable-specific tests that we call *profile-sparsity*, *profile-extent*, *near-surface-coverage*, and *sfc-physics*. Before those tests are run, we remove data above the drop height (*gpsalt-below-aircraft*). The tests are based on the JOANNE processing (George et al., 2021) with slight modifications and additions, described in section 3.5.

Filter: gpsalt-below-aircraft The ASPEN Level 1 data contains the altitude variable `gpsalt` obtained from GPS measurements. Depending on the GPS connectivity within the aircraft, the GPS measurements need an equilibration period of up to 10 seconds (Vömel and Goodstein, 2020) to build up a connection after a sonde launch. The ASPEN-processing removes this period from the u and v data, but not from the GPS-altitude (`gpsalt`) data points, such that some-
180 times erroneous measurements above the aircraft altitude are included before the connection is established. Therefore, `pydropsonde` removes the `gpsalt`, u , v , sonde latitude, and sonde longitude values for any measurement with a `gpsalt` above the aircraft altitude as measured by BAHAMAS (Konow et al., 2021). This is a valid approach because the sondes ~~are too heavy to be carried upwards~~ were not carried upwards in the ORCESTRA measurements. The data is removed before any other QC tests, such that their results are not influenced by a faulty `gpsalt`.

185 **QC: gps-valid** A second QC checks whether the u and v measurements have an unusually large variation. It is failed if

$$\sigma_{uv} := \sqrt{\sigma_{\Delta u}^2 + \sigma_{\Delta v}^2}, \quad (1)$$

exceeds three times the mean σ_{uv} of all sondes: $\sigma_{uv} > 3 \cdot \overline{\sigma_{uv}}$, because large σ_{uv} values indicate that the GPS measurements are faulty, either due to fast falls or large gaps in the GPS measurements.

Profile Fullness This is checked with three variable specific tests: The *profile-extent* is passed if the highest valid measurement
190 of a sonde is above 8000m. The *profile-sparsity* test is passed if less than 20% of the theoretically available data is missing. This is an adapted version of the *profile-fullness-test (sat-test)* in George et al. (2021). The *near-surface-coverage* test is passed if 50 or more measurements have been made in the lowest 1km. The altitude measurement used depends on the variable, since u and v have `gpsalt` as their reference altitude, while p , temperature, and RH have `alt` as their reference altitude in the Level 1 data. Figure 3 shows the histograms for the QC tests that determine the profile fullness
195 of a measured variable. It shows, that the horizontal wind measurements have a larger fraction of missing values than the PTU measurements and that the relative humidity sensor needs a longer time to equilibrate, resulting in lower profile extents.

QC: sfc-physics In addition to those QC tests which are aimed to check the fullness of a profile, the bottom-most value in each profile is checked for physical plausibility; i.e. the check is passed if the lowest RH measurement exceeds 0.3, the
200 lowest temperature measurement is above 293.15K, and the lowest pressure is between 1005hPa and 1020hPa. Those thresholds were chosen relatively lax and for a tropical atmosphere. They should be chosen differently for dropsondes in other locations, as for example during HALO-(AC)³. Two things can lead to a failed *sfc-physics* test: (i) the sonde did not send data until it reached the surface; and (ii) there was a calibration issue. In the first, more common, case the *near-surface-coverage* test is failed as well. The second case was only triggered for a single sonde (0bd0e322 on HALO-
205 20240924a), which was unintentionally factory-reset as described in 2.2, and consequentially has a shifted temperature profile and anomalously low RH measurements. The corresponding values have been masked in Level 3 so as to not adversely impact other calculations.

3.2.2 Variables in Level 2

For Level 2, all sondes are concatenated along the `time` dimension into a dataset with a ragged array data structure (Brian
210 Eaton et al., 2024) with dimensions `time` and `sonde`. The `times_per_sonde` variable contains the number of time
measurements per sonde. Only the measurements for temperature (`ta`), relative humidity (`rh`), pressure (`p`), and the wind
components u (`u`), v (`v`), and w (`w`) are transferred from the Level 1 output to the Level 2 dataset. Any other derived variables
are removed. Additionally, positional variables such as latitude (`lat`), longitude (`lon`), and sonde altitude obtained from GPS
(`gpsalt`) and pressure (`alt`) at each time point are included. The flight altitude, time, and position at drop are contained as
215 variables along the `sonde` dimension.

The four variable-specific QC flags are combined into one QC variable per physical variable, which is called `*_qc` and
contains all information in binary format (Brian Eaton et al., 2024, sec 3.5). In Level 2, the detailed results of the QC analysis
are stored in respective QC variables, that have the naming pattern `{variable}_{qc_name}_{value_type}`.

To help parse the results of the tests, an overall `sonde_qc` variable is introduced that is GOOD if the Profile Fullness and
220 Surface Physics tests are passed for all variables. A variable of a sonde is BAD if all individual QC tests are failed, or if the
sfc-physics is the only failed test as the sondes measurements are deemed unphysical then. Any other sondes are flagged as
UGLY, since they contain valid measurements for some purposes. Apart from one sonde that did not have any valid data for
any of the variables in Level 2, all sondes are at least Ugly.

After the Level 2 QC 976 sondes are GOOD, 139 sondes are UGLY, and no sonde is BAD.

225 3.3 Level 3 processing: A combined dataset

Level 3 (Gloeckner et al., 2026d) is a combined dataset of all GOOD and UGLY Level 2 dropsondes. It contains the data of
those sondes interpolated to the same altitude grid, as well as the QC flag for each variable for each sonde. There exists a
separate Level 3 QC dataset that contains all QC details. The data was split in this way, because for most use cases the QC
details are irrelevant and unnecessarily clutter the Level 3 product. The Level 3 dataset and the Level 3 QC dataset have the
230 same dimensions and coordinates so that they can be easily merged if necessary.

To obtain the same altitude grid, sondes are interpolated to the same 10m altitude grid. After the interpolation, 10m sections
that do not contain a measured value are masked. Instead of interpolating T directly, θ is calculated as

$$\theta = T \cdot \left(\frac{p_0}{p} \right)^{\frac{R_d}{c_p}}, \quad (2)$$

and T is recalculated on the interpolated data, because θ behaves more linear, to form a consistent dataset.

235 Before the interpolating in altitude for Level 3, `gpsalt` is linearly interpolated in `time`, because it can happen that a given
point in time has a PTU measurement, but no GPS-altitude measurement.




Figure 4. Distribution of surface pressure and gpsalt values from all sondes. (xrange alt: 100m, xrange p: 10hPa)

3.3.1 Defining a common altitude

By default the Level 2 data contains two separate altitude variables: `gpsalt`, which is derived from GPS measurements, and `alt`, which is calculated from p using the assumption of hydrostaticity. It is a priori not clear which altitude dimension should
 240 be used, so we will use a simple train of thought to justify our decision:

If a perfect sonde falls at roughly 10m s^{-1} , we would expect the last altitude measurement to be equally distributed between 0m and 10m. Everything outside of this range would then be either an error on the GPS measurement, or a sonde that did not send data up to its splash in the ocean. If we further assume that a 10m difference in altitude is roughly equal to a 1hPa difference in pressure, we would accordingly assume the surface pressure to roughly vary within 1hPa.

245 Figure 4 shows the probability histogram of the last pressure and last `gpsalt` measurements. The axes are chosen such that 10m in `gpsalt` are equivalent to 1hPa in pressure. For `gpsalt`, most of the values indeed fall within 0m to 10m, but there is an error of roughly 20m. The pressure measurements have a slightly larger dispersion, but that is expected since the surface pressure is not exactly the same everywhere. There is however no reason to assume that one altitude measurement is better than the other, because the range of the histograms is similar. Considering that the `alt` variable in addition to the pressure
 250 measurement error assumes hydrostasis, which is not always valid (especially at higher altitudes), we decided to use `gpsalt` as the default altitude coordinate in Level 3.

For sondes that do not have a valid `gpsalt` profile, i. e. if the *gps-valid* QC failed, the `alt` variable is used for the altitude if a valid surface-*p* measurement was taken. In addition, if the `gpsalt` profile is incomplete, i.e. if the *near-surface-count* or the *profile-extent* QC tests failed for *u* (meaning there was poor GPS signal), the `alt` variable is used for the altitude if

- 255 1. the QC that was failed for *u* was passed for *p* and
 2. a valid surface-*p* measurement was taken.

The first condition ensures that an incomplete `gpsalt` profile is not replaced by an equally incomplete `alt` profile. The second condition is required because the `alt` calculation needs a reference height, and the ASPEN software used for generating Level 1 assumes that the last measured pressure is the surface pressure for that purpose. Although we can never say with 100%
 260 certainty that a sonde sent data until its splash, we assume that it was close enough if the *sfc-physics* QC was passed for *p*.

The chosen altitude dimension for each sonde is renamed to `altitude` in Level 3 to indicate that it is a new variable, and an ancillary variable `altitude_source` is added to the QC dataset, which contains the name of the altitude variable from Level 2 that is used as the altitude in Level 3 for each sonde. For five sondes on HALO-20240827a, neither `gpsalt` nor `alt` provides a valid altitude. These sondes are dropped between Level 2 and 3.

265 3.3.2 Variables in Level 3

In addition to the Level 2 variables, Level 3 contains θ , q , integrated water vapor (IWV), wind direction and wind speed. Vertical velocity w of individual sondes as estimated by ASPEN, is removed between Level 2 and Level 3. The specific humidity q is calculated from RH and T using the saturation vapor pressure from Hardy (1998), because the same formulation is used by Vaisala for the calibration of the sondes:

$$270 \quad \ln e_s = \sum_{i=0}^6 g_i t^{i-2} + g_7 \ln(t) \quad (3)$$

with the coefficients given in Tab. 2

Table 2. Coefficients in Hardy (1998) formula for saturation vapor pressure.

$g_0 = -2.8365744 \cdot 10^3$	$g_1 = -6.028076559 \cdot 10^3$	$g_2 = 1.954263612 \cdot 10$	$g_3 = -2.73783018 \cdot 10^{-2}$
$g_4 = 1.6261698 \cdot 10^{-5}$	$g_5 = 7.0229056 \cdot 10^{-10}$	$g_6 = -1.8680009 \cdot 10^{-13}$	$g_7 = 2.7150305$

The integrated water vapor (IWV) is only added for sondes that have a GOOD quality control flag for RH, p , and T measurements and set to NaN otherwise to ensure an adequate representation of the actual IWV. We use

$$IWV = \int q \rho_v dz, \quad \text{where} \quad \rho_v(p, T, q) = \frac{p}{(R_d + (R_v - R_d)q)T}. \quad (4)$$

275 The time coordinate from Level 2 is also interpolated and stored in an `interpolated_time` variable. This time coordinate, as it is interpolated, is no longer useful to calculate fall speeds. These should be computed from the Level 2 data if necessary.

3.4 Level 4: Circle Variables

Level 4 (Gloeckner et al., 2026f) of the BEACH datasets contains the mesoscale divergence (\mathcal{D}), vorticity, vertical velocities, and pressure velocity (ω), all of which were derived from circles. Circles usually had a radius of either ca 133km and a duration of one hour, or of about 70km and a duration of 40 minutes (see Section 2).

The BEACH Level 3 sonde dataset is grouped into individual circles according to the flight segmentation. The segmentation provides the circle times, latitudes, longitudes, and radii. A sonde belongs to a given circle if it was dropped between the circle's start and end time. In addition sondes dropped inside the circle area and within 20 minutes from the circle start or end are tagged as `extra_sondes` in the flight segmentation for that circle and are also included.

We apply the linear regression method described in (Bony and Stevens, 2019) in order to obtain the gradient terms and the mean profiles. From these, we compute divergence, vorticity, vertical velocity, and ω . Before applying the regression, vertical gaps in the Level 3 profiles are interpolated using the Akima method (Akima, 1970). The details of the interpolation, the regression, the circle products, and corresponding errors are described in this section.

3.4.1 Circle Fit

The regression method is applied as described in Bony and Stevens (2019) and George et al. (2021): The solution to the equation

$$\phi(x, y) \approx \phi_0 + \frac{\partial\phi}{\partial x} \Delta x + \frac{\partial\phi}{\partial y} \Delta y, \quad (5)$$

where Δx and Δy are the eastward and northward distances to the circle center, is found by solving the least square problem

$$\min_x \|(Ax - b)\|_2, \quad (6)$$

where $x = \begin{bmatrix} 1 \\ \frac{\partial\phi}{\partial x} \\ \frac{\partial\phi}{\partial y} \end{bmatrix}$, $b = \begin{bmatrix} \phi_1 \\ \dots \\ \phi_k \end{bmatrix}$ and $A = \begin{bmatrix} 1 & \Delta x_1 & \Delta y_1 \\ \dots & \dots & \dots \\ 1 & \Delta x_k & \Delta y_k \end{bmatrix}$ for a circle with k sondes. This system can be solved with the

Moore-Penrose pseudo-inverse to derive ϕ_0 , which is the circle mean, $\frac{\partial\phi}{\partial x}$, which is the linear variation in the eastward direction, and $\frac{\partial\phi}{\partial y}$, which is the linear variation in the northward direction (George et al., 2021). Each of these variables are given at every altitude that contains values from six or more sondes, after gaps are vertically interpolated. The details for the interpolation are discussed in Section 3.4.3.

3.4.2 Circle Products

The above mentioned components for u and v on the circle scale can be used to derive the area-averaged horizontal divergence, \mathcal{D} , and vorticity ζ :

$$\mathcal{D} = \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \quad \text{and} \quad \zeta = \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y},$$

305 and the vertical velocity w , and the pressure velocity ω given as

$$w(z) = \int_0^z \mathcal{D}dz \quad \text{and} \quad \omega(z) = - \int_{p_{\text{stc}}}^{p_z} \mathcal{D}dp. \quad (7)$$

3.4.3 Vertical gap interpolation of sonde profiles

Although 12 to 15 sondes were dropped in a typical circle, not all circles contain full measurements from 12 or more sondes (see Section 2.2). This raises the questions of (1) how to handle sondes without valid data and (2) what to do with sondes that provide partial data, but don't pass all quality control checks (Section 3.2). Sondes that do not provide valid data, for example due to a launch detection failure, were ignored. Following an error analysis by Bony and Stevens (2019), the errors incurred should be tolerable if six or more sondes contribute to the products. In the case of measurements from fewer than six sondes circle products are not calculated. That is only the case for the ATR-coordinated circle on HALO-20240827a and the first circle on HALO-20240914a, due to a full dropsonde system failure on both flights.

315 Apart from sondes that do not contain data at all, there are many sondes with vertical gaps in the measurements. The profile sparsity is largest for u and v , which are used in the divergence calculation (Figure 3 left). Since the meteorological situation is very variable in the ITCZ (the domain of BEACH), dismissing information that could be interpolated with confidence might lead to larger errors in the divergence, vorticity, vertical velocity, and ω estimation than would arise from simply interpolating gaps. We tested this train of thought using circles with at least 12 sondes with GOOD u , v and p measurements, as defined in Section 3.2. For one of those circles with n sondes, ω was calculated $2 \cdot (n+1)$ times: (1) using all available data and no vertical interpolation (*no int*), (2) using all data and vertically interpolating gaps with the Akima method (*int*), (3) once for every sonde assuming that the sonde has an artificial gap at 500m with interpolation (*gap int*), and (4) once for every sonde assuming that the sonde has an artificial gap at 500m without interpolation (*gap no int*). For interpolation we use the Akima splines (Akima, 1970), which are similar to a cubic-spline interpolation but less prone to overshooting. In the boundary layer missing values are extrapolated by assuming constant u , v , and θ , and linear extrapolation for $\log(p)$ and RH.

325 Figure 5 shows the vertical sum of the differences between *no int* and *gap no int* on the x-axis and the vertical sum of the differences between *int* and *gap int* on the y-axis for different gap sizes in the columns. The difference to the full calculation becomes larger for larger gap sizes (left to right), which is not surprising as more information about the actual situation is missing. For small gaps, the vertical mean absolute error without interpolation ($\approx 0.01 \pm 0.02 \text{hPa hr}^{-1}$) is roughly two orders of magnitude larger than when an interpolation is applied ($\approx 2 \times 10^{-4} \pm 2 \times 10^{-4} \text{hPa hr}^{-1}$). For intermediate gaps, there are some calculations with a noticeably larger error with an interpolation, but overall the mean of absolute errors ($\approx 0.03 \pm 0.03 \text{hPa hr}^{-1}$) is approximately quartered as compared to no interpolation ($\approx 0.13 \pm 0.15 \text{hPa hr}^{-1}$). For large gaps, it does not make a difference whether an interpolation is applied ($\approx 0.41 \pm 0.44 \text{hPa hr}^{-1}$), which again is anticipated as we do not expect the measurements to be informative over such large gaps. Based on those results, we interpolate gaps up to 1500m.




Figure 5. Difference to full omega calculation for artificially introduced vertical gaps of 30m, 300m, and 1500m depth at 500m altitude for circles with 12 or more sondes with GOOD u , v and p measurements. Differences using the interpolation method are on the x-axis, differences without interpolation on the y-axis. Each color represents a circle and each large dot the median over altitude for one sonde. The grey area illustrates where the interpolation leads to better results than no interpolation. Be aware that the axes do not have the same scale in the different columns.

335 3.4.4 Error measures

We calculated the regression standard error for the circle products as described in Bony and Stevens (2019). In addition, we tested the sensitivity of ω to the removal of an individual sonde in the circle. To do so, we removed each sonde from its circle and calculated the circle products again while ignoring this sonde. The difference to the value calculated using all profiles in the circle is stored in a variable `omega_remove_sonde_qc`.

340 Figure 6 (middle and right) shows the ω of seven arbitrary circles with the two different error measures in shading. In the middle panel we show the regression standard error and on the right we show the span between the minimum and maximum value if a sonde is removed. The sonde-removal calculation was also done for \mathcal{D} , vorticity, and vertical velocity. The results are stored in the variables `*_remove_sonde_qc`.

345 Although the regression standard error is small ($\approx 6\%$ in the mean), individual sondes can have a large impact on the calculated omega. Figure 6 (left) shows a histogram of the errors that emerge if individual sondes are removed. It illustrates that the relevance of the sondes increases with height, which is a direct implication of the integration of divergence, and that the overall mean ω is reliable up to $3.1\text{hPa}\text{hr}^{-1}$ (2σ , averaged over altitude). For individual circles the importance of single sondes can be much larger, as indicated by the spread in the histogram. Hence, if studying individual circles it is useful to check the relevance of individual sondes before interpreting omega values. Encouragingly, the sign of ω does not change in
 350 roughly 90% of the points (and if so then for values of $\omega \approx 0$).




Figure 6. Histogram of the error ϵ_ω , with profiles from all circles where any sonde is removed from its circle in lower panel (left). Regression standard error for seven arbitrary circles (middle). Range of minimum and maximum estimate if a sonde is removed from circle calculation for the same circles (right).

3.4.5 Variables in Level 4

The Level 4 dataset contains the interpolated profiles of all sondes that belong to a circle as well as all circle products. Consequentially, in addition to the altitude dimension, it has a circle and a sonde dimension, but each variable is only dependent on one of them. To connect both, the dataset has a contiguous ragged array representation following the respective
 355 CF-conventions. It contains a `sondes_per_circle` variable which gives the number of sondes contained in each circle. As long as both dimensions remain sorted by `circle_time` and `launch_time` respectively, this structure allows to select all sondes from a circle.

For the sondes, only the `sonde_qc` information is kept. The interpolated time is also not in the Level 4 dataset. For each of p , RH , q , T , θ , u , and v the fit as described in Section 3.4.1 leads to the new variables `mean_*`, `d*dx` and `d*dy` on the circle
 360 level. The necessary x and y variables are also saved in the Level 4 dataset.

In addition to divergence, vorticity, ω , and vertical velocity, the standard errors and the remove-sonde-errors are added for those variables. The latter have `sonde` instead of `circle` as a dimension, because they are the difference in specific variables, if one sonde is removed.

3.5 Main differences between BEACH and JOANNE

365 Although the processing has been adapted from the JOANNE processing, which has been conducted on the EUREC⁴A drop-sonde data (George et al., 2021), many steps in the processing have changed.

From Level 2 onward, BEACH does not use the serial id from the manufacturer as a unique reference to a sonde since there can be multiple files containing the same serial id (Section 2.2). Instead, a hash derived from the first line in the D-file (including launch-time and manufacturer serial-id) was used.

370 For the calculation of Level 2, the QC tests have been rearranged: The *profile-fullness/sat-test* in JOANNE is renamed to *profile sparsity* and the fraction of missing values as compared to a hypothetical perfect sonde measurement is calculated instead of the fraction of measured values. JOANNE’s *surface-test* has been split into a surface count (*near-surface-count*) and a surface physics test (*sfc-physics*). They are both present in JOANNE but combined to the *low-altitude-measurement-test/low-test*. An additional test that checks the *profile-extent* was introduced to flag sondes with incomplete profiles, e.g. if the
375 parachute opened very late, and which are problematic when comparing integrated quantities such as integrated water vapor.

Contrarily to JOANNE, sondes that did not pass the tests were not discarded from the dataset, but flagged in the Level 2 and Level 3 datasets. Additionally, a `{var}_qc` variable is introduced that contains the QC information in binary format for all variables. Most of those changes have little impact on the overall QC flag, but were introduced to account for edge cases that occurred during PERCUSION and are not covered by the JOANNE QC framework.

380 During PERCUSION, all sondes were reconditioned on the morning of the flight that they were dropped. As a consequence there is no dry bias correction (as described in George et al. (2021)) and the humidity measurements are consistent across data levels.

In BEACH Level 3, the altitude derived from GPS measurements, `gpsalt`, is used as the default height coordinate instead of pressure altitude `alt`. However, if no `gpsalt` values were present or the `alt` measurements were better, the latter is used.

385 In JOANNE, q and θ are binned to a 10m grid, while we chose to interpolate to the same grid. The decision to bin was made as to not have values that are not measured in the dataset. This approach was changed, because binning instead of interpolating introduces an error in height of up to 5m per 10m bin. Although irrelevant for most applications, it creates an error of a couple of centimeters in the hydrostatic equation that adds up over the depth of the troposphere and is avoided by interpolating. To still maintain consistent RH and q values, BEACH interpolates in RH instead of q , because the former is more linear.

390 While JOANNE linearly interpolates gaps of up to 50m in altitude, BEACH Level 3 does not include any gap interpolation larger than 5m in Level 3. It does however use the Akima-splines interpolation on gaps in the measurements before the circle fits, assumes constant u , v , and θ , and linear RH and $\log(p)$ in the lowest 300m if there are no measurements for Level 4. For convenience, those interpolated profiles are contained in BEACH’s Level 4. BEACH also uses sondes that did not pass all QC tests for the Level 4 calculation if the QC of used variables were passed, while JOANNE only uses sondes that passed every
395 QC test.

JOANNE and BEACH use the same formula for vertical velocity w , but BEACH uses the integration over divergence in p for the vertical pressure velocity ω , while JOANNE uses

$$\omega_h = -\rho \cdot g \cdot w. \tag{8}$$




Figure 7. Mean wind direction in the lowest 50m for all sondes. Level 3 data is used for this plot.

Calculating w and ω independently comes at the cost that they cannot be easily transformed anymore, but it calculates each
 400 variable relative to its altitude coordinate, and hence is more physically accurate. For BEACH, the sonde relevance variables
 were calculated in addition to the regression standard error.

4 Data overview

The dropsondes' surface wind and integrated water vapor measurements give an indication of the conditions spanned by the
 BEACH data. The moist tropics are mostly defined by an integrated water vapor above 48mm (Mapes et al., 2018), with a
 405 peak near the southern edge of the ITCZ (Windmiller and Stevens, 2024). The integrated water vapor (Fig. 8) confirms that
 most measurements have been taken within the moist tropics, especially in the East, oftentimes with an IWV much greater than
 48mm.

Since the ITCZ is marked by strong convergence at the surface, we expect the surface wind direction to change at the edge(s)
 of the ITCZ. Figure 7 shows the surface wind direction of all sondes. Especially in the East, ~~the transition from southerlies and
 410 northerlies to westerlies is apparent, indicating the southerlies turn westward as they cross the equator to form the monsoon
 trough (Flohn, 1951; Praturi and Stevens, 2026). This indicates~~ that most of the dropsondes sampled the breadth of the ITCZ
~~also by this measure. Westerly surface winds in the Eastern Atlantic are in line with the idea of an equatorial westwind zone
 (Flohn, 1951). by measure of surface wind direction as well as IWV. Sometimes the ITCZ is distinguished from the monsoon
 trough through its absence of westerlies, a distinction we do not adopt in this paper.~~

415 In the West, neither the wind nor the integrated water vapor field follow the clear structure that is apparent in the East,
 consistent with the ITCZ being less well defined there (Stevens et al., 2026, Fig. 6).




Figure 8. IWV of all sondes. Colorbar is centered at 48mm. Level 3 data is used for this plot.

The vertical profile of the wind shows weak baroclinicity (Fig. 9), where the zonal wind changes sign above 10 km, with the predominant westerlies at the surface transform into strong easterlies higher up in the East, while the easterlies in the West become westerlies in the West. The larger spread in the near surface wind component v in the East is further evidence for the larger variety of conditions that were sampled there. In the North region, easterlies dominate the whole column, indicating that those sondes were mostly dropped in the trade wind region.

Although it is generally moist everywhere, the relative humidity profiles vary considerably. The mean profile in the East has two distinct peaks at the top of the boundary layer, and near the freezing level and a minimum in between. This structure fits to the trimodal characteristic of tropical convection (Johnson et al., 1999) and might be of interest because reanalyses and satellite observations struggle to represent the elevated moist layers in the mid-troposphere (Prange et al., 2023). This feature is even more pronounced in the North, which might be indicative of a role for dry Saharian air on this region, especially since the winds are predominantly north-easterlies above 2000m. In the West, the mean profile shows less evidence of a freezing level maximum, but the troposphere below the 0° isotherm is moister than the troposphere aloft. Note that the humidity profiles above the freezing level rarely reach 100% but might still be saturated, especially at higher altitudes, as relative humidity is calculated with respect to water instead of ice.

Measurements coordinated with the SAFIRE ATR-42 were in the North region. Although the mean conditions there indicate the trade wind region, some circles as well as a bimodal IWV distribution (see Appendix Figures B1, B2) point to several interesting cases in or at the edge of the ITCZ.

Mesoscale vertical air motion in the East, as shown in Fig. 10, supports the idea of the ITCZ being a region of mean ascent with convergence below ≈ 2000 m and divergence above 10000m. The West Atlantic shows mean surface divergence in the lower-troposphere similar to the divergence profile from the JOANNE data measured during the EUREC⁴A campaign, which sampled the winter trades in 2020 (yellow). However, whereas ω in JOANNE indicates subsidence through a deep layer,




Figure 9. Profiles of individual sondes from BEACH Level 3 for θ , RH, u and v from the East region (dark red), West region (blue), and North region (light red). Means are plotted with thick lines while the thin lines correspond to individual profiles (East of -40°E in red and West of -40°E in blue). The gray horizontal lines mark the mean freezing level and the lower relative humidity peak respectively. BEACH Level 3 data is used for this plot.

the mean ω of the Western measurements in BEACH shows rising air motion above $\approx 3000\text{m}$. The switch in the direction of vertical air motion however is caused by strong updrafts at these levels prevailing in some circles. Considering the median
 440 vertical velocity instead of the mean shows predominant subsidence in the West Atlantic above 3000m as well. The mean in the East Atlantic is similarly influenced by deep convective events, but the median still shows upward air motion in the whole column.

The East-West difference in mesoscale divergence is shown in more detail in Figure 11 for all circles in the BEACH dataset. Each column is a circle and flights are separated by black vertical lines with circles and flights being sorted in time from left to
 445 right (similar to George et al., 2023, Figure 1). The transfer from sampling the East Atlantic to the West Atlantic on September




Figure 10. Average divergence (left) and ω (right) for the full campaign, as well as split into East and West Atlantic. The average profiles from the JOANNE dataset showing typical characteristics for the winter trades in the Western Atlantic are added for comparison. BEACH Level 4 data is used for this plot.




Figure 11. Divergence for all circles during the campaign. Flights are separated by black lines. The transfer between East and West is marked at the top. Circles are ordered in time. Level 4 data is used for this plot.

6th is marked at the top. Most obvious are the stronger convergence and divergence patterns in the East (stronger red and blue colors) compared to less strong patterns in the West except for the flight on September 24th.

5 Summary

The PERCUSION and MAESTRO aircraft campaigns took place in August and September 2024 in the tropical Atlantic as part of ORCESTR450A. A main focus of ORCESTR450A was the influence of convective and mesoscale circulation systems on the mean structure of the ITCZ. As part of this effort, 1191 sondes were dropped from the research aircraft HALO. This paper presents the data from these sondes in the form of the BEACH datasets.

The BEACH datasets contain four Levels of processing of the raw data (Level 0), as schematically depicted in Fig. 2. This leads to the following data levels: ASPEN quality controlled data (Level 1); custom quality controlled data (Level 2); a combined dataset including all sondes that have at least partially valid data (Level 3); and a dataset containing divergence and vertical velocity on the mesoscale for all circles with sufficient valid sonde measurements (Level 4). All datasets are openly available on IPFS (Stevens et al., 2026). The general hierarchy of the levels and most parts of the processing were adapted from the EUREC455⁴A dropsonde processing (George et al., 2021).

The BEACH dropsonde data confirms that the PERCUSION flight tracks span the meridional extent of the ITCZ, with measurements within as well as at the edges as defined by the surface wind field and integrated water vapor (Figures 7, 8). It also samples zonal variations within the ITCZ. While the ITCZ is clearly outlined in both the surface wind direction and IWV in the East, in the West it is less structured. Another difference is that in the East there are two distinct peaks in the mean relative humidity – around the freezing level and above the subcloud layer. In the West, there is no distinct peak in RH at the freezing level. In the wind profiles, known dynamical features such as surface westerlies within the equatorial trough are captured as well as a weak imprint of an Atlantic Walker cell, and the African Easterly Jet.460465

A core objective of the flight strategy was the derivation of mesoscale divergence and vertical velocities from sondes dropped on circular flight patterns. We succeeded in processing 87 circles from all flights that show on average upward motion with stronger updrafts in the East compared to the West Atlantic. Furthermore, in the lowest 2000m the vertical velocity profile in the West is closer to the JOANNE measurements from the wintertime trades compared to the measurements from the summertime East Atlantic, leaving much room for further analysis.470

6 Code and data availability

All datasets in the hierarchy of BEACH are made available via IPFS and have a landing page in the ORCESTR475A-browser: <https://browser.orcestra-campaign.org/>

The respective dois are

L0 <https://doi.org/10.82246/bafybeif4n7lov7syd5ragolfosqf6t7zuawexxjhr5r2ezecvqvj7eje> (Gloeckner et al., 2026a)

L1 <https://doi.org/10.82246/bafybeieqqc5fktpmwbluppgb4vidrrx7vv44l5e7rol6zp3e3wlbjmq2q4> (Gloeckner et al., 2026b)

L2 <https://doi.org/10.82246/bafybeifi5pglGPCq6onwb3yixhvghcmenkvqmcqi7qgjbrbqlgoyrvj52i> (Gloeckner et al., 2026c)

L3 <https://doi.org/10.82246/bafybeiczbv7mycr2jois6t4dq3zwiltycomwo5xxvjqcz2ot3newzar6q> (Gloeckner et al., 2026d)

L3qc <https://doi.org/10.82246/bafybeidytxgupj3evqesvvuv3vsl1v6pbcu5gc2torqkz7yo6yuazwtmu> (Gloeckner et al., 2026e)

L4 <https://doi.org/10.82246/bafybeibgeeqs5uhmbqy4hz4v3pihrfisiklcetisxkf63d2r473szaprwi> (Gloeckner et al., 2026f)

Further information on the ORCESTRa data policy and concept can be found in the ORCESTRa overview paper (Stevens et al., 2026) and on the ORCESTRa campaign website. The dropsonde processing software generating the various data levels is available on GitHub in the pydropsonde repository as well as a Python package called *pydropsonde* via the Python Package Index (PyPI). For the processing and plots presented in this paper version 0.5.5 was used which includes the initial processing
480 with ASPEN v4.0.4. The ASPEN software is hosted in a docker image (on GitHub), making it independent from the operating system. The repository includes a Dockerfile and the respective GitHub workflows needed to generate the image and push it to the GitHub container registry. It can be used via its name ghcr.io/atmdrops/aspenc. The configuration file for running *pydropsonde* on the ORCESTRa dropsondes as well as all analysis scripts generating plots and tables for this paper are stored on GitHub in the orcestra-campaign-dropsondes repository.

485 The BEACH datasets are stored such that they can be easily accessed with a few lines of code. For example, one can access the Level 3 BEACH dataset directly using Python. This requires a working IPFS Gateway and the ipfsspec package to be installed.

```
1: import xarray as xr  
490 2: ds = xr.open_dataset(  
3:     "ipfs://bafybeiczbv7mycr2jois6t4dq3zwiltycomwo5xxvjqcjz2ot3newzar6q",  
4:     engine="zarr",  
5: )
```

495 **Appendix A: Glossary**

ASPEN Atmospheric Sounding Processing ENvironment

AVAPS Airborne Vertical Atmospheric Profiling System

BAHAMAS Basic Halo Measurement and Sensor System

BEACH Barbados and Eastern Atlantic Combined High-altitude dropsonde datasets

500 **EarthCARE** Earth Clouds, Aerosols and Radiation Explorer

EUREC⁴A Elucidating the role of clouds-circulation coupling in climate 2020

GATE GARP Atlantic Tropical Experiment 1974

GARP Global Atmospheric Research Project

GPS Global Positioning System

505 **HALO** High-Altitude and LOng range research aircraft

HALO-(AC)³ Arctic Air Mass Transformations During Warm Air Intrusions and Marine Cold Air Outbreaks 2021

ICON The ICOSahedral Non-hydrostatic model

IFS Integrated Forecasting Model (ECMWF)

INMG Instituto Nacional de Meteorologia e Geofísica (Cabo Verde)

510 **IPFS** Inter Planetary File System

ITCZ InterTropical Convergence Zone

JOANNE Joint dropsonde Observations of the Atmosphere in tropical North-atlaNtic meso-scale Environments

MAESTRO Mesoscale organisation of tropical convection subcampaign of ORCESTR

NARVAL2 Next-generation Aircraft Remote-sensing for VALidation studies (2) 2016

515 **NCAR** National Center of Atmospheric Research (US)

netCDF Network Common Data Format

ORCESTR Organized Convection and EarthCARE Studies over the Tropical Atlantic 2024

OTREC Organisation of Tropical East Pacific Convection 2019

PERCUSION Persistent EarthCARE underflight studies of the ITCZ and organized convection subcampaign of ORCESTR

520 **PTU sensor** Pressure Temperature and hUmidity sensor

RAPSODI Radiosonde Atmospheric Profiles from Ship and island platforms during ORCESTR, collected to Decipher the ITCZ

SAFIRE ATR-42 Service des Avions Français Instrumentés pour la Recherche en Environnemen (Avions de Transport Régional 42)

525 **Appendix B: Setting SAFIRE ATR-42coordinated measurements into the PERCUSION context**

During the ORCESTR campaign, 10 circles were flown in coordination with the SAFIRE ATR-42. Table B1 shows the segment ids of those circles as well as the SAFIRE ATR-42flight that was closest in time, and the number of sondes in Level 3 and Level 4 for those circles. Apart from one circle on HALO-20240827, where the system shut down (see Section 2.2), all SAFIRE ATR-42-coordinated circles have 10 or more sondes

530 Figure B1 is similar to Figure 10, but including the ATR divergence and omega estimates. Contrarily to the figure in the main text, here the mean omega and divergence for the East Atlantic does not include the ATR values. This illustrates that the

Table B1. PERCUSION ATR coordination statistics showing the closest ATR flight to each atr-coordinated HALO circle and the number of sondes in Level 3 and Level 4 for those circles.

flight ID	flight date	flight time	Level 3 sondes	Level 4 sondes	HALO circle ID
ATR-20240811	2024-08-11	14:33:45-18:13:42	12	12	HALO-20240811a_7a38
ATR-20240813b	2024-08-13	19:11:11-22:34:54	12	12	HALO-20240813a_63f7
ATR-20240816b	2024-08-16	16:06:24-19:38:05	12	12	HALO-20240816a_7241
ATR-20240822a	2024-08-22	13:55:27-17:32:49	10	10	HALO-20240822a_049e
ATR-20240822b	2024-08-22	19:33:26-22:42:09	10	10	HALO-20240822a_b5e4
ATR-20240825	2024-08-25	19:21:56-22:41:33	12	12	HALO-20240825a_64c5
ATR-20240827	2024-08-27	19:02:25-22:34:22	4	4	HALO-20240827a_107f
ATR-20240829	2024-08-29	13:52:13-17:40:57	12	12	HALO-20240829a_3585
ATR-20240831	2024-08-31	13:57:37-17:30:33	12	12	HALO-20240831a_e17f
ATR-20240903	2024-09-03	16:33:28-19:51:44	12	12	HALO-20240903a_71a0




Figure B1. Divergence for all circles during PERCUSION (dark blue), for the West Atlantic (light blue), compared to only circles in the East Atlantic (without ATR - red) and only the circles flown in coordination with SAFIRE ATR-42 (yellow).

ATR circles were mostly flown in a different environment than the larger circles. Thin yellow lines in the plot are individual ATR circles and demonstrate the spread in the measurements.

Figure B2 further shows the distribution in integrated water vapor and how it differs between SAFIRE ATR-42 coordinated sondes and the others. Again, the curve for the East Atlantic excludes SAFIRE ATR-42 coordinated measurements. All distributions have a peak close to 60mm IWV, which is well above the 48mm threshold assumed for the ITCZ in the long term

mean. The distribution of ATR dropsondes has a distinguished second peak at lower IWV values ($\approx 48\text{mm}$), indicating that the majority of those sondes were dropped at the edge or outside of the ITCZ.




Figure B2. Normalized probability density of integrated water vapor for the full PERCUSION campaign (dark blue), for the West Atlantic (light blue), from the Eastern Atlantic (without ATR - red), and only from measurements in coordination with the SAFIRE ATR-42 (yellow).

Appendix C: Variables in Level 3

Table C1. BEACH Level 3 Variables

object	variable	units	standard_name	dimensions
Coordinates	altitude	m	altitude	altitude
	launch_altitude	m		sonde
	launch_lat	degrees_north	deployment_latitude	sonde
	launch_lon	degrees_east	deployment_longitude	sonde
	launch_time			sonde
Variables	flight_id			sonde
	interpolated_time		time	sonde altitude
	iwv	kg m-2	atmosphere_mass_content_of_water_vapor	sonde
	lat	degrees_north	latitude	sonde altitude
	lon	degrees_east	longitude	sonde altitude
	p	Pa	air_pressure	sonde altitude
	p_qc		quality_flag	sonde
	platform_id			sonde
	q	kg kg-1	specific_humidity	sonde altitude
	rh	1	relative_humidity	sonde altitude
	rh_qc		quality_flag	sonde
	sonde_id			sonde
	sonde_qc		aggregate_quality_flag	sonde
	ta	K	air_temperature	sonde altitude
	ta_qc		quality_flag	sonde
	theta	K	air_potential_temperature	sonde altitude
	u	m s-1	eastward_wind	sonde altitude
	u_qc		quality_flag	sonde
	v	m s-1	northward_wind	sonde altitude
	v_qc		quality_flag	sonde
	vaisala_serial_id			sonde
wdir	degree	wind_from_direction	sonde altitude	
wspd	m s-1	wind_speed	sonde altitude	

Table D1. BEACH Level 4 Variables

object	variable	units	standard_name	dimensions
Coordinates	altitude	m	altitude	altitude
	circle_lat	degrees_north		circle
	circle_lon	degrees_east		circle
	circle_time			circle
	launch_time			sonde
	sondes_per_circle			circle
Variables	*_d*dx	* m-1	eastward_derivative_of_*	circle altitude
	*_d*dx_std_error	* m-1	eastward_derivative_of_* standard_error	circle altitude
	*_d*dy	* m-1	northward_derivative_of_*	circle altitude
	*_d*dy_std_error	* m-1	northward_derivative_of_* standard_error	circle altitude
	*_mean	*		circle altitude
	circle_altitude	m		circle
	circle_id			circle
	circle_radius	m		circle
	div	s-1	divergence_of_wind	circle altitude
	div_sonde_relevance			sonde altitude
	div_std_error	s-1	divergence_of_wind standard_error	circle altitude
	omega	Pa s-1	vertical_air_velocity_expressed_as_tendency_of_pressure	circle altitude
	omega_sonde_relevance			sonde altitude
	omega_std_error	Pa s-1	vertical_air_velocity_expressed_as_tendency_of_pressure standard_error	circle altitude
	vor	s-1	atmosphere_upward_relative_vorticity	circle altitude
	vor_sonde_relevance			sonde altitude
	vor_std_error	s-1	atmosphere_upward_relative_vorticity standard_error	circle altitude
	wvel	m s-1	upward_air_velocity	circle altitude
	wvel_sonde_relevance			sonde altitude
	wvel_std_error	m s-1	upward_air_velocity standard_error	circle altitude
	x	m		sonde altitude
	y	m		sonde altitude

Appendix E: Problematic sondes

Author contributions. SB, JW, and BS conceptualized the measurements and developed the flight and drop strategy for PERCUSION and the coordination with MAESTRO. HMG, TM, and NRB dropped the majority of sondes that form the BEACH datasets. GG initialized the halodrops repo and HMG, TM, NRB, GG, LK, and TK developed it into the pydropsonde package. HMG and TM prepared the data overview
545 (Section 4) and Figures with input from all other authors. HMG prepared the paper draft with contributions by all other authors.

Competing interests. No competing interests are known.

Acknowledgements. ORCESTRA was made possible thanks to the support of the Max Planck Society (MPG), the European Research Council through Grant No. 101098063, the German Research Foundation (DFG), the German Aerospace Center (DLR), the German Federal Ministry for Education and Research (BMBF), the French National Centre for Scientific Research (CNRS), the Dutch Research Council
550 (NWO), the European Space Agency (ESA), and the US National Science Foundation (NSF), through the Award No. 2331199, 2331202 and 2331200.

We thank Friedhelm Jansen and Lutz Hirsch that they handled the logistics for the sondes; and Felix Ament who provided an additional
50 sondes he found under his desk on Barbados.

We appreciate the help Holger Vömel offered regarding our problematic sondes during the processing.

555 We are grateful to James Ruppert and Allison Wing, because they pointed out a bug in the integrated water vapor calculation in an earlier version of the data, and Basile Pujol for pointing out an inaccuracy in the vertical pressure velocity calculation in JOANNE.

We further credit Jakob Deutloff, Allison Wing, and Marius Winkler who operated the sonde system for one of the flights, as well as Lutz Hirsch, Elina Plesca, Daniel Rowe, Luca Schmidt, and Yuting Wu who got up early in the morning to recondition the sondes.

We also recognize everyone who helped with the communication during one of the flights: Felix Ament, Romain Fievet, Henning Franke,
560 Silke Gross, Suelly Katiza, Daniel Klocke, Brett McKim, Chelsea Nam, Sebastian Ortega Arango, Elina Plesca, Basile Pujol, Chavez Pope, Divya Praturi, Rene Redler, Marius Rixen, Nicolas Rochetin, Daniel Rowe, Martin Singh, Lea Volkmer, Tristan Vostry, and Manfred Wendisch.

And of course Yuting Wu again, who came up with the name for BEACH.

NRB acknowledges support from the ERC starting grant ROTOR (grant no. 101116282)

565 GG acknowledges financial support from HALO DFG SPP 1294 to organize the HALODROPS workshop which kick-started the efforts to develop a dropsonde package that has evolved into the current pydropsonde package.

Table E1. Problematic sondes

Vaisala id (or init time)	Sonde id	Flight Id	problem	procedure
234140294	0ebbb62f	HALO-20240811a	no valid data in L3	removed after L2
231431370	df132c3f	HALO-20240818a	not enough values to interpolate altitude	removed after L2
140238 / 233825141	–	HALO-20240821a	empty A-file empty L1 file	removed after L1
140425 / 234020762	–	HALO-20240821a	empty A-file empty L1 file	removed after L1
140529	–	HALO-20240821a	empty A-file empty D-file	removed after L0
174603 / 233530211	8f96cae2	HALO-20240827a	empty A-file no valid altitude	removed after L2
174847 / 234030059	e77f2e8e	HALO-20240827a	empty A-file no valid altitude	removed after L2
175216 / 234021411	183cf442	HALO-20240827a	empty A-file no valid altitude	removed after L2
175543 / 231820683	672a747e	HALO-20240827a	empty A-file no valid altitude	removed after L2
175913 / 231220385	298e2b48	HALO-20240827a	empty A-file no valid altitude	removed after L2
180136 / 231220384	–	HALO-20240827a	empty A-file empty L1 file	removed after L1
180450 / 234030012	–	HALO-20240827a	empty A-file empty L1 file	removed after L1
180732	–	HALO-20240827a	empty A-file empty D-file	removed after L0
234030131	91823f21	HALO-20240829a	detected as floater	flagged
233824584	10514909 –	HALO-20240914a	serial id appears twice	in L3 no launch detect
233211701	8b593afa 2e760f22	HALO-20240921a	serial id appears twice	in L3 in L3
000007500	0bd0e322 –	HALO-20240924a	serial id appears twice	once in l3 (only winds) no launch detect

Table E2. Minisondes

Vaisala id (or init time)	Sonde id	Flight Id	Reason	procedure
233441164	92afabd1	HALO-20240831a	minisonde	processed with ASPEN minisonde config
233814578	6d733176	HALO-20240831a	minisonde	
233814590	3e5fce09	HALO-20240831a	minisonde	
233814531	df53b31c	HALO-20240831a	minisonde	
233814533	0c1b9857	HALO-20240907a	minisonde	
233814535	ad7d6167	HALO-20240831a	minisonde	
233814536	163b9bd6	HALO-20240831a	minisonde	
233814537	20c81165	HALO-20240906a	minisonde	
233814546	22521c8e	HALO-20240829a	minisonde	
233814577	7097c2fe	HALO-20240831a	minisonde	
233814584	09c7657e	HALO-20240829a	minisonde	
233814586	16fc3f82	HALO-20240829a	minisonde	
233814605	2d623ac0	HALO-20240831a	minisonde	
234141208	ff629abe	HALO-20240829a	minisonde	
234141209	f719c43f	HALO-20240829a	minisonde	
234141210	7dce7693	HALO-20240831a	minisonde	
234141211	7a2b1052	HALO-20240831a	minisonde	

References

- Aberson, S. D., Zhang, J. A., Zawislak, J., Sellwood, K., Rogers, R., and Cione, J. J.: The NCAR GPS dropwindsonde and its impact on hurricane operations and research, *Bulletin of the American Meteorological Society*, 104, E2134–E2154, 2023.
- 570 Akima, H.: A New Method of Interpolation and Smooth Curve Fitting Based on Local Procedures, *J. ACM*, 17, 589–602, <https://doi.org/10.1145/321607.321609>, 1970.
- AspenDocs 1.0: AspenDocs 1.0, https://ncar.github.io/aspendocs/man_qc.html, accessed: 28 May 2025.
- Back, L. and Bretherton, C.: Geographic variability in the export of moist static energy and vertical motion profiles in the tropical Pacific, *Geophysical research letters*, 33, 2006.
- 575 Bellamy, J. C.: Objective calculations of divergence, vertical velocity and vorticity, *Bulletin of the American Meteorological Society*, 30, 45–49, 1949.
- Bernardez, M. and Back, L.: Integrating thermodynamic and dynamic views on the control of the top-heaviness of convection in the Pacific ITCZ with weak temperature gradient simulations, *Journal of Advances in Modeling Earth Systems*, 16, e2022MS003455, 2024.
- Bony, S. and Stevens, B.: Measuring Area-Averaged Vertical Motions with Dropsondes, *Journal of the Atmospheric Sciences*, 76, 767–783, 580 <https://doi.org/10.1175/JAS-D-18-0141.1>, 2019.

- Bony, S., Stevens, B., Ament, F., Bigorre, S., Chazette, P., Crewell, S., Delanoë, J., Emanuel, K., Farrell, D., Flamant, C., et al.: EUREC 4 A: A field campaign to elucidate the couplings between clouds, convection and circulation, *Surveys in Geophysics*, 38, 1529–1568, 2017.
- Brian Eaton, Jonathan Gregory, Bob Drach, Karl Taylor, Steve Hankin, John Caron, Rich Signell, Phil Bentley, Greg Rappa, Heinke Höck, Alison Pamment, Martin Juckes, Martin Raspaud, Randy Horne, Jon Blower, Timothy Whiteaker, David Blodgett, Charlie Zender, Daniel Lee, David Hassell, Alan D. Snow, Tobias Kölling, Dave Allured, Aleksandar Jelenak, Anders Meier Soerensen, Lucile Gaultier, Sylvain Herlédan, Fernando Manzano, Lars Bärning, Christopher Barker, and Sadie Bartholomew: NetCDF Climate and Forecast (CF) Metadata Conventions, <https://doi.org/10.5281/zenodo.14275599>, 2024.
- Ceselski, B. F. and Sapp, L. L.: Objective wind field analysis using line integrals, *Monthly Weather Review*, 103, 89–100, 1975.
- Ehrlich, A., Crewell, S., Herber, A., Klingebiel, M., Lüpkes, C., Mech, M., Becker, S., Borrmann, S., Bozem, H., Buschmann, M., Clemen, H.-C., De La Torre Castro, E., Dorff, H., Dupuy, R., Eppers, O., Ewald, F., George, G., Giez, A., Grawe, S., Gourbeyre, C., Hartmann, J., Jäkel, E., Joppe, P., Jourdan, O., Jurányi, Z., Kirbus, B., Lucke, J., Luebke, A. E., Maahn, M., Mahernndl, N., Mallaun, C., Mayer, J., Mertes, S., Mioche, G., Moser, M., Müller, H., Pörtge, V., Risse, N., Roberts, G., Rosenburg, S., Röttenbacher, J., Schäfer, M., Schaefer, J., Schäfler, A., Schirmacher, I., Schneider, J., Schnitt, S., Stratmann, F., Tatzelt, C., Voigt, C., Walbröl, A., Weber, A., Wetzel, B., Wirth, M., and Wendisch, M.: A comprehensive in-situ and remote sensing data set collected during the HALO–(AC)³ aircraft campaign, *Earth System Science Data Discussions*, 2024, 1–49, <https://doi.org/10.5194/essd-2024-281>, 2024.
- Flohn, H.: Passatzzirkulation und äquatoriale Westwindzone, *Archiv für Meteorologie, Geophysik und Bioklimatologie, Serie B*, 3, 3–15, 1951.
- Fuchs-Stone, Ž., Raymond, D. J., and Sentić, S.: OTREC2019: Convection over the east Pacific and southwest Caribbean, *Geophysical Research Letters*, 47, e2020GL087564, 2020.
- George, G., Stevens, B., Bony, S., Pincus, R., Fairall, C., Schulz, H., Kölling, T., Kalen, Q. T., Klingebiel, M., Konow, H., et al.: JOANNE: Joint dropsonde Observations of the Atmosphere in tropical North Atlantic meso-scale Environments, *Earth System Science Data*, 13, 5253–5272, 2021.
- George, G., Stevens, B., Bony, S., Vogel, R., and Naumann, A. K.: Widespread shallow mesoscale circulations observed in the trades, *Nature Geoscience*, 16, 584–589, 2023.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 0), <https://doi.org/10.83509/bafybeif4n7lov7syd5ragolfosqf6t7zuawexxjhtr5r2ezecvqvj7eje>, 2026a.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 1), <https://doi.org/10.83509/bafybeieqqc5fktpmwblupgb4vidrrx7vv4415e7rol6zp3e3wlbjmq2q4>, 2026b.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 2), <https://doi.org/10.83509/bafybeifi5pplgpcq6onwb3yixhvhgcmenvqmcqi7qgjbrbqlgoyrvj52i>, 2026c.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 3), <https://doi.org/10.83509/bafybeiczbv7mycr2jois6t4dq3zwiltycomwo5xxvjqcz2ot3newzar6q>, 2026d.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 3) QC, <https://doi.org/10.83509/bafybeidytxgujp3evqesvvuvvc3vsllv6pbcu5gc2torqkz7yo6yuazwtmu>, 2026e.
- Gloeckner, H. M., Mieslinger, T., and Robbins-Blanch, N.: BEACH dropsonde dataset (Level 4), <https://doi.org/10.83509/bafybeibgeeqs5uhmbqy4hz4v3pihrfisiklcetisxkf63d2r473szaprwi>, 2026f.

- Hardy, B.: ITS-90 formulations for vapor pressure, frostpoint temperature, dewpoint temperature, and enhancement factors in the range–100 to+ 100 C, in: The proceedings of the third international symposium on Humidity & Moisture, Teddington, London, England, pp. 1–8, 1998.
- 620 Hock, T. F. and Franklin, J. L.: The near gps dropwindsonde, *Bulletin of the American Meteorological Society*, 80, 407–420, 1999.
- Huaman, L., Schumacher, C., and Sobel, A. H.: Assessing the vertical velocity of the East Pacific ITCZ, *Geophysical Research Letters*, 49, e2021GL096192, 2022.
- Johnson, R. H., Rickenbach, T. M., Rutledge, S. A., Ciesielski, P. E., and Schubert, W. H.: Trimodal characteristics of tropical convection, *Journal of climate*, 12, 2397–2418, 1999.
- 625 Konow, H., Ewald, F., George, G., Jacob, M., Klingebiel, M., Kölling, T., Luebke, A. E., Mieslinger, T., Pörtge, V., Radtke, J., et al.: EUREC 4 A's HALO, *Earth System Science Data*, 13, 5545–5563, 2021.
- Lenschow, D. H., Krummel, P. B., and Siems, S. T.: Measuring entrainment, divergence, and vorticity on the mesoscale from aircraft, *Journal of atmospheric and oceanic technology*, 16, 1384–1400, 1999.
- Lenschow, D. H., Savic-Jovicic, V., and Stevens, B.: Divergence and vorticity from aircraft air motion measurements, *Journal of atmospheric and oceanic technology*, 24, 2062–2072, 2007.
- 630 López Carrillo, C. and Raymond, D.: Retrieval of three-dimensional wind fields from Doppler radar data using an efficient two-step approach, *Atmospheric Measurement Techniques*, 4, 2717–2733, 2011.
- Mapes, B. E. and Houze Jr, R. A.: Diabatic divergence profiles in western Pacific mesoscale convective systems, *Journal of Atmospheric Sciences*, 52, 1807–1828, 1995.
- 635 Mapes, B. E., Chung, E. S., Hannah, W. M., Masunaga, H., Wimmers, A. J., and Velden, C. S.: The meandering margin of the meteorological moist tropics, *Geophysical Research Letters*, 45, 1177–1184, 2018.
- Martin, C. and Suhr, I.: NCAR/EOL Atmospheric Sounding Processing ENvironment (ASPEN) software, Version 3.4, 3, <https://www.eol.ucar.edu/content/aspn>, 2021.
- Orlanski, I.: A rational subdivision of scales for atmospheric processes, *Bulletin of the American Meteorological Society*, pp. 527–530, 1975.
- 640 Panofsky, H.: Methods of computing vertical motion in the atmosphere, *J. Meteor*, 3, 45–49, 1946.
- Pincus, R., Fairall, C. W., Bailey, A., Chen, H., Chuang, P. Y., de Boer, G., Feingold, G., Henze, D., Kalen, Q. T., Kazil, J., et al.: Observations from the NOAA P-3 aircraft during ATOMIC, *Earth System Science Data Discussions*, 2021, 1–25, 2021.
- Poujol, B. and Bony, S.: Measuring clear-air vertical motions from space, *AGU Advances*, 5, e2024AV001267, 2024.
- Prange, M., Buehler, S. A., and Brath, M.: How adequately are elevated moist layers represented in reanalysis and satellite observations?, *Atmospheric Chemistry and Physics*, 23, 725–741, 2023.
- 645 Praturi, D. S. and Stevens, B.: On the Meridional Asymmetry of the Poleward-Displaced Intertropical Convergence Zone, *Quarterly Journal of the Royal Meteorological Society*, 152, e70043, <https://doi.org/10.1002/qj.70043>, 2026.
- Reed, R. J. and Recker, E. E.: Structure and properties of synoptic-scale wave disturbances in the equatorial western Pacific, *Journal of Atmospheric Sciences*, 28, 1117–1133, 1971.
- 650 Stevens, B., Bony, S., Gross, S., Klocke, D., Windmiller, J. M., Wing, A. A., von Bismarck, J., Brito, E., David, R. O., Delanoë, J., Farrell, D., and Wu, Y.: Orcestra: Organized convection and earthcare studies over the tropical Atlantic, *Tellus*, <https://doi.org/10.16993/tellus.4123>, 2026.
- Vömel, H. and Goodstein, M.: Dropsonde Data Quality Report: Investigation of Microphysics and Precipitation for Atlantic Coast-Threatening Snowstorms (IMPACTS, 2020) Version 1.0, Tech. rep., UCAR/NCAR - Earth Observing Laboratory, <https://www.earthdata>.

- 655 nasa.gov/s3fs-public/2025-02/2020_impacts_dropsondes_readme_20200514.pdf?VersionId=Kn.ghjdNFkl9wGHZ2a8nsgSrpr5KVti,
2020.
- Vömel, H., Goodstein, M., Tudor, L., Witte, J., Fuchs-Stone, Ž., Sentić, S., Raymond, D., Martinez-Claros, J., Juračić, A., Maithel, V., et al.:
High-resolution in situ observations of atmospheric thermodynamics using dropsondes during the Organization of Tropical East Pacific
Convection (OTREC) field campaign, *Earth System Science Data Discussions*, 2020, 1–25, 2020.
- 660 Wendisch, M., Crewell, S., Ehrlich, A., Herber, A., Kirbus, B., Lüpkes, C., Mech, M., Abel, S. J., Akansu, E. F., Ament, F., et al.: Overview:
quasi-Lagrangian observations of Arctic air mass transformations—introduction and initial results of the HALO–(AC) 3 aircraft campaign,
Atmospheric Chemistry and Physics, 24, 8865–8892, 2024.
- Windmiller, J. M. and Stevens, B.: The inner life of the Atlantic Intertropical Convergence Zone, *Quarterly Journal of the Royal Meteorological
Society*, 150, 523–543, 2024.
- 665 Winkler, M., Rixen, M., Beucher, F., Couvreur, F., Nam, C. C., Peyrillé, P., Schmidt, H., Segura, H., Wieners, K.-H., Alkilani-Brown, E.,
Coly, A. A., Biagioli, G., Bell, M. M., Brito, E., Chauvin, E., Capo, J., Colón-Burgos, D., Dawes, A., da Luz, J. C., Demiralay, Z., Douet,
V., Ducastin, V., Dufaux, C., Dufresne, J.-L., Favot, F., Fiolleau, T., Fons, E., George, G., Gloeckner, H. M., Gonçalves, S., Gouttesoulard,
L., Hayo, L., Hsiao, W.-T., Kennison, S., Kopelman, M., Lee, T.-Y., Le Gall, E., Lothon, M., Lovato, M., Luschen, E., Maury, N., McKim,
B., Netz, L., Ousseynou, D., Peters-von Gehlen, K., Pope, C., Poujol, B., Rivera Maldonado, N., Robbins Blanch, N., Rochetin, N., Rowe,
670 D., Romero Jure, P., Ruppert Jr., J. H., Segura Bermudez, J., Starr, J. C., Stelzner, M., Stoll, C., Syrett, M., Tekoe, A., Trules, J., Welty, C.,
Klocke, D., Vogel, R., Bony, S., Wing, A. A., and Stevens, B.: RAPSODI: radiosonde atmospheric profiles from ship and island platforms
during ORCESTRAS, collected to Decipher the ITCZ, *Earth System Science Data*, 18, 1833–1854, 2026.
- Yanai, M.: A Detailed Analysis of Typhoon Formation, *Journal of the Meteorological Society of Japan. Ser. II*, 39, 187–214,
https://doi.org/10.2151/jmsj1923.39.4_187, 1961.
- 675 Yanai, M., Esbensen, S., and Chu, J.-H.: Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture
budgets, *Journal of Atmospheric Sciences*, 30, 611–627, 1973.