

SO SLICE: a 19-year gridded sea level anomaly dataset for the open and ice-covered Southern Ocean (2003–2021)

Cosme Mosneron Dupin¹, Jean-Baptiste Sallée¹, Casimir de Lavergne¹

¹Sorbonne Université, CNRS, IRD, MNHN, Laboratoire d'Océanographie et du Climat : Expérimentations et Approches Numériques, LOCEAN/IPSL, F-75005 Paris, France

Correspondence to: Cosme Mosneron Dupin (cosme.mosneron-dupin@locean.ipl.fr)

Abstract. Sea Level Anomalies (SLA) in the Southern Ocean remain poorly observed due to sea ice, which hampers conventional satellite altimetry. Recent advances in lead-based retracking techniques have enabled SLA estimates within ice-covered regions. However, existing datasets are temporally limited, with most available products covering periods shorter than 10 years. Here, we present SO SLICE (Southern Ocean Sea Level in the ICE), a gridded SLA dataset spanning both ice-free and ice-covered areas of the Southern Ocean from 2003 to 2021. This 19-year dataset provides the longest temporal SLA coverage currently available across the whole Southern Ocean. It is obtained by processing along-track measurements from four satellite missions: Envisat, CryoSat-2, SARAL/AltiKa, and Sentinel-3A. It relies on retrieving observations from the openings in the sea ice cover. The product shows continuous transitions across the sea ice edge, with a mean formal mapping error near 1.3 cm in the subpolar Southern Ocean. The gridded fields have an effective spatial resolution of approximately 200 km south of 65°S, mainly constrained by the input correlation length scales. It captures variability at periods longer than 30 days. Comparison with independent in situ observations yields mean correlation coefficients of 0.58 with tide gauges and 0.66 with bottom pressure recorders. Overall, SO SLICE fills a key observational gap and offers new opportunities to study trends and interannual variability in sea level and ocean circulation in the Southern Ocean, particularly under sea-ice cover.

1 Introduction

The Southern Ocean plays a pivotal role in the climate system. It stores large amounts of anthropogenic heat and carbon (Frölicher et al., 2015; Meredith et al., 2019; Sallée, 2018; The SO-CHIC consortium et al., 2023), ventilates more than half of the global ocean volume (Millet et al., 2024), and enables interbasin exchanges. However, it has undergone important changes over the last several decades. The Southern Ocean has been warming and freshening (Sallée, 2018; The SO-CHIC consortium et al., 2023). Westerly winds that force its circulation have intensified and shifted towards the poles (Thompson et al., 2011). Antarctic ice shelves are experiencing accelerated basal melting, increasing freshwater fluxes into the ocean, and contributing to sea level rise (Holland et al., 2020; The IMBIE team, 2018). In 2016, sea-ice extent reversed its decades-long increasing trend, giving way to a rapid decline in its observed area coverage (Hobbs et al., 2024). This suggests that the entire Southern Ocean might be experiencing a regime shift (Abram et al., 2025; Silvano et al., 2025).

a supprimé: Reconstructing Sea Level Anomalies in the open and ice-covered Southern Ocean from 2003 to 2021

- a supprimé: Antarctic Sea Level Anomalies (SLA)
- a supprimé: the presence of sea ice
- a supprimé: estimation
- a supprimé: products
- a supprimé: currently
- a supprimé: dataset
- a supprimé: In this study, we
- a supprimé: extend the time span of SLA reconstruction
- a supprimé: multiple
- a supprimé: k
- a supprimé: We produce a consistent and continuous SLA dataset spanning both ice-free and ice-covered areas of the Southern Ocean (south of 50°S) from 2003 to 2021. This 19-year product provides the longest temporal coverage to date for SLA under sea ice.
- a supprimé: The resulting SLA field resolves the large-scale variability and parts of the mesoscale signal, with smooth transitions across the sea-ice edge.
- a supprimé: We show that the product reliably captures physical signals on timescales longer than 10 days, with estimated uncertainties of 1.3 cm. in the subpolar ocean.
- a supprimé: Our satellite-based SLA reconstruction compares well with most independent in situ observations from tide gauges and bottom pressure recorders: mean correlation coefficients are 0.58 and 0.66, respectively.
- a supprimé: The reconstruction
- a supprimé:
- a supprimé: uptakes and
- a supprimé: in the past decades
- a supprimé: and ice sheet discharge
- a supprimé: , ultimately
- a supprimé: a regime shift in
- a supprimé: leading
- a supprimé: and reaching the lowest observed area coverage in 2023...

70 Despite these profound changes, interannual variability of ocean circulation remains largely uncharacterized in the ice-covered Southern Ocean, owing to observational challenges affecting both in-situ and satellite measurements (Newman et al., 2019; Smith et al., 2019). A key variable for tracking these changes is the Sea Level Anomalies (SLA) field, which represents sea level variations relative to a reference surface. Through geostrophic balance, the SLA field enables computation of anomalies in large-scale ocean currents. It also captures the combined effects of density-driven and mass-driven contributions on sea level, making it an integrated measure of change in the ocean. For the past thirty years, radar altimetry has enabled SLA

75 retrievals in the open ocean, revolutionizing the understanding of ocean dynamics and ocean state changes (Fu and Cazenave, 2000; Morrow and Le Traon, 2012). However, ocean altimetric processing degrades significantly in the presence of sea ice, which contaminates the measured signal. As a result, SLA retrievals in the subpolar Southern Ocean and the Arctic Ocean have historically been sparse and unreliable. Overcoming this limitation requires recovering ocean signal when the satellite overflies sea ice. In practice, this amounts to measuring SLA within the openings in the ice cover. Notable advances have

80 progressively made this possible. Drinkwater (1991) demonstrated that radar returns from leads, i.e. fractures between the ice floes where the ocean is directly visible, are highly specular. This makes their signatures readily distinguishable within the sea ice background. Laxon (1994) was the first to suggest using signal from leads. This idea was operationalized for oceanographic studies a decade later. Peacock and Laxon (2004) successfully measured SLA in leads and derived the first Mean Sea Surface (MSS) and a two-year record of SLA variability in the Arctic Ocean.

85 During the 2010s, most developments in ocean polar altimetry focused on the Arctic. Giles et al. (2012) measured sea level in leads from the missions ERS-1/-2 and Envisat to quantify freshwater accumulation in the Beaufort gyre, linking it to wind-driven convergence. Kwok and Morison (2011) used laser altimeter observations from ICESat to map the Arctic Dynamic Ocean Topography (DOT) in the winters of 2004 to 2008. They restricted the study to the ice-covered domain and highlighted

90 the importance of the baroclinic circulation in the Arctic Ocean. Bulczak et al. (2015) investigated Sea Surface Height (SSH) variability from 2003 to 2009 in the Nordic Seas using lead data from Envisat. These efforts culminated in basin-wide, multi-mission products. Armitage et al. (2016) produced the first publicly available monthly DOT estimates for the entire Arctic Ocean from Envisat and CryoSat-2 (2003–2014). Rose et al. (2019) extended the record back to 1991 by integrating ERS-1 and ERS-2. Doglioni et al. (2023) focused on resolving the variability of boundary currents in the Arctic at scales of 70 km with CryoSat-2 (2011–2020).

In contrast, efforts in the Southern Ocean have been more limited. Early studies focused on either the ice-covered season (e.g., Kwok and Morison, 2016) or the open-water season (e.g., Rye et al., 2014), and left the full seasonal cycle poorly characterized. A significant advance was made by Armitage et al. (2018), who extended their Arctic processing to the Southern Ocean. They

100 produced the first maps of SSH covering both the ice-covered and ice-free Southern Ocean, from 2011 to 2016. Their dataset revealed coherent seasonal variability and highlighted the influence of wind on large-scale features such as the Antarctic Slope

a supprimé:

a supprimé: ocean circulation...ariability of ocean circulation remains poorly characterized ...argely uncharacterized in the ice-covered Southern Ocean, owing due...to observational challenges affecting both in-situ and satellite measurements (Newman et al., 2019; Smith et al., 2019). A key variable for tracking these changes is the Sea Level Anomalies (SLA) field, which represents sea level variations across ...relative to a reference surface. Through equilibrium...alance, the SLA field enables computation of in ...arge-scale ocean currents. It also captures the combined effects of density-driven and mass-driven contributions on sea level, making it an integrated measure of change in the oceanlt also captures both density-driven (steric) and mass-driven (barystatic) components of sea level variability, providing insight into the ocean vertical structure and mass redistribution... Observing SLA therefore allows derivation of both structural and dynamical ocean changes. ...or the past thirty years, satellite ...adar altimetry has enabled SLA retrievals in the open ocean, revolutionizing oceanographic ...he understanding of circulation ...cean dynamics and ocean state changes (Fu and Cazenave, 2000; Morrow and Le Traon, 2012). However, this technique has remained largely ineffective in the subpolar Southern Ocean, as standard processing degrades significantly in the presence of sea ice. As a result, SLA retrievals in polar regions, especially in ice-covered waters, have historically been sparse or unreliable (Prandi et al., 2012).

... [1]

a mis en forme : Anglais (E.U.)

a supprimé:

Notable advances have progressively enabled the use of radar altimetry in ice-covered regions. Initial efforts date back to Dwyer and Godin (1980), who employed data from the GEOS-3 satellite to identify sea-ice boundaries and infer ice roughness from space. Subsequently, Drinkwater (1991) demonstrated that radar returns from leads—open-water holes within the ice pack that are flat and therefore extremely reflective—are highly specular. These distinct returns make lead signatures readily identifiable in a sea-ice covered area. Laxon (1994) advanced altimetry processing over sea-ice. Crucially, he was the first to suggest using returns from leads, not to estimate sea surface topography but to measure freeboard and infer sea-ice thickness. Nonetheless, it laid the groundwork for an eventual application of lead returns for sea level estimation. This concept was operationalized a decade later. Peacock and Laxon (2004) successfully measured SLA in leads and derived the first Mean Sea Surface (MSS) and a two-year record of SLA variability under Arctic sea ice.

a supprimé:

a mis en forme

a supprimé: Armitage et al. (2016) produced the first publicly available monthly estimates of Dynamic Ocean Topography (DOT) for the whole Arctic Ocean, using Envisat and CryoSat-2 data from 2003-2012. (Doglioni et al., 2023)

a supprimé:

... [4]

Current and the Weddell Gyre. Leveraging this product, Dotto et al. (2018) uncovered the seasonal cycle of the Ross gyre, while Naveira Garabato et al. (2019) linked circumpolar SSH variability to wind forcing modulated by sea-ice conditions. The most comprehensive SLA product to date was developed by Auger et al. (2022a), combining three altimetry missions: CryoSat-2, SARAL/AltiKa (referred to as SARAL in the rest of the article) and Sentinel-3A. It allows for an improved spatial resolution and spans from 2013 to 2019. This dataset enabled further investigation of seasonal variability in the subpolar Southern Ocean (Auger et al., 2022b) as well as mesoscale circulation features (Auger et al., 2023). It was recently extended to 2021 by Veillard et al. (2024). Dragomir (2024) further expanded the temporal baseline of Armiatge et al. (2018) by incorporating Envisat data, producing an SLA record spanning 2003–2018.

a supprimé: used a similar approach to uncover...ncovered the seasonal cycle of the Ross gyre, while Naveira Garabato et al. (2019) linked circumpolar SSH variability to wind forcing modulated by sea-ice conditions. The most comprehensive SLA product to date was developed by Auger et al. (2022a). It combines..., combining three altimetry missions: (...ryos...at-2, Saral...ARAL/AltiKa (referred to as SARAL in the rest of the article) and...Sentinel-3A)...It allows for anto...improved spatial resolution and spans from 2013 to 2019. It was used...his dataset enabled to ...urther investigation ofe the...seasonal variability of ...n the subpolar Southern Ocean (Auger et al., 2022b) as well as the small...esoscale...scale...irculation circulation ...eatures (Auger et al., 2023). This dataset...it was recently extended to 2021 by Veillard et al. (2024). (...ragom... [5]

a mis en forme ... [6]

While these studies have significantly advanced the understanding of the subpolar Southern Ocean, they often focused on limited domains, either ice-free or ice-covered, or relied on short temporal records. Here, we present SO SLICE (Southern Ocean Sea Level in the ICE), a 19-year SLA dataset covering the period 2003–2021. SO SLICE relies on the core principle of extracting SLA from radar returns in leads. It builds upon the products of Auger et al. (2022) and Veillard et al. (2024), using similar mapping procedure and processing framework. As in Dragomir (2024), it incorporates Envisat data to extend the temporal baseline back to 2003. Overall, SO SLICE integrates observations from four satellite missions: Envisat, CryoSat-2, SARAL, and Sentinel-3A. It is a publicly available product documented here and provides consistent, spatially continuous coverage of the Southern Ocean. Compared to earlier records, its main contributions are the length of the time-series which enables the study of interannual-to-decadal variability in the Southern Ocean, and the consistency of the multi-mission processing across two decades. In addition to describing SO SLICE and its methods, this paper provides a thorough characterization of its effective resolution and uncertainty. Existing altimetric SLA products in the Southern Ocean regions have generally been validated through limited in-situ comparisons, and a systematic assessment of their resolved scales and error has been lacking. We address this gap using the spectral validation framework of Ballarotta et al. (2019). Section 2 describes the satellite and in situ data used in this study. Section 3 details the sea-ice altimetry processing used to construct the dataset. Results and validation are presented in Sect. 4, and our findings are summarized in Sect. 5.

a supprimé: our ...he understanding of the subpolar Southern Ocean dynamics... they often focused on limited domains, (...ither ice-free or ice-covered,) ...r relied on short temporal records. Here, we present SO SLICE (Southern Ocean Sea Level in the ICE), a 19-year SLA dataset covering the period 2003–2021. build upon...O SLICE relies on the the ...ore principle of extracting SLA from radar returns in sea-ice ...eads to present the most extensive SLA dataset yet produced for the Southern Ocean... It builds upon products of (...uger et al. (...022) and...Veillard et al. (...024), using similar mapping procedure and processing framework. As in Dragomir (2024), it incorporates Envisat data to extend the temporal baseline back to 2003. Overall, . This new product...O SLICE integrates observations from four satellite altimetry ...issions: — Envisat, CryoSS...t-2, Saral...ARAL...and Sentinel-3A. It is —... publicly available product documented here and to ...rovides consistent, spatially continuous coverage of the Southern Ocean from from 2003 to 2021... Compared to earlier records, its main contributions are the length of the time-series Spanning 19 years, it...hich offers unprecedented potential for...nables the study of analyzing long-term trends and ...nterannual-to-decadal variability in the Southern Ocean, and the consistency of the multi-mission processing across two decades sea level and circulation... In addition to describing SO SLICE and its methods, this paper provides a thorough characterization of its effective resolution and uncertainty. Existing altimetric SLA products in the Southern Ocean regions have generally been validated through limited in-situ comparisons, and a systematic assessment of their resolved scales and error has been lacking. We address this gap using the spectral validation fram... [7]

2 Data

2.1 Satellite data

SO SLICE is derived from four satellite missions: Envisat (2002–2012), CryoSat-2 (2011–2021), SARAL (2013–2021), and Sentinel-3A (2016–2021). The starting and ending dates of each mission integration are indicated in Fig. 1. Table 1 summarizes their key characteristics, including orbital parameters, radar frequency, and acquisition mode.

a supprimé: Four satellite missions are integrated into the dataset...O SLICE is derived from four satellite missions: Envisat (2002–2012), Cryosat...ryoSat-2 (2011–2021), Sentinel-3A and Saral...ARAL (2013–2021), and Sentinel-3A (2016–2021). The product starts on the 1st of January 2003 (Fig. 1) with observations from Envisat. Cryosat-2 observations are used from 16/07/2010 (DD/MM/YYYY) onwards, allowing for a short overlap before Envisat stops acquisitions on 07/04/2012. Saral and Sentinel-3A are assimilated in the product starting 14/03/2013 and 01/07/2016, respectively. Periods of single mission coverage - spanning from 01/01/2003 to 16/07/2010 and 07/04/2012 to 14/03/2013...he starting and ending dates of each mission integration are indicated in F... [8]

Envisat was launched in March 2002. Its altimeter, RA-2, operates in Low Resolution Mode (LRM), at a nominal frequency of 13.575 GHz (Ku band). It has a beam-limited footprint of 18 km diameter (ESA, 2018) and a pulse-limited footprint of 1.7 km

a supprimé: Satellite missions and their associated altimeters have key differences (see Table 1) which require a dedicated processing for each. Envisat...ts altimeter, RA-2, operates in Low Resolution Mode (LRM), in the Ku frequency band (13 to 17 GHz...at a nominal frequency of 13.575 GHz (Ku band). It z). It ...as a surface ...eam-limited footprint of roughly ...85...km diameter (Anon...SA, 2018n.d.... [9]

diameter (Connor et al., 2009). Envisat data from the Fundamental Data Records for Altimetry project (European Space Agency, 2023) are used at the 18Hz frequency (~370 m along-track posting).

SARAL was launched in February 2013. Its altimeter, AltiKa, also operates in LRM but at a nominal frequency of 35.75 GHz (Ka band). This, coupled to a larger chirp bandwidth, yields smaller beam-limited and pulse-limited footprints (respectively 9 km and 1.4 km in diameter) than Envisat (Verron et al., 2021). The SARAL data come from the CNES PEACHI project (Valladeau et al., 2015), using the outputs at 40 Hz (~175 m along-track posting).

CryoSat-2 was launched in April 2010. Its altimeter SIRAL operates in three modes: LRM above the open ocean, Synthetic Aperture Radar Mode (SARM) above the sea ice and Synthetic Aperture Interferometric Mode (SARInM) over the land ice, the ice-sheet margins and parts of the coastal Southern Ocean (Wingham et al., 2006). The acquisition mask can be found in the CryoSat-2 product handbook (ESA, 2021). Only LRM and SARM data are used here. The SARM data are L1b data from ESA PDGS Ice Baseline E processor (ESA, 2021; Meloni et al., 2020), with measurements at 20 Hz (~350 m along-track posting). It includes zero padding and Hamming windowing which improves the SLA retrieval over specular surfaces such as leads (Quartly et al., 2019; Smith and Scharroo, 2015). Compared to the previous missions, CryoSat-2 has a longer repeat cycle of 369 days but a much finer footprint, yielding denser spatial coverage than repeat-track missions at the expense of temporal regularity.

Sentinel-3A was launched in February 2016. Its altimeter SRAL operates in SAR mode over both sea-ice covered and open ocean (Donlon et al., 2012). Data come from the S3A Processing Prototype (S3PP) with measurements at 20 Hz (~350 m along-track posting). It is an experimental SAR processing chain produced by the Centre National d'Etudes Spatiales, the French space agency, building on the SAR processing developments for CryoSat-2 described in Boy et al. (2017). S3PP applies zero-padding and Hamming windowing to the SAR waveforms, as in the CryoSat-2 Ice Baseline-E processing. More recent L2 Sentinel 3-A dataset exists, such as the ESA thematic LAN SI product, but the choice was made to use the L1 data from S3PP to apply the same classifier and retracking as CryoSat-2.

All four missions share the same fundamental measurement principle. The altimeters described above measure the range, i.e. the distance between the satellite antenna phase center and the surface. They do so by emitting short radar pulses towards the surface and recording the reflected echoes (Fig. 2a, top). These return signals, known as waveforms, represent the backscattered power as a function of time delay (Fig. 2a, bottom). The range is estimated from the waveform shape using dedicated algorithms called retrackers. Along-track SSHs are then derived from each mission following the standard altimetric equation (1):

$$SSH = H - R - \sum H_{corr}$$

- a supprimé: We
- a supprimé: (
- a supprimé: use its measurements sampled
- a supprimé: at a rate of 18 Hz, giving a data point every 370 m
- a supprimé: (Abileah et al., 2013). Saral
- a supprimé: Low Resolution Mode,
- a supprimé: in the Ka frequency band (35.75 GHz).
- a supprimé: allows for a smaller
- a supprimé: and a higher sampling frequency compared to Envisat (respectively 8 km and 40 Hz)
- a supprimé: (Verron et al., 2021)
- a supprimé:
- a supprimé: s
- a supprimé: 2
- a supprimé: LRM
- a supprimé: (in this paper we refer to the region not seas (... [10])
- a supprimé: SARM (
- a supprimé: SARInM (
- a supprimé: on
- a supprimé: and
- a supprimé: W
- a supprimé: (Wingham et al., 2006)
- a supprimé: . In the present paper
- a supprimé: ,
- a supprimé: is
- a supprimé: from Cryosat-2
- a supprimé: .
- a supprimé: (Prandi et al., 2021)
- a supprimé: SARM allows for a better effective along-tra (... [11])
- a supprimé: also
- a supprimé: M
- a supprimé: but
- a supprimé: (reaching an effective resolution of 300 m.
- a supprimé: (
- a supprimé: ,
- a supprimé:
- a supprimé: We use data processed and distributed by the (... [12])
- a mis en forme (... [13])
- a supprimé: Altitude
- a supprimé: ange

Where H is the altitude of the satellite, R is the range estimated from the radar waveform, and H_{corr} accounts for geophysical corrections. For a comprehensive treatment of the altimetric measurement principle and correction models, the reader is referred to Fu and Cazenave (2001). SLA are then obtained by removing a Mean Sea Surface (MSS) from the SSH:

$$SLA = SSH - MSS$$

The applied corrections and MSS used are consistent across all missions and are listed in Table 2.




Figure 1: Satellite missions timeline, with the start and end dates of data merging into SO SLICE indicated as colored text (dd/MM/YYYY).

Table 1. Satellite missions used in SO SLICE and their characteristics.

Satellite	Mode over open-ocean	Mode over sea-ice	Retracker for leads	Sampling rate	Inclination	Cycle duration
Envisat	LRM	LRM	Adaptive	18 Hz	99°	35
CryoSat-2	LRM	SARM	TFMRA	20 Hz	92°	369 (30-day pseudo cycle)
SARAL	LRM	LRM	Adaptive	40 Hz	99°	35
Sentinel-3A	SARM	SARM	TFMRA	20 Hz	99°	27

Table 2. Summary of geophysical corrections applied on altimeter measurements and used for SLA retrieval.

Orbit	POE-E (Ollivier et al., 2015)
Ocean Tide	FES22B (Carrère et al., 2022)
Polar Tide	Desai et al. (2015)
Earth Tide	Cartwright and Edden (1973)
Dry tropospheric	Model from ECMWF
Wet tropospheric	Model from ECMWF
Ionospheric correction	GIM (Iijima et al., 1999)
Dynamical Atmosphere correction	MOG2D (Carrère and Lyard, 2003)
Sea State Bias	Non-parametric (Tran et al., 2010), only for open ocean
MSS	CNESCLS22_Hybrid (Laloue et al., 2024)

a mis en forme le tableau

a mis en forme : Anglais (E.U.)

2.2 Ancillary and validation data

a supprimé: ¶

a supprimé: Open ocean satellite data

Several independent datasets are used in this study for cross-calibration, validation, and comparison purposes.

2.2.1 C3S open ocean data

a mis en forme : Police :Non Gras, Police de script complexe :Non Gras

a mis en forme : Espace Avant : 12 pt, Après : 12 pt, Interligne : simple

a supprimé: To validate our product in the open ocean, we use the Sea Level Anomalies from DUACS, available on the Copernicus Marine Environment Monitoring Service (CMEMS)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a supprimé: {Citation}

a mis en forme : Police :Gras, Police de script complexe :Gras

a mis en forme : Police :Gras, Police de script complexe :Gras, Anglais (E.U.)

a mis en forme : Police :Gras, Police de script complexe :Gras

a mis en forme : Espace Avant : 12 pt, Après : 12 pt, Interligne : simple

a mis en forme : Anglais (E.U.)

2.2.2 Other altimetry datasets covering the ice-covered regions

Two other altimetric SLA products covering the ice-covered Southern Ocean are used for intercomparison. Dragomir (2024) is an SSH dataset spanning 2003–2018 by combining Envisat and CryoSat-2 data, though the record relies on a single mission for most of its duration (Envisat only before 2010, CryoSat-2 only after 2012). Auger et al. (2022a) produced a multi-mission product spanning 2013–2019 by combining CryoSat-2, SARAL, and Sentinel-3A. Comparing with both allows for the evaluation of SO_SLICE in ice-covered regions where C3S is not always available (Sect. 4.3.4).

690 **2.2.3 In-situ data**

Two types of in-situ observations are used for validation: tide gauges and bottom pressure recorders (BPRs). All platforms presented here were deployed in ocean regions with a seasonal or permanent sea-ice cover. Deployment dates, geographic coordinates, and distance to coast are specified in the supplementary Table S1, and location are shown in Fig. 10. Hourly tide gauge records from stations on the Antarctic coast are taken from the Gloss/Clivar dataset (IOC, 2020). They have been corrected for Dynamical Atmospheric Correction (Carrère and Lyard, 2003), ocean tide (Carrère et al., 2022) and glacial isostatic adjustment (Peltier, 2004). BPRs record pressure at the seafloor, providing a proxy for the variations of the ocean mass above the instrument. Vinogradova et al. (2007) demonstrated with a modeling study that bottom pressure variations south of 60°S are significantly correlated to SLA on periods shorter than 60 days. Six BPRs are used. Four are in the Drake Passage - referred to as Drake Passage South (DPS), Drake Passage South Deep (DPSDeep), Myrtle B and Myrtle C (PSML, 2024). The other two are ANT-17.1, deployed on the Greenwich meridian (Androsov et al., 2020), and the JARE BPR, deployed near Syowa station (Hayakawa et al., 2012). Figure 10a shows the location of all these in-situ sources.

700





Figure 2: (a, top) Schematic of a satellite altimeter operating over the open ocean, recording a surface echo (i.e. waveform) characteristic of an open ocean surface (a, bottom). (b, top) Configuration with the same altimeter over sea ice, passing above a lead. (b, bottom) Typical specular waveform reflected by a lead. The retracking algorithm estimates the track point and derives the epoch (red point), i.e. the two-way travel time of the radar pulse, from which the range R, i.e. the distance between the satellite antenna and the surface, can be derived.

705

a supprimé: 1
a supprimé: 3

a supprimé: and ...eographic coordinates...ordinates, and distance to coast are specified in the supplementary Table S1, and location are shown in Fig. 10. Hourly tide gauge records from stations on the Antarctic coast are taken from the Gloss/Clivar dataset (IOC, 2020). They have been corrected for Dynamical Atmospheric Correction (Carrère and Lyard, 2003), ocean tide (Carrère...e et al., 2022) and glacial isostatic adjustment (Peltier, 2004). BPRs record pressure at the seafloor, providing a proxy for the variations of the ocean mass above the instrument. According to ...inogradova et al. (2007) demonstrated with a modeling study that bottom , these pressure variations south of 60°S are significantly correlated to sea level changes on timescales up to 100 days globally, and...LA on periods shorter than 60 days at high latitudes... Six BPRs are used. Four are located

... [14]



a supprimé:

a supprimé: Principle of satellite altimetry. ...a, upper...op) Schematic of the ... configuration ...atellite altimeter operating with a satellite operating above...ver the open ocean, with recording a surface echo (i.e. waveform) typical ...haracteristic of of an open ...cean echoes...urface (a, bottom). (a bottom) The waveform is the surface echo power recorded by the satellite as a function of time. ...b, top upper... Schematic of the configuration...onfiguration with the same satellite ...ltimeer operating ...ver the ...ea ...ce domain... passing just ...bove a lead. (b, bottom) Typical specular waveform of the echo reflected by...

... [15]

phase centre and the sea surface, is derived. SSH is the satellite altitude above a reference ellipsoid minus the range. Sea Level Anomaly (SLA) is the difference between SSH and Mean Sea Surface (MSS).

3 Methods

To retrieve SLA, the standard altimetric processing described in Section 2.1. assumes the radar signal is reflected by the ocean surface, as in Fig. 2a configuration. In ice-covered regions, this assumption breaks down. The altimeter footprint is predominantly occupied by sea ice. Measuring SLA in these conditions requires identifying locations where the ocean surface is directly visible, i.e. leads, and applying dedicated waveform retracking to estimate the associated range and SLA. This section describes the processing steps that allow SO_SLICE to integrate SLA measurements in ice-covered conditions. Figure 3 recapitulates the whole data pipeline.

3.1 Lead detection, retracking and editing

The first step of the processing chain in sea-ice regions is to identify measurements acquired within the ice-covered regions that correspond to leads rather than ice. As the ocean surface within leads is sheltered from winds and waves, its surface is extremely smooth and acts as a quasi-specular reflector for the radar pulse. Thus, leads produce highly peaked returns associated with high backscattering coefficients. Lead signatures dominate the waveform, even if they cover a small fraction of the footprint (Drinkwater, 1991). This configuration is shown in Fig. 2b. Based on this property, a neural-network classifier (Longép   et al., 2019; Poisson et al., 2018) is used to automatically identify lead waveforms and discard sea ice returns. The classifier categorizes each waveform into one of sixteen classes according to its shape. Lead echoes have their dedicated class. It is combined with a multiple criteria classification.

Once a waveform is classified as a lead waveform, a retracking algorithm is applied to derive the range. For CryoSat-2 and Sentinel-3A, which operate in SAR mode above sea ice, the TFMRA empirical retracker is applied on lead waveforms (Helm et al., 2014). For their open ocean waveforms, a different retracker, MLE4, is used. Sentinel-3A operates in SAR mode above the open ocean, but its waveforms are converted to pseudo LRM waveforms, making the use of MLE4 possible. Since two different retrackers are applied for the same mission, a bias between open-ocean and lead SLA arises and must be corrected (Sect. 3.2). For the LRM missions, Envisat and SARAL, the Adaptive retracker is used (Poisson et al., 2018; Tourain et al., 2021). It handles both open-ocean and lead waveforms within a single physical model. Theoretically, it should allow for the merging of open ocean and lead data without needing to correct for a bias. In practice, it reduces the bias between open ocean SLA and lead SLA by about 50 centimeters, but a residual offset remains, which needs to be removed.

- a supprim  : SSH
- a supprim  : the difference between
- a supprim  : Altitude and Range.Sea
- a supprim  : Sea Surface Height (
- a supprim  :)
- a supprim  : tate
- a supprim  :). SSH is the difference between Altitude and Range.




Figure 3: Data processing pipeline of SO SLICE, from the L1b waveforms to the final gridded product. Each key step has its associated box and is discussed in the referred subsection.

The altimetric equation (1) is then applied to retrieve the SLA, with the same set of geophysical corrections and MSS as for the open ocean (Table 2). The only difference is that Sea State Bias (SSB) is not corrected for, under the assumption that wind and wave effects on the SSB are negligible in the ice-covered domain. This hypothesis should hold except in the marginal ice zone. Sea ice inhibits wave propagation coming from the open ocean (Squire, 2020), and protects surface water from wind generated ripples. If high frequency waves are efficiently dissipated by sea ice, low frequency waves can propagate within the sea ice cover and may be a source of bias to our product. Unfortunately, it is hard to correct for as no reliable SSB estimates currently exist for specular waveforms. SSB is empirically derived from significant wave height (SWH) and wind speed (Tran

a supprimé: Satellite altimetry is a well-established technique for measuring sea level variations on a large scale. It relies on emitting radio waves from a satellite and recording the echoes reflected by the surface (Fig. 2, top). These return signals, known as waveforms, represent the echo power measured by the satellite as a function of time (Fig. 2, bottom). By analyzing the shape and timing of the waveform—a process called retracking—key geophysical parameters such as sea surface height (SSH) can be derived (Fu and Cazenave, 2000).

a supprimé: We use SSH data collected by the four satellite missions described above, both when sampling the open ocean and the sea-ice sector (Fig. 2a, 2b). Advances in processing techniques now enable the retrieval of the SSH in leads, i.e. narrow openings of water between sea-ice floes (Fig. 2b). Altimetric data from both leads and open-ocean collected over the span of 19 years are combined and mapped. This ultimately yields a Sea Level Anomaly (SLA) product consistent across the open ocean and ice-covered regions of the Southern Ocean. The sections below describe the methodology used in the open-ocean and ice-covered domain.

where ΣH_{corr} is an ensemble of geophysical corrections, listed in Table 2. It is the same set of corrections used and described in Veillard et al. (2024). These corrections take out certain geophysical contributions to the sea surface height that are not relevant for the study of oceanic dynamic variability and can alias the observed signal. They also correct for processes that modify the propagation speed of the radar wave through the atmosphere (dry tropospheric, wet tropospheric, ionospheric corrections). The physical variable of interest for our product is the SLA. It is obtained by subtracting the Mean Sea State (MSS) to the SSH:

The MSS used comes from Laloue et al. (2024). It combines three different MSS estimates with the aim of keeping the best attr...

a supprimé: Retracking algorithms are then applied to these lead waveforms to get the local SLA. For Cryosat-2 and Sentinel-3A, an empirical retracker called TFMRA (Helm et al., 2014) is used

a supprimé: Table 2. Summary of geophysical corrections applied on altimeter measurements and used for SLA retrieval.

a supprimé: We then

a supprimé: y

a supprimé: the same

a supprimé: we used

a supprimé: data

a supprimé: points

a supprimé: A key

a supprimé: well

a supprimé: (Squire, 2020), because

a supprimé: s

a supprimé: ,

et al., 2010), but SWH is not reliably retrieved from specular lead waveforms (Poisson et al., 2018). Hence, all existing polar altimetric SLA products similarly neglect SSB in the ice-covered domain (e.g., Armitage et al., 2016; Auger et al., 2022a).

After classification and retracking, the along-track SLA records still contain outliers. A series of editing steps is used to remove them. This editing is consistent across all four missions and similar to the one applied in Auger et al. (2022a) and Veillard et al. (2024). First, an empirical threshold based on sea ice concentration, backscatter coefficient and significant wave height is applied. Its aim is to discard measurements whose characteristics are inconsistent with a reliable ocean or lead return but were not removed by the classification step. Second, an iterative editing is applied to the open ocean data only. Along-track SLA are filtered with a 124-point low-pass Lanczos filter, and outliers are determined as points such that $|SLA - SLA_{filtered}| > 2 \cdot \text{std}(SLA - SLA_{filtered})$. This process is iterated until outliers represent less than 0.1% of the measurement points. Third, a dedicated filter following Poisson et al. (2018) is applied to lead data to mitigate the off-nadir hooking (or snagging) effect. As leads are highly specular and associated with a high backscatter coefficient, their returns can dominate the waveform even when the lead is located off-nadir but within the beam-limited footprint. In such cases, the measured range is longer than the true nadir range, introducing a negative bias in the estimated SLA (Armitage and Davidson, 2014; Quartly et al., 2019). This correction is only applied to the LRM altimeters (mounted on Envisat and SARAL) as altimeters operating in SAR mode are much less affected by the hooking effect due to their smaller footprint.

Figure S3 shows the number of lead measurements per $100 \text{ km} \times 100 \text{ km} \times 1\text{-month}$ bin before and after editing. For Envisat, the median number of lead observations per bin decreases from approximately 300 to 35 after editing, with similar values for SARAL. The hooking filter accounts for the largest fraction of removed points, making the editing particularly conservative for the LRM altimeters. The hooking filter is not applied to CryoSat-2 and Sentinel-3A, which consequently retain a higher fraction of lead measurements, at around 200 observations per bin after editing. Despite this substantial reduction for LRM missions, the remaining lead observation density is comparable to the open-ocean observation density or higher in the case of SAR altimeters. This is because the open-ocean along-track data are down-sampled to 5 Hz prior to interpolation to speed up the interpolation.

3.2 Ocean/Leads bias correction

For the SARM missions (CryoSat-2 and Sentinel-3A), different retrackers are applied to open ocean and lead waveforms. This induces a systematic offset between the two domains which needs to be corrected. To address this, we apply the ocean/lead (O/L) bias correction introduced by Giles et al. (2012). For each mission, SLA estimates from the two surface types are averaged into separate $75 \text{ km} \times 75 \text{ km} \times 10\text{-day}$ bins. The difference in SLA between the two types of surfaces is computed for each bin, then spatially averaged to obtain a monthly climatological value of the O/L bias. We interpolate it to a daily climatology, which is then removed from the lead dataset. Fig. S1 details this process, showing the time-averaged bias map and the climatological bias that is removed for each mission.

a supprimé: ¶

a supprimé: 3.3 Final editing of the data

a mis en forme : Normal

a supprimé: Once the SLA is estimated for every waveform, an extra editing is performed on the data similar to the one applied in

a supprimé: A

a supprimé: the

a supprimé: are used

a supprimé: still went through the

a supprimé: Then

a supprimé: iterative editing

a supprimé:

a supprimé: s

a supprimé: 3

a supprimé: For

a supprimé: leads(Armitage and Davidson, 2014; Quartly et al., 2019), a final treatment similar to

a supprimé: performed to prevent the “off-nadir hooking” phenomenon described by Quartly et al. (2019).

a supprimé: ¶

a mis en forme : Non Surlignage

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a supprimé: ¶

An example of an Envisat along-track is shown on Fig. 3a. The original leads and open ocean data are shown in gray. The leads data points kept after the editing steps are shown in red. The open ocean data points kept after the editing steps are shown in blue. Overall, fewer SLA measurements coming from leads are kept after the editing steps than in the open ocean domain. As in all filtering, a compromise has to be made between data quantity and data quality. Being too conservative may improve point wise precision, but would considerably reduce the number of leads kept and, thus, may provide insufficient coverage of sea level data. The remaining points are those in which we have the most confidence. The example along-track highlights the continuity between the open ocean domain and the ice-covered domain. A repetitive pattern on our along-tracks is that fewer data points, leads or ocean, pass through the editing in the transition zone between the open ocean and the sea ice. It corresponds to the region with a sea ice concentration ranging from 10 to 70%. This marginal ice band is where we have the least information and the highest uncertainties. Envisat is also equipped with the Medium Resolution Imaging Spectrometer (MERIS) ([19])

a supprimé: 4

a supprimé: s

a supprimé: s

The O/L bias correction is also applied to Envisat and SARAL data. Although the same retracker is used for open ocean and lead waveforms, we empirically observe a remaining residual bias. Applying this O/L bias correction also improves consistency across the missions. This O/L bias is identically applied to all lead data points and allows correcting for a homogeneous bias as depicted in Fig. S1. However, for Envisat we observe that a spatially heterogeneous O/L bias remains. This causes large negative anomalies to appear at the boundary between the open and ice-covered ocean. Empirically, we find a linear relationship between this spatially heterogeneous bias and the peakiness of the corresponding lead waveforms. Peakiness is defined as the ratio of the maximum waveform power to the mean power, and is a measure of the waveform specularity. Hence, we introduce a correction term amounting to the peakiness of the waveform times the slope regression between O/L bias and peakiness. This removes both the negative anomalies not observed with other satellites and reduces the heterogeneous bias to a homogeneous one (see Fig. S2).

3.3 Merging the missions

After correcting for the O/L bias of each individual mission, a cross-calibration is performed to account for inter-satellite offsets arising from differences in instruments, orbits and processing. CryoSat-2 is used as the reference mission as it overlaps with all three other missions. The strategy is analogous to the O/L bias correction. SLA estimates are aggregated in 75 km x 75 km x 10-days bins for each mission, and the mean offset relative to CryoSat-2 is computed over the overlapping time periods. These offsets are removed from each mission.

3.4 Optimal interpolation and gridding

After calibration, along-track data from all missions are combined into a daily gridded dataset using the DUACS mapping procedure (Taburet et al., 2019). It relies on an Optimal Interpolation algorithm (Bretherton et al., 1976; Ducet et al., 2000; Le Traon et al., 1998). SLA at each grid point is estimated as a weighted linear combination of nearby observations. The weights are determined by a space-time covariance function, which is controlled by three inputs: the spatial and temporal correlation scales, the expected signal variance, and the measurement noise. The correlation scales define the space-time radius within which the observations contribute to the grid point estimate. They thus control the smoothness of the output and the resolved scales. The signal variance sets the expected amplitude of SLA variability. The noise variance determines how much the optimal interpolation trusts individual observations. In the absence of any nearby observations, the interpolation returns zero SLA with an uncertainty equal to the full signal variance. SO_SLICE uses the inputs derived in Auger et al. (2022), which are presented in Fig. S4 and S5. The spatial correlation scales range from 100 km in the ACC to a 100 to 250 km in the subpolar Southern Ocean. The temporal correlation scales ranges from 30 days in the ACC to 10 days in the subpolar Southern Ocean. These parameters set a lower bound on the effective resolution of the gridded product. The interpolation cannot resolve features smaller than the prescribed correlation scales. The effective spatial and temporal resolution of SO_SLICE are assessed in Sect. 4.3.

- a supprimé: We choose to apply the
- a supprimé: same
- a supprimé: Saral
- a supprimé: Indeed, although
- a supprimé: in their case
- a supprimé: residual bias
- a supprimé: s
- a supprimé: F
- a supprimé: , an additional debiasing step is implemented depending on the peakiness of lead waveforms
- a supprimé: ¶
- a supprimé: ¶
- a supprimé: 5
- a supprimé: satellite
- a supprimé: n
- a supprimé: offset correction
- a supprimé: is applied across the different missions as they do not have the same baseline.
- a supprimé: s
- a supprimé: a
- a supprimé: and data from the other missions are adjusted towards its mean state value on their overlapping time range
- a supprimé: First,
- a supprimé: .
- a supprimé: Then, the mean inter-satellite offsets are computed (Table S2) and removed from the corresponding satellite. After, every satellite is centered on the Cryosat-2 mean value.
- a supprimé: 6
- a supprimé: Mapping the along-tracks
- a supprimé: After calibration, along-track data from all missions is combined into a daily gridded dataset using the DUACS-DT2022 mapping procedure (Taburet et al., 2019). It relies on an Optimal Interpolation (OI) algorithm (Bretherton et al., 1976; Ducet et al., 2000; Le Traon et al., 1998). It estimates the SLA at each grid point as a weighted linear combination of nearby observations. The weights are determined by a space-time covariance function, which is controlled by three inputs: the spatial and temporal correlation scales, the expected signal variance, and the measurement noise variance. The correlation scales define the space-time radius within which the observations contribute to the grid point estimate. They thus control the smoothness of the output and the resolved scales. The signal variance sets the expected amplitude of SLA variability. The (... [20]
- a mis en forme : Français
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)

Given the repeat cycles of the contributing missions (27 to 35 days) and the input temporal correlation scale, the daily fields are temporally interpolated rather than independent snapshots: the local effective temporal resolution is set by a combination of data density and the temporal correlation scale. Days with fewer nearby observations are associated with higher formal error. The final output is distributed on an EASE2 polar stereographic grid (Brodzik et al., 2012), with uniform 25 km x 25 km spatial grid cells, centered on the South Pole. Monthly and daily files are available.

3.5 Available variables

In addition to the gridded SLA fields, several variables are derived and distributed alongside the product, for user convenience. They are listed in Table 3. First the Absolute Dynamic Topography (ADT) is provided, computed as:

$$ADT = SLA + MDT$$

The MDT used is from Jousset (2023), derived from the CNES CLS22 MSS (Schaeffer et al., 2023) and the GOCO06S (Kvas et al., 2021). As the MSS used for the MDT computation differs from the Hybrid MSS (Laloue et al., 2024) used for the SLA computation, a static residual between the two MSS products may be present in the ADT field. Thus, users must use this field with caution. Future versions of the product will use a consistent MSS for both the SLA and MDT computations. Surface geostrophic velocities are computed from the ADT field using:

$$u_g = \frac{g}{f} (\hat{k} \times \nabla_H ADT)$$

where g is gravity, f is the Coriolis parameter, $\hat{k}$ is the unit vector normal to the geoid and ∇_H the horizontal gradient operator. These derived variables are provided for user convenience but are not analysed further in this study. The optimal interpolation also provides a formal mapping error at each grid point and time step, with the formula:

$$\sigma_{\text{map}}^2(x_0) = \sigma_s^2(x_0) - c^T A^{-1} c$$

Where c is the vector of signal covariances between the target grid point x_0 and the surrounding observations, $A = C_s + R$ is the sum of the inter-observation signal covariance and the observational noise covariance, and $\sigma^2(x_0)$ is the prior signal variance at x_0 . Hence, it can be defined as the prior signal variance minus the variance reduction achieved by the observations (Bretherton et al., 1976). It reflects the residual signal variance left unexplained by the observations after optimal interpolation and is driven by the density and distribution of nearby observations. It does not account for systematic error sources such as

a supprimé: 6

a supprimé: Mapping the along-tracks

a supprimé: In addition to the gridded SLA fields, several variables are derived and distributed alongside the product, for the user convenience. First the Absolute Dynamic Topography (ADT) is provided, computed as:

The MDT used is from Jousset (2023), derived from the CNES CLS22 MSS (Schaeffer et al., 2023) and the GOCO06S (Kvas et al., 2021). As the MSS used for the MDT computation differs from the Hybrid MSS (Laloue et al., 2024) used for the SLA computation, a static residual between the two MSS products may be present in the ADT field. Future versions of the product will use a consistent MSS for both the SLA and MDT computations. Surface geostrophic velocities are computed from the ADT field using:

a supprimé: the

a supprimé: .

a supprimé: ¶

a mis en forme : Police : Times New Roman, Non Italique

a mis en forme : Centré

a mis en forme : Police : Non Gras, Police de script complexe : Non Gras

residual biases or the omission of the SSB correction over leads. It should therefore be interpreted as a lower bound on the total uncertainty and is also distributed with the SLA field.

245 **Table 3.** Available datasets and associated fields

File	Field	Description
sla_25km_1M.nc	sla	Sea Level Anomaly (in m)
	MDT	Mean Dynamic Topography
	u_east	Zonal Geostrophic Current (in m s ⁻¹)
geostrophic_currents_25km_1M.nc	v_north	Meridional Geostrophic Current (in m s ⁻¹)
	formal_err	Formal Error (in m)
formal_error_25km_1M.nc	sla	Sea Level Anomaly (in m)
sla_25km_1d.nc	MDT	Mean Dynamic Topography

4 Results

4.1 Along track SLA data

Figure 4 shows an example of along-track SLA measurements (Fig.4a) used to produce SO_SLICE. They originate from an Envisat pass over the Weddell Sea (Fig.4b) on 11 October 2003. The thin grey line shows the SLA estimates before editing, from both open-ocean and lead waveforms. After the processing pipeline, the retained measurements are shown in blue (open ocean) and red (leads). 85% of the categorized open ocean waveforms are kept, and 10.5% of the lead waveforms. The resulting 178 lead measurements provide SLA information in the ice-covered domain. The along-track profile illustrates the continuity between the two surface types: SLA evolves smoothly from the open ocean into the sea-ice sector, with no jump at the ice edge.

255 To qualitatively illustrate that the retained lead points correspond to actual openings in the ice cover, we use the Medium Resolution Imaging Spectrometer (MERIS; Bézy et al., 2000), also mounted on Envisat. MERIS measures the radiance at the top of the atmosphere at several wavelengths. During clear-sky periods, it distinguishes highly reflective surfaces (sea ice) from poorly reflective ones (open water). A subset of along-track lead points (highlighted in orange) is colocized on the MERIS swath (Fig. 4c). A small number of points illustrate residual limitations of the classification. Observation 3 is classified as a lead but overlies a surface with a reflectance intermediate between neighbouring leads and ice floes. It may be a lead not

260

- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)
- a mis en forme : Anglais (E.U.)

265

visible due to MERIS resolution, but it could also corresponds to newly formed sea ice whose flat and snow-free surface produces a quasi-specular echo similar to a lead (Drinkwater, 1991). Observation 4 sits above sea ice, with a lead visible slightly off-nadir in the cross-track direction. These cross-track contaminations are not systematically detected by the hooking filter introduced in Sect. 3.1. Such misclassifications are discussed in detail by Poisson et al. (2018). However, the majority of retained lead points, as illustrated by observations 1 and 2, coincide with low-reflectance surfaces at 779 nm corresponding to open water exposed in leads. These cases represent the bulk of the retained observations in this example. The gridded product validation presented in Sect. 4.3 provides an integrated assessment of the impact of residual classification errors on the final fields.

270




Figure 4: (a) SLA from one Envisat pass, on 11-10-2003. Blue points are edited SLA from open-ocean waveform. Red points are edited SLA from leads waveforms. The grey line aggregates SLA estimates from open-ocean or lead waveforms, before the editing. (b) Trajectory of the along-track data shown in a. (c) Surface reflectance at 700 nm along the trajectory, measured by MERIS. SLA estimates shown in orange in panel a are colocalized on the MERIS swath. Points labelled 1 to 4 are discussed in the text.

a supprimé: (Poisson et al., 2018)

a mis en forme : Centré

a supprimé: 3

a supprimé: Example along-track

a supprimé: from the

a supprimé: mission

a supprimé: estimates

a supprimé: waveforms classified as open-ocean

a supprimé: estimates

a supprimé: waveforms

a supprimé: classified as leads

a supprimé: including the ones that were removed by the editing steps

a supprimé: A

a supprimé: Envisat along track

a supprimé: the

a supprimé: (Medium Resolution Imaging Spectrometer) instrument

a supprimé: MERIS captures spectral reflectance data, which can provide information on surface properties such as sea ice, water type, and biological activity.

a supprimé: A

a supprimé: collocated

a supprimé: ¶

4.2 Final dataset

The along-track measurements from the four satellite missions are combined using the DUACS procedure described in Sect. 3.4. The resulting gridded SLA dataset, SO_SLICE, has a grid resolution of 25 km and covers the period 2003–2021. It is publicly available (Mosneron Dupin, 2025).

Figure 5 presents monthly SLA fields for February and August 2008 and 2018, from C3S (Fig. 5a) and SO_SLICE (Fig. 5b). Two regions are considered: the ACC region and the subpolar Southern Ocean. The boundary between the two is defined as the -1.05 m MDT contour from MDT CNES/CLS22. In the ACC region, SO_SLICE displays strong agreement with C3S. Qualitatively, similar mesoscale patterns are visible in the monthly snapshots. Quantitatively, 95% of grid points located in the ACC display a correlation greater than 0.80 between SO_SLICE and C3S. The maps of their differences show residual discrepancies of approximately 1 cm. This may be due to C3S having an extra filtering on the open-ocean along-track data. In the subpolar region, SO_SLICE provides SLA estimates year-round and presents mesoscale features within ice-covered areas such as the Weddell Sea. By contrast, the C3S SLA field is heavily smoothed and contains missing data under sea ice. In seasonally or permanently ice-covered regions, SO_SLICE offers a substantial gain in spatial coverage and provides continuous SLA estimates across the entire subpolar Southern Ocean. The snapshots reveal spatial continuity between the open ocean and sea ice sectors: no sharp frontier exists at the transition, providing confidence in our processing and calibration. Where C3S does have data in the subpolar Southern Ocean, the two products have stronger differences than in the ACC. This arises from the computation of the climatological mean used to define the anomalies. For C3S, this mean is computed without wintertime observations in ice-covered regions, which biases the reference period and affects the anomaly values. SO_SLICE, by contrast, includes observations in all seasons yielding a different and more representative climatology.

Figure 6a shows the spatial power spectral density (PSD) of SO_SLICE and C3S. To compute the spectra, the spatial SLA field has been divided into 1600 km x 1600 km patches. The 2D Fast Fourier Transform isotropic PSD is computed in each patch and averaged over all directions. Spectra from patches in the ACC domain and in the subpolar Southern Ocean are separated and averaged together to get the characteristic PSD of both regions. In the ACC, both products are in close agreement at wavelengths longer than ~150 km. The spectral slope close to k^{-5} is consistent with interior quasi-geostrophic dynamics expected in an eddy-rich region. At shorter wavelengths, SO_SLICE retains slightly more energy. This reflects the additional along-track filtering performed in C3S, which suppresses energy and noise at small scales. In the subpolar SO, SO_SLICE contains substantially more energy at all wavelengths. The PSD ratio between the two products ranges from approximately 5 at ~1000 km to 17 at 50 km, illustrating the gain in resolved variability in ice-covered regions. This confirms that SO_SLICE and C3S resolve similar spatial scales in the ACC, and that SO_SLICE provides substantial added value in the subpolar Southern Ocean. Figure 6b shows the temporal PSD. In the ACC, both products are in close agreement at periods longer than 90 days, confirming the consistency of SO_SLICE with C3S for seasonal and interannual variability. Their associated spectra

a supprimé: 1

a supprimé: Following the procedures outlined in the Methods section,

a supprimé: a satellite altimetry dataset with a grid resolution of 25 km is obtained for the period 2003-2021

a supprimé: on ZENODO

a supprimé: The available variables are listed in Table 3. Both daily and monthly outputs are provided. Given the native temporal sampling of 10 to 30 days, the daily fields contain values temporally interpolated between observational acquisitions.

a supprimé: 4

a supprimé: an example of the

a supprimé: sea level anomaly (SLA)

a supprimé: 7

a supprimé: from our product

a supprimé: 4

a supprimé: -d

a supprimé: and from the CMEMS product (Fig. 4a-c).

a supprimé: open-ocean

a supprimé: s

a supprimé: our derived

a supprimé: SLA field shows

a supprimé: the product from Copernicus Marine Environment Monitoring Service (CMEMS

a supprimé:)

a supprimé: .

a supprimé: While the two products display similar mesoscale features in the ACC

a supprimé: .

a supprimé: some small-scale

a supprimé: can be observed

a supprimé: They

a supprimé: arise from

a supprimé: CMEMS incorporating additional satellite mission

a supprimé: s such as Jason-2, enabling improved resolution and spotting extra mesoscale features in dynamic open-ocean areas. CMEMS processing also incorporates a final smoothing step that is not included in the product presented in this paper. Quantitatively, when comparing time-series at each grid point between CMEMS and this study's product, we find t

a supprimé: hat 95% of grid points located in regions always ice-free exhibit a correlation greater than 0.80.¶

a mis en forme : Non Surlignage

are similar at time scales larger than 90 days. At shorter periods (90–20 days), SO_SLICE retains slightly more energy, which we impute again to differences in filtering. In the subpolar Southern Ocean, the SO_SLICE spectrum closely matches its ACC counterpart, indicating that the product captures comparable levels of temporal variability in both regions.




Figure 5: (a) Monthly SLA fields relative to the period 2003–2021 from C3S (b) Monthly SLA fields relative to the period 2003–2021 from SO_SLICE (c) Difference between C3S and SO_SLICE. Each row corresponds to a specific month, from top to bottom: February 2008, August 2008, February 2018 and August 2018.


a supprimé: overlap

a mis en forme : Non Surlignage

a supprimé: In both domains, energy decreases significantly at periods shorter than 30 days.

a supprimé: Table 3. Available datasets and associated fields

The August and February 2007 SLA maps (Fig. 4b-d) reveals spatial continuity between the open ocean and sea ice sectors: no sharp frontier exists at the transition, providing confidence in our processing. Mesoscale features are visible within ice-covered areas such as the Weddell Sea. It is not the case for the associated CMEMS maps (Fig. 4a-c). In seasonally or permanently ice-covered regions, the present product offers a substantial gain in spatial coverage. [21]



a supprimé:




Figure 6: (a) Spatial PSD of SO_SLICE and C3S SLA field in the ACC region and the subpolar Southern Ocean, as well as the ratio between SO_SLICE and C3S PSD. The linear fit between 250km and 70km is plotted, and the slope associated is given in the legend.

(b) Temporal PSD of SO_SLICE and C3S in the ACC and the subpolar Southern Ocean, with the PSD ratio for each region.

4.3 Validations

The validation of SO_SLICE proceeds in five steps, progressing from the raw input data to the final gridded product. First, we assess the consistency of the along-track observations between missions before any interpolation is applied (Sect. 4.3.1). Second, we evaluate the quality of the interpolated maps through leave-one-out comparisons with withheld along-track data (Sect. 4.3.2). Third, we examine the formal error output by the optimal interpolation as a measure of internal uncertainty (Sect. 4.3.3). Fourth, we compare the product with independent in situ observations from tide gauges and bottom pressure recorders (Sect. 4.3.4). Finally, we assess the consistency of SO_SLICE with existing altimetric SLA products in the ice-covered Southern Ocean (Sect. 4.3.5). Together, these diagnostics provide a comprehensive assessment of SO_SLICE accuracy, from the reliability of individual measurements through to the resolved scales of the final gridded fields.

a supprimé: Figure 4: (a-b) SLA field averaged over February 2007, from CMEMS (a) and from our product (b). (c-d) Same as before, for the month of August 2007. (e) Time series of the SLA averaged over five sectors of the subpolar Southern Ocean (south of 62°S; see inset). The daily data has been smoothed with a 6-month rolling average. [22]

a mis en forme [23]

a mis en forme [24]

a mis en forme [25]

a mis en forme [26]

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme [27]

a mis en forme [28]

a mis en forme [29]

a mis en forme : Normal

a mis en forme [30]

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme [31]

a mis en forme [32]

a mis en forme [33]

a mis en forme [34]

a mis en forme [35]

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme [36]

a mis en forme [37]

a mis en forme [38]

a mis en forme [39]

a supprimé: 2

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

4.3.1 Consistency between missions

As a first validation step, we compare the fully edited and calibrated outputs of each altimetry mission against one another. Each mission has different instrument characteristics, footprints and independent processing chains. A high level of agreement between them is therefore a strong indicator that the observed signal is geophysical rather than instrumental noise. For this comparison, SLA observations from each mission are averaged onto a 75 km EASE2 grid in 10-day bins, without optimal interpolation. Four sets of maps are obtained, one per mission. They are compared pairwise, using CryoSat-2 as the reference due to its temporal overlap with the three other satellites.

Figure 7a shows the spatial spectral coherence computed from these maps. As for the PSD analysis in Sect. 4.3, coherence is computed on 1600km x 1600km patches and averaged separately in the ACC and the subpolar Southern Ocean. The coherence $\gamma^2(k)$ measures the fraction of variance at each wavelength that is shared between two fields: $\gamma^2 = 1$ indicates perfect agreement, while $\gamma^2 < 0.5$ means that more than half the variance at that scale is independent between the two missions. For the two regions, all map pairs display coherence greater than 0.5 at wavelengths longer than 300 km, confirming that the satellites observe the same signal at these scales. This is corroborated by the time series of spatially averaged SLA in the subpolar Southern Ocean (Fig. 7b, c, d), which show strong mutual consistency at a 10-day time step. The Pearson correlation coefficient (r) and RMSD relative to CryoSat-2 are $r = 0.83$ and $\text{RMSD} = 1.18$ cm for Envisat, $r = 0.93$ and $\text{RMSD} = 0.75$ cm for SARAL, and $r = 0.87$ and $\text{RMSD} = 0.87$ cm for Sentinel-3A.

At shorter wavelengths, the scale at which coherence drops below 0.5 depends on both region and instrument type. In the ACC, all pairs cross the 0.5 threshold at approximately 180 km, indicating that the maps agree down to similar scales in the open ocean. In the subpolar Southern Ocean, the coherence between CryoSat-2 and Envisat (resp. SARAL) falls below 0.5 at 304 km (resp. 247 km), indicating weaker agreement than in the ACC. Several factors contribute to this degradation. LRM altimeters have higher measurement noise than SAR instruments (Boy et al., 2017), which reduces the signal-to-noise ratio at short wavelengths. In addition, the sparser sampling density of the LRM missions increases the likelihood that they observe a given grid cell at different times than CryoSat-2 within a 10-day window, introducing temporal representativeness errors that appear as decorrelation. By contrast, maps from CryoSat-2 and Sentinel-3A, the two SAR altimeters, maintain a similar coherence threshold in both regions. The higher sampling density of these missions produce more consistent measurements over leads, sustaining agreement to shorter wavelengths even under ice.


To complement the spectral analysis, the right-hand panels of Fig. 7 show grid-point RMSD and correlation distributions for each pair. These are computed on maps now coarsened to a 300 km grid, the scale above which all pairs show coherence greater than 0.5. For CryoSat-2 vs Envisat, the RMSD and correlation distributions are similar in the ACC and the subpolar Southern Ocean, with median RMSD above 3 cm and median correlation of ~ 0.64 in both regions. For the other two pairs

- a supprimé: 1
- a supprimé: 2
- a supprimé: confront
- a supprimé: s
- a supprimé: output with
- a mis en forme : Motif : Transparente
- a supprimé: utilizes
- a supprimé: s
- a supprimé: , retracking algorithms
- a supprimé: Hence, a
- a supprimé: missions
- a supprimé: in nature
- a supprimé: In Fig.
- a supprimé: 6
- a supprimé: Cryosat-2 is used as the reference due to its temporal overlap with the three other satellites.

- a mis en forme : Non Surlignage

- a supprimé: Time series of spatially averaged SLA south of 70°S derived from individual satellites show strong mutual consistency (Fig. 6a,d). The same monthly to seasonal variations are picked up. The Pearson correlation coefficient (R) and root-mean-square difference (RMSD) relative to Cryosat-2 are: R = 0.91 and RMSD = 1.20 cm for Envisat, R = 0.87 and RMSD = 1.18 cm for Saral, R = 0.83 and RMSD = 1.11 cm for Sentinel-3A
- a mis en forme : Non Surlignage
- a mis en forme : Non Surlignage
- a mis en forme : Non Surlignage
- a mis en forme : Non Surlignage


(CryoSat-2 vs SARAL, CryoSat-2 vs Sentinel-3A), median RMSD values are lower (~1.8–1.9 cm) and median correlations are higher (~0.8 in the subpolar SO, ~0.9 in the ACC), consistent with the lower range noise of SAR (Boy et al., 2017) and Ka-band (Verron et al., 2015) altimeters. Together, these diagnostics confirm robust inter-mission agreement at scales larger than 300 km across the entire domain. At smaller scales, the disagreement is driven by instrumental noise and spatially heterogeneous sampling. These limitations reflect the properties of the raw binned maps (75 km grid, 10-day time step) before any interpolation is applied. Merging multiple missions through the optimal interpolation is expected to mitigate them by increasing the effective observation density with correlation length scales larger than 75 km.



a mis en forme : Non Surlignage

a supprimé: . Additionally, monthly snapshots of SLA fields reveal good agreement in large-scale spatial patterns across missions (Fig. 6b,c and e,g). While discrepancies are evident at smaller spatial scales, they are largely attributable to differences in satellite orbital characteristics. Indeed, each satellite samples the sea surface at different times and locations within a given month, introducing variability across mission-specific SLA fields. This highlights the benefit of merging the missions to gain precision.

a mis en forme : Anglais (G.B.)



a supprimé:

interpolation blends both observations. The temporally mismatched observation introduces a representativeness error that in that instance would not be present in a single-mission map. The net effect is a more compact RMSE distribution: the extreme values are reduced, while the median remains similar. Similar RMSE values are obtained for the other mission combinations, with typical median RMSE values around 4 cm, and a slight tendency toward lower RMSE in the subpolar Southern Ocean (Fig. 8b). These leave-one-out RMSE values represent upper bounds on the errors of the final product, since SO SLICE integrates all four missions simultaneously rather than withholding one. Overall, the comparable RMSE values in the ACC and the subpolar Southern Ocean indicate that the processing and mapping of lead returns in the ice-covered domain achieves a quality comparable to the standard open-ocean processing.




Figure 8: (a) Winter (June-July-August) RMSE map of SARAL along-track values compared to interpolated data from a CryoSat-2 + Sentinel-3A map, during the period 2016-2021. The white dotted line is the ACC/subpolar Southern Ocean MDT contour. The black dashed line is the 3% contour of the climatological minimum of sea ice concentration for 2016-2021. (b) Distribution of grid-cell RMSE for the 5 sets of along-track data/map comparison. ACC grid cells are on blue and subpolar Southern Ocean cells in orange. The median of the distribution is indicated by the thick full lines, with their associated values. The thin dotted lines are the 95% quantile.

4.3.3 Formal error

The formal error associated with optimal interpolation represents the expected error variance of the interpolated sea level estimate at each grid point. It quantifies the confidence in the gridded field based on both the spatial and temporal distributions of observations. In practice, it provides a lower bound on the total uncertainty under the assumptions of the interpolation scheme, i.e. unbiased errors and known covariances. Here, we use the formal error as a proxy for the local uncertainty in the reconstructed SLA field.

a supprimé: Outside of these regions, the RMSE remains broadly consistent across both open-ocean and ice-covered domains, with median values of 4.0 cm and 3.7 cm, respectively.


a supprimé: results

a supprimé: under sea ice

a supprimé: 7b

a supprimé: These results indicate that our processing of lead returns from the sea-ice domain

a supprimé: is of a comparable quality to the more standard open ocean processing. As the latter performed well against DUACS, it provides confidence that our merged (open ocean and sea-ice) dataset behaves well. At face value, the computed RMSEs should ter ... [55]



a supprimé:

a supprimé: 7

a supprimé: Sentinel-3A

a supprimé:

a supprimé: Altika/

a supprimé: Cryosat

a mis en forme

a supprimé: ith

a supprimé: 5% contour of sea-ice cover for the climate ... [57]

a supprimé:

a supprimé: b

a supprimé: Open-ocean

a supprimé: i

a supprimé: (seasonally) ice-covered

a supprimé: They are categorized according to the whi ... [58]

a supprimé: 2

a supprimé: (OI)

a supprimé: distribution and density

a supprimé: Sea Level Anomaly (SLA)

a supprimé:)

Figure 9 presents daily maps of this formal error under different satellite observation scenarios. Figure 9a corresponds to a day in August 2008, during which only Envisat data are used. The average formal error at that time is 1.7 cm, with higher uncertainty concentrated within highly energetic regions of the ACC, the marginal ice zone, and certain coastal areas. Banded patterns stem from the uneven distribution of satellite tracks. Overall, uncertainty hotspots reflect the combined effects of lower observation density, stronger mesoscale activity, and signal aliasing due to the limitations of a single-mission dataset. Panel b of Fig. 9 illustrates the formal mapping error for a day in 2012 when only CryoSat-2 data are available. Despite the improved precision of its SAR mode over leads compared to what can be achieved by Envisat (Sect. 2.1.), having only CryoSat-2 SAR data has two intrinsic limitations for lead SLA mapping. Its SAR mode acquisition does not extend close to the coast in many regions, and its orbit results in daily maps with spatial gaps. These limitations manifest as increased domain averaged (i.e. south of 50°S) formal errors, with pronounced striping artifacts and degraded confidence near the coast, visible both in the map (Fig. 9b) and the time series (Fig. 9c). Panel c of Fig. 9 illustrates the formal error on a representative day in August 2018 when the multi-mission dataset includes CryoSat-2, SARAL, and Sentinel-3A. The incorporation of multiple satellite tracks improves spatial sampling and cross-track resolution, leading to a reduction in the average formal error to 1.1 cm. Although the same dynamical regions (ACC, marginal ice zone, and coastal zones) remain areas of enhanced uncertainty, the magnitude of these errors is significantly lower. This highlights the main benefit of combining several missions. It reduces the uncertainty of SLA estimates, especially in higher uncertainty regions.

Overall, the formal error analysis gives a typical uncertainty of approximately 1.5 cm for SO SLICE, in line with previous estimates by Armitage et al. (2016) or Giles et al. (2012). The subpolar Southern Ocean has an average uncertainty of 1.3 cm, with larger uncertainties on the continental shelf (average: 1.9 cm) than off the shelf (average: 1.2 cm). These are average numbers: the local uncertainty can be quite different as it is strongly affected by the density of observations, i.e. the number of satellites available and their orbit (Fig. 9e). The formal error time series also highlight the benefits of multi-mission merging. During the Envisat-only period (2003–2010), the subpolar formal error displays a seasonal cycle, peaking at 1.9 cm in winter when ice coverage is at its maximum and sampling density is at its lowest. From 2013 onward, the addition of SARAL and Sentinel-3A largely suppresses this seasonality and reduces the peak formal error to 1.1 cm in the subpolar Southern Ocean. We note that the formal error does not capture systematic errors, such as satellite-specific biases, retracking uncertainties, or residual geophysical corrections. As such, while the formal error maps are a valuable diagnostic, they must be interpreted as only one component of the total uncertainty.

a supprimé: ...98...presents daily maps of this formal error under different satellite observation scenarios. Panel ...figure 9a corresponds to a day in August 2008... during which only Envisat data is ...re used. The average formal error at that time is 1.7 cm, with higher uncertainty concentrated within highly energetic regions of the ACC, (i.e., oceanic storm tracks in the ACC), ...he marginal ice zone, and certain coastal areas. Banded patterns stem from the uneven distribution of satellite tracks. Overall, uncertainty hotspots reflect the combined effects of lower observation density, stronger mesoscale activity, and si...ignal aliasing due to the limitations of a single-mission dataset. Panel b of Fig. 8 ... illustrates the formal mapping error for a day in 2012 when only CryoSat-2 data is ...re available. Despite the improved precision of its Synthetic Aperture Radar...AR mode over leads compared to what can be achieved by e.g....Envisat (Sect. 2.1.) (REF)... having only CryoSat...ryoSat-2 SAR data has...two intrinsic limitations for lead SLA mapping:g: li...s SAR modeM...acquisition does not extend close to the coast in many regions, and its orbit results in non-uniform sampling...aily maps with temporal ...patial gaps. These limitations manifest as increased globally ...omain averaged (i.e. south of 53...°S) formal errors, with pronounced striping artifacts and degraded confidence near the coast, clearly ...isible both in the map (Fig. ...8...) and the time series (Fig. 8c...e). Panel c of Fig. 98...illustrates the formal error on a representative day in August 2018 9, ...hen the multi-mission dataset includes CryoSat-2, Saral...ARAL, and Sentinel-3A in addition to Envisat... The incorporation of multiple satellite tracks improves spatial sampling and cross-track resolution, leading to a reduction in the average formal error to 1.1 cm. Although the same dynamical regions (ACC, MIZ...arginal ice zone, and coastal zones) remain areas of enhanced uncertainty, the magnitude of these errors is significantly lower. This highlights the main benefit , confirming the benefit ...f combining multi-mission products (... [59])

a supprimé: about ...pproximately 1.5 cm for our product...O SLICE, in line with with ...revious estimates by Armitage et al. (2016) and ...r Giles et al. (2012) estimations... The subpolar Southern Oo...ean has an average uncertainty of 1.3 cm, with larger uncertainties on the continental shelf (average: 1.9 cm) than north of it...ff the shelf (average: 1.2 cm). These are average numbers: the local uncertainty can be quite different as it is strongly affected by the density of observations, i.e. the number of satellites available and their orbit (Fig. ...8...). The formal error time series also highlight the benefits of multi-mission merging During the Envisat-only period (2003–2010), the subpolar formal error displays a seasonal cycle, peaking at 1.9 cm in winter when ice coverage is at its maximum and sampling density is at its lowest. From 2013 onward, the addition of SARAL and Sentinel-3A largely suppresses this seasonality and reduces the peak formal error to 1.1 cm in the subpolar Southern Ocean. We note that the formal error does not capture systematic errors, such as satellite-specific biases, retracking uncertainties, or residual geophysical corrections. As such, while the formal error maps are a valuable diagnostic tools... they must be interpreted as only one component of the total uncertainty, not a complete representation of it (... [60])





Figure 9: Optimal interpolation formal error maps in August for three different years. (a) is in 2008, with only Envisat data. (b) is in 2012, with only CryoSat-2. (c) is in 2018, with CryoSat-2, SARAL and Sentinel-3A data. (d) Time average daily formal error map over the full period. (e) Spatially averaged formal error time series, for several subdomains: south of 50°S in blue, subpolar domain in orange, off-shelf subpolar domain in green and on-shelf subpolar domain in red. The subpolar Southern Ocean is south of the white dashed contour in the maps; the off-shelf/on-shelf separation is defined by the 1000 m bathymetric contour shown as a white full contour.

4.3.4 Comparisons with in-situ observations

An alternative method to validate SO SLICE is to compare it with observations from in-situ instruments. We use six tide gauges and five Bottom Pressure Recorders (BPRs) for this validation step (Fig. 10a). The majority of BPRs are located in the Atlantic sector of the subpolar Southern Ocean, while all tide gauges are deployed on the Antarctic coast, either in the Indian or Atlantic sector. There are several caveats to this comparison. First, the number of available instruments remains limited, reflecting the scarcity of sustained observational platforms in the Southern Ocean. Nevertheless, this study compiles the largest set of in situ records used for SLA validation in this region to date. Second, the spatial coverage is restricted to the Atlantic



a supprimé:

a supprimé: 8... (Top) ...ptimal interpolation formal error maps in August for three different days...ears. (a) is in 06/08/2004 ... [61]

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a supprimé: 04/08/2012

a supprimé: Cryosat...2. (c) is in 08/08/2019 ... [62]

a supprimé: Cryosat...2, Saral ...ARAL and Sentinel-3A data. (Bottom) ... [64]

a mis en forme : Police :9 pt, Gras, Police de script complexe :9 pt, Gras

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a supprimé: defined

a supprimé: by a -180cm MDT...contour shown as ...he continuous ...hite black ...ashed contour in the mapsd... the off-shelf/on-shelf separation is defined by the 1000 m bathymetric contour shown as a dashed ...hite black ...ull contour. in ... [65]

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Gras, Police de script complexe :9 pt, Gras

a supprimé: 2...4 In-situ comparisons ... [66]

a supprimé: validation ...ethod to validate that assesses the net error associated with the satellite-derived data...O_SLICE set ...s to compare it with observations from in-situ instruments. We use six tide gauges and five Bottom Pressure Recorders (BPRs) for t ... [67]

1990 and Indian Ocean sectors, with no data available from the Pacific sector. Finally, the operational periods of the instruments primarily overlap with the first half of the altimetry record. It is the period with only Envisat, hence the one with the highest uncertainties (Fig. 9e).


Despite these limitations, the comparison provides valuable insight into SO SLICE performance. The observations are filtered to make a reliable comparison. For tide gauges, both the product and in situ time series are smoothed using a 15-day moving average. For BPRs, the 60-day rolling average is removed from the 15-day rolling average to isolate variability within the 15–60 days frequency band. At frequencies lower than 60 days the BPR signals are more likely to not be associated with SLA signals (Vinogradova et al., 2007). In addition, SLA time series from the satellite-based dataset are spatially averaged within a 300 km radius of each in situ instrument. Correlations between the satellite-derived SLA and tide-gauge observations are all statistically significant ($p \ll 0.05$) with correlation coefficients ranging from 0.51 to 0.77, depending on the tide gauge (Table 4, Fig. S8–13). Two examples of time series are shown in Fig. 10(b,d) to illustrate the good agreement including under sea-ice cover (gray shading on Fig. 10). We find correlation coefficients of 0.77 and 0.6 and RMSE values of 4.2 and 6.4 cm for the time series shown in panels b and d, respectively. Seasonal cycles are well captured, although some discrepancies in amplitude are noted. Shorter-timescale variability (~15 days) is also partially resolved by the product. A regression analysis aggregating all tide gauge data points during sea ice conditions ($> 10\%$ ice cover) yields a correlation of 0.58 and an RMSE of 4.8 cm (Fig. 10c). The regression deviates from the 1:1 line, with extreme SLA values being biased low in the satellite-derived product. The slope of the linear fit between the aggregated tide gauges and our altimeter product is at 0.47. This may be attributed to several factors: the coastal proximity of tide gauges, where satellite altimetry performance is known to degrade; the temporal mismatch between satellite overpasses and short-lived coastal features; the smoothing and averaging applied to the product; and the uncertainty introduced by tidal aliasing and corrections in polar coastal regions. Similar to the comparison with the tide gauges, all correlations between satellite-based SLA and BPRs are statistically significant, with correlation coefficients ranging from 0.45 to 0.83 depending on the BPR (Table 5; Fig. S14–18). One example of time series shown in Fig. 10f demonstrates that SLA variability in the 15–60-day range is well captured, with a correlation of 0.52. A regression analysis combining all BPR data points yields a correlation of 0.66 and an RMSE of 1.1 cm for ice-covered periods (Fig. 10e). Similar to the tide gauge comparison, the regression slope deviates from the 1:1 line, confirming that the product underestimates the largest local SLA values.

Overall, these results confronting our satellite-based SLA estimate to in situ observations confirm that the product reliably captures sub-seasonal SLA variability even in ice-affected regions, with an error of 1–5 cm and a tendency to underestimate SLA extremes. Largest error estimates (4–5 cm) are associated with comparison with tide gauges that are all coastal. Further away from the coast, error estimates computed from comparisons with BPRs drop to 1–2 cm.

a supprimé: not the most accurate ...he one with the highest uncertainties (Fig. 9e8) ... [68]

a supprimé: the product ...erformance. Because in situ observations and the satellite product are associated with distinct temporal and spatial scales, we filter the observations ...he observations are filtered to make a reliable comparison. For tide gauges, both the product and in situ time series are smoothed using a 15-day moving average. For BPRs, the 60-day rolling average is removed from the 15-day rolling average to isolate variability within the 15–60 days frequency band. A; a... frequencies lower than 60 days the BPR signals are more likely to not be associated with SLA signals (Vinogradova et al., 2007). In addition, SLA time series from the satellite-based dataset are spatially averaged within a 150 ...00 km radius of each in situ instrument after applying a 150 km Gaussian smoothing filter... Correlations between the satellite-SLA and tide-gauge observations are all statistically significant ($p \dots 0.05$) with correlation coefficients ranging from 0.5144...to ... [69]

a supprimé: mesoscale and ...ub-seasonal SLA variability even in ice-affected regions, with an error of 1–5 cm and a tendency to underestimate the largest ... [70]



a supprimé:




Figure 10: (a) Locations of in-situ instruments (stars: tide gauges; circles: BPRs). (b) SLA time series from the Davis tide gauge (blue) and co-located SO_SLICE grid-point average within 150 km (orange), both smoothed with a 15-day rolling average. Grey shading indicates sea-ice concentration >10% from Ocean and Sea Ice Satellite Application Facility (EUMETSAT OSI SAF, 2017). (c) 2D histogram of product SLA vs. tide gauge SLA for all data points coinciding with sea-ice concentration >10%, with regression line (red) and 1:1 line (dashed black). (d) Same as (b) for Mawson. (e) Same as (c) for BPR data. (f) Same as (b) for the Myrtle C BPR, with an additional 60-day rolling average subtracted to isolate the 15–60 day variability band.

a mis en forme : Centré

a mis en forme : Anglais (E.U.)

a mis en forme : Justifié

a mis en forme : Anglais (E.U.)

a supprimé: Figure 9: (a) In-situ instrument locations. Stars are tide gauges, circles are BPRs. (b) Time series of in-situ SLA measured by the Davis tide gauge (blue) and the associated altimetric SLA (orange). The latter is obtained by averaging all grid points within a 150 km radius of the tide gauge. Both time series are smoothed with a 15-day rolling average. The grey shading corresponds to periods with a sea-ice concentration higher than 10% (obtained from OSISAF(EUMETSAT OSI SAF, 2017)(OSI SAF and EUMETSAT SAF on Ocean and Sea Ice, 2017)). (d) Same as (b) for the Mawson tide gauge. (f) Same as (b) for the Myrtle C Bottom Pressure Recorder. In this instance, both time series are smoothed with a 15-day rolling average. The 60-day rolling average is then subtracted to keep the variability in the 15–60 days band. (c) Product SLA vs Tide gauge SLA in a two-dimensional histogram obtained by aggregating all data points from the tide gauge comparisons that coincide with a period of sea-ice concentration higher than 10%. The associated regression is plotted in red, and the 1:1 line is shown in dashed black. (e) Same as (c) but for BPR data vs product SLA.

Table 4: Comparison metrics with tide gauges, computed with data points when sea-ice concentration is greater than 10%. Effective temporal resolution as defined by Ballarotta et al. (2019)

Tide gauges	Casey	Davis	Dumont d'Urville	Faraday	Mawson	Syowa	Total
Pearson R	0.56	0.61	0.57	0.51	0.77	0.6	0.58
RMSE (in cm)	4.2	4.3	4.8	5.0	4.2	6.4	4.8
Effective resolution (days)	35	35	23	33	45	48	36.5

Table 5: Comparison metrics with the full Bottom Pressure Recorders time series, computed with data points when sea-ice concentration is greater than 10%. Effective temporal resolution as defined by Ballarotta et al. (2019)

Bottom Pressure Recorders	Myrtle B	Myrtle C	DPS	DPS Deep	ANT17	JARE	Total
Pearson R	0.72	0.85	0.77	0.74	0.57	0.59	0.66
RMSE (in cm)	1.1	1.01	1.3	1.3	1.8	2.1	1.1
Effective resolution (days)	44	33	36	38			37.75

4.3.5 Comparison with recent altimetry products

Figure 11a shows the spatial PSD of SO_SLICE compared to Dragomir (2024) and Auger et al. (2022a) over the ACC and subpolar SO. Compared to Dragomir (2024), SO_SLICE contains more energy at all wavelengths except the very largest scales (>1000 km), in both regions. This is consistent with the fact that the Dragomir product relies on a single mission for most of its record, which limits the observation density and thus the resolved variability. Its mapping procedure is a 1°x1° spatial binning which acts as a low-pass filter and suppresses variability at a cutoff wavelength. SO_SLICE PSD matches well with Auger et al. (2022a). In the ACC, Auger et al. displays slightly more energy at short wavelengths, possibly reflecting differences in along-track editing or noise levels. In the subpolar SO, the two spectra are in close agreement across most scales.

Gridded maps of pointwise RMSD and correlation between SO_SLICE and the two products are shown in Fig. 11b and 11c. The associated boxplots can be found in Fig. S7. Once again, the region where the differences (weak correlation, large RMSD) are the most striking is the ACC for Dragomir. In the subpolar SO, the median RMSD is at ~2 cm in both cases. The temporal correlation is 0.8 with Auger et al. (2022a) and 0.7 with Dragomir (2024). The higher correlation with Auger et al. is expected, as the two products share a similar processing methodology and common satellite missions over the 2013–2019 period. The

a mis en forme... [71]

a mis en forme... [72]

a mis en forme... [73]

a supprimé: 1

a mis en forme... [74]

a mis en forme... [75]

a supprimé: 65

a supprimé: 59

a supprimé: 44

a supprimé: 1

a supprimé: 3

a supprimé: 57

a mis en forme... [76]

a supprimé: 3.5

a supprimé: 3.8

a supprimé: 5.0

a supprimé: 4.6

a supprimé: 3

a supprimé: 5.8

a supprimé: Table 4: Comparison metrics with tide gauges.

a supprimé: . In brackets, the metrics associated ...ith onl... [80]

a mis en forme... [77]

a mis en forme... [79]

a mis en forme... [78]

a mis en forme... [81]

a mis en forme le tableau... [82]

a mis en forme... [83]

a supprimé: 0.45 (...7245) ... [84]

a supprimé: 0.78 (...853) ... [85]

a supprimé: 0.55 (...7768) ... [86]

a supprimé: 0.65 (...76...) ... [87]

a supprimé: 0.52 (...56...7) ... [88]

a supprimé: 0.68 (...59) ... [89]

a supprimé: 0.59 (...66) ... [90]

a mis en forme... [91]

a supprimé: 1.1 (...1) ... [92]

a supprimé: 0.9 (...012) ... [93]

a supprimé: 1.1 (...3) ... [94]

a supprimé: 1.0 (...31) ... [95]

a supprimé: 0.9 (...80.9) ... [96]

a supprimé: 2.1 (...1) ... [97]

a supprimé: 1.0 (...1) ... [98]

a supprimé: Table 5: Comparison metrics with the full... [99]

a mis en forme... [100]

a supprimé: 2...5 Comparison with a... [101]

a supprimé: The SSH product presented here is compare... [102]

a mis en forme... [103]

a mis en forme... [104]

main differences with Auger et al. (2022a) are concentrated on the continental shelf, particularly in the Weddell Sea. These are attributed to a change in MSS between the two products. The main differences with Dragomir are found near the coast and in the most northern part of the subpolar domain. The coastal discrepancies likely reflect the inclusion of CryoSat-2 SARIn data for Dragomir, while the northern boundary differences may arise from the different spatial extent and mission coverage of the two products. Overall, these comparisons confirm that SO SLICE is consistent with existing independent products where they overlap, while offering extended temporal and spatial coverage.




Figure 11: (a) Spatial PSD of the SO_SLICE SLA field compared to the Dragomir (2024) and Auger et al. (2022a) datasets, computed separately in the ACC (left) and the subpolar Southern Ocean (centre). The right panel shows the ratio of the SO_SLICE PSD to each comparison product, for each region. (b) Grid-point correlation (left) and RMSD (right) maps between SO_SLICE and Dragomir et al. (2024). (c) Same as (b) but between SO_SLICE and Auger et al. (2022a).

4.4 Effective resolution of SO_SLICE

After validating SO_SLICE, its effective spatial and temporal resolution is computed. The effective spatial resolution of a gridded altimetric product is the shortest wavelength at which the mapped fields faithfully represent the ocean signal. To estimate it, we follow the procedure from Ballarotta et al. (2019). Independent along-track observations, withheld from the interpolation, serve as ground truth. The mapping error is defined as the difference between these withheld along-track measurements and the colocated gridded SO_SLICE values. The effective resolution is the wavelength at which the error spectrum reaches half the signal spectrum. It is the wavelength where the noise-to-signal ratio (NSR), defined as the PSD of the error divided by the PSD of the ground truth, crosses 0.5 when moving from large to small scales.

The NSR is computed on 500 km segments extracted from one year of CryoSat-2 along-track data withheld from the mapping. Figure 12a shows an example along-track pass with two such segments. The orange segment lies mainly in the ACC, yielding an estimated effective resolution of 105 km. The blue segment is located in the subpolar Southern Ocean and consists of lead measurements. It yields an effective resolution of 184 km. As illustrated by this example, individual spectra are noisy. Obtaining robust estimates requires averaging over many segments to obtain stable NSR spectra. To do so and regionalize our metrics, each segment is referenced by its median latitude and spectra are averaged within latitude bands. Figure 12c shows the resulting NSR curves for several bands. From 50°S to 65°S, the effective resolution is comprised between 124 and 133 km, consistent with both the input correlation scales prescribed in the interpolation (Sect. 3.4) and the values reported by Ballarotta et al. (2019) for the C3S product at comparable latitudes. South of 65°S, the effective resolution degrades to approximately 200 km, as observed for example in the Weddell Sea. This is coherent with the longer correlation scales prescribed in this region, and the sparser observation density that limits the amount of information the optimal interpolation can extract. Segments are also referenced by their longitude and binned in 10°x10° boxes to produce the effective resolution map of Fig. 12d. Overall, the spatial pattern of effective resolution closely follows the input correlation length scales, with lower resolution found in the Weddell Sea and in the north of the Ross Sea, where the correlation scales are the highest. This points toward these parameters being the primary control on the resolved scales of the gridded product.

The effective temporal resolution is estimated using the same NSR framework. The in-situ observations described in Sect. 4.3.4 serve as ground truth. The NSR is computed on 180-day segments between colocated SO_SLICE values and in situ records. Table 4 and 5 summarize the temporal effective resolution estimated at each station, and the corresponding NSR spectra are provided in the supplementary material. The estimated temporal resolution ranges from 23 to 48 days across stations, with a mean of 37 days. This is consistent with the temporal correlation scales prescribed in the OI, which range from 20 to 30 days in the coastal subpolar Southern Ocean and the Drake Passage. It is also comparable to the average temporal resolution of 34 days estimated by Ballarotta et al. (2019) for the C3S product. In both studies, the temporal resolution estimate relies on comparison with in-situ observations that are predominantly coastal. Altimetric performance is known to degrade

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (G.B.)

a mis en forme : Anglais (E.U.)

a mis en forme : Justifié

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

near the coast due to land contamination of the radar footprint and less accurate geophysical corrections. The values reported here may therefore represent a conservative estimate of the temporal resolution in the offshore ocean, where these coastal effects are absent.




Figure 12: (a) Example CrvoSat-2 along-track pass withheld from the OI: along-track SLA measurements (black) and collocated gridded SO SLICE values (red). The orange and blue shading indicate two 500 km segments used for spectral analysis. (b) Noise-to-signal ratio (NSR) computed on the orange and blue segments individually and averaged over all segments from this pass (black). The dashed line indicates the NSR = 0.5 threshold. The effective resolution estimated on these segments is indicated with the figures. (c) NSR spectra averaged over all segments split into five latitude bands, and the associated effective resolutions. (d) Map of effective spatial resolution, estimated from spectra obtained by averaging every spectrum segment within 10°x10° boxes.

5 Discussion

In this study, we presented SO SLICE, a 19-year SLA dataset covering the open-ocean and ice-covered Southern Ocean over the period 2003–2021. It is the longest altimetric record currently available for the entire Southern Ocean. The approach relies on detecting radar echoes reflected from leads and extracting SLA measurements from them. Certain assumptions are required to enable this retrieval, most notably the omission of the SSB correction within sea-ice. They represent known sources of residual bias. After data editing, the density of valid observations in ice-covered regions is equal or greater than in the open ocean. Along-track data from four satellite altimetry missions are combined through optimal interpolation to produce gridded SLA fields on a 25 km EASE2 grid. Monthly snapshots of SO SLICE show it provides spatially continuous coverage across both the open-ocean and ice-covered domains, with smooth transitions at the ice edge. The spectral content of the product is

a mis en forme : Anglais (E.U.)

a supprimé: ¶

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a mis en forme : Police :9 pt, Police de script complexe :9 pt

a supprimé: ¶

a supprimé: In this study, we presented the methodology behind the retrieval of Sea Level Anomalies (SLA) in ice-covered regions over the period 2003-2021. The approach is based on detecting and reprocessing previously acquired radar echoes that are coming from leads in the sea-ice cover. SLA measurements are extracted from them. While certain assumptions—such as neglecting sea state bias (SSB) in leads—are not entirely satisfactory, they are necessary compromises to enable this SLA estimation. After data editing, the density of valid observations in ice-covered regions is lower than in the open ocean. However, these retained points are of high quality, and the along-track profiles demonstrate good continuity across the sea-ice edge. Along-track data from four satellite altimetry missions are combined together to construct an SLA product. This dataset, presented here, is the longest time series currently available for the entire Southern Ocean, spanning 19 years. It provides consistent SLA fields across both open-ocean and ice-covered regions, with smooth transitions at the ice edge. It captures large-scale and some mesoscale variability, indicating the physical coherence of the signal throughout the domain.¶

a supprimé: , which could be an issue but is discussed later on

consistent with the expected dynamical regimes in the ACC and reveals a substantial gain in resolved variability in the subpolar Southern Ocean compared to the C3S reference product.

Multiple validation procedures are carried out, progressing from the raw input data to the final gridded product. Along-track observations from each satellite mission are first binned separately onto a 75 km grid in 10-day windows, without interpolation, and the resulting maps are cross-compared pairwise, using CryoSat-2 as the reference due to its temporal overlap with the three other missions. Despite differences in instrumentation and orbits, all mission pairs display spectral coherence greater than 0.5 at wavelengths longer than 300 km, confirming that the satellites retrieve the same large-scale physical signal from lead observations. At shorter scales, maps from LRM altimeters (Envisat, SARAL) show lower coherence than maps from SAR altimeters (Sentinel-3A) in the subpolar Southern Ocean. This is attributed to the higher measurement noise and sparser sampling density of LRM missions, which may introduce temporal representativeness errors within the 10-day binning window. To assess whether these limitations propagate through the interpolation into the final product, we perform leave-one-out comparisons between withheld along-track observations and gridded maps constructed from the remaining missions. The resulting RMSE values are comparable in the open-ocean and ice-covered domains (~4 cm), even for the Envisat-only period when sampling density is at its lowest. This demonstrates that the lead-based SLA retrievals carry sufficient geophysical information to constrain the interpolated fields at a level of accuracy comparable to the open ocean. The main benefit of adding missions is to reduce the uncertainty of the final dataset, as confirmed by the formal error analysis, which shows a reduction from 1.9 cm (single mission) to 1.1 cm (multi-mission) in the subpolar Southern Ocean.

SO SLICE is also compared with independent in situ observations from tide gauges and bottom pressure recorders. While these observations are scarce and mostly available for the first half of the time period (2003–2013), they provide a fully independent benchmark. Overall, the correlations are statistically significant ($p \ll 0.005$). Specifically, we find a mean correlation coefficient of 0.58 for tide gauges and 0.66 for BPRs. These results indicate that the product captures physical variability during the earlier period of the dataset. Nevertheless, some discrepancies remain. In particular, the greatest positive and negative SLA values observed in tide gauge records are not fully reproduced by the altimetry product. This is reflected in the slope of the linear fit between altimetry and tide gauge SLA, which is 0.47, deviating from the 1:1 relationship. These discrepancies likely stem from processes unresolved by satellite altimetry, which smooths out localized coastal variability, and to inaccuracies in tidal correction models in the complex coastal environments where the tide gauges are located. Despite these limitations, the in-situ comparisons support the reliability of our dataset and confirm its ability to capture meaningful physical variability in ice-covered regions of the Southern Ocean.

The effective resolution of SO SLICE was assessed in both the spatial and temporal domains. Using the noise-to-signal ratio (NSR) approach of Ballarotta et al. (2019), the effective spatial resolution is estimated at approximately 120 km between 50°S and 65°S, degrading to 200 km south of 65°S. This estimate is based on the year 2018, when SO SLICE incorporates three

a mis en forme : Non Surlignage

a supprimé: were

a supprimé: First, computed SLA in open-ocean regions is compared with the DUACS reference product. Both datasets show good agreement with high correlation for most grid points. Then, input satellite missions were separately grouped, binned and then cross-compared. This exercise showed that their outputs are consistent despite independent processing pipelines and differences in instrumentation, measurement modes, frequency bands. It shows that no mission-specific biases were introduced by mission-specific choices. This consistency supports the robustness of the global processing strategy. Furthermore, the difference between along-tracks and independent maps in both open-ocean and ice-covered domains was computed. The results suggest that our overall process extracts comparable levels of signal and variability from leads and open-water along-track data. Given the high confidence in the open-ocean component, this result indicates that the dataset provides a reliable estimate of SLA in the sea-ice-covered Southern Ocean.

a supprimé: Finally, we conducted a comparison with in situ observations from tide gauges and bottom pressure recorders.

a supprimé: nonetheless

a supprimé: n

a supprimé: and are in line with those reported by Armitage et al. (2016) for the Arctic.

a supprimé: comparable to Armitage's values of 0.58 and 0.64 from tide gauge comparisons in the Laptev and Kara Seas, respectively

a supprimé: on timescales longer than 10 days,

a supprimé: at least

a supprimé: As more satellite missions are incorporated in the later years, the product may improve its capacity to resolve shorter-timescale variability.

a supprimé: extreme

a supprimé: —

a supprimé: ideal

a supprimé: . Armitage et al. (2014) similarly report slopes of 0.54 and 0.49 in their Arctic comparisons

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Motif : Transparente

a supprimé: which is inherently optimized for capturing large-scale circulation. One potential contributor is the tidal correction model used to derive SLA: in complex coastal and semi-enclosed environments where tide gauges are often located, tide models typically perform poorly. Despite these limitations, the in-situ comparisons support the reliability of our dataset and confirm its ability to capture meaningful physical variability in ice-covered regions of the Southern Ocean.

a mis en forme : Anglais (E.U.)

2520 satellite missions. Figure S19 shows effective resolution results over the period 2010–2012, comparing Envisat daily maps
with independent along-track observations from CryoSat-2. The effective resolution estimates are slightly lower but of the
same order of magnitude, reaching 210 km in the 65°S–70°S latitude band. These values are consistent with the correlation
scales prescribed in the Optimal Interpolation, confirming that these parameters are the primary control on the resolved scales.
In the ACC, the 120 km estimate is comparable to the values reported by Ballarotta et al. (2019) for the C3S product at similar
latitudes, indicating that SO_SLICE achieves comparable performance where both products have sufficient observations. The
2525 effective temporal resolution, estimated from NSR analysis against in situ records, ranges from 23 to 48 days with a mean of
37 days, consistent with the prescribed temporal correlation scale of approximately 30 days. The 25 km daily gridded output
therefore oversamples the true information content of the product. Users should interpret the gridded fields accordingly:
SO_SLICE resolves ocean variability at wavelengths longer than approximately 120 km and periods longer than approximately
30 days in the ACC, with coarser resolution in the subpolar domain south of 65°S. Auger et al. (2023) used a similar product
2530 to identify mesoscale eddies in the ice-covered Southern Ocean. Our effective resolution analysis suggests that such features
are only reliably resolved at wavelengths longer than approximately 200 km in the subpolar Southern Ocean. Eddies smaller
than this scale may appear in the gridded fields if sampled at the right time, but should be interpreted with caution, as they fall
below the effective resolution of the interpolation. Despite these limitations, SO_SLICE resolves spatial and temporal scales
that no other gridded altimetric product currently captures in ice-covered regions of the Southern Ocean.

2535 This dataset represents a significant step forward in observing sea level variability in the Southern Ocean, particularly in
regions previously obscured by sea ice. With nearly two decades of consistent altimetry coverage across both open ocean and
ice-covered areas, it offers new opportunities to investigate large-scale and mesoscale ocean dynamics, trend analysis, and
interannual variability in sea level under Antarctic sea-ice. It can support studies of subpolar gyre variability (Dotto et al.,
2540 2018), Antarctic Bottom Water export, and ice-shelf–ocean interactions (Lauber et al., 2023), especially where other observing
systems (in-situ and other satellite observations) are limited. By extending the observational record and bridging the
observational gap across the ice edge, this dataset provides a valuable resource to better understand how the Southern Ocean
works and how it has been changing over the past two decades.

6 Data availability

2545 The altimetry product discussed in this article is publicly available on Zenodo at: <https://doi.org/10.5281/zenodo.17986180>.

Authors contribution

2550 CMD, J-BS and CdL designed the study. CMD developed the product. J-BS and CdL supervised the research. CMD wrote the
first draft. All the co-authors reviewed and revised the manuscript.

a supprimé: The uncertainty associated with the dataset was also evaluated. Multiple sources contribute to the total error budget, including instrumental noise, unresolved high-frequency SLA variability, orbit errors, and inaccuracies in the applied geophysical corrections. Due to the scarcity of independent reference data—particularly in ice-covered regions—it remains difficult to isolate and quantify the contribution of each component individually. Instead, we rely on the formal error output from the Optimal Interpolation (OI) process as a practical proxy for uncertainty. While this formal error provides a consistent and spatially explicit estimate, it should be considered a lower-bound uncertainty. It does not account for systematic biases in SLA retrievals or for potential inaccuracies in tide and atmospheric corrections. Based on this approach, the estimated uncertainty in the open ocean is approximately 1.5 cm, and decreases to 1.3 cm in the subpolar Southern Ocean. Off the continental shelf, the uncertainty remains relatively low (~1.2 cm), but can exceed 1.9 cm in coastal and shelf regions. These values are consistent with our independent validation using tide gauges, which show a root mean square error (RMSE) of 4.3 cm.

a supprimé: ¶

a mis en forme : Non Surlicnagne

a supprimé: ¶

a supprimé: enables

a supprimé: 's role in the climate system

a supprimé: h

a mis en forme : Anglais (E.U.)

a mis en forme : Police par défaut, Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a supprimé: <https://doi.org/10.5281/zenodo.17467408>

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

Altimeter data processing was made possible thanks to a strong collaboration with Collecte Localisation Satellite (CLS) teams, particularly Yannice Faugère, Marie-Isabelle Pujol, Pierre Veillard and Fanny Piras. Matthis Auger and Maxime Ballarotta provided valuable feedback. [Cosme Mosneron Dupin Ph.D. studentship is co-funded by the CNES \(Centre National d'Etudes Spatiales\), France. This research was supported by Ocean Cryosphere Exchanges in ANtarctica: Impacts on Climate and the Earth system, OCEAN ICE, which is funded by the European Union, Horizon Europe Funding Programme for research and innovation under grant agreement Nr. 101060452, 10.3030/101060452. OCEAN ICE contribution number 47.](#)

References

Abileah, R., Gómez-Enri, J., Scozzari, A., and Vignudelli, S.: Coherent ranging with Envisat radar altimeter: A new perspective in analyzing altimeter data using Doppler processing, *Remote Sens. Environ.*, 139, 271–276, <https://doi.org/10.1016/j.rse.2013.08.005>, 2013.

Abram, N. J., Purich, A., England, M. H., McCormack, F. S., Strugnell, J. M., Bergstrom, D. M., Vance, T. R., Stål, T., Wienecke, B., Heil, P., Doddridge, E. W., Sallée, J.-B., Williams, T. J., Reading, A. M., Mackintosh, A., Reese, R., Winkelmann, R., Klose, A. K., Boyd, P. W., Chown, S. L., and Robinson, S. A.: Emerging evidence of abrupt changes in the Antarctic environment, *Nature*, 644, 621–633, <https://doi.org/10.1038/s41586-025-09349-5>, 2025.

Androssov, A., Boebel, O., Schröter, J., Danilov, S., Macrandner, A., and Ivanciu, I.: Ocean Bottom Pressure Variability: Can It Be Reliably Modeled?, *J. Geophys. Res. Oceans*, 125, e2019JC015469, <https://doi.org/10.1029/2019JC015469>, 2020.

Armitage, T. W. K., Bacon, S., Ridout, A. L., Thomas, S. F., Aksenov, Y., and Wingham, D. J.: Arctic sea surface height variability and change from satellite radar altimetry and GRACE, 2003–2014, *J. Geophys. Res. Oceans*, 121, 4303–4322, <https://doi.org/10.1002/2015JC011579>, 2016.

Armitage, T. W. K., Kwok, R., Thompson, A. F., and Cunningham, G.: Dynamic Topography and Sea Level Anomalies of the Southern Ocean: Variability and Teleconnections, *J. Geophys. Res. Oceans*, 123, 613–630, <https://doi.org/10.1002/2017JC013534>, 2018.

Auger, M., Prandi, P., and Sallée, J.-B.: Southern ocean sea level anomaly in the sea ice-covered sector from multimission satellite observations, *Sci. Data*, 9, 70, <https://doi.org/10.1038/s41597-022-01166-z>, 2022a.

Auger, M., Sallée, J., Prandi, P., and Naveira Garabato, A. C.: Subpolar Southern Ocean Seasonal Variability of the Geostrophic Circulation From Multi-Mission Satellite Altimetry, *J. Geophys. Res. Oceans*, 127, e2021JC018096, <https://doi.org/10.1029/2021JC018096>, 2022b.

a mis en forme : Police :Non Italique, Police de script complexe :Non Italique

- Auger, M., Sallée, J., Thompson, A. F., Pauthenet, E., and Prandi, P.: Southern Ocean Ice-Covered Eddy Properties From Satellite Altimetry, *J. Geophys. Res. Oceans*, 128, e2022JC019363, <https://doi.org/10.1029/2022JC019363>, 2023.
- 2610 Bézy, J. L., Delwart, S., and Rast, M.: MERIS – A new generation of ocean-colour sensor onboard Envisat., *ESA Bull.* 103 Eur. Space Agency, 2000.
- Bretherton, F. P., Davis, R. E., and Fandry, C. B.: A technique for objective analysis and design of oceanographic experiments applied to MODE-73, *Deep Sea Res. Oceanogr. Abstr.*, 23, 559–582, [https://doi.org/10.1016/0011-7471\(76\)90001-2](https://doi.org/10.1016/0011-7471(76)90001-2), 1976.
- Brown, G.: The average impulse response of a rough surface and its applications, *IEEE J. Ocean. Eng.*, 2, 67–74, <https://doi.org/10.1109/JOE.1977.1145328>, 1977.
- 2615 Bulczak, A. I., Bacon, S., Naveira Garabato, A. C., Ridout, A., Sonnewald, M. J. P., and Laxon, S. W.: Seasonal variability of sea surface height in the coastal waters and deep basins of the Nordic Seas, *Geophys. Res. Lett.*, 42, 113–120, <https://doi.org/10.1002/2014GL061796>, 2015.
- Carrère, L. and Lyard, F.: Modeling the barotropic response of the global ocean to atmospheric wind and pressure forcing - comparisons with observations, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2002gl016473>, 2003.
- 2620 Carrere, L., Lyard, F., Cancet, M., Allain, D., Dabat, M.-L., Fouchet, E., Sahuc, E., Faugere, Y., Dibarboure, G., and Picot, N.: A new barotropic tide model for global ocean: FES2022, <https://doi.org/10.24400/527896/A03-2022.3287>, 2022.
- Cartwright, D. E. and Edden, A. C.: Corrected Tables of Tidal Harmonics, *Geophys. J. Int.*, 33, 253–264, <https://doi.org/10.1111/j.1365-246X.1973.tb03420.x>, 1973.
- 2625 Cocks, J., Silvano, A., Naveira Garabato, A. C., Dragomir, O., Schifano, N., Hogg, A. E., and Marzocchi, A.: Satellite-derived steric height in the Southern Ocean: trends, variability, and climate drivers, *Ocean Sci.*, 21, 1609–1625, <https://doi.org/10.5194/os-21-1609-2025>, 2025.
- Desai, S., Wahr, J., and Beckley, B.: Revisiting the pole tide for and from satellite altimetry, *J. Geod.*, 89, 1233–1243, <https://doi.org/10.1007/s00190-015-0848-7>, 2015.
- 2630 Donlon, C., Berruti, B., Buongiorno, A., Ferreira, M.-H., Féménias, P., Frerick, J., Goryl, P., Klein, U., Laur, H., Mavrocordatos, C., Nieve, J., Rebhan, H., Seitz, B., Stroede, J., and Sciarra, R.: The Global Monitoring for Environment and Security (GMES) Sentinel-3 mission, *Remote Sens. Environ.*, 120, 37–57, <https://doi.org/10.1016/j.rse.2011.07.024>, 2012.
- Dotto, T. S., Naveira Garabato, A., Bacon, S., Tsamados, M., Holland, P. R., Hooley, J., Frajka-Williams, E., Ridout, A., and Meredith, M. P.: Variability of the Ross Gyre, Southern Ocean: Drivers and Responses Revealed by Satellite Altimetry, *Geophys. Res. Lett.*, <https://doi.org/10.1029/2018GL078607>, 2018.
- 2635 Dragomir, O. C.: Dynamics of the subpolar Southern Ocean response to climate change, University of Southampton, <https://doi.org/10.5258/SOTON/T0085>, 2024.
- Drinkwater, M. R.: K_u band airborne radar altimeter observations of marginal sea ice during the 1984 Marginal Ice Zone Experiment, *J. Geophys. Res. Oceans*, 96, 4555–4572, <https://doi.org/10.1029/90JC01954>, 1991.
- 2640 Ducet, N., Le Traon, P. Y., and Reverdin, G.: Global high-resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and -2, *J. Geophys. Res. Oceans*, 105, 19477–19498, <https://doi.org/10.1029/2000jc900063>, 2000.

- Dwyer, R. E. and Godin, R. H.: Determining sea-ice boundaries and ice roughness using GEOS-3 altimeter data. NASA contract report CR-156862., 1980.
- Frölicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., and Winton, M.: Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Models, *J. Clim.*, 28, 862–886, <https://doi.org/10.1175/JCLI-D-14-00117.1>, 2015.
- Fu, L. L. and Cazenave, A.: *Satellite Altimetry and Earth Sciences: A Handbook of Techniques and Applications*, 2000.
- Giles, K. A., Laxon, S. W., Ridout, A. L., Wingham, D. J., and Bacon, S.: Western Arctic Ocean freshwater storage increased by wind-driven spin-up of the Beaufort Gyre, *Nat. Geosci.*, 5, 194–197, <https://doi.org/10.1038/ngeo1379>, 2012.
- Hayakawa, H., Shibuya, K., Aoyama, Y., Nogi, Y., and Doi, K.: Ocean bottom pressure variability in the Antarctic Divergence Zone off Lützow-Holm Bay, East Antarctica, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 60, 22–31, <https://doi.org/10.1016/j.dsr.2011.09.005>, 2012.
- Helm, V., Humbert, A., and Miller, H.: Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8, 1539–1559, <https://doi.org/10.5194/tc-8-1539-2014>, 2014.
- Hobbs, W., Spence, P., Meyer, A., Schroeter, S., Fraser, A. D., Reid, P., Tian, T. R., Wang, Z., Liniger, G., Doddridge, E. W., and Boyd, P. W.: Observational Evidence for a Regime Shift in Summer Antarctic Sea Ice, *J. Clim.*, 37, 2263–2275, <https://doi.org/10.1175/jcli-d-23-0479.1>, 2024.
- Holland, D. M., Nicholls, K. W., and Basinski, A.: The Southern Ocean and its interaction with the Antarctic Ice Sheet, *Science*, 367, 1326–1330, <https://doi.org/10.1126/science.aaz5491>, 2020.
- Hughes, C. W. and Ash, E. R.: Eddy forcing of the mean flow in the Southern Ocean, *J. Geophys. Res. Oceans*, 106, 2713–2722, <https://doi.org/10.1029/2000jc900332>, 2001.
- Iijima, B. A., Harris, I. L., Ho, C. M., Lindqwister, U. J., Mannucci, A. J., Pi, X., Reyes, M. J., Sparks, L. C., and Wilson, B. D.: Automated daily process for global ionospheric total electron content maps and satellite ocean altimeter ionospheric calibration based on Global Positioning System data, *J. Atmospheric Sol.-Terr. Phys.*, 61, 1205–1218, [https://doi.org/10.1016/s1364-6826\(99\)00067-x](https://doi.org/10.1016/s1364-6826(99)00067-x), 1999.
- IOC: Global Sea Level Observing System (GLOSS) Implementation Plan – 2020, IOC Tech. Ser. No 89, 2020.
- Jousset, S.: Mean Dynamic Topography MDT CNES_CLS 2022 (2022), <https://doi.org/10.24400/527896/A01-2023.003>, 2023.
- Kwok, R. and Morison, J.: Dynamic topography of the ice-covered Arctic Ocean from ICESat: DYNAMIC TOPOGRAPHY OF ARCTIC OCEAN, *Geophys. Res. Lett.*, 38, n/a–n/a, <https://doi.org/10.1029/2010GL046063>, 2011.
- Kwok, R. and Morison, J.: Sea surface height and dynamic topography of the ice-covered oceans from CryoSat-2: 2011–2014, *J. Geophys. Res. Oceans*, 121, 674–692, <https://doi.org/10.1002/2015JC011357>, 2016.
- Laloue, A., Schaeffer, P., Pujol, M.-I., Veillard, P., Andersen, O. B., Sandwell, D. T., Delepoulle, A., Dibarboure, G., and Faugère, Y.: Merging recent Mean Sea Surface into a 2023 Hybrid model (from Scripps, DTU, CLS and CNES), <https://doi.org/10.22541/au.171987154.42384510/v1>, 1 July 2024.

Lauber, J., Hattermann, T., De Steur, L., Darelius, E., Auger, M., Nøst, O. A., and Moholdt, G.: Warming beneath an East Antarctic ice shelf due to increased subpolar westerlies and reduced sea ice, *Nat. Geosci.*, 16, 877–885, <https://doi.org/10.1038/s41561-023-01273-5>, 2023.

Laxon, S.: Sea ice altimeter processing scheme at the EODC, *Int. J. Remote Sens.*, 15, 915–924, <https://doi.org/10.1080/01431169408954124>, 1994.

Le Traon, P. Y., Nadal, F., and Ducet, N.: An Improved Mapping Method of Multisatellite Altimeter Data, *J. Atmospheric Ocean. Technol.*, 15, 522–534, [https://doi.org/10.1175/1520-0426\(1998\)015%253C0522:AIMMOM%253E2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015%253C0522:AIMMOM%253E2.0.CO;2), 1998.

Longepe, N., Thibaut, P., Vadaine, R., Poisson, J.-C., Guillot, A., Boy, F., Picot, N., and Borde, F.: Comparative Evaluation of Sea Ice Lead Detection Based on SAR Imagery and Altimeter Data, *IEEE Trans. Geosci. Remote Sens.*, 57, 4050–4061, <https://doi.org/10.1109/TGRS.2018.2889519>, 2019.

Meredith, M., Sommerkon, M., and Cassotta, S.: Polar Regions, IPCC Spec. Rep. Ocean Cryosphere Chang. Clim., 2019.

Millet, B., Gray, W. R., De Lavergne, C., and Roche, D. M.: Oxygen isotope constraints on the ventilation of the modern and glacial Pacific, *Clim. Dyn.*, 62, 649–664, <https://doi.org/10.1007/s00382-023-06910-8>, 2024.

Mosneron Dupin, C. (2025). SO_SLICE : Southern Ocean Sea Level anomalies, including the ICE-covered regions (2003-2021) [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.17986180>.

Naveira Garabato, A. C., Dotto, T. S., Hooley, J., Bacon, S., Tsamados, M., Ridout, A., Frajka-Williams, E. E., Herraiz-Borreguero, L., Holland, P. R., Heorton, H. D. B. S., and Meredith, M. P.: Phased Response of the Subpolar Southern Ocean to Changes in Circumpolar Winds, *Geophys. Res. Lett.*, 46, 6024–6033, <https://doi.org/10.1029/2019GL082850>, 2019.

Newman, L., Heil, P., Trebilco, R., Katsumata, K., Constable, A., Van Wijk, E., Assmann, K., Beja, J., Bricher, P., Coleman, R., Costa, D., Diggs, S., Farneti, R., Fawcett, S., Gille, S. T., Hendry, K. R., Henley, S., Hofmann, E., Maksym, T., Mazloff, M., Meijers, A., Meredith, M. M., Moreau, S., Ozsoy, B., Robertson, R., Schloss, I., Schofield, O., Shi, J., Sikes, E., Smith, I. J., Swart, S., Wahlin, A., Williams, G., Williams, M. J. M., Herraiz-Borreguero, L., Kern, S., Lieser, J., Massom, R. A., Melbourne-Thomas, J., Miloslavich, P., and Spreen, G.: Delivering Sustained, Coordinated, and Integrated Observations of the Southern Ocean for Global Impact, *Front. Mar. Sci.*, 6, 433, <https://doi.org/10.3389/fmars.2019.00433>, 2019.

Ollivier, A., Phillips, S., Couhert, A., and Picot, N.: Assessment of Orbit Quality through the SSH calculation: POE-E orbit standards, 2015.

Peacock, N. R. and Laxon, S. W.: Sea surface height determination in the Arctic Ocean from ERS altimetry, *J. Geophys. Res. Oceans*, 109, 2001JC001026, <https://doi.org/10.1029/2001JC001026>, 2004.

Peltier, W. R.: GLOBAL GLACIAL ISOSTASY AND THE SURFACE OF THE ICE-AGE EARTH: The ICE-5G (VM2) Model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32, 111–149, <https://doi.org/10.1146/annurev.earth.32.082503.144359>, 2004.

Poisson, J.-C., Quartly, G. D., Kurekin, A. A., Thibaut, P., Hoang, D., and Nencioli, F.: Development of an ENVISAT Altimetry Processor Providing Sea Level Continuity Between Open Ocean and Arctic Leads, *IEEE Trans. Geosci. Remote Sens.*, 56, 5299–5319, <https://doi.org/10.1109/TGRS.2018.2813061>, 2018.

a supprimé: Mosneron Dupin, C.: Sea Level Anomalies in the subpolar Southern Ocean, 2003-2021, <https://doi.org/10.5281/ZENODO.17467407>, 2025.

a mis en forme : Police par défaut, Couleur de police : Texte 1

- Prandi, P., Ablain, M., Cazenave, A., and Picot, N.: A New Estimation of Mean Sea Level in the Arctic Ocean from Satellite Altimetry, *Mar. Geod.*, 35, 61–81, <https://doi.org/10.1080/01490419.2012.718222>, 2012.
- 2715 PSML: Permanent Service for Mean Sea Level (PSMSL) (2024). Tide Gauge Data. Available online at: <http://www.psml.org/data/obtaining/>. [August 1, 2024], 2024.
- Pujol, M.-I., Faugère, Y., Taburet, G., Dupuy, S., Pelloquin, C., Ablain, M., and Picot, N.: DUACS DT2014: the new multi-mission altimeter data set reprocessed over 20years, *Ocean Sci.*, 12, 1067–1090, <https://doi.org/10.5194/os-12-1067-2016>, 2016.
- 2720 Quartly, G., Rinne, E., Passaro, M., Andersen, O., Dinardo, S., Fleury, S., Guillot, A., Hendricks, S., Kurekin, A., Müller, F., Ricker, R., Skourup, H., and Tsamados, M.: Retrieving Sea Level and Freeboard in the Arctic: A Review of Current Radar Altimetry Methodologies and Future Perspectives, *Remote Sens.*, 11, 881, <https://doi.org/10.3390/rs11070881>, 2019.
- Rye, C. D., Naveira Garabato, A. C., Holland, P. R., Meredith, M. P., George Nurser, A. J., Hughes, C. W., Coward, A. C., and Webb, D. J.: Rapid sea-level rise along the Antarctic margins in response to increased glacial discharge, *Nat. Geosci.*, 7, 732–735, <https://doi.org/10.1038/ngeo2230>, 2014.
- 2725 Sallée, J.-B.: Southern Ocean Warming, *Oceanography*, 31, <https://doi.org/10.5670/oceanog.2018.215>, 2018.
- Scagliola, M., Tagliani, N., and Fornari, M.: Mesuring the effective along-track resolution of Cryosat., , March, 2013.
- Schaeffer, P., Pujol, M.-I., Veillard, P., Faugere, Y., Dagneaux, Q., Dibarboure, G., and Picot, N.: The CNES CLS 2022 Mean Sea Surface: Short Wavelength Improvements from CryoSat-2 and SARAL/AltiKa High-Sampled Altimeter Data, *Remote Sens.*, 15, 2910, <https://doi.org/10.3390/rs15112910>, 2023.
- 2730 Silvano, A., Narayanan, A., Catany, R., Olmedo, E., González-Gambau, V., Turiel, A., Sabia, R., Mazloff, M. R., Spira, T., Haumann, F. A., and Naveira Garabato, A. C.: Rising surface salinity and declining sea ice: A new Southern Ocean state revealed by satellites, *Proc. Natl. Acad. Sci.*, 122, <https://doi.org/10.1073/pnas.2500440122>, 2025.
- Smith, G. C., Allard, R., Babin, M., Bertino, L., Chevallier, M., Corlett, G., Crout, J., Davidson, F., Delille, B., Gille, S. T., 2735 Hebert, D., Hyder, P., Intrieri, J., Lagunas, J., Larnicol, G., Kaminski, T., Kater, B., Kauker, F., Marec, C., Mazloff, M., Metzger, E. J., Mordy, C., O’Carroll, A., Olsen, S. M., Phelps, M., Posey, P., Prandi, P., Rehm, E., Reid, P., Rigor, I., Sandven, S., Shupe, M., Swart, S., Smedstad, O. M., Solomon, A., Storto, A., Thibaut, P., Toole, J., Wood, K., Xie, J., Yang, Q., and the WWRP PPP Steering Group: Polar Ocean Observations: A Critical Gap in the Observing System and Its Effect on Environmental Predictions From Hours to a Season, *Front. Mar. Sci.*, 6, 429, <https://doi.org/10.3389/fmars.2019.00429>, 2019.
- 2740 Taburet, G., Sanchez-Roman, A., Ballarotta, M., Pujol, M.-I., Legeais, J.-F., Fournier, F., Faugere, Y., and Dibarboure, G.: DUACS DT2018: 25 years of reprocessed sea level altimetry products, *Ocean Sci.*, 15, 1207–1224, <https://doi.org/10.5194/os-15-1207-2019>, 2019.
- The IMBIE team: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 2745 The SO-CHIC consortium, Sallée, J. B., Abrahamsen, E. P., Allaire, C., Auger, M., Ayres, H., Badhe, R., Boutin, J., Brearley, J. A., De Lavergne, C., Ten Doeschate, A. M. M., Droste, E. S., Du Plessis, M. D., Ferreira, D., Giddy, I. S., Gülk, B., Gruber,

N., Hague, M., Hoppema, M., Josey, S. A., Kanzow, T., Kimmritz, M., Lindeman, M. R., Llanillo, P. J., Lucas, N. S., Madec, G., Marshall, D. P., Meijers, A. J. S., Meredith, M. P., Mohrmann, M., Monteiro, P. M. S., Mosneron Dupin, C., Naeck, K., Narayanan, A., Naveira Garabato, A. C., Nicholson, S.-A., Novellino, A., Ödalen, M., Østerhus, S., Park, W., Patmore, R. D., Piedagnel, E., Roquet, F., Rosenthal, H. S., Roy, T., Saurabh, R., Silvy, Y., Spira, T., Steiger, N., Styles, A. F., Swart, S., Vogt, L., Ward, B., and Zhou, S.: Southern ocean carbon and heat impact on climate, *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 381, 20220056, <https://doi.org/10.1098/rsta.2022.0056>, 2023.

Thibaut, P., Poisson, J. C., Bronner, E., and Picot, N.: Relative Performance of the MLE3 and MLE4 Retracking Algorithms on Jason-2 Altimeter Waveforms, *Mar. Geod.*, 33, 317–335, <https://doi.org/10.1080/01490419.2010.491033>, 2010.

2755 Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nat. Geosci.*, 4, 741–749, <https://doi.org/10.1038/ngeo1296>, 2011.

Tourain, C., Piras, F., Ollivier, A., Hauser, D., Poisson, J. C., Boy, F., Thibaut, P., Hermozo, L., and Tison, C.: Benefits of the Adaptive Algorithm for Retracking Altimeter Nadir Echoes: Results From Simulations and CFOSAT/SWIM Observations, *IEEE Trans. Geosci. Remote Sens.*, 59, 9927–9940, <https://doi.org/10.1109/TGRS.2021.3064236>, 2021.

2760 Tran, N., Labroue, S., Philipps, S., Bronner, E., and Picot, N.: Overview and Update of the Sea State Bias Corrections for the Jason-2, Jason-1 and TOPEX Missions, *Mar. Geod.*, 16th August, 348–362, 2010.

Veillard, P., Prandi, P., Pujol, M.-I., Daguzé, J.-A., Piras, F., Dibarboure, G., and Faugère, Y.: Arctic and Southern Ocean polar sea level maps and along-tracks from multi-mission satellite altimetry from 2011 to 2021, *Front. Mar. Sci.*, 11, 1419132, <https://doi.org/10.3389/fmars.2024.1419132>, 2024.

2765 Verron, J., Bonnefond, P., Andersen, O., Ardhuin, F., Bergé-Nguyen, M., Bhowmick, S., Blumstein, D., Boy, F., Brodeau, L., Crétaux, J.-F., Dabat, M. L., Dibarboure, G., Fleury, S., Garnier, F., Gourdeau, L., Marks, K., Queruel, N., Sandwell, D., Smith, W. H. F., and Zaron, E. D.: The SARAL/AltiKa mission: A step forward to the future of altimetry, *Adv. Space Res.*, 68, 808–828, <https://doi.org/10.1016/j.asr.2020.01.030>, 2021.

2770 Vinogradova, N. T., Ponte, R. M., and Stammer, D.: Relation between sea level and bottom pressure and the vertical dependence of oceanic variability, *Geophys. Res. Lett.*, 34, 2006GL028588, <https://doi.org/10.1029/2006GL028588>, 2007.

Wingham, D. J., Francis, C. R., Baker, S., Bouzinac, C., Brockley, D., Cullen, R., De Chateau-Thierry, P., Laxon, S. W., Mallow, U., Mavrocordatos, C., Phalippou, L., Ratier, G., Rey, L., Rostan, F., Viau, P., and Wallis, D. W.: CryoSat: A mission to determine the fluctuations in Earth’s land and marine ice fields, *Adv. Space Res.*, 37, 841–871, <https://doi.org/10.1016/j.asr.2005.07.027>, 2006.

a mis en forme : Français

a mis en forme : Anglais (E.U.)

a supprimé: