

Mapping Paddy Rice Distribution and Cropping Intensity in South and Southeast Asia (1995 - 2024) at 30m Resolution

Zizhang Zhao^{1,2}, Geli Zhang^{1*}, Jinwei Dong^{-2,3}, Jilin Yang⁻⁴, Chang Fan¹, Ruoqi Liu¹, Xiangming Xiao⁵

1 College of Land Science and Technology, China Agricultural University, Beijing 100193, China

2 Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

3 University of Chinese Academy of Sciences, Beijing 100049, China

4 College of Grassland Science and Technology, China Agricultural University, Beijing 100193, China

5 School of Biological Sciences, University of Oklahoma, Norman, OK 73019, USA

Correspondence to: Geli Zhang (geli.zhang@cau.edu.cn)

Abstract: South and Southeast Asia, a major global hub for paddy rice cultivation, exhibits the highest rice cropping intensity worldwide due to its favourable hydrothermal conditions, and also has experienced considerable spatiotemporal changes due to climate change and anthropogenic activities. However, the absence of long-term spatial distribution and cropping intensity of paddy rice hinders effective agricultural and environmental management. This gap is particularly critical especially in the 21st century, with enhanced impacts from changing climate, water resources, and food trade pattern. Using all the available Landsat and Sentinel-2 archives, we refined a phenology-based algorithm to generate 30-m rice maps and cropping intensity across South and Southeast Asia for the years 1995, 2005, 2015, and 2024. The algorithm overcomes the challenge of detecting rice cropping intensity in long time-series and comprises three core steps: (1) identifying pixel-level rice phenological peaks using an enhanced peak detection method, thereby defining potential transplanting windows and minimizing monsoon-induced cloud and precipitation interference; (2) detecting paddy flooding signals and delineating rice cultivation areas based on phenological rules derived from the relationship between the Land Surface Water Index (LSWI) and Enhanced Vegetation Index (EVI); (3) determining rice cropping intensity according to the number of valid crop peaks and associated flooding signals within a single year. The resulting maps were validated using 23,396 samples collectively derived from a field photo library, visual interpretation of Sentinel-1/2 satellite imagery, and a sample migration algorithm. Across the four periods, the maps achieved overall accuracies ranging from 83% to 87%. In addition, the resultant products were compared with existing regional and period-specific rice datasets (e.g., NESEA-RICE10 and Open-SEA-Rice-10) for further evaluation. The comparisons demonstrated that the refined approach achieved higher accuracy and robustness in mapping both rice distribution and cropping intensity, whereas the existing products performed well only in partial environments. When compared with the FAO official statistics for South and mainland Southeast Asian countries, the derived maps yielded R^2 values exceeding 0.9. This dataset holds great potential for applications such as methane emission

34 estimation, water resource management, and crop yield monitoring, thereby supporting sustainable agricultural
35 practices and policy development in the region.

1. Introduction

Rice is a staple food crop across South and Southeast Asia, underpinning food security in a region characterized by rapid population growth and escalating food demand (Xiao et al., 2006). This demographic pressure has inevitably driven the expansion of rice paddies and intensified cultivation practices (Godfray et al., 2010; Foley et al., 2005). However, the expansion of agricultural land, including rice paddies, exacerbates environmental challenges, such as freshwater depletion, deforestation, and increased methane emissions, all of which threaten ecosystem functions (Mehta et al., 2024; Wang et al., 2023; Zeng et al., 2018; Chen et al., 2024). Balancing food security with the preservation of ecosystem services is thus a critical component of achieving the United Nations' 2030 Sustainable Development Goals (SDGs) (Persaud and Dagher, 2021). Understanding the spatial extent and cropping intensification patterns of rice agriculture is essential to provide robust data support for sustainable development policies (Potapov et al., 2022; Zabel et al., 2019).

Satellite remote sensing has emerged as a powerful tool for accurately mapping rice agriculture over large spatial scales (Dong and Xiao, 2016; Zhang et al., 2017). Moderate Resolution Imaging Spectroradiometer (MODIS) data, with its 500-meter spatial resolution, has been effectively utilized to map rice distributions in monsoon Asia, including South and Southeast Asia (Zhang et al., 2020). However, the coarse resolution of MODIS limits its ability to capture fragmented rice paddies and introduces errors due to mixed-pixel effects, particularly in detecting multi-cropping systems (Han et al., 2021). Higher-resolution satellite data, such as those from Landsat (30 m) and Sentinel (10–20 m), enable more precise identification of rice paddies (Zhao et al., 2024). Current research employs phenology-based algorithms or machine learning approaches to leverage these higher-resolution datasets for rice mapping.

Phenology-based rice mapping methods primarily exploit the relationship between the Land Surface Water Index (LSWI) and the Enhanced Vegetation Index (EVI) derived from optical imagery to detect the unique flooding signals of rice paddies (Dong et al., 2015; Xiao et al., 2005). These methods have achieved high accuracy in regions such as Northeast China, Japan, and South Korea (Dong et al., 2016; Carrasco et al., 2022; Han et al., 2022). Recent studies have extended phenology-based approaches to Synthetic Aperture Radar (SAR) data, improving rice mapping in cloud-prone and rainy regions (Xu et al., 2023; Song et al., 2025). Due to their simplicity, computational efficiency, and robustness, phenology-based methods are particularly well-suited for large-scale applications and can be readily implemented on cloud-computing platforms, making them the most widely adopted approach for regional and continental-scale rice mapping studies (Zhao et al., 2025a).

Machine learning-based rice classification methods have also gained prominence, as they leverage extensive training samples to achieve high accuracy in mapping paddy rice distribution (Dong and Xiao, 2016). For instance, several studies have achieved high-precision rice mapping in regions with complex cropping patterns, such as the crop rotation systems in Northeast China and the multi-season rice planting patterns in the Jiangnan Plain, by integrating rice-specific phenological features with machine learning classifiers, such as One-Class Support Vector Machines (OC-SVM) and Random Forest (He et al., 2021; Ni et al., 2021). More recently, deep learning techniques, particularly convolutional neural networks like U-Net and eXplainable Mamba UNet, have been employed to process time-series Sentinel-1 SAR data, demonstrating remarkable robustness in capturing spatial patterns, even in cloud-prone regions where optical imagery is limited (Ge et al., 2025; Thorp and Drajat, 2021; Lin et al., 2022). Despite the achievements of the above two mapping approaches, none has yet provided a method capable of long-term, large-scale analysis with precise extraction of cropping intensity and spatial distribution to elucidate the spatiotemporal patterns of rice distribution in South and Southeast Asia. Traditional phenology-based rice mapping methods rely on prior expert knowledge to determine phenological stages, but interpreting phenological periods from several decades ago introduces considerable uncertainty. Machine learning methods, on the other hand, have shown potential for global-scale classification, yet the acquisition of large, high-quality historical training datasets remains time-consuming and resource-intensive (Zhao et al., 2025; Zhan et al., 2021). Even though a fully automated, sample-free rice-mapping framework has been developed, its dependence on SAR-derived rice features limits its applicability to periods beyond the operational lifetime of Sentinel-1 (Gao et al., 2023). This study addresses this gap by employing a refined phenology-based method that eliminates the need for extensive training samples while accurately identifying planting intensity and phenological windows. Through optical imagery fusion and false peak elimination techniques, we aim to precisely identify rice cropping systems and their spatial distributions, while quantifying changes over the past three decades. This approach provides critical data to reconcile the conflict between traditional rice cultivation expansion and sustainable development goals, offering robust support for informed policy-making in South and mainland Southeast Asia.

2. Materials

2.1 Study area

~~The study area encompasses South and mainland Southeast Asia countries, specifically Vietnam, Thailand, Cambodia, Myanmar, Laos, Bangladesh, India, and Pakistan (Fig. S1). These eight countries are predominantly~~

characterized by a tropical monsoon climate (with the exception of Balochistan Province), featuring distinct wet and dry seasons. Rice is the predominant crop in this region, with the highest cropping intensity globally. In 2024, the total rice harvested area in this region accounted for approximately 47% of the global rice harvested area, making it the world's most critical rice production base. The study area encompasses South and mainland Southeast Asia, including Vietnam, Thailand, Cambodia, Myanmar, Laos, Bangladesh, India, and Pakistan (Fig. 1). These eight countries are predominantly characterized by a tropical monsoon climate, with the exception of Balochistan Province, and feature distinct wet and dry seasons. Rice is the dominant crop in this region and exhibits the highest cropping intensity worldwide. The region also includes diverse rice production systems, such as the rice–winter wheat rotation system in northern India, flood-associated single-season rice cultivation in Bangladesh, and typical multi-season rice systems in the Mekong River Basin. In 2024, the total rice harvested area in this region accounted for approximately 47% of the global rice harvested area, making it the world's most important rice production base.




Figure 1. (a) Spatial distribution of rice harvested area in South and Southeast Asia ($\times 10^5$ ha). Insets show

设置了格式: 英语(英国)

带格式的: 居中

107 example rice fields at the locations marked by orange squares. (b–d) Representative NDVI (green) and
108 LSWI (gray) time series for (b) rice-wheat cropping system, (c) single-season rice, and (d) multi-season
109 rice. The locations of the example pixels are indicated by arrows on the map.

设置了格式: 字体: (中文) + 中文正文 (等线), (中文) 简体中文(中国大陆), (其他) 英语(美国)

111 **2.2 Data and preprocessing**

112 **2.2.1 Satellite imagery**

113 To develop long-term rice mapping products for South and mainland Southeast Asia, we acquired all available
114 Sentinel-2 (S2_HARMONIZED) and Landsat T1_L2 imagery data from the Google Earth Engine (GEE) platform
115 for specific time periods (Table 1). Invalid observations, including clouds, cloud shadows, and snow cover, were
116 filtered out using the Sentinel-2 QA60 and Landsat QA_PIXEL bands. However, because imagery from a single
117 year was insufficient to reliably support rice information extraction across South and mainland Southeast Asia, we
118 applied a multi-year median compositing approach for the periods 1993–1997, 2003–2007, 2014–2016, and 2023–
119 2025 to mitigate data gaps (see Table 1). Specifically, each year was segmented into half-monthly intervals (e.g.,
120 January 1–15 and January 16–31, with some intervals spanning 14 or 16 days due to monthly variations), and all
121 valid observations within the selected time periods for each interval were used to generate a single median
122 composite image. Additionally, linear regression was employed to harmonize the spectral bands of Landsat and
123 Sentinel-2 data (Yang et al., 2023), resulting in the generation of half-monthly time-series curves.

124 **Table 1:** Satellite sensors and composite periods for rice mapping in South and mainland Southeast Asia

Reference Year	Satellite Sensors	Composite Period
1995	Landsat-5	1993-1997
2005	Landsat-5, -7	2003-2007
2015	Landsat-5, -7, -8	2014-2016
2022 2024	Sentinel-2, Landsat-8	2023-2025

125
126 Based on the generated half-monthly time-series curves, we further calculated relevant vegetation indices to
127 identify the spatial distribution and cropping intensity of paddy rice, including NDVI, LSW, and EVI (Eqs. 1–3).

128
$$NDVI = \frac{NIR - Red}{NIR + Red} \tag{1}$$

129
$$LSWI = \frac{NIR - SWIR}{NIR + SWIR} \tag{2}$$

130
131
132
133
134
135
136
137
138
139

140
141
142
143
144
145
146
147
148
149

$$EVI = 2.5 \times \frac{NIR - Red}{NIR + 6 \times Red - 7.5 \times Blue + 1} \tag{3}$$

2.2.2 Cropland maskAuxiliary data

Cropland Mask. To investigate changes in rice planting patterns in South and mainland Southeast Asia, a region that has undergone significant land-use transformations, including cropland expansion and forest loss over recent decades, we selected high-quality land cover and cropland products aligned with the target mapping years. Four global products were chosen to mitigate uncertainties in rice mapping arising from cropland expansion. The specific cropland products selected for each year are detailed in Table 2 (Zhang et al., 2021; Karra et al., 2021; Potapov et al., 2022) (<https://zenodo.org/records/5571936>).

Table 2: Cropland Products Used for Cropland Mask in South and mainland Southeast Asia

Reference Year	Cropland Products
1995	GLC_FCS30 (1990, 1995)
2005	GLC_FCS30 (2005), GLAD 30 (2003)
2015	GLC_FCS30 (2015), GLAD 30 (2015)
2024	ESA (2021), GLAD (2019), ESRI (2021)

~~Rice products and statistical data.~~ To evaluate the reliability of our mapping results, we conducted spatial comparisons with several established rice mapping products, including: (1) the JAXA High-Resolution Land-Use and Land-Cover Map, including the 2020 Vietnam and 2023 Southeast Asia products (https://www.core.jaxa.jp/ALOS/en/dataset/hulc_e.htm); and (2) Li et al. (2025) and Sun et al. (2023), whose datasets (NESEA-RICE10 (Han et al., 2021) and OPEN-SEA-RICE10 (Ginting et al., 2025)) focus on regional-scale rice area mapping. The comparisons allowed us to assess the spatial consistency and intensity agreement across products. In addition, national-level statistical data from the Food and Agriculture Organization (FAO) were used to validate the mapped rice areas.

设置了格式: 字体: (中文) Times New Roman, 小四
设置了格式: 字体: 小四

带格式的: 缩进: 首行缩进: 0 字符

域代码已更改

150

151

152

154

带格式的: 无孤行控制

160

165

174

175




Figure 24. Flowchart of the mapping process for long-term rice spatial distribution and planting intensity in South and mainland Southeast Asia

2.3.2 Phenology-based approach for identifying rice fields and their cropping intensity

The phenology-based rice mapping method identifies rice by detecting the unique biophysical characteristic of fields being flooded during the transplanting period. In this study, we first selected imagery data within a 90-day time window preceding the peak date of each growing season. Subsequently, flood signal detection was performed on the imagery data within this time window by applying the rule $LSWI + 0.05 \geq EVI$ (Xiao et al., 2005; Xiao et al., 2006; Dong and Xiao, 2016). Notably, the LSWI and EVI used here differ from those employed in crop peak identification, as these vegetation indices (VIs) do not require smoothing. Additionally, to minimize interference from factors such as precipitation and soil background value, we applied the rule that, when a flood signal is detected, EVI must be ≤ 0.35 and LSWI must be > 0.05 (Zhao et al., 2025). Pixels meeting these conditions were classified as rice (pixel value = 1).

For rice cropping intensity, we integrated multi-season rice data by calculating the number of times a pixel was classified as rice (value = 1) within a year. This process generated the final rice cropping intensity distribution map, with values ranging from 1 to 3, representing single-, double-, or triple-season rice, respectively.

2.4 Confidence Layer Generation

Optical-based rice mapping is often affected by cloud contamination, which reduces the availability of valid observations during critical phenological stages. To support downstream applications and provide an estimate of data reliability, an observation-based confidence layer was generated. For each pixel, the number of valid observations used in the time-series analysis was calculated within the 90-day temporal window preceding the identified crop peak. Pixels with more valid observations generally provide more reliable phenological signals for rice detection; therefore, the observation frequency was used as a proxy for mapping confidence. The observation count was further normalized by the total number of compositing intervals to produce a confidence score ranging from 0 to 1.

$$confidence = \frac{N_{obs}}{N_{max}}$$
Where N_{obs} represents the number of valid observations within the 90-day window before the detected crop peak, and N_{max} is the total number of compositing intervals in that window.

2.3.35 Accuracy assessment and comparisons

Validation samples. Validation samplesSample points for 2024 were primarily derived collected through visual interpretation using multiple complementary data sources. First, false-color composites of Sentinel-2 imagery (R/G/B = SWIR1, NIR, RED) were generated to represent different rice growth stages, in which rice fields during the flooded transplanting period typically appear dark green.representing multiple rice growth stages were generated, in which rice fields during the flooded transplanting stage appeared dark green. Second, following the approach of (Sun et al., 2023), Sentinel-1 VH time series were used to calculate the maximum, minimum, and variance values, which were combined into RGB composites (R: VH_max, G: VH_min, B: VH_variance).we used Sentinel-1 VH time series to compute the maximum, minimum, and variance values, and composed them as R: VH_max, G: VH_min, and B: VH_variance. In these composites, flooded rice fields generallytypically appeared purple, providing an effective means of identifying rice in persistently cloudy and rainy tropical regions.which is

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 10 磅, 英语(英国)
带格式的: 行距: 1.5 倍行距, 孤行控制

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 非倾斜

设置了格式: 字体: (中文) Times New Roman, 10 磅, 英语(英国)

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅, 英语(英国)

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅

设置了格式: 字体: (中文) Times New Roman, 10 磅, 英语(英国)

设置了格式: 字体: (中文) + 中文正文 (等线), (中文) 简体中文(中国大陆)

设置了格式: 字体: 非加粗

217 ~~particularly useful for identifying rice in persistently cloudy and rainy tropical regions.~~ An example of this
218 visualization approach has been implemented and is available at the following link:
219 (<https://code.earthengine.google.com/7dd210998c6812da2e79ebeb1536822>).
220 Based on these two sets of composites, rice sample points were manually labeled and further cross-validated using
221 ultra-high-resolution Google Earth imagery and the Global Geo-Referenced Field Photo Library
222 (<http://www.eomf.ou.edu/photos/>). In total, 4,000 rice samples were collected across the study area.
223 For the remaining 4,000 non-rice samples, random sampling was first conducted within the study area, and each
224 point was then visually verified using the same image composites to ensure the absence of rice-related features.
225 Although the above approach was effective for sample generation, manually labeling samples for periods before
226 the launch of Sentinel-1 and Sentinel-2 satellites was labor-intensive and constrained by the limited availability of
227 valid imagery. Therefore, a sample migration algorithm (Huang et al., 2020) was employed to transfer the 8,000
228 manually labeled samples to other target years. ~~To ensure the reliability of the validation results, an additional set~~
229 ~~of 2,371 validation samples was independently generated through manual visual interpretation for the years 1995,~~
230 ~~2005, and 2015. These samples were used solely for independent accuracy assessment, and the corresponding~~
231 ~~validation results are presented in the Appendix (Table S1).~~
232 Rice products and statistical data. To evaluate the reliability of our mapping results, we conducted spatial
233 comparisons with several established rice mapping products, including: (1) the JAXA High-Resolution Land-Use
234 and Land-Cover Map, including the 2020 Vietnam and 2023 Southeast Asia products
235 (https://www.eorc.jaxa.jp/ALOS/en/dataset/lulc_e.htm); and (2) Li et al. (2025) and Sun et al. (2023), whose
236 datasets (NESEA-RICE10 (Han et al., 2021) and OPEN-SEA-RICE10 (Ginting et al., 2025)) focus on regional-
237 scale rice area mapping. The comparisons allowed us to assess the spatial consistency and intensity agreement
238 across products. In addition, national-level statistical data from the Food and Agriculture Organization (FAO) ~~were~~
239 ~~employed to evaluate the consistency between mapped rice areas and official statistical records.~~ ~~were used to~~
240 ~~validate the mapped rice areas.~~

设置了格式: 字体: (默认) Times New Roman, (中文) + 中文正文 (等线), 非加粗, 英语(美国)

设置了格式: 字体: (默认) Times New Roman, (中文) + 中文正文 (等线), 非加粗, 英语(美国)

设置了格式: 字体: (默认) Times New Roman, (中文) + 中文正文 (等线), 非加粗, 英语(美国)

设置了格式: 字体: 非加粗, 英语(美国)

域代码已更改

3. Results

3.1 Validation of the spatial distribution accuracy of rice in South and mainland Southeast Asia

The spatial distribution and cropping intensity of rice across South and mainland Southeast Asia are illustrated in Fig. 2. The interactive version of this map can also be accessed and visualized using the GEE platform at: <https://ee-zhaozhangcau.projects.earthengine.app/view/rice-planting-intensity--south--southeast-asia>.

Based on validation using 23,396 sample points, the mapping accuracies for different years are summarized in Table 3, with overall accuracies ranging from 83.74% to 87.60% and F1-scores between 0.7872 and 0.8628. The 2024 results achieved the highest accuracy, primarily because the higher temporal frequency of Sentinel-2 observations enabled more effective detection of flooding signals compared with the earlier Landsat sensors. Mapping accuracies for other years did not show a declining trend over time, mainly due to the aggregation of multi-year image archives and the use of biweekly composite imagery, which stabilized rice extraction performance. Although Landsat-8 data were incorporated for the 2015 (2014–2016) period, residual striping effects from Landsat-7 imagery slightly degraded the classification accuracy.





Figure 32. Long-term spatial distribution and planting intensity of rice in South and mainland Southeast Asia (1995, 2005, 2015, 2024)

260 **Table 3** Accuracy assessment of paddy rice mapping results in South and Southeast Asia

Year	Overall Accuracy	User Accuracy	Producer Accuracy	F1 Score	Sample size
2024	0.8760	0.9668	0.7788	0.8628	8000
2015	0.8374	0.9500	0.7129	0.8143	7821
2005	0.8461	0.9122	0.6936	0.7881	3814
1995	0.8378	0.9011	0.6993	0.7872	3761

261
262



263
264
265

Figure 43. Comparison of RS-based rice area estimates with FAO statistics for South and mainland Southeast Asia (1995, 2005, 2015, 2024)

266 To further evaluate the accuracy of our results, we compared the estimated rice area from our generated rice
267 product with FAO statistical data (Fig. 2). The results showed that the R^2 values for all four years exceeded 0.9,
268 with a multi-year average RMSE of approximately 4.15 million hectares. Based on remote sensing retrievals, the
269 rice planting area across South and mainland Southeast Asian countries has increased by approximately 22.5
270 million hectares since the 1990s, with the largest increases observed in India and Bangladesh. Among these, the
271 2024 results exhibited the smallest discrepancy with statistical data. In Bangladesh, the remote sensing estimates
272 were relatively lower, potentially due to the influence of frequent flooding and cloud cover in the region.

273 **3.2 Comparison with other rice maps**


274 ~~We selected typical rice-growing areas within the study region for visual comparison with other rice mapping~~
275 ~~product. Specifically, in Vietnam, we compared our results with the land cover product (including rice layer)~~
276 ~~released by JAXA. The spatial distributions were highly consistent across multiple years. In the 2005 results for~~
277 ~~northern Vietnam, our product reduced some noise through image compositing and phenological identification~~
278 ~~(Fig. S2). In the 2024 product for India, our results exhibited spatial distributions comparable to those of Li et al.~~
279 ~~(2025) (Fig. 4). However, in the 1995 mapping of Punjab and other northern Indian provinces, Li's results were~~
280 ~~notably affected by limited observation conditions, which manifested as strip-like artifacts in the output. By~~
281 ~~contrast, our approach, which employed multi-year data compositing, effectively mitigated this issue. Differences~~
282 ~~observed in the southeastern coastal regions of India further highlight the advantages of multi-year compositing~~
283 ~~for rice mapping in the 1990s.~~We selected several typical rice-growing areas within the study region for visual
284 comparison with other rice mapping products. In Vietnam, our results were compared with the JAXA land-cover
285 product, including its rice layer, and showed high spatial consistency across multiple years (Fig. S1).In addition,
286 our mapped rice distribution exhibits good spatial correspondence with rice patterns derived from Sentinel-1
287 observations (Fig. S2).
288 For India in 2024, the spatial distribution of rice in our product was generally consistent with that reported by Li
289 et al. (2025) (Fig. 4). However, for the 1995 mapping of Punjab and other northern Indian regions, Li's results
290 were visibly affected by limited observation availability, resulting in strip-like artifacts. In contrast, our multi-year
291 compositing approach largely mitigated this issue. In particular, rice pixels identified in our product show higher
292 spatial agreement with dark-green areas in Landsat false-color composites, which indicate flooding signals during
293 the transplanting stage.

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 英语(英国)

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 英语(英国)

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 英语(英国)

设置了格式: 字体: (默认) 宋体, 英语(美国)






Figure 54. Comparison of rice distribution maps: Our Results compared to Li et al. (2025) for South Asia (1995, 2024)

Notably, Fig. 5 presents a spatial comparison of five rice mapping products across Southeast Asia, including our results. Among them, products a1, b1, and e1 exhibit relatively similar spatial distributions, whereas the NESEA-Rice10 product captures the smallest rice extent. In contrast, c1 and d1 identified a greater number of rice pixels,

303 particularly in central–eastern Thailand, where both products detected extensive rice areas. We attribute these
304 differences mainly to the distinct responses of irrigated (paddy) and rainfed rice systems. The e2 panel, derived
305 from the JAXA LULC dataset, provides supporting evidence for this interpretation, as it delineates these regions
306 predominantly as paddy fields (see Section 4.1 for further discussion). Additional pixel-level visual comparisons
307 are provided in Fig. S3.

308 Panels a3–e3 illustrate the rice-growing regions in western Thailand, where three intensity-based products
309 consistently identified multi-season rice cultivation. However, the c3 product shows clear image boundary artifacts,
310 likely due to inconsistencies in image mosaicking. ~~Spatially, the b3 product—derived from a single-year~~
311 ~~dataset—identified a smaller extent of rice cultivation, while the d3 product appears to have overestimated rice~~
312 ~~coverage. In contrast, along the moisture-rich coastal areas of Myanmar, all products exhibited relatively consistent~~
313 ~~spatial patterns with minimal discrepancies. Spatially, the b3 product, which was derived from a single-year dataset,~~
314 ~~identified a smaller extent of rice cultivation, while the d3 product appears to have overestimated rice coverage.~~
315 In contrast, along the moisture-rich coastal areas of Myanmar, all products exhibited relatively consistent spatial
316 patterns with minimal discrepancies.




Figure. 65 Comparative analysis of single and multiple rice field Distribution in Southeast Asia. Panels

(a1)-(a4) display results from our proposed method, (b1)-(b4) from NESEA-Rice10, (c1)-(c4) from Open-SEA-

Rice-10, and (d1)-(d4) from Sun et al. (2023). Panels (e1)-(e4) show JAXA-LULC data, retaining the original

land cover classifications of Single Paddy Field and Multi Paddy Field.

3.3 Spatial patterns and change trends of rice cultivation over the past 30 years

In South and mainland Southeast Asia, rice is predominantly distributed in plains and riverine areas with abundant freshwater resources. Since 1990, there has been a significant expansion trend in rice planting areas, primarily in central-western India and Pakistan (Fig. 2). Rice cropping intensity is dominated by double and single rice (this study's statistics focus solely on rice cropping intensity, excluding other crops). The Mekong Delta region in Vietnam is predominantly characterized by triple rice cultivation.

Based on the analysis of Figs. 2 and 6, substantial changes in rice cultivation patterns have occurred across South and mainland Southeast Asia over the past three decades. In these regions, intensification within existing rice croplands—manifested as increased planting intensity in stable areas—has become a more prevalent trend than the expansion of rice-growing areas. For instance, regions previously dominated by single-cropping systems, such as southern Thailand, have transitioned toward double-cropping regimes. Likewise, northern Pakistan and eastern India have shown a pronounced increase in rice cropping intensity. In contrast, the Mekong Delta has experienced a shift from double- to triple-cropping systems. Areas with reduced rice extent are primarily associated with cropland degradation or conversion to other crops rather than a decline in planting intensity. The most notable reductions are concentrated in eastern Thailand and the southern Mekong Delta estuary, corresponding respectively to shifts from irrigated to rainfed systems and a loss of arable land.

As shown in Fig. 6, we further quantified changes in rice cultivation area and mean cropping intensity at the national scale. Among all countries, India exhibited the most substantial expansion in rice cultivation area, whereas most Southeast Asian countries experienced a general decline over the past three decades. In terms of cropping intensity, India showed an average increase of approximately 0.1, while most Southeast Asian countries exhibited a decrease of less than -0.05 , reflecting intensified cultivation in South Asia and widespread contraction in Southeast Asia.




Figure 76. Spatial Distribution and Planting Intensity Changes of Paddy Rice in South and mainland Southeast Asia (1995–2024). Orange represents pixels with decreased paddy rice area in 2024 compared to 1995 (contraction), green indicates increased areas (expansion), and gray-white denotes stable areas (paddy rice present in both periods). Rice cropping intensity changes are shown for stable areas, with red indicating increases (+1 for one additional season, +2 for two additional seasons) and blue indicating decreases (-1 for one season less, -2 for two seasons less).

4. Discussion

4.1 Differences between Rainfed and Irrigated Rice

In the results, substantial discrepancies among rice mapping products were observed in central–western Thailand. To investigate the underlying causes, we analyzed the time-series profiles of selected sample points over three consecutive years using the JAXA land-cover product, our phenology-based rice maps, and the Open-SEA-Rice-10 dataset (Fig. 8).

From the optical time series, it is evident that rainfed rice fields do not maintain stable surface water coverage during the “transplanting” stage. Consequently, they often fail to produce sufficiently low SWIR reflectance relative to the RED band, which is required for satisfying the condition $LSWI \geq NDVI$ —the key indicator of flooding used in phenology-based optical algorithms such as NESEA-Rice10. This explains why our optical-based

approach and other similar methods tend to identify fewer rice pixels compared with radar-based products shown in Fig. 5.

By contrast, radar observations capture a distinct VH backscatter trough, indicating an increase in soil moisture that alters the soil dielectric constant. As crop growth progresses, VH backscatter gradually rises, resulting in large VH time series variations. This enables radar-based methods to detect rainfed rice, even in areas without persistent standing water. Moreover, these VH minima typically coincide with the onset of the monsoon season, marking the beginning of the rice growth cycle (Fig. 7).

For Irrigated rice, each valid cropping cycle is accompanied by a strong flooding signal that can be effectively identified by all water-related optical indices (e.g., combinations of SWIR, RED, or GREEN bands; Zhao et al. (2025)). In radar observations, paddy fields exhibit deeper VH troughs than rainfed fields, and the subsequent increase after the trough (rice growing stage) reflects higher canopy water content (VWC) and a stronger dielectric response, which amplifies VH time series variations. Unlike rainfed rice, the timing of Irrigated rice cultivation is typically decoupled from monsoon onset, owing to irrigation management.

However, we also note a potential source of misclassification when relying solely on Sentinel-1 VH backscatter for rice mapping, especially in temperate regions outside tropical zones. Although radar-based algorithms perform well in persistently cloudy and humid environments, our previous findings—and those of other studies—indicate that several non-rice crops (both rainfed and irrigated, such as maize and soybean) can exhibit amplitude variations comparable to those of paddy rice (Zhao et al., 2024). Consequently, such V-shaped radar signatures are not unique to rainfed or paddy rice, but rather represent a general biophysical response to canopy development and surface moisture dynamics.




Figure 87. Spatial comparison and time-series analysis of vegetation indices, VH backscatter, and total precipitation between rainfed and irrigated (paddy) rice in Thailand. (a) JAXA land-cover product, where pink represents non-paddy croplands and blue indicates paddy fields. (b) Comparison between the Open-SEA-Rice-10

dataset (orange) and this study's paddy rice product (blue; overlaid on orange). The lower two panels present the corresponding time-series curves for rainfed and paddy rice, respectively.

4.2 Advantages of the proposed product


Recent advances in remote sensing have produced numerous high-accuracy rice mapping algorithms (Deng et al., 2025). However, their application to long-term, large-scale mapping remains constrained by data availability and computational demands. For instance, although SAR can effectively improve mapping accuracy (Adrian et al., 2021), its use is limited in earlier periods due to the absence of Sentinel-1 observations. In addition, the computational complexity of large-scale mapping remains a considerable challenge, even when implemented on cloud-based platforms such as Google Earth Engine.

To overcome these challenges, this study employs a streamlined and phenology-based rice mapping framework specifically optimized for the biophysical and climatic conditions of South and Southeast Asia. Instead of extracting detailed phenological phases (e.g., tillering or harvest), the method focuses on detecting key phenological peaks, providing a robust solution under conditions of high cropping intensity, short intervals between multiple rice cycles, and persistent monsoon-related cloud cover. For instance, in the Mekong Delta, the interval between two consecutive rice seasons is perhaps shorter than 15 days (Fig. S43), meaning that even a slight temporal offset in detecting the flooded transplanting stage could miss the short-lived flooding signal. Thus, identifying phenological peaks offers a more reliable and practical strategy than full-season tracking using Landsat imagery.

Another aspect that merits consideration is the effect of multi-year image compositing. As shown in the comparison with NESEA-Rice10 (Fig. 5), although both studies employ flooding-signal-based approaches, our results identify a larger number of rice pixels in some regions, mainly due to the increased availability of valid observations. This effect is particularly evident for the 1990s. As illustrated in Fig. 9, rice maps derived from single-year imagery show pronounced spatial instability, especially for 1995, owing to severe limitations in observation frequency. In contrast, the five-year composite produces a more spatially coherent and reliable rice distribution. Under such data-limited conditions, the uncertainty associated with single-year mapping exceeds the potential smoothing effect introduced by multi-year compositing.

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman



带格式的: 居中

414 Figure 9. Comparison of rice maps derived from single-year imagery and five-year composite imagery in the
415 mid-1990s.

416

417 **Example: limitation of fixed phenology windows.** Another advantage of the proposed framework is the use of
418 dynamically detected crop peaks rather than predefined phenological calendars. To illustrate this point, we
419 conducted a comparison using transplanting windows derived from the RiceAtlas phenology database (hereafter
420 referred to as the RiceAtlas approach).

421 As shown in Fig. 10, the RiceAtlas approach relies on fixed planting windows (e.g., DOY 196–227 for the first
422 season in the example location). However, during the monsoon period, temporary flooding events may cause LSWI
423 to exceed EVI even in non-rice areas. When flooding detection is applied within these predefined windows, such
424 transient signals can be incorrectly interpreted as rice transplanting events.

425 In contrast, the proposed method first identifies valid crop peaks from the vegetation index time series and then
426 restricts flooding detection to phenologically consistent crop growth cycles. This strategy effectively reduces the
427 influence of seasonal flooding signals. In the example shown in Fig. 10, the RiceAtlas-based approach identifies
428 an additional rice season, leading to an overestimation of cropping intensity, whereas our method avoids this flood-
429 induced overestimation.

- 设置了格式: 字体: (中文) Times New Roman
- 设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 10 磅, 英语(英国)
- 设置了格式: 字体: (中文) Times New Roman, 英语(英国)
- 带格式的: 正文, 居中, 段落间距段前: 0 磅, 段后: 0 磅, 行距: 1.5 倍行距
- 带格式的: 正文, 居中, 段落间距段前: 0 磅, 段后: 0 磅
- 设置了格式: 字体: 加粗




Fig. 10. Time-series comparison between the RiceAtlas-based approach and the proposed peak-detection framework at an example location affected by seasonal flooding.

4.3.2 Uncertainty Assessment

The primary source of uncertainty or error in the products generated by this study stems from the observation quality and the number of valid observations from the optical imagery data used. Specifically, the striping issue in Landsat-7 data led to omission errors in some rice fields. Despite using all available imagery over a five-year period for compositing, striped gaps persisted in certain areas, resulting in the failure to effectively identify rice fields, particularly in the results for 2005 and 2015 (Fig. S8).

Additionally, validation based on sample points revealed omission errors in northern India, due to limitations in valid optical observations. Based on the extracted rice phenological peaks, this region is characterized by a rice–wheat double-cropping system, with rice transplanting typically occurring between June and July. However, as shown in Fig. S4, many pixels in this region have very few valid observations during this period because of

设置了格式: 字体: 非加粗

带格式的: 居中

带格式的: 正文, 居中, 段落间距段前: 0 磅, 段后: 0 磅, 行距: 1.5 倍行距

设置了格式: 英语(美国)

443 persistent cloud cover. Consequently, optical imagery often fails to capture the flooding signals of rice, leading to
444 underestimation of rice extent and reduced accuracy in northern India.
445 To address this issue, we re-mapped the 2024 rice intensity and spatial distribution for northern India using our
446 previously developed Rice-Sentinel algorithm, which effectively integrates Sentinel-1 radar observations to
447 compensate for missing optical data. Nevertheless, the original optical-based results are also retained and provided
448 for comparison and reference by other researchers.
449

450 **5. Data availability**

451 The Spatial distribution and cropping intensity maps of rice in South and mainland Southeast Asia can be accessed
452 in the Zenodo data set from the following DOI: <https://doi.org/10.5281/zenodo.17615341> (Zhao et al., 2025b). We
453 welcome other researchers to validate these products and collaborate in improving the accuracy of rice mapping
454 in this region.

455 **6. Code availability**

456 The code independently developed of this study is available at the following link:
457 <https://code.earthengine.google.com/d410ce31604e93518ef9281097e71cc5>.

458 **7. Conclusions**

459 This study employed an improved phenology-based rice mapping algorithm, leveraging the full archive of Landsat
460 series and Sentinel-2 satellite data, to generate high-resolution maps of rice planting areas and cropping intensity
461 across South and mainland Southeast Asia for the years 1995, 2005, 2015, and 2024. By applying optical image
462 compositing and a fake-peak filtering technique, the challenges of extracting long-term rice cropping intensity in
463 tropical regions were effectively addressed, yielding relatively reliable results. The resulting products achieved
464 overall accuracies ranging from 83.74% to 87.60% across the four periods, validated against 23,396 independent
465 samples, and exhibited R^2 values exceeding 0.9 when compared with FAO statistical data for South and mainland
466 Southeast Asian countries. In conclusion, this study fills a critical gap in high-resolution, long-term rice mapping,
467 providing a robust foundation for sustainable agricultural practices and policy development in South and Southeast
468 Asia, with broader implications for global food security and environmental sustainability.

Adrian, J., Sagan, V., and Maimaitijiang, M.: Sentinel SAR-optical fusion for crop type mapping using deep learning and Google Earth Engine, *ISPRS Journal of Photogrammetry and Remote Sensing*, 175, 215-235, <https://doi.org/10.1016/j.isprsjprs.2021.02.018>, 2021.

Carrasco, L., Fujita, G., Kito, K., and Miyashita, T.: Historical mapping of rice fields in Japan using phenology and temporally aggregated Landsat images in Google Earth Engine, *ISPRS Journal of Photogrammetry and Remote Sensing*, 191, 277-289, <https://doi.org/10.1016/j.isprsjprs.2022.07.018>, 2022.

Chen, Z., Balasus, N., Lin, H., Nesser, H., and Jacob, D. J.: African rice cultivation linked to rising methane, *Nature Climate Change*, 14, 148-151, 10.1038/s41558-023-01907-x, 2024.

Deng, Y., Peng, B., Guan, K., Runkle, B. R. K., Moreno-García, B., Wu, X., Wang, S., Zhou, Q., and Reba, M. L.: Detecting the onset of rice field inundation in the Lower Mississippi River Basin via Harmonized Landsat Sentinel-2 (HLS) satellite time series, *ISPRS Journal of Photogrammetry and Remote Sensing*, 228, 28-43, <https://doi.org/10.1016/j.isprsjprs.2025.07.003>, 2025.

Dong, J. and Xiao, X.: Evolution of regional to global paddy rice mapping methods: A review, *ISPRS Journal of Photogrammetry and Remote Sensing*, 119, 214-227, <https://doi.org/10.1016/j.isprsjprs.2016.05.010>, 2016.

Dong, J., Xiao, X., Menarguez, M. A., Zhang, G., Qin, Y., Thau, D., Biradar, C., and Moore, B.: Mapping paddy rice planting area in northeastern Asia with Landsat 8 images, phenology-based algorithm and Google Earth Engine, *Remote Sensing of Environment*, 185, 142-154, <https://doi.org/10.1016/j.rse.2016.02.016>, 2016.

Dong, J., Xiao, X., Kou, W., Qin, Y., Zhang, G., Li, L., Jin, C., Zhou, Y., Wang, J., Biradar, C., Liu, J., and Moore, B.: Tracking the dynamics of paddy rice planting area in 1986–2010 through time series Landsat images and phenology-based algorithms, *Remote Sensing of Environment*, 160, 99-113, <https://doi.org/10.1016/j.rse.2015.01.004>, 2015.

Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P. K.: Global Consequences of Land Use, *Science*, 309, 570-574, 10.1126/science.1111772, 2005.

Gao, Y., Pan, Y., Zhu, X., Li, L., Ren, S., Zhao, C., and Zheng, X.: FARM: A fully automated rice mapping framework combining Sentinel-1 SAR and Sentinel-2 multi-temporal imagery, *Computers and Electronics in Agriculture*, 213, 108262, <https://doi.org/10.1016/j.compag.2023.108262>, 2023.

Ge, J., Zhang, H., Zuo, L., Xu, L., Jiang, J., Song, M., Ding, Y., Xie, Y., Wu, F., Wang, C., and Huang, W.: Large-scale rice mapping under spatiotemporal heterogeneity using multi-temporal SAR images and explainable deep learning, *ISPRS Journal of Photogrammetry and Remote Sensing*, 220, 395-412, <https://doi.org/10.1016/j.isprsjprs.2024.12.021>, 2025.

Ginting, F. I., Rudiyanto, R., Fatchurrachman, Mohd Shah, R., Che Soh, N., Eng Giap, S. G., Fiantis, D., Setiawan, B. I., Schiller, S., Davitt, A., and Minasny, B.: High-resolution maps of rice cropping intensity across Southeast Asia, *Scientific Data*, 12, 1408, 10.1038/s41597-025-05722-1, 2025.

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S. M., and Toulmin, C.: Food Security: The Challenge of Feeding 9 Billion People, 327, 812-818, doi:10.1126/science.1185383, 2010.

Han, J., Zhang, Z., Luo, Y., Cao, J., Zhang, L., Cheng, F., Zhuang, H., Zhang, J., and Tao, F.: NESEA-Rice10: high-resolution annual paddy rice maps for Northeast and Southeast Asia from 2017 to 2019, *Earth Syst. Sci. Data*, 13, 5969-5986, 10.5194/essd-13-5969-2021, 2021.

Han, J., Zhang, Z., Luo, Y., Cao, J., Zhang, L., Zhuang, H., Cheng, F., Zhang, J., and Tao, F.: Annual paddy rice planting area and cropping intensity datasets and their dynamics in the Asian monsoon region from 2000 to 2020,

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

513 Agricultural Systems, 200, 103437, <https://doi.org/10.1016/j.agsy.2022.103437>, 2022.

514 He, Y., Dong, J., Liao, X., Sun, L., Wang, Z., You, N., Li, Z., and Fu, P.: Examining rice distribution and cropping
515 intensity in a mixed single- and double-cropping region in South China using all available Sentinel 1/2 images,
516 International Journal of Applied Earth Observation and Geoinformation, 101, 102351,
517 <https://doi.org/10.1016/j.jag.2021.102351>, 2021.

518 Huang, H., Wang, J., Liu, C., Liang, L., Li, C., and Gong, P.: The migration of training samples towards dynamic
519 global land cover mapping, ISPRS Journal of Photogrammetry and Remote Sensing, 161, 27-36,
520 <https://doi.org/10.1016/j.isprsjprs.2020.01.010>, 2020.

521 Karra, K., Kontgis, C., Statman-Weil, Z., Mazzariello, J. C., Mathis, M., and Brumby, S. P.: Global land use / land
522 cover with Sentinel 2 and deep learning, 2021 IEEE International Geoscience and Remote Sensing Symposium
523 IGARSS, 11-16 July 2021, 4704-4707, 10.1109/IGARSS47720.2021.9553499,

524 Li, S., Shen, R., Jiang, J., Peng, Q., Chen, X., Dong, J., Dong, J., and Yuan, W.: A long-term paddy rice distribution
525 dataset in Asia at a 30 m spatial resolution, Scientific Data, 12, 1052, 10.1038/s41597-025-05374-1, 2025.

526 Lin, Z., Zhong, R., Xiong, X., Guo, C., Xu, J., Zhu, Y., Xu, J., Ying, Y., Ting, K. C., Huang, J., and Lin, T.: Large-
527 Scale Rice Mapping Using Multi-Task Spatiotemporal Deep Learning and Sentinel-1 SAR Time Series, 14, 699,
528 2022.

529 Mehta, P., Siebert, S., Kumm, M., Deng, Q., Ali, T., Marston, L., Xie, W., and Davis, K. F.: Half of twenty-first
530 century global irrigation expansion has been in water-stressed regions, Nature Water, 2, 254-261, 10.1038/s44221-
531 024-00206-9, 2024.

532 Ni, R., Tian, J., Li, X., Yin, D., Li, J., Gong, H., Zhang, J., Zhu, L., and Wu, D.: An enhanced pixel-based
533 phenological feature for accurate paddy rice mapping with Sentinel-2 imagery in Google Earth Engine, ISPRS
534 Journal of Photogrammetry and Remote Sensing, 178, 282-296, <https://doi.org/10.1016/j.isprsjprs.2021.06.018>,
535 2021.

536 Persaud, N. and Dagher, R.: The United Nations: 2030 Sustainable Development Goals Agenda, in: The Role of
537 Monitoring and Evaluation in the UN 2030 SDGs Agenda, edited by: Persaud, N., and Dagher, R., Springer
538 International Publishing, Cham, 1-41, 10.1007/978-3-030-70213-7_1, 2021.

539 Potapov, P., Turubanova, S., Hansen, M. C., Tyukavina, A., Zalles, V., Khan, A., Song, X.-P., Pickens, A., Shen,
540 Q., and Cortez, J.: Global maps of cropland extent and change show accelerated cropland expansion in the twenty-
541 first century, Nature Food, 3, 19-28, 10.1038/s43016-021-00429-z, 2022.

542 Song, M., Xu, L., Ge, J., Zhang, H., Zuo, L., Jiang, J., Ding, Y., Xie, Y., and Wu, F.: EARice10: a 10 m resolution
543 annual rice distribution map of East Asia for 2023, Earth Syst. Sci. Data, 17, 661-683, 10.5194/essd-17-661-2025,
544 2025.

545 Sun, C., Zhang, H., Xu, L., Ge, J., Jiang, J., Zuo, L., and Wang, C.: Twenty-meter annual paddy rice area map for
546 mainland Southeast Asia using Sentinel-1 synthetic-aperture-radar data, Earth Syst. Sci. Data, 15, 1501-1520,
547 10.5194/essd-15-1501-2023, 2023.

548 Thorp, K. R. and Drajat, D.: Deep machine learning with Sentinel satellite data to map paddy rice production
549 stages across West Java, Indonesia, Remote Sensing of Environment, 265, 112679,
550 <https://doi.org/10.1016/j.rse.2021.112679>, 2021.

551 Wang, Y., Hollingsworth, P. M., Zhai, D., West, C. D., Green, J. M. H., Chen, H., Hurni, K., Su, Y., Warren-Thomas,
552 E., Xu, J., and Ahrends, A.: High-resolution maps show that rubber causes substantial deforestation, Nature, 623,
553 340-346, 10.1038/s41586-023-06642-z, 2023.

554 Xiao, X., Boles, S., Froking, S., Li, C., Babu, J. Y., Salas, W., and Moore, B.: Mapping paddy rice agriculture in
555 South and Southeast Asia using multi-temporal MODIS images, Remote Sensing of Environment, 100, 95-113,
556 <https://doi.org/10.1016/j.rse.2005.10.004>, 2006.

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman
设置了格式: 字体: (默认) Times New Roman

557 Xiao, X., Boles, S., Liu, J., Zhuang, D., Frolking, S., Li, C., Salas, W., and Moore, B.: Mapping paddy rice
558 agriculture in southern China using multi-temporal MODIS images, *Remote Sensing of Environment*, 95, 480-
559 492, <https://doi.org/10.1016/j.rse.2004.12.009>, 2005.

560 Xu, S., Zhu, X., Chen, J., Zhu, X., Duan, M., Qiu, B., Wan, L., Tan, X., Xu, Y. N., and Cao, R.: A robust index to
561 extract paddy fields in cloudy regions from SAR time series, *Remote Sensing of Environment*, 285, 113374,
562 <https://doi.org/10.1016/j.rse.2022.113374>, 2023.

563 Yang, J., Dong, J., Liu, L., Zhao, M., Zhang, X., Li, X., Dai, J., Wang, H., Wu, C., You, N., Fang, S., Pang, Y., He,
564 Y., Zhao, G., Xiao, X., and Ge, Q.: A robust and unified land surface phenology algorithm for diverse biomes and
565 growth cycles in China by using harmonized Landsat and Sentinel-2 imagery, *ISPRS Journal of Photogrammetry*
566 *and Remote Sensing*, 202, 610-636, <https://doi.org/10.1016/j.isprsjprs.2023.07.017>, 2023.

567 Zabel, F., Delzeit, R., Schneider, J. M., Seppelt, R., Mauser, W., and Václavík, T.: Global impacts of future cropland
568 expansion and intensification on agricultural markets and biodiversity, *Nature Communications*, 10, 2844,
569 10.1038/s41467-019-10775-z, 2019.

570 Zeng, Z., Estes, L., Ziegler, A. D., Chen, A., Searchinger, T., Hua, F., Guan, K., Jintrawet, A., and F. Wood, E.:
571 Highland cropland expansion and forest loss in Southeast Asia in the twenty-first century, *Nature Geoscience*, 11,
572 556-562, 10.1038/s41561-018-0166-9, 2018.

573 Zhan, P., Zhu, W., and Li, N.: An automated rice mapping method based on flooding signals in synthetic aperture
574 radar time series, *Remote Sensing of Environment*, 252, 112112, <https://doi.org/10.1016/j.rse.2020.112112>, 2021.

575 Zhang, G., Xiao, X., Dong, J., Xin, F., Zhang, Y., Qin, Y., Doughty, R. B., and Moore, B.: Fingerprint of rice
576 paddies in spatial – temporal dynamics of atmospheric methane concentration in monsoon Asia, *Nature*
577 *Communications*, 11, 554, 10.1038/s41467-019-14155-5, 2020.

578 Zhang, G., Xiao, X., Biradar, C. M., Dong, J., Qin, Y., Menarguez, M. A., Zhou, Y., Zhang, Y., Jin, C., Wang, J.,
579 Doughty, R. B., Ding, M., and Moore, B.: Spatiotemporal patterns of paddy rice croplands in China and India from
580 2000 to 2015, *Science of The Total Environment*, 579, 82-92, <https://doi.org/10.1016/j.scitotenv.2016.10.223>,
581 2017.

582 Zhang, X., Liu, L., Chen, X., Gao, Y., Xie, S., and Mi, J.: GLC_FCS30: global land-cover product with fine
583 classification system at 30 m using time-series Landsat imagery, *Earth Syst. Sci. Data*, 13, 2753-2776,
584 10.5194/essd-13-2753-2021, 2021.

585 Zhao, Z., Dong, J., Yang, J., Liu, L., You, N., Xiao, X., and Zhang, G.: From rice planting area mapping to rice
586 agricultural system mapping: A holistic remote sensing framework for understanding China's complex rice systems,
587 *ISPRS Journal of Photogrammetry and Remote Sensing*, 224, 382-397,
588 <https://doi.org/10.1016/j.isprsjprs.2025.03.026>, 2025.

589 Zhao, Z., Dong, J., Zhang, G., Yang, J., Liu, R., Wu, B., and Xiao, X.: Improved phenology-based rice mapping
590 algorithm by integrating optical and radar data, *Remote Sensing of Environment*, 315, 114460,
591 <https://doi.org/10.1016/j.rse.2024.114460>, 2024.

592 Zhao, Z., Dong, J., Yang, J., Liu, L., You, N., Xiao, X., and Zhang, G.: From rice planting area mapping to rice
593 agricultural system mapping: A holistic remote sensing framework for understanding China's complex rice systems,
594 *ISPRS Journal of Photogrammetry and Remote Sensing*, 224, 382–397,
595 <https://doi.org/10.1016/j.isprsjprs.2025.03.026>, 2025a.

596 Zhao, Z., Dong, J., Zhang, G., Yang, J., and Xiao, X.: Mapping Paddy Rice Distribution and Cropping Intensity in
597 South and Southeast Asia (1995 - 2024) at 30m Resolution, <https://doi.org/10.5281/ZENODO.17615341>, 2025b.

598

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman