

The WoKaS-Iso Database: Workflow for a Global Compilation of Oxygen-18 and Deuterium Records in Karst Springs and Cave Drip Water for Enhanced Understanding of Karst Systems

Yining Zang^{1*}, Pauline C. Treble^{2,3*}, Kei Yoshimura^{4,5}, Jayson Gabriel Pinza^{6,7}, Fengbo Zhang⁸, Kübra Özdemir Çallı¹, Xiaojun Mei⁹, Admin Husic¹⁰, Alena Gessert¹¹, Andrej Stroj¹², Bartolomé Andreo¹³, Bernard Ladouche^{14,15}, Christine Stumpp¹⁶, Diana Mance¹⁷, Eleni Zagana¹⁸, Fen Huang¹⁹, Giuseppe Sappa²⁰, Harald Kunstmann^{21,22}, Heike Brielmann²³, Hong Zhou²⁴, Huaying Wu¹⁹, Jakob Garvelmann^{21,25}, James Berglund²⁶, Jean-Baptiste Charlier^{14,15}, Jens Lange²⁷, Juan Antonio Barberá Fornell¹³, Junbing Pu²⁸, Konstantina Katsanou²⁹, Kun Ren¹⁹, Laura Toran³⁰, Laurence Gill³¹, Maria Filippini³², Martin Kralik³³, Matías Mudarra Martínez¹³, Min Zhao³⁴, Mingming Luo²⁴, Nico Goldscheider³⁵, Nikolaos Lambrakis¹⁸, Pantaleone De Vita³⁶, Qiong Xiao¹⁹, Shi Yu¹⁹, Silvia Iacurto²⁰, Silvio Coda³⁶, Ted McCormack³⁷, Vincenzo Allocca³⁶, W. George Darling³⁸, Walter D'Alessandro³⁹, Xulei Guo²⁴, Yundi Hu⁴⁰, Zhijun Wang⁴¹, Eva Kaminsky^{42,43}, Jiří Faimon^{44,45}, Marek Lang⁴⁴, Pavel Pracný⁴⁴, Aurel Persoiu^{46,47,48}, Andreas Hartmann¹

* Corresponding authors

¹Institute of Groundwater Management, Technical University of Dresden, Dresden, 01069, Germany

²ANSTO, New Illawarra Road, Lucas Heights, NSW 2234, Australia

³School of Biological, Earth and Environmental Sciences, UNSW Sydney, Sydney NSW, 2052, Australia

⁴Institute of Industrial Science, The University of Tokyo, Kashiwa, Japan

⁵Japan Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Japan

⁶ECOSPHERE Research Group, University of Antwerp, Universiteitsplein 1C, 2610 Antwerp, Belgium

⁷Division of Soil and Water Management, KU Leuven, Celestijnenlaan 200E, 3001 Leuven, Belgium

⁸Ningbo River Management Center, Ningbo Water Resources Bureau, Ningbo 315000, China

⁹Merchant Marine College, Shanghai Maritime University, Shanghai 201306, China

¹⁰Department of Civil and Environmental Engineering, Virginia Tech, Blacksburg 24061, USA

¹¹Institute of Geography, Faculty of Sciences, University of P.J. Šafárik, 040 01 Košice, Slovakia

¹²Croatian Geological Survey, Sachsova 2, 10000 Zagreb, Croatia

¹³Department of Geology and Centre of Hydrogeology at the University of Malaga, 29071, Malaga, Spain

¹⁴BRGM, Univ Montpellier, Montpellier, France

¹⁵G-EAU, Univ Montpellier, AgroParisTech, BRGM, CIRAD, INRAE, Institut Agro, IRD, Montpellier, France

¹⁶Institute of Soil Physics and Rural Water Management, Department of Landscape, Water and Infrastructure, BOKU University, Vienna, Austria

¹⁷Faculty of Physics, University of Rijeka, 51000, Rijeka, Croatia

¹⁸Laboratory of Hydrogeology, Department of Geology, Faculty of Natural Sciences, University of Patras, GR-26504 Patras, Greece

¹⁹Key Laboratory of Karst Dynamics, MNR and GZAR, Institute of Karst Geology, Chinese Academy of Geological Sciences, Guilin, 541004, China

²⁰Department of Civil, Constructional and Environmental Engineering (DICEA), Sapienza University of Rome, 00184 Rome, Italy

²¹Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Campus Alpin, D-82467

Garmisch-Partenkirchen, Germany

²²Institute of Geography, University of Augsburg, D-86135 Augsburg, Germany

²³Environment Agency Austria, Groundwater Unit, Spittelauer Laende 5, A-1090 Vienna, Austria

- ²⁴School of Environmental Studies& Institute of Geological Survey, China University of Geosciences, Wuhan, 430074, Hubei, China
- ²⁵Boden & grundwasser Allgäu GmbH, D-87527 Sonthofen, Germany
- ²⁶Department of Civil and Environmental Engineering, University of Wisconsin – Platteville, Platteville, Wisconsin, USA
- ²⁷Hydrology, University of Freiburg, 79098 Freiburg, Germany
- ²⁸Chongqing Key Laboratory of Carbon Cycle and Carbon Regulation of Mountain Ecosystem, School of Geography and Tourism, Chongqing Normal University, Chongqing, 401331, China
- ²⁹Department of Water Resources and Ecosystems, IHE Delft Institute for Water Education, Delft, Netherlands
- ³⁰Department of Earth and Environmental Science, Temple University, Philadelphia, Pennsylvania, USA
- ³¹Department of Civil, Structural and Environmental Engineering, Trinity College Dublin, Ireland
- ³²Department of Biological, Geological and Environmental Sciences—BiGeA, Alma Mater Studiorum, University of Bologna, Via Selmi 3, 40126 Bologna, Italy
- ³³Department of Environmental Geosciences, Center for Microbiology and Environmental Systems Science, University of Vienna, 1090 Vienna, Austria
- ³⁴State Key Laboratory of Environmental Geochemistry (SKLEG), Institute of Geochemistry, Chinese Academy of Sciences (CAS), Guiyang, China
- ³⁵Institute of Applied Geosciences, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany
- ³⁶Department of Earth, Environmental and Resources Sciences (DiSTAR), University of Naples Federico II, Naples, Italy
- ³⁷Geological Survey Ireland, Booterstown Avenue, Booterstown, Blackrock, Ireland
- ³⁸British Geological Survey, Wallingford OX10 8BB, United Kingdom
- ³⁹Istituto Nazionale di Geofisica e Vulcanologia – Sezione di Palermo, Via La Malfa 153, 90146 Palermo, Italy
- ⁴⁰School of Geography and Environmental Science, Guizhou Normal University, Guiyang 550001, People's Republic of China
- ⁴¹Faculty of Arts and Sciences, Beijing Normal University, Zhuhai 519087, China
- ⁴²Karst and Cave Group, Geological Paleontological Department, Natural History Museum, Vienna, Austria
- ⁴³Institute of Soil Physics and Rural Water Management, Department of Landscape, Water and Infrastructure, BOKU University, Vienna, Austria
- ⁴⁴Department of Geological Sciences, Faculty of Science, Masaryk University, Brno, Czech Republic
- ⁴⁵Czech Geological Survey, Brno Branch, Brno, Czech Republic
- ⁴⁶Emil Racoviță Institute of Speleology, Romanian Academy, Cluj-Napoca, 400006, Romania
- ⁴⁷Stable Isotope Laboratory, Stefan cel Mare University, Suceava, 720229, Romania
- ⁴⁸School of Geology, Department of Physical Geography, Aristotle University of Thessaloniki, Thessaloniki, 54636, Greece

Correspondence to: Yining Zang (yining.zang@tu-dresden.de), Pauline C.Treble (ptr@ansto.gov.au)

Abstract

For analysing karst hydrogeological systems, observations of karst springs and cave drips are considered indispensable. In addition to hydrometric observations, knowing the oxygen and hydrogen stable isotope ratios has improved the understanding of vadose zone and aquifer dynamics, likewise supporting system characterisation and modelling. However, limited accessibility and high costs of the analysis of stable isotopes in karst aquifers have hindered progress in karst research and impeded the accurate understanding of karst processes especially when it comes to comparative or large-scale studies. In this study, we present our workflow to compile the WoKaS-Iso database, the first extensive collection of time series data for Oxygen-18 and Deuterium isotopes in karst springs and cave drip water from diverse sources, encompassing publications, theses, reports, online archives, and collaborative initiatives worldwide. The database incorporates data sourced from 241 springs and 74 caves, comprising in total 1007 time series including 389 spring time series and 618 cave drip water time series.

85 Of these sites, 148 springs and 64 caves contain both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ records, while 93 springs and 9 caves contain only one
86 isotope tracer. These datasets provide coverage across significant karst regions globally, spanning China, the USA, Europe,
87 the Middle East, and Australia. Within datasets, 79% for springs and 68% for cave drip water exhibit resolutions finer than
88 monthly intervals. In addition, by integrating isotopic records with ancillary environmental variables including spring
89 discharge, cave drip rate, precipitation, and rainwater isotopes, the database offers a more comprehensive perspective on
90 hydrological behaviours in karst aquifers, hence advancing hydrogeological characterisation and modelling. The WoKaS-Iso
91 database not only deepens the understanding of the complex systems but also promotes sustainable water resource management
92 as well as the potential to foster collaborative research. The database can be accessed at: [https://doi.org/10.25532/OPARA-](https://doi.org/10.25532/OPARA-909)
93 [909](https://doi.org/10.25532/OPARA-909) (Zang, 2025).

94 1 Introduction

95 Carbonate rock areas, particularly those with highly karstified systems, are home to 16.5% of the global population
96 (Goldscheider et al., 2020) and serve as a critical source of potable water for an estimated 9.2% of the global population
97 (Stevanović, 2019). Beyond water supply, these aquifers encompass extensive cave networks that preserve valuable
98 paleoclimate records, reflecting key variability patterns of past environmental conditions (McDermott, 2004). The
99 karstification leads to inherent structural heterogeneity and strong nonlinearities of hydraulic properties (Frank et al., 2021;
100 Jourde et al., 2018; Labat et al., 2016) that drive subsurface water flow regimes and storage mechanisms including fast conduit
101 turbulent flow and slow matrix storage (Bakalowicz, 2005; Goldscheider and Drew, 2007) while challenging characterization
102 of karst aquifer structures, hydraulic dynamics, and contamination pathways. Consequently, a comprehensive understanding
103 of hydrogeological behaviours in karst systems is essential for sustainable water resource management and protection. This is
104 also a matter of urgency for certain European regions that appear to be sensitive to the effects of global warming (Giese et al.,
105 2025).

106
107 Stable isotopes of oxygen and hydrogen, being part of the water molecule itself, are ideal tracers for identifying water sources
108 (Hartmann et al., 2014) delineating subsurface flow paths, estimating transit times (Liu and Yamanaka, 2012; McDonnell et
109 al., 2010), and quantifying mixing processes (Rusjan et al., 2019). In karst hydrogeology, spring isotope signatures integrating
110 responses from both saturated and unsaturated zones often give additional information about complex recharge patterns that
111 are not captured by discharge data alone. Isotope data thus provide critical complementary information to resolve flow and
112 transport dynamics. Meanwhile, in speleological research, stable isotopes are central to understanding recharge processes
113 through the epikarst, providing a distinctive window into unsaturated zone and recharge processes (Hartmann and Baker, 2017).
114 Cave drips act as a medium to monitor flow path through epikarst that is directly linked to climate signals preserved in
115 speleothem formed from cave drips which can serve as archives for paleoclimate reconstruction (Bradley et al., 2010;
116 (MacDermott et al., 2006). Long-term rainfall and drip water monitoring is crucial for distinguishing local or regional climatic

influences on cave system isotope variability (Pape et al., 2010). Moreover, stable isotopes provide critical information on water transit times through karst systems, allowing the identification of fast conduit flow versus slow matrix storage contributions. This capability is essential for understanding the temporal dynamics of recharge, flow pathways, and aquifer vulnerability to contamination, as it directly links precipitation events to spring or cave drip responses. Incorporating isotope-based transit time estimates into hydrological models enhances the characterization of flow heterogeneity and improves predictions of contaminant transport and residence times within karst aquifers.

More recently, isotope-based models have emerged to couple hydrological dynamics with stable isotopes (Birkel and Soulsby, 2015; Zhang et al., 2019). Andreo et al. (2004) applied $\delta^{18}\text{O}$ and $\delta^2\text{H}$ in precipitation and groundwater to determine the water origin and understand the aquifer dynamics in the Yunquera-Nieves karstic massif in Spain. Wang et al. (2022) coupled hydrochemical and stable isotopic compositions to identify the groundwater hydrochemistry evolution in the Western Yellow Sea Coast of China. Hartmann et al. (2012) utilized isotopic information ($\delta^{18}\text{O}$) to calibrate their recharge model. Hydrological functions such as water storage, flux and water age distributions are effectively quantified by these isotope-coupled models, providing essential indicators of understanding the behaviour of the karst aquifer (Mayer-Anhalt et al., 2022). Additionally, coupling isotopic and hydrological data has proven valuable for facilitating the multi-objective calibration and validation (Hartmann et al., 2013) enhancing the model evaluation, mitigating overparameterization and reducing the potential uncertainties (Li et al., 2022; Zhang et al., 2019). On the other hand, recent cave-based research has used stable water isotopes to advance the understanding of karst vadose zone hydrology. Priestley et al. (2023) combined $\delta^{18}\text{O}$ from speleothems and drip waters with modelled soil moisture data to show how regional drying in southwestern Australia disrupted rainfall recharge to shallow karst aquifers—an unprecedented shift over the past 800 years. Baker et al. (2019) analysed $\delta^{18}\text{O}$ from 163 global drip sites and found that in seasonal climates, drip water isotopes reflect recharge-weighted precipitation, refining the interpretation of speleothem records. Treble et al. (2022) demonstrated that isotopic variability within caves is largely driven by differences in flow path properties—fracture versus matrix flow—highlighting the importance of hydrological heterogeneity in shaping $\delta^{18}\text{O}$ signals. Together, these studies underline the growing potential of isotope-hydrology approaches for resolving spatial and temporal variability in karst recharge and for reconstructing past hydroclimatic conditions. All these applications highlight the growing value of isotopic datasets for both process understanding and model development in karst systems.

Several global databases have been established to archive stable water isotope data, most notably the Global Network of Isotopes in Precipitation (GNIP) and the Global Network of Isotopes in Rivers (GNIR) maintained by the International Atomic Energy Agency (IAEA) and World Meteorological Organization (WMO). In addition, Staudinger et al. (2020) developed a long-term $\delta^{18}\text{O}$ and $\delta^2\text{H}$ database for streamflow and precipitation for 23 catchments in Switzerland, and Li et al. (2025) established the first global surface water stable isotope database from measured data, web-sourced records and referenced data. While these databases provide essential resources for hydrological studies, they were not designed to systematically represent

karst environments, and stable isotope observations from karst springs and cave drip waters remain scattered across individual studies. $\delta^{18}\text{O}$ and $\delta^2\text{H}$ signatures in karst springs and cave drip waters as natural tracers have been proven invaluable, offering insights into characterizing recharge patterns, validating and enhancing the accuracy of hydrological models and calibrating the reconstruction of paleoclimate through speleothem records (Tremaine et al., 2011). Despite the significance, the broader use of isotopic applications remained constrained by the high cost of stable isotope analysis and limited accessibility to the existing datasets for large-scale modelling and comparative studies across karst regions.

To allow more a broader application and comparative studies, we developed the WoKaS-Iso - a global compilation of the time series of stable isotope ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) records from karst springs and cave drip water. As complementary information, this database also includes the karst discharge, drip rate, precipitation, and precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ observations. For springs, datasets were aggregated from collaborators, reviewed publications, technical reports and theses. For caves, data were sourced from the SISAL_mon_v1 cave drip monitoring database (Treble et al., to be submitted), under consistent formatting and quality control. Metadata considered karst springs and cave name, geological information of sites, monitoring starting and ending date, observation length and data source details. We present the methodology of WoKaS-Iso database construction, describe datasets records, datasets quality control, data usage notes, and discuss its applications and outlook. Furthermore, we evaluate the performance of three global precipitation isotope models, offering the perspectives of model selection in karst hydrology.

2 Data records and methodology

The WoKaS-Iso database is built upon the foundational WoKaS (World Karst Spring hydrograph) database, the first global repository of karst spring discharge records encompassing over 400 karst springs (Olarinoye et al., 2020), which however does not contain stable isotope records. WoKaS-Iso extends this framework by comprising stable isotope data ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) for 241 karst springs worldwide. Initially developed from a compilation of 119 stable isotope time series from Europe (Hartmann et al., 2021), the database has since expanded in scope to global coverage. Among these, 29 springs are also included in WoKaS, enabling their isotope records to be directly linked with the corresponding discharge data. For the remaining springs, available isotopic, hydrological and climatological data were compiled from peer-reviewed literature, technical reports, open-access data repositories, and direct contributions from research collaborators.

The cave component was developed in collaboration with the SISAL (Speleothem Isotope Synthesis and Analysis) working group (<https://pastglobalchanges.org/science/wg/sisal/intro>). Of the 74 caves incorporated in the database, 66 were derived from SISAL_mon_v1 (Treble et al., 2026), while the remaining 8 were compiled from research collaborators contributions. WoKaS-Iso extends SISAL_mon_v1 by integrating cave drip water stable isotope compositions and drip rate measurements with spring isotope records within a unified database framework. The dataset is further augmented by supplementary datasets including drip rate measurements and GNIP station records, that were recoverable from the original data sources but were not

incorporated in SISAL_mon_v1. All cave records were systematically standardised in accordance with the WoKaS-Iso data schema to ensure interoperability and to facilitate cross-referencing across database components. Detailed descriptions of the data recovery, standardisation procedures, and quality control measures applied to the cave monitoring records are provided in Sect. 2.1.2 ("Cave Monitoring") and Chapter 4.

WoKaS-Iso also synthesizes paired hydrological and climatic parameters datasets including karst spring discharge, rainfall, and rainwater isotopes to support interpretation of the water isotope signature. These datasets were gathered from peer-reviewed publications, theses, reports and research collaborators. In addition to site-based observations, we integrated modelled climatic datasets from gridded global products, including precipitation amount (MSWEP), air temperature (ERA5), evapotranspiration (GLEAM), and precipitation isotopes models (IsoGSM, Isoscape, Sine Curve). Where observed data were unavailable, the precipitation amount and precipitation isotopes from nearby GNIP stations were used as substitutes.

From a modelling perspective, the datasets in WoKaS-Iso are organized in two main categories: **system output data** (spring isotopes, spring discharge, cave drip isotopes, cave drip rate) and **system input data** (precipitation amounts, precipitation isotopes). The overview workflow of WoKaS-Iso database creation is illustrated in Fig.1.




Figure 1: Workflow of the karst spring and cave drip water stable isotopes database construction. The left branch illustrates the karst spring component, built upon the WoKaS global spring discharge database and an initial European stable isotope compilation (Hartmann et al., 2021), with data sourced from publications, reports, theses, and research collaborators via collection and digitization. The right branch shows the cave monitoring component, developed in collaboration with the SISAL working group and drawing primarily on SISAL_mon_v1 (Treble et al., 2026), supplemented by additional published and contributed cave datasets. The central panel depicts the system input data,

comprising observed and modelled precipitation amounts (MSWEP), precipitation isotopes (GNIP, IsoGSM, Isoscape, Sine Curve), evapotranspiration (GLEAM), and air temperature (ERA5). The main workflow is described in the preceding paragraphs, with further methodological details on output data compilation and input data integration provided in Section 2.1.2 and 2.2.1.

2.1 System output data

2.1.1 Available data types

The WoKaS-Iso database compiles observations of oxygen ($\delta^{18}\text{O}$) and hydrogen ($\delta^2\text{H}$), expressed in delta (δ) notation relative to the Vienna Standard Mean Ocean Water (VSMOW), from 241 karst springs and 74 caves accommodating in total 365 drip observation points (e.g., different speleothems). In addition, WoKaS-Iso includes associated karst spring discharge, cave drip rate, precipitation amounts observations and precipitation isotope observations for karst springs and cave sites. Table 1 details the number of time series included in the database for various variables.


Table 1: Summary of WoKaS-Iso database observations

Variables	Karst Springs	Cave Drips	Total
$\delta^{18}\text{O}$ Time Series	237	339	576
$\delta^2\text{H}$ Time Series	152	279	431
Flow Rate Time Series	122	186	308
Precipitation Amounts	119	58	177
Precipitation Isotopes	127	66	193

Among 241 karst springs, 122 (51%) are accompanied by discharge data. Of these, 95 springs include both discharge and output $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data, while 27 springs possess discharge data paired with either $\delta^{18}\text{O}$ or $\delta^2\text{H}$ measurements. The remaining 119 springs (49%) include only output isotope data: 53 springs with both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data, and 66 with either $\delta^{18}\text{O}$ or $\delta^2\text{H}$ data. Within 74 caves, one cave includes input data alone but lacks output data. 73 caves comprise 365 drip sites, among these, 161 drip sites (44%) combine drip rate with drip water isotope data, of 161 drip sites, 154 drip sites have drip rate along with $\delta^{18}\text{O}$ and $\delta^2\text{H}$ measurements, while 7 drip sites have drip rate and only $\delta^{18}\text{O}$ data. Additionally, 25 drip sites (7%) were measured containing drip rate only. Furthermore, 178 drip sites (49%) contain drip water isotope data alone, 125 drip sites with both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data, 53 drip sites with $\delta^{18}\text{O}$ only. In addition, the database includes 127 time series of rainwater isotopes, and 119 time series of rainfall associated with karst springs, as well as 66 time series of rainwater isotopes and 58 time series of rainfall associated with cave sites. The complete list of karst springs and caves included in the database, along with their coordinates and referenced literatures, are presented in Table S1 and Table S2 in the supplement.

229 **2.1.2 Origin of the collected data**

230 Fig.2 illustrates the sources of the karst spring and cave drip data, respectively. For the karst springs, 90% isotopes data are
 231 raw observations while 10% are digitized (extracted from a reference paper). The corresponding karst spring discharge data
 232 consists of 86% raw data from collaborators, WoKaS discharge database or online open-access national database. For the
 233 rainwater isotopes, 8% of the data are from local measurements, 45% from weather stations, and 47% from global isotope
 234 model products. Besides, 50% rainfall data are sourced from local and meteorological data. In terms of cave drip waters, 96%
 235 of drip water isotopes derive from direct field measurements, with only 4% digitized. A parallel distribution is observed for
 236 drip rate records, where 72% source from observations. In case of rainwater, the majority of rainwater isotopes are obtained
 237 from both local sampling campaigns and meteorological stations, with approximately 45% originating from local observations.
 238 Likewise, 78% of rainfall data are acquired from local records and meteorological station datasets.



239 **Figure 2:** Distribution of time series counts for hydrological and isotopic datasets in WoKaS-Iso database, categorized by source types. For
 240 karst springs, precipitation isotopes come from local observations (18), meteorological stations (109), and global model products (114),
 241 while precipitation amounts come from local observations (20), meteorological stations (99), and MSWEP (122). Spring water isotopes
 242 include collected (217) and digitized (24) data, and spring discharge includes collected (105) and digitized (17) data. For caves, precipitation
 243 isotopes come from local observations (33), meteorological stations (33), and global model products (8). Cave drip water isotopes include
 244 collected (324) and digitized (15) data, and cave drip rates include collected (133) and digitized (53) data.

247 **Karst Springs:** As summarized in the left branch of Fig. 1, the time series of $\delta^{18}\text{O}$ and $\delta^2\text{H}$ output data were collected from
248 (1) published data in peer-reviewed literatures, scientific reports and academic theses; (2) research collaborators from the
249 network. **Published data:** The publications containing stable isotopes time series data for karst springs were identified through
250 systematic searches of academic databases and web search engines using targeted keywords (e.g. “karst spring isotopes”).
251 Some isotope datasets were obtained directly from the supplementary documents accompanying these publications. For the
252 data not readily accessible, requests were sent to the corresponding authors to solicit contributions to the database. In cases
253 where no response was received, the datasets were digitized from figures in scientific articles, reports and theses using
254 WebplotDigitizer (Rohatgi, 2025), a web-based open-source tool for data extraction. To maximum digitization accuracy, high-
255 quality figures in articles were retrieved programmatically. The extracted datasets were then visually validated by replotting
256 and comparing them with the original figures to ensure the trends’ consistency. **Research collaborators:** To further enrich the
257 WoKaS-Iso database, additional data were acquired through communications with research project partners and institutional
258 collaborations. Advertisements were disseminated to relevant academic departments, organizations and researchers.
259 Furthermore, calls for contribution were made to the karst community through poster and oral presentations during the
260 international conferences and workshops. In addition to the sources above, the corresponding karst discharge data were
261 obtained from the online national database such as the U.S. Geological Survey (USGS)’s National Water Information System,
262 and Austria’s Bundesministerium für Nachhaltigkeit und Tourismus (eHYD) database.

263 **Cave Monitoring:** As illustrated in the right branch of Fig. 1, The compilation of cave drip water isotopes and drip rate for
264 the WoKaS-Iso database followed a structured workflow analogous to that of karst springs. In collaboration with SISAL
265 (Kaushal et al., 2024), data acquisition primarily drew upon the SISAL_mon_v1 (Treble et al., 2026), which provided a key
266 foundation for this component. GNIP station data excluded from SISAL_mon_v1 (as noted in the SISAL workbooks) are
267 retained in WoKaS-Iso to preserve the completeness and representativeness of the original monitoring datasets. For some caves,
268 SISAL_mon_v1 includes only the drip water isotope data provided by the authors, but the original drip rate measurements
269 were not available. To address this, WoKaS-Iso digitized the drip rate information from figures in the publications and
270 incorporated these data into the database, thereby enabling quantitative analyses of cave hydrology, drip water dynamics, and
271 speleothem formation processes. Furthermore, additional cave monitoring sites, independent of SISAL_mon_v1, have been
272 incorporated from published studies and research collaborators. All cave drip water datasets were systematically standardised
273 and additional quality control according to the WoKaS-Iso protocol (see chapter 4) was applied. For datasets originating from
274 SISAL_mon_v1, this quality control focused on additional harmonisation and alignment with the WoKaS-Iso structure,
275 building on the quality control already performed in Treble et al. (2026). The datasets were then organised within a structure
276 specifically designed to ensure interoperability with the broader WoKaS-Iso. To ensure transparency and facilitate cross-
277 referencing, the corresponding SISAL_mon_v1 cave identifiers are provided in the WoKaS-Iso metadata tables where
278 applicable. Additional information on the cave drip water data collection methodology and site selection in SISAL_mon_v1
279 is provided in Treble et al. (2026).

2.1.3 Description of available datasets

The spatial distributions of springs and caves within the database are shown in Fig.3 together with the World Karst Aquifer Map (WOKAM) (Chen et al., 2017). The locations in the database are primarily situated across key global karst regions including China, Europe, Australia, the Middle East and the USA. We summarised the detailed statistics of data type combinations of 241 springs and 339 cave drip sites with isotopic data records in Fig.4, stratified by the predominant karst areas identified in Fig.3. Springs are analysed for China, the USA, Europe for springs, while cave drip sites are grouped by China, Australia, and Europe. The stacked histograms display the distribution of dataset types across these regions and align with the regional distribution trend described in the preceding section. Fig. 5 further illustrates regional differences in isotope dynamics by presenting example $\delta^{18}\text{O}$ time series from three karst springs and three cave drip water systems across North America, Europe, and Asia.

Most karst spring records are collected in Europe with approximately half of them accompanied by discharge measurements (Fig. 4a). Similar proportions are observed in China, whereas only one-third of spring records in the USA include discharge data. In contrast, more than 80 % of spring records from other regions are associated with discharge measurements. Regarding stable isotope observations, datasets containing both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ are most common in China (86 %), followed by the USA (70 %) and other regions (62 %). For cave drip water systems (Fig. 4b), drip rate observations are most frequently available in China and are present in roughly half of the records from Australia and Europe. In addition, more than 95 % of cave drip water records in China and Australia include both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ measurements, while the proportion exceeds 70 % in Europe and other regions.




Figure 3: Geographic distribution of karst springs and cave sites included in the WoKaS-Iso database, along with the types of data available at each site. Blue areas indicate karst regions from the World Karst Aquifer Map (WOKAM, Chen et al. 2017). “Q” in the legend refers to karst spring discharge. ‘Green round’ symbols represent karst springs; ‘red star’ symbols represent caves. The symbol color intensity reflects the richness of available data combinations at each site, categorized into four subgroups: **flow rate+ $\delta^{18}\text{O}$ + $\delta^2\text{H}$** , **flow rate+ $\delta^{18}\text{O}$** , **$\delta^{18}\text{O}$ + $\delta^2\text{H}$** , **$\delta^{18}\text{O}$** , with darker color indicating more comprehensive datasets including both hydrological and isotopic data. Panels (a), (b), and (c)

show zoomed-in maps of karst springs in China, the USA, and Europe, respectively. Panels (d), (e), and (f) show zoomed-in maps of cave sites in China, Australia, and Europe. In the zoomed maps, the size of each symbol presents the sampling resolution of $\delta^{18}\text{O}$ and $\delta^2\text{H}$ time series at each spring and cave drip site. The temporal resolution is categorized into three classes: **less than biweekly, monthly and coarser than monthly**, corresponding to large, medium and small markers respectively.




Figure 4: Summaries of data types across the major karst regions for (a) karst springs and (b) cave drip water sites. In panel (a), “Q” in the legend represents karst spring discharge.

Stable isotope dynamics generally differ among regions due to differences in climatic conditions, precipitation isotope inputs, and karst hydrological processes. Therefore, three karst springs and three caves with overlapping observation periods were selected from North America, Europe, and Asia as illustrative examples to show the temporal dynamics of $\delta^{18}\text{O}$. Fig.5a shows considerable differences in both isotope values and temporal variability of three springs. BS1 Spring (North America) exhibits relatively enriched $\delta^{18}\text{O}$ values (approximately -5 to -2 ‰), whereas Anyak Spring (Europe) remains comparatively stable around -10 ‰. In contrast, Luxi No.1 Spring (Asia) displays larger temporal fluctuations, with $\delta^{18}\text{O}$ values ranging from approximately -14 to -9 ‰. Similar regional differences are also evident in cave drip water records (Fig.5b). Larga Cave (North America) is characterised by relatively enriched isotope values around -3 ‰, while Beke Cave (Europe) and Yangkou Cave (Asia) exhibit more depleted values between approximately -10 and -7 ‰. Moreover, multiple drip sites within individual caves show both common temporal behaviour but differ in absolute $\delta^{18}\text{O}$ values.





Figure 5: Representative $\delta^{18}\text{O}$ time series from selected karst springs and caves across North America, Europe, and Asia. (a) Karst spring records from BS1 Spring, Anyak Spring, and Luxi No.1 Spring. (b) Cave drip water records from Larga Cave, Beke Cave, and Yangkou Cave. Multiple drip sites are shown for each cave to illustrate intra-cave variability.

Stable isotopes in spring discharge and drip waters can show strong seasonal variability due to fluctuations in precipitation, and temperature under changing climatic conditions. Adequate temporal resolutions are necessary to quantify these seasonal dynamics in detail and capture short-term hydrological patterns during rainfall or snowmelt events. The regional maps in Fig. 3a-f provide the spatial context of isotope sampling resolution, showing that monthly sampling is common across the main

333 data-rich regions, particularly for cave drip water records. Higher resolution isotope records are mainly available for karst
 334 springs in Europe, whereas finer than biweekly resolution is rare for cave drip water records in China, Australia, and
 335 Europe. Fig. 6 provides a summary of temporal resolution for hydrological and isotope measurements in WoKaS-Iso. Spring
 336 discharge datasets generally have the highest temporal resolution, with nearly half available at daily or finer intervals,
 337 compared with just over 12 % of cave drip rate datasets (Fig. 6a). Cave drip rate records are mainly concentrated in the
 338 biweekly-to-monthly category, reaching approximately 74 %, whereas spring discharge records are more evenly distributed
 339 across the resolution classes. For stable isotope observations, resolution between biweekly and monthly is the dominant
 340 category for both springs and cave drip waters, accounting for around 64 % of cave drip isotope datasets and more than 40 %
 341 of spring isotope datasets (Fig. 6b). In addition, 36 % of spring isotope datasets have the resolution finer than biweekly, while
 342 27 % of cave drip water isotope records are sampled at intervals coarser than monthly.



343 **Figure 6:** (a) Proportions of temporal resolution categories of discharge and drip rate datasets for karst springs and cave drip sites; The
 344 datasets are divided into four resolution levels: **finer than daily**, **daily to biweekly**, **biweekly to monthly** and **coarser than monthly**. (b)
 345 Proportions of temporal resolution categories of stable isotopes for karst springs and caves. The resolutions are stratified into four categories:
 346 **finer than biweekly**, **biweekly to monthly**, **coarser than monthly** and **single-value records**.
 347

2.2 System input data

2.2.1 Available data types

The workflow for compiling the input datasets is summarized in the central panel of Fig. 1. The WoKaS-Iso database includes the precipitation amount and precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ observations for karst springs and cave sites, compiled from the same sources described in subsection 2.1.2 (also see Table 1 & Figure 2). To ensure adequate spatial and temporal coverage, karst springs and cave sites lacking local rainfall and rainwater isotopes records were supplemented with the data from Global Network of Isotopes in Precipitation (GNIP) database. Firstly, the geographic coordinates of karst springs, cave sites, and GNIP stations within the related countries were imported into ArcGIS where the closest GNIP station to each karst spring and cave site was identified. Secondly, the distances between the springs or caves and their nearest GNIP stations were measured using ArcGIS. With the aim of reflecting representative of the local climatic and isotopic characteristics for karst springs and caves, only GNIP stations within 25 km were retained. This threshold was selected to be consistent with the spatial resolution of the Isoscape and Sine Curve datasets (5 arcmin, approximately 10 km), corresponding to roughly twice the grid-cell diagonal (~28 km). It is therefore a conservative choice that is also compatible with the coarser ERA5 dataset (~31 km). Although this approach cannot fully account for topographic differences between GNIP stations and karst sites, it provides a consistent spatial matching criterion for all products evaluated in this study.

For karst springs, 20 sites possess local rainfall measurements, and 99 springs derive precipitation data from nearby meteorological stations including GNIP stations, while the remaining 122 springs lack rainfall data. Regarding precipitation isotope observations, only 18 springs are with auxiliary local isotope data, 109 springs use the data from meteorological or GNIP network, and 114 springs have no related isotope records. For cave sites, 21 caves include locally measured rainfall, 37 caves obtained rainfall data from adjacent weather stations, and 16 caves without rainfall records. As for precipitation isotopes, 33 caves are associated with local measurements, 33 caves rely on weather stations-based isotope datasets or GNIP-derived data, and 8 caves lack isotope observations. To offer comprehensive metadata for input observations, the distance from the precipitation and precipitation isotopes sampling sites to cave/spring sites were calculated where geographic coordinates were available, documented in the metadata tables for spring input and cave input datasets.

Since datasets on precipitation amount and precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$, observed locally or provided by GNIP, were not available at all or output locations or limited in their temporal extent, the WoKaS-Iso database includes complementary datasets from six global products for continuous rainfall, precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$, air temperature as well as both actual and potential evapotranspiration. The information of these global products used in this study are listed in Table 2, including relevant modelled variables, time span, spatial and temporal resolution and the source for the construction of models.

Precipitation data were obtained from MSWEP v2.8 (Beck et al., 2017, 2019a, b), a multi-source ensemble product offering daily and monthly estimates at 0.1° resolution from 1979–2020. MSWEP improves global precipitation accuracy by combining

gauge, satellite, and reanalysis data using a weighted ensemble based on network density and local performance (Liu et al., 2019). Evaporation estimates were sourced from GLEAM v4.1a (Hulsman et al., 2023; Koppa et al., 2022; Martens et al., 2017; Miralles et al., 2011, 2025; Zhong et al., 2022), which integrates satellite data and physical modelling with machine learning to estimate daily and monthly actual and potential evaporation at 0.1° resolution from 1980–2023. Air temperature time series were extracted from ERA5, the fifth-generation atmospheric reanalysis developed by ECMWF (Hersbach et al., 2020, 2023), which provides consistent daily climate data since 1940 at 0.25° resolution using 4D-Var data assimilation technique and the CY41R2 forecasting system. To characterize precipitation isotope variability, we used monthly $\delta^{18}\text{O}$ and $\delta^2\text{H}$ outputs from IsoGSM (Kanamitsu et al., 2002a, b; Nan et al., 2021; Yoshimura et al., 2008; Bong et al., 2024), a general circulation model incorporating isotopic fractionation and nudged with NCEP (National Centres for Environmental Prediction) reanalysis data, covering 1979–2021. Additionally, long-term spatial isotope patterns were derived from the Isoscape product (etopo5, 2025; Bowen et al., 2005; Bowen and Revenaugh, 2003; Bowen and Wilkinson, 2002), which interpolates GNIP data using topographic and geographic predictors at a 5-arcminute resolution. Finally, seasonal isotope dynamics were represented using the sinusoidal precipitation isotope model (Allen et al., 2019), which fits annual sine curves to global precipitation isotope records and maps the resulting parameters to derive daily and monthly isotope predictions from 60°S to 90°N. Together, these products provide a consistent climatological and isotopic framework for interpreting site-level karst and cave hydrology.

Table 2: Information of the global products included in the WoKas-Iso database

Variable	Model Name	Data Source	Spatial Resolution	Temporal Resolution	Spatial Coverage	Time Coverage	Reference
Precipitation	MSWEP ¹ v2.8	Gauge, Satellite, Reanalysis	0.1°	Daily/Monthly	Global	1979–2020	(Beck et al., 2019b)
Precipitation isotopes	IsoGSM-NCEP-R2 ²	Reanalysis	200 km	Monthly	Global	1979–2021	(Yoshimura et al., 2008)
Precipitation isotopes	Isoscape	GNIP ⁴	5 arcmin	Monthly	Global	Jan–Dec	(Bowen et al., 2005)
Precipitation isotopes	Sine Curve	GNIP, CNIP ⁵ , USNIP ⁶	5 arcmin	Daily/Monthly	60°S– 90°N	Jan–Dec	(Allen et al., 2019)

Air temperature	ERA 5	Reanalysis	31 km	Daily/Monthly	Global	1940–2024	(Hersbach et al., 2023)
Evapotranspiration	GLEAM ³ 4.1a	Satellite, Reanalysis	0.1°	Daily/Monthly	Global	1980–2023	(Miralles et al., 2025)

398 ¹Multi-Source Weighted-Ensemble Precipitation ⁵Canadian Network for Isotopes in Precipitation

399 ²Isotopes-integrated Global Spectral Model ⁶US Network for Isotopes in Precipitation

400 ³Global Land Evaporation Amsterdam Model

401 ⁴Global Network of Isotopes in Precipitation

402 2.2.2 Evaluation of different precipitation isotope data

403 While global precipitation and evapotranspiration data were evaluated with observations in their original studies (see Table 2),
404 an evaluation of the isotopic composition of rainfall provided by the three precipitation isotope models—IsoGSM, Isoscape,
405 and the Sine Curve model, is still required. For that, we compared monthly simulated precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ against site-
406 specific observations from both cave drip water and karst spring locations. For precipitation $\delta^{18}\text{O}$ comparison, a total of 55
407 sites were analysed comprising 26 cave locations, 15 spring locations and 14 nearby meteorological stations to compensate the
408 limited availability of isotopic measurements at spring locations. For precipitation $\delta^2\text{H}$, 40 sites were assessed including 25
409 caves and 15 springs. For sites with daily precipitation, the monthly amount-weighted precipitation isotopes were calculated
410 by weighting the available isotope measurements with the corresponding daily precipitation amounts and aggregating them to
411 monthly values. For sites lacking daily precipitation, the stepwise interpolation of precipitation isotopes was applied to derive
412 monthly means.

413 We quantified the agreement of the global products and the observed isotopic composition of precipitation by the root-mean-
414 square error (RMSE) and the coefficient of determination (R^2). Fig.7 presents the performance metrics in predicting $\delta^{18}\text{O}$.
415 Based on RMSE, IsoGSM displays the lower median RMSE value ($\sim 2\text{‰}$) with a narrow interquartile and minimum outliers,
416 indicating the most consistent predictions across both caves and springs. The Isoscape model exhibits a comparable median
417 RMSE, though with a greater variability and higher outliers, reflecting less robustness. In contrast, the Sine Curve model yields
418 the highest median error ($>3\text{‰}$) and broader distributions, highlighting significantly inferior performance. Concerning R^2 ,
419 IsoGSM achieves the highest median R^2 (~ 0.6), however, its R^2 values span a wide range. The Isoscape model, having a
420 slightly lower median R^2 value (~ 0.5), shows a tighter distribution. The Sine Curve model performs worst, with median R^2 less
421 than 0.2 and a concentration of low values, indicating poor explanatory power.




Figure 7: Statistical comparison of three model performance for $\delta^{18}\text{O}$ in precipitation at cave (stars) and springs sites (circles). (a) Root Mean Square Error (RMSE); (b) Coefficient of determination (R^2). Violin plots show the distribution of performance metrics for each model: IsoGSM (red), Isoscape (green), and Sine Curve (blue). Boxplots within violin indicate the median, interquartile range and outliers. Lower RMSE and higher R^2 indicate better model performance.

Fig.8 presents the similar performance trends for $\delta^2\text{H}$ predictions. The IsoGSM achieves the lowest RMSE median ($\sim 17\%$) with a relatively narrow spread, reflecting the most accurate and consistent predictions. The Isoscape model follows closely with a slightly higher RMSE ($\sim 19\%$) and a greater variability. The Sine Curve model shows the highest RMSE median value ($\sim 29\%$) and the broadest RMSE distribution. For R^2 , there is a slight shift in performance ranking compared to $\delta^{18}\text{O}$. The Isoscape model has the highest R^2 value (~ 0.5). The IsoGSM model has a lower R^2 (~ 0.4) with a more balanced distribution. The Sine Curve model exhibits the poorest fit with a median R^2 below 0.2.




Figure 8: Statistical comparison of three model performance for $\delta^2\text{H}$ in precipitation at cave (stars) and springs sites (circles). (a) Root Mean Square Error (RMSE); (b) Coefficient of determination (R^2). Violin plots show the distribution of performance metrics for each model: IsoGSM (red), Isoscape (green), and Sine Curve (blue). Boxplots within violin indicate the median, interquartile range and outliers. Lower RMSE and higher R^2 indicate better model performance.

To evaluate the different performances of the global datasets, we compare time series of observed monthly $\delta^{18}\text{O}$ and $\delta^2\text{H}$ with model outputs from the Sine Curve, IsoGSM, and Isoscape models across all selected locations. Fig. 9 shows the $\delta^{18}\text{O}$ comparisons at four representative sites: Liangfeng Cave, Yongxing Cave, Chaoshuidong Spring and the Met-II.uz meteorological station near Anyak Spring (see supplement Section S1 for a complete comparison for other 51 sites). Overall, the Sine Curve model shows limited ability in reproducing the temporal course of the observations. In contrast, the IsoGSM model closely aligns the observed $\delta^{18}\text{O}$ dynamics at four sites, yielding high R^2 values (0.85, 0.81, 0.94 and 0.78 for Liangfeng Cave, Yongxing Cave, Chaoshuidong Spring and the Met-II.uz meteorological station near Anyak Spring, respectively) alongside relatively low RMSE values, which reflects robust performance in both correlation and prediction accuracy. The Isoscape model attains moderate predictions with R^2 exceeding 0.5 at all four sites. Although it outperforms the Sine Curve model, it tends to be less reliable than IsoGSM in capturing the isotopic variability.

Fig.10 presents $\delta^2\text{H}$ time series comparisons between observations and simulations from three models at four sites: Heshang Cave, Yongxing Cave, Krbavica Spring and Laolongshui Spring(comparison for other sites shown in supplement Figure S2). Similarly, the Sine Curve model shows the poorest agreement with observations at Heshang Cave, Yongxing Cave and Krbavica Spring as reflected by low R^2 values of 0.37, 0.57 and 0.1 respectively and high RMSEs (up to 50.11 ‰). Although it shows relatively higher R^2 values of 0.85 at Laolongshui Spring, its corresponding large RMSE value of 46.55 reveals substantial absolute predictive errors. The IsoGSM model shows the best performance at Heshang Cave and Yongxing Cave, with the highest R^2 (0.8 and 0.82) and lowest RMSE (15.53 ‰ and 10.56 ‰). These highlight IsoGSM model's strong predictive accuracy in tracking $\delta^2\text{H}$ variability at both sites. At Krbavica Spring, the IsoGSM and Isoscape exhibit comparable performance in terms of correlation with R^2 values of 0.71 and 0.7. However, IsoGSM shows a lower RMSE (21.3 ‰) compared to Isoscape (35.21 ‰), indicating that although both models reproduce the temporal variation similarly, IsoGSM offers more precise estimates of $\delta^2\text{H}$ magnitudes. At Laolongshui Spring (Fig.10d), the Isoscape model outperforms the IsoGSM model and Sine Curve model with the highest R^2 and lowest RMSE values.

Overall, the comparisons indicate that IsoGSM shows slightly superior performances over Isoscape and pronounced superior performance over the Sine Curve model. Different phase shifts between $\delta^{18}\text{O}$ and $\delta^2\text{H}$ in their Sine Curve model indicate a poor representation of their physical covariance which is given through the global/local meteoric waterline. Based on these results, IsoGSM is recommended as the preferred product for cave and spring locations where no local observations are available.




Figure 9: Time series comparison of modelled and observed monthly $\delta^{18}O$ in precipitation at two cave sites (a–b) and two spring sites (c–d). Observations (red stars) are compared with estimates from the Sine Curve model (dashed blue line), IsoGSM (purple crosses), and the Isoscape model (green circles). Each panel shows the number of monthly observations (n.obs) and model performance statistics (R² and RMSE).




Figure 10: Time series comparison of modelled and observed monthly δ^2H in precipitation at two cave sites (a–b) and two spring sites (c–d). Observations (red stars) are compared with estimates from the Sine Curve model (dashed blue line), IsoGSM (purple crosses), and the Isoscape model (green circles). Each panel shows the number of monthly observations (n_{obs}) and model performance statistics (R^2 and RMSE).

3 Database structure and usage notes

The datasets compiled in the WoKaS-Iso database are publicly archived as a zipped package titled “WoKaS_Iso_Data_Records” in OPARA, an open-access repository of research data from Saxon Universities (<https://doi.org/10.25532/OPARA-909>; Zang, 2025). The zip file is organized into three main folders: WoKaS_Iso_Input_Data, WoKaS_Iso_Output_Data, Global_Products_Data. Each folder is subdivided into several subfolders, with their structure and contents described in the following section:

3.1 WoKaS_Iso_Input_Data

Contains precipitation amount and precipitation isotopes ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) metadata tables and corresponding datasets for karst springs and cave sites:

- i. **WoKaS_Iso_Spring_Input_Metadata** provides summarised information on the precipitation and precipitation isotopes sites associated with each spring included in the database. Each spring is assigned a unique identifier as *wokas_iso_id*, which is constructed by using ISO country code (e.g., *AT* for Austria), a data type indicator (*S* for spring and *C* for cave) and a serial number (e.g., *AT-S-0001*). Some springs are associated with multiple precipitation or precipitation isotopes sites sharing the same *wokas_iso_id*, but each site is assigned its own *precipitation_entity_name* and *precip_iso_entity_name* as keys to link the metadata with their corresponding dataset. Furthermore, the *precip_distance_to_spring* and *precip_iso_distance_to_spring* indicate the distance in meters from each site to the specific spring. This structure allows users to flexibly select the desirable sites for their use. The full list of variables included in the spring input metadata table are explained in the Table 3.
- ii. **WoKaS_Iso_Spring_Input_Datasets** consists of 140 csv files. Each csv file represents the time series of precipitation amount and precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ for a specific spring. The csv file naming format is *Input-wokas_iso_id@spring_name* (e.g., *Input-AT-S-0002@Wasseralmquelle(A)*). Each csv file contains *wokas_iso_id*, *precipitation_entity_name*, *date_precip* (precipitation observation date), *precipitation*, *unit* (precipitation amount unit), *precip_iso_entity_name*, *date_precip_iso* (precipitation isotopes sampling date), *d18O*, *dD* and *note*.
- iii. **WoKaS_Iso_Cave_Input_Metadata** offers metadata summaries for cave sites input data, structured in the same way as the spring input metadata table. Similarly, each cave identifier is assigned to the cave site consisting of the ISO country code, the cave data type indicator *C* and its serial number (e.g., *AT-C-0001*). The entity name and distance fields align with the same conventions as springs. All variables are consistent with those used for spring metadata and are presented in Table 3.
- iv. **WoKaS_Iso_Cave_Input_Datasets** contains 67 csv files; each refers to a specific cave site. The csv file naming convention and internal structure follow the same pattern (e.g., *Input-AT-C-0001@Obir Cave*) as described in the spring input datasets section.

Table 3: Description of variables in the spring/cave input data metadata files

Variable	Description	Variable	Description
country	country the spring /cave is in	precip_iso_distance_to_spring/cave (m)	distance between precipitation isotope site and spring/cave (m)
wokas_iso_id	WoKaS-Iso identifier	precip_iso_entity_name	identifier assigned to precipitation isotopes site linked to a specific spring/cave
sisal_monv1_id/wokas_id	cave/spring identifier in SISAL_monv1/WoKaS database	precip_iso_latitude	precipitation isotopes measurements site latitude in decimal degrees. (N+, S-)

spring/cave_name	spring/cave name	precip_iso_longitude	precipitation isotopes measurements site longitude in decimal degrees. (E+, W-)
precipitation_site_name	location of precipitation measurements	precip_iso_elevation	meters above mean sea level
precip_distance_to_spring/cave(m)	distance between precipitation site and spring/cave	precip_d18O_start_date	start day of the oxygen-18 measurement in rainwater sample(dd/mm/yyyy)
precip_entity_name	identifier assigned to precipitation site linked to a specific spring/cave	precip_d18O_end_date	end day of the oxygen-18 measurement in rainwater sample(dd/mm/yyyy)
precip_latitude	P site latitude in decimal degrees. (N+, S-)	precip_d18O_observation_length	number of oxygen-18 measurements in rainwater
precip_longitude	P site longitude in decimal degrees. (E+, W-)	precip_dD_start_date	start day of the deuterium measurement in rainwater sample(dd/mm/yyyy)
precip_elevation	meters above mean sea level	precip_dD_end_date	end day of the deuterium measurement in rainwater sample(dd/mm/yyyy)
precipitation_start_date	start day of precipitation measurement(dd/mm/yyyy)	precip_dD_observation_length	number of deuterium measurements in rainwater
precipitation_end_date	end day of precipitation measurement(dd/mm/yyyy)	precip_iso_data_source	isotopic datasets in precipitation from who or where
precip_observation_length	number of precipitation measurements	reference	relevant literatures
precip_data_source	precipitation datasets from who or where	additional information	free-text column
precip_isotopes_site_name	location of precipitation Isotopes measurements		

3.2 WoKaS_Iso_Output_Data

Includes output metadata tables and output datasets for karst springs and cave drip sites:

- i. **WoKaS_Iso_Spring_Output_Metadata** summarizes attributes for karst springs documented in the WoKaS-Iso database including the geographic information of springs, an overview of karst spring discharge and isotopes for each spring. Each spring has the consistent identification through unique *wokas_iso_id*, same identifier used in the spring input data records. In the same way, the specific fields *discharge_entity_name* and *isotope_entity_name* are designed to establish the connection to their corresponding datasets. The complete list of parameters contained in the spring output metadata table are shown in Table 4.
- ii. **WoKaS_Iso_Spring_Output_Datasets** presents karst spring discharge, $\delta^{18}\text{O}$ and/or $\delta^2\text{H}$ datasets for 241 springs, organized as separate 241 csv files. Every csv file follows a standardized naming format “*Output-*

wokas_iso_id@spring_name” (e.g., *Output-AT-S-0002@Wasseralmquelle(A)*). The uniform internal structure of workbook includes the following columns: *wokas_iso_id*, *discharge_entity_name*, *date_discharge* (measurements date), *discharge*, *discharge_unit* (the unit of discharge), *iso_entity_name*, *date_iso* (isotope data sampling date), *d18O* and *dD*.

Table 4: Description of variables in the spring output metadata table

Variable	Description	Variable	Description
country	country the spring is in	discharge_data_source	spring discharge datasets from who or where
wokas_iso_id	WoKaS-Iso identifier	isotope_entity_name	identifier assigned to link the metadata to its isotopic datasets
wokas_id	spring identifier in WoKaS database	d18O_start_date	start day of oxygen-18 data collection of spring water sample(dd/mm/yyyy)
spring_name	spring name	d18O_end_date	end day of oxygen-18 data collection of spring water sample(dd/mm/yyyy)
latitude	spring latitude in decimal degrees. (N+, S-)	d18O_observation_length	number of oxygen-18 measurements
longitude	spring longitude in decimal degrees. (E+, W-)	d2H_start_date	start day of deuterium data collection of spring water sample(dd/mm/yyyy)
elevation(m)	meters above mean sea level	d2H_end_date	end day of deuterium data collection of spring water sample(dd/mm/yyyy)
type	spring	d2H_observation_length	number of deuterium measurements
discharge_entity_name	identifier assigned to link the metadata to its discharge datasets	isotope_data_source	spring isotopic datasets from who or where
discharge_start_date	start day of spring discharge measurement(dd/mm/yyyy)	reference	relevant literatures
discharge_end_date	end day of spring discharge measurement(dd/mm/yyyy)	additional_information	free-text column
discharge_observation_length	number of spring discharge observations		

- iii. **WoKaS_Iso_Cave_Output_Metadata** comprises a comprehensive overview of cave drip rate and cave drip isotopic records ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) for all cave drip sites. Each cave includes multiple drip sites which are assigned the same *wokas_iso_id* which is assigned in the cave input section but are distinguished by the unique *drip_rate_entity_name* and *drip_iso_entity_name* entries. Explanations for each variable in the cave output meta table is presented in Table 5.

iv. **WoKaS_Iso_Cave_Output_Datasets** contains 73 csv files. Each csv file stores the drip rate and drip isotopes datasets for the individual cave. The csv file name follows the format *Output-wokas_iso_id@cave_name* (e.g., *Output-AT-C-0001@Obir Cave*). Within each csv, multiple drip sites are documented using their unique entity names. The content includes the columns: *wokas_iso_id*, *drip_rate_entity_name*, *date_rate* (drip rate measurement date), *unit* (the unit of drip rate), *drip_iso_entity_name*, *date_iso* (sampling date of drip water isotopes), *d18O* and *dD*.

Table 5: Description of variables in the cave output metadata table

Variable	Description	Variable	Description
country	where the spring is located	drip_rate_observation_length	number of cave drip rate observations
wokas_iso_id	WoKaS-Iso identifier	drip_isotopes_entity_name	assigned identifier to link the metadata to related drip water isotopic data
sisal_monv1_id	cave identifier in SISAL_monv1 database	drip_d18O_start_date	start day of oxygen-18 data collection of drip water sample(dd/mm/yyyy)
name	cave name	drip_d18O_end_date	end day of oxygen-18 data collection of drip water sample(dd/mm/yyyy)
latitude	cave latitude in decimal degrees. (N+, S-)	drip_d18O_observation_length	number of drip oxygen-18 measurements
longitude	cave longitude in decimal degrees. (E+, W-)	drip_d2H_start_date	start day of deuterium data collection of drip water sample(dd/mm/yyyy)
elevation(m)	meters above mean sea level	drip_d2H_end_date	end day of deuterium data collection of drip water sample(dd/mm/yyyy)
type	cave	drip_d2H_observation_length	number of drip deuterium measurements
drip_site	cave drip site name	data_source	drip rate and drip isotopes datasets from where or who
drip_rate_entity_name	assigned identifier to link the metadata to related drip rate data	reference	relevant literatures
drip_rate_start_date	start day of cave drip rate measurement(dd/mm/yyyy)	additional_information	free-text column
drip_rate_end_date	end day of cave drip rate measurement(dd/mm/yyyy)		

3.3 Global_Products_Data

Contains gridded global product data and associated extraction tools used in WoKaS-Iso. It includes extracted datasets for WoKaS-Iso Spring and cave sites from six global products, global IsoGSM2 datasets prepared for user-defined extraction, and MATLAB scripts for extracting selected global products at specified geographic coordinates. The folder is organized into the

544 following subfolders: WoKaS_Iso_MSWEF, WoKaS_Iso_GLEAM, WoKaS_Iso_ERA5, WoKaS_Iso_IsoGSM,
 545 WoKaS_Iso_Isoscape, WoKaS_Iso_Sine_Curve, IsoGSM2_Global_DAT_for_Extraction, and WoKaS_Iso_Scripts.

- 546 • WoKaS_Iso_MSWEF, WoKaS_Iso_GLEAM, WoKaS_Iso_ERA5, WoKaS_Iso_Sine_Curve: the modelled data
 547 from these products are systematically sorted into four subfolders according to location type (spring or cave) and time
 548 resolution (daily or monthly). For example, WokaS_Iso_MSWEF folder includes
 549 WoKaS_Iso_Daily_MSWEF_Caves, WoKaS_Iso_Daily_MSWEF_Springs, WoKaS_Iso_Monthly_MSWEF_Caves,
 550 WoKaS_Iso_Monthly_MSWEF_Springs. Each subfolder contains compatible csv files storing continuous time series
 551 for individual spring or cave. The csv file names adhere to a standardized naming rule, combining the product name,
 552 temporal resolution, the *wokas_iso_id* and *spring /cave name*. For instance, *MSWEF_Daily_AT-C-0001@Obir Cave*
 553 refers to the daily gridded datasets from MSWEF at Obir Cave in Austria.
- 554 • WoKaS_Iso_IsoGSM, WoKaS_Iso_Isoscape: Since IsoGSM2 and Isoscape provide monthly $\delta^{18}\text{O}$ and $\delta^2\text{H}$ data only,
 555 each folder is subdivided into two categories for caves and springs: WoKaS_Iso_IsoGSM/Isoscape_Monthly_Caves
 556 and WoKaS_Iso_IsoGSM/Isoscape_Monthly_Springs. While, the csv file naming is slightly different from other
 557 products, constructed using *IsoGSM/Isoscape_wokas_iso_id@spring/cave_name* (e.g., *IsoGSM_AT-C-0001@Obir*
 558 *Cave*; *Isoscape_AT_C_0001@Obir Cave*).
- 559 • IsoGSM2_Global_DAT_for_Extraction: this folder contains the global IsoGSM2 precipitation isotopes datasets in
 560 DAT format. These data are not limited to WoKaS-Iso sites and can be used with the provided MATLAB extraction
 561 script to extract monthly precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values at any geographic coordinates of interest.
- 562 • WoKaS_Iso_Scripts: this folder contains MATLAB scripts developed to aid users in data extraction from selected
 563 global products. The scripts include for extracting daily and monthly hydroclimate data from NetCDF files of
 564 MSWEF, GLEAM, and ERA5, as well as routines for extracting precipitation isotope data from IsoGSM2 DAT files
 565 and Sine Curve model TIFF files. The accompanying README.md file provides detailed instructions on obtaining
 566 and preparing the required input datasets, including links to download or access the relevant global product data,
 567 preparing coordinate files for user-specified extraction, setting file paths and running the scripts, and interpreting the
 568 resulting CSV outputs. In addition, detailed comments are also included within each script to guide users through
 569 extraction workflow.

570 3.4 Usage notes

- 571 • **Metadata:** The columns *precip_data_source*, *precip_iso_data_source* (in the spring and cave input meta tables),
 572 *discharge_data_source*, *isotope_data_source* (in the spring output meta table), *data_source* (in the cave output meta
 573 table) specify the origins of respective data: precipitation, precipitation isotopes, discharge, spring isotopic data, cave
 574 drip rate and cave drip isotopes respectively. These sources can include data contributors, or digitization from specific
 575 figures or tables in publications, reports or theses, or retrieval from online repositories with links provided if
 576 applicable. All meta tables include the *reference* column listing all related literature that:(1) the spring/cave drip site

was studied; (2) the data were originally published; (3) contained figures and tables were digitized. These references enable users to get further context regarding datasets sampling method, climate conditions, hydrogeological settings of the studied spring/cave system. For peer-reviewed publications, DOIs are offered, while URLs of reports or theses are provided if available.

- **Machine-readability:** All csv files storing input and output datasets are designed in a machine-readable format to facilitate efficient and automated data processing.
- **Link the metadata to datasets:** Each csv file name incorporates the unique *wokas_iso_id* that is organized in the metadata tables. This identifier enables location of the related spring or cave dataset by matching with file name (e.g. *Output-wokas_iso_id@spring_name*). Additionally, within each csv file, the entity name fields (*precip_entity_name*, *precip_iso_entity_name*, *discharge_entity_name*, *isotope_entity_name*, *drip_rate_entity_name* and *drip_iso_entity_name*) can work as the identified keys to connect with the corresponding metadata entries. These identifiers can be utilized to retrieve specific time series data from the measurement sites of interest.
- **Global products datasets:** Model-derived data corresponding to each site are provided in CSV files named according to their respective *wokas_iso_id* for easy reference. Monthly datasets, dates are separated into two columns: year and month, while daily data use a single dd/mm/yyyy format. These data represent the closest grid cell to each site; therefore, users should be aware of potential spatial differences when comparing with local observations. Proper citation of original data sources is required when utilizing these datasets.
- **Global product scripts:** The MATLAB scripts allow users to extract selected global product data for additional coordinates or updated input datasets. For MSWEP, GLEAM, and ERA5, the extractable time period depends on the NetCDF files provided by the user. IsoGSM2 extraction is based on the provided monthly DAT-format datasets covering 1979–2021, while the Sine Curve model can generate daily or monthly values for free years. Automated data download is not included; users should follow README.md file for data preparation, path settings, and script execution.
- **Isoscape data extraction:** The Isoscape product provides gridded raster datasets in Geo Tiff format. In WoKaS-Iso, the precipitation isotope values were extracted from the global gridded maps using ArcGIS Pro. The global gridded maps of precipitation $\delta^{18}\text{O}$ and $\delta^2\text{H}$ are available from the WaterIsotopes.org (Bowen, 2026). For the mean monthly precipitation isotope grids, the interpolation method follows the Bowen et al. (2005), with GNIP data used as the main data source (IAEA/WMO, 2015). The monthly Isoscape products consist of separate gridded maps for each month from January to December, for both $\delta^{18}\text{O}$ and $\delta^2\text{H}$. To extract values for new locations, users should first download the relevant monthly maps, load them into ArcGIS, and import corresponding coordinates as a CSV file. After extraction, the resulting table provides monthly $\delta^{18}\text{O}$ and $\delta^2\text{H}$ estimates for each site from January to December. Since this procedure relies on GIS-based raster extraction, no MATLAB extraction routine is provided for Isoscape in WoKaS_Iso_Scripts.

- Example use case:** A typical workflow starts by selecting a spring or cave site from the output metadata tables and identifying its *wokas_iso_id* and *spring/cave name*, for example CN-S-0006 and Banzhai Spring. Users can then locate the matching output CSV file Output-CN-S-0006@Banzhai Spring in the WoKaS_Iso_Spring_Output_Datasets folder (or WoKaS_Iso_Cave_Output_Datasets folder for a selected cave site) using this identifier CN-S-0006 or spring name Banzhai Spring. In this output file, the columns *date_iso*, *d18O*, and *dD* provide the isotope time series, while *date_discharge* and *discharge* provide the discharge record where available. If precipitation or precipitation isotope input data are required, users can check the input metadata tables and locate corresponding input CSV file Input-CN-S-0006@Banzhai Spring in WoKaS_Iso_Spring_Input_Datasets (or WoKaS_Iso_Cave_Input_Datasets folder for a cave).

If measured input data are unavailable, global product data can be used as an alternative. For Banzhai Spring, users first select the relevant product folder according to the required variable, such as WoKaS_Iso_MSWEF for precipitation, WoKaS_Iso_GLEAM for AET and PET, WoKaS_Iso_ERA5 for air temperature, and WoKaS_Iso_IsoGSM, WoKaS_Iso_Isoscape, or WoKaS_Iso_Sine_Curve for precipitation isotopes. Within each product folder, users then select the subfolder corresponding to spring sites and the required temporal resolution. For example, daily precipitation data can be found in WoKaS_Iso_MSWEF/WoKaS_Iso_Daily_MSWEF_Springs/ with the file MSWEF_Daily_CN-S-0006@Banzhai Spring, while monthly precipitation data can be found in WoKaS_Iso_MSWEF/WoKaS_Iso_Monthly_MSWEF_Springs/ with the file MSWEF_Monthly_CN-S-0006@Banzhai Spring. The same logic applies to other products, for example GLEAM_Daily_CN-S-0006@Banzhai Spring or GLEAM_Monthly_CN-S-0006@Banzhai Spring in the corresponding GLEAM spring subfolders, ERA5_Daily_CN-S-0006@Banzhai Spring or ERA5_Monthly_CN-S-0006@Banzhai Spring in the corresponding ERA5 spring subfolders, and Sine_Curve_Daily_CN-S-0006@Banzhai Spring or Sine_Curve_Monthly_CN-S-0006@Banzhai Spring in the corresponding Sine Curve spring subfolders. IsoGSM2 and Isoscape are provided at monthly resolution only and can be accessed through their spring subfolders using files such as IsoGSM_CN-S-0006@Banzhai Spring and Isoscape_CN-S-0006@Banzhai Spring.

For cave sites, users can follow the same procedure using the corresponding cave metadata tables, cave input and output dataset folders, cave subfolders within Global_Products_Data, and cave identifiers or names in the CSV filenames.

4 Data quality control and evaluation

4.1 Data quality control

A series of quality control measures were implemented throughout the large datasets synthesized in the WoKaS-Iso database to ensure data precision, consistency and integrity:

641 **Site information verification:** All spring and cave drip site names, coordinates, elevation, data units, precipitation sites and
642 precipitation isotope sampling sites were initially cross-checked against the original sources: supplementary materials, tables,
643 original data contributed by collaborators or digitized from related figures. Additionally, all the time series were reviewed for
644 consistency with the original datasets, ensuring correct sampling date accurately matching its associated measurements to
645 avoid any discrepancies.

646 **Standardized formatting:**

- 647 i. Naming convention: All *wokas_iso_id* entries were validated to confirm conformity with the standardized naming
648 convention including verification of the ISO country code, data type indicator (C for cave and S for spring) and
649 sequential numbering within each country. The csv file names were systematically reviewed folder by folder to ensure
650 compliance with the naming patterns stated in section 3.
- 651 ii. CSV file structure: All files were checked to be machine-readable in comma-delimited csv format. Each csv file was
652 examined to guarantee a consistent internal structure with uniform column headers.
- 653 iii. Date and coordinates formatting: The dates were formatted consistently in dd/mm/yyyy across all datasets. The
654 latitude and longitude were formatted with four decimals.

655 **Consistency checks:**

- 656 i. Data consistency: The dates and corresponding time series of all datasets were checked to confirm they were sorted
657 chronologically.
- 658 ii. Spatial consistency: Latitude and longitude were validated to fall in valid ranges (N+, S-, E+, W-), and validation of
659 spatial distribution on map were verified by visualization.
- 660 iii. Metadata and datasets matching: The identifiers *wokas_iso_id*, *precip_entity_name*, *precip_iso_entity_name*,
661 *discharge_entity_name*, *iso_entity_name*, *drip_rate_entity_name* and *drip_iso_entity_name* were cross validated
662 against metadata to ensure correct linkage with the corresponding datasets. In addition, the start date, end date and
663 observation length of all time series were checked to align with actual datasets.

664 **Data source verification:** Checking data sources recorded in the meta tables was implemented to ensure proper citation and
665 traceability. The indicated figures and tables were checked to be correctly represented. All provided URLs access to online
666 database were tested to verify datasets retrieval. The DOIs (provided as web links) were confirmed working and accurately
667 linked to the cited publications. Where available, links of reports and theses were also tested for accessibility.
668 For cave drip water datasets originating from SISAL_mon_v1, WoKaS-Iso builds on the automated quality control already
669 applied in Treble et al. (to be submitted). Within WoKaS-Iso quality control, the SISAL_mon_v1 records were therefore not
670 re-processed from scratch, but underwent additional harmonisation and validation steps: mapping to the WoKaS-Iso file and
671 naming conventions, cross-checking identifiers and time ranges against the new metadata structure, and ensuring consistent
672 linkage and interoperability with associated precipitation and spring datasets and references as described in the following.

4.2 Data quality attribution

The datasets quality for each spring and cave site was assessed using a scoring scheme that considers data availability and data source reliability for both system output and input variables (Table 6). For system output data, **collected spring discharge/drip rate** and **collected spring/drip water isotope** were obtained from research collaborators, original authors, online databases or supplementary materials of publications. These sources are generally of high quality and were hence assigned a score of 3. **digitized spring discharge/drip rate** and **digitized spring/drip water isotope** were digitized from the plots in publications or reports or theses. Although the extraction process was carried out as precisely as possible, some inaccuracies and uncertainties inherently remained due to the resolution and temporal coverage of the original plots. Thus, the score of 2 was assigned to the digitized data. For those locations without relevant data were given a score of 0. For system input data, three quality levels were defined according to the sources of precipitation isotopes and precipitation information: local measurements, meteorological stations and global models. The **local observation (precip. isotope/amount)** acquired from on-site monitoring, which are rare and valuable particularly for karst springs, were thus assigned the highest score of 3. The **meteorological stations (precip.isotope/amount)** data sourced from meteorological stations, while still based on measurements, are deemed less directly representative of site-specific conditions and were scored 2. In cases where neither local nor station-based input were available, the input data from **Global products** were provided as substitutes and assigned a score of 1. The total grade for each spring and cave site was calculated from the sum of its output and input data scores, based on the assessment criteria stated above. The sites evaluation was then divided into three classes that reflect the overall reliability and completeness of the data collection.

- **Class I** (scores from 10 to 12): indicates robust datasets with high quality, generally obtained from raw output origins and/or locally measured inputs. The site in this category keeps the most complete and reliable datasets, providing comprehensive information for interpretation and modelling of karst systems.
- **Class II** (scores from 7 to 9): reflects the datasets with moderate quality. These sites usually retain the main output and input information required for analysis, but some records may be digitized or supplemented by less site-specific input sources. Although less strong, these data are still valuable for studies in both karst geohydrology and speleology.
- **Class III** (scores below 7): represents datasets with limited data availability and lower source reliability. Sites in this category are generally characterized by missing variables and/or stronger reliance on digitized records or global modelled data. These datasets are helpful in identifying data-scarce regions and highlighting the need for more data collection.

Table 6: The grading system for evaluation of the WoKaS-Iso database for karst springs and caves. *Example:* a spring with collected spring discharge (3 points), collected spring isotopes (3 points), local observation of precipitation isotopes (3 points), and local precipitation amount (3 points) receives a total score of 12. *Note:* the “No data” category is assigned to sites without flow rate data or aligned with SISAL_mon_v1 database at Shenqi Cave, as no cave drip water data are available and only input data are provided.

System output data		System input data	
Variable	Assigned Score	Variable	Assigned Score
Collected spring discharge/drip rate	3	Local observation (precip. isotope/amount)	3
Collected spring /drip water isotope			
Digitized spring discharge/drip rate	2	Meteorological stations (precip.isotope/amount)	2
Digitized spring/drip water isotope			
No data	0	Global products	1

Attributing the quality classes to all our sites shows that most of them fall within Class I and Class II (Figure 11). For springs, 21% are categorized into Class I, and are predominantly distributed in China, Europe and the Middle East. Class II springs, the most common category, occupied 46%, due to scarcity and limited availability of locally measured rainfall and rainwater isotopes for karst springs; while Class III accounts for 33%, with a high proportion due to the absence of paired precipitation and precipitation isotopes measurements for karst springs, the substitute modelled data were used. Class I and Class II are spread relatively evenly globally, while Class III springs are primarily found in Europe where karst spring monitoring is more extensive in the WoKaS-Iso database. For cave sites, a high proportion of 50% is classified as Class I, Class II reached 46%, with only a small proportion of 4% falling into Class III. The distribution characteristics reflect the more complete and robust monitoring systems available for caves compared to karst springs. The Class I cave sites are mainly suited in Australia, China and Europe. The Class II caves are mostly located in Europe and the USA. Only 4 caves are grouped as Class III, scattered over various regions.




Figure 11: Spatial distribution of karst springs and cave sites in the WoKaS-Iso database according to data quality evaluation classes. Blue areas indicate karst regions as delineated in the World Karst Aquifer Map (WOKAM, Chen et al., 2017). Star symbols refer to caves and circle symbols represent springs. Colours represent the three quality tiers: Class I (red), Class II (orange), and Class III (blue). Accompanying pie charts illustrate the proportion of each quality class for springs and caves, respectively.

4.3 Measurement gap characteristics of isotope records

In addition to the source-based quality control and completeness checks, we provide an additional characterization of isotope measurements intervals within WoKaS-Iso. Since stable isotope records are often collected at irregular frequencies, the time interval between consecutive observations provides useful complementary information on the temporal structure of the dataset. For this purpose, time gaps were calculated between successive sampling dates within each individual isotope record karst springs and cave drip water using all available $\delta^{18}\text{O}$ and $\delta^2\text{H}$ observations. The resulting gap distributions were summarized separately for karst spring and cave drip water records using empirical cumulative distribution functions (ECDFs; Fig. 12). For karst spring records, the ECDF curve (Fig.12a) rises markedly around the 7-day threshold, showing a substantial contribution of weekly observations. The median gap is 15 days, and the interquartile range of 7–30 days indicates that the middle 50% of spring observation intervals span from weekly to monthly timescales, reflecting heterogeneous observation frequencies across sites. Accordingly, 34 % of spring gaps are shorter than or equal to 7 days, while nearly 80 % are shorter than or equal to 31 days.




Figure 12: Empirical cumulative distribution functions (ECDFs) of time gaps between consecutive stable isotope measurements for (a) karst spring and (b) cave drip water records included in WoKaS-Iso. Statistics within each panel show the median gap, interquartile range (IQR), and the proportion of observations sampled within 7, 31, and 90 days. Dashed lines denote the corresponding temporal thresholds. The cave drip water ECDF remains low at weekly timescales but rises sharply around 31 days. This pattern indicates that cave drip water isotope measurements are strongly concentrated around monthly intervals. The median gap of 31 days and the narrower interquartile range of 26–45 days further suggest more regular observation intervals compared with karst springs. Only 3.5 % of cave drip-water gaps are shorter than or equal to 7 days, whereas 51 % are shorter than or equal to 31 days. Despite these differences at weekly-to-monthly timescales, both systems converge near the 90-day threshold, with more than 94 % of gaps shorter than 90 days, indicating that most records were sampled at frequencies finer than seasonal timescales.

5 Data availability

The dataset is available in the public repository at: <https://doi.org/10.25532/OPARA-909> (Zang, 2025).

6 Recommended usage and outlook

The data quality classification provides guidance for users on the proper application and interpretation of each site. **Class I** with the most complete observational records are recommended for process-based model calibration and validation, characterization of the karst system and paleoclimate construction. **Class II** consisting of partial measurements or digitized output data or station-based input data/ modelled sources are appropriate for regional-scale comparative studies, trend analysis

and model sensitivity testing. Although less precise than Class I, these datasets remain valuable in regions with limited monitoring. **Class III**, which rely highly on digitised outputs and global products inputs, are useful for global exploratory studies, particularly in data-scarce regions. Global products offer critical alternatives where observations are unavailable and support long-term climate analysis. Based on our analysis, we recommend **IsoGSM** as the preferred substitute of precipitation isotope input. Future development of WoKaS-Iso efforts will focus on expanded spatial coverage and improving data quality through encouraging the sharing of raw data from literatures, reports, internal databases and unpublished records by researchers, institutions and monitoring programs, especially in underrepresented regions such as Africa, South America and Southeast Asia. Enhancements will include more detailed metadata, uncertainty estimates, data quality flags to improve usability. A web-based interface with visualizations is planned to assist data access.

7 Author contributions

YZ and AH collected the karst spring datasets. PCT and YZ contacted data contributors to obtain cave drip water data and aligned this dataset with the SISAL_mon_v1 dataset. YZ designed the structure of WoKaS-Iso database and organized the datasets. YZ digitized data from publication figures. YZ, AH, JGP and FZ contributed to data quality control. KY extracted IsoGSM datasets for each site and YZ performed data extraction from other global products. YZ developed extraction scripts for MSWEP, GLEAM and ERA5, with KOC and XM assisting in code testing. YZ drafted the manuscript under the supervision of AH. All other authors contributed data to the WoKaS-Iso database.

8 Competing interests

The authors declare that they have no conflict of interest.

9 Acknowledgment

Yining Zang acknowledges the financial support from the China Scholarship Council (202207720061). Yining Zang and Andreas Hartmann were supported by the German Research Foundation (DFG, grant number: HA 8113/6–1, project “Robust Conceptualisation of KARst Transport (ROCKAT)”). We want to thank Nikita Kaushal (American Museum of Natural History) for her help in aligning WoKaS-Iso with SISAL_mon_v1, and Qi Li (Robotics Research Lab, University Kaiserslautern-Landau) for assistance in writing global products extraction scripts. We also extend our gratitude to all researchers and institutions who provided data essential for this compilation. In addition, we acknowledge Albert Goede (Retired, formerly University of Tasmania, Australia), Chaojun Chen (No.768, Juxian Street, Chenggong District, Kunming, Yunnan, 650500, China), Juan Pablo Bernal (Instituto de Geociencias, Campus UNAM Juriquilla, Querétaro, México, 76230), Paul W. Williams (Retired, formerly University of Auckland, School of Environment, Auckland, New Zealand), Quan Wang, Sebastian

780 Breitenbach (Department of Geography and Environmental Sciences, Ellison D116, Northumbria University, Newcastle upon
 781 Tyne, NE1 8ST, UK), Yongjin Wang (School of Geography, Nanjing Normal University, Nanjing 210023, China), Yunxia Li
 782 (Hunan Normal University, No. 36 Lushan South Road, Changsha 410081, China) for their valuable contribution to both the
 783 SISAL_mon_v1 and WoKaS-Iso databases.

784 References

- 785
- 786 Allen, S. T., Jasechko, S., Berghuijs, W. R., Welker, J. M., Goldsmith, G. R., and Kirchner, J. W.: Global sinusoidal seasonality
 787 in precipitation isotopes, *Hydrol Earth Syst Sci*, 23, 3423–3436, <https://doi.org/10.5194/hess-23-3423-2019>, 2019.
- 788 Andreo, B., Liñán, C., Carrasco, F., Jiménez de Cisneros, C., Caballero, F., and Mudry, J.: Influence of rainfall quantity on
 789 the isotopic composition (^{18}O and ^2H) of water in mountainous areas. Application for groundwater research in the Yunqueira-
 790 Nieves karst aquifers (S Spain), *Applied Geochemistry*, 19, 561–574, <https://doi.org/10.1016/J.APGEOCHEM.2003.08.002>,
 791 2004.
- 792 Ankit Rohatgi: WebPlotDigitizer, version 4.6, <https://apps.automeris.io>, 2023.
- 793 etopo5: <https://pubs.usgs.gov/of/1998/of98-801/bathy/metadata/etopo5.htm>, last access: 9 April 2025.
- 794 Bakalowicz, M.: Karst groundwater: A challenge for new resources, *Hydrogeol J*, 13, 148–160,
 795 <https://doi.org/10.1007/S10040-004-0402-9/FIGURES/6>, 2005.
- 796 Baker, A., Hartmann, A., Duan, W., Hankin, S., Comas-Bru, L., Cuthbert, M. O., Treble, P. C., Banner, J., Genty, D., Baldini,
 797 L. M., Bartolomé, M., Moreno, A., Pérez-Mejías, C., and Werner, M.: Global analysis reveals climatic controls on the oxygen
 798 isotope composition of cave drip water, *Nat Commun*, 10, 1–7, <https://doi.org/10.1038/S41467-019-11027-W>;SUBJMETA,
 799 2019.
- 800 Beck, H. E., van Dijk, A. I. J. M., Levizzani, V., Schellekens, J., Miralles, D. G., Martens, B., and de Roo, A.: MSWEP: 3-
 801 hourly 0.25° global gridded precipitation (1979–2015) by merging gauge, satellite, and reanalysis data, *Hydrol. Earth Syst.*
 802 *Sci.*, 21, 589–615, <https://doi.org/10.5194/hess-21-589-2017>, 2017.
- 803 Beck, H. E., Pan, M., Roy, T., Weedon, G. P., Pappenberger, F., van Dijk, A. I. J. M., Huffman, G. J., Adler, R. F., and Wood,
 804 E. F.: Daily evaluation of 26 precipitation datasets using Stage-IV gauge-radar data for the CONUS, *Hydrol. Earth Syst. Sci.*,
 805 23, 207–224, <https://doi.org/10.5194/hess-23-207-2019>, 2019a.
- 806 Beck, H. E., Wood, E. F., Pan, M., Fisher, C. K., Miralles, D. G., van Dijk, A. I. J. M., McVicar, T. R., and Adler, R. F.:
 807 MSWEP V2 Global 3-Hourly 0.1° Precipitation: Methodology and Quantitative Assessment, *Bull. Amer. Meteor. Soc.*, 100,
 808 473–500, <https://doi.org/10.1175/BAMS-D-17-0138.1>, 2019b.
- 809 Birkel, C. and Soulsby, C.: Advancing tracer-aided rainfall–runoff modelling: a review of progress, problems and unrealised
 810 potential, *Hydrol Process*, 29, 5227–5240, <https://doi.org/10.1002/HYP.10594>, 2015.

811 Bong, H., Cauquoin, A., Okazaki, A., Chang, E.-C., Werner, M., Wei, Z., Yeo, N., and Yoshimura, K.: Process-based
812 intercomparison of water isotope-enabled models and reanalysis nudging effects, *J. Geophys. Res.-Atmos.*, 129,
813 e2023JD038719, <https://doi.org/10.1029/2023JD038719>, 2024.

814 Bowen, G. J. and Revenaugh, J.: Interpolating the isotopic composition of modern meteoric precipitation, 39, 1299,
815 <https://doi.org/10.1029/2003WR002086>, 2003.

816 Bowen, G. J. and Wilkinson, B.: Spatial distribution of $\delta^{18}\text{O}$ in meteoric precipitation, *Geology*, 315–318,
817 [https://doi.org/10.1130/0091-7613\(2002\)030](https://doi.org/10.1130/0091-7613(2002)030), 2002.

818 Bowen, G. J., Wassenaar, L. I., and Hobson, K. A.: Global application of stable hydrogen and oxygen isotopes to wildlife
819 forensics, *Oecologia*, 143, 337–348, <https://doi.org/10.1007/S00442-004-1813-Y>, 2005.

820 Bowen, G. J.: Gridded maps of the isotopic composition of meteoric waters, *WaterIsotopes.org*, available at:
821 https://wateriso.utah.edu/waterisotopes/pages/data_access/ArcGrids.html, last access: 17 June 2026, 2026.

822 Bradley, C., Baker, A., Jex, C. N., and Leng, M. J.: Hydrological uncertainties in the modelling of cave drip-water $\delta^{18}\text{O}$ and the implications for
823 stalagmite palaeoclimate reconstructions, *Quat Sci Rev*, 29, 2201–2214, <https://doi.org/10.1016/J.QUASCIREV.2010.05.017>,
824 2010.

825 Chen, Z., Auler, A. S., Bakalowicz, M., Drew, D., Griger, F., Hartmann, J., Jiang, G., Moosdorf, N., Richts, A., Stevanovic,
826 Z., Veni, G., and Goldscheider, N.: The World Karst Aquifer Mapping project: concept, mapping procedure and map of Europe.
827 *Hydrogeol J* 25, 771–785, <https://doi.org/10.1007/s10040-016-1519-3>, 2017.

828 Frank, S., Goeppert, N., and Goldscheider, N.: Improved understanding of dynamic water and mass budgets of high-alpine
829 karst systems obtained from studying a well-defined catchment area, *Hydrol Process*, 35, e14033,
830 <https://doi.org/10.1002/HYP.14033>, 2021.

831 Giese, M., Caballero, Y., Hartmann, A., and Charlier, J. B.: Trends in long-term hydrological data from European karst areas:
832 Insights for groundwater recharge evaluation, *Hydrol Earth Syst Sci*, 29, 3037–3054, [https://doi.org/10.5194/HESS-29-3037-](https://doi.org/10.5194/HESS-29-3037-2025)
833 2025, 2025.

834 Goldscheider, N. and Drew, D.: Methods in karst hydrogeology. *International Contribution to Hydrogeology*, 264, 2007.

835 Hartmann, A. and Baker, A.: Modelling karst vadose zone hydrology and its relevance for paleoclimate reconstruction, *Earth*
836 *Sci Rev*, 172, 178–192, <https://doi.org/10.1016/J.EARSCIREV.2017.08.001>, 2017.

837 Goldscheider, N., Chen, Z., Auler, A. S., Bakalowicz, M., Broda, S., Drew, D., Hartmann, J., Jiang, G., Moosdorf, N.,
838 Stevanović, Z., and Veni, G.: Global distribution of carbonate rocks and karst water resources, *Hydrogeol. J.*, **28**, 1661–1677,
839 <https://doi.org/10.1007/s10040-020-02139-5>, 2020.

840 Hartmann, A., Kralik, M., Humer, F., Lange, J., and Weiler, M.: Identification of a karst system's intrinsic hydrodynamic parameters: Upscaling from single springs to the whole aquifer,
841 *Environ Earth Sci*, 65, 2377–2389, <https://doi.org/10.1007/s12665-011-1033-9>, 2012.

842 Hartmann, A., Wagener, T., Rimmer, A., Lange, J., Brielmann, H., and Weiler, M.: Testing the realism of model structures to
843 identify karst system processes using water quality and quantity signatures, *Water Resour Res*, 49, 3345–3358,
844 <https://doi.org/10.1002/wrcr.20229>, 2013.

845 Hartmann, A., Goldscheider, N., Wagener, T., Lange, J., and Weiler, M.: Karst water resources in a changing world: Review
846 of hydrological modeling approaches, *Rev. Geophys.*, 52, 218–242, <https://doi.org/10.1002/2013RG000443>, 2014.

847 Hartmann, A., Jasechko, S., Gleeson, T., Wada, Y., Andreo, B., Barberá, J. A., Brielmann, H., Bouchaou, L., Charlier, J.,
848 Darling, W. G., Filippini, M., Garvelmann, J., Goldscheider, N., Kralik, M., Kunstmann, H., Ladouche, B., Lange, J.,
849 Lucianetti, G., Martín, J. F., Mudarra, M., Sánchez, D., Stumpp, C., Zagana, E., and Wagener, T.: Risk of groundwater
850 contamination widely underestimated because of fast flow into aquifers, *Proc. Natl. Acad. Sci. U.S.A.*, 118, e2024492118,
851 <https://doi.org/10.1073/pnas.2024492118>, 2021.

852 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,
853 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara,
854 G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L.,
855 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P.,
856 Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal*
857 *Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

858 Hersbach, Hans., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum,
859 I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: Copernicus Climate Change Service, Climate Data Store,
860 (2023): ERA5 hourly data on single levels from 1940 to present., Copernicus Climate Change Service (C3S) Climate Data
861 Store (CDS), <https://doi.org/10.24381/cds.adbb2d47>, 2023.

862 Hulsman, P., Keune, J., Koppa, A., Schellekens, J., and Miralles, D. G.: Incorporating Plant Access to Groundwater in Existing
863 Global, Satellite-Based Evaporation Estimates, *Water Resour Res*, 59, <https://doi.org/10.1029/2022WR033731>,
864 2023.IAEA/WMO: Global Network of Isotopes in Precipitation, The GNIP Database, available at:
865 <https://nucleus.iaea.org/wiser>, last access: 17 June 2026, 2015.

866 Jourde, H., Massei, N., Mazzilli, N., Binet, S., Batiot-Guilhe, C., Labat, D., Steinmann, M., Bailly-Comte, V., Seidel, J.L.,
867 Arfib, B., Charlier, J.B., Guinot, V., Jardani, A., Fournier, M., Aliouache, M., Babic, M., Bertrand, C., Brunet, P., Boyer, J.F.,
868 Bricquet, J.P., Camboulive, T., Carrière, S.D., Celle-Jeanton, H., Chalikakis, K., Chen, N., Cholet, C., Clauzon, V., Soglio,
869 L.D., Danquigny, C., Défargue, C., Denimal, S., Emblanch, C., Hernandez, F., Gillon, M., Gutierrez, A., Sanchez, L.H., Hery,
870 M., Houillon, N., Johannet, A., Jouves, J., Jozja, N., Ladouche, B., Leonardi, V., Lorette, G., Loup, C., Marchand, P., de
871 Montety, V., Muller, R., Ollivier, C., Sivelles, V., Lastennet, R., Lecoq, N., Maréchal, J.C., Perotin, L., Perrin, J., Petre, M.A.,
872 Peyraube, N., Pistre, S., Plagnes, V., Probst, A., Probst, J.L., Simler, R., Stefani, V., Valdes-Lao, D., Viseur, S. and Wang, X.:
873 SNO KARST: A French Network of Observatories for the Multidisciplinary Study of Critical Zone Processes in Karst
874 Watersheds and Aquifers. *Vadose Zone Journal*, 17: 1-18 180094. <https://doi.org/10.2136/vzj2018.04.0094>, 2018.

875 Kanamitsu, M., Kumar, A., Juang, H. H., Schemm, J., Wang, W., Yang, F., Hong, S., Peng, P., Chen, W., Moorthi, S., and Ji,
876 M.: NCEP dynamical seasonal forecast system 2000, *Bull. Amer. Meteor. Soc.*, 83, 1019–1038, [https://doi.org/10.1175/1520-0477\(2002\)083<1019:NDSFS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2002)083<1019:NDSFS>2.3.CO;2), 2002a.

878 Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S., Hnilo, J. J., Fiorino, M., & Potter, G. L.: NCEP–DOE AMIP-II
879 Reanalysis (R-2), *Bull. Amer. Meteor. Soc.*, 83,1631–1644. <https://doi.org/10.1175/BAMS-83-11-1631>, 2002b.

880 Kaushal, N., Lechleitner, F. A., Wilhelm, M., Azennoud, K., Bühler, J. C., Braun, K., Ait Brahim, Y., Baker, A., Burstyn, Y.,
881 Comas-Bru, L., Fohlmeister, J., Goldsmith, Y., Harrison, S. P., Hatvani, I. G., Rehfeld, K., Ritzau, M., Skiba, V., Stoll, H. M.,
882 Szűcs, J. G., Tanos, P., Treble, P. C., Azevedo, V., Baker, J. L., Borsato, A., Chawchai, S., Columbu, A., Endres, L., Hu, J.,
883 Kern, Z., Kimbrough, A., Koç, K., Markowska, M., Martrat, B., Masood Ahmad, S., Nehme, C., Novello, V. F., Pérez-Mejías,
884 C., Ruan, J., Sekhon, N., Sinha, N., Tadros, C. V., Tiger, B. H., Warken, S., Wolf, A., Zhang, H., and SISAL Working Group
885 members: SISALv3: a global speleothem stable isotope and trace element database, *Earth Syst. Sci. Data*, 16, 1933–1963,
886 <https://doi.org/10.5194/essd-16-1933-2024>, 2024.

887 Koppa, A., Rains, D., Hulsman, P., Poyatos, R., and Miralles, D. G.: A deep learning-based hybrid model of global terrestrial
888 evaporation, *Nat Commun*, 13, <https://doi.org/10.1038/s41467-022-29543-7>, 2022.

889 Labat, D., Sivakumar, B., and Mangin, A.: Evidence for deterministic chaos in long-term high-resolution karstic streamflow
890 time series, *Stoch. Environ. Res. Risk Assess.*, 30, 2189–2196, <https://doi.org/10.1007/s00477-015-1175-5>, 2016.

891 Li, G., Chen, X., Zhang, Z., Wang, L., and Soulsby, C.: Effects of passive-storage conceptualization on modeling hydrological
892 function and isotope dynamics in the flow system of a cockpit karst landscape, *Hydrol Earth Syst Sci*, 26, 5515–5534,
893 <https://doi.org/10.5194/hess-26-5515-2022>, 2022.

894 Li, R., Zhu, G., Chen, L., Qi, X., Lu, S., Meng, G., Wang, Y., Li, W., Zheng, Z., Yang, J., and Gun, Y.: Global Stable Isotope
895 Dataset for Surface Water, *Earth Syst Sci Data*, 17, 2135–2145, <https://doi.org/10.5194/ESSD-17-2135-2025>, 2025.

896 Liu, J., Shangguan, D., Liu, S., Ding, Y., Wang, S., and Wang, X.: Evaluation and comparison of CHIRPS and MSWEP daily-
897 precipitation products in the Qinghai-Tibet Plateau during the period of 1981–2015, *Atmos Res*, 230, 104634,
898 <https://doi.org/10.1016/J.ATMOSRES.2019.104634>, 2019.

899 Liu, Y. and Yamanaka, T.: Tracing groundwater recharge sources in a mountain–plain transitional area using stable isotopes
900 and hydrochemistry, *J Hydrol (Amst)*, 464–465, 116–126, <https://doi.org/10.1016/J.JHYDROL.2012.06.053>, 2012.

901 Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W.
902 A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10,
903 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.

904 Mayer-Anhalt, L., Birkel, C., Sánchez-Murillo, R., and Schulz, S.: Tracer-aided modelling reveals quick runoff generation and
905 young streamflow ages in a tropical rainforest catchment, *Hydrol Process*, 36, e14508, <https://doi.org/10.1002/HYP.14508>,
906 2022.

907 McDermott, F.: Palaeo-climate reconstruction from stable isotope variations in speleothems: a review, *Quat Sci Rev*, 23, 901–
908 918, <https://doi.org/10.1016/J.QUASCIREV.2003.06.021>, 2004.

909 McDermott, F., Schwarcz, H., Rowe, P.J.: Isotopes in speleothems. In: Leng, M.J. (eds) *Isotopes in Palaeoenvironmental*
910 *Research. Developments in Palaeoenvironmental Research*, vol 10. Springer, Dordrecht. [https://doi.org/10.1007/1-4020-2504-](https://doi.org/10.1007/1-4020-2504-1_05)
911 [1_05](https://doi.org/10.1007/1-4020-2504-1_05), 2006.

McDonnell, J. J., McGuire, K., Aggarwal, P., Beven, K. J., Biondi, D., Destouni, G., Dunn, S., James, A., Kirchner, J., Kraft, P., Lyon, S., Maloszewski, P., Newman, B., Pfister, L., Rinaldo, A., Rodhe, A., Sayama, T., Seibert, J., Solomon, K., Soulsby, C., Stewart, M., Tetzlaff, D., Tobin, C., Troch, P., Weiler, M., Western, A., Wörman, A., and Wrede, S.: How old is streamwater? Open questions in catchment transit time conceptualization, modelling and analysis, *Hydrol Process*, 24, 1745–1754, <https://doi.org/10.1002/HYP.7796>, 2010.

Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., and Dolman, A. J.: Global land-surface evaporation estimated from satellite-based observations, *Hydrol Earth Syst Sci*, 15, 453–469, <https://doi.org/10.5194/hess-15-453-2011>, 2011.

Miralles, D. G., Bonte, O., Koppa, A., Baez-Villanueva, O. M., Tronquo, E., Zhong, F., Beck, H. E., Hulsman, P., Dorigo, W., Verhoest, N. E. C., and Haghdoust, S.: GLEAM4: global land evaporation and soil moisture dataset at 0.1° resolution from 1980 to near present, *Scientific Data* 12, 416, <https://doi.org/10.1038/s41597-025-04610-y>, 2025.

Nan, Y., He, Z., Tian, F., Wei, Z., and Tian, L.: Can we use precipitation isotope outputs of isotopic general circulation models to improve hydrological modeling in large mountainous catchments on the Tibetan Plateau? *Hydrol Earth Syst Sci*, 25, 6151–6172, <https://doi.org/10.5194/hess-25-6151-2021>, 2021.

Olarinoye, T., Gleeson, T., Marx, V., Seeger, S., Adinehvand, R., Allocca, V., Andreo, B., Apaéstegui, J., Apolit, C., Arfib, B., Auler, A., Barberá, J. A., Batiot-Guilhe, C., Bechtel, T., Binet, S., Bittner, D., Blatnik, M., Bolger, T., Brunet, P., Charlier, J. B., Chen, Z., Chiogna, G., Coxon, G., De Vita, P., Doummar, J., Epting, J., Fournier, M., Goldscheider, N., Gunn, J., Guo, F., Guyot, J. L., Howden, N., Huggenberger, P., Hunt, B., Jeannin, P. Y., Jiang, G., Jones, G., Jourde, H., Karmann, I., Koit, O., Kordilla, J., Labat, D., Ladouche, B., Liso, I. S., Liu, Z., Massei, N., Mazzilli, N., Mudarra, M., Parise, M., Pu, J., Ravbar, N., Sanchez, L. H., Santo, A., Sauter, M., Sivelle, V., Skoglund, R. Ø., Stevanovic, Z., Wood, C., Worthington, S., and Hartmann, A.: Global karst springs hydrograph dataset for research and management of the world’s fastest-flowing groundwater, *Sci Data*, 7, <https://doi.org/10.1038/s41597-019-0346-5>, 2020.

Pape, J. R., Banner, J. L., Mack, L. E., Musgrove, M. L., and Guilfoyle, A.: Controls on oxygen isotope variability in precipitation and cave drip waters, central Texas, USA, *J Hydrol (Amst)*, 385, 203–215, <https://doi.org/10.1016/J.JHYDROL.2010.02.021>, 2010.

Priestley, S. C., Treble, P. C., Griffiths, A. D., Baker, A., Abram, N. J., and Meredith, K. T.: Caves demonstrate decrease in rainfall recharge of southwest Australian groundwater is unprecedented for the last 800 years, *Commun Earth Environ*, 4, 206, <https://doi.org/10.1038/S43247-023-00858-7>, 2023.

Rusjan, S., Sapač, K., Petrič, M., Lojen, S., and Bezak, N.: Identifying the hydrological behavior of a complex karst system using stable isotopes, *J Hydrol (Amst)*, 577, <https://doi.org/10.1016/j.jhydrol.2019.123956>, 2019.

Staudinger, M., Seeger, S., Herbstritt, B., Stoelzle, M., Seibert, J., Stahl, K., and Weiler, M.: The CH-IRP data set: A decade of fortnightly data on $\delta^2\text{H}$ and $\delta^{18}\text{O}$ in streamflow and precipitation in Switzerland, *Earth Syst Sci Data*, 12, 3057–3066, <https://doi.org/10.5194/ESSD-12-3057-2020>, 2020.

945 Stevanović, Z.: Karst waters in potable water supply: a global scale overview, *Environ Earth Sci*, 78, 662,
 946 <https://doi.org/10.1007/s12665-019-8670-9>, 2019.

947 Treble, P. C., Baker, A., Abram, N. J., Hellstrom, J. C., Crawford, J., Gagan, M. K., Borsato, A., Griffiths, A. D., Bajo, P.,
 948 Markowska, M., Priestley, S. C., Hankin, S., and Paterson, D.: Ubiquitous karst hydrological control on speleothem oxygen
 949 isotope variability in a global study, *Communications Earth & Environment* 3, 29, [https://doi.org/10.1038/s43247-022-00347-](https://doi.org/10.1038/s43247-022-00347-3)
 950 3, 2022.

951 Treble, P. C., Zang, Y., Kaushal, N., Hatvani, I. G., Tanos, P., Kern, Z., Baker, A., Hartmann, A., Lechleitner, F. A., Belanger,
 952 B., Daudt de Oliveira, B., Sinclair, W., Riechelmann, D. F. C., Hidas, A., Yoshimura, K., Comas-Bru, L., and the SISAL
 953 working group members: SISAL_monv1: a global database of cave monitoring observations, *Earth Syst. Sci. Data Discuss.*
 954 [preprint], <https://doi.org/10.5194/essd-2026-31>, in review, 2026.

955 Tremaine, D. M., Froelich, P. N., and Wang, Y.: Speleothem calcite formed in situ: Modern calibration of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$
 956 paleoclimate proxies in a continuously-monitored natural cave system, *Geochim Cosmochim Acta*, 75, 4929–4950,
 957 <https://doi.org/10.1016/J.GCA.2011.06.005>, 2011.

958 Wang, Z., Su, Q., Wang, S., Gao, Z., and Liu, J.: Coupling hydrochemistry and stable isotopes ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) to identify the
 959 major factors affecting the hydrochemical evolution of groundwater in the Western Yellow Sea Coast, China, *Applied*
 960 *Geochemistry*, 138, 105221, <https://doi.org/10.1016/J.APGEOCHEM.2022.105221>, 2022.

961 Yoshimura, K., Kanamitsu, M., Noone, D., and Oki, T.: Historical isotope simulation using Reanalysis atmospheric data,
 962 *Journal of Geophysical Research Atmospheres*, 113, <https://doi.org/10.1029/2008JD010074>, 2008.

963 Zang, Y.: The WoKaS-Iso database: workflow for a global compilation of oxygen-18 and deuterium records in karst springs
 964 and cave drip water for enhanced understanding of karst systems, OPARA [data set], <https://doi.org/10.25532/OPARA-909>,
 965 2025.

966 Zhang, Z., Chen, X., Cheng, Q., and Soulsby, C.: Storage dynamics, hydrological connectivity and flux ages in a karst
 967 catchment: Conceptual modelling using stable isotopes, *Hydrol Earth Syst Sci*, 23, 51–71, [https://doi.org/10.5194/HESS-23-](https://doi.org/10.5194/HESS-23-51-2019)
 968 51-2019, 2019.

969 Zhong, F., Jiang, S., Van Dijk, A. I. J. M., Ren, L., Schellekens, J., and Miralles, D. G.: Revisiting large-scale interception
 970 patterns constrained by a synthesis of global experimental data, *Hydrol Earth Syst Sci*, 26, 5647–5667,
 971 <https://doi.org/10.5194/hess-26-5647-2022>, 2022.