

TundraFlux: A database of ecosystem respiration with biotic and abiotic metadata from Arctic and alpine tundra warming experiments

Sarah Schwieger^{1,2}, Jan Dietrich^{1,2}, Mats P. Björkman^{3,4}, Judith M. Sarneel², Bowen Li³, Joel White³,
5 Inge H.J. Althuizen^{5,6}, Christina Biasi^{7,8}, Robert G. Björk^{4,9}, Hanna Böhner¹⁰, Brage Bremset Hansen^{11,12},
Michele Carbognani¹³, Giorgio Chiari¹³, Casper T. Christiansen^{14,15}, Elisabeth J. Cooper¹⁶, Hans
Cornelissen¹⁷, Ludovica D'Imperio^{15,18}, Ellen Dorrepaal^{1,2}, Bo Elberling¹⁵, Patrick Faubert¹⁹, Ned
Fetcher²⁰, T'ai G. W. Forte¹³, Joseph Gaudard²¹, Konstantin Gavazov^{22,1}, Zhen-huan Guan²³, Jón
Guðmundsson²⁴, Siri V. Haugum²⁵, Jin-Sheng He^{26,23}, Caitlin Hicks Pries²⁷, Mark Hovenden^{28,29}, Simone
10 I. Lang³⁰, Gus Jespersen³¹, Ingibjörg S. Jónsdóttir³², Ji Young Jung³³, Olga Khitun³⁴, Birgitte Kortegaard
Danielsen¹⁵, Richard Lamprecht⁷, Mathilde Le Moullec^{35,36}, Hanna Lee³⁷, Maija E. Marushchak⁷, Anders
Michelsen¹⁴, Tariq Munir³⁸, Eero Myrsky^{39,40}, Kevin K. Newsham⁴¹, Marion Nyberg²⁸, Steven F.
Oberbauer^{42,43}, Paulo Olivas^{44,43}, Johan Olofsson², Hlynur Óskarsson²⁴, Thomas C. Parker⁴⁵, Matteo Petit
Bon⁴⁶, Alessandro Petraglia¹³, Emily Pickering Pedersen^{1,14}, Katrine Raundrup³⁵, Nynne R. Ravn¹⁴,
15 Riikka Rinnan⁴⁷, Heidi Rodenhizer^{48,49}, Ingvild Ryde^{14,15}, Alejandro Salazar²⁴, Niels M. Schmidt⁵⁰, Ted
Schuur^{48,51}, Sofie Sjögersten⁵², Cecilie Skov Nielsen^{53,15}, Sari Stark⁵⁴, Maria Strack⁵⁵, Jianwu Tang (纟)⁵⁶,
Sylvia Toet⁵⁷, Anne Tolvanen⁵⁸, Maria Väisänen⁵⁹, Richard Van Logtestijn¹⁷, Vigdis Vandvik²¹, Carolina
Voigt^{7,60,61}, Josefine Walz¹, Jeffrey M. Welker^{31,62,63}, Yuanhe Yang^{64,65}, Henni Yläne⁶⁶, Sybryn L.
Maes^{1,67}

20

Affiliations:

- ¹ Climate Impacts Research Centre, Department of Ecology and Environmental Science, Umeå University, Abisko, Sweden
² Department of Ecology, Environment and Geoscience, Umeå University, Umeå, Sweden
25 ³ Department of Biological and Environmental Sciences, University of Gothenburg, Box 463, SE-405 30 Gothenburg, Sweden
⁴ Gothenburg Global Biodiversity Centre, SE-405 30 Gothenburg, Sweden
⁵ NORCE Research AS, Bergen, Norway
⁶ Bjerknes Centre for Climate Research, University of Bergen, Norway
⁷ University of Eastern Finland, Department of Environmental and Biological Sciences, PO Box 1627, 70211 Kuopio, Finland
30 ⁸ University of Innsbruck, Department of Ecology, Sternwartestraße 15, 6020 Innsbruck, Austria
⁹ Department of Earth Sciences, University of Gothenburg, SE-405 30 Gothenburg, Sweden
¹⁰ Department of Arctic and Marine Biology, UiT – the Arctic University of Norway, N-9037 Tromsø, Norway
¹¹ Department of Terrestrial Ecology, Norwegian Institute for Nature Research, Trondheim, Norway
¹² Gjærevoll Centre for Biodiversity Foresight Analysis, Norwegian University of Science and Technology, Høgskoleringen
35 5, 7491 Trondheim, Norway
¹³ Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parma, Italy
¹⁴ Terrestrial Ecology Section, Department of Biology, University of Copenhagen, Denmark
¹⁵ Center for Permafrost, Department of Geosciences and Natural Resource Management, University of Copenhagen, Denmark
¹⁶ Department of Arctic and Marine Biology, UiT – the Arctic University of Norway, N-9037 Tromsø, Norway
40 ¹⁷ Amsterdam Institute for Life and Environment (A-LIFE), Vrije Universiteit, Amsterdam, The Netherlands
¹⁸ Department of Biotechnology and Biomedicine, Technical University of Denmark, Kgs. Lyngby

- ¹⁹ Carbone boréal, Département des sciences fondamentales, Université du Québec à Chicoutimi, 555 boulevard de l'Université, Chicoutimi, QC, G7H 2B1, Canada
- ²⁰ Institute for Environmental Science and Sustainability, Wilkes University, Wilkes-Barre, PA 18766
- 45 ²¹ Department of Biological Sciences and Bjerknes Centre for Climate Research, University of Bergen, Norway
- ²² Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Lausanne, Switzerland
- ²³ State Key Laboratory of Herbage Improvement and Grassland Agro-Ecosystems, and College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou 730000, China
- ²⁴ Agricultural University of Iceland, Árleyni 22, 112 Reykjavik, Iceland
- 50 ²⁵ The Heathland Centre, Alver, Norway
- ²⁶ Institute of Ecology, College of Urban and Environmental Sciences, Key Laboratory for Earth Surface Processes of the Ministry of Education, Peking University, Beijing 100871, China
- ²⁷ Department of Biological Sciences, Dartmouth College, Hanover, NH, USA 03755
- ²⁸ Biological Sciences, School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia
- 55 ²⁹ Australian Mountain Research Facility
- ³⁰ Department of Arctic Biology, The University Centre in Svalbard, PO Box 156, 9171 Longyearbyen, Norway
- ³¹ Department of Biological Sciences University of Alaska Anchorage
- ³² University of Iceland, Life and Environmental Sciences, Sturlugata 2, 102 Reykjavík, Iceland
- ³³ Division of Life Sciences, Korea Polar Research Institute, Incheon, South Korea
- 60 ³⁴ Department of Biological and Environmental Sciences, University of Gothenburg, Box 463, SE-405 30 Göteborg, Sweden
- ³⁵ Greenland Institute of Natural Resources, Kivioq 2, 3900 Nuuk, Greenland
- ³⁶ Department of Biology, Norwegian University of Science and Technology, Høgskoleringen 5, 7491 Trondheim, Norway
- ³⁷ Department of Biology, NTNU Norwegian University of Science and Technology, Trondheim, Norway
- ³⁸ Department of Geography, University of Calgary, Alberta, Canada
- 65 ³⁹ Arctic Centre, University of Lapland, Rovaniemi, Finland
- ⁴⁰ University of Helsinki, Finland
- ⁴¹ British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge, CB3 0ET, United Kingdom
- ⁴² Department of Biological Sciences, Florida International University, Miami, FL, USA
- ⁴³ Institute of Environment, Florida International University, Miami, FL, USA
- 70 ⁴⁴ Department of Earth and Environment, Florida International University, Miami, FL, USA
- ⁴⁵ Ecological Sciences, The James Hutton Institute, Aberdeen, Scotland, UK AB15 8QH
- ⁴⁶ Department of Applied Ecology | College of Agriculture and Life Sciences, North Carolina State University, NC-27695 Raleigh, North Carolina, USA
- ⁴⁷ Center for Volatile Interactions, Department of Biology, University of Copenhagen, Denmark
- 75 ⁴⁸ Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff AZ USA
- ⁴⁹ Woodwell Climate Research Center, Falmouth MA, USA
- ⁵⁰ Department of Ecoscience, Aarhus University, Roskilde, Denmark
- ⁵¹ Department of Biological Sciences, Northern Arizona University, Flagstaff AZ USA
- ⁵² School of Biosciences, University of Nottingham, Sutton Bonington Campus, LE12 5ND, Loughborough, UK
- 80 ⁵³ SEGES Innovation P/S, Agro Food Park 15, Aarhus, Denmark
- ⁵⁴ Department of Arctic Biology, University Centre in Svalbard, PO Box 156, Longyearbyen, Norway
- ⁵⁵ Department of Geography and Environmental Management, University of Waterloo, Waterloo, ON, Canada
- ⁵⁶ The Ecosystems Center, Marine Biological Laboratory, Woods Hole, Massachusetts
- ⁵⁷ Department of Biology, University of York, UK
- 85 ⁵⁸ Natural Resources Institute Finland
- ⁵⁹ Water, Energy and Environmental Engineering Research Unit, University of Oulu, Oulu, Finland
- ⁶⁰ Permafrost Research Section, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, 14473 Potsdam, Germany
- ⁶¹ University of Hamburg, Department of Earth System Sciences, Allende-Platz 2, 20146 Hamburg, Germany
- 90 ⁶² Ecology and Genetics Research Unit, University of Oulu Finland
- ⁶³ University of the Arctic

⁶⁴ State Key Laboratory of Forage Breeding-by-Design and Utilization

⁶⁵ Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, China

⁶⁶ School of Forest Sciences, University of Eastern Finland, Joensuu, Finland

95 ⁶⁷ Forest Ecology and Management Group (FORECOMAN), Department of Earth and Environmental Sciences, KU Leuven, Leuven, Belgium

✝ “deceased”

Correspondence to: Sarah Schwieger (sarah.schwieger@umu.se)

100

Abstract

Empirical *in-situ* measurements of ecosystem carbon dioxide respiration (R_{eco}) in high-latitude ecosystems remain limited, yet they are essential for understanding how tundra carbon cycling responds to climate warming across different environmental contexts and for reducing uncertainties in upscaled carbon budgets and carbon–climate feedbacks. Here, we present the TundraFlux Database, which to date is the most comprehensive synthesis of tundra R_{eco} responses to experimental warming. The database compiles over 24,000 daily-aggregated *in-situ* R_{eco} measurements from control and warmed plots with open-top chambers at 64 Arctic and alpine tundra sites across 12 countries. By coupling R_{eco} measurements with extensive metadata on climate, vegetation, and soil characteristics, the TundraFlux Database enables the integration of field-scale ecological processes into large-scale models, offering new opportunities to refine global carbon budgets and test predictions of tundra ecosystem responses to warming. Open access to the TundraFlux Database will empower the research community to better quantify and predict how warming alters carbon cycling in Arctic and alpine tundra ecosystems.

Keywords: Arctic tundra, alpine tundra, ecosystem respiration, open-top chambers (OTCs), warming experiments, database, carbon dioxide flux, vegetation data, soil data, climate data

1 Introduction

Ecosystem carbon dioxide (CO_2) respiration, the sum of autotrophic respiration from plants and heterotrophic respiration from soil organisms (i.e., fauna and microbes), constitutes the largest natural carbon flux from terrestrial ecosystems to the atmosphere (Jones et al., 1999; Lu et al., 2013; Oberbauer et al., 1998; Schuur et al., 2008; Tarnocai et al., 2009). The Arctic and alpine tundra biome stores one-third of global soil organic carbon, which is nearly twice the atmospheric carbon pool (Schuur et al., 2015), much of it locked in permafrost (i.e., soil that remains consecutively frozen for at least 2 years), organic-rich mineral soils, and peat (Gorham, 1991; Hugelius et al., 2020; Park et al., 2025; Schuur et al., 2022; Tarnocai et al., 2009; Zimov et al., 2006). As ecosystem CO_2 respiration (R_{eco}) is a temperature-sensitive process (Davidson et al., 2006; Gudas et al., 2021; Mahecha et al., 2010; Niu et al., 2024), understanding the consequences of the current rapid warming in Arctic and alpine tundra (Rantanen et al., 2022; Tingley and Huybers, 2013; Welker et al., 1999) is crucial for understanding climate-driven shifts in soil processes and global carbon cycling. Rising air and soil temperatures are expected to thaw permafrost, release previously stored soil organic carbon, and accelerate microbial decomposition of soil organic matter, thereby increasing CO_2 emissions to the atmosphere (Dorrepaal et al., 2009; Friggens et al., 2025; Karhu et al., 2014; Maes et al., 2024; Rustad et al., 2001; Schimel et al., 2004, 2006), which could significantly amplify global climate change (Cox et al., 2000; Welker et al., 2004).

Ecosystem respiration plays a central role in the global carbon cycle, making it essential to predict how R_{eco} responds to climate change. However, accurately forecasting the extent, as well as the spatial and temporal variability of these responses, remains

a major scientific challenge (Karhu et al., 2014; Maes et al., 2024; Rustad et al., 2001; Schuur et al., 2022; Sulman et al., 2018; Virkkala et al., 2021). Spatially, addressing these challenges requires moving beyond isolated case studies toward synthesising empirical data across diverse tundra sites and microclimates. Temporally, interannual variability is high and data from the non-growing season are sparse, even though respiration during this period can contribute substantially to annual carbon budgets (Fahnestock et al., 1998, 1999; Natali et al., 2019; Welker et al., 2000). Here, we present the TundraFlux Database, which compiles R_{eco} measurements derived from open-top chamber (OTC) warming experiments. These experiments use small greenhouses to passively increase air temperatures during the snow-free season while allowing relatively free entry of precipitation (Hollister et al., 2023; Marion et al., 1997; Welker et al., 1997). They are commonly used to simulate climate warming at a plot-scale in low-stature Arctic and alpine tundra systems, e.g., in the International Tundra Experiment network (ITEX; <https://www.gvsu.edu/itex/>; Henry and Molau, 1997; Hollister et al., 2023). These experiments provide a unique opportunity to analyse patterns and drivers of CO_2 respiration under warming conditions across bioclimatic gradients, which arise from differences in climate, vegetation, and soil characteristics (Maes et al., 2024). Experimental warming studies uniquely integrate multiple, interacting ecosystem responses, including vegetation dynamics, microbial activity, soil processes, and snow-mediated microclimate effects (Hollister et al., 2023; Leffler et al., 2016; Welker et al., 1997, 1999), all of which jointly regulate R_{eco} (Niu et al., 2024).

The TundraFlux Database currently includes 24,951 R_{eco} observations, aggregated to daily values (whenever multiple measurements were made within the same day) from experimentally warmed and associated control plots across 64 Arctic and alpine tundra sites (Fig. 1A), with measurements conducted between 2000 and 2024 (Fig. 1B). Here, we describe the data sources, the database structure and variables (Sect. 2), as well as potential applications (Sect. 3), data coverage and resolution (Sect. 4), future directions (Sect. 5), and availability (Sect. 6) of the TundraFlux Database.




Figure 1 A. Map showing the locations of the 64 open-top chamber experiments from which the ecosystem CO₂ respiration (R_{eco}) measurements in the TundraFlux Database were derived. Temporal spread of the data showing **B**, the distribution of the year of R_{eco} measurements, **C**, the duration of R_{eco} measurements (start of flux measurements - start of the experiment), and **D**, the distribution of day-of-year for R_{eco} measurements, with shaded regions highlighting winter (Dec–Mar, light blue), shoulder (Oct–Nov, Apr–May, grey), and growing (Jun–Sep, white) seasons. **E**. Boxplots showing the daily average R_{eco} ($\text{g C-CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) for each site, for both the unmanipulated control (i.e., ambient conditions, **top**) and the warmed treatment with open-top chambers (OTC, **bottom**). Boxes show the median and interquartile range (IQR, 25th–75th percentiles); whiskers extend to $1.5 \times \text{IQR}$, and values beyond the whiskers are plotted as outliers. Colours for Site ID are the same for all panels. The database includes occasional negative and exact-zero R_{eco} values. Negative values can result from instrument noise or brief net CO₂ uptake, while exact zeros may reflect contributor preprocessing (e.g., rounding or thresholding small fluxes).

2 Description and structure

2.1 Scope and purpose

Warming alters tundra ecosystems through a suite of interacting biotic and abiotic pathways, including changes in vegetation composition (Bjorkman et al., 2020; Collins et al., 2021; García Criado et al., 2025; Wilson and Nilsson, 2009), plant productivity (Hollesen et al., 2015; Myers-Smith et al., 2019), microbial activity (Frossard et al., 2021), decomposition rates (Sarnecki et al., 2020; Schwieger et al., 2025), nutrient cycling (Weedon et al., 2012), growing season length (Barichivich et al., 2013; Collins et al., 2021; Myers-Smith et al., 2019; Oberbauer et al., 1998), and snow-mediated microclimate conditions (Morgner et al., 2010; Pattison and Welker, 2014; Rixen et al., 2022), as well as increased thermokarst activity and permafrost degradation, which can strongly alter soil structure and carbon availability and thereby contribute to ecosystem respiration (R_{eco}) (Abbott and Jones, 2015; Lewkowicz and Way, 2019; Olefeldt et al., 2016; Vogel et al., 2009). These processes jointly regulate R_{eco} , making it an integrative indicator of tundra ecosystem responses to climate warming.

As these responses occur at multiple spatial and temporal scales and are significantly influenced by the local environmental context, a robust evaluation of the effects of warming on R_{eco} requires long-term, spatially diverse datasets that link flux measurements with detailed site and plot-level metadata. The TundraFlux Database was developed to address this issue, systematically compiling chamber-derived R_{eco} measurements from experimental warming studies in Arctic and alpine tundra ecosystems. Fin these efforts by specifically compiling R_{eco} observations from both ambient and experimentally manipulated tundra ecosystems, as well as including extensive metadata on vegetation (e.g., plant community composition, functional traits, biomass), soils (e.g., pH, organic carbon and nitrogen content, soil organic matter), and microclimate (e.g., air and soil temperature, soil moisture). This structure enables users to categorise analyses according to ecosystem properties, environmental factors or experimental design. This supports hypothesis testing, model evaluation and the large-scale synthesis of tundra carbon–climate feedback.

2.2 Data sources and data collection

The TundraFlux Database compiles *in-situ* measurements of daily-aggregated terrestrial ecosystem-level CO_2 fluxes ($g\ C-185\ CO_2\ m^{-2}\ day^{-1}$) from Arctic and alpine tundra ecosystems to assess warming effects on ecosystem CO_2 respiration (R_{eco}). We compiled data from experiments that used open-top chambers (OTCs) from across the Arctic and alpine tundra biome.

To achieve this, we contacted potential data contributors through established research networks like ITEX, WaRM, INTERACT and the Permafrost Carbon Network, and identified relevant contact information from authors of previously published meta-analyses on R_{eco} responses in warming experiments (Table S2).

190 We included data from experiments situated within the Arctic and alpine tundra biome (i.e., treeless regions beyond the climatic limit for tree growth) that reported *in-situ* measurements of ecosystem respiration (R_{eco}) in both warmed open-top chamber (OTC) and unmanipulated control plots. Based on these criteria, we compiled 40,160 individual R_{eco} measurements,

encompassing repeated observations across multiple plots, years, days, and, in some cases, multiple measurement times within a day at each site.

195 To maximize data usability and harmonize temporal resolution across studies, we provide two interlinked versions of the R_{eco} dataset. The raw dataset retains all individual measurements at their original temporal resolution, including multiple measurements per day and associated quality-control information (in total 40,160 observations), allowing users full flexibility in data filtering and aggregation. In addition, we provide a daily-aggregated dataset in which R_{eco} values were averaged within each site (`site_id`), treatment (OTC or control), plot (`plot_id`), year (`flux_year`), and day of year (`flux_doy`) when multiple
200 measurements occurred per day. This aggregation reduces short-term variability and preprocessing effort for synthesis applications, resulting in a total of 24,951 daily mean observations (OTC, $n = 11,046$; control, $n = 13,905$; Fig. 1E).

We define a *dataset* as all R_{eco} measurements from a given site and year (`site_id` $\times$ `flux_year`), including both OTC and control treatments, all replicate plots, and one or more measurement dates (Table S3). Plot-level data were retained by design to preserve within-site replication and enable flexible, user-defined aggregation and statistical modeling approaches, including
205 treatment-wise averaging or hierarchical analyses.

Alongside R_{eco} , we compiled *in-situ* metadata on microclimate conditions, geolocation, vegetation cover, plant traits, and soil characteristics. These ancillary variables were measured at both plot and site levels, with some recorded concurrently with flux measurements (and provided within the Carbon Flux data, Fig. 2), and others representing broader site-level environmental context that can be linked to the relevant flux measurements (Fig. 2, Fig. 3, Table S4).

210 Part of the database compiled through this effort was previously presented in Maes et al. (2024), which focused exclusively on growing season measurements (i.e., June to August) and included data up to the year 2020. This current published version of the TundraFlux Database extends the earlier synthesis by incorporating updated measurements through 2024 (Fig. 1B) and by including shoulder season fluxes from April to May and September to November (Fig. 1D).

2.3 Data structure and variables

215 The TundraFlux Database comprises 76 variables describing R_{eco} from 64 tundra sites spanning 2000–2024 (Fig. 1A, B). A comprehensive data dictionary with a description of the variables of the site, vegetation, soil and method metadata, including data-type and unit is provided in the Supplements (Table S4).

The TundraFlux Database			
1. Carbon Flux Data (plot-level)		2. Metadata	
site_id flux_date flux_year flux_doy reco treatment plot_id par air_temperature soil_temperature soil_moisture water_table		2a. Site Metadata (site-level)	2b. Vegetation Cover & Traits (plot-level)
		site_name country_code latitude longitude altitude mean_annual_temperature mean_annual_precipitation active_layer_depth parent_material vegetation_type PI_contact add_contact	vascular_plant_cover deciduous_shrubs_cover evergreen_shrubs_cover forbs_cover graminoids_cover lichen_cover mosses_cover vegetation_method plant_height plant_height_method biomass biomass_method
3. Method Metadata (site-level)		2c. Soil Properties (plot-level)	
measurement_method machine machine_type flux_measurement_start_year warming_start_year plot_size flux_plot_size otc_height chamber_temperature		organic_layer_depth bulk_density_organic bulk_density_mineral bulk_density_mixture c_n_organic c_n_mineral soc_organic soc_mineral soc_mixture son_organic son_mineral son_mixture som_organic som_mineral ph_h2o_organic ph_h2o_mineral ph_h2o_mixture soil_moisture_plot soil_moisture_plot_type	

Figure 2. Data structure of the TundraFlux Database. Boxes represent the main data types: (1) Carbon flux data, including plot-level daily averages of ecosystem CO₂ respiration (R_{eco}) in g C-CO₂ m⁻² d⁻¹ and data measured along with flux measurements; (2) Metadata, including information on (a) site, (b) vegetation cover and plant traits at plot-level, and (c) Soil properties at plot level; and (3) Method data including documentation of experimental setups. Variables listed inside the boxes correspond to column names in the database (identical to Table S4).




Figure 3. Percentage of available observations for metadata variables presented in Figure 2 relevant for CO₂ modelling. Soil organic carbon (SOC), soil organic nitrogen (SON), carbon-to-nitrogen ratio (c_n), soil organic matter (SOM).

2.4 Data standardization and quality control

Data were processed in R version 4.5.1 (R Core Team 2025). We used the R package TIDYVERSE (v. 2.0.0) for data handling. For each dataset, defined by a unique combination of site_id and flux_year, we calculated treatment- and plot-specific daily mean R_{eco} values, whenever multiple measurements were made within the same day. We provide an R script on Zenodo (DOI: 10.5281/zenodo.17976235) that documents the full aggregation procedure, and the un-aggregated dataset (see the ‘Data

Availability' section). All individual flux measurements were standardized to a common unit ($\text{g C-CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) to ensure comparability across studies. We screened the data for inconsistencies in unit or sign conventions, contacting data contributors in cases of uncertainty. Spatial coordinates were validated by visualizing all sites on a map, and any imprecise or conflicting locations were corrected in consultation with site researchers.

235 All experiments measured daytime R_{eco} using dark or opaque chambers, except for the CiPEHR site in Alaska (ALA_1), where automated chambers continuously measured CO_2 respiration using clear chambers. To obtain comparable respiration estimates for this site, we extracted only night-time fluxes, defined by photosynthetically active radiation (PAR) values below $5 \mu\text{mol m}^{-2} \text{ s}^{-1}$, as these best approximated dark chamber conditions. These night-time measurements were therefore used as the site's R_{eco} values.

240 The same quality-control procedures were applied to associated metadata to ensure consistency in spatial reference and measurement units across sites. Soil moisture data, reported at the plot level, were provided either as volumetric or gravimetric values and were subsequently converted to percentages. Bulk density was standardized to g dwt cm^{-3} . Soil organic carbon (SOC) and soil organic nitrogen (SON) were standardized to g C kg^{-1} soil and g N kg^{-1} soil, respectively, while soil organic matter (SOM) was standardized to percentages.

245 The TundraFlux Database includes occasional negative and exact-zero R_{eco} values (Table S3). These originate from the heterogeneity of measurement techniques and pre-processing procedures used in the studies that contribute to the database. Negative fluxes can occur in chamber-based measurements due to short-term CO_2 concentration fluctuations, instrument noise, pressure-induced artefacts or brief periods of apparent net CO_2 uptake when R_{eco} is close to the detection limit. We retained such values exactly as reported in the original datasets in order to preserve data fidelity. Exact-zero values appear when the
250 net change in chamber CO_2 concentration falls within instrument noise, or when contributors apply local preprocessing (e.g. rounding small negative fluxes to zero or applying minimum detection thresholds) prior to submission. As preprocessing conventions differed between studies, both negative and zero values are present in the compiled dataset. Users may apply additional filtering or thresholding, in line with their research objectives, for instance by removing negative fluxes, setting minimum detection limits or utilizing the quality control flags provided in the non-aggregated dataset.

255 **2.4.1 Outlier detection and handling**

We chose not to filter outliers from the database before calculating daily averages to preserve the original data structure. However, recognizing the substantial variability in carbon fluxes across sites (Fig. 1E) and years (Table S3), we developed a tailored outlier detection and flagging procedure for R_{eco} data. This method applies the Median Absolute Deviation (MAD, Leys et al., 2013) to the non-aggregated R_{eco} values. This resulted in 285 flagged outliers (0.71% of the original 40,160 data
260 points) and the user should consider whether to remove them or to use other criteria for outlier detection. We added a column to the non-aggregated dataset that indicates the flagged outliers. A full description of this method is provided in the

Supplements. The R scripts used for data processing are archived on Zenodo (DOI: 10.5281/zenodo.17976235, Schwieger et al., 2026) and are developed and maintained in a public GitHub repository.

3 Applications of the TundraFlux database

265 3.1 Identifying extent and drivers of ecosystem respiration response to warming through meta-analysis

The TundraFlux Database is based on a global synthesis of how ecosystem respiration in Arctic and alpine tundra responds to experimental warming (Maes et al., 2024). Using 136 datasets from 56 OTC experiments at 28 tundra sites worldwide (60% of the data presented here), the study quantified that a mean rise of 1.4 °C [confidence interval (CI) 0.9–2.0 °C] in air and 0.4 °C [CI 0.2–0.7 °C] in soil temperatures causes a 30% [CI 22–38%] increase in growing season R_{eco} . It further showed that this stimulation persisted for up to 25 years of warming treatment, with evidence pointing to enhanced plant and microbial activity as the underlying drivers. By linking respiration responses to changes in local soil conditions (i.e., nitrogen concentration and carbon-to-nitrogen ratio), Maes et al. (2024) demonstrated that tundra sites with stronger nitrogen limitations and sites in which warming had stimulated plant and microbial nutrient turnover seemed particularly sensitive in their R_{eco} response to warming. This example highlights the power of standardized, long-term experimental data to uncover generalizable patterns in climate responses. It also showcases how the TundraFlux Database enables large-scale syntheses that identify not only the direction but also the mechanism of possible climate change feedbacks on tundra ecosystems, which are essential to improve carbon–climate feedback projections in Earth System Models.

The expanded TundraFlux Database, which now includes spring and autumn (‘shoulder-season’) measurements and updated observations through 2024, enables new high-impact research questions that could not be addressed in Maes et al. (2024). For example, users can quantify how experimental warming affects R_{eco} beyond the peak growing season and evaluate whether the environmental controls of R_{eco} differ between early-season, peak-season, and late-season periods. These additions substantially improve the capacity to analyze seasonal dynamics and long-term trajectories of tundra carbon cycling, thereby supporting more robust evaluations of climate–carbon feedbacks.

3.2 Carbon model parameterization and validation

285 Accurately predicting carbon release from permafrost soils under warming scenarios remains a major challenge in climate science (Knoblauch et al., 2021; Schuur et al., 2015) due to the high variability in ecosystem responses and limited availability of long-term data (Swindles et al., 2015). The Warming Permafrost Model Intercomparison Project (WrPMIP) led by Woodwell Climate Research Center is using the TundraFlux Database to bridge the gap between experimental warming studies and large-scale carbon modeling (warmingpermafrost.nau.edu). In this project, multi-model simulations are being run at both regional and site scales to align with the spatial and temporal dimensions of experimental data (Wells et al., 2023). By aligning field-based warming measurements from the TundraFlux Database with model simulations, the project will enhance our ability to project the magnitude and timing of carbon release from permafrost regions under climate change (Schädel et al., 2018).

4 Data coverage and resolution

As with any large database, not all biotic and abiotic variables, seasons, plant functional types, habitats, regions, or bioclimatic settings within the Arctic and alpine biomes are equally represented in the TundraFlux Database. Here, we identify some limitations related to gaps in the spatial and temporal resolution the database, which highlight clear priorities for future data collection efforts.

4.1 Spatial coverage and bias

The TundraFlux Database inherently reflects a sampling imbalance in the field, particularly the underrepresentation of key Arctic regions such as the Canadian High Arctic archipelago and Siberia (López-Blanco et al., 2024; Metcalfe et al., 2018; Virkkala et al., 2019; Fig. 1A, Table S5). Still it represents the most comprehensive effort that is currently available of R_{eco} data from warming experiments in the tundra. In particular, high-latitude North America and northern Europe are the best-represented regions (Table 1). This geographical concentration reflects a well-documented spatial bias in Arctic field sampling toward long-established research hubs with good accessibility and infrastructure (Metcalfe et al., 2018).

There is substantial variability in the number of measurements across sites (median = 120, IQR = 133.5; total measurements = 24,951). In particular, the CiPEHR site (ALA_1) near Eight Mile Lake, Alaska, USA, with 13,572 daily-aggregated observations, contributed 52.5% of the R_{eco} data in the TundraFlux Database (Table S5). A large proportion of observations from a single, well-studied site increases temporal and treatment-level detail but may disproportionately influence cross-site or pan-Arctic analyses if not accounted for. How to address this imbalance depends on the user's research question and analytical framework, but several approaches can help prevent disproportionate influence of high-density sites. These include applying hierarchical or mixed-effects models that treat site as a random effect, using equal or inverse-effort site weighting, aggregating fluxes to common temporal resolutions, conducting sensitivity analyses (e.g., subsampling or excluding dominant sites), or using site-level bootstrapping or partial-pooling approaches (Choi and Kang, 2025). At the same time, these long-term, high-density datasets provide valuable opportunities for method development, uncertainty quantification, and benchmarking models at sites with robust metadata.

By systematically compiling data from OTC experiments across Alaska, Greenland, Svalbard, Iceland, Fennoscandia, Canada, and Russia, the TundraFlux Database is the most comprehensive resource currently available for evaluating warming effects on tundra carbon cycling. Importantly, tundra ecosystems exhibit substantial spatial variability in both abiotic and biotic conditions (Aalto et al., 2022; Magnússon et al., 2023), meaning that each site, regardless of data quantity, contributes distinct information about ecosystem responses under different environmental and vegetation contexts.

Table 1. Number and percentages of daily R_{eco} observations by region. See Table S5 for classification of sites into regions and distribution of individual sites across the regions.

Region	Daily_obs	%	Nb of sites
High-latitude North American	15,906	61.6	13
Oceanic	3,083	11.9	8
High-latitude European	4,682	18.1	30
Alpine European	479	1.9	5
High-latitude Asian	542	2.1	3
Alpine Asian	1,136	4.4	4

325 4.2 Longer-term data on R_{eco} response to experimental warming

In the TundraFlux Database, 93% of the averaged daily data points ($n = 22,534$) come from warming experiments that lasted 11 years or less, and over half (51%, ($n = 12,239$)) lasted fewer than 4 years, at the time of R_{eco} measurements. Out of the 64 sites in total, 7 sites (11.1%) represent long-term experiments with continuous measurements (≥ 5 years), while 56 sites (88.9%) represent short-term experiments (< 5 years) with often only single measurements (Table S5). Consequently, longer-term measurements (> 11 -24 years) are rare ($n = 1,573$), and data from warming experiments that lasted longer than 24 years are absent (Fig. 1C). This clearly highlights the importance of maintaining long-term experiments, particularly given that changes in soil processes or vegetation composition driven by warming may unfold over decades in Arctic and alpine tundra (Hollister et al., 2005; Jónsdóttir et al., 2023; Wei et al., 2025). At the same time, both OTCs and manual chamber measurements may introduce experimental disturbance effects (e.g., trampling, vegetation damage, soil compaction) that can accumulate over time (Hollister et al., 2023). Such disturbance may increasingly influence ecosystem functioning the longer an experiment runs. Thus, while longer-term data would help capture slow ecological changes, extended experiment duration may simultaneously amplify disturbance-related artefacts, complicating the interpretation of long-term trends.

4.3 Shoulder and winter season respiration

Our database contains 21,830 daily-aggregated R_{eco} observations from the growing season (defined here as June-September in the Northern Hemisphere, and December-March in the Southern Hemisphere), and 3,297 observations from outside the growing season, highlighting a gap in our understanding of carbon fluxes in the shoulder (i.e., April, May, October, and November, $n = 3,330$) and winter seasons (i.e., December-March in the Northern Hemisphere, $n=0$, Australia excluded) (Fig. 1D, Fig. 4).

As Arctic soils are usually covered in snow during the winter months, the spring and autumn shoulder seasons are particularly vulnerable to the effects of global warming, as these transitional periods experience rapid changes in snow cover and soil temperatures (Hassol, 2004; Shukla et al., 2019). Despite its length and significance, winter remains the most understudied season in Arctic ecosystems, even though cold-season processes can contribute substantially to annual carbon budgets through continued microbial activity and ecosystem respiration beneath the snowpack (Natali et al., 2019; Zona et al., 2016). Climate warming is altering Arctic winters by increasing air and soil temperatures, changing snow accumulation and duration, and

350 increasing the frequency of winter warming events and rain-on-snow events, all of which affect soil thermal conditions, microbial activity, and carbon fluxes (Cooper, 2014; Rixen et al., 2022). Similarly, alpine ecosystems are experiencing reduced snowpack, earlier snowmelt, and glacier retreat, with important consequences for soil temperature regimes, hydrology, and ecosystem functioning (Ernakovich et al., 2014). Warming effects and mechanisms identified during the growing season (Maes et al., 2024) may not apply to the winter season, when factors such as snow depth and duration exert great control over carbon
355 fluxes (Björkman et al., 2010; Grogan, 2012; Morgner et al., 2010; Rixen et al., 2022; Semenchuk et al., 2016; Slatyer et al., 2022).

More research focusing on the effects of warming on carbon fluxes during the underrepresented winter and shoulder seasons while developing a mechanistic understanding of winter carbon dynamics is therefore essential to improve predictions of future CO₂ emissions from Arctic soils. Overlooking these seasons risks underestimating both the extent and variability of carbon
360 release from rapidly warming Arctic soils.




Figure 4. Monthly R_{eco} availability aggregated across all years. White/ light grey/ light blue cells = 0 observations. Color intensity based on $\text{Log}_{10}(\text{Number of Observations} + 1)$ due to strong bias of ALA_1. Growing season = June–September in the Northern Hemisphere, December – March in the Southern Hemisphere. Except for AUS_1, shaded areas indicate winter months (light blue) and shoulder months (light grey).

4.4 Partitioning data

R_{eco} responses to climate change depend on the dynamics of its two main components: autotrophic (plant-derived) and heterotrophic (microbial and faunal) respiration (Bond-Lamberty et al., 2004; Hicks Pries et al., 2013). Because these two components can respond differently to changes in climate (Borken et al., 2006; Muhr & Borken, 2009; Gomez-Casanovas et al., 2012), understanding their individual contributions is crucial to accurately predict ecosystem carbon dynamics and modeling carbon–climate feedback processes. However, partitioning R_{eco} into these source fluxes remains methodologically challenging. As respiration partitioning measurements are usually destructive, altering ecosystem dynamics and making long-term measurements difficult, such data remain scarce. For example, in the meta-analysis on the drivers of R_{eco} in Arctic and alpine tundra by Maes et al. (2024), only nine out of 136 R_{eco} datasets included partitioning data on the autotrophic and heterotrophic respiration components.

5 Future directions

In the coming years, our aim is to expand the scope of the TundraFlux Database beyond R_{eco} by incorporating methane and nitrous oxide fluxes, net ecosystem exchange (NEE), and gross primary productivity (GPP). We also plan to expand the scope of warming manipulations represented in the database by including additional climate change treatments, such as the use of snow fences to manipulate snow depth. This will enable us to distinguish between the effects of summer (OTC) and winter (snow fences) warming on R_{eco} , as well as the combined effects of these treatments (Hermesdorf et al., 2024). It will also allow us to account for cross-seasonal carry-over and the effects of coupled air-soil warming, whereby winter soil warming persists into summer and summer air warming can indirectly modify winter soil conditions via changes to vegetation (Kropp et al., 2020).

In recent years, remote sensing using unmanned aerial vehicles (UAVs) and satellites has advanced significantly, providing an opportunity to quantify landscape heterogeneity in the tundra biome at high spatio-temporal resolutions (Assmann et al., 2020; Myers-Smith et al., 2020). With this data, it is possible to bridge the gap between the TundraFlux plot-scale field measurements and large-scale remote sensing mapping products. In addition, we will incorporate remote sensing-derived metadata, such as tundra type and landform classification (e.g., peatlands, thaw slumps and coastal systems), as well as other derived products (Niittynen, 2026; Virkkala et al., 2024; Wagner and Hugelius, 2026), into future syntheses. Finally, we aim to link the TundraFlux Database with other available databases on terrestrial carbon fluxes (Table S1), including the Tundra Trait Team database (Bjorkman et al., 2018), the Manipulation Experiments Synthesis Initiative (MESI; Van Sundert et al., 2023), COSORE (Bond-Lamberty et al., 2020), and the ABCflux database (Leffler et al., 2025; Virkkala et al., 2022). Although

integration remains challenging due to differences in data formats, identifiers, and metadata standards, establishing common
395 protocols will be crucial to advance syntheses across databases.
To follow updates on ongoing and future projects related to our Tundra flux Database, please visit our website
<https://arcticflux.org/>. To contribute new datasets, please contact us via our mail tundrafluxdatabase@lists.umu.se.

6 Conclusions

The TundraFlux Database provides the most comprehensive synthesis of tundra R_{eco} responses to experimental warming,
400 integrating over 40,000 individual *in-situ* R_{eco} observations into 24,951 daily-aggregated R_{eco} measurements from open-top
chamber and control plots across 64 Arctic and alpine sites. By combining these data with extensive environmental metadata,
the database enables cross-scale analyses that link ecological processes to global carbon modeling. Although long-term (>24
years) data and measurements from outside the growing season remain limited, the TundraFlux Database establishes a
foundation for coordinated synthesis and future expansion to include methane fluxes, NEE, and GPP. When linked with other
405 ecological datasets, it will contribute to forming an unprecedented platform for cross-network analyses of Arctic and alpine
carbon dynamics.

Data availability

The TundraFlux Database is organized as a set of interlinked R data files (.rds) that can be merged using shared identifiers
such as site_id, flux_year, and, for plot-level data, plot_id. For users who prefer a modular workflow, separate metadata files
410 are available, including site (site_metadata_v1.rds), vegetation (plant_metadata_v1.rds), soil properties
(soil_metadata_plot_v1.rds), and methodological details (method_metadata_site_v1.rds). These metadata files can be linked
to R_{eco} -specific datasets, such as Reco_microclimate_daily_v1.rds, using the shared identifiers mentioned above.
For users who want a ready-to-use dataset, two fully integrated data files are provided: TundraFlux_daily_v1.rds, which
contains daily aggregated Reco measurements along with site, vegetation, soil, and methodological metadata (Table S4), and
415 TundraFlux_raw_v1.rds, which contains non-aggregated individual R_{eco} measurements with quality-control flags.
Missing values are consistently represented as NA across all files.
All data can be provided upon request from the corresponding author and will be made publicly available on Zenodo (DOI:
10.5281/zenodo.17976235, Schwieger et al., 2026) after acceptance of the paper.

Code availability

420 The code associated with this publication will be made publicly available on Zenodo (DOI: 10.5281/zenodo.17976235, Schwieger et al., 2026). The R scripts used for data processing are archived alongside the data and are also maintained in a public GitHub repository and can be provided upon request from the corresponding author.

Supplement link

Author contribution

425 The TundraFlux database was conceptualized by JD, JS, ED, MB, and SLM. JD and SLM compiled the data in 2020. JD and SS updated the data in 2023 and 2025. Data screening and curation by JD, SLM and SS. SS drafted and coordinated the manuscript in close collaboration with SLM, JD, JS, BL, JW and MB. SS prepared the code and data files for the repository, revised by BL and JD. All authors contributed to the realization of the TundraFlux Database and participated in reviewing and editing of the manuscript.

430 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to acknowledge the numerous students and field assistants involved in the collection of ecosystem respiration and connected metadata measurements. In particular, we would like to thank Minna Männistö and Elina Kaarlejärvi. We would
435 also like to thank the two anonymous reviewers for their helpful and constructive comments. We would also like to thank the station managers for their assistance and for granting us access to the field sites. We would also like to thank Jérémy Monsimet for his support in setting up our repository. Furthermore, we acknowledge the posthumous contribution of Jianwu Tang, whose work inspired the collection of the datasets ALA_2 and ALA_3.

Recognising the importance of Indigenous lands, we acknowledge that parts of our fieldwork were conducted on territories
440 historically and presently belonging to Indigenous peoples. We express our respect and gratitude to these communities. Special thanks are extended to the residents of Utqiagvik and Atkasuk, Alaska, for their cooperation and understanding during research activities in the Arctic region. This research would not have been possible without the collective efforts and support of these individuals and communities.

Financial support

445 This work was supported by the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS; grants 2016-01187, 2018-04202, 2021-02449, 2022-00786, 2023-04048, and 2024-00244); Biodiversity and Ecosystem Services in a Changing Climate (BECC); the Swiss National Science Foundation (grant PZ00P2_174047); the Australian Research Council (DP220100915); the U.S. Department of Energy (grants DE-SC0006982, DE-SC0014085, and DE-SC0020227); Rannís – The Icelandic Centre for Research (grants 70255021 and 1931333); Danmarks Grundforskningsfond (DNR100, DNR168, and VOLT); the National Research Foundation of Korea (RS-2025-24683148 and KOPRI-PN26010); the National Natural Science Foundation of China (32201358); the Research Council of Norway (grants 223257, 250740, 269957, 274712, 276080, and 294948); the Research Council of Finland (grants 332196, 337550, 341348, 218121, and 130507); Fonds Wetenschappelijk Onderzoek (12ZZV21N); the U.S. National Science Foundation (grants 0632184, 0856728, 0119279, 09321730, 09617643, 1417763, 1418010, 1504381, 1931333, and 2113641); the 455 Svalbard Environmental Protection Fund (15/128); the Finnish Cultural Foundation (grants 00170598 and 00180628); the European Commission Directorate-General for Research (EVK2-2001-00235); the European Research Council Starting Grant (101163177); the European Union’s Horizon 2020 research and innovation programme (grants 657627, 819202, and 771012); the European Union Seventh Framework Programme (FP7); the DG XII of the European Commission as part of the 'Environment and Climate' programme of the '4th Framework (ENV4-CT97-0586); the Oskari Huttunen Foundation; the 460 Academy of Finland; Alberta Innovates – Technology Futures and the Canada Research Chairs Program; Greenland Ecosystem Monitoring; and the Danish Environmental Protection Agency.

References

- Aalto, J., Tyystjärvi, V., Niittynen, P., Kemppinen, J., Rissanen, T., Gregow, H., and Luoto, M.: Microclimate temperature variations from boreal forests to the tundra, *Agric. For. Meteorol.*, 323, 109037, <https://doi.org/10.1016/j.agrformet.2022.109037>, 2022.
- 465 Abbott, B. W. and Jones, J. B.: Permafrost collapse alters soil carbon stocks, respiration, CH₄, and N₂O in upland tundra, *Glob. Change Biol.*, 21, 4570–4587, <https://doi.org/10.1111/gcb.13069>, 2015.
- Assmann, J. J., Myers-Smith, I. H., Kerby, J. T., Cunliffe, A. M., and Daskalova, G. N.: Drone data reveal heterogeneity in tundra greenness and phenology not captured by satellites, *Environ. Res. Lett.*, 15, 125002, <https://doi.org/10.1088/1748-9326/abbf7d>, 2020.
- 470 Barichivich, J., Briffa, K. R., Myneni, R. B., Osborn, T. J., Melvin, T. M., Ciais, P., Piao, S., and Tucker, C.: Large-scale variations in the vegetation growing season and annual cycle of atmospheric CO₂ at high northern latitudes from 1950 to 2011, *Glob. Change Biol.*, 19, 3167–3183, <https://doi.org/10.1111/gcb.12283>, 2013.
- Bjorkman, A., Criado, M., Myers-Smith, I., Ravolainen, V., Jonsdottir, I., Westergaard, K., Lawler, J., Aronsson, M., Bennett, B., Gardfjell, H., Heidmarsson, S., Stewart, L., and Normand, S.: Status and trends in Arctic vegetation: Evidence from experimental warming and long-term monitoring, *AMBIO*, 49, 678–692, <https://doi.org/10.1007/s13280-019-01161-6>, 2020.

- Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Thomas, H. J. D., Alatalo, J. M., Alexander, H., Anadon-Rosell, A., Angers-Blondin, S., Bai, Y., Baruah, G., te Beest, M., Berner, L., Björk, R. G., Blok, D., Bruelheide, H., Buchwal, A., Buras, A., Carbognani, M., Christie, K., Collier, L. S., Cooper, E. J., Cornelissen, J. H. C., Dickinson, K. J. M., Dullinger, S., Elberling, B., Eskelinen, A., Forbes, B. C., Frei, E. R., Iturrate-Garcia, M., Good, M. K., Grau, O., Green, P., Greve, M., Grogan, P., Haider, S., Hájek, T., Hallinger, M., Happonen, K., Harper, K. A., Heijmans, M. M. P. D., Henry, G. H. R., Hermanutz, L., Hewitt, R. E., Hollister, R. D., Hudson, J., Hülber, K., Iversen, C. M., Jaroszynska, F., Jiménez-Alfaro, B., Johnstone, J., Jorgensen, R. H., Kaarlejärvi, E., Klady, R., Klimešová, J., Korsten, A., Kuleza, S., Kulonen, A., Lamarque, L. J., Lantz, T., Laval, A., Lembrechts, J. J., Lévesque, E., Little, C. J., Luoto, M., Macek, P., Mack, M. C., Mathakutha, R., Michelsen, A., Milbau, A., Molau, U., Morgan, J. W., Mörsdorf, M. A., Nabe-Nielsen, J., Nielsen, S. S., Ninot, J. M., Oberbauer, S. F., Olofsson, J., Onipchenko, V. G., Petraglia, A., Pickering, C., Prevéy, J. S., Rixen, C., Rumpf, S. B., Schaepman-Strub, G., Semenchuk, P., Shetti, R., Soudzilovskaia, N. A., Spasojevic, M. J., Speed, J. D. M., Street, L. E., Suding, K., Tape, K. D., Tomaselli, M., Trant, A., Treier, U. A., Tremblay, J.-P., Tremblay, M., Venn, S., et al.: Tundra Trait Team: A database of plant traits spanning the tundra biome, *Glob. Ecol. Biogeogr.*, 27, 1402–1411, <https://doi.org/10.1111/geb.12821>, 2018.
- Björkman, M. P., Morgner, E., Björk, R. G., Cooper, E. J., Elberling, B., and Klemetsson, L.: A comparison of annual and seasonal carbon dioxide effluxes between sub-Arctic Sweden and High-Arctic Svalbard, *Polar Res.*, 29, 75–84, <https://doi.org/10.3402/polar.v29i1.6051>, 2010.
- Bond-Lamberty, B., Wang, C., and Gower, S. T.: A global relationship between the heterotrophic and autotrophic components of soil respiration?, *Glob. Change Biol.*, 10, 1756–1766, <https://doi.org/10.1111/j.1365-2486.2004.00816.x>, 2004.
- Bond-Lamberty, B., Christianson, D. S., Malhotra, A., Pennington, S. C., Sihi, D., AghaKouchak, A., Anjileli, H., Altaf Arain, M., Armesto, J. J., Ashraf, S., Ataka, M., Baldocchi, D., Andrew Black, T., Buchmann, N., Carbone, M. S., Chang, S.-C., Crill, P., Curtis, P. S., Davidson, E. A., Desai, A. R., Drake, J. E., El-Madany, T. S., Gavazzi, M., Görres, C.-M., Gough, C. M., Goulden, M., Gregg, J., Gutiérrez del Arroyo, O., He, J.-S., Hirano, T., Hoppo, A., Hughes, H., Järveoja, J., Jassal, R., Jian, J., Kan, H., Kaye, J., Kominami, Y., Liang, N., Lipson, D., Macdonald, C. A., Maseyk, K., Mathes, K., Mauritz, M., Mayes, M. A., McNulty, S., Miao, G., Migliavacca, M., Miller, S., Miniati, C. F., Nietz, J. G., Nilsson, M. B., Noormets, A., Norouzi, H., O’Connell, C. S., Osborne, B., Oyónarte, C., Pang, Z., Peichl, M., Pendall, E., Perez-Quezada, J. F., Phillips, C. L., Phillips, R. P., Raich, J. W., Renchon, A. A., Ruehr, N. K., Sánchez-Cañete, E. P., Saunders, M., Savage, K. E., Schrumpp, M., Scott, R. L., Seibt, U., Silver, W. L., Sun, W., Szutu, D., Takagi, K., Takagi, M., Teramoto, M., Tjoelker, M. G., Trumbore, S., Ueyama, M., Vargas, R., Varner, R. K., Verfaillie, J., Vogel, C., Wang, J., Winston, G., Wood, T. E., Wu, J., Wutzler, T., Zeng, J., Zha, T., Zhang, Q., and Zou, J.: COSORE: A community database for continuous soil respiration and other soil-atmosphere greenhouse gas flux data, *Glob. Change Biol.*, 26, 7268–7283, <https://doi.org/10.1111/gcb.15353>, 2020.
- Choi, G. J. and Kang, H.: Heterogeneity in meta-analyses: an unavoidable challenge worth exploring, *Korean J. Anesthesiol.*, 78, 301–314, <https://doi.org/10.4097/kja.25001>, 2025.
- Collins, C. G., Elmendorf, S. C., Hollister, R. D., Henry, G. H. R., Clark, K., Bjorkman, A. D., Myers-Smith, I. H., Prevéy, J. S., Ashton, I. W., Assmann, J. J., Alatalo, J. M., Carbognani, M., Chisholm, C., Cooper, E. J., Forrester, C., Jónsdóttir, I. S., Klanderud, K., Kopp, C. W., Livensperger, C., Mauritz, M., May, J. L., Molau, U., Oberbauer, S. F., Ogburn, E., Panchen, Z. A., Petraglia, A., Post, E., Rixen, C., Rodenhizer, H., Schuur, E. A. G., Semenchuk, P., Smith, J. G., Steltzer, H., Totland, Ø., Walker, M. D., Welker, J. M., and Suding, K. N.: Experimental warming differentially affects vegetative and reproductive phenology of tundra plants, *Nat. Commun.*, 12, 3442, <https://doi.org/10.1038/s41467-021-23841-2>, 2021.
- Cooper, E. J.: Warmer Shorter Winters Disrupt Arctic Terrestrial Ecosystems, *Annu. Rev. Ecol. Evol. Syst.*, 45, 271–295, <https://doi.org/10.1146/annurev-ecolsys-120213-091620>, 2014.
- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., and Totterdell, I. J.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 184–187, <https://doi.org/10.1038/35041539>, 2000.

- Davidson, E. A., Janssens, I. A., and Luo, Y.: On the variability of respiration in terrestrial ecosystems: moving beyond Q10, *Glob. Change Biol.*, 12, 154–164, <https://doi.org/10.1111/j.1365-2486.2005.01065.x>, 2006.
- Dorrepaal, E., Toet, S., van Logtestijn, R. S. P., Swart, E., van de Weg, M. J., Callaghan, T. V., and Aerts, R.: Carbon respiration from subsurface peat accelerated by climate warming in the subarctic, *Nature*, 460, 616–619, <https://doi.org/10.1038/nature08216>, 2009.
- Ernakovich, J. G., Hopping, K. A., Berdanier, A. B., Simpson, R. T., Kachergis, E. J., Steltzer, H., and Wallenstein, M. D.: Predicted responses of arctic and alpine ecosystems to altered seasonality under climate change, *Glob. Change Biol.*, 20, 3256–3269, <https://doi.org/10.1111/gcb.12568>, 2014.
- Fahnestock, J. T., Jones, M. H., Brooks, P. D., Walker, D. A., and Welker, J. M.: Winter and early spring CO₂ efflux from tundra communities of northern Alaska, *J. Geophys. Res. Atmospheres*, 103, 29023–29027, <https://doi.org/10.1029/98JD00805>, 1998.
- Fahnestock, J. T., Jones, M. H., and Welker, J. M.: Wintertime CO₂ efflux from Arctic soils: Implications for annual carbon budgets, *Glob. Biogeochem. Cycles*, 13, 775–779, <https://doi.org/10.1029/1999GB900006>, 1999.
- Friggens, N. L., Hugelius, G., Kokelj, S. V., Murton, J. B., Phoenix, G. K., and Hartley, I. P.: Positive rhizosphere priming accelerates carbon release from permafrost soils, *Nat. Commun.*, 16, 3576, <https://doi.org/10.1038/s41467-025-58845-9>, 2025.
- Frossard, A., De Maeyer, L., Adamczyk, M., Svenning, M., Verleyen, E., and Frey, B.: Microbial carbon use and associated changes in microbial community structure in high-Arctic tundra soils under elevated temperature, *Soil Biol. Biochem.*, 162, 108419, <https://doi.org/10.1016/j.soilbio.2021.108419>, 2021.
- García Criado, M., Myers-Smith, I. H., Bjorkman, A. D., Elmendorf, S. C., Normand, S., Aastrup, P., Aerts, R., Alatalo, J. M., Baeten, L., Björk, R. G., Björkman, M. P., Boulanger-Lapointe, N., Butler, E. E., Cooper, E. J., Cornelissen, J. H. C., Daskalova, G. N., Fadrique, B., Forbes, B. C., Henry, G. H. R., Hollister, R. D., Høye, T. T., Jacobsen, I. B. D., Jägerbrand, A. K., Jónsdóttir, I. S., Kaarlejärvi, E., Khitun, O., Klanderud, K., Kolari, T. H. M., Lang, S. I., Lecomte, N., Lenoir, J., Macek, P., Messier, J., Michelsen, A., Molau, U., Muscarella, R., Nielsen, M.-L., Petit Bon, M., Post, E., Raundrup, K., Rinnan, R., Rixen, C., Ryde, I., Serra-Diaz, J. M., Schaepman-Strub, G., Schmidt, N. M., Schrodt, F., Sjögersten, S., Steinbauer, M. J., Stewart, L., Strandberg, B., Tolvanen, A., Tweedie, C. E., and Vellend, M.: Plant diversity dynamics over space and time in a warming Arctic, *Nature*, 642, 653–661, <https://doi.org/10.1038/s41586-025-08946-8>, 2025.
- Gorham, E.: Northern Peatlands: Role in the Carbon Cycle and Probable Responses to Climatic Warming, *Ecol. Appl.*, 1, 182–195, <https://doi.org/10.2307/1941811>, 1991.
- Grogan, P.: Cold season respiration across a low arctic landscape: the influence of vegetation type, snow depth, and interannual climatic variation, *Arct. Antarct. Alp. Res.*, 44, 446–456, <https://doi.org/10.1657/1938-4246-44.4.446>, 2012.
- Gudas, C., Karlsson, J., and Bastviken, D.: When does temperature matter for ecosystem respiration?, *Environ. Res. Commun.*, 3, 121001, <https://doi.org/10.1088/2515-7620/ac3b9f>, 2021.
- Hassol, S. J.: Impacts of a warming Arctic: Arctic Climate Impact Assessment, Cambridge University Press, Cambridge, U.K. ; New York, N.Y., 2004.
- Henry, G. h. r. and Molau, U.: Tundra plants and climate change: the International Tundra Experiment (ITEX), *Glob. Change Biol.*, 3, 1–9, <https://doi.org/10.1111/j.1365-2486.1997.gcb132.x>, 1997.

- Hermesdorf, L., Liu, Y., Michelsen, A., Westergaard-Nielsen, A., Mortensen, L. H., Jepsen, M. S., Sigsgaard, C., and Elberling, B.: Long-term changes in the daytime growing season carbon dioxide exchange following increased temperature and snow cover in arctic tundra, *Glob. Change Biol.*, 30, e17087, <https://doi.org/10.1111/gcb.17087>, 2024.
- Hicks Pries, C. E., Schuur, E. A. G., and Crummer, K. G.: Thawing permafrost increases old soil and autotrophic respiration in tundra: Partitioning ecosystem respiration using $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$, *Glob. Change Biol.*, 19, 649–661, <https://doi.org/10.1111/gcb.12058>, 2013.
- Hollesen, J., Buchwal, A., Rachlewicz, G., Hansen, B. U., Hansen, M. O., Stecher, O., and Elberling, B.: Winter warming as an important co-driver for *Betula nana* growth in western Greenland during the past century, *Glob. Change Biol.*, 21, 2410–2423, <https://doi.org/10.1111/gcb.12913>, 2015.
- Hollister, R. D., Webber, P. J., and Tweedie, C. E.: The response of Alaskan arctic tundra to experimental warming: differences between short- and long-term responses, *Glob. Change Biol.*, 11, 525–536, <https://doi.org/10.1111/j.1365-2486.2005.00926.x>, 2005.
- Hollister, R. D., Elphinstone, C., Henry, G. H. R., Bjorkman, A. D., Klanderud, K., Björk, R. G., Björkman, M. P., Bokhorst, S., Carbognani, M., Cooper, E. J., Dorrepaal, E., Elmendorf, S. C., Fetcher, N., Gallois, E. C., Guðmundsson, J., Healey, N. C., Jónsdóttir, I. S., Klarenberg, I. J., Oberbauer, S. F., Macek, P., May, J. L., Mereghetti, A., Molau, U., Petraglia, A., Rinnan, R., Rixen, C., and Wookey, P. A.: A review of open top chamber (OTC) performance across the ITEX Network, *Arct. Sci.*, 9, 331–344, <https://doi.org/10.1139/as-2022-0030>, 2023.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw, *Proc. Natl. Acad. Sci.*, 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020.
- Jones, M. H., Fahnestock, J. T., and Welker, J. M.: Early and Late Winter CO_2 Efflux from Arctic Tundra in the Kuparuk River Watershed, Alaska, U.S.A., *Arct. Antarct. Alp. Res.*, 31, 187–190, <https://doi.org/10.1080/15230430.1999.12003296>, 1999.
- Jónsdóttir, I. S., Halbritter, A. H., Christiansen, C. T., Althuizen, I. H. J., Haugum, S. V., Henn, J. J., Björnsdóttir, K., Maitner, B. S., Malhi, Y., Michaletz, S. T., Roos, R. E., Klanderud, K., Lee, H., Enquist, B. J., and Vandvik, V.: Intraspecific trait variability is a key feature underlying high Arctic plant community resistance to climate warming, *Ecol. Monogr.*, 93, e1555, <https://doi.org/10.1002/ecm.1555>, 2023.
- Karhu, K., Auffret, M. D., Dungait, J. A. J., Hopkins, D. W., Prosser, J. I., Singh, B. K., Subke, J.-A., Wookey, P. A., Ågren, G. I., Sebastià, M.-T., Gouriveau, F., Bergkvist, G., Meir, P., Nottingham, A. T., Salinas, N., and Hartley, I. P.: Temperature sensitivity of soil respiration rates enhanced by microbial community response, *Nature*, 513, 81–84, <https://doi.org/10.1038/nature13604>, 2014.
- Knoblauch, C., Beer, C., Schuett, A., Sauerland, L., Liebner, S., Steinhof, A., Rethemeyer, J., Grigoriev, M. N., Faguet, A., and Pfeiffer, E.-M.: Carbon Dioxide and Methane Release Following Abrupt Thaw of Pleistocene Permafrost Deposits in Arctic Siberia, *J. Geophys. Res. Biogeosciences*, 126, e2021JG006543, <https://doi.org/10.1029/2021JG006543>, 2021.
- Kropp, H., Loranty, M. M., Natali, S. M., Kholodov, A. L., Rocha, A. V., Myers-Smith, I., Abbot, B. W., Abermann, J., Blanc-Betes, E., Blok, D., Blume-Werry, G., Boike, J., Breen, A. L., Cahoon, S. M. P., Christiansen, C. T., Douglas, T. A., Epstein, H. E., Frost, G. V., Goeckede, M., Høye, T. T., Mamet, S. D., O'Donnell, J. A., Olefeldt, D., Phoenix, G. K., Salmon, V. G., Sannel, A. B. K., Smith, S. L., Sonnentag, O., Vaughn, L. S., Williams, M., Elberling, B., Gough, L., Hjort, J., Lafleur, P. M., Euskirchen, E. S., Heijmans, M. M., Humphreys, E. R., Iwata, H., Jones, B. M., Jorgenson, M. T., Grünberg, I., Kim, Y., Laundre, J., Mauritz, M., Michelsen, A., Schaepman-Strub, G., Tape, K. D., Ueyama, M., Lee, B.-Y., Langley, K., and Lund,

M.: Shallow soils are warmer under trees and tall shrubs across Arctic and Boreal ecosystems, *Environ. Res. Lett.*, 16, 015001, <https://doi.org/10.1088/1748-9326/abc994>, 2020.

600 Leffler, A. J., Klein, E. S., Oberbauer, S. F., and Welker, J. M.: Coupled long-term summer warming and deeper snow alters species composition and stimulates gross primary productivity in tussock tundra, *OECOLOGIA*, 181, 287–297, <https://doi.org/10.1007/s00442-015-3543-8>, 2016.

605 Leffler, A. J., Serrone, A., Bolek, A., Rocha, A. V., Korrensalo, A., Vähä, A., Tremblay, A., Barr, A., Salazar, A., Sabrekov, A. F., Rasanen, A., Mavrovic, A., Roy, A., Panov, A. V., Townsend-Small, A., Niyazova, A., Prokushkin, A., Jonsson, A., Lindroth, A., Christen, A., Westergaard-Nielsen, A., Varlagin, A., Black, T. A., Rutgersson, A., Sytiuk, A., Virkkala, A.-M., Lohila, A., Ojala, A., Thiboult, A., Sepulveda-Jauregui, A., Malhotra, A., Kotani, A., Holmes, B., Elberling, B., Rogers, B., Johannesson, C., Trotta, C., Voigt, C., Christiansen, C. T., Dieleman, C., Jones, C. P., Wille, C., Biasi, C., Minions, C., Cunada, C., Arp, C. D., Schulze, C., Treat, C., Edgar, C., Emmerton, C. A., Nadeau, D. F., Ilyasov, D. V., Papale, D., Bastviken, D., Holl, D., Lipson, D., Olefeldt, D., Zona, D., Schuur, T., Tuittila, E. S., López-Blanco, E., Koster, E., Dyukarev, E., Blanc-Betes, E., Lapshina, E., Veretennikova, E., Canfora, E., Männisto, E., Humphreys, E., Stanley, E. H., Lundin, E., Sahlée, E., Nicholls, E., Kyzivat, E. D., Euskirchen, S. E., Kane, E. S., Tigreros, F. G., Domine, F., Clayer, F., Parmentier, F., Bouchard, F., Thalasso, F., Hould-Gosselin, G., Hensgens, G., Kling, G. W., Rocher-Ros, G., Pastorello, G., Iwahana, G., Clark, G., Falvo, G., Henry, G., Granath, G., Webb, H., Alcock, H., Thorgeirsson, H., Lee, H., Silvennoinen, H. M., Nykanen, H., Rautakoski, H., et al.: Arctic-Boreal Vulnerability Experiment (ABOve)ABCFlux v2: Arctic-Boreal CO₂ and CH₄ In-situ Flux and Environmental Data, <https://doi.org/10.3334/ORNLDAAAC/2448>, 1 January 2025.

615 Lewkowicz, A. G. and Way, R. G.: Extremes of summer climate trigger thousands of thermokarst landslides in a High Arctic environment, *Nat. Commun.*, 10, 1329, <https://doi.org/10.1038/s41467-019-09314-7>, 2019.

López-Blanco, E., Topp-Jørgensen, E., Christensen, T. R., Rasch, M., Skov, H., Arndal, M. F., Bret-Harte, M. S., Callaghan, T. V., and Schmidt, N. M.: Towards an increasingly biased view on Arctic change, *Nat. Clim. Change*, 14, 152–155, <https://doi.org/10.1038/s41558-023-01903-1>, 2024.

620 Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726–738, <https://doi.org/10.1890/12-0279.1>, 2013.

625 Maes, S. L., Dietrich, J., Midolo, G., Schwieger, S., Kummu, M., Vandvik, V., Aerts, R., Althuizen, I. H. J., Biasi, C., Björk, R. G., Böhner, H., Carbognani, M., Chiari, G., Christiansen, C. T., Clemmensen, K. E., Cooper, E. J., Cornelissen, J. H. C., Elberling, B., Faubert, P., Fetcher, N., Forte, T. G. W., Gaudard, J., Gavazov, K., Guan, Z., Guðmundsson, J., Gya, R., Hallin, S., Hansen, B. B., Haugum, S. V., He, J.-S., Hicks Pries, C., Hovenden, M. J., Jalava, M., Jónsdóttir, I. S., Juhanson, J., Jung, J. Y., Kaarlejärvi, E., Kwon, M. J., Lamprecht, R. E., Le Moullec, M., Lee, H., Marushchak, M. E., Michelsen, A., Munir, T. M., Myrsky, E. M., Nielsen, C. S., Nyberg, M., Olofsson, J., Óskarsson, H., Parker, T. C., Pedersen, E. P., Petit Bon, M., Petraglia, A., Raundrup, K., Ravn, N. M. R., Rinnan, R., Rodenhizer, H., Ryde, I., Schmidt, N. M., Schuur, E. A. G., Sjögersten, S., Stark, S., Strack, M., Tang, J., Tolvanen, A., Töpper, J. P., Väisänen, M. K., Van Logtestijn, R. S. P., Voigt, C., Walz, J., Weedon, J. T., Yang, Y., Yläne, H., Björkman, M. P., Sarneel, J. M., and Dorrepaal, E.: Environmental drivers of increased ecosystem respiration in a warming tundra, *Nature*, 629, 105–113, <https://doi.org/10.1038/s41586-024-07274-7>, 2024.

630 Magnússon, R. Í., Sass-Klaassen, U., Limpens, J., Karsanaev, S. V., Ras, S., van Huissteden, K., Blok, D., and Heijmans, M. M. P. D.: Spatiotemporal variability in precipitation-growth association of *Betula nana* in the Siberian lowland tundra, *J. Ecol.*, 111, 1882–1904, <https://doi.org/10.1111/1365-2745.14165>, 2023.

635 Mahecha, M. D., Reichstein, M., Carvalhais, N., Lasslop, G., Lange, H., Seneviratne, S. I., Vargas, R., Ammann, C., Arain, M. A., Cescatti, A., Janssens, I. A., Migliavacca, M., Montagnani, L., and Richardson, A. D.: Global Convergence in the Temperature Sensitivity of Respiration at Ecosystem Level, *Science*, 329, 838–840, <https://doi.org/10.1126/science.1189587>, 2010.

- Marion, G. M., Henry, G. H. R., Freckman, D. W., Johnstone, J., Jones, G., Jones, M. H., Lévesque, E., Molau, U., Mølgaard, P., Parsons, A. N., J. Svoboda, and Virginia, R. A.: Open-top designs for manipulating field temperature in high-latitude ecosystems, *Glob. Change Biol.*, 3, 20–32, <https://doi.org/10.1111/j.1365-2486.1997.gcb136.x>, 1997.
- Metcalfe, D. B., Hermans, T. D. G., Ahlstrand, J., Becker, M., Berggren, M., Björk, R. G., Björkman, M. P., Blok, D., Chaudhary, N., Chisholm, C., Classen, A. T., Hasselquist, N. J., Jonsson, M., Kristensen, J. A., Kumordzi, B. B., Lee, H., Mayor, J. R., Prevéy, J., Pantazatou, K., Rousk, J., Sponseller, R. A., Sundqvist, M. K., Tang, J., Uddling, J., Wallin, G., Zhang, W., Ahlström, A., Tenenbaum, D. E., and Abdi, A. M.: Patchy field sampling biases understanding of climate change impacts across the Arctic, *Nat. Ecol. Evol.*, 2, 1443–1448, <https://doi.org/10.1038/s41559-018-0612-5>, 2018.
- Morgner, E., Elberling, B., Strebel, D., and Cooper, E. J.: The importance of winter in annual ecosystem respiration in the High Arctic: effects of snow depth in two vegetation types, *Polar Res.*, 29, 58–74, <https://doi.org/10.3402/polar.v29i1.6052>, 2010.
- Myers-Smith, I. H., Grabowski, M. M., Thomas, H. J. D., Angers-Blondin, S., Daskalova, G. N., Bjorkman, A. D., Cunliffe, A. M., Assmann, J. J., Boyle, J. S., McLeod, E., McLeod, S., Joe, R., Lennie, P., Arey, D., Gordon, R. R., and Eckert, C. D.: Eighteen years of ecological monitoring reveals multiple lines of evidence for tundra vegetation change, *Ecol. Monogr.*, 89, e01351, <https://doi.org/10.1002/ecm.1351>, 2019.
- Myers-Smith, I. H., Kerby, J. T., Phoenix, G. K., Bjerke, J. W., Howard E. Epstein, Assmann, J. J., John, C., Andreu-Hayles, L., Angers-Blondin, S., Beck, P. S. A., Berner, L. T., Bhatt, U. S., Bjorkman, A. D., Blok, D., Bryn, A., Christiansen, C. T., Cornelissen, J. H. C., Cunliffe, A. M., Elmendorf, S. C., Forbes, B. C., Goetz, S. J., Hollister, R. D., de Jong, R., Loranty, M. M., Macias-Fauria, M., Maseyk, K., Normand, S., Olofsson, J., Parker, T. C., Parmentier, F.-J. W., Post, E., Schaepman-Strub, G., Stordal, F., Sullivan, P. F., Thomas, H. J. D., Tømmervik, H., Treharne, R., Tweedie, C. E., Walker, D. A., Wilmking, M., and Wipf, S.: Complexity revealed in the greening of the Arctic, *Nat. Clim. Change*, 10, 106–117, <https://doi.org/10.1038/s41558-019-0688-1>, 2020.
- Natali, S. M., Watts, J. D., Rogers, B. M., Potter, S., Ludwig, S. M., Selbmann, A.-K., Sullivan, P. F., Abbott, B. W., Arndt, K. A., Birch, L., Björkman, M. P., Bloom, A. A., Celis, G., Christensen, T. R., Christiansen, C. T., Commene, R., Cooper, E. J., Crill, P., Czimczik, C., Davydov, S., Du, J., Egan, J. E., Elberling, B., Euskirchen, E. S., Friborg, T., Genet, H., Göckede, M., Goodrich, J. P., Grogan, P., Helbig, M., Jafarov, E. E., Jastrow, J. D., Kalhori, A. A. M., Kim, Y., Kimball, J. S., Kutzbach, L., Lara, M. J., Larsen, K. S., Lee, B.-Y., Liu, Z., Loranty, M. M., Lund, M., Lupascu, M., Madani, N., Malhotra, A., Matamala, R., McFarland, J., McGuire, A. D., Michelsen, A., Minions, C., Oechel, W. C., Olefeldt, D., Parmentier, F.-J. W., Pirk, N., Poulter, B., Quinton, W., Rezanezhad, F., Risk, D., Sachs, T., Schaefer, K., Schmidt, N. M., Schuur, E. A. G., Semenchuk, P. R., Shaver, G., Sonnentag, O., Starr, G., Treat, C. C., Waldrop, M. P., Wang, Y., Welker, J., Wille, C., Xu, X., Zhang, Z., Zhuang, Q., and Zona, D.: Large loss of CO₂ in winter observed across the northern permafrost region, *Nat. Clim. Change*, 9, 852–857, <https://doi.org/10.1038/s41558-019-0592-8>, 2019.
- Niittynen, P.: *snowman*: an open-source R package for automated 30-m snow and ice cover mapping using the Landsat archive, *EGUsphere*, 1–0, <https://doi.org/10.31223/X5FF32>, 2026.
- Niu, S., Chen, W., Liáng, L. L., Sierra, C. A., Xia, J., Wang, S., Heskell, M., Patel, K. F., Bond-Lamberty, B., Wang, J., Yvon-Durocher, G., Kirschbaum, M. U. F., Atkin, O. K., Huang, Y., Yu, G., and Luo, Y.: Temperature responses of ecosystem respiration, *Nat. Rev. Earth Environ.*, 5, 559–571, <https://doi.org/10.1038/s43017-024-00569-3>, 2024.
- Oberbauer, S., Starr, G., and Pop, E.: Effects of extended growing season and soil warming on carbon dioxide and methane exchange of tussock tundra in Alaska, *J. Geophys. Res.-ATMOSPHERES*, 103, 29075–29082, <https://doi.org/10.1029/98JD00522>, 1998.

- Olefelt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P., McGuire, A. D., Romanovsky, V. E., Sannel, A. B. K., Schuur, E. a. G., and Turetsky, M. R.: Circumpolar distribution and carbon storage of thermokarst landscapes, *Nat. Commun.*, 7, 13043, <https://doi.org/10.1038/ncomms13043>, 2016.
- Park, S.-W., Mun, J.-H., Lee, H., Steinert, N. J., An, S.-I., Shin, J., and Kug, J.-S.: Continued permafrost ecosystem carbon loss under net-zero and negative emissions, *Sci. Adv.*, 11, eadn8819, <https://doi.org/10.1126/sciadv.adn8819>, 2025.
- Pattison, R. R. and Welker, J. M.: Differential ecophysiological response of deciduous shrubs and a graminoid to long-term experimental snow reductions and additions in moist acidic tundra, Northern Alaska, *Oecologia*, 174, 339–350, <https://doi.org/10.1007/s00442-013-2777-6>, 2014.
- Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, 3, 1–10, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Rixen, C., Høye, T. T., Macek, P., Aerts, R., Alatalo, J. M., Anderson, J. T., Arnold, P. A., Barrio, I. C., Bjerke, J. W., Björkman, M. P., Blok, D., Blume-Werry, G., Boike, J., Bokhorst, S., Carbognani, M., Christiansen, C. T., Convey, P., Cooper, E. J., Cornelissen, J. H. C., Coulson, S. J., Dorrepaal, E., Elberling, B., Elmendorf, S. C., Elphinstone, C., Forte, T. G. W., Frei, E. R., Geange, S. R., Gehrmann, F., Gibson, C., Grogan, P., Halbritter, A. H., Harte, J., Henry, G. H. R., Inouye, D. W., Irwin, R. E., Jespersen, G., Jónsdóttir, I. S., Jung, J. Y., Kluges, D. H., Kudo, G., Lämsä, J., Lee, H., Lembrechts, J. J., Lett, S., Lynn, J. S., Mann, H. M. R., Mastepanov, M., Morse, J., Myers-Smith, I. H., Olofsson, J., Paavola, R., Petraglia, A., Phoenix, G. K., Semenchuk, P., Siewert, M. B., Slatyer, R., Spasojevic, M. J., Suding, K., Sullivan, P., Thompson, K. L., Väisänen, M., Vandvik, V., Venn, S., Walz, J., Way, R., Welker, J. M., Wipf, S., and Zong, S.: Winters are changing: snow effects on Arctic and alpine tundra ecosystems, *Arct. Sci.*, 8, 572–608, <https://doi.org/10.1139/as-2020-0058>, 2022.
- Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., Gurevitch, J., and GCTE-NEWS: A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming, *OECOLOGIA*, 126, 543–562, <https://doi.org/10.1007/s004420000544>, 2001.
- Sarneel, J. M., Sundqvist, M. K., Molau, U., Björkman, M. P., and Alatalo, J. M.: Decomposition rate and stabilization across six tundra vegetation types exposed to >20 years of warming, *Sci. Total Environ.*, 724, 138304, <https://doi.org/10.1016/j.scitotenv.2020.138304>, 2020.
- Schädel, C., Koven, C. D., Lawrence, D. M., Celis, G., Garnello, A. J., Hutchings, J., Mauritz, M., Natali, S. M., Pegoraro, E., Rodenhizer, H., Salmon, V. G., Taylor, M. A., Webb, E. E., Wieder, W. R., and Schuur, E. A.: Divergent patterns of experimental and model-derived permafrost ecosystem carbon dynamics in response to Arctic warming, *Environ. Res. Lett.*, 13, 105002, <https://doi.org/10.1088/1748-9326/aae0ff>, 2018.
- Schimel, J. P., Bilbrough, C., and Welker, J. M.: Increased snow depth affects microbial activity and nitrogen mineralization in two Arctic tundra communities, *Soil Biol. Biochem.*, 36, 217–227, <https://doi.org/10.1016/j.soilbio.2003.09.008>, 2004.
- Schimel, J. P., Fahnstock, J., Michaelson, G., Mikan, C., Ping, C.-L., Romanovsky, V. E., and Welker, J.: Cold-season Production of CO₂ in Arctic Soils: Can Laboratory and Field Estimates Be Reconciled through a Simple Modeling Approach?, *Arct. Antarct. Alp. Res.*, 38, 249–256, [https://doi.org/10.1657/1523-0430\(2006\)38%5B249:CPOCIA%5D2.0.CO;2](https://doi.org/10.1657/1523-0430(2006)38%5B249:CPOCIA%5D2.0.CO;2), 2006.
- Schuur, E., Bockheim, J., Canadell, J. G., Euskirchen, E., Field, C. B., Goryachkin, S. V., Hagemann, S., Kuhry, P., Lafleur, P. M., Lee, H., Mazhitova, G., Nelson, F. E., Rinke, A., Romanovsky, V. E., Shiklomanov, N., Tarnocai, C., Venevsky, S., Vogel, J. G., and Zimov, S. A.: Vulnerability of Permafrost Carbon to Climate Change: Implications for the Global Carbon Cycle, *BioScience*, 58, 701–714, <https://doi.org/10.1641/B580807>, 2008.

- Schuur, E. a. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- 720 Schuur, E. A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven, C., Leshyk, V., Lawrence, D., Lorant, M. M., Mauritz, M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C., Strauss, J., Treat, C., and Turetsky, M.: Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic, *Annu. Rev. Environ. Resour.*, 47, 343–371, <https://doi.org/10.1146/annurev-environ-012220-011847>, 2022.
- 725 Schwieger, S., Dorrepaal, E., Petit Bon, M., Vandvik, V., le Roux, E., Strack, M., Yang, Y., Venn, S., van den Hoogen, J., Valiño, F., Thomas, H. J. D., te Beest, M., Suzuki, S., Petraglia, A., Myers-Smith, I. H., Munir, T. M., Michelsen, A., Løkken, J. O., Li, Q., Koike, T., Klanderud, K., Karr, E. H., Jónsdóttir, I. S., Hollister, R. D., Hofgaard, A., Hassan, I. A., Genxu, W., Filippova, N., Crowther, T. W., Clark, K., Christiansen, C. T., Casanova-Katny, A., Carbognani, M., Bokhorst, S., Björnsdóttir, K., Asplund, J., Althuisen, I., Alonso, R., Alatalo, J., Agathokleous, E., Aerts, R., and Sarneel, J. M.: Environmental Conditions Modulate Warming Effects on Plant Litter Decomposition Globally, *Ecol. Lett.*, 28, e70026, <https://doi.org/10.1111/ele.70026>, 2025.
- 730 Semenchuk, P. R., Christiansen, C. T., Grogan, P., Elberling, B., and Cooper, E. J.: Long-term experimentally deepened snow decreases growing-season respiration in a low- and high-arctic tundra ecosystem, *J. Geophys. Res.-BIOGEOSCIENCES*, 121, 1236–1248, <https://doi.org/10.1002/2015JG003251>, 2016.
- Shukla, P. R., Skeg, J., Calvo Buendia, E., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D. C., Zhai, P., Slade, R., Connors, S., van Diemen, S., Ferrat, M., Haughey, E., Luz, S., Pathak, M., Petzold, J., Portugal Pereira, J., Vyas, P., Huntley, E., Kissick, K., Belkacemi, M., and Malley, J.: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems, 2019.
- 735 Slatyer, R. A., Umbers, K. D. L., and Arnold, P. A.: Ecological responses to variation in seasonal snow cover, *Conserv. Biol.*, 36, e13727, <https://doi.org/10.1111/cobi.13727>, 2022.
- 740 Sulman, B. N., Moore, J. A. M., Abramoff, R., Averill, C., Kivlin, S., Georgiou, K., Sridhar, B., Hartman, M. D., Wang, G., Wieder, W. R., Bradford, M. A., Luo, Y., Mayes, M. A., Morrison, E., Riley, W. J., Salazar, A., Schimel, J. P., Tang, J., and Classen, A. T.: Multiple models and experiments underscore large uncertainty in soil carbon dynamics, *Biogeochemistry*, 141, 109–123, <https://doi.org/10.1007/s10533-018-0509-z>, 2018.
- 745 Swindles, G. T., Morris, P. J., Mullan, D., Watson, E. J., Turner, T. E., Roland, T. P., Amesbury, M. J., Kokfelt, U., Schoning, K., Pratte, S., Gallego-Sala, A., Charman, D. J., Sanderson, N., Garneau, M., Carrivick, J. L., Woulds, C., Holden, J., Parry, L., and Galloway, J. M.: The long-term fate of permafrost peatlands under rapid climate warming, *Sci. Rep.*, 5, 17951, <https://doi.org/10.1038/srep17951>, 2015.
- Tarnocai, C., Canadell, J. G., Schuur, E. a. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Glob. Biogeochem. Cycles*, 23, <https://doi.org/10.1029/2008GB003327>, 2009.
- 750 Tingley, M. P. and Huybers, P.: Recent temperature extremes at high northern latitudes unprecedented in the past 600 years, *Nature*, 496, 201–205, <https://doi.org/10.1038/nature11969>, 2013.
- 755 Van Sundert, K., Leuzinger, S., Bader, M. K. -F., Chang, S. X., De Kauwe, M. G., Dukes, J. S., Langley, J. A., Ma, Z., Mariën, B., Reynaert, S., Ru, J., Song, J., Stocker, B., Terrer, C., Thoresen, J., Vanuytrecht, E., Wan, S., Yue, K., and Vicca, S.: When things get MESI: The Manipulation Experiments Synthesis Initiative—A coordinated effort to synthesize terrestrial global change experiments, *Glob. Change Biol.*, 29, 1922–1938, <https://doi.org/10.1111/gcb.16585>, 2023.

- Virkkala, A.-M., Abdi, A. M., Luoto, M., and Metcalfe, D. B.: Identifying multidisciplinary research gaps across Arctic terrestrial gradients, *Environ. Res. Lett.*, 14, 124061, <https://doi.org/10.1088/1748-9326/ab4291>, 2019.
- 760 Virkkala, A.-M., Aalto, J., Rogers, B. M., Tagesson, T., Treat, C. C., Natali, S. M., Watts, J. D., Potter, S., Lehtonen, A., Mauritz, M., Schuur, E. A. G., Kochendorfer, J., Zona, D., Oechel, W., Kobayashi, H., Humphreys, E., Goeckede, M., Iwata, H., Lafleur, P. M., Euskirchen, E. S., Bokhorst, S., Marushchak, M., Martikainen, P. J., Elberling, B., Voigt, C., Biasi, C., Sonnentag, O., Parmentier, F.-J. W., Ueyama, M., Celis, G., St. Louis, V. L., Emmerton, C. A., Peichl, M., Chi, J., Järveoja, J., Nilsson, M. B., Oberbauer, S. F., Torn, M. S., Park, S.-J., Dolman, H., Mammarella, I., Chae, N., Poyatos, R., López-Blanco, E., Christensen, T. R., Kwon, M. J., Sachs, T., Holl, D., and Luoto, M.: Statistical upscaling of ecosystem CO₂ fluxes across the terrestrial tundra and boreal domain: Regional patterns and uncertainties, *Glob. Change Biol.*, 27, 4040–4059, <https://doi.org/10.1111/gcb.15659>, 2021.
- 765
- 770 Virkkala, A.-M., Natali, S. M., Rogers, B. M., Watts, J. D., Savage, K., Connon, S. J., Mauritz, M., Schuur, E. A. G., Peter, D., Minions, C., Nojeim, J., Commane, R., Emmerton, C. A., Goeckede, M., Helbig, M., Holl, D., Iwata, H., Kobayashi, H., Kolari, P., López-Blanco, E., Marushchak, M. E., Mastepanov, M., Merbold, L., Parmentier, F.-J. W., Peichl, M., Sachs, T., Sonnentag, O., Ueyama, M., Voigt, C., Aurela, M., Boike, J., Celis, G., Chae, N., Christensen, T. R., Bret-Harte, M. S., Dengel, S., Dolman, H., Edgar, C. W., Elberling, B., Euskirchen, E., Grelle, A., Hatakka, J., Humphreys, E., Järveoja, J., Kotani, A., Kutzbach, L., Laurila, T., Lohila, A., Mammarella, I., Matsuura, Y., Meyer, G., Nilsson, M. B., Oberbauer, S. F., Park, S.-J., Petrov, R., Prokushkin, A. S., Schulze, C., St. Louis, V. L., Tuittila, E.-S., Tuovinen, J.-P., Quinton, W., Varlagin, A., Zona, D., and Zyryanov, V. I.: The ABCflux database: Arctic–boreal CO₂ flux observations and ancillary information aggregated to monthly time steps across terrestrial ecosystems, *Earth Syst. Sci. Data*, 14, 179–208, <https://doi.org/10.5194/essd-14-179-2022>, 2022.
- 775
- 780 Virkkala, A.-M., Niittynen, P., Kemppinen, J., Marushchak, M. E., Voigt, C., Hensgens, G., Kerttula, J., Happonen, K., Tyystjärvi, V., Biasi, C., Hultman, J., Rinne, J., and Luoto, M.: High-resolution spatial patterns and drivers of terrestrial ecosystem carbon dioxide, methane, and nitrous oxide fluxes in the tundra, *Biogeosciences*, 21, 335–355, <https://doi.org/10.5194/bg-21-335-2024>, 2024.
- 785 Vogel, J., Schuur, E. A. G., Trucco, C., and Lee, H.: Response of CO₂ exchange in a tussock tundra ecosystem to permafrost thaw and thermokarst development, *J. Geophys. Res. Biogeosciences*, 114, 2008JG000901, <https://doi.org/10.1029/2008JG000901>, 2009.
- Wagner, J. and Hugelius, G.: Comparing Multiscale Object-Based Image Analysis and Deep Learning for High-Resolution Classification of Ice Wedge Polygon Tundra, *Can. J. Remote Sens.*, 52, 2654952, <https://doi.org/10.1080/07038992.2026.2654952>, 2026.
- 785 Weedon, J. T., A. Kowalchuk, G., Aerts, R., van Hal, J., van Logtestijn, R., Taş, N., F. M. Röling, W., and M. van Bodegom, P.: Summer warming accelerates sub-arctic peatland nitrogen cycling without changing enzyme pools or microbial community structure, *Glob. Change Biol.*, 18, 138–150, <https://doi.org/10.1111/j.1365-2486.2011.02548.x>, 2012.
- 790 Wei, B., Zhang, D., Voigt, C., Zhou, W., Bai, Y., Zheng, Z., Xie, Y., Zhao, C., Wang, F., Huang, L., Yang, G., Kou, D., Peng, Y., Luo, Y., Peñuelas, J., and Yang, Y.: Progressive decline in soil nitrogen stocks with warming in a Tibetan permafrost ecosystem, *Nat. Geosci.*, 18, 997–1004, <https://doi.org/10.1038/s41561-025-01786-1>, 2025.
- Welker, J. M., Molau, U., Parsons, A. N., Robinson, C. H., and Wookey, P. A.: Responses of *Dryas octopetala* to ITEX environmental manipulations: a synthesis with circumpolar comparisons, *Glob. Change Biol.*, 3, 61–73, <https://doi.org/10.1111/j.1365-2486.1997.gcb143.x>, 1997.

- 795 Welker, J. M., Brown, K. B., and Fahnestock, J. T.: CO₂ Flux in Arctic and Alpine Dry Tundra: Comparative Field Responses Under Ambient and Experimentally Warmed Conditions, *Arct. Antarct. Alp. Res.*, 31, 272–277, <https://doi.org/10.1080/15230430.1999.12003309>, 1999.
- Welker, J. M., Fahnestock, J. T., and Jones, M. H.: Annual CO₂ Flux in Dry and Moist Arctic Tundra: Field Responses to Increases in Summer Temperatures and Winter Snow Depth, *Clim. Change*, 44, 139–150,
800 <https://doi.org/10.1023/A:1005555012742>, 2000.
- Welker, J. M., Fahnestock, J. T., Henry, G. H. R., O’Dea, K. W., and Chimner, R. A.: CO₂ exchange in three Canadian High Arctic ecosystems: response to long-term experimental warming, *Glob. Change Biol.*, 10, 1981–1995, <https://doi.org/10.1111/j.1365-2486.2004.00857.x>, 2004.
- Wells, J., Schaedel, C., Riley, W. J., Tao, J., Mekonnen, Z. A., Lawrence, D. M., Swenson, S. C., Schuur, E., Huntzinger, D.
805 N., Poe, J. M., Maes, S., Dietrich, J., Schwieger, S., and Dorrepaal, E.: State-of-the-Arctic: Confronting State-of-the-Art Carbon Cycle Models with Pan-Arctic Warming Responses, AGU Fall Meeting Abstracts, ADS Bibcode: 2023AGUFM.B51C..03W, B51C-03, 2023.
- Wilson, S. D. and Nilsson, C.: Arctic alpine vegetation change over 20 years, *Glob. Change Biol.*, 15, 1676–1684, <https://doi.org/10.1111/j.1365-2486.2009.01896.x>, 2009.
- 810 Zimov, S. A., Schuur, E. A. G., and Chapin, F. S.: Permafrost and the Global Carbon Budget, *Science*, 312, 1612–1613, <https://doi.org/10.1126/science.1128908>, 2006.
- Zona, D., Gioli, B., Commane, R., Lindaas, J., Wofsy, S. C., Miller, C. E., Dinardo, S. J., Dengel, S., Sweeney, C., Karion, A., Chang, R. Y.-W., Henderson, J. M., Murphy, P. C., Goodrich, J. P., Moreaux, V., Liljedahl, A., Watts, J. D., Kimball, J. S., Lipson, D. A., and Oechel, W. C.: Cold season emissions dominate the Arctic tundra methane budget, *Proc. Natl. Acad. Sci.*, 113, 40–45, <https://doi.org/10.1073/pnas.1516017113>, 2016.
- R Core Team (2025). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <<https://www.R-project.org/>>