

Development of the Long-term Harmonized multi-satellite SIF (LHSIF) dataset at 0.05° resolution (1995–2023)

Chu Zou^{1,2,3}, Shanshan Du^{1,2}, Xinjie Liu^{1,2}, Liangyun Liu^{1,2,3}

¹ Key Laboratory of Digital Earth Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing, 100094, China

² University of Chinese Academy of Sciences, Beijing 100049, China

³ International Research Center of Big Data for Sustainable Development Goals, Beijing 100094, China

Correspondence to: Liangyun Liu (liuly@radi.ac.cn)

Supplementary Materials





Figure S1 Global distribution of FLUXNET sites labeled by their climate zones and IGBP vegetation types.

Table S 1 Information for the ChinaSpec sites

Site_Name	ID	Location	Landcover Type	Time range*
XiaoTangshan	XTS	40.1786°N, 116.4432°E	Cropland (winter wheat and maize rotation)	2017–2023
HuaiLai	HL	40.3489°N, 115.7882°E	A mixture of grassland and forest	2017–2023
DaMan	DM	38.8555°N, 100.3722°E	Cropland (maize)	2017–2022
ARou	AR	38.0444°N, 100.4647°E	Grassland	2019–2023
GuCheng	GC	39.1486°N, 115.7351°E	Cropland (winter wheat and maize rotation)	2020–2022

Figure S2 The relationship between the re-aggregated SCIAMACHY SIF (SIF_{reagg}) and coarse spatial resolution SCIAMACHY SIF ($SIF_{original}$) for different months in 2007.




Figure S3 The relationship between the re-aggregated GOME-2 SIF (SIF_reagg) and coarse spatial resolution GOME-2 SIF(SIF_original) for different months in 2007.




Figure S4 The relationship between the re-aggregated OCO-2 SIF (SIF_reagg) and coarse spatial resolution OCO-2 SIF(SIF_original) for different months in 2015.