



META4.0: a new mesoscale eddy network atlas derived from altimetry

Juliette Gamot ¹, Antoine Delepoulle ², Francesco Nencioli ¹, Marie-Isabelle Pujol ¹, and Gerald Dibarboure ³

¹Collecte Localisation Satellite (CLS), 11 rue Hermès, Ramonville Saint Agne, 31250, France

²Datlas, 70 rue de la Physique, Saint-Martin-d'Hères, France, 38421, France

³Centre Nationale d'Études Spatiales (CNES), 18 avenue Edouard Belin, Toulouse, 31400, France

Correspondence: Juliette Gamot (jgamot@groupcls.com)

Abstract. This study introduces the new global Mesoscale Eddy Trajectory Atlases (META4.0), which provide eddy detections, trajectories, and interaction networks derived from satellite altimetry. Eddy detection relies on the `pyeddytracker` (PET) algorithm (Mason et al., 2014), further optimized by Pegliasco et al. (2022), and represents a substantial improvement over the previous META3.2 product (SSALTO/DUACS, distributed by AVISO+ with CNES support).

5 The main advance of META4.0 is the explicit identification of eddy merging and splitting events. By combining grouping, which links detections across consecutive days, and segmentation, which tracks continuity through interaction events, trajectories are organized into networks of interconnected eddies. This network-based representation complements the classical single-trajectory view of eddy life cycles by explicitly accounting for eddy interactions.

The paper presents both diagnostic tools designed to explore individual eddy networks (e.g., timelines, spatial trajectories, and eddy properties such as effective radius or shape error) and the results of a global statistical analysis over more than three decades. These analyses provide new insights into network properties, eddy lifetimes, and the spatial and temporal distribution of merging and splitting events. Clustering analyses reveal recurrent interaction patterns and identify regions where eddy networks are particularly active. Independent datasets, including sea surface temperature and chlorophyll concentration, are used for qualitative validation of selected events. Finally, Lagrangian advection of synthetic particles highlights coherent forward and backward transport signatures associated with interaction events, providing a physical validation of the reconstructed networks.

Overall, META4.0 offers a novel and physically consistent framework to characterize mesoscale eddy interactions and to better understand their role in shaping ocean dynamics.

1 Introduction

20 1.1 General context

Mesoscale eddies are key drivers of ocean dynamics, playing a central role in the transport of water masses, energy, heat, salt, and biogeochemical tracers across both horizontal and vertical scales. These ubiquitous structures, typically spanning tens to



hundreds of kilometers and persisting from days to several years (Morrow and Le Traon, 2012), also exert a strong influence on marine biological and chemical processes (Beal et al., 2011; Chaigneau et al., 2011; Gaube et al., 2014; Gruber et al., 2011; Sosa-Gutierrez et al., 2025; Zhang et al., 2014). Coupling eddy-resolving physical observations with ecological datasets has substantially advanced our understanding of marine animal behavior (Braun et al., 2019; Chambault et al., 2019; Christie et al., 2010; Siegel et al., 2008; Staaterman et al., 2012) as well as the dispersion and fate of pollutants (Brach et al., 2018; Gilchrist et al., 2020). Continuous improvements in the accuracy of satellite altimetry over the past three decades have enabled global monitoring of mesoscale eddies, providing unprecedented insights into their role in ocean circulation and ecosystem functioning.

1.2 Eddy detection and atlases

Satellite altimetry provides near-global, weather-independent observations of mesoscale ocean variability, primarily through daily, multi-satellite gridded fields of absolute dynamic topography (ADT) and sea level anomaly (SLA). Although the spatiotemporal interpolation of nadir altimeter observations limits the effective resolution of these gridded products to spatial scales larger than ~ 100 km, they remain the backbone of global mesoscale eddy detection. Traditionally, eddy detection methods have relied on Eulerian approaches — based on fixed, time-invariant surface fields — using either geometric diagnostics (Chaigneau et al., 2008, 2009; Faghmous et al., 2013; Ioannou et al., 2017; Liu et al., 2016) or rotational metrics (Isern-Fontanet et al., 2003; Le Vu et al., 2018; Mkhinini et al., 2014; Morrow et al., 2004; Nencioli et al., 2010). Over the past decade, Lagrangian approaches — based on time-evolving surface fields — have also been introduced (Abernathy and Haller, 2018; Beron-Vera et al., 2008; El Aouni, 2021; Haller, 2016); however, their application at the global scale remains computationally intensive.

Several global atlases have been developed, beginning with the pioneering work of Chelton et al. (2011), which was later refined through collaborations between OSU, CLS, and CNES, leading to the Mesoscale Eddy Trajectory Atlas (META; Pegliasco et al. (2022)). Successive releases (META1.0exp, META2.0, META3.1exp, META3.2) progressively improved detection and tracking by refining search schemes, handling missing eddies, and extending coverage to coastal and island regions. These efforts established META as a reference global product, freely distributed via AVISO+ (https://data.aviso.altimetry.fr/aviso-gateway/data/META3.2_DT/). Other datasets followed similar principles, such as the global atlas by Tian et al. (2020), GOMEAD (Dong et al. (2022)), and ToEddies (Ioannou et al. (2024)), offering complementary approaches to eddy detection and tracking.

1.3 Single-trajectory approaches

Most global atlases, including META and others (Chelton et al. (2011); Tian et al. (2020); Dong et al. (2022)), rely on single-trajectory tracking, where eddies are followed over time by identifying candidates within a localized search window. The META2.0 version refined this method to handle the “missing eddy” problem through short temporal extrapolations and virtual eddies. Later versions (i.e., META3.x) adopted a contour-overlap strategy, linking detections across timesteps when their intersection exceeded a given threshold (Keppler et al. (2018); Laxenaire et al. (2018); Li et al. (2016); Pegliasco et al. (2015, 2022)).



This approach proved more robust, enabling consistent trajectories even during partial occlusion or deformation, and reducing the need for virtual eddies.

1.4 Eddy networks and interactions

Eddies were initially viewed as isolated coherent structures driven by baroclinic instabilities or topographic forcing (Stammer and Wunsch (1999)). However, observational and modeling evidence has revealed that mesoscale dynamics also involve frequent eddy–eddy interactions such as merging and splitting (Fang and Morrow (2003); Adcock and Marshall (2000); Trieling et al. (2005); Han et al. (2025); Wang et al. (2019)). These processes are now recognized as a fundamental component of oceanic energy transfer and water mass transport.

Recent algorithms have been developed to explicitly identify such interactions, treating eddies as nodes within a dynamic network rather than independent trajectories. Examples include AMEDA (Le Vu et al. (2018)), which integrates merging/splitting detection into tracking, the multicore Gaussian approach of Cui et al. (2019), the graph-based framework of ToEddies (Ioannou et al. (2024); Laxenaire et al. (2018, 2020)), and the topological tree approach of EddyGraph (Tian et al. (2021)). Together, these methods have demonstrated the potential of network-based descriptions to capture the full complexity of eddy life cycles.

Building upon this progress, we present the META4.0-Networks atlas, the new global version of the Mesoscale Eddy Trajectory Atlas distributed by AVISO+. META4.0 introduces eddy networks into the dataset, providing a unified representation of eddy evolution, merging, and splitting events. This update represents a key advancement for the META eddy products, ensuring continuity with previous versions while adding a new dimension for studying eddy connectivity and interactions. The construction of the networks follows three stages:

- Daily eddy detection from DT2021 ADT fields;
- Network grouping based on eddy overlap between consecutive days, accounting for missing detections;
- Network segmentation to identify merging and splitting events within each group.

Beyond the methodological improvement, the increasing complexity of network representations requires appropriate tools for data manipulation, visualization, and analysis. For this reason, the full Python source code is released under a GPL v3 license (<https://doi.org/10.5281/zenodo.7197432>; Delepoulle et al., 2022) and integrated into the open-source py-eddy-tracker (PET) library (Delepoulle et al. (2022); Mason et al. (2014); Pegliasco et al. (2022)). The toolbox is freely available and documented online (https://py-eddy-tracker.readthedocs.io/en/latest/python_module/index.html), providing ready-to-use functionalities for detecting eddies, analyzing trajectories, building networks, and reproducing the results presented in this paper.

This paper is organized as follows. Section 2 introduces the META4.0-Networks algorithm and describes the data sources used. Section 3 presents the toolbox and main validation diagnostics. Section 4 details data and code availability, and Section 5 concludes with a summary and perspectives.



2 Data and methods

2.1 Eddy detection

90 2.1.1 Altimetric fields for eddy detection

Following (Pegliasco et al. (2022)), the detection is performed on the absolute dynamic topography (ADT) field rather than on the sea level anomaly (SLA), as done in earlier META versions. Using ADT provides better detection in dynamically energetic regions characterized by strong sea surface height (SSH) gradients or recurrent mesoscale structures such as current retroflections, coastal eddies, and topographic recirculations. Many of these features are imprinted in the mean dynamic topography (MDT), and would therefore be partially masked in SLA fields. The ADT fields used here are the global daily products from the DT2021 reprocessing (Faugère et al. (2022)), with a $1/4^\circ$ grid resolution.

Unlike early approaches such as Chelton et al. (2011), which relied on the Okubo–Weiss parameter to isolate rotational structures from a non-rotative background, spatial filtering to the ADT fields is no longer applied. Filtering was originally introduced to reduce large-scale gradients and noise that affected vorticity-based methods. However, the contour-based approach adopted here is inherently less sensitive to background flow variations, as it directly identifies closed SSH contours enclosing extrema. In line with the methodology of Ioannou et al. (2024) (ToEddies), we therefore retain the full, unfiltered ADT signal, preserving the complete range of observed altimetric variability and ensuring a more faithful representation of mesoscale and submesoscale dynamics.

2.1.2 Detecting mesoscale eddies

105 The detection algorithm is based on the py-eddy-tracker (PET) software (Mason et al. (2014)), itself derived from the methods of Chelton et al. (2011); Kurian et al. (2011); Penven et al. (2005). It proceeds as follows:

1. SSH contour extraction: closed contours are traced around SSH extrema, downward from maxima for anticyclones, upward from minima for cyclones.
2. Validation tests: each contour must
 - 110 – contain a single extremum;
 - enclose between 4 and 1000 pixels;
 - exceed an SSH amplitude of 0.4 cm;
 - have a shape error (deviation from a best-fit circle) below 70%, ensuring realistic geometries while allowing elongated shapes.
- 115 3. Interpolation and diagnostics: validated contours and mean radial speed profiles are interpolated over 30 points using the visvalingam algorithm (Visvalingam (2016)), facilitating further analyses and data colocation (SST, color, salinity, Argo, etc.).



The resulting daily detections of anticyclonic and cyclonic eddies are stored separately in NetCDF files.

2.2 Network grouping

120 Network construction is performed separately for anticyclonic and cyclonic eddies. While cyclonic–anticyclonic interactions do exist (Amores et al. (2017); Chang and Park (2015); Shi and Nof (1993)), they are comparatively rare and involve complex dynamical processes. Restricting the analysis to eddies of the same polarity allows for a more robust and computationally tractable description of eddy interactions, while retaining the dominant merging and splitting dynamics observed globally.

125 The grouping step links daily detections into spatio-temporal networks based on their geometric overlap across consecutive days. Two eddies are connected if their normalized overlap area exceeds 10%. To account for occasional missing detections, the search is extended up to seven days forward, provided that eddies remain geographically consistent. No constraint is imposed on radius or amplitude variations, allowing the detection of rapid size changes associated with merging or splitting.

The process can be summarized as follows:

1. Preprocessing of files

- 130
- Input daily eddy detections (polygons) are sorted chronologically;
 - Missing days are checked to ensure temporal continuity.

2. Pairwise comparison across days

- 135
- For each day, all eddies are compared with those of subsequent days within a temporal window (i.e., the next 7 days);
 - Candidate pairs are first filtered using bounding box intersection to quickly eliminate eddies that are too far apart;
 - For the remaining pairs, the geometric overlap between polygons is computed. If the normalized overlap exceeds a threshold (i.e., 10%), the two eddies are considered connected.

3. Group construction

- 140
- Each set of linked detections defines a group (a temporal network of eddies);
 - If a new match connects two existing groups, they are merged into a single group.

Each eddy observation is thus assigned to a group, corresponding either to a simple trajectory or to a more complex network involving multiple interacting eddies. These steps are illustrated through dedicated Jupyter notebooks available in the documentation (https://py-eddy-tracker.readthedocs.io/en/latest/python_module/16_network/pet_group_anim.html#sphx-glr-python-module-16-network).

2.3 Network segmentation

145 The segmentation step decomposes each group into coherent trajectory segments bounded by an event. An event can be a birth, death, merging, or splitting, defining the start or end of one or several eddy segments. Figure 1 illustrates an example



of a network composed of three segments connected through two successive events. The first segment is composed of four observations represented by the dots. It ends by merging with the second observations of segment 2 resulting in a larger eddy belonging to segment 2. The fifth observation of this segment splits and give birth to the third segment.




Figure 1. Example of a network composed of three segments linked by two successive events (merging and splitting).

150 The segmentation algorithm proceeds as follows:

1. Initialization

- Each observation in a group is assigned an identifier and a time index;
- Links to potential predecessors and successors are initialized.

2. Forward and backward association

- 155
- Starting from an observation, the algorithm searches for its most likely successor in the following days (up to a window of tolerance for missing detections), based on maximal polygon overlap;
 - Similarly, the best predecessor is searched in earlier days;
 - This creates chains of linked observations, each chain defining a segment (trajectory).

3. Event handling

- 160
- Birth: observations without a predecessor → start of a new segment;
 - Death: observations without a successor → end of a segment;
 - Splitting: one observation links to multiple successors → trajectory bifurcation into several segments;
 - Merging: multiple observations link to the same successor → trajectories converge, one segment continues while others terminate;



- 165 – Segment reindexing: when an interaction occurs, segments involved in merging or splitting events are reindexed to ensure a consistent identification of the main trajectory. The main segment is selected based on continuity and overlap criteria, rather than on the order in which observations are processed.

4. Postprocessing

- 170 – Short segments (i.e., less than 3 observations) that don't connect several segments are removed from their networks in order to prevent event detection mistakes.

5. Output

- Each group is segmented into individual trajectories, capturing the full life cycle of eddies, including births, deaths, splits, and merges.

175 It should be noted that the segmentation algorithm does not impose any restriction on the number of eddies involved in merging or splitting events. In principle, more than two eddies may merge, and a single eddy may split into more than two offspring eddies.

 The segmentation procedure is illustrated in the PET documentation notebooks (https://py-eddy-tracker.readthedocs.io/en/latest/python_module/16_network/pet_segmentation_anim.html#sphx-glr-python-module-16-network-pet-segmentation-anim-py), which provide visual examples and interactive tools for reproducing this step.

180 3 Results

 In this section, the results of eddy networks detections and characterizations are presented. First, the diagnostic tools available for exploring individual networks are introduced, before moving to a global statistical analysis over three decades of data. Then, a focus is made on the dynamics of merging and splitting events, studying their spatial and temporal distributions, as well as the eddies properties during such events, and finally qualitatively validate the results with independent oceanographic
185 observations (i.e., chlorophyll products). Eventually, the typology of networks are explored through clustering.

3.1 Exploratory tools for network analysis

 In this section, the capabilities of the `pyeddytracker` toolbox are illustrated through a representative case study. This local example highlights some of the graphical diagnostics available to investigate the structure and dynamics of individual eddy networks.

190 A single network was selected from the META4.0 anticyclonic networks atlas to demonstrate the diagnostic possibilities. Networks can be extracted using flexible selection tools:

- Spatio-temporal criteria (e.g., specifying a time window and/or longitude–latitude box);
- Intrinsic characteristics such as network size or lifetime;



- Targeted searches, where the user tracks the history of a specific eddy of interest.

195 The chosen network characteristics are summarized in Table 1.

Characteristics	Number of observations	Number of segments	Number of merging events	Number of splitting events	Lifetime (days)	Longitude/latitude range
Values	601	6	3	2	445	(92.15°;106.90°) (-37.14°;-30.15°)

Table 1. Internal characteristics of the demo network.

The temporal evolution of the network is represented as timelines in Figure 2. Timelines can be ordered either by segment identifiers or by latitude, both as a function of time. Merging and splitting events are symbolized by black hexagons and stars, respectively. Such representations are useful for quickly assessing the internal complexity of a network: the number of coexisting segments, their relative durations, and the timing of interaction events.


200 To complement this view, the geographical trajectories of the segments are shown in Figure 3. In this map-based representation, the merging and splitting events are explicitly localized, allowing for a joint spatial and temporal understanding of network dynamics.

Timelines can also be enriched with eddy properties such as effective radius or shape error (i.e., quantifying the deviation from an idealized circular eddy), as illustrated in Figure 4. On this timeline, each dot corresponds to an observed eddy. 205 Marker color indicates the shape error (i.e., ellipticity), while marker size is proportional to the effective radius. This combined representation directly links the temporal organization of eddy segments to the evolution of their physical characteristics.


For example, Figure 4 reveals that the effective radius tends to increase after merging events and before splitting events, suggesting a reorganization of the eddy structure when mass and energy are redistributed. At the same time, the timeline indicates that eddies become more distorted around interaction events, corresponding to increased dynamic stress during merging 210 or division. Together, these diagnostics provide physically interpretable signatures of merging and splitting: growth in size accompanied by shape deformation is a strong indicator of structural reorganization. Similar diagnostics could also be performed with other eddy properties such as amplitude or mean translation speed, however, they are omitted here for conciseness.

Overall, these diagnostics can be particularly informative for:

- Detecting structural changes in eddy properties (e.g., increase in radius, deformation) around merging and splitting events;
- Tracing the continuity of segments and checking whether apparent gaps in detection correspond to real interruptions or short-lived tracking issues;
- Visually linking eddy trajectories with interaction events to assess the temporal ordering of merging and splitting within a network.




(a) Demo network timeline function of the segments' orders.



(b) Demo network timeline function of latitude.

Figure 2. Timelines of the demo network.



(a) Segments' tracks and one example of merging event.

(b) Segments' tracks and one example of splitting event.

Figure 3. Demo networks segments' tracks and merging and splitting events.




Figure 4. Timeline of eddy trajectories. Marker color indicates shape error, while marker size is proportional to eddy radius.

220 These exploratory diagnostics therefore provide a practical toolkit for investigating the structure and dynamics of individual eddy networks. While they are demonstrated here on a single representative case, the tendencies observed will be assessed and statistically confirmed at the global scale in the following sections.

3.2 Global statistics

3.2.1 General information

225 The main properties of the META4.0-Networks dataset are summarized in Table 2. These numbers provide a first overview of the database, including the total number of eddy observations, the number of reconstructed networks and segments, and the occurrence of merging and splitting events. For completeness, the number of lonely eddies (i.e., eddies that were never associated with any network) is also reported.

Properties	Number of observations	Number of networks	Number of segments	Number of merging events	Number of splitting events	Number of lonely eddies
Values	60,691,435	1,939,458	2,281,543	188,260	186,610	997,977

Table 2. Global characteristics of the META4.0-Networks dataset.

230 The lonely eddies correspond to individual eddies that could not be attached to any trajectory or network. Their spatial distribution is shown in Figure 5, expressed as the number of lonely eddy detections per year and per degree², in 0.25° boxes. This diagnostic reveals that lonely eddies occur throughout the global ocean, but with enhanced frequencies in specific regions: along continental slopes and coasts (e.g., the Indonesian seas), at high and low latitudes (e.g., the Bering Strait), and within the equatorial band (e.g., the western Pacific). These patterns are consistent with areas where mesoscale detection is more challenging, either because of energetic small-scale variability, proximity to land, or reduced altimetric resolution.




Figure 5. Frequency of lonely eddies expressed as number of observations per year and per degree².

235 To further document the distribution of networks, Figure 6 presents histograms of (a) the number of observations per network and (b) the number of segments per network. These two metrics quantify the internal complexity of networks and provide insight into the range of eddy interactions captured in the dataset.




Figure 6. Histograms of META4.0 networks: (a) number of observations and (b) number of segments.

Figures 6a and 6b indicate that most networks remain relatively simple, composed of a limited number of observations and segments. Nevertheless, the dataset also contains a non-negligible fraction of very large networks, with more than 20,000



240 observations and over 300 segments. This wide range illustrates the strong variability of eddy life cycles and the added value of the network representation in capturing both simple, isolated structures and highly interactive, long-lived systems.

3.2.2 Comparison with the META3.2 single-trajectory dataset

In this section, the META4.0 network-based atlas is compared with the META3.2 single-trajectory dataset.

245 Figures 7 and 8 display frequency maps of birth and death events, respectively. In both cases, the single-trajectory dataset shows a larger number of events than the network-based approach. This difference arises because, in the single-trajectory framework, each merging or splitting tends to artificially create new births and deaths, whereas the network formalism connects these trajectories into a continuous system.




Figure 7. Comparison of birth event frequencies between META4.0 networks and META3.2 single trajectories.





Figure 8. Comparison of death event frequencies between META4.0 networks and META3.2 single trajectories.

250 As both datasets are derived from the same set of eddy observations, this comparison highlights the main strength of the network approach: it reduces artificial discontinuities and allows for a more consistent tracking of eddy lifecycles. In practice, the META4.0 networks extend the effective lifetime of many trajectories compared with META3.2, while preserving the underlying physical signal.



To quantify these differences, we use Complementary Cumulative Distribution Functions (CCDF). The CCDF represents the fraction of eddies surviving longer than a given lifetime. This diagnostic avoids binning effects and highlights differences in the tail of the distributions. Figure 9a compares the META3.2 lifetime CCDF with the META4.0 lifetime CCDF of segments, while Figure 9b compares META3.2 with the META4.0 CCDF of whole networks. Both comparisons show that network segments are statistically longer than single trajectories, and that networks themselves capture even longer lifetimes by connecting segments across merging and splitting events.



(a) Lifetime CCDF of META3.2 single-trajectories and (b) Lifetime CCDF of META3.2 single-trajectories and META4.0 networks' segments.

Figure 9. Comparison of lifetime CCDF between META4.0 networks' segments and networks and META3.2 single trajectories.

To further illustrate these results, the spatial distribution of long-lived trajectories is also compared. Figure 10 shows the trajectories of eddies persisting longer than one year (365 days) for both datasets.

META4.0 detects and connects a larger number of eddies above this threshold compared with META3.2, especially in regions of intense mesoscale activity such as the western boundary currents and the Southern Ocean. This visual comparison confirms that the network approach enhances the detection of persistent eddy structures and reduces artificial fragmentation.

Overall, these diagnostics demonstrate that the network-based tracking extends eddy lifetimes beyond what is observed with isolated trajectories, providing a more robust representation of mesoscale eddy dynamics.




Figure 10. Comparison of eddy trajectories persisting longer than 365 days in META3.2 and META4.0. META4.0 exhibits more long-lived trajectories than the single-trajectory dataset, particularly in regions of strong eddy activity.

3.3 Merging and splitting events

This section analyzes the characteristics of merging and splitting events identified with the network-based approach.

3.3.1 Spatial distribution of interaction events

Figure 11 shows frequency maps of merging and splitting events. These maps are computed as the fraction of pixels covered by eddy contours involved in merging or splitting. This approach emphasizes the spatial regions where such interactions most frequently occur.




Figure 11. Frequency maps of merging and splitting events.

Both maps confirm that interactions are concentrated in high-energy regions, such as western boundary currents and the equatorial Pacific. The spatial signatures of merging and splitting are broadly similar, suggesting that both processes tend to occur in dynamically active areas where eddy–eddy interactions are frequent.



275 3.3.2 Temporal variability

Figure 12 shows the temporal evolution of the number of merging and splitting events over the 30-year record, with four points per year.




Figure 12. Temporal evolution of merging and splitting events.

The number of events remains relatively stable, between 1300 and 1700 per trimester, without evidence of long-term trends. Abrupt variations mainly coincide with changes in the altimeter constellation and the associated observing system, as described
 280 in (Pegliasco et al., 2022).

3.3.3 Eddy properties around interaction events

To better understand how merging and splitting unfold, eddy properties are analyzed before and after interaction events. Figures 13 and 14 show the median and interquartile range of normalized effective radius and shape error in the days surrounding merging and splitting events, respectively. Eddy properties are normalized by the maximum value reached by each eddy over
 285 its lifetime. This normalization allows comparison of the temporal evolution of properties within an eddy's life cycle, but does not preserve absolute size or amplitude relationships between interacting eddies. As a result, relative values before and after interactions should be interpreted as indicators of the eddy's stage within its own life cycle, rather than as direct physical comparisons between parent and resulting eddies.




Figure 13. Median and interquartile range of eddy normalized shape error and effective radius around merging events.




Figure 14. Median and interquartile range of eddy normalized shape error and effective radius around splitting events.

The results reveal systematic and consistent patterns when interpreted in a relative, life-cycle framework:

- **Merging events:** prior to merging, parent eddies exhibit increasing shape distortion while remaining relatively far from their maximum size within their own life cycle. Following the interaction, the resulting eddy rapidly reaches a large fraction of its lifetime maximum radius and subsequently evolves toward a more stable, less distorted configuration.
- **Splitting events:** eddies undergoing splitting tend to approach their maximum relative size and deformation shortly before the event. After splitting, the resulting eddies begin their evolution at relatively high normalized radii and shape errors, indicating that they inherit dynamically active structures which reorganize over the following days.



Because the diagnostics are expressed in normalized units, these figures do not quantify absolute size or amplitude transfers during interactions. Absolute radius and shape errors diagnostics (not shown for synthesis purpose) confirm that merging leads to larger structures and splitting to smaller ones. The normalized analysis presented here instead highlights the timing of interaction events within the relative life cycle of eddies and their associated transient deformations.

300 These tendencies generalize the case-study diagnostics shown in Figure 4, providing robust statistical evidence that merging and splitting are associated with transient structural deformation followed by a reorganization into more stable eddies.

3.3.4 Interaction events in the eddy lifecycle

To put these results into perspective, Figure 15, inspired from Dong et al. (2012), illustrates the evolution of eddy normalized effective radius and shape errors along their normalized lifetime in the META 3.2 single-trajectory framework. This classical
 305 view highlights that eddies typically undergo an initial adjustment phase after their birth, followed by a mature stage with nearly stable properties, and finally a weakening phase characterized by decreasing radius and increasing deformation before death.





Figure 15. Evolution of normalized effective radii and deformation along normalized lifetime for the META3.2 single-trajectory framework.

The META4.0 network-based diagnostics confirm these birth–maturity–decay patterns by reproducing the statistical signatures of eddy properties around birth and death events, as shown on Figures 16 and 17. At birth, eddies are smaller and deformed before stabilizing within weeks, while at death, they decay in size and lose coherence, consistent with Dong et al.
 310 (2012).


Beyond this canonical picture, however, the network framework reveals intermediate reorganizations that are absent from the single-trajectory view. The diagnostics of merging and splitting (Figures 13 and 14) demonstrate that long-lived eddies are not only shaped by their formation and decay, but also by episodic interactions that transiently alter their structure before they
 315 restabilize.

Together, these results show that merging and splitting are frequent and robust features of mesoscale dynamics: they occur in energetic regions, at stable rates, and are systematically associated with transient deformation followed by reorganization. By



(a) Evolution of eddy shape errors around birth events. (b) Evolution of eddy effective radii around birth events.

Figure 16. Evolution (median and IQR) of eddy shape error and effective radius around birth events in META4.0.



(a) Evolution of eddy properties around birth events. (b) Evolution of eddy properties around death events.

Figure 17. Evolution (median and IQR) of eddy shape error and effective radius around death events in META4.0.

embedding interaction events into the classical birth–maturity–decay framework, META4.0 provides a more complete picture of the eddy lifecycle.

320 3.3.5 Qualitative validation with external observations

Finally, a qualitative validation of merging and splitting detections is provided using independent chlorophyll concentration (CHL) fields. The CLS CatSat chlorophyll product is used to cross-check the occurrence of interaction events. One representative example of each interaction type is shown.

Figure 18 illustrates a merging event observed both in the ADT field (Figure 18a) and in the CHL field (Figure 18b), across
 325 four dates capturing the situation before, during, and after the event. Similarly, Figure 19 presents a splitting event documented in both ADT (Figure 19a) and CHL (Figure 19b). For each case, the corresponding network timelines are also displayed, providing additional confirmation of the detected interactions.




Figure 18. Merging event observed in both ADT and CHL fields (between purple and light green eddies).




Figure 19. Splitting event observed in both ADT and CHL fields (purple and light green eddies).

These examples confirm that merging and splitting events detected from altimetry are also visible in independent fields, reinforcing the robustness of the methodology. However, conducting systematic detections at the global scale from CHL fields remains challenging due to their high sensitivity to factors such as cloud cover and atmospheric corrections.



3.4 Clustering and typology of networks

This section explores whether mesoscale eddy networks can be grouped into a limited number of recurrent types based on their main properties. The objective is to identify regions of the global ocean where networks share similar structural and dynamical characteristics, thus providing a typology of mesoscale organization. This investigation is performed using a clustering analysis based on some chosen network attributes, as follows:

- The number of observations;
- The number of segments;
- The mean lifetime of segments within a network;
- The mean effective radius of eddies within a network;
- The spatial extent of networks, computed as the maximum distance between two eddies in the same network.

Different clustering algorithms and parameterizations were tested (e.g., hierarchical clustering, Gaussian Mixture Models). The *k*-Means algorithm (Lloyd (1982)) with five clusters was retained for its convergence speed, accuracy and robustness. For the sake of conciseness, this exploratory phase is not reported here.

Figure 20 shows the spatial frequency of networks assigned to each cluster. To better interpret the properties of these groups, Figure 21 presents a radar chart of their mean attributes, while Figure 22 displays the distribution of each attribute of the five clusters.

The five groups can be interpreted as distinct dynamical regimes of mesoscale organization:

- **Group 1:** Small networks of small eddies, distributed nearly everywhere. These networks likely represent the background turbulent field, with limited impact on large-scale transport.
- **Group 2:** Small networks of relatively large eddies, concentrated in the equatorial band. This is consistent with equatorial dynamics, where Rossby waves and instabilities generate large but weakly connected eddies.
- **Group 3:** Small networks of long-lived, large eddies, found in low-energy mid-latitudes. These networks reflect solitary, persistent eddies that act as local reservoirs of water masses rather than active transporters.
- **Group 4:** Medium-size networks of large eddies, mainly located in high-energy mid-latitude regions such as western boundary currents. These zones of strong baroclinic and barotropic instabilities foster frequent eddy interactions and exchanges.
- **Group 5:** The largest and longest-lived networks, concentrated in the Agulhas, the Antarctic Circumpolar Current, and other major currents (but not in the Gulf Stream which appear in the fourth group). These networks correspond to highly connected systems with strong transport capacity across ocean basins.




Figure 20. Spatial frequency of networks assigned to each cluster.




Figure 21. Radar chart of mean attributes for the five clusters.




Figure 22. Box plots of the attributes for the five network clusters.

360 This typology extends the single-eddy perspective of previous atlases (e.g., META3.2) by explicitly considering the collec-
 tive organization of eddies into networks. Rather than only identifying where eddies are dense, the clustering highlights where
 eddies evolve as isolated, long-lived structures versus where they form complex, interacting systems. From a dynamical point
 of view, the results distinguish between regions dominated by transient, localized structures (Groups 1–2), zones of persistent
 solitary eddies (Group 3), regions of intense mesoscale interactions (Group 4), and major corridors of inter-basin transport
 365 (Group 5).

Overall, the clustering highlights the heterogeneous roles of eddy networks in ocean circulation: from local stirring and
 storage to long-distance connectivity and redistribution of water masses.

370 In the next section, Lagrangian particle advection is used to complement the spatial clustering analysis by providing a
 dynamical assessment of water mass transport and retention within eddy networks.

3.5 Particle advection and network coherence

The coherence of eddy segments and their associated networks is assessed using Lagrangian particle advection experiments.
 Virtual particles are advected using surface geostrophic velocities derived from the same altimetric sea level anomaly (ADT)
 product employed for eddy detection. An eddy (or eddy segment) is considered coherent when a significant fraction of the
 375 particles initially enclosed within its contour remain trapped within coherent eddy structures over time, whereas it is deemed
 incoherent when particles rapidly escape the eddy boundaries.



To perform this assessment, particles are uniformly initialized at a spatial resolution of $1/50^\circ$ within the eddy speed contour at a reference date d . Particle trajectories are then integrated both forward and backward in time over a 7-day window. At $d \pm 7$ days, particle positions are analyzed and assigned to the detected eddies at those dates. The coherence score is defined as the sum of the fractions of particles contained within the two eddy effective contours gathering the largest proportions of the initial particle population. This metric provides a quantitative estimate of the retentive capacity of individual segments and their associated networks, expressed as a percentage of retained particles.

Performing particle advection both forward and backward in time allows to characterize the directionality of coherence and to assess the intrinsic irreversibility of eddy interactions. Forward advection evaluates the ability of eddies to retain particles in the future, while backward advection measures the extent to which particles observed within a given eddy originate from a coherent structure in the past. Differences between forward and backward coherence therefore provide insight into the irreversible nature of eddy dynamics and particle transport.

Figure 23 presents the raw and cumulative distributions of segment autocohérence scores for anticyclonic and cyclonic eddy networks. In both atlases, the raw distributions exhibit a pronounced maximum at high autocohérence values (around 90%), indicating that most detected segments behave as strongly Lagrangian coherent structures. In particular, more than 50% of the segments display autocohérence scores exceeding 75%, which supports the robustness of the segmentation procedure and highlights the persistence of coherent transport pathways within eddy networks.




Figure 23. Raw and cumulative distributions of segment autocohérence scores.

Interaction events are further evaluated using the same framework, by computing the fraction of particles that remain involved in merging or splitting processes, as opposed to escaping from interacting structures.



As described in Figure 24a, a merging event is characterized by a minor eddy M_2 merging into a dominant eddy M_1 , resulting in a single eddy M that follows the trajectory of M_1 , based on overlapping criteria. Conversely, a splitting event is modeled as an initial eddy S separating into a main eddy S_1 , which retains the original trajectory, and a secondary eddy S_2 .

Figure 24b (respectively Figure 24c) shows the cumulative distributions of coherence scores for merging (respectively splitting) events.




Figure 24. Cumulative distributions of coherence scores for merging and splitting events.

Forward advection reveals a strong coherence of merging processes, as particles initially enclosed within interacting eddies remain largely confined within the merged structure. In contrast, backward advection exhibits significantly lower coherence, reflecting the irreversible mixing of particle trajectories once the eddies have merged. More than 60% of detected merging events display forward coherence scores above 80% for the dominant eddy M_1 , whereas the secondary eddy M_2 generally exhibits lower coherence, with only about 40% exceeding the same threshold. This asymmetry likely reflects the weaker dynamical integrity of M_2 , which is typically smaller and more rapidly distorted during the merging process.



Conversely, Figure 24c shows that splitting events display higher backward autocohere-
410 nce. Backward integration highlights the convergence of particle trajectories toward a single coherent structure in the past, while forward advection results in a dispersion of particles among multiple offspring eddies. More than 60% of detected splitting events exhibit backward coherence scores above 80% for the main eddy S_1 . The secondary eddy S_2 shows systematically lower coherence, with only about 35% of events exceeding this threshold, consistent with its more recent formation and reduced ability to retain particles.

Overall, these diagnostics demonstrate that most detected interaction events exhibit significant Lagrangian coherence, thereby validating the identification of segments, merging and splitting processes. They further provide a quantitative characterization
415 of particle transport and redistribution within eddy networks, contributing to a refined description of mesoscale connectivity.

It should be noted, however, that the velocity fields used for particle advection are derived from the same altimetric product as that employed for eddy detection. As a result, the coherence assessment is not fully independent of the detection methodology, unlike diagnostics based on external tracers such as chlorophyll or sea surface temperature. Nevertheless, the particle-based analysis remains semi-independent, as it probes the Lagrangian behavior of the flow rather than the Eulerian properties of the
420 eddy contours themselves.

Additional limitations include the finite 7-day advection window, which may miss shorter-lived interactions or intermediate events occurring within this period. Despite these limitations, the consistency of the results across multiple diagnostics supports the robustness of the proposed framework.

Finally, tools to perform Lagrangian particle advection analyses on individual eddy networks are provided within the
425 `pyeddytracker` toolbox (https://py-eddy-tracker.readthedocs.io/en/v3.6.1/python_module/16_network/pet_follow_particle.html#sphx-glr-python-module-16-network-pet-follow-particle-py). These tools allow, for instance, the advection of particles within a given network, the visualization of their trajectories, and the analysis of interaction events along their temporal evolution.

4 Code and datasets availability

430 The detection, tracking and networking algorithms have been freely released under a GPL V3 license (<https://doi.org/10.5281/zenodo.6333989>;Delepoulleetal.,2022) in the Python language.

The META4.0-Networks delayed-time (DT) dataset is available in an AVISO repository at DOI 10.24400/527896/a01-2026.001 (CLS and CNES (2026)). An associated handbook describes the atlas as well as the variables stored in the NetCDF files and gives an insight into data manipulation. A large gallery of illustrated and documented routines to help with the data
435 manipulation and visualization is also available, and is updated when new methods are developed (<https://py-eddy-tracker.readthedocs.io/en/v3.3.1/py-eddy-tracker,2022>)).



5 Conclusions

In this article, the new global Mesoscale Eddy Trajectory Atlas META4.0-Networks has been introduced. The dataset was computed using the `pyeddytracker` framework, developed in Python, and the routines for data manipulation and visualization have been fully documented and made publicly available.

The main novelty of META4.0 lies in the introduction of eddy networks as a new framework for eddy tracking. Unlike the previous single-trajectory strategy, the network-based approach enables the identification of merging and splitting events and organizes trajectories into segments belonging to larger interacting structures. In this way, networks group together all eddies linked through interaction events. This representation adds a new layer of complexity to eddy tracking and allows for a more realistic description of eddy life cycles, explicitly accounting for their interactions and temporal connectivity.

A series of exploratory diagnostics was conducted to assess the robustness and relevance of the dataset. Global statistics and a comparison with the META3.2 single-trajectory atlas demonstrated that the network approach extends eddy lifetimes and refines the reconstruction of their pathways. Merging and splitting events were characterized in terms of their spatial distribution, temporal variability, and impact on eddy properties throughout their life cycles. These analyses revealed that interaction events play a fundamental role in shaping eddy evolution, a dimension that cannot be captured within a single-trajectory framework. Indeed, the results highlight that merging and splitting are frequent and robust features of mesoscale dynamics that occur in energetic regions, at stable rates, and are systematically associated with transient deformation followed by reorganization. A qualitative validation using independent chlorophyll concentration data further confirmed the realism of the detected interactions for selected case studies. By adding interaction events into the classical birth–maturity–decay framework, META4.0 provides a more complete picture of the eddy lifecycle.

Beyond individual events, the network representation provides a natural framework to investigate the collective organization of mesoscale eddies. A clustering analysis identified five distinct typologies of eddy networks, revealing contrasted regimes of ocean circulation ranging from localized, short-lived structures to highly connected systems associated with major current regions. These results highlight the emergence of large-scale connectivity patterns resulting from repeated eddy interactions.

Altogether, the results demonstrate that network-based tracking constitutes a significant advance in the description of mesoscale dynamics. By extending the representation beyond isolated trajectories, META4.0-Networks enables a more comprehensive reconstruction of eddy origins, pathways, interactions, and cross-basin connectivity. The Lagrangian diagnostics further demonstrate that eddy interactions are intrinsically irreversible processes, with distinct signatures in forward and backward particle transport. Beyond this dynamical interpretation, the consistency of these signatures with the expected behavior of merging and splitting events provides a coherent validation of the detected interactions and of the network-based representation. Although the advection relies on the same velocity fields as those used for eddy detection, these results offer a physically meaningful assessment of the realism of the reconstructed networks.

Future developments will include the application of the detection and tracking methods to next-generation sea surface height products currently under preparation, enabling eddy identification at higher spatial and temporal resolution. In addition, collocations with temperature and salinity Argo float profiles will allow the reconstruction of the three-dimensional structure of eddies,



thereby enriching the current datasets and enabling a more complete characterization of their dynamical and biogeochemical impacts. Ultimately, the combination of eddy networks with in situ observations and high-resolution numerical models opens new perspectives to quantify mesoscale connectivity and its role in the oceanic system.

Author contributions. Conceptualization: AD, JG, FN, MIP. Data curation: AD, JG. Formal analysis: JG, FN, AD. Funding acquisition: MIP, GD. Investigation: JG, FN, AD. Methodology : AD, JG, FN. Project administration: MIP, GD. Resources: MIP, GD. Software: AD, JG, FN. Supervision: MIP, GD. Validation: JG, FN, AD, MIP, GD. Visualization: JG, FN, AD. Writing (original draft preparation): JG, AD, FN.

Competing interests. The contact author has declared that none of the authors has any competing interests.



References

- Abernathy, R. and Haller, G.: Transport by Lagrangian Vortices in the Eastern Pacific, *Journal of Physical Oceanography*, 48, 667–685, <https://doi.org/10.1175/JPO-D-17-0102.1>, 2018.
- Adcock, S. T. and Marshall, D. P.: Interactions between geostrophic eddies and the mean circulation over large-scale bottom topography, *Journal of Physical Oceanography*, 30, 3223–3238, [https://doi.org/10.1175/1520-0485\(2000\)030<3223:IBGEAT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2000)030<3223:IBGEAT>2.0.CO;2), 2000.
- Amores, A., Monserrat, S., Melnichenko, O., and Maximenko, N.: On the shape of sea level anomaly signal on periphery of mesoscale ocean eddies, *Geophysical Research Letters*, 44, 6926–6932, <https://doi.org/10.1002/2017GL073978>, 2017.
- Beal, L. M., de Ruijter, W. P. M., Biastoch, A., Zahn, R., Cronin, M., Hermes, J., Lutjeharms, J., Quartly, G., Tozuka, T., Baker-Yeboah, S., Bornman, T., Cipollini, P., Dijkstra, H., Hall, I., Park, W., Peeters, F., Penven, P., Ridderinkhof, H., and Zinke, J.: On the role of the Agulhas system in ocean circulation and climate, *Nature*, 472, 429–436, <https://doi.org/10.1038/nature09983>, 2011.
- Beron-Vera, F. J., Olascoaga, M. J., and Goni, G. J.: Oceanic mesoscale eddies as revealed by Lagrangian coherent structures, *Geophysical Research Letters*, 35, L12 603, <https://doi.org/10.1029/2008GL033957>, 2008.
- Brach, L., Deixonne, P., Bernard, M.-F., Durand, E., Desjean, M.-C., Perez, E., van Sebille, E., and ter Halle, A.: Anticyclonic eddies increase accumulation of microplastic in the North Atlantic subtropical gyre, *Marine Pollution Bulletin*, 126, 191–196, <https://doi.org/10.1016/j.marpolbul.2017.10.077>, 2018.
- Braun, C. D., Gaube, P., Sinclair-Taylor, T. H., Skomal, G. B., and Thorrold, S. R.: Mesoscale eddies release pelagic sharks from thermal constraints to foraging in the ocean twilight zone, *Proceedings of the National Academy of Sciences*, 116, 17 187–17 192, <https://doi.org/10.1073/pnas.1903067116>, 2019.
- Chaigneau, A., Gizolme, A., and Grados, C.: Mesoscale eddies off Peru in altimeter records: Identification algorithms and eddy spatio-temporal patterns, *Progress in Oceanography*, 79, 106–119, <https://doi.org/10.1016/j.pocean.2008.10.013>, 2008.
- Chaigneau, A., Eldin, G., and Dewitte, B.: Eddy activity in the four major upwelling systems from satellite altimetry (1992–2007), *Progress in Oceanography*, 83, 117–123, <https://doi.org/10.1016/j.pocean.2009.07.012>, 2009.
- Chaigneau, A., Le Texier, M., Eldin, G., Grados, C., and Pizarro, O.: Vertical structure of mesoscale eddies in the eastern South Pacific Ocean, *Journal of Geophysical Research: Oceans*, 116, C11 025, <https://doi.org/10.1029/2011JC007134>, 2011.
- Chambault, P., Baudena, A., Bjørndal, K. A., Santos, M. A. R., Bolten, A. B., and Vandeperre, F.: Swirling in the ocean: Immature loggerhead turtles seasonally target old anticyclonic eddies at the fringe of the North Atlantic gyre, *Progress in Oceanography*, 175, 345–358, <https://doi.org/10.1016/j.pocean.2019.05.005>, 2019.
- Chang, Y. S. and Park, Y. G.: Variation of flow properties during a collision event of two mesoscale eddies in the Gulf Stream region from numerical simulation, *Ocean Science Journal*, 50, 567–579, <https://doi.org/10.1007/s12601-015-0052-y>, 2015.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91, 167–216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- Christie, M. R., Tissot, B. N., Albins, M. A., Beets, J. P., Jia, Y., Ortiz, D. M., Thompson, S. E., and Hixon, M. A.: Larval connectivity in an effective network of marine protected areas, *PLOS ONE*, 5, e15 715, <https://doi.org/10.1371/journal.pone.0015715>, 2010.
- CLS and CNES: Meta4.0 DT allsat, <https://doi.org/10.24400/527896/a01-2026.001>, 2026.
- Cui, W., Wang, W., Zhang, J., and Yang, J.: Multicore structures and the splitting and merging of eddies in global oceans from satellite altimeter data, *Ocean Science*, 15, 413–430, <https://doi.org/10.5194/os-15-413-2019>, 2019.



- Delepoulle, A., evanmason, Clément, CoriPegliasco, Capet, A., Troupin, C., and Koldunov, N.: AntSimi/py-eddy-tracker: v3.6.1,
 515 <https://doi.org/10.5281/zenodo.6333989>, 2022.
- Dong, C., Lin, Y., Liu, F., Nencioli, F., Chao, Y., Guan, Y., Chen, D., Dickey, T., and McWilliams, J.-C.: Three-dimensional oceanic
 eddy analysis in the Southern California Bight from a numerical product, *Journal of Geophysical Research: Oceans*, 117, C00H14,
<https://doi.org/10.1029/2011JC007354>, 2012.
- Dong, C., Liu, L., Nencioli, F., Bethel, B. J., Liu, Y., Xu, G., and Zou, B.: The near-global ocean mesoscale eddy atmospheric-oceanic-
 520 biological interaction observational dataset, *Scientific Data*, 9, 436, <https://doi.org/10.1038/s41597-022-01550-9>, 2022.
- El Aouni, A.: A hybrid identification and tracking of Lagrangian mesoscale eddies, *Physics of Fluids*, 33, 036604,
<https://doi.org/10.1063/5.0038761>, 2021.
- Faghmous, J. H., Le, M., Uluyol, M., Kumar, V., and Chatterjee, S. B.: A parameter-free spatio-temporal pattern mining model
 to catalog global ocean dynamics, in: *Proceedings of the IEEE International Conference on Data Mining*, pp. 151–160,
 525 <https://doi.org/10.1109/ICDM.2013.162>, 2013.
- Fang, F. and Morrow, R.: Evolution, movement and decay of warm-core Leeuwin Current eddies, *Deep Sea Research Part II: Topical Studies
 in Oceanography*, 50, 2245–2261, [https://doi.org/10.1016/S0967-0645\(03\)00055-9](https://doi.org/10.1016/S0967-0645(03)00055-9), 2003.
- Faugère, Y., Taburet, G., Ballarotta, M., Pujol, I., Legeais, J. F., Maillard, G., and Dibarboure, G.: DUACS DT2021: 28 years of reprocessed
 sea level altimetry products, *EGU General Assembly Conference Abstracts*, pp. EGU22–7479, <https://doi.org/10.5194/egusphere-egu22-7479>,
 530 7479, 2022.
- Gaube, P., McGillicuddy, D. J., Chelton, D. B., Behrenfeld, M. J., and Strutton, P. G.: Regional variations in the influence of mesoscale eddies
 on near-surface chlorophyll, *Journal of Geophysical Research: Oceans*, 119, 8195–8220, <https://doi.org/10.1002/2014JC010111>, 2014.
- Gilchrist, R. M., Hall, R. A., Bacon, J. C., Rees, J. M., and Graham, J. A.: Increased dispersion of oil from a deep water seabed release by
 energetic mesoscale eddies, *Marine Pollution Bulletin*, 156, 111258, <https://doi.org/10.1016/j.marpolbul.2020.111258>, 2020.
- 535 Gruber, N., Lachkar, Z., Frenzel, H., Marchesiello, P., Münnich, M., McWilliams, J. C., Nagai, T., and Plattner, G.-K.: Eddy-induced reduction
 of biological production in eastern boundary upwelling systems, *Nature Geoscience*, 4, 787–792, <https://doi.org/10.1038/ngeo1273>,
 2011.
- Haller, G.: Dynamic rotation and stretch tensors from a dynamic polar decomposition, *Journal of the Mechanics and Physics of Solids*, 86,
 70–93, <https://doi.org/10.1016/j.jmps.2015.10.002>, 2016.
- 540 Han, G., Zhu, W., Wang, Y., Lin, X., Xie, J., and Liu, Y.: The Role of Eddy Merging in Maintaining the Quasi-Permanent Ulleung Eddy,
Deep Sea Research Part I: Oceanographic Research Papers, p. 104542, <https://doi.org/10.1016/j.dsr.2025.104542>, 2025.
- Ioannou, A., Stegner, A., Le Vu, B., Taupier-Letage, I., and Speich, S.: Dynamical Evolution of Intense Ierapetra Eddies on a 22 Year Long
 Period, *Journal of Geophysical Research: Oceans*, 122, 9276–9298, <https://doi.org/10.1002/2017JC013158>, 2017.
- Ioannou, A., Guez, L., Laxenaire, R., and Speich, S.: Global assessment of mesoscale eddies with TOEddies: comparison between multiple
 545 datasets and colocation with in situ measurements, *Remote Sensing*, 16, 4336, <https://doi.org/10.3390/rs16224336>, 2024.
- Isern-Fontanet, J., García-Ladona, E., and Font, J.: Identification of Marine Eddies from Altimetric Maps, *Journal of Atmospheric and
 Oceanic Technology*, 20, 772–778, [https://doi.org/10.1175/1520-0426\(2003\)20<772:IOMEFA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2003)20<772:IOMEFA>2.0.CO;2), 2003.
- Keppler, L., Cravatte, S., Chaigneau, A., Pegliasco, C., Gourdeau, L., and Singh, A.: Observed Characteristics and Vertical
 Structure of Mesoscale Eddies in the Southwest Tropical Pacific, *Journal of Geophysical Research: Oceans*, 123, 2731–2756,
 550 <https://doi.org/10.1002/2017JC013712>, 2018.



- Kurian, J., Colas, F., Capet, X., McWilliams, J. C., and Chelton, D. B.: Eddy properties in the California Current System, *Journal of Geophysical Research - Oceans*, 116, C08 027, <https://doi.org/10.1029/2010JC006895>, 2011.
- Laxenaire, R., Speich, S., Blanke, B., Chaigneau, A., Pegliasco, C., and Stegner, A.: Anticyclonic eddies connecting the western boundaries of Indian and Atlantic Oceans, *Journal of Geophysical Research: Oceans*, 123, 7651–7677, <https://doi.org/10.1029/2018JC014270>, 2018.
- 555 Laxenaire, R., Speich, S., , and Stegner, A.: Agulhas ring heat content and transport in the South Atlantic estimated by combining satellite altimetry and Argo profiling floats data, *Journal of Geophysical Research: Oceans*, 125, e2019JC015511, <https://doi.org/10.1029/2019JC015511>, 2020.
- Le Vu, B., Stegner, A., and Arsouze, T.: Angular Momentum Eddy Detection and Tracking Algorithm (AMEDA) and Its Application to Coastal Eddy Formation, *Journal of Atmospheric and Oceanic Technology*, 35, 739–762, <https://doi.org/10.1175/JTECH-D-17-0010.1>,
 560 2018.
- Li, Q.-Y., Sun, L., and Lin, S.-F.: GEM: a dynamic tracking model for mesoscale eddies in the ocean, *Ocean Science*, 12, 1249–1267, <https://doi.org/10.5194/os-12-1249-2016>, 2016.
- Liu, Y., Chen, G., Sun, M., Liu, S., and Tian, F.: A Parallel SLA-Based Algorithm for Global Mesoscale Eddy Identification, *Journal of Atmospheric and Oceanic Technology*, 33, 2743–2754, <https://doi.org/10.1175/JTECH-D-16-0033.1>, 2016.
- 565 Lloyd, S. P.: Least squares quantization in PCM, *IEEE Transactions on Information Theory*, 28, 129–136, <https://doi.org/10.1109/TIT.1982.1056489>, 1982.
- Mason, E., Pascual, A., and McWilliams, J. C.: A New Sea Surface Height-Based Code for Oceanic Mesoscale Eddy Tracking, *Journal of Atmospheric and Oceanic Technology*, 31, 1181–1188, <https://doi.org/10.1175/JTECH-D-14-00019.1>, 2014.
- Mkhinini, N., Coimbra, A. L. S., Stegner, A., Arsouze, T., Taupier-Letage, I., and Béranger, K.: Long-lived mesoscale eddies in the eastern
 570 Mediterranean Sea: Analysis of 20 years of AVISO geostrophic velocities, *Journal of Geophysical Research: Oceans*, 119, 8603–8626, <https://doi.org/10.1002/2014JC010176>, 2014.
- Morrow, R. and Le Traon, P.-Y.: Recent advances in observing mesoscale ocean dynamics with satellite altimetry, *Advances in Space Research*, 50, 1062–1076, <https://doi.org/10.1016/j.asr.2011.09.033>, 2012.
- Morrow, R., Birol, F., Griffin, D., and Sudre, J.: Divergent pathways of cyclonic and anti-cyclonic ocean eddies, *Geophysical Research
 575 Letters*, 31, L24 311, <https://doi.org/10.1029/2004GL020974>, 2004.
- Nencioli, F., Dong, C., Dickey, T., Washburn, L., and McWilliams, J. C.: A Vector Geometry–Based Eddy Detection Algorithm and Its Application to a High-Resolution Numerical Model Product and High-Frequency Radar Surface Velocities, *Journal of Atmospheric and Oceanic Technology*, 27, 564–579, <https://doi.org/10.1175/2009JTECHO725.1>, 2010.
- Pegliasco, C., Chaigneau, A., and Morrow, R.: Main eddy vertical structures observed in the four major Eastern Boundary Upwelling Systems,
 580 *Journal of Geophysical Research: Oceans*, 120, 6008–6033, <https://doi.org/10.1002/2015JC010950>, 2015.
- Pegliasco, C., Delepoulle, A., Mason, E., Morrow, R., Faugère, Y., and Dibarboure, G.: META3.1exp: A new global mesoscale eddy trajectory atlas derived from altimetry, *Earth System Science Data*, 14, 1087–1107, <https://doi.org/10.5194/essd-14-1087-2022>, 2022.
- Penven, P., Echevin, V., Pasapera, J., Colas, F., and Tam, J.: Average circulation, seasonal cycle, and mesoscale dynamics of the Peru Current System: A modeling approach, *Journal of Geophysical Research: Oceans*, 110, C10 021, <https://doi.org/10.1029/2005JC002945>, 2005.
- 585 Shi, C. and Nof, D.: The splitting of eddies along boundaries, *Journal of Marine Research*, 51, 771–795, 1993.
- Siegel, D. A., Court, D. B., Menzies, D. W., Peterson, P., Maritorena, S., and Nelson, N. B.: Satellite and in situ observations of the bio-optical signatures of two mesoscale eddies in the Sargasso Sea, *Deep-Sea Research Part II*, 55, 1218–1230, <https://doi.org/10.1016/j.dsr2.2008.01.012>, 2008.



- Sosa-Gutierrez, R., Jouanno, J., and Berline, L.: Sargassum spp. accumulation and transport by mesoscale eddies, *Ocean Science*, 21, 1505–1514, <https://doi.org/10.5194/os-21-1505-2025>, 2025.
- Staaterman, E., Paris, C. B., and Helgers, J.: Orientation behavior in fish larvae: A missing piece to Hjort’s critical period hypothesis, *Journal of Theoretical Biology*, 304, 188–196, <https://doi.org/10.1016/j.jtbi.2012.03.016>, 2012.
- Stammer, D. and Wunsch, C.: Temporal changes in eddy energy of the oceans, *Deep-Sea Research Part II*, 46, 77–108, [https://doi.org/10.1016/S0967-0645\(98\)00106-4](https://doi.org/10.1016/S0967-0645(98)00106-4), 1999.
- 595 Tian, F., Wu, D., Yuan, L., and Chen, G.: Impacts of the efficiencies of identification and tracking algorithms on the statistical properties of global mesoscale eddies, *International Journal of Remote Sensing*, 41, 2835–2860, <https://doi.org/10.1080/01431161.2019.1694724>, 2020.
- Tian, F., Li, Z., Yuan, Z., and Chen, G.: EddyGraph: The tracking of mesoscale eddy splitting and merging events in the northwest Pacific Ocean, *Remote Sensing*, 13, 3435, <https://doi.org/10.3390/rs13173435>, 2021.
- 600 Trieling, R.-R., Fuentes, O.-U., and van Heijst, G.-J.-F.: Interaction of two unequal corotating vortices, *Physics of Fluids*, 17, <https://doi.org/10.1063/1.1993887>, 2005.
- Visvalingam, M.: The Visvalingam algorithm: Metrics, measures and heuristics, *The Cartographic Journal*, 53, 242–252, <https://doi.org/10.1080/00087041.2016.1151097>, 2016.
- Wang, Z.-F., Sun, L., Li, Q.-Y., and Cheng, H.: Two typical merging events of oceanic mesoscale anticyclonic eddies, *Ocean Science*, 15, 1545–1559, <https://doi.org/10.5194/os-15-1545-2019>, 2019.
- 605 Zhang, Z., Wang, W., and Qiu, B.: Oceanic mass transport by mesoscale eddies, *Science*, 345, 322–324, <https://doi.org/10.1126/science.1252418>, 2014.