

1 **A random forest isoscape model of bioavailable Sr for South America:**
2 **a focus on southern Brazil**

3
4 Cinzia Scaggion¹, Tommaso Giovanardi^{1*}, László Palcsu², Daniel Loponte³, Mirian Carbonera⁴,
5 Sara Bernardini^{5,1}, Stefano Benazzi⁵, Giulia Marciani⁵, Marcos Cesar Pereira Santos⁶, Eugenio
6 Bortolini⁵, Anna Cipriani^{1,7}, Federico Lugli^{1,8}

¹ Department of Chemical and Geological Sciences, University of Modena and Reggio Emilia,
via Campi 103, I-41125 Modena, Italy.

² ICER Centre, HUN-REN Institute for Nuclear Research, Bem square, 18/C 4026 Debrecen,
Hungary

³ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto Nacional de
Antropología y Pensamiento Latinoamericano (INAPL), 3 de Febrero 1370/78 Buenos Aires,
Argentina.

⁴ Programa de Pós-graduação em Ciências Ambientais e Centro de Memória do Oeste de Santa
Catarina da Universidade Comunitária da Região de Chapecó (Unochapecó), Rua Senador Atilio
Fontana, 591 E Efapi, 89809000 - Chapecó, SC Brazil.

⁵ Department of Cultural Heritage, University of Bologna, via degli Ariani 1, I-48121 Ravenna,
Italy.

⁶ Instituto de Ciências da Sociedade, Curso de Arqueologia, Universidade Federal do Oeste do
Pará, Santarém, Brazil.

⁷ Lamont-Doherty Earth Observatory of Columbia University, Palisades, New York, United
States of America.

⁸ Institut für Geowissenschaften, Goethe-Universität Frankfurt, Frankfurt am Main, Germany.

7 *corresponding author

8 E-mail: tommaso.giovanardi@unimore.it (TG)

9

10 **Abstract**

11 In recent years, advances in machine learning have greatly improved the generation of maps showing
12 the geographic distribution of isotope ratios (isoscapes), which have become essential tools for
13 environmental, mobility and provenance studies in both modern and archaeological contexts. Among
14 the various isotopic systems employed, strontium (Sr) is particularly useful because its ⁸⁷Sr/⁸⁶Sr ratio
15 in the environment is largely controlled by the underlying geology through the composition of local
16 soils and rocks.

Definizione stile: Titolo 1

Definizione stile: Titolo 2

Tabella formattata

In this work, we present a new dataset of bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ ratios derived from $n = 233$ plant samples collected across southern Brazil, covering the states of Santa Catarina and Rio Grande do Sul (c.a. 370,000 km²). The measured ratios span from 0.70521 to 0.76039 and capture the bioavailable Sr isotope signatures over all major geological units in the region.

We combined these new data with an extensive compilation of published bioavailable Sr measurements from across South America (including plants, fauna, ancient human remains, shells, snails, lichens, water and soils) to construct three random forest Sr isoscapes using different subsets of the combined dataset at the regional and continental scales. The first model incorporates the entire dataset ('All' dataset, $n = 883$ sites), the second is based on plant+fauna+lichen+human ($n = 661$ sites) and the third is limited to plant+lichen samples ($n = 531$ sites). Among the three models, the full dataset model shows lower predictive power, while the plant+fauna+lichen+human and the plant+lichen models yield better results, with similar RMSE (0.0049 and 0.0054) and R^2 values (ca. 0.76). Compared to existing Sr isoscapes of South America, our models significantly enhance both spatial coverage and resolution of bioavailable Sr predictions, particularly in southern Brazil.

The new bioavailable Sr isotope dataset from Santa Catarina and Rio Grande do Sul states is available at <https://doi.org/10.5281/zenodo.17988601> (Scaggion et al., 2025a) and the compiled literature dataset is reported as supplementary material.

1 Introduction

Maps of isotopic variability through the landscape are powerful tools used for answering questions related to migration, trade, diet, provenance, forensic investigations, and environmental changes in both modern and ancient contexts (e.g. Hobson, 1999; Benson et al., 2006; Makarewicz and Sealy, 2015; Bataille et al., 2018, 2020; Hoogerwerff et al., 2019; Lugli et al., 2022; Gigante et al., 2023; Armaroli et al., 2024; Asrat et al., 2025; Dosseto et al., 2025; Scaggion et al., 2025b).

Among the various isotopic systems, strontium (Sr) isotopes play a central role in geotracing and provenance studies of biological materials because biota obtains Sr primarily from soils derived through weathering and disaggregation of local rocks. As a result, the so-called bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ act as a proxy for the underlying geology of a region.

Radiogenic ^{87}Sr originates by β -decay of rubidium-87 (^{87}Rb), while ^{86}Sr is stable. Consequently, the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in rocks and minerals changes depending on their ^{87}Rb content, their ages and geological history providing characteristic isotopic signatures for different geological domains (Faure and Mensing, 2005; Dickin, 2018). Through alteration and pedogenetic processes, Sr ions are leached from rocks into soils and waters, where they mix with other Sr pools derived from rainfall, snow, groundwater, atmospheric deposition or anthropic activities (Vitousek et al., 1999; Voerkelius et al.,


51 2010). These processes can shift the local Sr isotopic signature away from the original bedrock. From
 52 soil and water, Sr enters the local ecosystem (Bullen et al., 1996; Capo et al., 1998). This bioavailable
 53 Sr is passively absorbed by plant roots and transported to leaves, where Sr^{2+} substitutes for calcium²⁺
 54 (Ca) in metabolic processes (Poszwa, 2000; White and Broadley, 2003). From plants and water, the
 55 bioavailable Sr enters the local trophic chain. In vertebrates, Sr replaces Ca in the crystalline structure
 56 of carbonated-hydroxyapatite (Pors Nielsen, 2004; Bentley, 2006), the main inorganic constituent of
 57 bones and teeth (Weiner and Wagner, 1998; Hughes and Rakovan, 2002). Thus, the $^{87}\text{Sr}/^{86}\text{Sr}$ ratios
 58 of bone and dental tissues reflect those of water, plants and animals consumed, which, in turn, mirror
 59 the bioavailable isotopic signature of the environment (Schwarcz et al., 2010; Crowley et al., 2017).
 60 Although isotopic fractionation can occur along the trophic chain and through metabolic processes,
 61 its enrichment is generally minimal (Lahtinen et al., 2020) and can be effectively corrected using
 62 standard normalization protocols during mass spectrometry analysis (Fietzke and Eisenhaure, 2006;
 63 Frei and Frei, 2011; Spies et al., 2025). For this reason, the isotopic composition of low-mobility
 64 fauna and plants represents a valuable, though not exclusive, *proxy* for estimating the local
 65 bioavailable Sr isotope signatures (Spies et al., 2025). However, deciphering environmental
 66 heterogeneity and trophic complexity is not always straightforward; therefore, integrating multiple
 67 isotopic and environmental datasets is crucial for constructing robust isoscapes.
 68 Over the last decade, advances in spatial modelling and machine learning techniques such as kriging
 69 and random forest algorithms (e.g. Bataille et al., 2018, 2020; Hoogerwerff et al., 2019; Gigante et
 70 al., 2023; Armaroli et al., 2024; Scaggion et al., 2025b; Spies et al., 2025), have significantly
 71 increased the predictive potential of isoscapes models, by linking spatial isotopic variability to
 72 environmental predictors (Bataille et al., 2020). One of the main challenges, however, remains the
 73 acquisition of spatially representative and geologically diverse samples, making well designed
 74 sampling strategies crucial for developing accurate predictive models.
 75 Brazil represents a particularly interesting case study for Sr isoscape research, due to its geological
 76 diversity and rich archaeological and ecological history. In southern Brazil, the states of the Santa
 77 Catarina (SC) and Rio Grande do Sul (RGS) encompass various geological formations, from ancient
 78 Precambrian crystalline rocks to younger sedimentary basins, which provide a good opportunity to
 79 test the spatial resolution, robustness, and applicability of Sr isoscapes. Despite the relevance of Brazil
 80 for provenance studies in archaeology, ecology, and food authentication, to date a comprehensive
 81 isoscape of bioavailable Sr has not been produced.
 82 In this study, we present a newly generated dataset of $n = 233$ plant samples, collected across SC and
 83 RGS, representing all major geological units of the region (c.a. 370,000 km²). This dataset was
 84 integrated with a literature-based compilation of bioavailable Sr data from across South America to

~~developed~~develop new isoscapes of bioavailable strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) for South America and southern Brazil, with a focus on SC and RGS. Using a random forest machine learning model following Bataille et al. (2020), we incorporated global environmental variables described in Bataille et al. (2018, 2020) and Reich et al. (2024) to estimate spatial patterns of bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ ratios. The sampled region represents a pivotal area for understanding the expansion of pre-Columbian populations, constituting pass-through areas that facilitated the spread of natives such as the Guaraní, who from approximately the 5th c. AD gradually occupied the continent from the Amazon to the Paraná Delta (Loponte et al., 2024) as well as the movements of Southern Brazilian Highland populations toward the coast. The geological heterogeneity of these states provides a unique natural context for investigating large-scale dynamics of mobility, settlement, and cultural interaction.

2 Geology

We focused our sampling on the southern portion of Brazil, targeting the main geological units in detail to increase the reliability and spatial resolution of the resulting Sr database and isoscape. The study area, comprising the states of RGS and SC, represents a key segment of the South American continent. The region preserves a long and complex geological evolution, spanning from the Archean to the Cenozoic, encompassing processes of crustal growth, continental collision, sedimentary basin development, and large igneous province magmatism. A simplified overview of this geological complexity is provided in Fig. 1, modified from the 1:5000000 geological map of Brazil (Medeiros et al. 2025) downloaded from the Serviço Geológico do Brasil (<https://geosgb.sgb.gov.br/geosgb/downloads.html>) for use in QGIS software.

106



107




Figure 1: simplified geological map of southern Brazil, modified from Medeiros et al. (2025) with sample locations collected in this study. The lithology codes are simplified from Medeiros et al. (2025) and ~~divided for~~grouped according to the two main geological domains of the area: the Mantiqueira Province and the Paraná Basin. The supergroups of the ~~latter~~Paraná Basin are ~~divided according to~~subdivided following Milani et al. (2007).

Two major geological provinces dominate this area: the southern Brazilian Shield, representing the Precambrian crystalline basement, and the Paraná Basin, a large Paleozoic–Mesozoic sedimentary and volcanic basin. In addition to these two major geological provinces, Pleistocene and Holocene sediments outcrop along the coast and the southern lagoons of RGS.

The southern Brazilian Shield, also known as the Mantiqueira Province (Heilbron et al., 2004) is composed of a mosaic of cratonic nuclei, accreted terranes, and mobile belts that were amalgamated during the Neoproterozoic, culminating with the Gondwana accretion (Brito Neves et al., 2014). In the state of RGS, the principal tectonic domains of the Mantiqueira Province are the Dom Feliciano Belt and the São Gabriel Terrane (Brito Neves et al., 2014; Philipp et al., 2018). The Dom Feliciano Belt extends in a NE–SW orientation and consists of high-grade metamorphic complexes, syn- to post-tectonic granitic intrusions (650–540 Ma), and metavolcanic–sedimentary sequences (2.35–0.78 Ga). It is interpreted as a Neoproterozoic collisional orogen, developed during the closure of oceanic domains between the Rio de la Plata Craton and the Kalahari–Congo Craton (Brito Neves et al., 2014; Philipp et al., 2018). The belt includes the voluminous granites of the Pelotas Batholith dated between 650 and 580 Ma. The São Gabriel Terrane, located west of the Dom Feliciano Belt, comprises volcanic-arc sequences, ophiolitic remnants, and accretionary complexes. Geochemical data indicate that it represents an intra-oceanic arc formed between 900 and 700 Ma, subsequently accreted to the continental margin during the Brasiliano Orogeny (Cerva-Alvez et al., 2021). The Dom Feliciano Belt is also exposed along the coast in the eastern part of SC where it includes the Florianópolis granitic batholiths and the metavolcanic–sedimentary sequences (Brito Neves et al., 2014; Philipp et al., 2018).

The Paraná Basin is one of the world's largest intracratonic sedimentary basins, covering approximately 1.4 million km² across Brazil, Paraguay, Uruguay, and Argentina (Holz et al., 2010; Scherer et al., 2023). It developed over the consolidated Precambrian basement during a long-lived post-orogenic to extensional tectonic regime that lasted from the Ordovician to the Cretaceous.

Milani et al. (2007) divided the Paraná Basin succession into six different second-order supersequences (from oldest to youngest): Rio Ivaí, Paraná, Gondwana I, Gondwana II, Gondwana III (which includes the Serra Geral) and Bauru. In the studied area, the exposed units comprise: (1)

Carboniferous-Permian volcanic-sedimentary units of the Gondwana I Supersequence (Itararé group and the Rio Bonito, Palermo and Irati formations); (2) Triassic sedimentary rocks of the Santa Maria Formation belonging to the Gondwana II Supersequence; and (3) Jurassic volcanic rocks of the Serra Geral Formation and aeolian sandstones of the Botucatu Formation, representing the Gondwana III Supersequence.

The volcanic-sedimentary units outcrop along the eastern margin of the basin, in contact with the rocks of Dom Feliciano Belt, while the volcanic rocks of Serra Geral Formation outcrop in the western part of the studied area (Fig. 1). The stratigraphic record documents alternating marine, glacial, and continental depositional environments. The major stratigraphic units include: the Itararé Group (Late Carboniferous–Early Permian) composed of glaciogenic diamictites, sandstones, and shales deposited during the Gondwanan glaciation (Holz et al., 2010); the Rio Bonito and Palermo formations (Permian) composed of coal-bearing strata deposited in deltaic and coastal-plain environments, marking post-glacial transgression (Holz et al., 2010); the Irati Formation (Late Permian) composed of organic-rich black shales and carbonates deposited under restricted marine conditions and considered one of the principal source rocks for hydrocarbons in the basin (Holz et al., 2010); the Santa Maria Formation (Triassic) composed of fluvial and aeolian sandstones (Scherer et al., 2023); and the Botucatu Formation (Jurassic) composed of cross-bedded aeolian sandstones indicative of an extensive desert environment prior to the onset of volcanism (Scherer et al., 2023).

One of the main geological domains in southern Brazil is the Serra Geral Formation which records the bimodal volcanism of the Paraná-Etendeka Large Igneous Province (LIP). This volcanic province covers an area of approximately 1,200,000 km² in South America, extending across Brazil, Argentina, Uruguay and Paraguay, and constitutes the upper volcanic portion of the Paraná Basin (e.g. Peccerillo and Melfi, 1988). The magmatism has a bimodal character, with tholeiitic basalts and andesites as the dominant lithologies coupled with subordinate more silicic rocks (rhyodacite and rhyolite, also occurring as high-grade ignimbrites; Peate et al., 1992; Luchetti et al., 2018). The eruptive history is short, with ages constrained between 135.0 ± 0.6 Ma and 132.0 ± 0.2 Ma, with a time span of 1.6–3.0 Ma for the extrusive activity (Gomes and Vasconcelos, 2021). The Sr isotopic composition of this magmatism is extremely variable (e.g. ⁸⁷Sr/⁸⁶Sr₁₃₀ between 0.705–0.728; Peate et al., 1992), reflecting significant crustal contamination during magmas ascent and/or derivation from metasomatized mantle sources (e.g. Giovanardi et al., 2022). These volcanic rocks form a plateau in the western part of the studied area separated from other geological domains by a stiff cliff.

173

3 Material and Methods

3.1 Materials

176 A total of $n = 233$ samples of plant leaves were collected across the study area (Fig. 1), covering all
177 of the geological units identified on state geological maps (downloaded from the Serviço Geológico
178 do Brasil website https://geosgb.sgb.gov.br/geosgb/downloads_en.html; geological map of Santa
179 Catarina from Wildner et al., 2014; Rio Grande do Sul map from Wildner et al., 2013).
180 Sampling was designed to minimize potential contamination from agriculture, waste disposal or other
181 anthropogenic sources. To this end, leaves were taken from trees as far as possible from anthropic
182 activities (Spies et al., 2025). The sampled taxa include: *Eucalyptus* sp., *Phytolacca dioica*, *Neltume*
183 *nigra*, *Smilax campestris*, *Celtis australis*, *Fraxinus excelsior* (Common ash) and *Nerium oleander*.
184 Different plant taxa were intentionally collected at each location to account for different rooting
185 depths at each location.

186

187 3.2 Isotopic analyses

188 Chromatographic separation of Sr was conducted at the MeGic lab
189 (<https://www.geochem.unimore.it/>) of the Department of Chemical and Geological Sciences
190 (Unimore, Italy). Leaf samples were cleaned to remove major impurities using ultrapure MilliQ®
191 water and then dried for 24 h on a hotplate at 50°C. Approximately 20 g of dried clean plant material
192 was reduced to ash in a porcelain crucible in a muffle oven at 650 °C for 6 h. Five milligrams of ash
193 were then dissolved in 0.5 ml of 3M HNO₃ and centrifuged prior to loading into the chromatographic
194 column. Sr separation was performed using a standard procedure in ~30 µl columns filled with the
195 100-150 µm bead size Eichrom Sr-spec Resin (Eichrom Technologies, LLC) as described in
196 Argentino et al. (2021). The Sr was eluted with MilliQ® water. Through the chromatographic
197 separation runs, $n = 5$ blanks were produced to monitor possible contamination during the procedure.
198 Strontium isotope ratios ($^{87}\text{Sr}/^{86}\text{Sr}$) of 197 samples were measured with the Thermo Scientific
199 NEPTUNE Plus multi-collector ICP-MS (MC-ICPMS) at the HUN-REN Institute for Nuclear
200 Research (ATOMKI), Debrecen, Hungary. Isotopic ratios were corrected for instrumental mass
201 discrimination using $^{88}\text{Sr}/^{86}\text{Sr} = 8.375209$ and by applying an interference correction for $^{87}\text{Rb}^+$, $^{84}\text{Kr}^+$
202 and $^{86}\text{Kr}^+$ with $^{85}\text{Rb}^+$ and $^{83}\text{Kr}^+$, respectively. The same method, shared between the two laboratories
203 (Cavazzuti et al., 2021), was used to analyse $n = 36$ samples at the laboratory of the Centro
204 Interdipartimentale Grandi Strumenti of Unimore using a Thermo Scientific NEPTUNE MC-ICPMS.
205 Data reduction was performed with a modified version of the 'SrDR' spreadsheet (Lugli et al., 2020).
206 Data are available at <https://doi.org/10.5281/zenodo.17988601> (Scaggion et al., 2025a).

207

208 3.3 — Quality assessment of isotopic data

Codice campo modificato

Formattato: Car. predefinito paragrafo, Tipo di carattere: Calibri, 11 pt

209 ~~A poor~~Poor chromatographic removal of Rb from the sample could ~~lead~~lead to
210 ~~overestimate~~overestimation of the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio due to the isobaric interference of ^{87}Rb on ^{87}Sr .
211 Before isotopic analyses, the Rb and Sr concentration of ~~separate~~the separated solutions were
212 determined using a triple quadrupole ICPMS iCAP TQ (Thermo Fisher) at the laboratory of the
213 Centro Interdipartimentale Grandi Strumenti of Unimore. ~~The instrument was calibrated using a~~
214 ~~calibration curve for Rb, Sr and In from 1 to 1000 ppb, based on the international reference material~~
215 ~~IV-ICPMS-71A-A. ^{115}In solution was employed as an internal standard. Calibration curves were~~
216 ~~obtained using the multi-element standard IVICP-MS-71A (Inorganic Venture), in the range of 1-~~
217 ~~1000 ppb.~~ Samples containing Rb higher than 0.5 ppb and with Rb/Sr ratio higher than 0.005 were
218 reprocessed through a second chromatographic separation.

219 ~~Possible~~Potential contamination during the chromatographic separation ~~has been~~was evaluated with
220 ~~laboratory~~ blank samples, which were monitored ~~using their signals V for ion counting~~during the
221 isotopic analyses at the Centro Interdipartimentale Grandi Strumenti of Unimore. The ~~resulting~~
222 ~~signals of masses~~ were indistinguishable from the ~~instrument background~~analytical blank (i.e. 4%
223 ~~suprapur HNO_3~~).

224 To ~~provide~~ensure homogeneous reproducibility of the data, both laboratories ~~share~~followed the same
225 analytical protocol, already used in ~~other~~previous publications (e.g. Cavazzuti et al., 2021). The
226 international Sr carbonate isotope Standard Reference Material NBS987, produced by the National
227 Institute of Standard and Technology (NIST), was used ~~as~~by both laboratories as an external standard
228 for instrumental bias correction ~~by both laboratories~~. Each session includes the analysis of three
229 samples bracketed by the NBS-987 reference material. The instrumental bias was corrected
230 independently ~~in~~for each analytical session ~~by~~normalizing to ~~an~~the accepted NBS-987 value of
231 0.710248 (McArthur et al., 2001). The average $^{87}\text{Sr}/^{86}\text{Sr}$ value of the NBS-987 ~~of~~across all the
232 sessions ~~in HUN-REN laboratory is~~was 0.710329 $\pm$ 0.000006 ~~at the HUN-REN laboratory and the~~
233 ~~average value of Unimore is~~0.710250 $\pm$ 0.000014 ~~at Unimore~~.

234 3.4 Statistical analysis and geospatial modelling

236 We built the Sr isoscapes of South America by combining the new Sr isotope dataset of 233 plant
237 samples with literature data. The latter includes the Isoarch South America dataset (Salesse et al.,
238 2018, 2020), excluding rock samples, and additional published sources (see Supplement Table S1)
239 for a total of 1588 data points. ~~The final dataset used for the model includes 1821 data from 883 sites~~
240 ~~(i.e. For the continental-scale compilation, we included both environmental samples (plants, lichens,~~
241 ~~soils, and water) and archaeological human and faunal samples identified as local in the original~~
242 ~~publications. Non-local individuals were excluded. Metadata relating to coordinates and material~~

Formattato: Colore carattere: Automatico


Formattato: Colore carattere: Automatico

Formattato: Colore carattere: Automatico

243 categories were harmonized, and duplicates were consolidated by site. Uncertainty values not
 244 reported in the original publications or datasets were excluded. All primary datasets used in the
 245 compilation, including the open-access IsoArch repository, Borges et al. (2021), Plomp (2021),
 246 Stantis et al. (2024) and Seferidou (2025), have been integrated with all additional sources and
 247 reported in Table S1. Overall data are from : Brass (1976), Palmer and Edmond (1989), Edmond et
 248 al. (1995; 1996), Hieronymus (1995), Henry et al. (1996), Gaillardet et al. (1997), Négrel and
 249 Lachassagne (2000), Grove et al. (2003), Knudson et al. (2004; 2013), Brunet et al. (2005), Pasquini
 250 et al. (2005), Knudson and Price (2007), Knudson (2008), Martins et al. (2008), Andrushko et al.
 251 (2009), Bastos (2009; 2014), Fiege et al. (2009), Hermenegildo (2009), Poszwa et al. (2009), Bastos
 252 et al. (2011; 2014; 2015; 2016; 2019; 2021), Laffoon et al. (2012), Lee et al. (2013), Machado (2013),
 253 Malaspinas et al. (2014), Pouilly et al. (2014), Santos et al. (2014), Oppitz (2015), Strauss et al.
 254 (2016), Barberena et al. (2017; 2019; 2020; 2021), Chala-Aldana et al. (2018), Duran et al. (2018),
 255 Slovak et al. (2018), Plomp et al. (2019), Moquet et al. (2020), Serna et al. (2020), Azevedo et al.
 256 (2021), Washburn et al. (2021), Fernandez et al. (2022), Quaggio et al. (2022), Torres-Rouff et al.
 257 (2022), Avigliano et al. (2023), Seferidou et al. (2023), Silva et al. (2023), Kafino et al. (2024),
 258 Martinelli et al. (2025), Loponte et al. (2025), Scaggion et al. (2025a). The final dataset used for the
 259 model includes 1821 data from 883 sites (i.e., same coordinate samples) divided in 9 categories as
 260 follows: plant, fauna (archaeological remains, non-local excluded based on publication results),
 261 human (archaeological remains, non-local excluded), shell, snail, lichen, water and soil. The dataset
 262 is updated to September 2025. We constructed three different models, the first utilizing the whole
 263 dataset (hereafter ‘All’ dataset), the second limited to plant+fauna+lichen+human (661 sites) and the
 264 third using only the plant+lichen samples (531 sites). After a preliminary run of the random forest
 265 model, we noticed ~~high prediction errors~~remarkably large residuals for $^{87}\text{Sr}/^{86}\text{Sr}$ values higher than
 266 0.900. We thus excluded from the models $n = 5$ ‘anomalous’ values of river waters from the Parguaza
 267 Batholith (Venezuela; $^{87}\text{Sr}/^{86}\text{Sr} > 0.900$) from Edmond et al. (1995). ~~Here~~In the main text, we mainly
 268 focus on the plant+lichen isoscape, while the other models are reported in the Supplement Text S1.
 269 Note that the quantile colour scale is the same for each model, but rescaled to the values of the
 270 highlighted areas.
 271 The dataset was regressed using the random forest machine-learning model from Bataille et al. (2020)
 272 in the R free software (version 4.0.5), which reconstructs the Sr isotopic ratio variability with a
 273 resolution of 1 km^2 . The method uses external predictors variables to generate multiple decision trees
 274 using each time a random subset of data and covariates. For the models generation we used the global
 275 variables presented in Bataille et al. (2018, 2020) and Reich et al. (2024) as external predictors; (Table
 276 1), filtering for each model the variables with high correlation ($-0.80 > r > 0.80$, Fig. 2) to avoid

277 multicollinearity in the model and inflation of the variables importance (Strobl et al., 2008), as in
278 Scaggion et al. (2025b). This resulted in the use of n = 14 external predictors, which were identical
279 for the three models: *r.elevation*, *r.bulk*, *r.fert*, *r.dust*, *r.pet*, *r.cec*, *r.meanage_geol*, *r.ssaw*, *r.ph*,
280 *r.srsrq1*, *r.clay*, *r.age*, *r.volc* and *r.GUM*.
281 We thus optimized the ~~mtmtry~~ parameter for the random trees construction at 5 random variables
282 at the time. The spatial-uncertainty map was generated with the *ranger* package of Wright and
283 Ziegler (2017), which calculates a quantile random forest regression, and then halving the random
284 forest q0.84 - q0.16 difference (i.e., lower and upper limits of a ~68% interval: Funck et al., 2021 and
285 Armaroli et al., 2024).
286 The models were evaluated with a 10-fold cross validation by calculating the Root Mean Squared
287 Error (RMSE). This method requires partitioning of the data into multiple subsets (folds) which are
288 used as unknown to train and test the model, ensuring that it is not biased by single subsets. Using
289 this approach, the model's accuracy and precision are tested across multiple subsets of data, providing
290 a comprehensive assessment of the model's predictive capabilities.

Formattato: Tipo di carattere: Corsivo



Variables	Description	Resolution	Type	Source
r.m1	Median bedrock model	1 km	D	Bataille et al. 2018
r.srsrq1	1st quartile bedrock model	1 km	D	Bataille et al. 2018
r.srsrq3	3rd quartile bedrock model	1 km	D	Bataille et al. 2018
r.age	Terrane age attribute	1 km	D	Mooney et al. 1998
r.dust	Multi-models average (g.m-2.yr-1)	1°x1°	C	Mahowald et al., 2006

<u>r.map</u>	<u>Mean annual precipitation (mm.yr-1)</u>	<u>30-arc sec</u>	<u>C</u>	<u>Hijmans et al., 2005</u>
<u>r.salt.t</u>	<u>CCSM.3 simulation</u>	<u>1.4°×1.4°</u>	<u>C</u>	<u>Mahowald et al. 2006</u>
<u>r.pet</u>	<u>Global Potential Evapo-Transpiration</u>	<u>30-arc sec</u>	<u>C</u>	<u>Zomer et al. 2008</u>
<u>r.ai</u>	<u>Global Aridity Index</u>	<u>30-arc sec</u>	<u>C</u>	<u>Zomer et al. 2008</u>
<u>r.elevation</u>	<u>SRTM (m)</u>	<u>90 m</u>	<u>C</u>	<u>Jarvis et al., 2008</u>
<u>r.ph</u>	<u>Soil pH in H₂O solution (x10)</u>	<u>250 m</u>	<u>C</u>	<u>Hengl et al., 2017</u>
<u>r.clay</u>	<u>Clay (weight %)</u>	<u>250 m</u>	<u>C</u>	<u>Hengl et al., 2017</u>
<u>r.bulk</u>	<u>Bulk density (kg m⁻³)</u>	<u>250 m</u>	<u>C</u>	<u>Hengl et al., 2017</u>
<u>r.GUM</u>	<u>Global unconsolidated sediment map</u>	<u>1 km</u>	<u>C</u>	<u>Börker et al., 2018</u>
<u>r.ssaw</u>	<u>Multi-models average sea salt wet deposition (kg.ha⁻¹.yr⁻¹)</u>	<u>1° x 1°</u>	<u>C</u>	<u>Vet et al., 2014</u>
<u>r.ssa</u>	<u>Multi-models average Sea salt wet+dry deposition (kg.ha⁻¹.yr⁻¹)</u>	<u>1° x 1°</u>	<u>C</u>	<u>Vet et al., 2014</u>
<u>r.cec</u>	<u>Cation Exchange Capacity</u>	<u>250 m</u>	<u>C</u>	<u>Hengl et al., 2017</u>
<u>r.bouguer</u>	<u>WGM2012_Bouguer</u>	<u>2 min</u>	<u>C</u>	<u>Balmino et al., 2012</u>
<u>r.mat</u>	<u>Mean annual temperature (°C)</u>	<u>30-arc sec</u>	<u>C</u>	<u>Hijmans et al., 2005</u>
<u>r.maxage_geol</u>	<u>GLiM age attribute (Myrs)</u>	<u>1 km</u>	<u>D</u>	<u>Hartmann and Moosdorf, 2012</u>
<u>r.minage_geol</u>	<u>GLiM age attribute (Myrs)</u>	<u>1 km</u>	<u>D</u>	<u>Hartmann and Moosdorf, 2012</u>

<u>r.meanage_geol</u>	<u>GLiM age attribute (Myrs)</u>	<u>1 km</u>	<u>D</u>	<u>Hartmann and</u> <u>Moosdorf,</u> <u>2012</u>
<u>r.nfert</u>	<u>Global Nitrogen Fertilization</u>	<u>30-arc</u>	<u>C</u>	<u>Potter et al.,</u> <u>2010</u>
<u>r.volc</u>	<u>Atmospheric deposition of volcanic</u> <u>material (kg.m-2.s-1)</u>	<u>0.5°</u>	<u>C</u>	<u>Brahney et al.</u> <u>2015</u>

Table 1: list of global variables used in the model as external predictors before filtering.





Figure 2: Pearson's correlation coefficients (r) of the external predictors used in the random forest model.

301 **4 Results**

302 *4.1 $^{87}\text{Sr}/^{86}\text{Sr}$ of novel plant samples*

303 Collected samples from the studied area yielded a broad range of $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, between 0.70521
304 and 0.76039 with a median of 0.71360 (Fig. 3). The dataset shows an asymmetric distribution
305 (kurtosis = 13.03, skewness = 2.77) with a long tail towards more radiogenic values (Fig. 3). A
306 Shapiro-Wilk test indicates significant deviation from normality ($W = 0.733$, $p < 0.01$). To facilitate
307 interpretation, results of novel plant samples are presented here according to the underlying
308 geological units or rock type as in Fig. 1 and Table [42](#). We stress however that plants were collected
309 from soils developed on these units.






Figure 3: histogram of the whole dataset utilized to construct the isoscape model and of the new plant data. Boxplots of the plant samples from SC and RGS clustered by geological units as in Fig. 1: Q4-Holocene sediments; Q1-Pleistocene sediments; EN-Paleogene sedimentary; K1 α -Serra Geral Formation, felsic; K1 β -Serra Geral Formation, basic; K1-Cretaceous sedimentary; J-Jurassic sedimentary; T-Triassic sedimentary; P3-Late Permian sedimentary; P1-Early Permian sedimentary; Ppd-Permian andesite; C2-Carboniferous sedimentary; Np3-Ediacaran metasedimentary; Np3 γ -

318 Ediacaran metaigneous; Np2γ-Cryogenian metaigneous; Np1-Tonian metamorphic; PP-
 319 Paleoproterozoic basement.
 320

Geological Unit	Sample Number	⁸⁷ Sr/ ⁸⁶ Sr avg	st St . dev.	Median	Lithology	Age
Q4	21	0.71569	0.00870	0.71155	Sediments	Holocene
Q1	11	0.71306	0.00481	0.71156	Sedimentary	Pleistocene
EN	1	0.71273	0.00002	0.71273	Sedimentary	Paleogene
K1α	15	0.71421	0.00412	0.71436	Magmatic - felsic	Cretaceous
K1β	46	0.71156	0.00226	0.71173	Magmatic - basic	Cretaceous
K1	3	0.71351	0.00202	0.71284	Sedimentary	Cretaceous
J	8	0.71335	0.00151	0.71387	Sedimentary	Jurassic
T	15	0.71432	0.00203	0.71359	Sedimentary	Triassic
Ppd	2	0.71336	0.00309	0.71336	Magmatic	Permian
P3	13	0.71429	0.00292	0.71358	Metasedimentary	Permian - Late
P1	14	0.71516	0.00204	0.71456	Metasedimentary	Permian - Early
C2	5	0.72604	0.01263	0.72181	Metasedimentary	Carboniferous
Np3γ	51	0.71892	0.00830	0.71659	Metaigneous	Ediacaran
NP3	8	0.71364	0.00428	0.71345	Metasedimentary	Ediacaran
NP2γ	10	0.72522	0.01727	0.71850	Metaigneous	Criogenian
Np1	9	0.71601	0.00732	0.71537	Metamorphic	Tonian
PP	1	0.70521	0.00002	0.70521	Metamorphic	Paleoproterozoic

321 Table 42: bioavailable Sr isotopic average composition of this work plant samples divided for the
 322 different geological super-units of the region as reported in Fig. 1. The order follows the stratigraphic
 323 order.

324
 325 Plants from Holocene sediments (Q4) show values between 0.70843 and 0.74054, with an average of
 326 0.71569 ± 0.00870 (n = 21; error calculated as standard deviation; Fig. 3; Table 42) while those from
 327 Pleistocene sediments (Q1, Q2, and Q3) have a narrower range from 0.70940 to 0.72650, with an
 328 average of 0.71306 ± 0.00481 (n=11), similar to only one Paleogene sample (EN) with a value of
 329 0.71273 ± 0.00002 (Table 42). Plants from both these sedimentary groups ~~show~~include anomalous
 330 values, as outliers of the distribution (Fig. 3).

331 Samples from the Cretaceous felsic volcanics units of the Paraná LIP (K1α) show higher values
 332 (between 0.70766 and 0.72147; average of 0.71421 ± 0.00412 , n =15) than those from the basic
 333 volcanics units (K1β; between 0.70666 and 0.71741; average of 0.71156 ± 0.00226 , n =46; Fig. 3,
 334 Table 42). Plants from Cretaceous sediments of the Paraná Basin (K1) have similar K1α values,

Formattato: Colore carattere: Automatico

(0.71191 and 0.71578; average of 0.71351 ± 0.00202 ; $n = 3$; Table 42), almost indistinguishable from those from Jurassic sediments (J; between 0.71045 and 0.71515; average of 0.71335 ± 0.00151 ; $n = 8$; Fig. 3, Table 42). Plants from other sedimentary units of the Paraná Basin (i.e., T, P1, and P3 units) show progressive radiogenic enrichment over time (averages at 0.71432 ± 0.00203 ; 0.71429 ± 0.00292 ; 0.71516 ± 0.00204 , respectively). Samples from Permian volcanics (Ppd) have comparable values, between 0.71118 and 0.71554 with an average of 0.71336 ± 0.00309 ($n = 2$; Table 42). In contrast, plants from Carboniferous sedimentary rocks (C2), have a distinct and more radiogenic signature between 0.71796 and 0.74832, average at 0.72604 ± 0.01263 ($n = 5$; Fig. 3, Table 42). Samples collected from units of the Mantiqueira Province show broader ranges and generally higher values than those from Paraná Basin samples (except for C2 unit; Fig. 3). Plants from Ediacaran sediments of the Dom Feliciano Belt (Np3) yielded $^{87}\text{Sr}/^{86}\text{Sr}$ values between 0.70733 and 0.71871, with an average of 0.71364 ± 0.00428 ($n = 8$), while plants from metagneous units (Np3γ) show higher and more variable values (0.7064 to 0.74790, with an average of 0.71892 ± 0.00830 ; $n = 51$; Fig. 3, Table 42). Samples from Cryogenian metagneous units from the São Gabriel Terrane (Np2γ) show the widest range in the dataset (from 0.70987 to 0.76039; average of 0.72522 ± 0.01727 ; $n = 10$; Fig. 3, Table 42). Plants from the Tonian metasedimentary units (Np1) show values between 0.70769 and 0.72978 with an average of 0.71601 ± 0.00732 ($n = 9$; Fig. 3, Table 42). Finally, the sample from the Paleoproterozoic basement (PP) yielded the lowest value of the entire dataset (0.70521 ± 0.00002 ; Table 42).

4.2 Random forest models

The presence of isotopic outliers in our plant dataset, with ratios higher than those expected for unconsolidated coastal deposits, can be explained by the heterogeneity of the Holocene and Pleistocene deposits (fluvial, marine, aeolian) on which the plants grew, and the characteristics of the parental lithologies. Since these values reflect geological variability, they were retained in the model. In contrast, the extremely radiogenic values for river waters (>0.900) were excluded.

The three constructed isoscape models clearly distinguish the most radiogenic areas of the SC and RGS, which often correspond to the oldest geological units in the region (Figs 4 and 5). In all models, the associated errors are higher in these radiogenic areas (Figs 4 and 5), indicating lower predictive power of the models in these regions.




Figure 4: Spatial distribution of strontium isotopes in South America based on plant+lichen samples (n = 531 sites). Isoscape random forest model (a), associated spatial uncertainty map (b) and sample distribution (c).




Figure 5: Spatial distribution of strontium isotopes in Santa Catarina and Rio Grande do Sul states based on plant+lichen samples ($n = 531$ sites). Isoscape random forest model (a) and associated spatial uncertainty map (b).

Evaluation metrics reported in Table 23 show significant differences between the datasets: the model based on the ‘All’ dataset exhibits the lowest predictive power, with a RMSE of 0.0086 and an R^2 of 0.639, while the models built on the plant+fauna+lichen+human and the plant+lichen datasets show similar RMSE values of 0.0049 and 0.0054 respectively, and an R^2 of c.a. 0.76 each. The lower variance observed in these latter models likely reflects the smaller number and lower variability of

382 samples compared to the ‘All’ dataset, which includes outliers with $^{87}\text{Sr}/^{86}\text{Sr}$ higher than 0.818 in
 383 water samples (Fig. 3).

384


Model	n. of sites	n. of predictors	RMSE	R ²	MAE
‘All’ dataset	883	14	0.0086	0.639	0.0037
Plants+Fauna+Human	661	14	0.0049	0.760	0.0023
Plants+lichens	531	14	0.0054	0.764	0.0027

Tabella formattata

385 Table 23: evaluation metrics of the models.

386

387 The evaluation of the predictive power of the independent variables using the Increase in Node Purity
 388 (IncNodePurity; Fig. 6; Supplement Text S1), shows that in the plant+fauna+lichen+human and
 389 plant+lichen models, the most relevant variables are *r.srsrq1* (predicted first quartile of the global
 390 $^{87}\text{Sr}/^{86}\text{Sr}$ bedrock model; Bataille et al., 2018) and *r.ph* (soil pH in H₂O solution x10; Bataille et al.,
 391 2020), both with IncNodePurity values just below 0.012 (Fig. 6), followed by *r.meanage_geol* (GLiM
 392 age attribute in Myrs; Bataille et al., 2020) and *r.cec* (cation exchange capacity; Bataille et al., 2020)
 393 with values of around 0.009, while all other variables have a contribution lower than 0.004 (Fig. 6).
 394 Notably, a 2D-Partial Dependence Plot (PDP, Fig. 7) shows a nonlinear interaction between *r.ph* and
 395 *r.srsrq1*, and indicates that the RF model predicts higher $^{87}\text{Sr}/^{86}\text{Sr}$ ratios when pH is low and - as
 396 expected - when the *r.srsrq1* bedrock model value is high. (Fig. 7).



397





Figure 6: Predictive variables and model accuracy for random forest model based on plant+lichen samples. Increase in Node Purity (IncNodePurity) of the $n = 14$ external predictors (a); cross-validation scatterplot of observed vs. predicted $^{87}\text{Sr}/^{86}\text{Sr}$ values for the test-validation sample splits (b); model residuals vs. predicted $^{87}\text{Sr}/^{86}\text{Sr}$ ratios (c).






Figure 7: 2D Partial Dependence Plot for predictive variables *r.srsrq1* and *r.ph* for random forest model based on plant+lichen samples.

In contrast, within the ‘All’ model the most predictive variable is *r.pet* (global potential evapotranspiration; Bataille et al., 2018) with an IncNodePurity value equal to 0.038, followed by *r.ph* at 0.020, *r.age* (Terrane age attribute in Myrs; Bataille et al., 2020) at 0.018, and *r.cec* and *r.srsrq1* at 0.015 (Supplement Text S1).

5 Discussion

5.1 Validation of the novel dataset against bedrock geology

Several inputs and processes govern the bioavailable Sr isotopic signature of local areas. However, considering a regional dataset, the bedrock geology is one of the main factors which govern the isotopic distribution (Vitousek et al., 1999; Voerkelius et al., 2010), and thus its relationship with the dataset could be a useful proxy to validate it.

The asymmetric statistical distribution of $^{87}\text{Sr}/^{86}\text{Sr}$ values observed in the novel dataset (Fig. 3), characterized by a long tail toward more radiogenic ratios, reflects the geological complexity of the area and the variety of contributing lithologies (Cryogenian igneous rocks, Ediacaran sedimentary sequences, Cretaceous volcanic flows, etc.). The presence of anomalous values, i.e., ratios higher than those expected for unconsolidated coastal deposits and identified as outliers of the distribution, among the plants growing on Holocene and Pleistocene sediments can be explained by the heterogeneity of the deposits (fluvial, marine, aeolian) and the characteristics of the parental lithologies.

428 The more radiogenic values of plants samples growth on soils from felsic igneous and metamorphic
429 units compared to igneous basic and sedimentary units (e.g. units K1 α , magmatic felsic, and K1 β ,
430 magmatic basic with the same age), are consistent with well-known geological differentiation
431 processes, as Rb commonly behaves as a more incompatible element in magmatic and metamorphic
432 processes than Sr, thus becoming enriched during fractionation, forming rocks with higher or lower
433 Rb content, which over time produce higher or lower $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, as in K1 α and K1 β , respectively.
434 Another aspect related to geology, is the increase of the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in rocks with time, which we
435 expect to be reflected in the derived soil and then in the local vegetation. The plant dataset presented
436 here follows this rule, with a general increase of the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio as the bedrock is older in age (Fig.
437 3; Table 42) as observed for the Paranà Basin samples vs. Mantiqueira Province ones.
438 All ~~these evidences suggest~~this evidence suggests that the dataset is in overall agreement with
439 bedrock geology which is the primary source of bioavailable Sr in nature.

440 441 5.2 Limits, advantages and predicting variables of the random forest models

442 Across the developed models, higher spatial uncertainties are associated with the most radiogenic
443 areas of SC and RGS, generally corresponding to the oldest geological units in the area (Figs. 4, 5).
444 This pattern, also reported in random forest models from the literature (e.g., Bataille et al., 2018;
445 Gigante et al., 2023; Scaggion et al., 2025b), reflects the pronounced contrast in isotope composition
446 between radiogenic bedrock and less radiogenic end-members such as aerosols, seawater, and
447 precipitation. Such contrast complicates the mixing relationships processes that govern the
448 bioavailable Sr isotope signature. Another parameter that influences model accuracy and associated
449 errors is the uneven distribution of samples on a continental scale. Nonuniform sampling tends to
450 introduce bias towards specific geological contexts, thereby limiting the ability to generalize
451 predictions in unrepresented areas.

452 The improved performance observed in the plant+fauna+lichen+human and plant+lichen models,
453 compared to the 'All' dataset, can be attributed to the reduced variability of the data, which limits the
454 influence of outliers and improves the overall model stability. We reiterate that these differences in
455 RMSE and R² more accurately reflect the isotopic range and internal heterogeneity of the training
456 datasets, rather than intrinsic differences in model quality. The absence of significant differences in
457 the predictiveness between these two best performing models may reflect i) nonuniform and
458 incomplete geographic coverage associated with water and soil sampling strategies, and ii) the limited
459 representativeness of inorganic samples compared to biological and bioderived materials in
460 representing the local terrestrial bioavailable Sr signature, as recently reported by Spies et al. (2025).
461 In this regard, for random forest models, a smaller but more geologically representative dataset of

biological samples is preferable over a larger but uneven dataset. Indeed, the latter reduces the representativeness of the isotopic variability and, consequently, compromises the accuracy and reliability of the models. Conversely, a smaller dataset anchored to the local geological context is more stable, robust, and useful for modelling (Marsh et al., 2025).

The predicted distribution map of the bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ isotope ratio shows a consistent correlation with the data obtained as a function of the variables *r.srsrq1* and *r.ph*. The combined marginal effect (Fig.7) between these two independent variables reveals that isotopic predictions are influenced by non-linear interactions between strontium isotope ratios (*r.srsrq1*) and soil pH (*r.ph*). Higher $^{87}\text{Sr}/^{86}\text{Sr}$ ratios correlate with more acidic soils, a pattern consistent with pedogenesis on ancient silicate igneous-metamorphic rocks. In the Amazon region, low pH values result from intense decomposition of organic matter, which leads to strong leaching of the bedrock and, consequently, increases Sr availability in the soil. The PDP (Fig. 7) explains how the geological context, pedogenic processes, and environmental conditions shape the spatio-temporal distribution of Sr isotope values. Compared to existing Sr isoscapes, this study substantially improves both the spatial coverage and resolution of bioavailable predictions for South America, particularly southern Brazil. Although global or continental-scale isoscapes are essential for broader applications, developing regional models based on dense, geologically informed sampling offer much higher resolution and are essential for fine-scale provenance studies, such as intra-site archaeological mobility or ecological habitat use. Despite being focused on SC and the RGS regions, our dataset allowed us to obtain a reasonably robust continental-scale model with an R^2 of 0.764 (Fig.4). Our model compares positively with other continental-scale isoscapes (e.g. the bioavailable Sr isoscape of Australia, with an R^2 of 0.590; Dossetto et al., 2025). With this comparison, we do not intend to downplay the importance of large-scale modeling approaches, which are essential for global applications, but rather to highlight how carefully planned sampling strategies at the local level can provide robust, useful, and informative models at larger scales.

This highlights how carefully designed local sampling strategies not only strengthen regional provenance studies but also provide a reliable base for modeling at broader geographical scales.

6 Data availability

The bioavailable Sr plant isotope dataset of Santa Catarina and Rio Grande do Sul states analysed in this work, [the R script](#) and the raster files of the plant+lichen model are available at <https://doi.org/10.5281/zenodo.17988601> <https://doi.org/10.5281/zenodo.17988601> (Scaggion et al., 2025a), The literature dataset is reported [as in the](#) Supplement Table S1.

496 7 Conclusion

497 This work presents a comprehensive dataset of 233 bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values from plant samples
498 collected across the main geological units of Santa Catarina and Rio Grande do Sul states (southern
499 Brazil). The measured $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in the terrestrial biosphere are consistent with lithological
500 patterns, reflecting the pronounced geological heterogeneity of the region.

501 By integrating these new data with a multi-material dataset of literature data, we developed three
502 isoscapes of bioavailable Sr distribution for South America and the studied region (c.a. 370,000 km²)
503 using advanced random forest modelling. Among the tested approaches, the
504 plant+fauna+lichen+human and plant+lichen models achieved the best predictive performance, while
505 the addition of water and soil reduced model performance due to their lower representativeness of
506 locally bioavailable Sr.

507 Despite the limited dataset, the resulting models show robust predictive performance even at the
508 continental scale. This result demonstrates how well-designed local sampling strategies can ensure
509 reliable regional models that maintain their relevance and applicability at larger spatial scales. Beyond
510 providing a new baseline for environmental, archaeological, and palaeoecological research in
511 southern Brazil, our study highlights an important methodological insight: strategically constructed
512 regional datasets can play a key role in refining continental-scale isoscapes. Future efforts should
513 focus on expanding high resolution datasets across underrepresented geological domains.

514

515 Author Contribution

516 C.S. was in charge of Conceptualization, Data curation, Formal analysis, Investigation, Methodology,
517 Roles/Writing - original draft and Writing - review & editing;

518 T.G. was in charge of Conceptualization, Data curation, Formal analysis, Funding acquisition,
519 Investigation, Methodology, Project administration, Resources, Supervision, Visualization,
520 Roles/Writing - original draft and Writing - review & editing;

521 L.P. was in charge of Data curation, Formal analysis;

522 D.L. was in charge of Data curation;

523 M.C. was in charge of Data curation;

524 S.B. was in charge of Formal analysis;

525 S.B. was in charge of Data curation;

526 G.M. was in charge of Data curation;

527 M.C.P.S. was in charge of Data curation;

528 E.B. was in charge of Funding acquisition, Project administration;

529 A.C. was in charge of Formal analysis, Methodology, Supervision, Roles/Writing - original draft and
530 Writing - review & editing;

531 F.L. was in charge of Conceptualization, Formal analysis, Data curation, Investigation, Methodology,
532 Software, Visualization, Roles/Writing - original draft and Writing - review & editing;

533

534 Acknowledgement

535 CS, TG and EB were financially supported by European Union – Next Generation EU, Mission 4
536 Component 1 through grant PRIN2022 – HABITS 2022BC2Z5F project, CUP E53D23000130001
537 and CUP J53D23000140001. Sber is financed by the European Union - NextGenerationEU through
538 the National Recovery and Resilience Plan (PNRR) - Mission 4, Component 2, Investment 1.3, titled:
539 CHANGES - Cultural Heritage Active Innovation for Sustainable Society (PE0000020 - CUP
540 J33C22002850006). AC received financial support from the European Union - Next-GenerationEU
541 - National Recovery and Resilience Plan (NRRP) – MISSION 4 COMPONENT 2, INVESTIMENT
542 N. 1.1, CALL PRIN 2022 PNRR D.D. 1409 14-09-2022 – “CAST Copper Accumulation, Supply,
543 and Technology among Italian prehistoric societies”, Codice progetto: P202275A2M, Decreto di
544 finanziamento MUR D.D. n. 1388 del 1 settembre 2023 (CUP N. E53D23020120001).The funders
545 had no role in study design, data collection and analysis, decision to publish, or preparation of the
546 manuscript. SB and GM want to thank Antonino Vazzana and Owen Higgins.

547

548 References

549 [Andrushko, V. A., Buzon, M. R., Simonetti, A., and Creaser, R. A.: Strontium isotope evidence for](#)
550 [prehistoric migration at Chokepukio, valley of Cuzco, Perú, Lat. Am. Antiq. 20\(1\), 57-75, 2009.](#)

551

552 Argentino, C., Lugli, F., Cipriani, A., and Panieri, G.: Testing miniaturized extraction
553 chromatography protocols for combined $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{88}/^{86}\text{Sr}$ analyses of pore water by MC-ICP-
554 MS, ~~Limnology and Oceanography~~[Limnol. Oceanogr.](#) Methods, 19, 431-440,
555 <https://doi.org/10.1002/lom3.1043>, 2021.

556

557 Armaroli, E., Lugli, F., Cipriani, A., and Tütken, T.: Spatial ecology of moose in Sweden:
558 Combined Sr-O-C isotope analyses of bone and antler. PLoS One, 19(4):e0300867.
559 <https://doi.org/10.1371/journal.pone.0300867> PMID: 38598461, 2024.

560

561 Asrat, S., Lucchini, F., Tafuri, M. A., Aureli, C., Gallinaro, M., Zerboni, A., Fusco, M., and
562 Spinapolice, E. E.: A strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) isoscape of Southern Ethiopia: implications for hominin

land use and faunal mobility patterns, [Frontiers in Environmental Archaeology](#), [Front. Environ. Archaeol.](#) 4, DOI=10.3389/fearc.2025.1499291, 2025.

[Avigliano, E., Chung, M.-T., Pouilly, M., Huang, K.-F., Casalnuovo, M., Dominino, J., Silva, N., Sánchez, S., Facetti, J. F., and Volpedo, A. V.: Strontium isotope mapping and its application to study the fish life history \(Salminus brasiliensis\) in semi-fragmented rivers \(La Plata Basin, South America\), Fish. Res. 265, 106741, 10.1016/j.fishres.2023.106741, 2023.](#)

[Azevedo, V., Strikis, N., M., Novello, V. F., Roland, C. L., Cruz, F. W., Santos, R. V., Vuille, M., Utida, G., De Andrade, F. R. D., Cheng, H., and Edwards, R. L.: Paleovegetation seesaw in Brazil since the Late Pleistocene: A multiproxy study of two biomes, Earth Planet. Sci. Lett. 563, 116880, 10.1016/j.epsl.2021.116880, 2021.](#)

[Balmino, G., Vales, N., Bonvalot, S., and Briaies, A.: Spherical harmonic modelling to ultra-high degree of Bouguer and isostatic anomalies, J. Geod. 86, 499-520. <https://doi.org/10.1007/s00190-011-0533-4>, 2012.](#)

[Barberena, R., Durán, V. A., Novellino, P., Winocur, D., Benítez, A., Tessone, A., Quiroga, M. N., Marsh, E. J., Gasco, A., Cortegoso, V., Lucero, G., Llano, C., and Knudson, J. K.: Scale of human mobility in the southern Andes \(Argentina and Chile\): A new framework based on strontium isotopes, Am. J. Biol. Anthropol. 164, 305-320, <https://doi.org/10.1002/ajpa.23270>, 2017.](#)

[Barberena, R., Tessone, A., Cagnoni, M., Gasco, A., Durán, V., Winocur, D., Benítez, A., Lucero, G., Trillas, D., Zonana, I., Novellino, P., Fernández, M., Bavio, M. A., Zubillaga, E., and Gautier, E. A.: Bioavailable Strontium in the Southern Andes \(Argentina and Chile\): A Tool for Tracking Human and Animal Movement, Environ. Archaeol. 26\(3\), 323-335, 10.1080/14614103.2019.1689894, 2019.](#)

[Barberena, R., Menéndez, L., Novellino, P., Lucero, G., Luyt, J., Sealy, J., Cardillo, M., Gasco, A., Llano, C., Frigolé, C., Guevara, D., Da Pena, G., Winocur, D., Benítez, A., Cornejo, L., Falabella, F., Mendez, C., Nuevo-Delaunay, A., Sanhueza, L., Sagredo, F. S., Troncoso, A., Zarate, S., Duran, V. A., and Cortegoso, V.: Multi-isotopic and morphometric evidence for the migration of farmers leading up to the Inka conquest of the southern Andes, Sci. Rep. 10, 21171, <https://doi.org/10.1038/s41598-020-78013-x>, 2020.](#)

Barberena, R., Cardillo, M., Lucero, G., le Roux, P. J., Tessone, A., Llano, C., Gasco, A., Marsh, E. J., Nuevo-Delaunay, A., Novellino, P., Frigolé, C., Winocur, D., Benítez, A., Cornejo, L., Falabella, F., Sanhueza, L., Santana Sagredo, F., Troncoso, A., Cortegoso, V., Durán, V. A., and Méndez, C.: Bioavailable Strontium, Human Paleogeography, and Migrations in the Southern Andes: A Machine Learning and GIS Approach, *Front. Ecol. Evol.* 9, 584325. doi: [10.3389/fevo.2021.584325](https://doi.org/10.3389/fevo.2021.584325), 2021.

Bastos, M. Q. R.: Mobilidade humana na pré-história do litoral brasileiro: Análise de isótopos instáveis de estrôncio no sambaqui do Forte Marechal Luz, Fundação Oswaldo Cruz, Rio de Janeiro, 2009.

Bastos, M. Q. R.: Dos sambaquis do sul do Brasil à diáspora africana: Estudos de geoquímica isotópica de séries esqueléticas humanas escavadas de sítios arqueológicos brasileiros, Universidade de Brasília, Brasília, 2014.

Bastos, M. Q. R., Souza, S. M. F. M. D., Santos, R. V., Lima, B. A. F., Santos, R. V., and Rodrigues-Carvalho, C.: Human mobility on the Brazilian coast: an analysis of strontium isotopes in archaeological human remains from Forte Marechal Luz Sambaqui, *An. Acad. Bras. Ciênc.* 83, 731-743, [10.1590/s0001-37652011000200030](https://doi.org/10.1590/s0001-37652011000200030), 2011.

Bastos, M. Q. R., Lessa, A., Rodrigues-Carvalho, C., Tykot, R. H., and Santos, R. V.: Análise de isótopos de carbono e nitrogênio: A dieta antes e após a presença de cerâmica no sítio Forte Marechal Luz, *Revista do Museu de Arqueologia e Etnologia* 0(24), 137-151, [10.11606/issn.2448-1750.revmae.2014.109329](https://doi.org/10.11606/issn.2448-1750.revmae.2014.109329), 2014.

Bastos, M. Q. R., Santos, R. V., Tykot, R. H., Souza, S. M. F. M. D., Rodrigues-Carvalho, C., and Lessa, A.: Isotopic evidences regarding migration at the archeological site of Praia da Tapera: New data to an old matter, *J. Archaeol. Sci. Rep.* 4, 588-595, <https://doi.org/10.1016/j.jasrep.2015.10.028>, 2015.

Bastos, M. Q. R., Santos, R. V., Souza, S. M. F. M. D., Rodrigues-Carvalho, C., Tykot, R. H., Cook, D. C., and Santos, R. V.: Isotopic study of geographic origins and diet of enslaved africans buried in two Brazilian cemeteries, *J. Archaeolog. Sci.* 70, 82-90, [10.1016/j.jas.2016.04.020](https://doi.org/10.1016/j.jas.2016.04.020), 2016.

Bastos, M. Q. R., Solari, A., da Silva, S. F. S. M., and Martin, G.: Estudo preliminar de dieta à partir de isótopos em grupos caçadores-coletores do agreste pernambucano (Holoceno recente - Nordeste do Brasil), *FUMDHAMentos* 16(1), 3-18, 2019.

Bastos, M. Q. R., Murilo; Souza, S. M. F. M. D., Santos, R. V., Cook, D.C., Rodrigues-Carvalho, C., and Santos, R.V.: Da África ao Cemitério dos Pretos Novos, Rio de Janeiro, *Revista de Arqueologia da SAB* 24(1), 66-81, 10.24885/sab.v24i1.315, 2021.

Bataille, C. P., von Holstein, I. C. C., Laffoon, J. E., Willmes, M., Liu, X.-M., and Davies, G. R.: A bioavailable strontium isoscape for Western Europe: A machine learning approach, *PLoS One*, 13(5):, e0197386, <https://doi.org/10.1371/journal.pone.0197386> PMID: 29847595, 2018.

Bataille, C. P., Crowley, B. E., Wooller, M. J., and Bowen, G. J.: Advances in global bioavailable strontium isoscapes, *Palaeogeography, Palaeoclimatology, Palaeoecology, Palaeogeogr. Palaeoclimatol. Palaeoecol.* 555:_, 109849-, <https://doi.org/10.1016/j.palaeo.2020.109849>, 2020.

Benson, S., Lennard, C., Maynard, P., and Roux, C.: Forensic applications of isotope ratio mass spectrometry--a review, *Forensic Science International, Sci. Int.* 157(1), 1--22, <https://doi.org/10.1016/j.forsciint.2005.03.012> PMID: 15919168, 2006.

Bentley, A. R.: Strontium Isotopes from the Earth to the Archaeological Skeleton: A Review, *Journal of Archaeological, J. Archaeol. Method and Theory*, 13(3):_, 135--87, <https://doi.org/10.1007/s10816-006-9009-x>, 2006.

Borges, C., Chanca, I., and Salesse, K.: Stable and radiogenic isotope data from Brazilian bioarchaeological samples: a synthesis [data set], <https://doi.org/10.48530/isoarch.2021.005>, 2021.

Börker, J., Hartmann, J., Amann, T., and Romero-Mujalli, G.: Terrestrial sediments of the earth: development of a global unconsolidated sediments map database (gum), *Geochem. Geophys. Geosyst.* 19, 997-1024. <https://doi.org/10.1002/2017GC007273>, 2018.

Brahney, J., Ballantyne, A. P., Kocielek, P., Leavitt, P. R., Farmer, G. L., and Neff, J. C.: Ecological changes in two contrasting lakes associated with human activity and dust transport in

Formattato: Italiano (Italia)

western Wyoming: Dust-P controls on alpine lake ecology, *Limnol. Oceanogr.* 60, 678-695, <https://doi.org/10.1002/lno.10050>, 2015.

Brass, G. W.: The variation of the marine ratio during Phanerozoic time: interpretation using a flux model, *Geochim. Cosmochim. Acta* 40(7), 721-730, [10.1016/0016-7037\(76\)90025-9](https://doi.org/10.1016/0016-7037(76)90025-9), 1976.

Brito Neves, B. B., Fuck, R. A., and Pimentel, M. M.: The Brasiliano collage in South America: a review, *Brazilian Journal of Geology, Braz. J. Geol.* 44, 493-518, 2014.

Brunet, F., Gaiero, D., Probst, J. L., Depetris, P. J., Gauthier Lafaye, F., and Stille, P.: $\delta^{13}\text{C}$ tracing of dissolved inorganic carbon sources in Patagonian rivers (Argentina), *Hydrol. Process.* 19(17), 3321-3344 [10.1002/hyp.5973](https://doi.org/10.1002/hyp.5973), 2005.

Bullen, T. D., Krabbenhoft, D. P., and Kendall, C.: Kinetic and mineralogic controls on the evolution of groundwater chemistry and $^{87}\text{Sr}/^{86}\text{Sr}$ in a sandy silicate aquifer, northern Wisconsin, USA, *Geochimica et Cosmochimica Acta*, 60(10), 1807-21, [https://doi.org/10.1016/0016-7037\(96\)00052-x](https://doi.org/10.1016/0016-7037(96)00052-x), 1996.

Capo, R. C., Stewart, B. W., and Chadwick, O. A.: Strontium isotopes as tracers of ecosystem processes: theory and methods, *Geoderma*, 82(1-3), 197-225, [https://doi.org/10.1016/S0016-7061\(97\)00102-X](https://doi.org/10.1016/S0016-7061(97)00102-X), 1998.

Cavazzuti, C., Hajdu, T., Lugli, F., Sperduti, A., Vicze, M., Horváth A, Major, I., Molnár, M., Palcsu, L., and Kiss, V.: Human mobility in a Bronze Age Vátya 'urnfield' and the life history of a high-status woman, *PLoS ONE*, 16(7), e0254360, <https://doi.org/10.1371/journal.pone.0254360>, 2021.

Cerva-Alves, T., Hartmann, L. A., Lana, C., Queiroga, G. N., Maciel, L. A. C., Leandro, C. G., and Savian, J. F.: Rutile and zircon age and geochemistry in the evolution of the juvenile São Gabriel Terrane early in the Brasiliano Orogeny, *Journal of South American Earth Sciences*, 112(1), 103505 ISSN 0895-9811, <https://doi.org/10.1016/j.jsames.2021.103505>, 2021.

Chala-Aldana, D., Bocherens, H., Miller, C., Moore, K., Hodgins, G., Rademaker, K.: Investigating mobility and highland occupation strategies during the Early Holocene at the Cuncaicha rock shelter

Formattato: Colore carattere: Automatico

Formattato: Colore carattere: Automatico

Formattato: Colore carattere: Automatico

through strontium and oxygen isotopes, *J. Archaeol. Sci. Rep.* **19**, 811-827, [10.1016/j.jasrep.2017.10.023](https://doi.org/10.1016/j.jasrep.2017.10.023), 2018.

Crowley, B. E., Miller, J. H., and Bataille, C. P.: Strontium isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$) in terrestrial ecological and palaeoecological research: empirical efforts and recent advances in continental-scale models, *Biological Reviews, Biol. Rev.* **92**(1), 43–59, <https://doi.org/10.1111/brv.12217> PMID: 26392144, 2017.

Dickin, A. P.: Radiogenic Isotope Geology, third edition. McMaster University, editor. Cambridge: Cambridge University Press; 2018.

Dosseto, A., Dux, F., Bataille, C., and de Caritat, P.: A bioavailable strontium isoscape of Australia, *Earth System Science Syst. Sci. Data*, **17**, 4865–4880, <https://doi.org/10.5194/essd-17-4865-2025>, 2025.

Durán, V., Novellino, P., Menéndez, L., Gasco, A., Marsh, E., Barberena, R., and Frigolé, C.: Barrio Ramos I. Prácticas funerarias en el inicio del período de dominación Inca del valle de Uspallata (Mendoza, Argentina). *Relaciones de la Sociedad Argentina de Antropología XLIII* (1): xx-xx.

Edmond, J. M., Palmer, M. R., Measures, C. I., Grant, B., and Stallard, R. F.: The fluvial geochemistry and denudation rate of the Guayana Shield in Venezuela, Colombia, and Brazil, *Geochimica et Cosmochimica Acta*, **59**, (16), 3301-3325. [10.1016/0016-7037\(95\)00128-m](https://doi.org/10.1016/0016-7037(95)00128-m), 1995.

Edmond, J. M., Palmer, M. R., Measures, C. I., Brown, E. T., and Huh, Y.: Fluvial geochemistry of the eastern slope of the northeastern Andes and its foredeep in the drainage of the Orinoco in Colombia and Venezuela, *Geochim. Cosmochim. Acta* **16**, 2949-2976, [10.1016/0016-7037\(96\)00142-1](https://doi.org/10.1016/0016-7037(96)00142-1), 1996.

Faure, G., and Mensing, T. M.: Isotopes: Principles and Applications, third edition, editor, New York: John Wiley & Sons INC; 2005.

Fernandez, M. V., Gordon, F., Le Roux, P. J., Winocur, D., Lucero, G., Benitez, A., Rindel, D., Della Negra, C., Bernal, V., and Barberena, R.: Scale of human mobility in northwestern Patagonia: An

Formattato: Italiano (Italia)

733 [approach based on regional geology and strontium isotopes in human remains, *Geoarchaeology* 37,](#)
734 [227-241, 10.1002/gea.21881, 2022.](#)

735

736 [Fiege, K., Miller, C. A., Robinson, L. F., Figueroa, R., and Peucker-Ehrenbrink, B.: Strontium](#)
737 [isotopes in Chilean rivers: The flux of unradiogenic continental Sr to seawater, *Chem. Geol.* 268,](#)
738 [337-343, 10.1016/j.chemgeo.2009.09.013, 2009.](#)

739

740 Fietzke, J., and Eisenhauer, A.: Determination of temperature-dependent stable strontium isotope
741 ($^{88}\text{Sr}/^{86}\text{Sr}$) fractionation via bracketing standard MC-ICP-MS, ~~*Geochemistry—Geophysics*~~
742 ~~*Geosystems, Geochem. Geophys. Geosyst.*~~ 7(8), <https://doi.org/10.1029/2006gc001243>, 2006.

743

744 Frei, K. M., and Frei, R.: The geographic distribution of strontium isotopes in Danish surface waters
745 — A base for provenance studies in archaeology, hydrology and agriculture, ~~*Applied*~~
746 ~~*Geochemistry, Appl. Geochem.*~~ 26(3), 326–40, <https://doi.org/10.1016/j.apgeochem.2010.12.006>,
747 2011.

748

749 [Gaillardet, J., Dupre, B., Allegre, C. J., and Négrel, P.: Chemical and physical denudation in the](#)
750 [Amazon River Basin, *Chem. Geol.* 142, 141-173, 10.1016/s0009-2541\(97\)00074-0, 1997.](#)

751

752 Gigante, M., Mazzariol, A., Bonetto, J., Armaroli, E., Cipriani, A., and Lugli, F.: Machine learning-
753 based Sr isoscape of southern Sardinia: A tool for bio-geographic studies at the Phoenician-Punic site
754 of Nora, ~~*PLoS One*~~, 18(7), e0287787, <https://doi.org/10.1371/journal.pone.0287787> PMID:
755 37467179, 2023.

756

757 Giovanardi, T., da Costa, P. C. C., Girardi, V. A. V., Weska, R. K., Vasconcelos, P. M., Thiede, D.
758 S., Mazzucchelli, M., and Cipriani, A.: Age, geochemistry and mantle source of the Alto Diamantino
759 basalts: Insights on NW Paraná Magmatic Province, *Lithos*, 426-427, 106797, 2022.

760

761 Gomes, A. S., and Vasconcelos, P. M.: Geochronology of the Paraná-Etendeka large igneous
762 province, ~~*Earth—Sciences Review, Sci. Rev.*~~ 220, 103716
763 <https://doi.org/10.1016/j.earscirev.2021.103716><https://doi.org/10.1016/j.earscirev.2021.103716>,
764 2021.

765

Formattato: Colore carattere: Automatico

Formattato: Colore carattere: Automatico

Formattato: Colore carattere: Automatico

Grove, M. J., Baker, P. A., Cross, S. L., Rigsby, C. A., and Seltzer, G. O.: Application of strontium isotopes to understanding the hydrology and paleohydrology of the Altiplano, Bolivia-Peru, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 194, 281-297, 10.1016/s0031-0182(03)00282-7, 2003.

Hartmann, J., and Moosdorf, N.: The new global lithological map database GLiM: a representation of rock properties at the Earth surface, *Geochem. Geophys. Geosyst.* 13, Q12004 <https://doi.org/10.1029/2012GC004370>, 2012.

Heilbron, M., Pedrosa-Soares, A. C., Campos Neto, M. C., Silva, L. C., Trouw, R. A. J., and Janasi, V. A.: Província Mantiqueira. In: Mantesso-Neto V., Bartorelli A., Carneiro C.D.R., Brito Neves B.B. (eds.) *Geologia do Continente Sul-Americano: Evolução da Obra de Fernando Flávio Marques de Almeida*, São Paulo, Beca, 203-234, 2004.

Hengl, T., Mendes de Jesus, J., Heuvelink, G.B.M., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M.N., Geng, X., Bauer-Marschallinger, B., Guevara, M.A., Vargas, R., MacMillan, R.A., Batjes, N.H., Leenaars, J.G.B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS One* 12, e016974, <https://doi.org/10.1371/journal.pone.0169748>, 2017.

Hermenegildo, T.: *Reconstituição da dieta e dos padrões de subsistência das populações pré-históricas de caçadores-coletores do Brasil Central através da ecologia isotópica*, Universidade de São Paulo, thesis dissertation, 10.11606/D.91.2009.tde-14092009-084156, 2009.

Hieronymus, B., Godot, J. M., Boulègue, J., Bariac, T., Negrel, P., Dupré, B.: *Chimie du fleuve Tocantins et de rivières côtières de l'est du Para (Brésil)*, Grands Bassins Fluviaux, Nov 1993, Paris, France. (hal-04002094).

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *Int. J. Climatol.* 25, 1965-1978, <https://doi.org/10.1002/joc.1276>, 2005.

Hobson, K. A.: Tracing origins and migration of wildlife using stable isotopes: a review, *Oecologia* 120(3), 314-26, <https://doi.org/10.1007/s004420050865> PMID: 28308009, 1999.

Holz, M., França, A. B., Souza, P. A., Iannuzzi, R., and Rohn, R.: A stratigraphic chart of the Late Carboniferous/Permian succession of the eastern border of the Paraná Basin, Brazil, South America, *Journal of South American Earth Sciences*, 29(2), 381-399, ISSN 0895-9811, <https://doi.org/10.1016/j.jsames.2009.04.004>, 2010.

Hoogewerff, J. A., Reimann, C., Ueckermann, H., Frei, R., Frei, K. M., van Aswegen, T., Stirling, C., Reid, M., Clayton, A., and Ladenberger, A.: Bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ in European soils: A baseline for provenancing studies, *Science of The Total Environment*, 672, 1033-1044, <https://doi.org/10.1016/j.scitotenv.2019.03.387> PMID: 30999220, 2019.

Hughes, J. M., and Rakovan, J.: The Crystal Structure of Apatite, $\text{Ca}_5(\text{PO}_4)_3(\text{F}, \text{OH}, \text{Cl})$, *Reviews in Mineralogy and Geochemistry*, 48(1), 1-12, <https://doi.org/10.2138/rmg.2002.48.1>, 2002.

Jarvis, A., Reuter, A., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database, CGIAR CSI Consort. Spat. Inf., 1-9, 2008.

Kafino, C. V., de Sousa, I. M. C., Barbieri, C. B., de Amorim, A. M., Santos, R. V.: A proof-of-concept study: Determining the geographical origin of Brazilwood, (*Paubrasilia echinata*) with the use of strontium isotopic fingerprinting, *Sci Justice* 64, 159-165, [10.1016/j.scijus.2023.12.006](https://doi.org/10.1016/j.scijus.2023.12.006), 2024.

Knudson, K. J.: Tiwanaku influence in the South Central Andes: strontium isotope analysis and Middle Horizon migration, *Lat. Am. Antiq.* 19, 3-23, <https://doi.org/10.2307/25478206>, 2008.

Knudson, K. J. and Price, T. D.: Utility of multiple chemical techniques in archaeological residential mobility studies: case studies from Tiwanaku- and Chiribaya-affiliated sites in the Andes, *Am. J. Phys. Anthropol.* 132, 25-39, <https://doi.org/10.1002/ajpa.20480>, 2007.

Knudson, K. J., Price, T. D., Buikstra, J. E., and Blom, D. E.: The use of strontium isotope analysis to investigate tiwanaku migration and mortuary ritual in Bolivia and Peru, *Archaeometry* 46, 5-18, <https://doi.org/10.1111/j.1475-4754.2004.00140.x>, 2004.

- Knudson, K. J., Webb, E., White, C., and Longstaffe, F. J.: Baseline data for Andean paleomobility research: a radiogenic strontium isotope study of modern Peruvian agricultural soils, *Archaeol. Anthropol. Sci.* 6(3), 205-219, 10.1007/s12520-013-0148-1, 2013.
- Laffoon, J. E., Davies, G. R., Hoogland, M. L. P., and Hofman, C. L.: Spatial variation of biologically available strontium isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$) in an archipelagic setting: a case study from the Caribbean, *J. Archaeol. Sci.* 39, 2371-2384, 10.1016/j.jas.2012.02.002, 2012.
- Lahtinen, M., Arppe, L., and Nowell, G.: Source of strontium in archaeological mobility studies—marine diet contribution to the isotopic composition, *Archaeological and Anthropological Sciences*, 13(1), <https://doi.org/10.1007/s12520-020-01240-w>, *Archaeol. Anthropol. Sci.* 13, 1, <https://doi.org/10.1007/s12520-020-01240-w>, 2020.
- Lee, B., Han, Y., Huh, Y., Lundstrom, C., Siame, L. L., Lee, J. I., Park, and B.-K.: Chemical and physical weathering in south Patagonian rivers: A combined Sr-U-Be isotope approach, *Geochim. Cosmochim. Acta* 101, 173-190, 10.1016/j.gca.2012.09.054, 2013.
- Loponte, D., Carbonera, M., and Radaeski, J.: The Guaraní expansion in the Upper Uruguay River. Chronology, colonization strategies, social impacts and environmental changes, *Journal of Archaeological Science: Reports*, *J. Archaeol. Sci.: Rep.* 60, 104826, <https://doi.org/10.1016/j.jasrep.2024.104826>, 2024.
- Luchetti, A. C. F., Nardy, A. J. R., and Madeira, J.: Silicic, high- to extremely high-grade ignimbrites and associated deposits from the Paraná Magmatic Province, southern Brazil, *Journal of Volcanology and Geothermal Research*, *J. Volcanol. Geotherm. Res.* 355, 270–286, 2018.
- Lugli, F., Weber, M., Giovanardi, T., Arrighi, S., Bortolini, E., Figus, C., Marciani, G., Oxilia, G., Romandini, M., Silvestrini, S., Jochum, K. P., Benazzi, S., and Cipriani, A.: Fast offline data reduction of laser ablation MC-ICP-MS Sr isotope measurements: Via an interactive Excel-based spreadsheet 'SrDR', *Journal Of Analytical Atomic Spectrometry*, 35:5, *J. Anal. At. Spectrom.* 35(5), 852-862, <https://doi.org/10.1039/C9JA00424F>, 2020.

Lugli, F., Cipriani, A., Bruno, L., Ronchetti, F., Cavazzuti, C., and Benazzi, S.: A strontium isoscape of Italy for provenance studies, *Chemical Geology, Chem. Geol.* 587, 120624, <https://doi.org/10.1016/j.chemgeo.2021.120624>, 2022.

Machado, M. C.: Metodologias isotópicas Sr87/Sr86, δC13 e δ18O em estudos geológicos e arqueológicos, Universidade Federal do Rio Grande do Sul. Instituto de Geociências, Programa de Pós-Graduação em Geociências, LUME repository, <http://hdl.handle.net/10183/76142>, 2013.

Mahowald N. M., Muhs D. R., Levis S., Rasch P. J., Yoshioka M., Zender C. S., and Luo, C.: Change in atmospheric mineral aerosols in response to climate: Last glacial period, preindustrial, modern, and doubled carbon dioxide climates, *J. Geophys. Res. Atmos.* 111, D10202, <https://doi.org/10.1029/2005JD006653>, 2006.

Makarewicz, C. A., and Sealy, J.: Dietary reconstruction, mobility, and the analysis of ancient skeletal tissues: Expanding the prospects of stable isotope research in archaeology, *Journal of Archaeological Science, J. Archaeol. Sci.* 56, 146–58, <https://doi.org/10.1016/j.jas.2015.02.035>, 2015.

Malaspinas, A.-S., Lao, O., Schroeder, H., Rasmussen, M., Raghavan, M., Moltke, I., Campos, P. F., Sagredo, F. S., Rasmussen, S., Gonçalves, V. F., Albrechtsen, A., Allentoft, M. E., Johnson, P. L. F., Li, M., Reis, S., Bernardo, D. V., DeGiorgio, M., Duggan, A. T., Bastos, M., Wang, Y., Stenderup, J., Moreno-Mayar, J. V., Brunak, S., Sicheritz-Ponten, T., Hodges, E., Hannon, G. J., Orlando, L., Price, T. D., Jensen, J. D., Nielsen, R., Heinemeier, J., Olsen, J., Rodrigues-Carvalho, C., Lahr, M., M., Neves, W. A., Kayser, M., Higham, T., Stoneking, M., Pena, S. D.J., and Willerslev, E.: Two ancient human genomes reveal Polynesian ancestry among the indigenous Botocudos of Brazil, *Curr. Biol.* 24, R1035-R1037, [10.1016/j.cub.2014.09.078](https://doi.org/10.1016/j.cub.2014.09.078), 2014.

Marsh, E. J., Cardillo, M., Knudson, K. J., Dahlstedt, A., and Barberena, R.: A bioavailable strontium isoscape for the Andes based on machine learning, *Journal of Archaeological Science, J. Archaeol. Sci.* 184, 106410, ISSN 0305-4403, <https://doi.org/10.1016/j.jas.2025.106410>, 2025.

Martinelli, L. A., Bataille, C. P., Batista, A. C., Souza-Silva, I. M., Araujo, M. G., Abdalla Filho, A. L., Brunello, A., Tommasiello Filho, M., Higuchi, N., Barboas, A. C., Costa, F., and Nardoto, G. B.: Bioavailable strontium isoscape for the Amazon region using tree wood, *For. Ecol. Manag.* 594, 122963, <https://doi.org/10.1016/j.foreco.2025.122963>, 2025.

[Martins, V., Babinski, M., Ruiz, I., Sato, K., Souza, S., and Hirata, R.: Analytical procedures for determining Pb and Sr isotopic compositions in water samples by ID-TIMS, Quím. Nova 31\(7\), 1836-1842, 10.1590/s0100-40422008000700040, 2008.](#)

McArthur, J. M., Howarth, R. J., and Bailey, T. R.: Strontium isotope stratigraphy: LOWESS version 3: best fit to the marine Sr-isotope curve for 0–509 Ma and accompanying look-up table for deriving numerical age, [The Journal of Geology, J. Geol.](#) 109(2), 155-170, 2001.

Medeiros, V. C., Santos, F. G., Quadros, M. L. E. S., Espirito Santo, E. B. S., Silva, M. A., Moreira, G., Santana, J. S., Domingos, N. R. R., Barros, T. S. C., Santos, P. A., Almeida, M. E., and Wanderley, A. A.: Mapa Geológico do Brasil. Escala 1:5.000.000. Salvador: SGB-CPRM, 2025.

Milani, E. J., Melo, J. H. G., de Souza, P. A., de Fernandes, L. A., and França, A. B.: Bacia do Paraná: [Boletim de Geociências da, Bol. Geociênc.](#) Petrobras, 15(2), 265-287, 2007.

[Mooney, W. D., Laske, G., and Masters, T. G.: CRUST 5.1: A global crustal model at 5° × 5°, J. Geophys. Res. Solid Earth 10, 727-747. <https://doi.org/10.1029/97JB02122>, 1998.](#)

[Moquet, J.-S., Morera, S., Turcq, B., Poitrasson, E., Roddaz, M., Moreira-Turcq, P., Espinoza, J. C., Guyot, J.-L., Takahashi, K., Orrillo-Vigo, J., Petrick, S., Mounic, S., and Sondag, F.: Control of seasonal and inter-annual rainfall distribution on the Strontium-Neodymium isotopic compositions of suspended particulate matter and implications for tracing ENSO events in the Pacific coast \(Tumbes basin, Peru\), Glob. Planet. Change 185, 103080, 10.1016/j.gloplacha.2019.103080, 2020.](#)

[Négrel, P., and Lachassagne, P.: Geochemistry of the Maroni River \(French Guiana\) during the low water stage: implications for water-rock interaction and groundwater characteristics, J. Hydrol. 237, 212-233, 10.1016/s0022-1694\(00\)00308-5, 2000.](#)

[Oppitz, G.: Coisas que mudam: os processos de mudança nos sítios conchíferos catarinenses e um olhar isotópico sobre o caso do sítio Armação do Sul, Florianópolis/SC, Revista de Arqueologia 28\(1\), 166-170, 10.24885/sab.v28i1.421, 2015.](#)

931 [Palmer, M. R., and Edmond, J. M.: The strontium isotope budget of the modern ocean, Earth Planet.](#)
932 [Sci. Lett. 92, 11-26, 10.1016/0012-821x\(89\)90017-4, 1989.](#)
933
934 [Pasquini, A. I., Depetris, P. J., Gaiero, D. M., and Probst, J.-L.: Material Sources, Chemical](#)
935 [Weathering, and Physical Denudation in the Chubut River Basin \(Patagonia, Argentina\): Implications](#)
936 [for Andean Rivers, J. Geol. 113, 451-469, 10.1086/430243, 2005.](#)
937
938 Peate, D. W., Hawkesworth, C. J., and Mantovani, M. S. M.: Chemical stratigraphy of the Paraná
939 lavas (South America): classification of magma types and their spatial distribution. ~~Bulletin of~~
940 ~~Volcanology,~~ [Bull. Volcanol.](#) 55, 119–139, 1992.
941
942 Philipp, R. P., Pimentel, M. M., and Basei, M. A. S.: The Tectonic Evolution of the São Gabriel
943 Terrane, Dom Feliciano Belt, Southern Brazil: The Closure of the Charrua Ocean, S. Siegesmund
944 (Ed.), *Geology of Southwest Gondwana*, Regional Geology Reviews, 243-265, 2018.
945
946 Piccirillo, E. M., and Melfi, A. J.: The Mesozoic Flood Volcanism of the Paraná Basin: Petrogenetic
947 and Geophysical Aspects. Universidade de São Paulo, São Paulo, p. 600, 1988.
948
949 [Plomp, E.: Neodymium isotopes in modern human dental enamel: an exploratory dataset \[data set\],](#)
950 <https://doi.org/10.48530/isoarch.2021.011>, 2021.
951
952 [Plomp, E., von Holstein, I. C. C., Koornneef, J. M., Smeets, R. J., Baart, J. A., Forouzanfar, T., and](#)
953 [Davies, G. R.: Evaluation of neodymium isotope analysis of human dental enamel as a provenance](#)
954 [indicator using 1013 Ω amplifiers \(TIMS\), Sci. Justice 59, 322-331, 10.1016/j.scijus.2019.02.001,](#)
955 [2019.](#)
956
957 Pors Nielsen, S.: The biological role of strontium, *Bone*, 35(3), 583–8. ~~2~~
958 <https://doi.org/10.1016/j.bone.2004.04.026> PMID: 15336592, 2004.
959
960 Poszwa, A., Dambrine, E., Pollier, B., and Atteia, O.: A comparison between Ca and Sr cycling in
961 forest ecosystems, *Plant and Soil*, 225, 299–310, <https://doi.org/10.1023/a:1026570812307>,
962 2000.
963

964 [Poszwa, A., Ferry, B., Pollie, B., Grimaldi, C., Charles-Dominique, P., Loubet, M., and Dambrine,](#)
965 [E.: Variations of plant and soil 87Sr/86Sr along the slope of a tropical inselberg, Ann. For. Sci. 66,](#)
966 [512, 10.1051/forest/2009036, 2009.](#)

967

968 [Potter, P., Ramankutty, N., Bennett, E.M., and Donner, S.D.: Characterizing the spatial patterns of](#)
969 [global fertilizer application and manure production, Earth Interact. 14, 1-22,](#)
970 <https://doi.org/10.1175/2009EI288.1>, 2010.

971

972 [Pouilly, M., Point, D., Sondag, F., Henry, M., and Santos, R. V.: Geographical Origin of Amazonian](#)
973 [Freshwater Fishes Fingerprinted by 87Sr/86Sr Ratios on Fish Otoliths and Scales, Environ. Sci.](#)
974 [Technol. 48\(16\), 8980-8987, 10.1021/es500071w, 2014.](#)

975

976 [Quaggio, C. S., Gastmans, D., Martins, V. T. de S., and Gilmore, T. E.: Combined use of statistical](#)
977 [Bayesian model and strontium isotopes deciphering the high complexity groundwater flow in the](#)
978 [Guarani Aquifer System \(GAS\), Appl. Geochem. 146, 105473, 10.1016/j.apgeochem.2022.105473,](#)
979 [2022.](#)

980

981 Reich, M.S., Ghouri, S., Zabudsky, S., Hu, L., Le Corre, M., Ng'iru I, Benyamini, D., Shipilina, D.,
982 Collins, S. C., Martins, D. J., Vila, R., Talavera, G., and Bataille, C. P.: Trans-Saharan migratory
983 patterns in *Vanessa cardui* and evidence for a southward leapfrog migration, *iScience*, 27(12),
984 111342, <https://doi.org/10.1016/j.isci.2024.111342> PMID: 39654635, 2024.

985

986

987 Salesse, K., Fernandes, R., de Rochefort, X., Brůžek, J., Castex, D., and Dufour, É.: isoarch.org: An
988 open-access and collaborative isotope database for bioarcheological samples from Graeco-Roman
989 World and its margins, ~~Journal of Archaeological Science: Reports~~, [J. Archaeol. Sci.: Rep.](#), 19, 1050-
990 1055, 2018.

991

992 Salesse, K., Fernandes, R., de Rochefort, X., Brůžek, J., Castex, D., and Dufour, É.: isoarch.org
993 (v.1.1), accessed 10/15/2025, 2020.

994

995 [Santos, R. V., Sondag, F., Cochonneau, G., Lagane, C., Brunet, P., Hattingh, K., and Chaves, J. G.](#)
996 [S.: Source area and seasonal 87Sr/86Sr variations in rivers of the Amazon basin, Hydrol. Process.](#)
997 [29\(2\), 187-197, 10.1002/hyp.10131, 2015.](#)

- Scaggion, C., Giovanardi, T., Palcsu, L., Cipriani, A., and Lugli, F.: [Bioavailable Sr isotopes of plants from Rio Grande do Sul and Santa Catarina states \(Brazil, South America\), related isoscape R script and raster files \(Version 02\)](#)~~Bioavailable Sr isotopes of plants from Rio Grande do Sul and Santa Catarina states (Brazil, South America) and related isoscape raster files~~ (Version 01) [Data set]. [Zenodo](#), 2025a. <https://doi.org/10.5281/zenodo.17988601>[Zenodo](#), <https://doi.org/10.5281/zenodo.17988601>, 2025a.
- Scaggion, C., Giovanardi, T., Loponte, D., Carbonera, M., Armaroli, E., Bernardini, S., Benazzi, S., Gascue, A., Acosta, A., Marciani, G., Bortolini, E., Cipriani, A., and Lugli, F.: Random forest-based bioavailable strontium isoscape for environmental and archaeological applications in central eastern Argentina and western Uruguay, *Plos One*, 20(7), e0326047, <https://doi.org/10.1371/journal.pone.0326047>, 2025b.
- Scherer, C. M S., Reis, A. D., Horn, B. L. D., Bertolini, G., Lavina, E. L. C., Kifumbi, C., and Aguilar, C. G.: The stratigraphic puzzle of the permo-mesozoic southwestern Gondwana: The Paraná Basin record in geotectonic and palaeoclimatic context, *Earth-Science Reviews*, Volume *Sci. Rev.* 240, 104397, ISSN 0012-8252, <https://doi.org/10.1016/j.earscirev.2023.104397>, 2023.
- Schwarcz, H. P., White, C. D., and Longstaffe, F. J.: Stable and Radiogenic Isotopes in Biological Archaeology: Some Applications. Isoscapes. Dordrecht: Springer Netherlands; p. 335–356. https://doi.org/10.1007/978-90-481-3354-3_16, 2010.
- [Seferidou, E.: Isotopic analysis in pre-colonial Bonaire, Dutch Caribbean \[data set\], https://doi.org/10.48530/isoarch.2025.006, 2025.](#)
- [Seferidou, E., Knippenberg, S., and Laffoon, J. E.: Reconstructing past lifeways of Indigenous individuals in pre-colonial Bonaire, through multi-isotope analysis, J. Isl. Coast. Archaeol. 20, 346-370, 10.1080/15564894.2023.2289190, 2023.](#)
- [Serna, A., Prates, L., Mange, E., Salazar-García, D. C., and Bataille, C. P.: Implications for paleomobility studies of the effects of quaternary volcanism on bioavailable strontium: A test case in North Patagonia \(Argentina\), J. Archaeol. Sci. 121, 105198, 10.1016/j.jas.2020.105198, 2020.](#)

1032 [Silva, C., dos Santos, E. A., Dussin, I. A., Montibeller, C. C., de Avelar Las Casas Rebelo, V., da](#)
1033 [Costa Pereira Lavalle Heilbron, M., Pimentel, L. C. G., and Landau, L.: Spatial distribution of](#)
1034 [strontium and neodymium isotopes in South America: a summary for provenance research, Environ.](#)
1035 [Earth Sci. 82, 348, 10.1007/s12665-023-11028-5, 2023.](#)

1036

1037 [Slovak, N. M., Paytan, A., Rick, J. W., and Chien, C.-T.: Establishing radiogenic strontium isotope](#)
1038 [signatures for Chavín de Huántar, Peru, J. Archaeol. Sci. Rep. 19, 411-419,](#)
1039 [10.1016/j.jasrep.2018.03.014, 2018.](#)

1040

1041 Spies, M. J., Alblas, A., Ambrose, S. H., Barakat, S., Barberena, R., Bataille, C. P., Bowen, G. J.,
1042 Britton, K., Cawthra, H., Diamond, R., Dosseto, A., Evans, J. A., Fisher, E., Gray, K., Heddell-
1043 Stevens, P., Holt, E., James, H. F., Janzen, A., Le Corré, M., le Roux, P., Lee-Thorp, J., Mackay, A.,
1044 McNeill, P. J., Montgomery, J., Mugabe, B., Oelze, V. M., Pfab, M., Richards, M. P., Samec, C. T.,
1045 Santana-Sagredo, F., Serna, A., Stantis, C., Snoeck, C., Stewart, B., Stuurman, C., Tarrant, D., West,
1046 A. G., Winter-Schuh, C., and Sealy, J.: Strontium isoscapes for provenance, mobility and migration:
1047 the way forward, ~~Royal Society~~[R. Soc. Open Science](#), 12(6), 12250283,
1048 <http://doi.org/10.1098/rsos.250283>, 2025.

1049

1050 [Stantis, C., Bataille, C., Bowen, G., James, H., Kafino, C., Salesse, K., Verostick, K., and Willmes,](#)
1051 [M.: The ARDUOUS dataset \[data set\], <https://doi.org/10.48530/isoarch.2024.002>, 2024.](#)

1052

1053 [Strauss, A., Oliveira, R. E., Villagran, X. S., Bernardo, D. V., Salazar-García, D. C., Bissaro, M.](#)
1054 [C., Pugliese, F., Hermenegildo, T., Santos, R., Barioni, A., de Oliveira, E.C., de Sousa, J.C.M.,](#)
1055 [Jaouen, K., Ernani, M., Hubbe, M., Inglez, M., Gratao, M., Rockwell, H., Machado, M., de Souza,](#)
1056 [G., Chemale, F., Kawashita, K., O'Connell, T.C., Israde I., Feathers, J., Campi, C., Richards, M.,](#)
1057 [Wahl, J., Kipnis, R., Araujo, A., and Neves, W.: Early Holocene ritual complexity in South America:](#)
1058 [the archaeological record of Lapa do Santo \(east-central Brazil\), Antiquity 90, 1454-1473,](#)
1059 [10.15184/aqy.2016.220, 2016.](#)

1060

1061 Strobl, C., Boulesteix, A.-L., Kneib, T., Augustin, T., and Zeileis, A.: Conditional variable
1062 importance for random forests, BMC Bioinformatics, 9, 307, [https://doi.org/10.1186/1471-2105-9-](https://doi.org/10.1186/1471-2105-9-307)
1063 [307](#) PMID: 18620558, 2008.

1064

1065 [Torres-Rouff, C., Pimentel, G., Pestle, W. J., Ugarte, M., and Knudson, K. J.: The Life and Death of](#)
1066 [a Child: Mortuary and Bodily Manifestations of Coast-Interior Interactions during the Late Formative](#)
1067 [Period \(AD 100-400\), Northern Chile, Lat. Am. Antiq. 33, 187-204, 10.1017/laq.2021.56, 2022.](#)
1068
1069 [Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C. U., Aas, W., Baker, A., Bowersox, V. C., Dentener,](#)
1070 [F., Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher,](#)
1071 [T., Mahowald, N. M., Nickovic, S., Rao, P. S. P., and Reid, N. W.: A global assessment of](#)
1072 [precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity](#)
1073 [and pH, and phosphorus, Atmos. Environ. 93, 3-100,](#)
1074 <https://doi.org/10.1016/j.atmosenv.2013.10.060>, 2014.
1075
1076 Vitousek, P. M., Kennedy, M. J., Derry, L. A., and Chadwick, O. A.: Weathering versus atmospheric
1077 sources of strontium in ecosystems on young volcanic soils, *Oecologia*, 121(2), 255–97,
1078 <https://doi.org/10.1007/s004420050927> PMID: 28308565, 1999.
1079
1080 Voerkelius, S., Lorenz, G. D., Rummel, S., Quérel, C. R., Heiss, G., Baxter, M., Brach-Papa, C.,
1081 Deters-Itzelsberger, P., Hoelzl, S., Hoogewerff, J., Ponzevera, E., Van Bockstaele, M., and
1082 Ueckermann, H.: Strontium isotopic signatures of natural mineral waters, the reference to a simple
1083 geological map and its potential for authentication of food, *Food Chemistry, Chem.* 118(4), 933–40,
1084 <https://doi.org/10.1016/j.foodchem.2009.04.125>, 2010.
1085
1086 [Washburn, E., Nesbitt, J., Ibarra, B., Fehren-Schmitz, L., and Oelze, V. M.: A strontium isoscape for](#)
1087 [the Conchucos region of highland Peru and its application to Andean archaeology, Plos One,](#)
1088 [e0248209, 10.1371/journal.pone.0248209, 2021](https://doi.org/10.1371/journal.pone.0248209)
1089
1090 Weiner, S., and Wagner, H. D.: THE MATERIAL BONE: Structure-Mechanical Function Relations,
1091 ~~*Annual Review of Materials Research*~~, *Annu. Rev. Mater. Res.*, 28(1), 271–98,
1092 <https://doi.org/10.1146/annurev.matsci.28.1.271>, 1998.
1093
1094 Wildner, W., Camozzato, E., Toniolo, J. A., Binotto, R. B., Iglesias, C. M. F., and Laux, J. H.: Mapa
1095 geologica do estado de Santa Catarina. Porto Alegre: CPRM. Escala 1:500000. Programa geologia
1096 do Brasil. Subprograma de Cartografia Geologica Regional, 2014.
1097

Formattato: Italiano (Italia)

Formattato: Italiano (Italia)

Formattato: Italiano (Italia)

Formattato: Italiano (Italia)

Codice campo modificato

1098 Wildner, W., Ramgrab, G. E., Lopes, R. C., Iglesias, C. M. F., and Laux, J. H.: Mapa geológico do
1099 estado de Rio Grande do Sul. Porto Alegre: CPRM. Escala 1:750000. Programa geologia do Brasil.
1100 Subprograma de Cartografia Geologica Regional, 2013.

1101

1102 White, P. J., and Broadley, M. R.: Calcium in plants, ~~Annals of Botany~~, Ann. Bot. 92(4), 487–511,
1103 <https://doi.org/10.1093/aob/mcg164> PMID: 12933363, 2003.

1104

1105 Wright, M. N., and Ziegler, A.: ranger: A Fast Implementation of Random Forests for High
1106 Dimensional Data in C++ and R, ~~Journal of Statistical Software~~, J. Stat. Softw. 77(1), 1-17,
1107 <https://doi.org/10.18637/jss.v077.i01>, 2017.

1108

1109 Zomer, R. J., Trabucco, A., Bossio, D. A., and Verchot, L. V.: Climate change mitigation: A spatial
1110 analysis of global land suitability for clean development mechanism afforestation and reforestation,
1111 Agric. Ecosyst. Environ. 126, 67-80, <https://doi.org/10.1016/j.agee.2008.01.014>, 2008.

1112

1113 **Supplement**

1114 Supplement Table S1: literature dataset of ⁸⁷Sr/⁸⁶Sr from South America used to construct the random
1115 forest models. The dataset is updated in September 2025.

1116

1117 Supplement Text S1: models and parameters for the ‘All’ and plant+fauna+lichen+human dataset.

Formattato: Bordo: Superiore: (Nessun bordo),
Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A
destra: (Nessun bordo), Tra : (Nessun bordo)