

A Pan-Arctic Pigment Database for Phytoplankton and Sea-Ice Algae

Authors

Asta C. Heidemann¹, Alexander Hayward², Philipp Assmy³, Atreya Basu⁴, Astrid Bracher^{5,6}, Giulia Castellani⁶, Giacomo Ditullio⁷, Katarzyna Dragańska-Deja⁸, Amane Fujiwara⁹, Glaucia Moreira Fragoso¹⁰, Jacob Høyer², Jiwoon Hwang¹¹, Morten Iversen⁵, Anabel von Jackowski¹², Thomas Juul-Pedersen¹³, Piotr Kowalczyk⁸, Youngju Lee¹⁴, Maria A. van Leeuwe¹⁵, Atsushi Matsuoka¹⁶, Alenya Merz^{13, 17}, C. J. Mundy¹⁸, Else Ostermann¹³, Ilka Peeken⁵, Matt Pinkerton¹⁹, Joanna Stoń-Egiert⁸, Jacqueline Stefels¹⁵, Antonia U. Thielecke⁵, Gaëlle Veyssiere²⁰, Hongyan Xi⁵, Eun Jin Yang¹⁴ & Rafael Gonçalves-Araujo¹

Institution

¹National Institute of Aquatic Resources, Technical University of Denmark (DTU Aqua), Kgs. Lyngby, Denmark

²Danish Meteorological Institute

³Norwegian Polar Institute, Fram Centre, Tromsø, Norway

⁴Centre for Earth Observation Science, University of Manitoba, Winnipeg, Canada

⁵Alfred-Wegener-Institute Helmholtz Center for Polar and Marine Research (AWI), Bremerhaven, Germany

⁶Institute of Environmental Physics (IUP), University Bremen (UB), Bremen, Germany

⁷University of Charleston, Grice Marine Laboratory

⁸Institute of Oceanology Polish Academy of Sciences, Sopot, Poland

⁹Institute of Arctic Climate and Environment Research, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Japan

¹⁰SINTEF Ocean, Department of Climate and Environment, Trondheim, Norway, 7041

¹¹University of California, Berkeley, USA

¹²Stockholm University, Stockholm, Sweden

¹³Greenland Climate Research Centre (GCRC), Greenland Institute of Natural Resources (GINR), Nuuk, Greenland

¹⁴Korea Polar Research Institute, Incheon, Republic of Korea

¹⁵University of Groningen, Groningen, the Netherlands

¹⁶Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, USA

¹⁷Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, Netherlands

¹⁸Centre for Earth Observation Science, Faculty of Environment, Earth and Resources, University of Manitoba, Winnipeg, Manitoba, Canada

¹⁹Earth Sciences New Zealand, Wellington, New Zealand

²⁰British Antarctic Survey, Cambridge, Cambridgeshire, UK

Correspondence to: Asta C. Heidemann (aheidem@aqu.dtu.dk), Alexander Hayward (algh@dm.dk) and Rafael Gonçalves-Araujo (rafgo@aqu.dtu.dk)

Abstract

Climate change has dramatically altered the Arctic seas with significant decrease in sea ice extent and thickness and warming water temperature. The ecological impacts of such change have been described for many parts of the Arctic Ocean, but long-term records of biological indicators are still missing. Among those, photosynthetic and accessory pigments are one of the key tools that aid quantification of phytoplankton and sea-ice algae biomass and characterisation of community composition. To address this gap, we present the first pan-Arctic compilation of in situ algal pigment data obtained exclusively by High-Performance Liquid Chromatography (HPLC), containing 10,798 samples collected across 77 Arctic research cruises between 2000 and 2024. As a result of large-scale collaborative effort, this database covers both open water and sea-ice environments across coastal, shelf and open domains. The database (<https://doi.org/10.11583/DTU.29445104>) includes measures of up to 26 pigments, with 8 major marker/accessory pigments being considered in this study, namely Alloxanthin (Allo), 19'-Butanoyloxyfucoxanthin (But-fuco), Chlorophyll a (Chl-a), Chlorophyll b (Chl-b), Fucoxanthin (Fuco), 19'-Hexanoyloxyfucoxanthin (Hex-fuco), Peridinin (Peri), and Zeaxanthin (Zea). This publicly available database provides crucial data that can be used to assess phytoplankton dynamics, validating remote sensing observations and can serve as a resource for future Arctic ecological- and modelling studies.

1. Introduction

Climate change has disproportionately affected the Arctic, shown by drastic increases in temperature and ongoing sea-ice decline (England et al., 2021; Stroeve and Notz, 2018). Changing conditions in the Arctic have already altered the food web, with studies indicating further impacts in the near future (Ardyna and Arrigo, 2020; Arrigo and van Dijken, 2015; Flores et al., 2023; Freer et al., 2022; Quinlan et al., 2005). As the foundation of marine food webs, phytoplankton and sea-ice algae are critical for sustaining ecosystems and contribute to large parts of global primary production, leading to energy transfer across trophic levels and ultimately carbon sequestration (Falkowski, 1994; Serra-Pompei et al., 2022). Environmental heterogeneity and changes may alter community composition, with the potential to impact large-scale ecological processes. For example, in the Arctic and sub-Arctic regions, increased open water areas has led to a longer phytoplankton growing season likely associated with changes in community composition. Studies have demonstrated that climate change may favor increased biomass of smaller flagellated phytoplankton species, leading to reduced dominance of large diatoms species (Blais et al., 2017; Coupel et al., 2012; Vonnahme et al., 2025). This can impact the structure of the food web and overall carbon cycling by having an immediate effect on higher trophic levels, potentially influencing the abundance, distribution and feeding preferences of zooplankton (Campbell et al., 2009; Negrete-García et al., 2024).

Characterising phytoplankton community composition is therefore essential for better understanding ecosystem structure, dynamics and services, especially in the under-sampled Arctic seas, which have experienced the greatest impact of climate change in recent years (England et al., 2021). It is clear that long-term data on microalgae physiology and biomass are vital to enhance our ability to assess and detect spatial and temporal patterns in phytoplankton and sea-ice algae community structure and physiology in response to environmental change. Many studies have used pigment-based measurements, but datasets are often scattered across many sources. This makes it difficult to obtain, access and synthesize into a larger spatial and temporal scale. To address this gap, our paper provides baseline information regarding phytoplankton and sea-ice algae biomass and photosynthetic pigments in the Arctic Ocean aiming at providing a framework to support such large-scale Arctic algae studies across two decades of data.

Phytoplankton pigments, particularly chlorophyll a (Chl-a), drives photosynthesis by harvesting light and protecting the cells (Falkowski and Kiefer, 1985; Pereira and Gonçalves, 2022). Advances in pigments analysis through High-Performance Liquid Chromatography (HPLC) and ultra HPLC (UHPLC), have enabled precise identification and quantification of a suite of photosynthetic and photoprotective microalgal pigments. These pigments can serve as chemotaxonomic markers for major functional

and taxonomic phytoplankton groups, including the commonly used diagnostic pigments, Alloxanthin (Allo), 19'-Butanoyloxyfucoxanthin (But-fuco), Chlorophyll *b* (Chl-*b*), Fucoxanthin (Fuco), 19'-Hexanoyloxyfucoxanthin (Hex-fuco), Peridinin (Peri), and Zeaxanthin (Zea) used in this study. Additionally and when available, the database presented in this paper also includes α -/ β -carotene ($\alpha\beta$ -Car), Bacteriochlorophyll *a*, Chlorophyllide *a* (Chlide *a*), Chl-*c*1, Chl-*c*1+*c*2, Chl-*c*3, Diadinoxanthin (Diadino), Diatoxanthin (Diato), Divinyl Chlorophyll *a* and *b* (DV-Chl-*a* and DV-Chl-*b*), Lutein (Lut), Neoxanthin (Neo), Pheophorbide *a* (Phide *a*), Pheophytin *a* (Phytin *a*), Prasinoxanthin (Pras) and Violaxanthin (Viola) (Jeffrey et al., 1999) (**Table 1**). These additional accessory pigments, some of which are also commonly used as marker pigments (Coupel et al., 2015; Fragoso et al., 2017), enable phytoplankton to adapt and/or acclimate to various light regime changes that occur in the environment (Brunet et al., 2011). Variability in pigment composition can reflect strategies such as photoprotection, which involves the production of pigments that mitigate damage from excessive light and photo acclimation that refers to the ability of algae to adjust their pigment composition in response to light intensity changes that hereby can optimise photosynthetic efficiency (Bonilla et al., 2009; Gosselin et al., 2017).

Absorption and scattering of light by phytoplankton pigments and other cellular constituents influence the reflectance, which allow their detection by radiometers, making them an invaluable source for remote sensing applications (see overview in Bracher et al., 2017). Multispectral satellite sensors such as SeaWiFS, MODIS, MERIS and OLCI have paved the way and demonstrated the ability to distinguish phytoplankton functional types on large scales (Miller, 2004; Nieke et al., 2015). Recent development have focused on hyperspectral sensors, capable of measuring more narrow absorption and reflectance properties, that can be missed by multispectral sensors (Hu, 2022). Successful applications include ENVISAT's Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY) using the PhytoDOAS method (Bracher et al., 2009; Sadeghi et al., 2012) and the DLR Earth Sensing Imaging Spectrometer Mission (DESIS) using the WebAssembly System Interface (WASI) (Gege, 2004, 2014) by (Bracher et al., 2021). Most recently NASA's Plankton, Aerosol, Clouds, ocean Ecosystem (PACE) mission have been launched (Werdell et al., 2024) and the European Space Agency (ESA) is preparing the CHIME hyperspectral to be launched in the coming decade (Nieke et al., 2023).

Since satellites provide long-term synoptic observations of phytoplankton groups high-quality in situ pigment data is invaluable for product validation and algorithm refinement (Cetinić et al., 2024). Similarly, several studies based on self-organizing maps, spectral composition or machine-learning-based methods applied to satellite data have leveraged phytoplankton pigments to understand taxonomic shifts in phytoplankton communities (El Hourany et al., 2019, 2024; Hayward et al., 2025; Xi et al., 2020).

Table 1. Alphabetically ordered list of the major diagnostic pigments and additional accessory pigments considered in this study (including abbreviations, their full name and associated phytoplankton groups) (Fragoso et al., 2017; Jeffrey et al., 1999). Notice that Dinoflagellates contains two types, reflecting different major pigments.

Major diagnostic pigments		
Abbreviation	Pigment name	Associated phytoplankton groups
Allo	Alloxanthin	Cryptophytes
But-fuco	19'-Butanoyloxyfucoxanthin	Haptophytes, Pelagophytes
Chl- <i>a</i>	Chlorophyll <i>a</i>	All phytoplankton groups
Chl- <i>b</i>	Chlorophyll <i>b</i>	Green algae (chlorophytes and prasinophytes)
Fuco	Fucoxanthin	Diatoms, Pelagophytes, Dinoflagellates-2, Haptophytes (certain species of <i>Phaeocystis</i>)
Hex-fuco	19'-Hexanoyloxyfucoxanthin	Haptophytes, Dinoflagellates-2
Peri	Peridinin	Dinoflagellates-1
Zea	Zeaxanthin	Cyanobacteria, Green algae
Additional accessory pigments		
Abbreviation	Pigment name	
$\alpha\beta$ -Car	α -/ β -carotene	
Bacteriochlorophyll <i>a</i>	Bacteriochlorophyll <i>a</i>	
Chlide <i>a</i>	Chlorophyllide <i>a</i>	
Chl- <i>c</i> 1	Chlorophyll <i>c</i> 1	
Chl- <i>c</i> 1+ <i>c</i> 2	Chlorophyll <i>c</i> 1+ <i>c</i> 2	
Chl- <i>c</i> 3	Chlorophyll <i>c</i> 3	
Diadino	Diadinoxanthin	
Diato	Diatoxanthin	
DV-Chl- <i>a</i>	Divinyl Chlorophyll <i>a</i>	
DV-Chl- <i>b</i>	Divinyl Chlorophyll <i>b</i>	
Lut	Lutein	
Neo	Neoxanthin	
Phide <i>a</i>	Pheophorbide <i>a</i>	
Phytin <i>a</i>	Pheophytin <i>a</i>	
Pras	Prasinoxanthin	
Viola	Violaxanthin	

While satellite data is becoming increasingly important, it remains limited by its ability to capture only the first optical depth under cloud- and ice-free conditions.

In situ pigment data from the Arctic have compared to other parts of the world historically been limited due to the region's remoteness, harsh weather conditions and logistical challenges of conducting fieldwork in such remote regions. Advances in HPLC technology, together with standardised protocols (Hooker et al., 2005) and round-robins

intercomparison of methodologies (Canuti, 2023), have contributed to the growing availability of pigment data globally, which also includes data from the Arctic and Sub-arctic regions (Kramer et al., 2022; Losa et al., 2017; Mattei and Scardi, 2021; Swan et al., 2016; Xi et al., 2023). Related efforts to obtain pigment data, such as those using UPLC methods (Hwang, 2025), also represent a valuable contribution, though they cannot be incorporated into the current database for inter-comparability reasons with HPLC-based measurements. Despite advances, spatial and temporal coverage across the Arctic remains uneven, underscoring persistent knowledge gaps, highlighting the need for coordinated, collective efforts to compile and expand existing data in this rapidly changing region.

To address this gap, we present the first pan-Arctic overview of phytoplankton and sea-ice algae pigment distributions, based on a comprehensive database of Arctic and sub-Arctic in situ pigment measurements collected between 2000 and 2024. The database is compiled from 77 cruises conducted across the Arctic Ocean, covering coastal, shelf, open-ocean and sea-ice domains (see **Table 2**). Importantly, this database represents a collective effort involving multiple institutions, research teams and international collaborations. By consolidating data from diverse sources, it provides a baseline for assessing long-term trends in phytoplankton pigments, communities and their ecological responses to environmental change. Additionally, such a database can support a pan-Arctic taxonomic characterisation of phytoplankton communities by employing pigment-based taxonomy techniques (Hayward et al., 2023, 2024; Vidussi et al., 2001; Wright and Jeffrey, 2006).

Methodology

2.1. Database Compilation

Phytoplankton and ice algae pigment data was compiled from 77 research cruises conducted between 2000 and 2024. The datasets were sourced from a combination of publicly available repositories (e.g., PANGAEA), institutional databases (e.g., NASA) and personal correspondence with scientists. Datasets were accompanied by relevant ancillary metadata, including information on sampling time, location and depth. A total of 10,798 pigment datapoints were compiled (**Table 2**). Where possible, cruise reports and original publications were consulted to verify the context and methods of data collection. Part of the data has already been published individually for the respective research cruises and/or projects, however, the compiled database presented in this article (including datasets not yet publicly available) has been stored in a public repository (<https://doi.org/10.11583/DTU.29445104>).

189
190
191

Table 2. Alphabetically ordered summary of Arctic and sub-Arctic cruises (= 77) with key dataset information such as year, sampling region(s) (as defined in section 3.1.), environment, sample count and associated DOI, where possible. Note that few samples were from melt-ponds, here assigned as ice samples.

Dataset	Year	Region(s)	Environment(s)	Number of Samples	DOI
ARA06B	2015	North Pacific Ocean, East Siberian Sea, Chukchi Sea, CAO	Water	112	https://dx.doi.org/doi:10.22663/KOPRI-KPDC-00002844
ARA07B	2016	North Pacific Ocean, East Siberian Sea, Chukchi Sea	Water	50	
ARA08B	2017	North Pacific Ocean, East Siberian Sea, Chukchi Sea	Water	98	
ARA09B	2018	East Siberian Sea, Chukchi Sea	Water	110	
ARA10B	2019	North Pacific Ocean, East Siberian Sea, Chukchi Sea	Water	70	
ARA10B_Ice	2018	East Siberian Sea	Ice	65	
ARA11B	2020	North Pacific Ocean, East Siberian Sea, Chukchi Sea	Water	85	
AREX2000	2000	Norwegian Sea, Barents Sea, Fram Strait	Water	77	https://doi.org/10.48457/IOPAN.2025.523
AREX2001	2001	Norwegian Sea, Barents Sea, Fram Strait	Water	55	
AREX2002	2002	Norwegian Sea, Fram Strait	Water	20	
AREX2003	2003	Norwegian Sea, Barents Sea, Fram Strait	Water	50	
AREX2004	2004	Norwegian Sea, Fram Strait	Water	27	
AREX2005	2005	Norwegian Sea, Barents Sea, CAO, Fram Strait	Water	51	
AREX2014	2014	Norwegian Sea, Barents Sea, CAO, Fram Strait	Water	133	
AREX2019	2019	Barents Sea, CAO, Fram Strait	Water	113	
AREX2021	2021	Norwegian Sea, Barents Sea, Fram Strait	Water	59	
AREX2022	2022	Norwegian Sea, Barents Sea, CAO, Fram Strait	Water	140	

Academik_loffe_2003	2003	Greenland Sea, Labrador Sea, Norwegian Sea	Water	34	
Arc2002	2002	Beaufort Sea, Chukchi Sea	Water	53	
Arc2004	2004	Beaufort Sea, Chukchi Sea	Water	90	
Arctic_Ocean_2024	2024	CAO	Ice, Water	61	https://doi.org/10.48457/IOPAN.2025.386
Arex2017	2017	CAO, Fram Strait	Water	33	
Assmy_ICE2010	2010	CAO	Water	42	
BODC/NERC_2001	2001	Greenland Sea	Water	169	http://doi.pangaea.de/10.1594/PANGAEA.793246
BODC/NERC_2002	2002			419	
BaySys	2018	Hudson Bay, Labrador Sea	Ice, Water	131	https://canwin-datahub.ad.umanitoba.ca/data/dataset/algae-pigments-icewater
Bracher2021_PS126	2021	Norwegian Sea, Fram Strait	Water	175	https://doi.pangaea.de/10.1594/PANGAEA.982954
Bracher_2017_PS106	2017	Norwegian Sea, Barents Sea, CAO, Fram Strait	Ice, Water	40	https://doi.org/10.1594/PANGAEA.899284
Bracher_2019_PS121	2019	Norwegian Sea, Fram Strait	Water	298	https://doi.org/10.1594/PANGAEA.941011
Bracher_2020_MSM93	2020	Norwegian Sea, Greenland Sea, Fram Strait	Water	219	
Bracher_2022_PS131_A TWAICE	2022	Norwegian Sea, Greenland Sea, CAO, Fram Strait	Water	241	https://doi.pangaea.de/10.1594/PANGAEA.983242
Bracher_2023_PS136_HAUSGARTEN	2023	Norwegian Sea, Fram Strait	Water	137	https://doi.pangaea.de/10.1594/PANGAEA.983243
Bracher_2024_PS143.2	2024	Norwegian Sea, CAO, Fram Strait	Water	212	https://doi.pangaea.de/10.1594/PANGAEA.983244
CFL	2008	Beaufort Sea, Canadian Archipelago	Water	75	
DANA_ECOTIP_2021	2021	Baffin Bay, Labrador Sea	Water	16	https://doi.org/10.48457/IOPAN.2025.523
DSOS_GINR	2024	Baffin Bay, Labrador Sea	Water	234	
DiTuillio_2018	2018	CAO	Water	132	10.18739/A2028PD2H
DFO-Fragoso_2005	2005	Labrador Sea	Water	25	https://doi.pangaea.de/10.1594/PANGAEA.871872 For DFO-Fragoso (2005-2013), also see: https://waves-vagues.dfo-mpo.gc.ca/Library/354345.pdf
DFO-Fragoso_2006	2006	Labrador Sea	Water	12	
DFO-Fragoso_2007	2007	Labrador Sea	Water	32	
DFO-Fragoso_2008	2008	Labrador Sea	Water	25	
DFO-Fragoso_2009	2009	Labrador Sea	Water	26	
DFO-Fragoso_2010	2010	Labrador Sea	Water	27	
DFO-Fragoso_2011	2011	Labrador Sea	Water	33	
DFO-Fragoso_2012	2012	Labrador Sea	Water	30	
DFO-Fragoso_2013	2013	Labrador Sea	Water	27	
Fragoso_2014	2014	Greenland Sea, Labrador Sea	Water	16	
GreenEdge_Amundsen	2016	Baffin Bay, Labrador Sea	Water	589	10.17882/86417
Icescape_2010	2010	Chukchi Sea	Water	100	

Icescape_2011	2011	Beaufort Sea, Chukchi Sea	Water	455	https://seabass.gsfc.nasa.gov/experiment/ICESCAPE
Liu2015_PS93.2	2015	Norwegian Sea, Fram Strait	Water	241	
Liu_2016_PS99.1	2016	Norwegian Sea, Fram Strait	Water	96	https://doi.pangaea.de/10.1594/PANGAEA.905502
Liu_2016_PS99.2	2016	Fram Strait	Water	230	https://doi.pangaea.de/10.1594/PANGAEA.894874
Liu_2017_PS107	2017	Norwegian Sea, Fram Strait	Water	262	https://doi.pangaea.de/10.1594/PANGAEA.894860
MERIAN_ECOTIP2022	2022	Greenland Sea, Fram Strait	Water	40	https://doi.org/10.48457/IOPAN.2025.523
MOSAIC_Algal	2019, 2020	CAO, Fram Strait	Ice, Water	260	FYI samples: https://doi.pangaea.de/10.1594/PANGAEA.967448 SYI samples: https://doi.pangaea.de/10.1594/PANGAEA.967450 Open water samples: https://doi.org/10.1594/PANGAEA.955763
MOSAIC_SYI	2019, 2020	CAO	Ice	275	
MOSAIC_FYI	2019, 2020	CAO, Fram Strait	Ice	285	
MR17-05	2017	Beaufort Sea, North Pacific Ocean, Chukchi Sea	Ice, Water	74	https://www.godac.jamstec.go.jp/darwin_cruise/view/metadata?key=MR17-05C&lang=en
Malina	2009	Baffin Bay, Labrador Sea, Beaufort Sea, Canadian Archipelago, Laptev Sea, North Pacific Ocean, Chukchi Sea	Water	677	
Malina_Barge	2009	Beaufort Sea	Water	53	https://www.seanoe.org/data/00641/75345/
N-ICE2015	2015	CAO	Water	145	https://doi.org/10.21334/NPOLAR.2016.3EBB7F64
NunaWP4Mackenzie	2019	Beaufort Sea	Water	140	https://doi.org/10.1594/PANGAEA.937585
Thielecke_PS138_ArcWatch	2023	CAO	Ice, Water	151	
Peeken, PS74	2009	Norwegian Sea, Greenland Sea, Fram Strait	Water	130	
Peeken, PS76	2010	Norwegian Sea, Greenland Sea, Fram Strait	Water	549	
Peeken, PS78	2011	Fram Strait	Water	264	
Peeken, PS80	2012	Laptev Sea, CAO, Fram Strait	Water	576	
Peeken, PS85	2014	Fram Strait	Water	82	
Peeken, PS92	2015	CAO	Water	106	

Polarstern2003	2003	Greenland Sea, Norwegian Sea	Water	26	https://doi.org/10.48457/IOPAN.2025.523
SeaBASS_2004	2004	Beaufort Sea, Chukchi Sea	Water	59	
SeaBASS_2008	2008	Greenland Sea	Water	43	
SeaBASS_2013	2013	Baffin Bay, Labrador Sea, Norwegian Sea, Beaufort Sea, Barents Sea, East Siberian Sea, Canadian Archipelago, Chukchi Sea, Greenland Sea, Kara Sea	Water	77	
Sublce	2014	Chukchi Sea	Water	155	
Tara_Arctic	2013	Baffin Bay, Labrador Sea, Norwegian Sea, East Siberian Sea, Barents Sea, Canadian Archipelago, Greenland Sea, Kara Sea	Water	229	
Woodfjorden	2024	Fram Strait	Water	50	https://doi.org/10.48457/IOPAN.2025.504

2.2. Data Quality Measures

This dataset integrated HPLC pigment measurements from 77 research cruises, each carried out by laboratories using their own established laboratory procedures, including different HPLC set-ups, calibration standards and water filtration volume.

Methodological descriptions associated with each dataset can be accessed through the provided DOIs in **Table 2** or by contacting the original data providers. To ensure consistency and comparability across sources, we applied a set of inclusion/exclusion criteria and harmonisation procedures to ensure data consistency and comparability across sources. Only datasets with measurements of seven out of the eight marker pigments described in **Table 1** were included in the database, allowing for reliable inter-sample comparison of pigment composition (criterion 1, see **Figure 1**). Although zeaxanthin is normally included in the analysis of phytoplankton functional types, it was excluded as a formal requirement in this database compilation because one source lacked measurement of this pigment and reliable distinction of green algae from other phytoplankton groups remain possible without zeaxanthin (see available pigments for each cruise in Supplementary **S1**). Zeaxanthin is also a major diagnostic pigment for *Synechococcus*, which is largely absent from the Arctic region due to its thermal dependence on warmer waters (Six et al., 2021). Samples were excluded from the final database if they reported Chlorophyll (Chl-*a*) concentration of zero (criterion 2, see

Figure 1), lacked measurements of the required marker pigments and/or were missing essential metadata such as geographic coordinates (criterion 3, see **Figure 1**).




Figure 1. Workflow diagram displaying the major decisions in the data compilation process.

In addition to data filtering, multiple data harmonisation steps were also undertaken: longitudes reported in degrees East (0° to 360°E) were converted to signed longitude (-180° to 180°E/W) to ensure spatial compatibility with standard mapping tools ; datasets were also checked for true 0s and NaNs and that the distinction between these were standardised across datasets. This process required manual inspection of each dataset: If a pigment was entirely absent from a dataset, it was treated as NaNs (empty cells), whereas if a pigment was present but only had a single non-zero value reported, the remaining empty cells were interpreted as true 0s; finally, concentration units for accessory pigments and Chl-a were also standardised to the common unit, µg/L. There was considerable variability in methods used to calculate total Chl-a was observed across the database, and these methods were not always reported. Consequently, this parameter was excluded. In addition, only a limited number of melt pond samples were available. These were therefore treated as part of the ice sample category, instead of a category by itself.

2.3. Limitations and Uncertainties of the Dataset

Despite efforts to minimize both limitations and uncertainties, this dataset consists of HPLC measurements from multiple laboratories. We acknowledge that methodological

variability between laboratories exists and will introduce variability to the dataset. The following sections are provided to summarise the main known sources of limitations and uncertainties associated with compiling HPLC data from various laboratories, as well as general uncertainties inherent to pigments measurements.

Analytical

All datasets presented in this paper have followed standard protocols for HPLC analysis of phytoplankton pigments (see Supplementary Material S2). However, analytical uncertainties can arise from multiple stages of the workflow. As outlined by Wright & Jeffrey (2006), these can broadly be categorised into:

- **Collection and storage**

Filtration volume, filter type and size, storage temperature and storage time

- **Extraction procedures**

Sample disruption, duration and temperature, and choice of solvents

- **HPLC methods and techniques** (see review of HPLC pigment methods in Roy et al., 2011)

Variations of column type and flow rate etc.

- **Peak detection, identification, integration and quantification**

Differences in baseline correction, peak threshold, co-elution handling and pigment standards used for calibration

To minimize bias in each of these steps, substantial effort has been made through inter-comparability studies. Most notably are the SeaWiFS HPLC Analysis Round-Robin Experiments (SeaHarre-1 to SeaHarre-5) (Hooker et al., 2012) and HPLC/DAD intercomparison on phytoplankton pigments (HIP-1 to HIP-4) (Artuso, F et al., 2016; Canuti et al., 2016), which efforts has been to eliminate differences in pigment quantification across laboratories and quantify these uncertainties in terms of what can be expected, even when conducting analysis according to strict protocols.

Pigment-Specific

In addition to the analytical sources of uncertainties, pigment-specific sources also exist because of a varying degree of sensitivity to the above-described workflow. Any chosen HPLC method can therefore increase or reduce the variance experienced for a specific pigment. However, defining robust “uncertainty values” for each individual pigment remains a challenge across analytical methods. Recent inter-laboratory comparisons highlight this challenge, while also underscoring the need of such estimates in applications such as remote sensing validation (e.g., Canuti, 2023).

A main source of pigment-specific behavior comes from differences in the chemical stability of the pigments. This will affect their susceptibility to degradation, co-elution and extraction efficiency. For example, Chlorophylls (a, b, c₁ and c₂) are prone to degradation, particularly during storage and extraction, producing degradation products such as pheophytins and chlorophyllides. This can lead to biased and unprecise chlorophyll estimations as well as co-elution issues. This have often been reported for chlorophyll c₁ and chlorophyll c₂ as well as for the Fuco derivatives (But-fuco and Hex-fuco) (Roy et al., 2011; Simmons et al., 2016; Wright and Jeffrey, 2006; Zapata et al., 2000). Additionally, some chlorophyll degradation products, for example Phide a and Phytin a, can co-elute with carotenoids due to polarity changes (Mendes et al., 2007).

Finally, pigment concentration itself can be a major source of uncertainty. This is particularly true for low-abundance accessory pigments, which often approach the instrumental limits of detection (LOD) (Bidigare et al., 2002; Canuti, 2023; Canuti et al., 2016; Claustre et al., 2004; Hooker et al., 2005). For instance, when pigment concentrations fall below 0.01 mg m⁻³, the relative standard deviation can reach up to nearly double that of pigments present at higher concentrations (Claustre et al., 2004).

3. Data Description

3.1. Sampling Coverage

To classify the sampling efforts across the Arctic and sub-Arctic seas into common groups, the study area was subdivided into 15 regions (**Figure 2**). A total of 10,095 water samples and 703 sea-ice samples were compiled and mapped according to their geographic location. Water samples displayed extensive sampling across most regions, while ice samples were comparatively sparse and primarily concentrated in the Central Arctic Ocean (CAO), Canadian Archipelago and Hudson Bay. These samples originate from a limited number of cruises and are not consistently represented throughout the years. Regional sampling frequencies are summarised in **Figure 2**, highlighting the dominance of water samples and the uneven spatial distribution of sea-ice samples.




Figure 2. Map showing the spatial distribution of water and sea-ice samples across 15 defined regions of the Arctic and sub-Arctic seas. Each point represents an individual sampling location.

Overall, the water sample collection also varied substantially across years, with higher coverage in 2002, 2009, 2016 and 2020 (**Figure 3**). These periods correspond to large-scale, multidisciplinary campaigns such as MALINA (Coupel et al., 2015), GreenEdge (Bruyant et al., 2022; Massicotte et al., 2020), the AR7W transect during the MOSAiC cruise and others, including global initiatives such as the International Polar Year (IPY). Additionally, the MAREDAT pigment database represents a major synthesis effort (Peloquin et al., 2013) (see **Table 1**).




Figure 3. (Left) yearly sampling frequency for water column and sea-ice samples, (middle) monthly sampling frequency and (right) number of water samples observations per depth bin. Note that for sea-ice samples, the vertical distribution is not presented due to inconsistencies in the ice core sampling methods among the different campaigns.

Seasonally, sampling frequency was the highest in spring and summer (May-August). Sampling frequency was lower in September, followed by a notable secondary drop in December. The lowest sampling frequency is observed during winter and early spring (December-March), when there is extensive sea ice extent. Sampling frequency also varied vertically in the water column (**Figure 3**). Most samples were collected within the upper 1-15 meters range. Sampling frequency consistently declines from 21-30 meters onward, before a slight increase in the 91-100 meters range.

3.2. Pigment Distribution

Pigment concentration in water and sea-ice samples exhibited considerable variability in the biomass and composition of algal communities as well as in the spatiotemporal structure of the database (**Table 3**). In addition, the observed variability can also be attributed to adaptive pigment strategies.

Since direct comparison of absolute pigments concentrations between sea-ice and water samples should be interpreted with caution, only a brief description of absolute pigment concentration is provided for context (**Table 3a**). Chl-*a*, the primary indicator of phytoplankton and sea-ice algae biomass, had a mean concentration of 0.997 $\mu\text{g/L}$ in water samples and 0.639 $\mu\text{g/L}$ in sea-ice samples although the highest individual values in both environments point to episodic blooms. These values represent the mean of all individual concentration measurements in the compiled dataset. The highest concentration across the database, 32.76 $\mu\text{g/L}$, was recorded in the Chukchi Sea originating from the Ice Scape dataset in August 2010. In general, such high concentrations are rare, with the majority of samples falling below 0.5 $\mu\text{g/L}$ across seasons. Overall, concentrations were highly variable with 90% of values falling between 0.017 and 3.789 $\mu\text{g/L}$ in water samples and between 0.022 and 0.292 $\mu\text{g/L}$ in sea-ice samples (**Table 3a**). It should be recognised that there is a potential bias when comparing open-water and bulk-ice samples due to neglect of the dilution of pigments

when bulk sea-ice cores are melted, which can underestimate concentrations from sympagic algae (Chamberlain et al., 2022; Gradinger, 2009; Miller et al., 2015).

In terms of the relative ratios to Chl-*a* (**Table 3b and Figure 4**) the database also showed marked seasonal variability. Among the eight analysed marker pigments, Fuco consistently had the highest ratios, particularly during winter months, followed by a decline through summer and early autumn. The Fuco-derivatives, But-fuco and Hex-fuco, both considered diagnostic markers of haptophytes, had a stable distribution throughout the year, aside from notable peaks in November.

The ratio of Allo, a distinct marker pigment of cryptophytes, started to increase in late winter, prior to the main growing season. Peri, which is a marker pigment of dinoflagellates, had low overall ratios. These ratios increased during summer, peaking in October, before they exhibited a sharp decline toward the onset of December. As expected, Zea and Chl-*b* were relatively aligned throughout most of the year, consistent with their shared role as a marker pigment of green algae. However, a marked dissimilarity in their patterns occurred in November, when Chl-*b* declined while Zea increased.

For the sea-ice samples, the only accessory pigments with high ratios are Fuco and Chl-*b*.

These differences suggest distinct pigment signatures between sea-ice and water samples, underscoring the importance of expanding sea-ice sampling efforts to better capture algae dynamics in this environment.

Table 3. Summary of 5th, 95th, mean, median and std statistics across all cruises (= 77), for (a) pigment concentration categorized by sample type and (b) relative ratio of accessory pigments vs. Chl-*a*. For detailed pigment information by individual cruises, see Supplementary **Table S1**. For detailed information about all pigments across environments, see Appendices **Table A1**

(a) Water Samples (µg/L)						Sea-ice Samples (µg/L)				
Parameter / Pigment	5 th percentile	95 th percentile	Mean	Median	Std	5 th percentile	95 th percentile	Mean	Median	Std
Allo	0.000	0.061	0.014	0.003	0.031	0.000	0.016	0.004	0.000	0.020
But-fuco	0.000	0.080	0.020	0.007	0.040	0.000	0.005	0.001	0.000	0.008
Chl- <i>a</i>	0.017	3.789	0.997	0.417	1.855	0.022	2.292	0.639	0.179	1.746
Chl- <i>b</i>	0.000	0.207	0.059	0.033	0.091	0.000	0.158	0.037	0.000	0.109
Fuco	0.003	1.493	0.357	0.106	0.853	0.000	1.392	0.367	0.063	1.120
Hex-fuco	0.000	0.434	0.092	0.019	0.180	0.000	0.011	0.002	0.000	0.010
Peri	0.000	0.105	0.021	0.001	0.072	0.000	0.057	0.016	0.000	0.078

Zea	0.000	0.030	0.008	0.001	0.025	0.000	0.009	0.003	0.000	0.022
(b) Water Samples (ratio)						Sea-ice Samples (ratio)				
Parameter / Pigment	5 th percentile	95 th percentile	Mean	Median	Std	5 th percentile	95 th percentile	Mean	Median	Std
Allo vs Chl- <i>a</i>	0.000	0.078	0.100	0.009	1.634	0.000	0.034	0.006	0.000	0.021
But-fuco vs Chl- <i>a</i>	0.000	0.160	0.263	0.024	4.663	0.000	0.013	0.003	0.000	0.030
Chl- <i>b</i> vs Chl- <i>a</i>	0.000	0.358	1.094	0.077	15.482	0.000	0.414	0.087	0.000	0.138
Fuco vs Chl- <i>a</i>	0.068	0.666	2.504	0.290	26.620	0.000	0.741	0.428	0.443	0.235
Hex-fuco vs Chl- <i>a</i>	0.000	0.489	0.681	0.080	9.694	0.000	0.007	0.004	0.000	0.018
Peri vs Chl- <i>a</i>	0.000	0.118	0.221	0.004	5.792	0.000	0.150	0.028	0.000	0.109
Zea vs Chl- <i>a</i>	0.000	0.056	0.056	0.003	1.062	0.000	0.027	0.005	0.000	0.021

Pigment:Chl-a ratio




Figure 4. Display of the seasonal variation of individual accessory pigment concentration to Chl-a ratio. Water samples are shown as monthly ratios in blue and sea-ice samples are shown overall across months per pigment in green. Only values within 5th to 95th percentile are included. Data points are displayed as open circles, the mean ratio for each month is shown as a red dot and an orange line connects means.

Spatial and Temporal Dynamics

In our dataset, Chl-a exhibited strong spatial and temporal (seasonal) and inter-annual variability, reflecting the dynamic nature of phytoplankton and sea-ice algae in polar and subpolar environments. Elevated concentrations were observed in areas such as the Fram Strait, Barents Sea, Beaufort Sea, Chukchi Sea and Labrador Sea (**Figure 5**). However, as the database is opportunistic by nature, covering only discrete regional samplings, spatial and seasonal comparisons should be made with caution.

In spring, the Labrador Sea recorded the seasonal maximum (18.75 $\mu\text{g/L}$), with the highest values occurring mostly in surface waters (0-5 meters). As previously mentioned, summer exhibited the maximum database record for Chl-*a*. Contrary to spring, highest averages were found between 16-30 meters. By autumn, Chl-*a* had declined across most regions. However, localised high biomass persisted in some areas such as Beaufort Sea, Chukchi Sea and Fram Strait. Winter Chl-*a* was generally low.




Figure 5. (a) Spatial distribution of all measured Chl-*a* water samples in each season, shown as subplots for spring, summer, autumn and winter. Color mapping uses logarithmic scale, but the values themselves are not log-transformed. (b) Mean Chl-*a* concentration as function of depth range (rows) and season (columns). Color indicates the average value for each depth/season bin. Missing color indicates that data is not available. (c) Spatial distribution of all measured Chl-*a* for sea-ice samples.

The concentration frequency distribution plots of water samples (**Figure 6**, blue plots) showed that both Chl-*a*, Chl-*b* and Fuco exhibited approximately log-normal distribution, with peak concentrations near central values and tails on either side. This interpretation was supported by the log-normal distribution of data. However, formal normality tests such as Kolmogorov-Smirnov (K-S test) indicated significant deviations from perfect log-normality, likely due to large sample size and/or the presence of outliers (see Appendices **Figure A1**). In contrast, pigments such as Zea, Peri and the Fuco derivatives exhibited more variable distributions and greater heterogeneity among samples. The histograms for the sea-ice samples (**Figure 6**, grey plots), had much more variability in the distribution of concentrations. This is likely also related to smaller sample size for sea-ice samples.




Figure 6. Frequency histograms displaying log-scaled pigment concentrations. Water samples are shown in blue and sea-ice samples are shown in grey. x-axis represents actual pigment concentrations. Note the different ranges for y-axis comparing water and sea-ice samples.

4. Data Availability

All datasets used in this paper have been compiled into a single repository, accessible using DOI: <https://doi.org/10.11583/DTU.29445104> (Heidemann et al., 2026).

5. Conclusions

Photosynthetic and accessory pigments serve as indicators of community structure and biomass, and are important oceanographic variables for understanding ocean ecology, as they can also support assessment of biogeochemical cycling. Yet, datasets reporting concentrations of those pigments remain sparse in the Arctic region. To address this gap, we compiled, filtered, standardised and harmonised 10,798 pigment samples from HPLC, including water (n = 10,095) and sea-ice samples (n = 703), collected during 77 research cruises spanning from 2000 to 2024 across 15 defined Arctic and sub-Arctic regions. Sampling frequency varied by year, with peak efforts in 2002, 2009, 2016 and 2020. These years correspond to major multidisciplinary research projects and international initiatives such as the International Polar Year (IPY). Seasonally, most samples were collected during spring and summer months in the upper water column.

Within the available dataset, Chl-a was found in higher concentrations in sea-ice samples compared to water samples, on average. However, for accessory pigments, concentrations were generally found to be lower in sea-ice samples. Despite this, the

highest concentration of Chl-*a* across the dataset, 32.76 µg/L, was recorded for a water sample in the Chukchi Sea. Other areas that exhibited elevated Chl-*a* concentrations were spread across the Arctic marginal seas such as the Fram Strait, Barents Sea, Beaufort Sea and Labrador Sea. Interestingly, among the eight marker pigments, Fuco consistently had the highest pigment to Chl-*a* ratios, particularly during winter months, followed by a decline through summer and early autumn. For the sea-ice samples, the only accessory pigments with high ratios were Fuco and Chl-*b*.


Our comprehensive phytoplankton and sea-ice algal HPLC-based pigment dataset provides valuable insights into the research effort in the Arctic Ocean. Furthermore, it offers important observations on spatial and temporal patterns of primary producers supporting large-scale environmental research in the Arctic across over two decades of sampling efforts.



Appendices

Table A1. *Pigment concentration statistics for all included measured pigments, separated by sample type and summarised across all cruises. For each pigment and environment, the table reports the 5th percentile, 95th percentile, mean, median and standard deviation (Std).*



Parameter / Pigment	Water Samples (µg/L)					Sea-ice Samples (µg/L)				
	5 th percentile	95 th percentile	Mean	Median	Std	5 th percentile	95 th percentile	Mean	Median	Std
Chl c3	0.000	0.232	0.054	0.014	0.116	0.000	0.051	0.014	0.000	0.053
Chl c1c2	0.000	0.652	0.146	0.048	0.295	0.000	0.365	0.101	0.035	0.254
Chl c1	0.000	0.031	0.009	0.000	0.061	0.000	0.000	0.000	0.000	0.007
Chlide <i>a</i>	0.000	0.062	0.015	0.000	0.078	0.000	0.007	0.002	0.000	0.022
Peri	0.000	0.105	0.021	0.001	0.072	0.000	0.057	0.016	0.000	0.078
Phide <i>a</i>	0.000	0.132	0.028	0.000	0.147	0.000	0.437	0.132	0.000	0.821
But-fuco	0.000	0.080	0.020	0.007	0.040	0.000	0.005	0.001	0.000	0.008
Fuco	0.003	1.493	0.357	0.106	0.853	0.000	1.392	0.367	0.063	1.120
Neo	0.000	0.022	0.005	0.000	0.010	0.000	0.014	0.002	0.000	0.010
Pras	0.000	0.053	0.011	0.000	0.025	0.000	0.005	0.001	0.000	0.006
Viola	0.000	0.027	0.007	0.002	0.012	0.000	0.034	0.010	0.000	0.049
Hex-fuco	0.000	0.434	0.092	0.019	0.180	0.000	0.011	0.002	0.000	0.010
Diadino	0.000	0.232	0.058	0.020	0.108	0.000	0.239	0.052	0.000	0.208
Allo	0.000	0.061	0.014	0.003	0.031	0.000	0.016	0.004	0.000	0.020
Diat	0.000	0.025	0.005	0.000	0.016	0.000	0.063	0.010	0.000	0.036
Zea	0.000	0.030	0.008	0.001	0.025	0.000	0.009	0.003	0.000	0.022
Lut	0.000	0.011	0.003	0.000	0.012	0.000	0.025	0.009	0.000	0.069
bacterio chl <i>a</i>	0.000	0.000	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.000
DV Chl <i>b</i>	0.000	0.000	0.000	0.000	0.004	0.000	0.000	0.000	0.000	0.000
Chl- <i>b</i>	0.000	0.207	0.059	0.033	0.091	0.000	0.158	0.037	0.000	0.109
DV Chl <i>a</i>	0.000	0.000	0.001	0.000	0.007	0.000	0.000	0.000	0.000	0.000
Chl- <i>a</i>	0.017	3.789	0.997	0.417	1.855	0.022	2.292	0.639	0.179	1.746
Phytin <i>a</i>	0.000	0.034	0.007	0.000	0.047	0.000	0.000	0.007	0.000	0.048
alphabetaCar	0.000	0.067	0.018	0.006	0.032	0.000	0.085	0.017	0.000	0.064

Figure A1. Histogram and Q-Q plots performed for each marker pigment on standardised and log-transformed data. For each pigment, a histogram and a Q-Q plot are visualized to assess normality. All Kolmogorov-Smirnov (K-S) *p*-values were below 0.05 test, indicating that the data deviates from a normal distribution.









461



462



Author contributions

ACH consolidated the data, harmonized and standardised the dataset, performed the data analysis, generated plots and developed the manuscript.

AH conceived the study idea, provided relevant expertise, contributed data, facilitated key contacts and data sources and contributed to revising and improving the data analysis and manuscript.

RG-A provided relevant expertise and contributed to conceiving the study idea, revising and improving the data analysis and manuscript.

PA, AB (Astrid Bracher), GC, GD, KDD, AF, GMF, JH (Jacob Høyer), MI, AJ, TJP, PK, YL, MAVL, AM (Atsushi Matsuoka), AM (Alenya Merz), CJM, EO, IP, MP, JSE, JS, AT, GV, HX and EJY contributed data, provided relevant expertise and revised the manuscript.

AB (Atreya Basu) and JH (Jiwoon Hwang) provided relevant expertise and revised the manuscript.

Acknowledgement

ACH and RG-A were supported by the Independent Research Fund Denmark, through the project WaterColor (Gran No. 4251-00058B). RG-A was also supported by the European Union's Horizon Europe research and innovation program under Grant Agreement No. 101136480 (SEA-Quester).

PA was supported by the Research Council of Norway (project no. 244646) and the Ministry of Foreign Affairs, Norway, through the ID Arctic project.

AM was supported by NASA PACE (grant #: 80NSSC26K0091) and JAXA's GCOM-C/SGLI (25RT000334) projects.

YL was supported by Korea Institute of Marine Science and Technology Promotion (KIMST) grant funded by the Ministry of Oceans and Fisheries (KIMST RS-2021-KS211500, Korea-Arctic Ocean Warming and Response of Ecosystem, KOPRI).

KDD, JSE and PK were supported by a project funded by the Polish National Science Centre (NCN) OPUS26 project OptiCal-Green (2023/51/B/ST10/01344) granted to PK. Additional funding has been provided by the European Union's Horizon Europe research and innovation program under Grant Agreement No. 101136480 (SEA-Quester) and under Grant Agreement No. 101136748 (BioEcoOcean), as well as the statutory research program of the Institute of Oceanology, Polish Academy of Sciences (tasks I.2 and II.5). The ship time on board of r/v Kronprins Haakon was provided by the Norwegian Polar Institute, Tromsø, Norway.

GMF was supported by Fisheries and Oceans Canada (DFO). We acknowledge the Atlantic Zone Offshelf Monitoring Program, the captains and crew of the Canadian Coast Guard vessel CCGS *Hudson*, and DFO staff for their contributions to data collection during the Labrador Sea cruises and laboratory analysis at the Bedford Institute of Oceanography

AB and HX were supported by the Helmholtz Infrastructure Initiative FRAM, German Research Foundation (DFG) "Transregional Collaborative Research Center ArctiC Amplification: Climate Relevant Atmospheric and SurfaCe Processes and Feedback Mechanisms (AC)3" (Project C03), Copernicus Marine – Innovation Service Evolution R&D Project ML-PhyTAO (23138L03D-COP-INNO SCI-9000), and the ESA project Phyto-CCI(4000147645/25/I-LR). We thank the captain, the crew and all scientists of RV Polarstern expeditions PS93-2, PS99-2, PS107, PS121, PS126, PS131, PS136 and PS143-2 for their support in preparation and on board the expeditions.

IP was supported by the PoF III & IV program providing ship time on RV Polarstern [Changing Earth –Sustaining our Future, Topic 6.1 of the Helmholtz Association]

515 AT was supported by the Helmholtz Young Investigator Group SiDe-Effect (VH-NG-1600)
516 awarded to Prof. Mar Fernández-Méndez. PS138 (ArcWatch I) was supported by the
517 Helmholtz Research Programme “Changing Earth – Sustaining our Future” Topic 6,
518 Subtopics 1, 2 and 3 (Grant No. AWI_PS138_02).

519 JS and MAVL were supported by the Dutch Research Council (NOW), through the
520 Netherlands Polar Programme (NPP), Project no 866. 18. 002.

521

522 References

523 Ardyna, M. and Arrigo, K. R.: Phytoplankton dynamics in a changing Arctic Ocean, *Nat.*
524 *Clim. Change*, 10, 892–903, <https://doi.org/10.1038/s41558-020-0905-y>, 2020.

525 Arrigo, K. R. and van Dijken, G. L.: Continued increases in Arctic Ocean primary
526 production, *Synth. Arct. Res. SOAR*, 136, 60–70,
527 <https://doi.org/10.1016/j.pocean.2015.05.002>, 2015.

528 Artuso, F, Canuti, E, Cataldi, D, Costa Goela, P, Grung, M, Ras, J, and Röttgers, R:
529 HPLC/DAD intercomparison on phytoplankton pigments (HIP-1, HIP-2, HIP-3 and HIP-4),
530 Publications Office of the European Union, 2016.

531 Bidigare, R. R., Van Heukelem, L., and Trees, C. C.: HPLC phytoplankton pigments:
532 sampling, laboratory methods, and quality assurance procedures, *Ocean Opt. Protoc.*
533 *Satell. Ocean Color Sens. Valid. Revis.*, 3, 258–268, 2002.

534 Blais, M., Ardyna, M., Gosselin, M., Dumont, D., Bélanger, S., Tremblay, J.-É., Gratton, Y.,
535 Marchese, C., and Poulin, M.: Contrasting interannual changes in phytoplankton
536 productivity and community structure in the coastal Canadian Arctic Ocean, *Limnol.*
537 *Oceanogr.*, 62, 2480–2497, <https://doi.org/10.1002/lno.10581>, 2017.

538 Bonilla, S., Rautio, M., and Vincent, W. F.: Phytoplankton and phytobenthos pigment
539 strategies: implications for algal survival in the changing Arctic, *Polar Biol.*, 32, 1293–
540 1303, <https://doi.org/10.1007/s00300-009-0626-1>, 2009.

541 Bracher, A., Vountas, M., Dinter, T., Burrows, J. P., Röttgers, R., and Peeken, I.:
542 Quantitative observation of cyanobacteria and diatoms from space using PhytoDOAS
543 on SCIAMACHY data, *Biogeosciences*, 6, 751–764, [https://doi.org/10.5194/bg-6-751-](https://doi.org/10.5194/bg-6-751-2009)
544 2009, 2009.

545 Bracher, A., Bouman, H. A., Brewin, R. J. W., Bricaud, A., Brotas, V., Ciotti, A. M.,
546 Clementson, L., Devred, E., Di Cicco, A., Dutkiewicz, S., Hardman-Mountford, N. J.,
547 Hickman, A. E., Hieronimi, M., Hirata, T., Losa, S. N., Mouw, C. B., Organelli, E., Raitsos,
548 D. E., Uitz, J., Vogt, M., and Wolanin, A.: Obtaining Phytoplankton Diversity from Ocean
549 Color: A Scientific Roadmap for Future Development, *Front. Mar. Sci.*, Volume 4-2017,
550 <https://doi.org/10.3389/fmars.2017.00055>, 2017.

551 Bracher, A., Soppa, M. A., Gege, P., Losa, S. N., Silva, B., Steinmetz, F., and Droscher, I.:
 552 Extension of Atmospheric Correction Polymer to Hyperspectral Sensors: Application to
 553 HICO and First Results for DESIS Data,
 554 <https://doi.org/10.1109/IGARSS47720.2021.9553568>, 2021.

555 Brunet, C., Johnsen, G., Lavaud, J., and Roy, S.: Pigments and photoacclimation
 556 processes, in: *Phytoplankton Pigments: Characterization, Chemotaxonomy and*
 557 *Applications in Oceanography*, 2011.

558 Bruyant, F., Amiraux, R., Amyot, M.-P., Archambault, P., Artigue, L., Barbedo de Freitas,
 559 L., Bécu, G., Bélanger, S., Bourgain, P., Bricaud, A., Brouard, E., Brunet, C., Burgers, T.,
 560 Caleb, D., Chalut, K., Claustre, H., Cornet-Barthaux, V., Coupel, P., Cusa, M., Cusset, F.,
 561 Dadaglio, L., Davelaar, M., Deslongchamps, G., Dimier, C., Dinasquet, J., Dumont, D.,
 562 Else, B., Eulaers, I., Ferland, J., Filteau, G., Forget, M.-H., Fort, J., Fortier, L., Galí, M.,
 563 Gallinari, M., Garbus, S.-E., Garcia, N., Gérikas Ribeiro, C., Gombault, C., Gourvil, P.,
 564 Goyens, C., Grant, C., Grondin, P.-L., Guillot, P., Hillion, S., Husserr, R., Joux, F., Joy-
 565 Warren, H., Joyal, G., Kieber, D., Lafond, A., Lagunas, J., Lajeunesse, P., Lalande, C.,
 566 Larivière, J., Le Gall, F., Leblanc, K., Leblanc, M., Legras, J., Lévesque, K., Lewis, K.-M.,
 567 Leymarie, E., Leynaert, A., Linkowski, T., Lizotte, M., Lopes dos Santos, A., Marec, C.,
 568 Marie, D., Massé, G., Massicotte, P., Matsuoka, A., Miller, L. A., Mirshak, S., Morata, N.,
 569 Moriceau, B., Morin, P.-I., Morisset, S., Mosbech, A., Mucci, A., Nadaï, G., Nozais, C.,
 570 Obernosterer, I., Paire, T., Panagiotopoulos, C., Parenteau, M., Pelletier, N., Picheral, M.,
 571 Quéguiner, B., Raimbault, P., Ras, J., Rehm, E., Ribot Lacosta, L., Rontani, J.-F., Saint-
 572 Béat, B., Sansoulet, J., Sardet, N., Schmechtig, C., Sciandra, A., Sempéré, R., et al.: The
 573 Green Edge cruise: investigating the marginal ice zone processes during late spring and
 574 early summer to understand the fate of the Arctic phytoplankton bloom, *Earth Syst. Sci.*
 575 *Data*, 14, 4607–4642, <https://doi.org/10.5194/essd-14-4607-2022>, 2022.

576 Campbell, R. G., Sherr, E. B., Ashjian, C. J., Plourde, S., Sherr, B. F., Hill, V., and
 577 Stockwell, D. A.: Mesozooplankton prey preference and grazing impact in the western
 578 Arctic Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56, 1274–1289,
 579 <https://doi.org/10.1016/j.dsr2.2008.10.027>, 2009.

580 Canuti, E.: Phytoplankton pigment in situ measurements uncertainty evaluation: an
 581 HPLC interlaboratory comparison with a European-scale dataset, *Front. Mar. Sci.*, 10,
 582 <https://doi.org/10.3389/fmars.2023.1197311>, 2023.

583 Canuti, E., Grung, J., Röttgers, M., Goela, C., and Artuso, P.: HPLC/DAD Intercomparison
 584 on Phytoplankton Pigments (HIP-1, HIP-2, HIP-3 and HIP-4),
 585 <https://doi.org/10.2788/47099>, 2016.

586 Cetinić, I., Rousseaux, C. S., Carroll, I. T., Chase, A. P., Kramer, S. J., Werdell, P. J., Siegel,
 587 D. A., Dierssen, H. M., Catlett, D., Neeley, A., Soto Ramos, I. M., Wolny, J. L., Sadoff, N.,
 588 Urquhart, E., Westberry, T. K., Stramski, D., Pahlevan, N., Seegers, B. N., Sirk, E., Lange,
 589 P. K., Vandermeulen, R. A., Graff, J. R., Allen, J. G., Gaube, P., McKinna, L. I. W.,
 590 McKibben, S. M., Binding, C. E., Calzado, V. S., and Sayers, M.: Phytoplankton
 591 composition from sPACE: Requirements, opportunities, and challenges, *Remote Sens.*
 592 *Environ.*, 302, 113964, <https://doi.org/10.1016/j.rse.2023.113964>, 2024.

593 Chamberlain, E. J., Balmonte, J. P., Torstensson, A., Fong, A. A., Snoeijs-Leijonmalm, P.,
 594 and Bowman, J. S.: Impacts of sea ice melting procedures on measurements of
 595 microbial community structure, *Elem. Sci. Anthr.*, 10, 00017,
 596 <https://doi.org/10.1525/elementa.2022.00017>, 2022.

597 Claustre, H., Hooker, S. B., Van Heukelem, L., Berthon, J.-F., Barlow, R., Ras, J.,
 598 Sessions, H., Targa, C., Thomas, C. S., van der Linde, D., and Marty, J.-C.: An
 599 intercomparison of HPLC phytoplankton pigment methods using in situ samples:
 600 application to remote sensing and database activities, *Mar. Chem.*, 85, 41–61,
 601 <https://doi.org/10.1016/j.marchem.2003.09.002>, 2004.

602 Coupel, P., Jin, H. Y., Joo, M., Horner, R., Bouvet, H. A., Sicre, M.-A., Gascard, J.-C.,
 603 Chen, J. F., Garçon, V., and Ruiz-Pino, D.: Phytoplankton distribution in unusually low
 604 sea ice cover over the Pacific Arctic, *Biogeosciences*, 9, 4835–4850,
 605 <https://doi.org/10.5194/bg-9-4835-2012>, 2012.

606 Coupel, P., Matsuoka, A., Ruiz-Pino, D., Gosselin, M., Marie, D., Tremblay, J.-É., and
 607 Babin, M.: Pigment signatures of phytoplankton communities in the Beaufort Sea,
 608 *Biogeosciences*, 12, 991–1006, <https://doi.org/10.5194/bg-12-991-2015>, 2015.

609 El Hourany, R., Abboud-Abi Saab, M., Faour, G., Aumont, O., Crépon, M., and Thiria, S.:
 610 Estimation of Secondary Phytoplankton Pigments From Satellite Observations Using
 611 Self-Organizing Maps (SOMs), *J. Geophys. Res. Oceans*, 124, 1357–1378,
 612 <https://doi.org/https://doi.org/10.1029/2018JC014450>, 2019.

613 El Hourany, R., Pierella Karlusich, J., Zinger, L., Loisel, H., Levy, M., and Bowler, C.:
 614 Linking satellites to genes with machine learning to estimate phytoplankton community
 615 structure from space, *Ocean Sci.*, 20, 217–239, [https://doi.org/10.5194/os-20-217-](https://doi.org/10.5194/os-20-217-2024)
 616 2024, 2024.

617 England, M. R., Eisenman, I., Lutsko, N. J., and Wagner, T. J. W.: The Recent Emergence
 618 of Arctic Amplification, *Geophys. Res. Lett.*, 48, e2021GL094086,
 619 <https://doi.org/https://doi.org/10.1029/2021GL094086>, 2021.

620 Falkowski, P. and Kiefer, D. A.: Chlorophyll a fluorescence in phytoplankton: relationship
 621 to photosynthesis and biomass, *J. Plankton Res.*, 7, 715–731,
 622 <https://doi.org/10.1093/plankt/7.5.715>, 1985.

623 Falkowski, P. G.: The role of phytoplankton photosynthesis in global biogeochemical
 624 cycles, *Photosynth. Res.*, 39, 235–258, <https://doi.org/10.1007/BF00014586>, 1994.

625 Flores, H., Veyssi re, G., Castellani, G., Wilkinson, J., Hoppmann, M., Karcher, M.,
 626 Valcic, L., Cornils, A., Geoffroy, M., Nicolaus, M., Niehoff, B., Priou, P., Schmidt, K., and
 627 Stroeve, J.: Sea-ice decline could keep zooplankton deeper for longer, *Nat. Clim.*
 628 *Change*, 13, 1122–1130, <https://doi.org/10.1038/s41558-023-01779-1>, 2023.

629 Fragoso, G. M., Poulton, A. J., Yashayaev, I. M., Head, E. J. H., and Purdie, D. A.: Spring
 630 phytoplankton communities of the Labrador Sea (2005–2014): pigment signatures,

631 photophysiology and elemental ratios, *Biogeosciences*, 14, 1235–1259,
632 <https://doi.org/10.5194/bg-14-1235-2017>, 2017.

633 Freer, J. J., Daase, M., and Tarling, G. A.: Modelling the biogeographic boundary shift of
634 *Calanus finmarchicus* reveals drivers of Arctic Atlantification by subarctic zooplankton,
635 *Glob. Change Biol.*, 28, 429–440, <https://doi.org/https://doi.org/10.1111/gcb.15937>,
636 2022.

637 Gege, P.: The water color simulator WASI: an integrating software tool for analysis and
638 simulation of optical in situ spectra, *Comput. Geosci.*, 30, 523–532,
639 <https://doi.org/10.1016/j.cageo.2004.03.005>, 2004.

640 Gege, P.: WASI-2D: A software tool for regionally optimized analysis of imaging
641 spectrometer data from deep and shallow waters, *Comput. Geosci.*, 62, 208–215,
642 <https://doi.org/10.1016/j.cageo.2013.07.022>, 2014.

643 Gosselin, M., Rysgaard, S., Lavaud, J., Else, B., Galindo, V., Mundy, C. J., Ehn, J., and
644 Babin, M.: Pigment composition and photoprotection of Arctic sea ice algae during
645 spring, *Mar. Ecol. Prog. Ser.*, 585, 49–69, 2017.

646 Gradinger, R.: Sea-ice algae: Major contributors to primary production and algal
647 biomass in the Chukchi and Beaufort Seas during May/June 2002, *Deep Sea Res. Part II*
648 *Top. Stud. Oceanogr.*, 56, 1201–1212, <https://doi.org/10.1016/j.dsr2.2008.10.016>, 2009.

649 Hayward, A., Pinkerton, M. H., and Gutierrez-Rodriguez, A.: phytoclass: A pigment-
650 based chemotaxonomic method to determine the biomass of phytoplankton classes,
651 *Limnol. Oceanogr. Methods*, 21, 220–241,
652 <https://doi.org/https://doi.org/10.1002/lom3.10541>, 2023.

653 Hayward, A., Pinkerton, M. H., Wright, S. W., Gutiérrez-Rodriguez, A., and Law, C. S.:
654 Twenty-six years of phytoplankton pigments reveal a circumpolar Class Divide around
655 the Southern Ocean, *Commun. Earth Environ.*, 5, 92, [https://doi.org/10.1038/s43247-](https://doi.org/10.1038/s43247-024-01261-6)
656 [024-01261-6](https://doi.org/10.1038/s43247-024-01261-6), 2024.

657 Hayward, A., Wright, S. W., Carroll, D., Law, C. S., Wongpan, P., Gutiérrez-Rodriguez, A.,
658 and Pinkerton, M. H.: Antarctic phytoplankton communities restructure under shifting
659 sea-ice regimes, *Nat. Clim. Change*, <https://doi.org/10.1038/s41558-025-02379-x>,
660 2025.

661 Heidemann, A. C., Hayward, A., Assmy, P., Basu, A., Bracher, A., Castellani, G., Ditullio,
662 G., Dragańska-Deja, K., Fujiwara, A., Fragoso, G. M., Høyer, J., Hwang, J., Iversen, M.,
663 von Jackowski, A., Juul-Pedersen, T., Kowalczyk, P., Lee, Y., Matsuoka, A., Merz, A.,
664 Mundy, C. J., Ostermann, E., Pinkerton, M. H., Peeken, I., Stoń-Egiert, J., Thielecke, A. U.,
665 Veyssiere, G., Xi, H., Yang, E. J., and Gonçalves-Araujo, R.: Consolidated Arctic Pigments
666 (2000 to 2024), <https://doi.org/https://doi.org/10.11583/DTU.29445104>, 2026.

667 Hooker, S. B., Heukelem, L., Thomas, C. S., Claustre, H., Ras, J., Barlow, R., Sessions,
668 H., Schlüter, L., Perl, J., Trees, C., Stuart, V., Head, E., Clementson, L., Fishwick, J.,
669 Llewellyn, C., and Aiken, J.: The Second SeaWiFS HPLC Analysis Round-robin

670 Experiment (SeaHARRE-2), National Aeronautics and Space Administration, Goddard
671 Space Flight Center, 124 pp., 2005.

672 Hooker, S. B., Clementson, L., Thomas, C. S., Schlüter, L., Allerup, M., Ras, J., Claire, N.,
673 Cullen, J., Kienast, M., Kozlowski, W., Vernet, M., Chakraborty, S., Lohrenz, S., Tuel, M.,
674 Redalje, D., Cartaxana, P., Mendes, C. R., Brotas, V., Matondkar, S. G. P., Parab, S. G.,
675 Neeley, A., and Egeland, E. S.: The Fifth SeaWiFS HPLC Analysis Round-Robin
676 Experiment (SeaHARRE-5), 2012.

677 Hu, C.: Hyperspectral reflectance spectra of floating matters derived from
678 Hyperspectral Imager for the Coastal Ocean (HICO) observations, *Earth Syst. Sci. Data*,
679 14, 1183–1192, <https://doi.org/10.5194/essd-14-1183-2022>, 2022.

680 Hwang, J.: Preliminarily annotated phytoplankton lipids from Polar Cod Connectivity
681 Cruise 2022, <https://doi.org/10.5281/zenodo.15085544>, 2025.

682 Jeffrey, S. W., Wright, S. W., and Zapata, M.: Recent advances in HPLC pigment analysis
683 of phytoplankton, *Mar. Freshw. Res.*, 50, 879–896, 1999.

684 Kramer, S. J., Siegel, D. A., Maritorena, S., and Catlett, D.: Modeling surface ocean
685 phytoplankton pigments from hyperspectral remote sensing reflectance on global
686 scales, *Remote Sens. Environ.*, 270, 112879,
687 <https://doi.org/https://doi.org/10.1016/j.rse.2021.112879>, 2022.

688 Losa, S. N., Soppa, M. A., Dinter, T., Wolanin, A., Brewin, R. J. W., Bricaud, A., Oelker, J.,
689 Peeken, I., Gentili, B., Rozanov, V., and Bracher, A.: Synergistic Exploitation of Hyper-
690 and Multi-Spectral Precursor Sentinel Measurements to Determine Phytoplankton
691 Functional Types (SynSenPFT), *Front. Mar. Sci.*, 4,
692 <https://doi.org/10.3389/fmars.2017.00203>, 2017.

693 Massicotte, P., Amiraux, R., Amyot, M.-P., Archambault, P., Ardyna, M., Arnaud, L.,
694 Artigue, L., Aubry, C., Ayotte, P., Bécu, G., Bélanger, S., Benner, R., Bittig, H. C., Bricaud,
695 A., Brossier, É., Bruyant, F., Chauvaud, L., Christiansen-Stowe, D., Claustre, H., Cornet-
696 Barthaux, V., Coupel, P., Cox, C., Delaforge, A., Dezutter, T., Dimier, C., Domine, F.,
697 Dufour, F., Dufresne, C., Dumont, D., Ehn, J., Else, B., Ferland, J., Forget, M.-H., Fortier,
698 L., Galí, M., Galindo, V., Gallinari, M., Garcia, N., Gérikas Ribeiro, C., Gourdal, M.,
699 Gourvil, P., Goyens, C., Grondin, P.-L., Guillot, P., Guilmette, C., Houssais, M.-N., Joux,
700 F., Lacour, L., Lacour, T., Lafond, A., Lagunas, J., Lalande, C., Laliberté, J., Lambert-
701 Girard, S., Larivière, J., Lavaud, J., LeBaron, A., Leblanc, K., Le Gall, F., Legras, J., Lemire,
702 M., Lévasseur, M., Leymarie, E., Leynaert, A., Lopes dos Santos, A., Lourenço, A., Mah,
703 D., Marec, C., Marie, D., Martin, N., Marty, C., Marty, S., Massé, G., Matsuoka, A.,
704 Matthes, L., Moriceau, B., Muller, P.-E., Mundy, C.-J., Neukermans, G., Oziel, L.,
705 Panagiotopoulos, C., Pangrazi, J.-J., Picard, G., Picheral, M., Pinczon du Sel, F.,
706 Pogorzelec, N., Probert, I., Quéguiner, B., Raimbault, P., Ras, J., Rehm, E., Reimer, E.,
707 Rontani, J.-F., Rysgaard, S., Saint-Béat, B., Sampei, M., Sansoulet, J., Schmechtig, C.,
708 Schmidt, S., et al.: Green Edge ice camp campaigns: understanding the processes
709 controlling the under-ice Arctic phytoplankton spring bloom, *Earth Syst. Sci. Data*, 12,
710 151–176, <https://doi.org/10.5194/essd-12-151-2020>, 2020.

711 Mattei, F. and Scardi, M.: Collection and analysis of a global marine phytoplankton
 712 primary-production dataset, *Earth Syst. Sci. Data*, 13, 4967–4985,
 713 <https://doi.org/10.5194/essd-13-4967-2021>, 2021.

714 Mendes, C. R., Cartaxana, P., and Brotas, V.: HPLC determination of phytoplankton and
 715 microphytobenthos pigments: comparing resolution and sensitivity of a C18 and a C8
 716 method, *Limnol. Oceanogr. Methods*, 5, 363–370,
 717 <https://doi.org/10.4319/lom.2007.5.363>, 2007.

718 Miller, L. A., Fripiat, F., Else, B. G. T., Bowman, J. S., Brown, K. A., Collins, R. E., Ewert, M.,
 719 Fransson, A., Gosselin, M., Lannuzel, D., Meiners, K. M., Michel, C., Nishioka, J.,
 720 Nomura, D., Papadimitriou, S., Russell, L. M., Sørensen, L. L., Thomas, D. N., Tison, J.-
 721 L., van Leeuwe, M. A., Vancoppenolle, M., Wolff, E. W., and Zhou, J.: Methods for
 722 biogeochemical studies of sea ice: The state of the art, caveats, and recommendations,
 723 *Elem. Sci. Anthr.*, 3, 000038, <https://doi.org/10.12952/journal.elementa.000038>, 2015.

724 Miller, P.: Multi-spectral front maps for automatic detection of ocean colour features
 725 from SeaWiFS, *Int. J. Remote Sens.*, 25, 1437–1442,
 726 <https://doi.org/10.1080/01431160310001592409>, 2004.

727 Negrete-García, G., Luo, J. Y., Petrik, C. M., Manizza, M., and Barton, A. D.: Changes in
 728 Arctic Ocean plankton community structure and trophic dynamics on seasonal to
 729 interannual timescales, *Biogeosciences*, 21, 4951–4973, [https://doi.org/10.5194/bg-21-](https://doi.org/10.5194/bg-21-4951-2024)
 730 4951-2024, 2024.

731 Nieke, J., Mavrocordatos, C., Donlon, C., Berruti, B., Garnier, T., Riti, J.-B., and Delclaud,
 732 Y.: Ocean and Land Color Imager on Sentinel-3, in: *Optical Payloads for Space Missions*,
 733 John Wiley & Sons, Ltd, 223–245, <https://doi.org/10.1002/9781118945179.ch10>, 2015.

734 Nieke, J., Despoisse, L., Gabriele, A., Weber, H., Strese, H., Ghasemi, N., Gascon, F.,
 735 Alonso, K., Boccia, V., Tsonevska, B., Choukroun, P., Ottavianelli, G., and Celesti, M.:
 736 The copernicus hyperspectral imaging mission for the environment (CHIME): an
 737 overview of its mission, system and planning status, in: *Sensors, Systems, and Next-*
 738 *Generation Satellites XXVII*, *Sensors, Systems, and Next-Generation Satellites XXVII*,
 739 21–40, <https://doi.org/10.1117/12.2679977>, 2023.

740 Peloquin, J., Swan, C., Gruber, N., Vogt, M., Claustre, H., Ras, J., Uitz, J., Barlow, R.,
 741 Behrenfeld, M., Bidigare, R., Dierssen, H., Ditullio, G., Fernandez, E., Gallienne, C., Gibb,
 742 S., Goericke, R., Harding, L., Head, E., Holligan, P., Hooker, S., Karl, D., Landry, M.,
 743 Letelier, R., Llewellyn, C. A., Lomas, M., Lucas, M., Mannino, A., Marty, J.-C., Mitchell, B.
 744 G., Muller-Karger, F., Nelson, N., O’Brien, C., Prezelin, B., Repeta, D., Jr. Smith, W. O.,
 745 Smythe-Wright, D., Stumpf, R., Subramaniam, A., Suzuki, K., Trees, C., Vernet, M.,
 746 Wasmund, N., and Wright, S.: The MAREDAT global database of high performance liquid
 747 chromatography marine pigment measurements, *Earth Syst. Sci. Data*, 5, 109–123,
 748 <https://doi.org/10.5194/essd-5-109-2013>, 2013.

749 Pereira, L. and Gonçalves, A. M. M.: *Plankton Communities*, BoD – Books on Demand,
 750 200 pp., 2022.

751 Quinlan, R., Douglas, M. S. V., and Smol, J. P.: Food web changes in arctic ecosystems
 752 related to climate warming, *Glob. Change Biol.*, 11, 1381–1386,
 753 <https://doi.org/10.1111/j.1365-2486.2005.00981.x>, 2005.

754 Roy, S., Llewellyn, C. A., Egeland, E. S., and Johnsen, G.: *Phytoplankton Pigments:*
 755 *Characterization, Chemotaxonomy and Applications in Oceanography*, Cambridge
 756 University Press, 891 pp., 2011.

757 Sadeghi, A., Dinter, T., Vountas, M., Taylor, B. B., Altenburg-Soppa, M., Peeken, I., and
 758 Bracher, A.: Improvement to the PhytoDOAS method for identification of
 759 coccolithophores using hyper-spectral satellite data, *Ocean Sci.*, 8, 1055–1070,
 760 <https://doi.org/10.5194/os-8-1055-2012>, 2012.

761 Serra-Pompei, C., Ward, B. A., Pinti, J., Visser, A. W., Kiørboe, T., and Andersen, K. H.:
 762 Linking Plankton Size Spectra and Community Composition to Carbon Export and Its
 763 Efficiency, *Glob. Biogeochem. Cycles*, 36, e2021GB007275,
 764 <https://doi.org/https://doi.org/10.1029/2021GB007275>, 2022.

765 Simmons, L. J., Sandgren, C. D., and Berges, J. A.: Problems and pitfalls in using HPLC
 766 pigment analysis to distinguish Lake Michigan phytoplankton taxa, *J. Gt. Lakes Res.*, 42,
 767 397–404, <https://doi.org/10.1016/j.jglr.2015.12.006>, 2016.

768 Six, C., Ratin, M., Marie, D., and Corre, E.: Marine *Synechococcus* picocyanobacteria:
 769 Light utilization across latitudes, *Proc. Natl. Acad. Sci.*, 118, e2111300118,
 770 <https://doi.org/10.1073/pnas.2111300118>, 2021.

771 Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons,
 772 <https://doi.org/10.1088/1748-9326/aade56>, 2018.

773 Swan, C. M., Vogt, M., Gruber, N., and Laufkoetter, C.: A global seasonal surface ocean
 774 climatology of phytoplankton types based on CHEMTAX analysis of HPLC pigments,
 775 *Deep Sea Res. Part Oceanogr. Res. Pap.*, 109, 137–156,
 776 <https://doi.org/https://doi.org/10.1016/j.dsr.2015.12.002>, 2016.

777 Vidussi, F., Claustre, H., Manca, B. B., Luchetta, A., and Marty, J.-C.: Phytoplankton
 778 pigment distribution in relation to upper thermocline circulation in the eastern
 779 Mediterranean Sea during winter, *J. Geophys. Res. Oceans*, 106, 19939–19956,
 780 <https://doi.org/10.1029/1999JC000308>, 2001.

781 Vonnahme, T. R., Chitkara, C., Krawczyk, D., Meire, L., Skogseth, R., Vader, A., and Juul-
 782 Pedersen, T.: Abrupt decline of microplankton species richness linked to coastal inflow
 783 in an Arctic fjord, *Limnol. Oceanogr.*, 70, 2688–2702, <https://doi.org/10.1002/lno.70159>,
 784 2025.

785 Werdell, P. J., Franz, B., Poulin, C., Allen, J., Cairns, B., Caplan, S., Cetinić, I., Craig, S.,
 786 Gao, M., Hasekamp, O., Ibrahim, A., Knobelspiesse, K., Mannino, A., Martins, J. V.,
 787 McKinna, L., Meister, G., Patt, F., Proctor, C., Rajapakshe, C., Ramos, I. S., Rietjens, J.,
 788 Sayer, A., and Sirk, E.: Life after launch: a snapshot of the first six months of NASA's
 789 Plankton, Aerosol, Cloud, Ocean Ecosystem (PACE) mission, in: *Sensors, Systems, and*

790 Next-Generation Satellites XXVIII, Sensors, Systems, and Next-Generation Satellites
 791 XXVIII, 70–84, <https://doi.org/10.1117/12.3033830>, 2024.

792 Wright, S. W. and Jeffrey, S. W.: Pigment Markers for Phytoplankton Production, in:
 793 Marine Organic Matter: Biomarkers, Isotopes and DNA, edited by: Volkman, J. K.,
 794 Springer Berlin Heidelberg, Berlin, Heidelberg, 71–104,
 795 https://doi.org/10.1007/698_2_003, 2006.

796 Xi, H., Losa, S. N., Mangin, A., Soppa, M. A., Garnesson, P., Demaria, J., Liu, Y., d’Andon,
 797 O. H. F., and Bracher, A.: Global retrieval of phytoplankton functional types based on
 798 empirical orthogonal functions using CMEMS GlobColour merged products and further
 799 extension to OLCI data, *Remote Sens. Environ.*, 240, 111704,
 800 <https://doi.org/10.1016/j.rse.2020.111704>, 2020.

801 Xi, H., Peeken, I., Gomes, M., Brotas, V., Tilstone, G. H., Brewin, R. J. W., Dall’Olmo, G.,
 802 Tracana, A., Alvarado, L. M. A., Murawski, S., Wiegmann, S., and Bracher, A.:
 803 Phytoplankton pigment concentrations and phytoplankton groups measured on water
 804 samples collected from various expeditions in the Atlantic Ocean from 71°S to 84°N,
 805 <https://doi.org/10.1594/PANGAEA.954738>, 2023.

806 Zapata, M., Rodríguez, F., and Garrido, J. L.: Separation of chlorophylls and carotenoids
 807 from marine phytoplankton: a new HPLC method using a reversed phase C8 column
 808 and pyridine-containing mobile phases, *Mar. Ecol. Prog. Ser.*, 195, 29–45,
 809 <https://doi.org/10.3354/meps195029>, 2000.

810

811

812