

Satellite-based inversion of global methane fluxes: Capabilities and implications of GOSAT-2 measurements

Makoto Saito^{1*}, Yosuke Niwa¹, Yukio Yoshida¹, Hiroshi Suto², Kei Shiomi², Akihide Kamei¹, Fumie Kataoka³, Isamu Morino¹, Hibiki M. Noda¹, Hirofumi Ohyama¹, Tazu Saeki¹, Yu Someya¹, Hisashi Yashiro¹, and Tsuneo Matsunaga¹

¹Earth System Division, National Institute for Environmental Studies, Tsukuba, 305-8506, Ibaraki, Japan

²Japan Aerospace Exploration Agency, Tsukuba, 305-8505, Ibaraki, Japan

³Remote Sensing Technology Center of Japan, Minato-ku, 105-0001, Tokyo, Japan

Correspondence: Makoto Saito (saito.makoto@nies.go.jp)

Abstract. Methane (CH₄) is a key greenhouse gas with a strong climate impact and a relatively short atmospheric lifetime, making accurate monitoring essential for mitigation strategies. Satellite observations provide global coverage and independent constraints on CH₄ emission estimates, and GOSAT-2, launched in 2018 as the successor to GOSAT, was designed to improve retrieval accuracy and enhance flux estimation. This study presents an evaluation of the GOSAT-2 Level 4 (G2L4) CH₄ flux product, supported by analysis of the underlying Level 2 (L2) XCH₄ retrievals, and summarizes key findings on global and regional CH₄ budgets. Using an atmospheric inversion framework, we generated G2L4 posterior CH₄ fluxes and assessed their consistency by comparing them with inversions constrained by alternative observational datasets, including GOSAT L2 retrievals and ground-based and aircraft measurements. GOSAT-2 achieved substantial improvements in observational coverage and data density compared to its predecessor, particularly in tropical and high-latitude regions. Posterior flux estimates derived from G2L4 are broadly consistent with global CH₄ budgets reported in synthesis studies, while prior-to-posterior differences reveal positive corrections in tropical regions and negative adjustments in several mid-latitude industrial areas. A preliminary sector-focused assessment further demonstrates the potential of GOSAT-2 to inform anthropogenic CH₄ emission evaluations in regions where such sources dominate. These findings highlight the value of GOSAT-2 observations for improving regional and global CH₄ emission estimates and underscore priorities for future improvements in retrieval algorithms, observation strategies, and integration with complementary datasets.

1 Introduction

Methane (CH₄) is one of the most important greenhouse gases and a major contributor to anthropogenic climate forcing, the atmospheric amount of which has increased to more than 2.5 times its pre-industrial level. This rapid increase has made CH₄ a major contributor to climate change, ranking just behind carbon dioxide (CO₂) in terms of radiative forcing (Shindell et al., 2021). Its concentration is now increasing faster than at any time since the 1980s (Nisbet et al., 2019). CH₄ is a short-lived climate forcer with an atmospheric lifetime of roughly a decade, which means that taking action now can quickly reduce atmospheric concentrations and lead to similarly rapid reductions in climate forcing. Moreover, CH₄ contributes to the formation of

ground-level ozone, so mitigation would also help decrease ozone-related air pollution. A better understanding of the CH₄ budget at various scales from local to global is essential to constrain atmospheric concentrations and to design effective mitigation strategies that limit global warming to 1.5°C this century.

However, efforts to understand the CH₄ budgets remain challenging because atmospheric variability reflects a complex interplay of anthropogenic sources, natural emissions such as wetlands, and chemical sinks. For instance, the renewed increase in the global CH₄ burden since 2007, accompanied by a significant shift toward more depleted ¹³C in atmospheric CH₄, has yet to be fully explained, with multiple contributing factors still under debate (e.g., Worden et al., 2017; Nisbet et al., 2019; Li et al., 2022). These uncertainties highlight the need for comprehensive observations. However, existing ground-based and aircraft observations are unevenly distributed and provide limited coverage in regions where CH₄ emissions are large or highly variable. Capturing the full spatial and temporal variability of CH₄ therefore requires observational systems that can deliver consistent measurements at the global scale. Consequently, satellite measurements potentially provide a unique opportunity to deliver global coverage and independent constraints on CH₄ emissions.

Instruments aboard Earth-observing satellites retrieve CH₄ concentrations by measuring solar radiation reflected from the Earth's surface and attenuated by gas absorption at specific wavelengths. Instruments are designed to target absorption bands of CO₂ and CH₄ in the shortwave infrared (SWIR) region, which provides sensitivity to variations in column abundance, including contributions from the lower troposphere that are relevant for surface flux estimation. To reduce uncertainties caused by scattering from clouds and aerosols, absorption bands of oxygen (O₂), whose atmospheric abundance is nearly constant, are also observed to correct for variations in optical path length. Satellite-based CH₄ monitoring began with the SCIAMACHY (SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY) instrument aboard ENVISAT, which was launched in 2002 and provided the first global measurements of greenhouse gases from space (Frankenberg et al., 2006). This was followed by the launch of GOSAT (Greenhouse gases Observing SATellite) in 2009, the world's first dedicated greenhouse gas observation satellite, which delivered high-precision column-averaged dry-air mole fraction of CO₂ and CH₄ (XCO₂ and XCH₄) and enabled robust global CH₄ budget assessments (Yokota et al., 2009). Subsequent missions, such as the TROPOMI (TROPOspheric Monitoring Instrument) instrument aboard Sentinel-5P offering wide coverage and frequent observations, have further advanced CH₄ monitoring (Lorente et al., 2021).

These satellite-based CH₄ observations have significantly advanced our understanding of CH₄ sources and sinks. Early studies demonstrated that XCO₂ and XCH₄ data can be retrieved with high precision from SWIR measurements (Yoshida et al., 2013), and validation frameworks using ground truth provided by TCCON (Total Carbon Column Observing Network) were established to ensure data accuracy (Wunch et al., 2011). Building on these capabilities, atmospheric inversion ensembles incorporating satellite observations have contributed to constraining both natural and anthropogenic CH₄ emissions, particularly in regions with sparse ground-based coverage such as developing countries (Deng et al., 2025). Satellite data have also provided essential measurements that support the quantification of regional and sectoral contributions to the global CH₄ budget (Zhang et al., 2021) and evaluation of national emissions from anthropogenic activities (Chen et al., 2022). Beyond large-scale assessments, satellite observations have revealed critical events and natural source dynamics, including extreme CH₄ leakage from a natural gas well blowout (Pandey et al., 2019) and the spatial and temporal distribution of global wetland emissions

(Chen et al., 2025). These advances underscore the pivotal role of satellite measurements in improving CH₄ emission inventories and guiding mitigation strategies.

Motivated by the successful achievements of space-based XCH₄ measurements and the need for continuous long-term monitoring of greenhouse gases, GOSAT-2 was launched in 2018 as the successor to GOSAT (Imasu et al., 2023). The mission was designed to improve the accuracy of greenhouse gas flux estimates and to enable more robust assessments of anthropogenic emissions, including those from large urban areas. To meet these objectives, GOSAT-2 introduced several enhancements: improved signal-to-noise ratio (SNR) for higher retrieval precision, an expansion of the spectral coverage to include absorption by carbon monoxide (CO) to better characterize fossil fuel emissions, and expanded target observation strategies to increase the number of quality-screened measurements for national-scale emission estimates. Following its initial public release, the GOSAT-2 XCH₄ data product has been continuously improved through updates to spectral radiance data and retrieval algorithms. These continuous enhancements and the growing long-term record enable the use of GOSAT-2 XCH₄ data for estimating CH₄ sources and sinks at both regional and global scales.

This study focuses on two key GOSAT-2 data products. The Level 2 SWIR XCH₄ product (version V02.10) provides an updated long-term record of satellite-based CH₄ columns with improved retrieval performance and substantially expanded spatial coverage compared to GOSAT. In addition, we present the first public release of the GOSAT-2 Level 4 CH₄ flux product, which delivers monthly CH₄ surface fluxes at 1° resolution derived exclusively from GOSAT-2 observations. Together, these datasets offer a new basis for evaluating global and regional CH₄ budgets using the enhanced capabilities of the GOSAT-2 mission.

Building on these datasets, this paper provides an overview of the GOSAT-2 mission and evaluates its XCH₄ data product and CH₄ flux estimates. Section 2 describes the sensor specifications of GOSAT-2 as well as the data and modeling system used in this study, while Section 3 presents key findings on the global distribution of XCH₄ and methane budgets. Finally, we discuss the implications of these results for regional source and sink estimates and for assessments of anthropogenic emissions.

2 Methods

2.1 GOSAT-2 satellite system and instruments

2.1.1 Satellite bus design

The GOSAT-2 satellite bus system follows the basic design of the first mission, GOSAT, with several modifications introduced based on operational experience. Key enhancements include an enlarged solar array and increased power generation capacity, enabling stable full-operation even under partial system failure. The sensor was repositioned to reduce stray light from the solar array, and the overall bus configuration was adjusted accordingly to maintain thermal stability and instrument performance. These design updates contribute to the broader observational coverage provided by GOSAT-2 compared with its predecessor.

2.1.2 Orbital configuration

A major change from the GOSAT mission is the adoption of a six-day repeat cycle for GOSAT-2, compared with the three-day cycle of GOSAT. This configuration reduces the spacing between adjacent ground tracks at the equator, providing denser spatial sampling along the orbit. The improved sampling supports more uniform global coverage for SWIR observations of CH₄, especially in low-latitude regions where cloudiness often limits data availability. The repeat cycle is determined by the satellite's altitude and inclination, and for GOSAT-2 the orbit was set to 613 km, corresponding to 89 revolutions in six days. The local solar time at the descending node is maintained at 13:00 ± 15 minutes. In addition, the orbit was designed to pass directly over the Lamont TCCON site to enable regular cross-validation under frequent clear-sky conditions.

2.1.3 Primary instruments

GOSAT-2 is equipped with two primary instruments:

- TANSO-FTS-2 (Thermal And Near infrared Sensor for carbon Observation-Fourier Transform Spectrometer-2)
- TANSO-CAI-2 (Cloud and Aerosol Imager-2)

Similar to the TANSO-FTS instrument onboard GOSAT, TANSO-FTS-2 performs spectroscopic measurements using a Fourier Transform Spectrometer (FTS). It is equipped with five detector bands covering wavelengths from SWIR to thermal infrared (TIR), enabling the observation of atmospheric concentrations of CO₂, CH₄, water vapor (H₂O), O₂, ozone (O₃), and CO. A detailed summary of the wavelength coverage for each band is provided in Table 1.

TANSO-FTS-2 employs a double-pendulum interferometer. In this configuration, the intersection of the flexure blades serves as the rotation axis, and the scanner arm moves in either the clockwise (forward) or counterclockwise (backward) direction. Each forward or backward movement constitutes one scan, during which a single observation dataset is acquired. Because the instrument characteristics differ between forward and backward scans, the data processing algorithm identifies the scan direction and handles the data accordingly. In addition, the forward/backward viewing geometry is used in coordination with TANSO-CAI-2 to avoid sun-glint over the ocean and to improve cloud discrimination, contributing to increased numbers of usable ocean observations.

2.1.4 Observation geometry and Intelligent Pointing

The detector employs a circular point sensor, producing a footprint with a diameter of 9.7 km at nadir. This field of view can be steered using mirrors within ±40° in the along-track (AT) direction and ±35° in the cross-track (CT) direction using a two-axis pointing mirror.

To increase the number of successful observations, TANSO-FTS-2 incorporates an "Intelligent Pointing" function that uses an onboard field-of-view camera to capture visible images of the area surrounding the planned observation points. When clouds are detected at the nominal footprint, the system automatically searches for nearby cloud-free locations within the camera's

Table 1. Summary of TANSO-FTS-2 instrument.

	Band 1	Band 2	Band 3	Band 4	Band 5
Wavelength range (cm^{-1}) [μm]	12 950–13 250 [0.75–0.77]	5 900–6 400 [1.56–1.69]	4 200–5 200 [1.92–2.33]	1 188–1 800 [5.5–8.4]	700–1 188 [8.4–14.3]
Out-of-band characteristics (cm^{-1})	< 12 750 > 13 450	< 5 100 > 6 800	< 4 100 > 5 500	< 1 000 > 3 800	< 600 > 1 300
Polarization	○	○	○	–	–
Spectral resolution (cm^{-1})	0.2 (Double-sides scan) (MOPD ± 2.5 cm)				
FWHM of instrument function (cm^{-1})	< 0.4	< 0.27			

imaging area and adjusts the pointing direction toward these clear-sky targets. If a clear-sky target cannot be identified, the instrument observes the center of the scene.

120 Observations are conducted at 1,246 discrete points along each orbit, with 4.024 seconds allocated for interferogram acquisition and approximately 0.65 seconds for the turnaround phase during which the pointing mirror is reoriented toward the next observation target.

2.1.5 Performance improvement

TANSO-FTS-2 performs routine on-orbit calibration operations, including solar irradiance, blackbody, nighttime, deep-space,
125 instrument-function, lunar, and electrical calibration. Vicarious calibration using actual observation data is also applied (Kuze et al., 2010).

To fully utilize the available dynamic range and suppress quantization noise, TANSO-FTS-2 employs an adaptive gain-control system that adjusts detector gain according to signal intensity and observational conditions. This mechanism ensures appropriate sensitivity across diverse surface types, including bright desert regions and sun-glint areas. Detailed description of
130 the gain-control architecture and its operational configurations are provided in Suto et al. (2021, 2022).

2.1.6 Data downlink and processing

Data acquired in orbit are downlinked to a ground station located in the Svalbard Islands. The downlinked data are then transmitted to the Japan Aerospace Exploration Agency (JAXA) Tsukuba Space Center, where Level 0 and Level 1 processing are performed.

- 135 – Level 0 processing: involves unpacking the data packets received from the satellite and arranging them in sequence.
- Level 1A: conversion of Level 0 data into interferograms.

- Level 1B: inverse Fourier transformation to generate radiance spectra, followed by radiometric corrections and related adjustments.

The Level 1B product is then transmitted to the National Institute for Environmental Studies (NIES), where higher-level
140 processing, including Level 2 and beyond, is performed.

2.1.7 Level 1B product release and updates

The Level 1B product was first made publicly available in 2019 and has since undergone multiple updates to its calibration and processing algorithms. For details on the instrument characteristics and operational aspects of GOSAT-2, see Suto et al. (2021, 2022).

145 2.2 SWIR Column-averaged dry air mole fraction product

2.2.1 Input data and lookup tables

The XCH_4 retrieval in this study is based on the GOSAT-2 Level 1B radiance product, supplemented by auxiliary information required for the full-physics forward model. These inputs include instrument characterization data, the optical depth of the reference spectrum, solar irradiance, and lookup tables (LUTs) describing gas absorption and aerosol optical properties.
150 Gas absorption cross-sections, aerosol parameters, and solar Fraunhofer lines follow the datasets described in the G2L2 Algorithm Theoretical Basis Document (G2L2ATBD; Yoshida and Oshio, 2025), with aerosol information based on the Spectral Radiation-Transport Model for Aerosol Species (SPRINTARS; Takemura et al., 2003) and solar spectral data following Toon (2015).

Although the G2L2 retrieval algorithm is capable of accounting for cirrus cloud scattering, the V02.XX product applies a
155 pre-screening procedure that retains only cloud-free observations. As a result, clouds are not treated explicitly in the radiative transfer calculations for the version used in this study.

2.2.2 Preprocessing of radiance spectra

As part of the preprocessing, radiance correction, wavenumber-interval correction, and polarization synthesis are applied to the Level 1B spectra. Radiance correction accounts for sensitivity variations derived from on-orbit solar, lunar, and vicarious
160 calibration data. The wavenumber axis is adjusted to correct for small variations in the spectral sampling interval contained in the Level 1B product. Polarization synthesis is used to derive the total incident radiance on TANSO-FTS-2 from the two polarized components of the observed spectra.

2.2.3 Retrieval approach: Full physics method

The GOSAT-2 TANSO-FTS-2 SWIR Level 2 column-averaged dry-air mole fraction product (hereinafter referred to as G2L2
165 product) is retrieved using a full-physics approach in which multiple spectral bands are fitted simultaneously while accounting

for scattering by aerosols and surface reflection. Observations are first screened prior to the retrieval to remove soundings with instrumental quality issues or indications of cloud and aerosol contamination. After the retrieval, an additional quality screening based on convergence diagnostics and retrieval quality criteria is applied, and only successful retrievals are included in the final product.

170 The retrieval is formulated as a maximum a posteriori (MAP) optimization (Rodgers, 2000), in which the state vector includes concentrations of CO₂, CH₄, H₂O, and CO in 15 vertical layers, aerosol profile parameters, surface pressure, temperature-profile adjustment, surface albedo, and chlorophyll fluorescence. Instrument-related parameters include radiometric offsets and instrument line-shape scaling factors.

The forward model combines a radiative-transfer module and an instrument-response module. The radiative-transfer module
175 computes monochromatic radiances using a fast non-polarized radiative-transfer scheme; although the algorithm is capable of treating polarization, polarization is not considered in the operational retrieval. The instrument-response module applies the TANSO-FTS-2 spectral response to generate simulated spectra and Jacobians consistent with the Level 1B product.

Further methodological and numerical details of the LUT construction, preprocessing steps, and the full-physics retrieval formulation are documented comprehensively in the G2L2ATBD.

180 2.2.4 G2L2 product release and updates

The G2L2 product was first released as a preliminary version (V01.01) for research announcement users in November 2019, followed by a general release (V01.04) in November 2020. A major update, including improvements to both the retrieval algorithm and input data, was implemented in July 2022 (V02.00). Subsequently, a minor update (V02.10) addressing a bug in the *a priori* aerosol optical thickness setting and revising post-screening criteria was released in February 2025. This version
185 uses Level 1B product V220.220 as input data. The update from V02.00 to V02.10 significantly improved spatial distribution, particularly over desert regions; therefore, this study employs V02.10 for global surface CH₄ flux estimation. Since then, the product has been updated to V02.21 due to changes in Level 1B inputs, but no further modifications to the processing algorithm have been made since V02.10.

2.3 Global surface CH₄ flux estimates

190 2.3.1 Overview of GOSAT-2 Level 4 CH₄ flux estimation

The GOSAT project has provided global surface CO₂ and CH₄ flux estimates as Level 4 (G1L4) products since 2012. GOSAT Level 4 products were generated on a sub-continental scale using the NIES atmospheric tracer transport model (NIES-TM) together with a fixed-lag Kalman smoother (Maksyutov et al., 2013). While this framework improved flux accuracy, it relied heavily on in situ observations and could not fully isolate the contribution of satellite data. The GOSAT-2 Level 4 CH₄ flux
195 product (G2L4) was developed to address these limitations. G2L4 differs from G1L4 in two fundamental aspects: (1) it uses the Non-hydrostatic Icosahedral Atmospheric Model (NICAM)-based inversion system described in Niwa et al. (2025), which provides higher spatial resolution and fully mass-conservative transport, and (2) it assimilates only the G2L2 product, allowing

a satellite-only assessment of global and regional CH₄ budgets. Except for these differences in observational constraints and prior specifications, the inversion framework used in this study follows Niwa et al. (2025). The key components of the inversion framework are summarized below, and further details are provided in the following subsections.

In this study, global surface CH₄ fluxes are estimated using an atmospheric inversion framework based on a Bayesian approach, in which prior information on surface fluxes is optimized by assimilating atmospheric CH₄ observations. The inversion is formulated within a four-dimensional variational (4D-Var) framework, in which the optimal flux vector is obtained by minimizing a cost function that accounts for both prior flux constraints and observation-model mismatch. The cost function J is defined as:

$$J = \frac{1}{2} \delta \mathbf{x}^T \mathbf{B}^{-1} \delta \mathbf{x} + \frac{1}{2} (\mathbf{M} \delta \mathbf{x} - \mathbf{d})^T \mathbf{R}^{-1} (\mathbf{M} \delta \mathbf{x} - \mathbf{d}), \quad (1)$$

where $\delta \mathbf{x} = \mathbf{x} - \mathbf{x}_{\text{pri}}$, with $\mathbf{x}$ and $\mathbf{x}_{\text{pri}}$ denoting vectors of modeled and *a priori* source/sink strengths, respectively. $\mathbf{M}$ is the linear forward transport operator that maps flux perturbations to concentration changes at observation locations. $\mathbf{d} = \mathbf{y} - \mathbf{M} \mathbf{x}_{\text{pri}}$ where with $\mathbf{y}$ is the vector of observed concentrations. $\mathbf{B}$ and $\mathbf{R}$ are the error covariance matrices for the *a priori* flux estimates and the observation-model misfit, respectively. The superscript T denotes the transpose.

The control variables consist of adjustments to prior CH₄ fluxes, including anthropogenic emissions, biomass burning, terrestrial biosphere fluxes, and other natural sources. The global surface CH₄ fluxes are estimated on a 1° grid with monthly temporal resolution.

2.3.2 Model system: NISMON-CH₄

The inversion system used to generate the G2L4 product is based on the NICAM-based Inverse Simulation for Monitoring CH₄ (NISMON-CH₄; Niwa et al., 2025). NISMON-CH₄ integrates the NICAM-TM transport model with a four-dimensional variational (4D-Var) inversion system, providing mass-conservative global transport and high-resolution flux estimates on a 1° grid.

Because the present study focuses on evaluating the impact of GOSAT-2 observations, only the modifications specific to G2L4—such as the use of G2L2 as the sole observational constraint and updates to prior fluxes—are summarized in the following sections.

2.3.3 Prior fluxes

The prior CH₄ fluxes used in the G2L4 inversion largely follow the framework of Niwa et al. (2025), with updates specific to this study. Anthropogenic emissions are taken from the Emissions Database for Global Atmospheric Research (EDGAR version 6.0; Crippa et al., 2021; Ferrario et al., 2021) and prescribed as annual totals for five source sectors: coal mining, oil and gas, landfill and waste, biofuel use, and enteric fermentation and manure management.

Monthly biogenic flux components—rice paddies, wetlands, and soil oxidation—are derived from a prognostic biosphere model, the Vegetation Integrative Simulator for Trace gases (VISIT; Ito and Inatomi, 2012), using the Cao et al. (1996) scheme for

wetland and rice-paddy CH₄ production. Biomass-burning emissions are taken from the Global Biomass Burning Emissions
230 Inventory (GBEI version "2022a"; Shiraishi et al., 2021; Saito et al., 2022) and aggregated to a 1° grid.

Natural fluxes include oceanic, termite, and geological emissions, following Weber et al. (2019), Ito (2023), and Etiope et al. (2019). Geological emissions are scaled to a global total of 23 Tg CH₄ yr⁻¹ after Canadell et al. (2021), as in Niwa et al. (2025). The global surface CH₄ fluxes are estimated at a spatial resolution of 1.0° and a monthly time step.

As in Niwa et al. (2025), EDGAR v6.0 anthropogenic emissions and VISIT-derived biospheric fluxes were available only
235 through their final release years. For subsequent years, the fluxes from the final available year were used. Consequently, prior fluxes other than biomass-burning emissions do not contain interannual variability during the study period, and interannual variations in posterior flux estimates are primarily driven by observational constraints.

The control vector consists of scaling factors and flux adjustments applied to the prior CH₄ flux components. Different optimization approaches are applied depending on the source category. Anthropogenic emissions, biomass-burning emissions,
240 and other natural emissions are optimized using multiplicative scaling factors that preserve their prescribed spatial patterns. In contrast, rice-paddy, wetland, and soil-oxidation fluxes are optimized using additive flux adjustments, allowing both the magnitude and spatial distribution of these biosphere-related fluxes to vary. This treatment reflects the larger uncertainty in the spatial distribution of biospheric CH₄ fluxes and provides greater flexibility for the inversion to adjust these sources in response to atmospheric observations.

245 **2.3.4 Observational and error settings**

In the G2L4 inversion, the observation-model mismatch error for each GOSAT-2 XCH₄ retrieval is set to 20 ppb, corresponding to the square root of the diagonal elements of the observation-error covariance matrix, following the configuration used in Niwa et al. (2025). This value is comparable to the uncertainty estimated from validation against TCCON observations. For G2L2, comparisons with TCCON typically show standard deviations of approximately 10-20 ppb depending on site, surface
250 type, and collocation criteria (Yoshida et al., 2023; NIES GOSAT-2 Project, 2025). The use of a uniform value provides a consistent weighting of observations across regions and facilitates direct assessment of the influence of satellite observations on the inversion results. Only G2L2 retrievals that pass all internal pre- and post-screening criteria are assimilated, and an additional selection is applied based on the quality criteria provided in the G2L2 product. Thus, all analyses use XCH₄ data that satisfy both the retrieval-level filters and the product-level quality flags.

255 Prior flux uncertainties follow the sector-dependent settings of Niwa et al. (2025): anthropogenic emissions are assigned 50% uncertainty, and biomass burning and natural emissions 100%. For rice paddies, wetlands, and soil oxidation, prior covariances are derived from a 120-year ensemble using the VISIT terrestrial biosphere model, with spatial localization applied to suppress unrealistic long-range correlations.

2.3.5 Optimization framework

260 The optimization of surface CH₄ fluxes follows the 4D-Var scheme implemented in NISMON-CH₄ (Niwa et al., 2025). The inversion is solved using the Preconditioned Optimizing Utility for Large-dimensional analyses (POpULar) quasi-Newton

method (Fujii, 2005; Niwa et al., 2017), which iteratively updates flux adjustments through forward and adjoint model integrations.

2.3.6 Model implementation details

265 The G2L4 system adopts the standard implementation of NISMON-CH₄ described by Niwa et al. (2025). The inversion is conducted on the native NICAM icosahedral grid, while posterior fluxes are provided on a regular 1° latitude-longitude grid using the grid-conversion scheme described by Niwa et al. (2022).

To ensure consistency between simulated and retrieved atmospheric CH₄ concentrations, the averaging kernels and a priori profiles distributed with the G2L2 product are applied in the observation operator. In addition, the penalty formulation following
270 Niwa et al. (2022) is incorporated to suppress physically unrealistic negative flux estimates during the optimization.

Further methodological and numerical details, including covariance construction, adjoint implementation, grid conversion procedures, and penalty function settings, are documented in Niwa et al. (2025) and the G2L4 Algorithm Theoretical Basis Document (G2L4ATBD; Saito and Niwa, 2025).

2.4 Additional data

275 For comparison with the G2L2 product, we used the GOSAT TANSO-FTS SWIR Level 2 CH₄ column-amount product (version V03.05; hereinafter G1L2). The V03 series provides XCO₂ and XCH₄ retrieved from GOSAT Level 1B spectra using the full-physics method (Someya et al., 2023). Version V03.05 corresponds to the bias-corrected product validated against TCCON measurements, and only data meeting the product quality criteria were used in this study.

3 Results

280 This section presents the results of the GOSAT-2 Level 4 CH₄ flux estimates (G2L4), which constitute the primary objective of this study. To provide the necessary context for interpreting the inversion results, we first summarize key characteristics of the G2L2 XCH₄ product and briefly compare it with the G1L2 product. The flux estimates are evaluated for the period from May 2019 to October 2022, using G2L2 version V02.10 covering March 2019 to December 2022.

3.1 Observation coverage and spatial distribution of G2L2 product

285 Following the launch of GOSAT-2 in October 2018 and completion of its initial calibration phase, the G2L2 product has been available since March 2019. Several short interruptions in TANSO-FTS-2 operations occurred during the period, with the full list provided by the EORC-JAXA archive.

Monthly variations in the number of quality-screened XCH₄ observations are shown in Fig. 1. To ensure comparability between TANSO-FTS and TANSO-FTS-2, repeated soundings over the same location within short time intervals were counted
290 only once per day. Such repeated soundings mainly arise from targeted observation operations, in which the same location may be observed multiple times to increase the likelihood of obtaining cloud-free retrievals. After this adjustment, TANSO-FTS-2




Figure 1. Monthly count of XCH_4 observations meeting the quality criteria for (a) TANSO-FTS and (b) TANSO-FTS-2 in March 2019 and December 2022. Blue and green bars represent observations over ocean and land, respectively.

still yielded roughly 2-3 times more valid observations than TANSO-FTS over both land and ocean, reflecting improvements in sensor performance, pointing strategy, and observational coverage.

Figure 2 shows the spatial distribution of G2L2 observations. TANSO-FTS-2 achieved substantially enhanced global coverage, particularly over the oceans, where nearly all 2.5° grid cells contain valid retrievals. Over land, the Intelligent Pointing system increased the sampling of cloud-free scenes relative to TANSO-FTS, although persistent cloudiness in the deep tropics remains a limiting factor. These improvements reduce spatial sampling gaps and provide a more uniform global dataset, forming a stronger observational basis for the CH_4 flux estimation presented in later sections.

The retrieval counts shown here represent only successful observations included in the GOSAT and GOSAT-2 Level 2 products and therefore reflect the combined effects of orbital sampling, observing strategy, cloud screening, and retrieval performance.

3.2 Validation of G2L2 product

Uncertainties in GOSAT-2 XCH_4 retrievals arise from surface reflectance, aerosol and cirrus properties, instrument characteristics, and limitations in the forward-model representation. Validation of the G2L2 retrievals has been reported previously. Yoshida et al. (2023) evaluated version V02.00 through comparisons with TCCON and the G1L2 product, while the NIES GOSAT-2 Project (2025) provides validation results for the version used in this study, V02.10.

According to the NIES GOSAT-2 Project (2025), validation of the G2L2 V02.10 product against TCCON shows small negative biases over both land and ocean, with larger biases over ocean sites. Typical mean differences fall within a few ppb for land and around -5 to -8 ppb for ocean, with standard deviations of approximately 10-15 ppb depending on the site and collocation criteria. These results indicate that G2L2 V02.10 generally underestimates XCH_4 relative to TCCON, particularly




Figure 2. Spatial distribution of XCH₄ observations meeting the quality criteria, aggregated on a 2.5° grid, and their latitudinal totals for (a, b) TANSO-FTS and (c, d) TANSO-FTS-2 in March 2019 and December 2022. Blue and green lines in panels (b) and (d) indicate observations over ocean and land, respectively.

over ocean scenes. No bias correction is applied in this study; therefore, the flux estimates presented below reflect the original characteristics and remaining retrieval biases of the G2L2 V02.10 product.

3.3 Comparison between G1L2 and G2L2 products

The comparison of G1L2 and G2L2 products is subject to inherent limitations. First, the observation locations of TANSO-FTS and TANSO-FTS-2 do not perfectly coincide, making a direct one-to-one comparisons challenging. Second, differences in their retrieval algorithms can produce systematic variations in retrieved XCH₄ even when the measurements are taken at nearly the same time and location.

Figure 3 presents two-dimensional histograms comparing XCH₄ retrieved from TANSO-FTS (x-axis) and TANSO-FTS-2 (y-axis). Matchups were selected when the fields of view were within 20 km and the observation times were within 30 minutes.




Figure 3. Two-dimensional histograms of XCH_4 concentrations (ppb) retrieved from TANSO-FTS and TANSO-FTS-2 for (a) land and (b) ocean, based on matchup data only.

320 Repeated soundings at the same location within consecutive time intervals were averaged, and only the closest matchup pair was retained.

Over land, the mean difference ($\text{G2L2} - \text{G1L2}$) is -2.66 ppb with a standard deviation of 11.53 ppb ($N = 12,799$), whereas over ocean the mean difference is -6.84 ppb with a standard deviation of 10.74 ppb ($N = 3,614$). Correlation coefficients exceed 0.93 for both surface types, indicating that the two products show high consistency in XCH_4 variability. Combined with the
 325 TCCON validation results in Section 3.2, these comparisons confirm that G2L2 XCH_4 is generally slightly lower than G1L2 and TCCON, with the negative bias being more pronounced over the ocean.

Figure 4 shows the spatial distributions of G1L2 and G2L2 XCH_4 on a 2.5° grid, averaged over four consecutive three-month periods from December 2021 to November 2022. G2L2 provides significantly improved coverage, particularly over oceans, high latitudes, and tropical regions where cloud contamination often limits the number of valid GOSAT retrievals. A notable
 330 exception is the central-eastern region of South America, where the spatial coverage of G2L2 is degraded. Although the cause has not yet been fully identified, reduced radiance levels in Band 1 have been observed in this region, and a possible influence of the South Atlantic Anomaly has been suggested.

The seasonal distributions are broadly consistent between G1L2 and G2L2, showing higher XCH_4 values during boreal fall and lower values from spring to early summer, which is likely influenced by the seasonality of Northern Hemisphere wetland
 335 emissions (East et al., 2024). Regions of enhanced XCH_4 include South and East Asia, the Arabian Peninsula, central Africa, and parts of the Americas, and these patterns are largely reproduced across both products.




Figure 4. Three-month averaged XCH_4 distributions for G1L2 (left) and G2L2 (center) from December 2021 to November 2022, and their differences (G2L2 - G1L2) (right). Color scales represent XCH_4 concentrations (ppb) for G1L2 and G2L2, and differences in ppb for the bottom panels. The rightmost panels additionally show latitudinal mean differences (ppb), with blue and green lines indicating ocean and land regions, respectively.

Mean spatial differences (G2L2 - G1L2) averaged over three-month periods range from -2.6 to -5.4 ppb over land and from -5.2 to -10.8 ppb over ocean, with standard deviations of 12-14 ppb across both surface types. A systematic latitudinal pattern is also evident: negative differences are larger in low-latitude regions and diminish toward higher latitudes, implying a reduced north-south contrast within each hemisphere in G2L2 relative to G1L2. To investigate the influence of bias correction, the same analysis was repeated using G1L2 without bias correction (V03.00). In this case, the latitudinal trend largely disappears, suggesting that the G1L2 bias-correction scheme is a major contributor. Therefore, part of the spatial differences between G1L2 and G2L2 may reflect differences in the bias-correction strategy rather than differences in the underlying satellite observations themselves. Accordingly, caution is required when interpreting the spatial differences between the two products. Because assessing this correction is beyond the scope of this study, no further analysis is conducted, but users should be aware that the presence or absence of bias correction can introduce substantial spatial differences in XCH_4 .

3.4 Regional and global CH₄ flux estimates by G2L4 product

The global CH₄ budget synthesized by the Global Carbon Project (GCP; Saunois et al., 2025) provides the most authoritative benchmark for evaluating top-down CH₄ flux estimates. Against this benchmark, our satellite-only inversion using G2L2 retrievals reproduces the global budget with remarkable accuracy. The posterior global totals are 582.9 and 589.5 Tg CH₄ yr⁻¹ for 2020 and 2021, respectively—well within the uncertainty range of the GCP top-down estimate (572 Tg CH₄ yr⁻¹ for 2020). This high level of agreement suggests that GOSAT-2 SWIR observations are capable of reproducing the global-scale CH₄ budget, despite inherent retrieval uncertainties in satellite-based measurements (e.g., O'Dell et al., 2018). However, agreement at the global scale does not necessarily imply uniformly strong observational constraints at regional scales, where the influence of prior assumptions and observational coverage may vary substantially.

Figure 5a shows the spatial distribution of posterior CH₄ fluxes for 2021. Large emissions appear across regions where elevated XCH₄ was identified in Fig. 4, including South and East Asia, the Arabian Peninsula, central Africa, and the Americas. These hotspots reflect contributions from both natural wetlands and anthropogenic activities such as fossil-fuel extraction and waste management. The G2L4 flux distributions broadly align with established global CH₄ source patterns.

Figure 5b displays posterior-prior differences, highlighting regions where the G2L2 product induces substantial adjustments. Positive corrections occur mainly in tropical and subtropical regions—South Asia, central Africa, and northern to central South America—where prior estimates are typically uncertain and where ground-based observations are sparse. Negative adjustments, in contrast, appear over parts of China, the Persian Gulf region, Europe, and central North America. It is important to note that large posterior adjustments do not necessarily indicate strong observational constraints. Substantial corrections may arise either from (i) strong satellite constraints correcting prior biases, or (ii) weak observational constraints in regions with persistent cloud cover or limited viewing opportunities, where the inversion becomes more sensitive to residual retrieval biases or prior uncertainties. The degraded G2L2 coverage over central-eastern South America is one such example.

Overall, regions with large adjustments generally coincide with areas where prior uncertainties are large and observational coverage is limited, indicating where GOSAT-2 observations have the potential to provide valuable additional information beyond the prior estimates. These results emphasize both the value of G2L2 for improving flux estimates and the need for continued satellite observations to strengthen constraints in regions where current observational coverage remains limited.

Figure 6 presents monthly CH₄ flux variations for six latitude bands between 70°N and 50°S for the period May 2019–October 2022. Prior fluxes show only weak seasonal variability across all latitude bands, whereas posterior fluxes exhibit pronounced seasonal cycles, particularly north of 10°N. In the high northern latitudes (70°–50°N), posterior fluxes increase from around 2 Tg CH₄ month⁻¹ during winter to more than 10 Tg CH₄ month⁻¹ in July–August, reflecting well-known wetland-driven seasonality. Similar enhancements in seasonal amplitude are found in the 50°–30°N band and in the tropics (10°N–10°S), where prior fluxes show only limited variability.

A notable feature in Fig. 6 is the clear phase shift in the 30°N–10°N latitude band, where the posterior flux peak moves from May–June in the prior to July–August. This change is consistent with independent studies showing that CH₄ emissions from wetlands and rice paddies in South and East Asia tend to peak during the boreal summer monsoon (e.g., Zhang et al., 2020;




Figure 5. Spatial distribution of (a) annual posterior CH_4 fluxes for 2021 and (b) posterior-minus-prior differences ($\text{g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$).

East et al., 2024). Although satellite sampling in this region decreases during the monsoon season due to persistent cloud cover, the available G2L2 retrievals still exhibit enhanced XCH_4 in July-August, suggesting that the inversion adjusts the seasonal phase to align more closely with observational signals and to compensate for known biases in the prior. At the same time, the limited sampling also implies that part of the adjustment may reflect prior uncertainties rather than strong observational constraints. The posterior phase shift therefore likely results from a combination of satellite-derived information and structural

Overall, posterior fluxes peak in tropical and subtropical regions (30°N - 10°S), whereas prior fluxes show maximum emissions in the mid-latitude to tropical Northern Hemisphere. Averaged over 2020 and 2021, posterior fluxes increase by roughly 20-40 $\text{Tg CH}_4 \text{ yr}^{-1}$ in the tropics and subtropics relative to the prior, while high-latitude Northern Hemisphere fluxes remain broadly similar. The stronger seasonality and spatial redistribution evident in the posterior estimates suggest that GOSAT-2 observations contribute useful information for adjusting flux patterns in regions where prior constraints are relatively weak, particularly in tropical regions dominated by wetlands and agricultural activity.

4 Discussion

4.1 Comparison with CH_4 flux estimates derived from other observations

To evaluate the impact of the G2L2 product, we compared CH_4 flux estimates derived from inversions using the same model system as G2L4 but based on different observational datasets. Two datasets were employed: (1) fluxes estimated using only the G1L2 V03.05 satellite product, as in the Results section, and (2) fluxes from Niwa et al. (2025), which were derived from ground-based and aircraft (SURF+AIR) observations. The ground-based data include ObsPack GLOBALVIEWplus version 6.0 (Schuldt et al., 2023) and measurements from NIES and collaborative networks (Tohjima et al., 2002, 2014; Sasakawa et al., 2010, 2017; Terao et al., 2011; Nara et al., 2017; Nomura et al., 2017, 2021; Okamoto et al., 2018; Umezawa et al., 2025). Aircraft data consist of flask samples collected through the CONTRAIL program (Machida et al., 2008; Matsueda et al., 2015; Sawa et al., 2015; Umezawa et al., 2012) and Tohoku University campaigns (Umezawa et al., 2014). Although Niwa et al.




Figure 6. Monthly CH_4 flux variations ($\text{Tg CH}_4 \text{ month}^{-1}$) for six latitude bands between 70°N and 50°S , estimated from prior (broken line) and posterior (solid line) fluxes for May 2019–October 2022.

(2025) used the same inversion framework as this study, the prior biomass burning emissions differ between the two approaches. Niwa et al.’s dataset provides a benchmark based on dense in situ measurements, offering a complementary perspective to
 405 satellite-driven inversions and enabling assessment of consistency and potential biases in regional and seasonal CH_4 flux estimates.

We first compared the global annual CH_4 budgets among the three inversions. The G1L2-based inversion yielded 594.3 and 599.2 $\text{Tg CH}_4 \text{ yr}^{-1}$, and SURF+AIR-based inversion produced 593.2 and 595.4 $\text{Tg CH}_4 \text{ yr}^{-1}$ for 2020 and 2021, respectively. Differences across datasets remain within $\sim 10 \text{ Tg CH}_4 \text{ yr}^{-1}$, indicating that the choice of observational constraint does not
 410 materially change the global CH_4 budget in this inversion framework. This cross-dataset convergence provides confidence that the regional analyses below primarily reflect the characteristics of the observational constraints rather than structural instabilities of the inversion system itself.

Seasonal patterns of CH₄ fluxes are broadly consistent across the three inversions (Fig. A2), as noted in previous studies (e.g., Saunio et al., 2020; Niwa et al., 2025). Comparisons between G2L4 and G1L2-based inversions show small mean differences across latitude bands (-0.12 to 0.58 Tg CH₄ month⁻¹) and high correlations ($r = 0.83-0.99$), indicating similar seasonal cycles. Variance ratios (posterior-to-reference standard deviations; values >1 indicate stronger seasonal amplitude) range from 1.05 to 1.99—largest in 10°N–10°S—reflecting the combined influence of (i) differences in retrieval algorithms and product post-processing (including G1L2 bias-correction), (ii) differences in sampling and viewing geometry between missions (targeting strategies, off-nadir angles, cloud screening), and (iii) mission-dependent L1B radiance characteristics. In contrast, differences relative to the SURF+AIR-based inversion are more pronounced: the mean difference reaches -2.03 Tg CH₄ month⁻¹ in the tropics, root-mean-square difference exceeds 2.38 Tg CH₄ month⁻¹, and variance ratios rise to 4.36, indicating changes in both amplitude and phase. Correlations drop south of 10°S ($r = 0.55-0.31$), consistent with sparse in-situ coverage there.

Next, we extend our analysis to the regional scale using the 42 land regions defined in the GOSAT Level 4 product (Fig. A3). This framework enables a systematic comparison of CH₄ flux estimates derived from different observational datasets and helps identify where observational constraints most strongly affect inversion outcomes. At the regional scale (Fig. 7), the three inversions show broadly consistent spatial patterns, with all methods identifying similar high-emission regions and capturing the major continental-scale features of the CH₄ budget. This overall agreement indicates that G2L4, G1L2-based, and SURF+AIR-based inversions share a common large-scale structure and that the differences among them reflect regional observational characteristics rather than fundamental inconsistencies. Within this general consistency, regional differences become apparent. For example, in Southern Central Asia (CAS, RGN 30), the annual mean flux from G2L4 (50.5 Tg CH₄ yr⁻¹) is ~20% lower than the SURF+AIR-based estimate (60.2 Tg CH₄ yr⁻¹). Similar gaps (4–11 Tg CH₄ yr⁻¹) appear in North-eastern Siberia (SIB-NE, RGN 28) and the Arabian Peninsula-Western Middle East (WME, RGN 29). While these differences broadly correspond to sparse in-situ observational coverage, we cannot rule out contributions from region-dependent retrieval challenges—such as high aerosol loading, low surface albedo, or large solar zenith angles—that may influence satellite column retrievals. Thus, both observational distribution and potential retrieval systematics likely contribute to the regional spread.

Beyond differences in regional observational coverage, mission-specific observation strategies and retrieval characteristics also contribute to divergences among inversions. Both the GOSAT and GOSAT-2 missions allocate a considerable fraction of their observations to developed regions in order to target large emission sources such as megacities and power plants (Kuze et al., 2020). While this targeting strategy enhances the ability to capture urban greenhouse-gas enhancements (Ohyama et al., 2024), it can introduce a sampling bias toward high-concentration areas and may reduce the representativeness of regional fluxes. In addition, the two missions do not necessarily observe identical footprints, leading to differences in spatial sampling even within the same region. The case of Eastern Asia (RGN 32) illustrates this complexity. Although multiple TCCON sites are available in the region (Ohyama et al., 2020) and are used in the bias correction of G1L2, its flux estimates still differ substantially from those derived from the SURF+AIR-based inversion. This divergence arises despite both inversions being strongly constrained by extensive observational data. The differences likely reflect the fundamentally distinct nature of the observational datasets—satellite column retrievals versus in-situ and aircraft measurements—as well as differences in coverage, bias-correction approaches, data-selection criteria, and the spatial representativeness of observation sites. These factors can




Figure 7. Mean annual CH_4 fluxes ($\text{Tg CH}_4 \text{ yr}^{-1}$) for 2020 and 2021 across 42 land regions, estimated from prior (purple), G2L4 (orange), G1L2-based inversions (green), and ground-based inversions (blue). The geographical definitions and abbreviations of the regions are provided in Fig. A3.

lead to meaningful discrepancies in regional flux estimates even when the overall inversion frameworks are robust. Rather than indicating inconsistencies among the inversions, these results underscore the importance of considering observation-system characteristics when interpreting regional CH_4 budgets. They also highlight the need for a balanced combination of satellite, ground, and aircraft observations to improve representativeness and reduce structural uncertainties in future CH_4 monitoring systems.

Figure 8 also highlights where the three inversion systems show noticeable differences in their regional estimates. While G1L2-based inversions generally show amplitudes and correlations closer to those of G2L4, the SURF+AIR-based inversions display larger differences in regions where observational coverage is limited or where surface networks do not sufficiently capture the dominant emission regimes. This pattern is consistent with previous studies reporting that the performance of ground-based-driven inversions is strongly dependent on the spatial density and placement of in-situ sites (e.g., Stavert et al., 2022; Deng et al., 2025). Our results reflect this behavior across several regions, including Pampas (PAM, RGN 14), Central South America (SAM-, RGN 15), Northwestern Northern Africa (AFR-NNW, RGN 19), Maritime Southeast Asia (SEA-MAR, RGN 33), and Southeastern Australia (AUS-SE, RGN 36), where correlations with G2L4 are comparatively low. Amplitude differences also emerge in high-latitude regions such as Southwestern Siberia (SIB-SW, RGN 25) and Northeastern Siberia (SIB-NE, RGN 28), as well as in well-monitored areas like the Nordic Region (NOR, RGN 41) and mid-latitude North America (NAM-SW and NAM-NE; RGNs 5 and 8). These discrepancies do not undermine the overall consistency among the inversions; rather, they show how regional flux estimates respond differently to the distinct observational constraints available in each system. These regions illustrate where satellite observations offer broader spatial sampling that complements limited in-situ coverage. Looking ahead, integrating GOSAT-2 with complementary satellite missions-together with targeted in-situ




Figure 8. Correlation coefficients versus ratio of standard deviations for seasonal CH₄ flux variations across 42 land regions. Ratios are computed as the standard deviation of G1L2-based or ground-based (SURF+AIR) inversion divided by that of G2L4, and correlations are calculated between G2L4 and each of other two inversions.

deployments where feasible—will help reduce regional uncertainty and support the development of a more balanced, globally consistent CH₄-monitoring framework.

4.2 Insights into anthropogenic-dominated regions from satellite observations

Recent studies have demonstrated the capability of satellite observations to quantify national-scale CH₄ emissions using atmospheric inversion frameworks and to compare these estimates with national inventories (e.g., Shen et al., 2023; Janardanan et al., 2024; Deng et al., 2025). Inversion-based estimates, however, are generally sensitive to the total atmospheric CH₄ field, which integrates contributions from multiple source sectors, CH₄ sinks, and atmospheric transport processes. This property is not unique to satellite measurements; it is inherent to all top-down inversion systems. As a result, satellite-based inversions primarily optimize the net surface flux and cannot directly isolate sector-specific emissions. To gain insight into sectoral contributions despite this limitation, previous studies have adopted an indirect approach: in regions where the net prior flux is strongly dominated by a limited number of anthropogenic sectors, the direction and magnitude of posterior adjustments can help infer whether emissions from those dominant sectors are likely over- or underestimated. While this approach does not fully resolve

sector-specific budgets, it can provide useful qualitative information on sectoral patterns. Building on this concept, we assess whether the GOSAT-2 Level 4 (G2L4) product offers meaningful constraints on sector-dominated regions. We focus on regions where aggregated anthropogenic emissions account for more than 80% of prior total net surface flux, allowing us to examine whether posterior adjustments inferred from satellite observations align with expected sectoral contributions. Here, anthropogenic emissions refer to the sum of coal mining, oil and gas, landfill and waste, biofuel, enteric fermentation, manure management, and rice paddy fluxes.

Figure 9 illustrates that the G2L4 product provides useful observational information for anthropogenic-dominated regions, offering new insight into regional CH₄ flux patterns. In East Asia (EAS), where anthropogenic emissions constitute the majority of the prior total flux, the G2L4 posterior suggests a reduction of 12.3 Tg CH₄ yr⁻¹ relative to the prior. Although this posterior estimate (34.4 Tg CH₄ yr⁻¹) is lower than values reported in previous satellite-based studies such as Deng et al. (2025) and Chen et al. (2022), those studies also emphasize that East Asia exhibits large variability across inversion systems due to differences in observational datasets, retrieval configurations, and bias correction methods. The spread among inversion estimates therefore likely reflects this region's well-known sensitivity to differences in observational data sources, rather than any inconsistency among the inversion systems themselves. In this context, the G2L4 result contributes an independent satellite-based constraint that is useful for assessing the plausible range of anthropogenic emissions in East Asia.

In the Western Middle East (WME), where emissions are dominated by oil and gas production, the G2L4 posterior (18.2 Tg CH₄ yr⁻¹) falls within the range of previous estimates, including the Persian Gulf fossil emissions inferred by Deng et al. (2025). This consistency-combined with the region's frequent clear-sky conditions, which enable dense satellite sampling-demonstrates that GOSAT-2 observations provide useful observational information for fossil-fuel-dominated regions. At the same time, the strong influence of dust aerosols on SWIR retrievals (e.g., Yoshida et al., 2011) highlights the importance of continued refinement of aerosol correction methods to fully leverage the strengths of satellite data. Overall, the G2L4 results in EAS and WME illustrate how satellite-based inversions can provide valuable and region-specific insight into sectoral emissions, while also identifying observational and algorithmic factors that should be prioritized to further enhance the robustness of future CH₄ flux assessments.

Beyond East Asia and the Western Middle East, several other regions-such as Northern Africa and Central Asia-show only modest differences between prior and posterior estimates. Importantly, small posterior adjustments in these regions should not be interpreted as evidence that prior inventories are highly accurate. Rather, these areas are characterized by sparse ground-based observations and limited socioeconomic activity data, making the underlying inventories inherently uncertain (e.g., Ehret et al., 2022; Tibrewal et al., 2024). In such contexts, even small satellite-driven adjustments provide independent and valuable constraints that would otherwise be difficult to obtain. These regions also underscore a broader challenge: observational gaps persist in many parts of the world where establishing and maintaining surface networks is logistically difficult and financially costly (e.g., Velasco et al., 2017; Morino et al., 2018). Satellite observations offer spatial coverage that complements limited in-situ measurements and helps identify where additional observations would most improve regional constraints. Looking ahead, expanding the use of multi-satellite datasets-along with targeted deployments of ground-based sensors where




Figure 9. Anthropogenic CH₄ emissions (Tg CH₄ yr⁻¹) for nine regions—NAM-SW, AFR-NNW, AFR-NNE, WME, CAS, CAN, EAS, SEA-MAIN, and EUR-C—where anthropogenic sources in the a priori estimates account for more than 80% of prior fluxes. Bars show prior (blue) and posterior (orange) estimates derived from GOSAT-2 Level 4 product for 2020-2021.

feasible-offers a promising pathway for improving the sectoral attribution of CH₄ emissions in regions that have historically been under-constrained.

515 4.3 Limitations and interpretation of the G2L4 product

The G2L4 product provides monthly global CH₄ flux estimates derived from a satellite-only atmospheric inversion system. While the results presented in this study suggests that GOSAT-2 observations can improve the spatial and temporal distribution of CH₄ flux estimates, several limitations should be considered when interpreting the data product.

First, the current G2L4 release does not include formal posterior uncertainty estimates, posterior covariance matrices, inversion averaging kernels, or degrees-of-freedom-for-signal (DFS) diagnostics (Rodgers, 2000). The inversion is performed using a global high-dimensional 4D-Var framework based on NISMON-CH₄ (Niwa et al., 2017, 2025), and explicit estimation of posterior covariance matrices would require additional covariance calculations or large ensemble-based experiments that are not included in the operational G2L4 processing system. Consequently, quantitative estimates of uncertainty reduction and observation impact are not presently provided as part of the G2L4 product.

Second, it is important to distinguish between retrieval-level and inversion-level diagnostics. Averaging kernels and *a priori* profiles distributed with the G2L2 product are incorporated into the observation operator in order to account for retrieval smoothing effects and to ensure consistency between modeled and retrieved XCH₄ (Rodgers, 2000). However, these retrieval

averaging kernels should not be interpreted as inversion averaging kernels, nor do they provide information on posterior uncertainty reduction within the flux inversion framework.

530 Therefore, posterior-minus-prior flux differences presented in this study should primarily be interpreted as inversion adjustments rather than direct quantitative measures of observational constraint or uncertainty reduction. Interpretation of regional flux estimates should also take into account variations in observational coverage, prior uncertainty, retrieval characteristics, and the prescribed inversion error statistics.

535 Despite these limitations, comparisons with independent inversion systems indicate that the G2L4 product reproduces the large-scale characteristics of the global CH₄ budget and provides useful observational information in many regions where conventional surface observations remain sparse. Future developments of the G2L4 framework will investigate practical approaches for characterizing posterior uncertainty and information content while maintaining computational feasibility for global high-resolution inversions.

5 Conclusions

540 This study provides an overview of the GOSAT-2 mission and evaluates its XCH₄ data product, retrieved using absorption bands in the SWIR region, as well as Level 4 CH₄ flux estimates. Now that GOSAT-2 has been operating for more than seven years, the accumulated data volume enables a series of updates and improvements that were not previously possible.

545 The TANSO-FTS-2 instrument aboard GOSAT-2 incorporates several enhancements, including improved signal-to-noise ratio (SNR) and the Intelligent Pointing function. These improvements have led to a substantial increase in observational coverage and the volume of valid data acquisitions compared to the GOSAT mission. In particular, TANSO-FTS-2 has achieved significant progress in regions where observations were previously challenging, such as tropical areas and high-latitude zones. This expanded coverage and higher data density provide a improved observational basis for estimating national and regional CH₄ emissions through atmospheric inversion frameworks.

550 Using the G2L2 product, we estimated global CH₄ sources and sinks at a spatial resolution of 1.0° and a monthly time step. The global CH₄ budget derived from the posterior estimates is broadly consistent with annual budgets reported by the Global Carbon Project, confirming the reliability of the inversion framework. Compared to the prior estimates, the largest adjustments occur in regions with sparse ground-based coverage: positive corrections in tropical and subtropical areas and negative adjustments in parts of East Asia, the Middle East, and mid-latitude industrial regions. Posterior fluxes also exhibit pronounced seasonal cycles, particularly north of 10°S, reflecting improved representation of surface processes. These results
555 highlight the capability of GOSAT-2 to refine global and regional methane budgets.

To evaluate the G2L4 product, we compared posterior flux estimates with inversion results generated using the same model framework but constrained by different observational datasets: G1L2 product and a combination of ground-based and aircraft measurements. At the global and latitudinal scales, these comparisons show broad consistency, indicating that the accuracy of the G2L2 product is well supported. Regional assessments reveal that most areas exhibit good agreement among inversions,
560 although some regions display noticeable differences in annual budgets depending on the observational constraints applied.

In addition, we conducted a preliminary evaluation of anthropogenic CH₄ emissions at the regional scale. While this analysis demonstrates the potential of G2L2 for sectoral assessments, it also highlights regions where further investigation and methodological refinement are required.

The differences identified among inversion systems in this study indicate several directions for future improvement that directly follow from our regional analyses. First, the large spread observed in East Asia-where posterior estimates are particularly sensitive to the choice of satellite product and bias-correction approach-highlights the need for greater consistency among retrieval algorithms and for systematic cross-validation using independent datasets. Second, the strong agreement across inversions in the Western Middle East suggests that satellite observations can provide useful observational information for fossil-fuel-dominated regions when clear-sky conditions allow dense sampling, while also emphasizing the importance of refining aerosol-related radiance corrections to maximize the utility of SWIR measurements. Third, regions such as Northern Africa, Southeast Asia, and parts of South America, where inventories remain poorly constrained due to sparse ground-based observations, benefit substantially from the spatial coverage of satellite data. Coordinated use of multiple satellite missions, together with targeted in-situ deployments where feasible, would help reduce structural uncertainty in these under-observed areas. Overall, the results presented here highlight the value of GOSAT-2 observations for improving regional and sector-dominated CH₄ flux estimates and identify key areas where methodological harmonization and expanded observational coverage would further enhance confidence in future CH₄ assessments and satellite mission planning.

6 Code availability

7 Data availability

The NIES GOSAT-2 products, including the FTS-2 SWIR Level 2 Column-averaged Dry-air Mole Fraction Product and the Level 4A Global CH₄ Flux Product, are available from the NIES GOSAT-2 Product Archive (<https://prdct.gosat-2.nies.go.jp/index.html.en>). A detailed description of the variables, units, dimensions, missing-value conventions, and products included in the G2L4 dataset is provided in Appendix (Tables A1–A3). The NIES FTS SWIR Level 2 Column-averaged Dry-air Mole Fraction Product from GOSAT is available from the GOSAT Data Archive Service (<https://data2.gosat.nies.go.jp/>). EORC-JAXA Current Status: GOSAT-2 Operation Status. Available at: (https://www.eorc.jaxa.jp/GOSAT/GOSAT-2/gosat2_operationStatus.html).

Author contributions. MS conceptualized the study and performed the formal analysis. TM acquired funding. YN provided the model system. YY developed the G2L2 retrieval algorithm. Satellite operations and data acquisition were supported by HS, KS, and FK. The GOSAT-2 mission activities at NIES were supported by MS, YY, AK, IM, HN, HO, TS, YS, HY, and TM. MS wrote the initial draft of the manuscript. All authors contributed to reviewing and editing the final version of the manuscript.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Acknowledgements. This research has been supported by the GOSAT-2 project of NIES. The model simulations were completed using the NIES supercomputer.

Appendix A: Description of the G2L4 data product

To facilitate reuse of the G2L4 dataset, this appendix summarizes the principal variables included in the product, the associated data conventions, and information that is not currently provided in the operational release.

Table A1. Principal variables included in the GOSAT-2 Level 4 CH₄ flux product.

Variable	Unit	Description
lon	degrees_east	Longitude coordinate
lat	degrees_north	Latitude coordinate
time	hours since 01-01 00:00:00	Monthly time coordinate
flux_apri_ant_exricep	mg CH ₄ m ⁻² day ⁻¹	A priori anthropogenic emissions excluding rice paddies
flux_apri_ricep	mg CH ₄ m ⁻² day ⁻¹	A priori rice-paddy flux
flux_apri_wetl	mg CH ₄ m ⁻² day ⁻¹	A priori wetland flux
flux_apri_term	mg CH ₄ m ⁻² day ⁻¹	A priori termite flux
flux_apri_volc	mg CH ₄ m ⁻² day ⁻¹	A priori geological/volcanic flux
flux_apri_soilo	mg CH ₄ m ⁻² day ⁻¹	A priori soil oxidation flux
flux_apri_bmb	mg CH ₄ m ⁻² day ⁻¹	A priori biomass-burning flux
flux_apri_ocn	mg CH ₄ m ⁻² day ⁻¹	A priori ocean flux
flux_apos_tot	mg CH ₄ m ⁻² day ⁻¹	Posterior total surface CH ₄ flux

Table A2. General conventions used in the G2L4 product.

Item	Description
Format	NetCDF (CF-1.6 compliant)
Horizontal grid	1.0° latitude-longitude grid
Temporal resolution	Monthly
Missing value	-9999
Flux unit	mg CH ₄ m ⁻² day ⁻¹
Product version	V01.01
Data source	GOSAT-2 TANSO-FTS-2 SWIR L2 XCH ₄

Table A3. Information not included in the current G2L4 release.

Item	Availability
Posterior uncertainty estimates	Not provided
Posterior covariance matrix	Not provided
Inversion averaging kernels	Not provided
Degrees of freedom for signal (DFS)	Not provided




Figure A1. Global map of GOSAT-2 orbital path.

References

Canadell, J. G., Monteiro, P. M., Costa, M. H., Da Cunha, L. C., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogeli, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global carbon and other biogeochemical cycles and feedbacks, IPCC AR6 WGI, final government distribution, pp. 673–816, 2021.

600 Cao, M., Marshall, S., and Gregson, K.: Global carbon exchange and methane emissions from natural wetlands: Application of a process-based model, *Journal of Geophysical Research: Atmospheres*, 101, 14 399–14 414, 1996.

Chen, A., Zhang, Z., Poulter, B., Feng, L., Liu, H., Murray, L. T., Yu, X., Huang, Z., Gaeta, D. C., Morgan, K. L., Zhu, Q., and Miller, S. M.: Spatial and temporal distribution of global wetland methane emissions during 2019–2020 estimated from satellite observations, *Journal of Geophysical Research: Atmospheres*, 130, e2025JD044 354, 2025.

605 Chen, Z., Jacob, D. J., Nesser, H., Sulprizio, M. P., Lorente, A., Varon, D. J., Lu, X., Shen, L., Qu, Z., Penn, E., and Yu, X.: Methane emissions from China: a high-resolution inversion of TROPOMI satellite observations, *Atmospheric Chemistry and Physics*, 22, 10 809–10 826, <https://doi.org/10.5194/acp-22-10809-2022>, 2022.




Figure A2. Same as Fig. 6, but showing monthly CH₄ flux variations (Tg CH₄ month⁻¹) estimated from G2L4 (orange), G1L2-based inversions (green), and ground-based inversions (blue).

Crippa, M., Guizzardi, D., Solazzo, E., Muntean, M., Schaaf, E., Monforti-Ferrario, F., Banja, M., Olivier, J., Grassi, G., Rossi, S., and Vignati, E.: GHG emissions of all world - 2021 report, Publications Office of the European Union, <https://doi.org/doi/10.2760/173513>, 2021.

Deng, Z., Ciais, P., Hu, L., Martinez, A., Saunois, M., Thompson, R. L., Tibrewal, K., Peters, W., Byrne, B., Grassi, G., Palmer, P. I., Luijkx, I. T., Liu, Z., Liu, J., Fang, X., Wang, T., Tian, H., Tanaka, K., Bastos, A., Sitch, S., Poulter, B., Albergel, C., Tsuruta, A., Maksyutov, S., Janardanan, R., Niwa, Y., Zheng, B., Thanwerdas, J., Belikov, D., Segers, A., and Chevallier, F.: Global greenhouse gas reconciliation 2022, Earth System Science Data, 17, 1121–1152, <https://doi.org/10.5194/essd-17-1121-2025>, 2025.

East, J. D., Jacob, D. J., Balasus, N., Bloom, A. A., Bruhwiler, L., Chen, Z., Kaplan, J. O., Mickley, L. J., Mooring, T. A., Penn, E., Poulter, B., Sulprizio, M. P., Worden, J. R., Yantosca, R. M., and Zhang, Z.: Interpreting the seasonality of atmospheric methane, Geophysical Research Letters, 51, e2024GL108494, 2024.




Figure A3. Global land regions used in this study. The land surface is subdivided into 42 spatial units based on geographic location and broad climatic-biogeographic characteristics, following the regional classification adopted in the GOSAT Level 4 product.

Ehret, T., De Truchis, A., Mazzolini, M., Morel, J.-M., D'aspremont, A., Lauvaux, T., Duren, R., Cusworth, D., and Facciolo, G.: Global tracking and quantification of oil and gas methane emissions from recurrent sentinel-2 imagery, *Environmental science & technology*, 56, 10 517–10 529, 2022.

Etiope, G., Ciotoli, G., Schwietzke, S., and Schoell, M.: Gridded maps of geological methane emissions and their isotopic signature, *Earth System Science Data*, 11, 1–22, 2019.

Ferrario, F. M., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Lo Vullo, E., Solazzo, E., Olivier, J., and Vignati, E.: EDGAR v6.0 Greenhouse Gas Emissions, European Commission, Joint Research Centre (JRC)[data set] <http://data.europa.eu/89h/97a67d67-c62e-4826-b873-9d972c4f670b> (last access: 13 July 2022), 2021.

Frankenberg, C., Meirink, J. F., Bergamaschi, P., Goede, A. P. H., Heimann, M., Körner, S., Platt, U., van Weele, M., and Wagner, T.: Satellite cartography of atmospheric methane from SCIAMACHY on board ENVISAT: Analysis of the years 2003 and 2004, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2005JD006235>, 2006.

Fujii, Y.: Preconditioned optimizing utility for large-dimensional analyses (PopULar), *Journal of Oceanography*, 61, 167–181, 2005.

Imasu, R., Matsunaga, T., Nakajima, M., Yoshida, Y., Shiomi, K., Morino, I., Saitoh, N., Niwa, Y., Someya, Y., Oishi, Y., Hashimoto, M., Noda, H., Hikosaka, K., Uchino, O., Maksyutov, S., Takagi, H., Ishida, H., Nakajima, T. Y., Nakajima, T., and Shi, C.: Greenhouse gases Observing SATellite 2 (GOSAT-2): mission overview, *Progress in Earth and Planetary Science*, 10, 33, 2023.

- Ito, A.: Global termite methane emissions have been affected by climate and land-use changes, *Scientific Reports*, 13, 17 195, 2023.
- Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global terrestrial ecosystems and evaluation of
635 uncertainty, *Biogeosciences*, 9, 759–773, 2012.
- Janardanan, R., Maksyutov, S., Wang, F., Nayagam, L., Sahu, S. K., Mangaraj, P., Saunio, M., Lan, X., and Matsunaga, T.: Country-level methane emissions and their sectoral trends during 2009–2020 estimated by high-resolution inversion of GOSAT and surface observations, *Environmental Research Letters*, 19, 034 007, 2024.
- Kuze, A., O’Brien, D. M., Taylor, T. E., Day, J. O., O’Dell, C. W., Kataoka, F., Yoshida, M., Mitomi, Y., Bruegge, C. J., Pollock, H., Basilio,
640 R., Helmlinger, M., Matsunaga, T., Kawakami, S., Shiomi, K., Urabe, T., and Suto, H.: Vicarious calibration of the GOSAT sensors using the Railroad Valley desert playa, *IEEE transactions on geoscience and remote sensing*, 49, 1781–1795, 2010.
- Kuze, A., Kikuchi, N., Kataoka, F., Suto, H., Shiomi, K., and Kondo, Y.: Detection of methane emission from a local source using GOSAT target observations, *Remote Sensing*, 12, 267, 2020.
- Li, Q., Fernandez, R. P., Hossaini, R., Iglesias-Suarez, F., Cuevas, C. A., Apel, E. C., Kinnison, D. E., Lamarque, J.-F., and Saiz-Lopez, A.:
645 Reactive halogens increase the global methane lifetime and radiative forcing in the 21st century, *Nature Communications*, 13, 2768, 2022.
- Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., aan de Brugh, J., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., et al.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmospheric Measurement Techniques*, 14, 665–684, 2021.
- Machida, T., Matsueda, H., Sawa, Y., Nakagawa, Y., Hirokuni, K., Kondo, N., Goto, K., Nakazawa, T., Ishikawa, K., and Ogawa, T.: World-
650 wide measurements of atmospheric CO₂ and other trace gas species using commercial airlines, *Journal of Atmospheric and Oceanic Technology*, 25, 1744–1754, 2008.
- Maksyutov, S., Takagi, H., Valsala, V., Saito, M., Oda, T., Saeki, T., Belikov, D., Saito, R., Ito, A., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., and Yokota, T.: Regional CO₂ flux estimates for 2009–2010 based on GOSAT and ground-based CO₂ observations, *Atmospheric Chemistry and Physics*, 13, 9351–9373, 2013.
- 655 Matsueda, H., Machida, T., Sawa, Y., and Niwa, Y.: Long-term change of CO₂ latitudinal distribution in the upper troposphere, *Geophysical Research Letters*, 42, 2508–2514, <https://doi.org/https://doi.org/10.1002/2014GL062768>, 2015.
- Morino, I., Velasco, V. A., Hori, A., Uchino, O., Ohshima, H., Sakai, T., Izumi, T., Nagai, T., Bagtasa, G., Yoshida, Y., Kiel, M., Bukosa, B., Deutscher, N. M., Fisher, J. A., and Griffith, D. W. T.: Philippines TCCON Project: One-year Measurement Results and Future, in: *Light, Energy and the Environment 2018 (E2, FTS, HISE, SOLAR, SSL)*, p. FW5A.3, Optica Publishing Group,
660 <https://doi.org/10.1364/FTS.2018.FW5A.3>, 2018.
- Nara, H., Tanimoto, H., Tohjima, Y., Mukai, H., Nojiri, Y., and Machida, T.: Emission factors of CO₂, CO and CH₄ from Sumatran peatland fires in 2013 based on shipboard measurements, *Tellus B: Chemical and Physical Meteorology*, 69, 1399 047, 2017.
- NIES GOSAT-2 Project: Summary of the validation on GOSAT-2 TANSO-FTS-2 SWIR L2 Column-averaged Dry-air Mole Fraction Product (Ver. 02.10), 2025.
- 665 Nisbet, E. G., Manning, M., Dlugokencky, E., Fisher, R., Lowry, D., Michel, S., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., et al.: Very strong atmospheric methane growth in the 4 years 2014–2017: Implications for the Paris Agreement, *Global Biogeochemical Cycles*, 33, 318–342, 2019.
- Niwa, Y., Fujii, Y., Sawa, Y., Iida, Y., Ito, A., Satoh, M., Imasu, R., Tsuboi, K., Matsueda, H., and Saigusa, N.: A 4D-Var inversion system based on the icosahedral grid model (NICAM-TM 4D-Var v1. 0)–Part 2: Optimization scheme and identical twin experiment of
670 atmospheric CO₂ inversion, *Geoscientific Model Development*, 10, 2201–2219, 2017.

- Niwa, Y., Ishijima, K., Ito, A., and Iida, Y.: Toward a long-term atmospheric CO₂ inversion for elucidating natural carbon fluxes: technical notes of NISMON-CO₂ v2021.1, *Progress in Earth and Planetary Science*, 9, 42, 2022.
- Niwa, Y., Tohjima, Y., Terao, Y., Saeki, T., Ito, A., Umezawa, T., Yamada, K., Sasakawa, M., Machida, T., Nakaoka, S.-I., Nara, H., Tanimoto, H., Mukai, H., Yoshida, Y., Morimoto, S., Takatsuji, S., Tsuboi, K., Sawa, Y., Matsueda, H., Ishijima, K., Fujita, R., Goto, D., Lan, X., Schuldt, K., Heliasz, M., Biermann, T., Chmura, L., Necki, J., Xueref-Remy, I., and Sferlazzo, D.: Multi-observational estimation of regional and sectoral emission contributions to the persistent high growth rate of atmospheric CH₄ for 2020–2022, *Atmospheric Chemistry and Physics*, 25, 6757–6785, <https://doi.org/10.5194/acp-25-6757-2025>, 2025.
- Nomura, S., Mukai, H., Terao, Y., Machida, T., and Nojiri, Y.: Six years of atmospheric CO₂ observations at Mt. Fuji recorded with a battery-powered measurement system, *Atmospheric Measurement Techniques*, 10, 667–680, <https://doi.org/10.5194/amt-10-667-2017>, 2017.
- Nomura, S., Naja, M., Ahmed, M. K., Mukai, H., Terao, Y., Machida, T., Sasakawa, M., and Patra, P. K.: Measurement report: Regional characteristics of seasonal and long-term variations in greenhouse gases at Nainital, India, and Comilla, Bangladesh, *Atmospheric Chemistry and Physics*, 21, 16 427–16 452, <https://doi.org/10.5194/acp-21-16427-2021>, 2021.
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., Frankenberg, C., Kiel, M., Lindqvist, H., Mandrake, L., Merrelli, A., Natraj, V., Nelson, R. R., Osterman, G. B., Payne, V. H., Taylor, T. E., Wunch, D., Drouin, B. J., Oyafo, F., Chang, A., McDuffie, J., Smyth, M., Baker, D. F., Basu, S., Chevallier, F., Crowell, S. M. R., Feng, L., Palmer, P. I., Dubey, M., García, O. E., Griffith, D. W. T., Hase, F., Iraci, L. T., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Sussmann, R., Te, Y., Uchino, O., and Velasco, V. A.: Improved retrievals of carbon dioxide from Orbiting Carbon Observatory-2 with the version 8 ACOS algorithm, *Atmospheric Measurement Techniques*, 11, 6539–6576, <https://doi.org/10.5194/amt-11-6539-2018>, 2018.
- Ohyama, H., Morino, I., Velasco, V. A., Klausner, T., Bagtasa, G., Kiel, M., Frey, M., Hori, A., Uchino, O., Matsunaga, T., Deutscher, N. M., DiGangi, J. P., Choi, Y., Diskin, G. S., Pusede, S. E., Fiehn, A., Roiger, A., Lichtenstern, M., Schlager, H., Wang, P. K., Chou, C. C.-K., Andrés-Hernández, M. D., and Burrows, J. P.: Validation of XCO₂ and XCH₄ retrieved from a portable Fourier transform spectrometer with those from in situ profiles from aircraft-borne instruments, *Atmospheric Measurement Techniques*, 13, 5149–5163, <https://doi.org/10.5194/amt-13-5149-2020>, 2020.
- Ohyama, H., Yoshida, Y., and Matsunaga, T.: CH₄ and CO emission estimates for megacities: deriving enhancement ratios of CO₂, CH₄, and CO from GOSAT-2 observations, *Environmental Research Letters*, 19, 124 025, <https://doi.org/10.1088/1748-9326/ad89e0>, 2024.
- Okamoto, S., Tanimoto, H., Hirota, N., Ikeda, K., and Akimoto, H.: Decadal Shifts in Wind Patterns Reduced Continental Outflow and Suppressed Ozone Trend in the 2010s in the Lower Troposphere Over Japan, *Journal of Geophysical Research: Atmospheres*, 123, 12,980–12,993, <https://doi.org/https://doi.org/10.1029/2018JD029266>, 2018.
- Pandey, S., Gautam, R., Houweling, S., Van Der Gon, H. D., Sadavarte, P., Borsdorff, T., Hasekamp, O., Landgraf, J., Tol, P., Van Kempen, T., Hoogeveena, R., van Heesa, R., Hamburg, S. P., Maasakkers, J. D., and Aben, I.: Satellite observations reveal extreme methane leakage from a natural gas well blowout, *Proceedings of the National Academy of Sciences*, 116, 26 376–26 381, 2019.
- Rodgers, C. D.: Inverse methods for atmospheric sounding: theory and practice, World scientific Singapore, 2000.
- Saito, M. and Niwa, Y.: GOSAT-2 Level 4 CH₄ Algorithm Theoretical Basis Document, 2025.
- Saito, M., Shiraishi, T., Hirata, R., Niwa, Y., Saito, K., Steinbacher, M., Worthy, D., and Matsunaga, T.: Sensitivity of biomass burning emissions estimates to land surface information, *Biogeosciences*, 19, 2059–2078, 2022.
- Sasakawa, M., Shimoyama, K., Machida, T., Tsuda, N., Suto, H., Arshinov, M., Davydov, D., Fofonov, A., Krasnov, O., Saeki, T., Koyama, Y., and Maksyutov, S.: Continuous measurements of methane from a tower network over Siberia, *Tellus B*, 62, 403–416, 2010.

- Sasakawa, M., Machida, T., Ishijima, K., Arshinov, M., Patra, P. K., Ito, A., Aoki, S., and Petrov, V.: Temporal Characteristics of CH₄ Vertical Profiles Observed in the West Siberian Lowland Over Surgut From 1993 to 2015 and Novosibirsk From 1997 to 2015, *Journal of Geophysical Research: Atmospheres*, 122, 11,261–11,273, <https://doi.org/https://doi.org/10.1002/2017JD026836>, 2017.
- Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., et al.: The global methane budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, 2020.
- Saunio, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgersson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguia-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., van der Werf, G. R., Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, *Earth System Science Data*, 17, 1873–1958, <https://doi.org/10.5194/essd-17-1873-2025>, 2025.
- Sawa, Y., Machida, T., Matsueda, H., Niwa, Y., Tsuboi, K., Murayama, S., Morimoto, S., and Aoki, S.: Seasonal changes of CO₂, CH₄, N₂O, and SF₆ in the upper troposphere/lower stratosphere over the Eurasian continent observed by commercial airliner, *Geophysical Research Letters*, 42, 2001–2008, <https://doi.org/https://doi.org/10.1002/2014GL062734>, 2015.
- Schuldt, K. N., Mund, J., Aalto, T., Arlyn Andrews, Apadula, F., Jgor Arduini, Arnold, S., Baier, B., Bani, L., Bartyzel, J., Bergamaschi, P., Biermann, T., Biraud, S. C., Pierre-Eric Blanc, Boenisch, H., Brailsford, G., Brand, W. A., Brunner, D., Bui, T. P. V., Van Den Bulk, P., Francescopiero Calzolari, Chang, C. S., Huilin Chen, Lukasz Chmura, St. Clair, J. M., Sites Climadat, Coletta, J. D., Colomb, A., Condori, L., Conen, F., Conil, S., Couret, C., Cristofanelli, P., Cuevas, E., Curcoll, R., Daube, B., Davis, K. J., Dean-Day, J. M., Delmotte, M., Ankur Desai, DiGangi, E., DiGangi, J. P., Van Dinther, D., Elkins, J. W., Elsasser, M., Emmenegger, L., Fischer, M. L., Forster, G., Frumau, A., Fuente-Lastra, M., Galkowski, M., Gatti, L. V., Gehrlein, T., Gerbig, C., Francois Gheusi, Gloor, E., Goto, D., Hammer, S., Hanisco, T. F., Haszpra, L., Hatakka, J., Heimann, M., Heimann, M., Heliasz, M., Heltai, D., Henne, S., Hensen, A., Hermans, C., Hermansen, O., Hintsä, E., Hoheisel, A., Holst, J., Di Iorio, T., Iraci, L. T., Ivakhov, V., Jaffe, D. A., Jordan, A., Joubert, W., Kang, H.-Y., Karion, A., Kazan, V., Keeling, R. F., Keronen, P., Kers, B., Jooil Kim, Klausen, J., Kneuer, T., Ko, M.-Y., Kolari, P., Kominkova, K., Kort, E., Kozlova, E., Krummel, P. B., Kubistin, D., Kulawik, S. S., Kumps, N., Labuschagne, C., Lan, X., Langenfelds, R. L., Lanza, A., Laurent, O., Laurila, T., Lauvaux, T., Lavric, J., Choong-Hoon Lee, Haeyoung Lee, Lee, J., Lehner, I., Lehtinen, K., Leppert, R., Leskinen, A., Leuenberger, M., Levin, I., Levula, J., Lindauer, M., Lindroth, A., Loh, Z. M., Lopez, M., Lowry, D., Lunder, C. R., Machida, T., Mammarella, I., Manca, G., Manning, A., Marek, M. V., Marklund, P., Marrero, J. E., Martin, D., Martin, M. Y., Giordane A. Martins, Matsueda, H., McKain, K., Meinhardt, F., Menoud, M., Jean-Marc Metzger, Miles, N. L., Miller, C. E., Miller, J. B., Monteiro, V., Montzka, S., Moore, F., Moossen, H., Morgan, E., Josep-Anton Morgui, Morimoto, S., Munro, D., Mutuku, M., Myhre, C. L., Mölder, M., Jaroslaw Necki, Nichol, S., Nisbet, E., Niwa, Y., Njiru, D. M., Noe, S. M., O'Doherty, S., Obersteiner, F., Parworth, C. L., Peltola, O., Peters, W., Philippon, C., Piacentino, S., Pichon, J. M., Pickers, P., Pitt, J., Pittman, J., Plass-Dülmer, C., Platt, S. M., Popa, M. E., Prinzivalli, S., Ramonet, M., Richardson, S. J., Louis-Jeremy Rigouleau, Rivas, P. P., Röckmann, T., Rothe, M., Yves-Alain Roulet, Ju-Mee Ryoo, Santoni, G., Di Sarra, A. G., Sasakawa, M., Scheeren, B., Schmidt, M., Schuck, T., Schumacher, M., Seifert, T., Sha, M. K., Shepson, P., Sloop, C. D., Smith, P. D., Sørensen, L. L., De Souza, R. A. F., Spain, G., Steger, D., Steinbacher, M., Stephens, B., Sweeney, C., Taipale, R., Takatsuji, S., Thoning, K., Timas, H., Torn, M., Trisolino, P., Turnbull, J., Van Der Veen, C., Vermeulen, A., Vimont, I., Viner, B., Vitkova, G., De Vries, M., Watson, A., Weiss, R., Weyrauch, D., Wofsy, S. C., Worsey, J., Worthy, D., Xueref-Remy, I., Yates, E. L., Dickon Young, Yver-Kwok,

- C., Zaehle, S., Zahn, A., Zazzeri, G., Zellweger, C., and Mirosław Zimnoch: Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1983–2022; obspack_ch4_1_GLOBALVIEWplus_v6.0_2023-12-01, <https://doi.org/10.25925/20231001>, 2023.
- Shen, L., Jacob, D. J., Gautam, R., Omara, M., Scarpelli, T. R., Lorente, A., Zavala-Araiza, D., Lu, X., Chen, Z., and Lin, J.: National quantifications of methane emissions from fuel exploitation using high resolution inversions of satellite observations, *Nature Communications*, 14, 4948, 2023.
- Shindell, D., Ravishankara, A., Kuylensstierna, J. C., Michalopoulou, E., Höglund-Isaksson, L., Zhang, Y., Seltzer, K., Ru, M., Castelino, R., Faluvegi, G., et al.: Global methane assessment: Benefits and costs of mitigating methane emissions, Tech. rep., United Nations Environment Programme, 2021.
- Shiraishi, T., Hirata, R., and Hirano, T.: New inventories of global carbon dioxide emissions through biomass burning in 2001–2020, *Remote Sensing*, 13, <https://doi.org/10.3390/rs13101914>, 2021.
- Someya, Y., Yoshida, Y., Ohyama, H., Nomura, S., Kamei, A., Morino, I., Mukai, H., Matsunaga, T., Laughner, J. L., Velazco, V. A., Herkommer, B., Té, Y., Sha, M. K., Kivi, R., Zhou, M., Oh, Y. S., Deutscher, N. M., and Griffith, D. W. T.: Update on the GOSAT TANSO-FTS SWIR Level 2 retrieval algorithm, *Atmospheric Measurement Techniques*, 16, 1477–1501, <https://doi.org/10.5194/amt-16-1477-2023>, 2023.
- Stavert, A. R., Saunio, M., Canadell, J. G., Poulter, B., Jackson, R. B., Regnier, P., Lauerwald, R., Raymond, P. A., Allen, G. H., Patra, P. K., Bergamaschi, P., Bousquet, P., Chandra, N., Ciais, P., Gustafson, A., Ishizawa, M., Ito, A., Kleinen, T., Maksyutov, S., McNorton, J., Melton, J. R., Müller, J., Niwa, Y., Peng, S., Riley, W. J., Segers, A., Tian, H., Tsuruta, A., Yin, Y., Zhang, Z., Zheng, B., and Zhuang, Q.: Regional trends and drivers of the global methane budget, *Global Change Biology*, 28, 182–200, <https://doi.org/https://doi.org/10.1111/gcb.15901>, 2022.
- Suto, H., Kataoka, F., Kikuchi, N., Knuteson, R. O., Butz, A., Haun, M., Buijs, H., Shiomi, K., Imai, H., and Kuze, A.: Thermal and near-infrared sensor for carbon observation Fourier transform spectrometer-2 (TANSO-FTS-2) on the Greenhouse gases Observing SATellite-2 (GOSAT-2) during its first year in orbit, *Atmospheric Measurement Techniques*, 14, 2013–2039, 2021.
- Suto, H., Kataoka, F., Knuteson, R. O., Shiomi, K., Kikuchi, N., and Kuze, A.: Updated spectral radiance calibration on TIR bands for TANSO-FTS-2 onboard GOSAT-2, *Atmospheric Measurement Techniques*, 15, 5399–5413, <https://doi.org/10.5194/amt-15-5399-2022>, 2022.
- Takemura, T., Nakajima, T., Higurashi, A., Ohta, S., and Sugimoto, N.: Aerosol distributions and radiative forcing over the Asian Pacific region simulated by Spectral Radiation-Transport Model for Aerosol Species (SPRINTARS), *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Terao, Y., Mukai, H., Nojiri, Y., Machida, T., Tohjima, Y., Saeki, T., and Maksyutov, S.: Interannual variability and trends in atmospheric methane over the western Pacific from 1994 to 2010, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/https://doi.org/10.1029/2010JD015467>, 2011.
- Tibrewal, K., Ciais, P., Saunio, M., Martinez, A., Lin, X., Thanwerdas, J., Deng, Z., Chevallier, F., Giron, C., Albergel, C., Tanaka, K., Patra, P., Tsuruta, A., Zheng, B., Belikov, D., Niwa, Y., Janardanan, R., Maksyutov, S., Segers, A., Tzompa-Sosa, Z. A., Bousquet, P., and Sciare, J.: Assessment of methane emissions from oil, gas and coal sectors across inventories and atmospheric inversions, *Communications Earth & Environment*, 5, 26, <https://doi.org/10.1038/s43247-023-01190-w>, 2024.
- Tohjima, Y., Machida, T., Utiyama, M., Katsumoto, M., Fujinuma, Y., and Maksyutov, S.: Analysis and presentation of in situ atmospheric methane measurements from Cape Ochi-ishi and Hateruma Island, *Journal of Geophysical Research: Atmospheres*, 107, ACH 8–1–ACH 8–11, <https://doi.org/https://doi.org/10.1029/2001JD001003>, 2002.

785 Tohjima, Y., Kubo, M., Minejima, C., Mukai, H., Tanimoto, H., Ganshin, A., Maksyutov, S., Katsumata, K., Machida, T., and Kita, K.: Temporal changes in the emissions of CH₄ and CO from China estimated from CH₄ / CO₂ and CO / CO₂ correlations observed at Hateruma Island, *Atmospheric Chemistry and Physics*, 14, 1663–1677, <https://doi.org/10.5194/acp-14-1663-2014>, 2014.

Toon, G. C.: Solar Line List for the TCCON 2014 Data Release, <https://doi.org/10.14291/TCCON.GGG2014.SOLAR.R0/1221658>, 2015.

Umezawa, T., Machida, T., Ishijima, K., Matsueda, H., Sawa, Y., Patra, P. K., Aoki, S., and Nakazawa, T.: Carbon and hydrogen isotopic
790 ratios of atmospheric methane in the upper troposphere over the Western Pacific, *Atmospheric Chemistry and Physics*, 12, 8095–8113, <https://doi.org/10.5194/acp-12-8095-2012>, 2012.

Umezawa, T., Goto, D., Aoki, S., Ishijima, K., Patra, P. K., Sugawara, S., Morimoto, S., and Nakazawa, T.: Variations of tropospheric methane over Japan during 1988–2010, *Tellus B: Chemical and Physical Meteorology*, 66, 23 837, 2014.

Umezawa, T., Tohjima, Y., Terao, Y., Sasakawa, M., Müller, A., Saeki, T., Machida, T., Nakaoka, S.-I., Nara, H., Nomura, S., Nishihashi,
795 M., Mukai, H., Frey, M. M., Morino, I., Ohyama, H., Yoshida, Y., Zeng, J., Noda, H., Saito, M., Matsunaga, T., Sugita, T., Tanimoto, H., Niwa, Y., Ito, A., Yamashita, Y., Shirai, T., Ishizawa, M., Ishijima, K., Tsuboi, K., Sawa, Y., and Matsueda, H.: Long-term and interannual variations of atmospheric methane observed by the NIES and collaborative observation networks, *Progress in Earth and Planetary Science*, 12, 39, 2025.

Velazco, V. A., Morino, I., Uchino, O., Hori, A., Kiel, M., Bukosa, B., Deutscher, N. M., Sakai, T., Nagai, T., Bagtasa, G., Izumi, T., Yoshida,
800 Y., and Griffith, D. W. T.: TCCON Philippines: First Measurement Results, Satellite Data and Model Comparisons in Southeast Asia, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9121228>, 2017.

Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nature communications*, 10, 4584, 2019.

Worden, J. R., Bloom, A. A., Pandey, S., Jiang, Z., Worden, H. M., Walker, T. W., Houweling, S., and Röckmann, T.: Reduced biomass
805 burning emissions reconcile conflicting estimates of the post-2006 atmospheric methane budget, *Nature communications*, 8, 2227, 2017.

Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The total carbon column observing network, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 369, 2087–2112, 2011.

Yokota, T., Yoshida, Y., Eguchi, N., Ota, Y., Tanaka, T., Watanabe, H., and Maksyutov, S.: Global concentrations of CO₂ and CH₄ retrieved
810 from GOSAT: first preliminary results, *SOLA*, 5, 160–163, 2009.

Yoshida, Y. and Oshio, H.: GOSAT-2 TANSO-FTS-2 SWIR L2 Retrieval Algorithm Theoretical Basis Document, https://prdct.gosat-2.nies.go.jp/documents/pdf/ATBD_FTS-2_L2_SWL2_en_03.pdf, 2025.

Yoshida, Y., Ota, Y., Eguchi, N., Kikuchi, N., Nobuta, K., Tran, H., Morino, I., and Yokota, T.: Retrieval algorithm for CO₂ and CH₄ column
815 abundances from short-wavelength infrared spectral observations by the Greenhouse gases observing satellite, *Atmospheric Measurement Techniques*, 4, 717–734, 2011.

Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmospheric Measurement Techniques*,
820 6, 1533–1547, 2013.

- Yoshida, Y., Someya, Y., Ohyama, H., Morino, I., Matsunaga, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Iraci, L. T., Kivi, R., Notholt, J., Pollard, D. F., Té, Y., Velazco, V. A., and Wunch, D.: Quality Evaluation of the Column-Averaged Dry Air Mole Fractions of Carbon Dioxide and Methane Observed by GOSAT and GOSAT-2, *SOLA*, 19, 173–184, <https://doi.org/10.2151/sola.2023-023>, 2023.
- 825 Zhang, G., Xiao, X., Dong, J., Xin, F., Zhang, Y., Qin, Y., Doughty, R. B., and Moore III, B.: Fingerprint of rice paddies in spatial–temporal dynamics of atmospheric methane concentration in monsoon Asia, *Nature Communications*, 11, 554, 2020.
- Zhang, Y., Jacob, D. J., Lu, X., Maasakkers, J. D., Scarpelli, T. R., Sheng, J.-X., Shen, L., Qu, Z., Sulprizio, M. P., Chang, J., Bloom, A. A., Ma, S., Worden, J., Parker, R. J., and Boesch, H.: Attribution of the accelerating increase in atmospheric methane during 2010–2018 by inverse analysis of GOSAT observations, *Atmospheric Chemistry and Physics*, 21, 3643–3666, <https://doi.org/10.5194/acp-21-3643-2021>, 2021.