

Point-by-Point Responses to Referee Comments

Manuscript Number: essd-2026-179

Title: Grounded Icebergs around Antarctica: A High-Resolution Dataset Derived from Deep Learning and Sentinel-1 Synthetic Aperture Radar

Author(s): Kaihong Jiao et al.

Note: All page and line numbers mentioned in our responses below refer directly to the marked-up (tracked-changes) manuscript version.

Part 1: Responses to Anonymous Referee #1

General Comments

Reviewer: *This is an excellent paper! The authors present a very detailed, thoroughly thought-through algorithm for the detection of grounded icebergs and icebergs captured in fast ice, which includes efficient suggestions for handling problems such as look-alikes (small icebergs versus patches of rough sea ice or ocean clutter) and clusters of densely spaced smaller icebergs. The algorithm also considers additional information such as seabed topography and fast ice extent. Technical issues (different polarizations and different spatial resolutions of the SAR images available for the study) are discussed as well. The paper is very well written, and the figure captions are sufficiently detailed. Although this ML algorithm was developed only for use on Sentinel-1 images, it is a valuable basis also for applying it on images from other SAR sensors after corresponding training and testing. I recommend this manuscript for publication after my comments/suggestions below have been taken into account by the authors.*

Author's Response: We sincerely thank the reviewer for the highly positive assessment and constructive feedback. We are particularly encouraged that the reviewer recognises the value and thoroughness of our automated grounded iceberg detection algorithm. Below, we address each comment point-by-point and outline the specific changes made in the revised manuscript.

Specific Comments

Comment 1: *Page 2 line 21: "picket fence effect" related to which process?*

Author's Response: We thank the reviewer for pointing this out. The "picket fence effect" refers to the process where high-density clusters of grounded icebergs act as a series of physical anchors that stabilise landfast sea ice and restrict its drift and export. Since this term first appears in the Abstract, a brief clarifying phrase directly in the Abstract was added to ensure immediate clarity. The detailed physical mechanism of this effect has been further elaborated in our Results section.

Author's Changes in Manuscript: We have added a brief clarifying phrase ("acting as physical anchors to stabilise fast ice") to the Abstract at [Page 2, Line 21-22]. Furthermore, the physical mechanisms have been expanded in Section 4.2.2 at [Page 28, Line 568-570].

Comment 2: *Page 3 lines 45-48: I do not see the immediate relation between obstruction of sea ice export on the one hand and the formation of a coastal latent heat polynya which requires free sea ice transport away from the coast. The sentence on lines 51-54 points to a reference that seems to support the formulation lines 45-48, but intuitively it is not clear. Is it possible to add a SHORT explanation?*

Author's Response: We appreciate the opportunity to clarify this mechanism. The formation of a latent heat polynya requires winds to blow ice away from the coast or another barrier (e.g., fast ice, grounded bergs, glacier tongues). Grounded icebergs can act as topographic barriers that impede the advection of sea ice from (generally) east to west in the westward Antarctic Coastal Current (Massom et al., 2001; Fraser et al., 2023). This blocking effect prevents external pack ice from drifting into the polynya area, allowing offshore winds to clear the downstream (leeward) side of the icebergs, i.e., the polynya area, thereby sustaining the

polynya (e.g., see the conceptual schematic in Figure 11 of Fraser et al., 2023; Ohshima et al., 2013). An intuitive explanation has been added to clarify this dynamic.

Author's Changes in Manuscript: We have incorporated a short explanation regarding the barrier effect of grounded icebergs on pack ice advection and its relation to sustaining latent heat polynyas in Section 1 at [Page 3, Line 48-51].

Comment 3 & 4: Page 6 lines 133-135: The pixel size (I guess that is what is meant here) is 20 m for IWS and 40 m for EWS. (Which product was used: GRD-HR or GRD-MR?) However, it was found that the EWS mode reveals a stronger detection capability than IWS (page 30 line 608-610). Any idea what the reason is? The nominal NESZ is the same for both modes (-22dB). For the ESA S1 products, the ENL is smaller for IWS than for EWS (HR: 4.4 versus 2.8, MR: 81.8 versus 10.7, first number is IWS) (<https://sentiwiki.copernicus.eu/web/s1-products>). In your case it may be different, see question below. The main issue is that a higher ENL should intuitively increase the detection performance.

Page 6 lines 135-136: which are the default values for range and azimuth multi-looking in pyroSAR?

Author's Response: We thank the reviewer for the detailed technical questions regarding the Sentinel-1 SAR products and processing parameters. Regarding the processing details, we would like to clarify that the original manuscript contained an error where the IW mode data were mistakenly described as GRD products. Upon a thorough review of our data workflow, we confirmed that we utilized GRD-MR products for the EW mode (197 scenes) and Single Look Complex (SLC) products for the IW mode (77 scenes). During this data verification process, we discovered that 15 out of the 77 IW scenes had inadvertently been processed as GRD data in our earlier iteration. To maintain strict methodological consistency across the entire study, we have re-acquired the original SLC files for these 15 scenes, re-run them through our dedicated IW processing pipeline, and updated the circum-Antarctic dataset. Consequently, the Data Section of the manuscript has been thoroughly modified to accurately reflect these data types, software packages, and workflows. As a result of updating these 15 scenes, some numerical values and statistics in the results section have been slightly adjusted in the revised manuscript. We emphasize that these changes are minor and have no material impact on the overall trends, model performance, or conclusions of our study.

To fully answer your specific questions regarding the product parameters and the Equivalent Number of Looks (ENL):

- **EW GRDM Data:** The original EW GRDM product is distributed with 40 m pixel spacing and a nominal ENL of 10.7 (approximately 6 x 2 looks in range and azimuth, see Table below). We did not apply any additional multi-looking during pre-processing.
- **IW SLC Data:** Rather than using ESA's standard IW GRD products (with nominal ENLs of 4.4 or 81.8 for GRDH and GRDM), we processed the SLC data (nominal ENL = 1) directly to a 20 m pixel spacing Radiometric Terrain Corrected (RTC) product. This was undertaken using the Geoscience Australia sar-pipeline, which implements the NASA JPL RTC and ISCE3 software packages. The ISCE3 software applies adaptive multi-looking to achieve the 20 m pixel spacing. This adaptive approach results in a spatially varying ENL based on local terrain, yielding an approximate scene-level ENL of ~16 (estimated from resampling the nominal $\sim 2.3 \times 14.1$ m pixel spacing to 20×20 m). We have added the correct description to the manuscript.

For your convenience and reference, we have also attached the table outlining the Sentinel-1 product specifications to this response.

Acq. Mode	Product Type	Resolution Class	Resolution Rng x Azi [m]	Pixel Spacing Rng x Azi [m]	Num Looks Rng x Azi	ENL
SM	SLC	-	1.7x4.3 to 3.6x4.9	1.5x3.6 to 3.1x4.1	1x1	1
	GRD	FR	9x9	3.5x3.5	2x2	3.7
		HR	23x23	10x10	6x6	29.7
		MR	84x84	40x40	22x22	398.4
IW	SLC	-	2.7x22 to 3.5x22	2.3x14.1	1x1	1
	GRD	HR	20x22	10x10	5x1	4.4
		MR	88x87	40x40	22x5	81.8
EW	SLC	-	7.9x43 to 15x43	5.9x19.9	1x1	1
	GRD	HR	50x50	25x25	3x1	2.8
		MR	93x87	40x40	6x2	10.7
WV	SLC	-	2.0x4.8 to 3.1x4.8	1.7x4.1 to 2.7x4.1	1x1	1
	GRD	MR	52x51	25x25	13x13	123.7

Regarding the observation that the EW mode exhibits stronger detection capability than the IW mode, we agree with the reviewer that a higher ENL should intuitively increase detection performance. However, in our specific application, this discrepancy is primarily driven by data availability and model training rather than the inherent physics of the SAR modes. First, the EW data available for our study primarily consisted of cross-polarisation (HV), whereas the IW data were largely restricted to co-polarisation (HH). As cross-polarisation provides a significantly higher backscatter contrast between icebergs and surrounding sea ice compared to co-polarisation, the model naturally achieved better segmentation performance on the EW (HV) imagery. Second, our deep learning model was trained on a dataset predominantly composed of EW imagery, making the network inherently more attuned to the feature representations of EW data. A brief discussion has been added to clarify these specific reasons.

Author's Changes in Manuscript: We have thoroughly updated the text in Section 2 at [Page 6, Line 133-156] to clarify the use of EW GRD-MR and IW SLC data products, and corrected the description of the processing pipelines. We have added the following explanatory summary paragraph at [Page 6, Line 155-158]: *"In summary, the final gamma-nought intensity images obtained after pre-processing exhibit comparable visual characteristics for both EW and IW acquisition modes. However, IW images feature half the pixel spacing and slightly higher ENL compared to EW images. These differences stem from the intrinsic properties of the respective acquisition modes, where EW prioritizes broader spatial coverage at the expense of spatial and/or radiometric resolution."* Furthermore, the statistical outcomes in Section 4 and the relevant figures/tables have been updated to account for the re-processed 15 IW scenes. The structural analysis on EW vs IW mode detection performance has been embedded into Section 5.2 at [Page 31, Line 638-646].

Comment 5: Page 7 Fig 1: Only one region was used for validation with the goal of assessing "the generalisation of our proposed method across different seasons and possible limitations caused by surface melt during the summer" (page 17 lines 68-69). Was this region chosen because of the optimal availability of suitable data which was worse in other regions? Would you expect similar results in other regions?

Author's Response: We thank the reviewer for this important question. To clarify, the Adélie Land region was selected for the full-year, cross-seasonal validation not due to a lack of data elsewhere, but because it presents a highly complex and representative environmental background throughout the year. This region consistently features a challenging mix of open water, landfast sea ice, and fragmented pack ice, making it a rigorous testbed for assessing seasonal variations such as summer melt. Regarding other regions, we indeed expect similar results across the Antarctic margin. For the specific temporal window of our dataset (February–April), we have already demonstrated the algorithm's spatial generalisation across multiple other distinct coastal regions (e.g., Princess Ragnhild Coast, Princess Martha Coast), as detailed in Section 3.5.1 and Table 5. In terms of cross-seasonal performance, performance drops are generally as expected during the summer surface melt season, or under specific extreme conditions, such as severe wind-driven sea clutter (mainly in IWS mode, co-polarisation) or within heavily deformed sea ice rubble, where the inherent radar contrast between icebergs and the background is highly reduced. Because these factors are not unique to

Adélie Land, we expect the model to maintain a similar level of year-round stability across other regions, provided those specific extreme conditions are not present.

Author's Changes in Manuscript: We have expanded our justification for choosing the Adélie Land region in Section 3.5.2 at [Page 18, Line 395-397], and included a comprehensive discussion detailing expected model generalisation and constraints across other Antarctic circum-continental zones under extreme clutter conditions.

Comment 6: Page 10, section 3.1.3: I suggest to mention here the possible combinations of the image pairs (e.g. EWS with IWS, like-pol with cross-pol), and whether image sequences ≥ 3 are used.

Author's Response: We thank the reviewer for pointing this out. We realise that while the overall data combinations are mapped in Figure 1, the specific combinations used during the multi-temporal matching process could be stated more explicitly. Because our tracking framework analyses repeat-pass imagery along the same orbital track, the image pairs used for cross-temporal matching within any given trajectory almost exclusively consist of the same acquisition mode and polarisation (e.g., matching EW-HV to EW-HV, or IW-HH to IW-HH). While our algorithm operates on extracted shapes and could theoretically match targets across different modes in spatially overlapping areas, cross-mode matching does not occur in our dataset. We specifically opted for this approach because maintaining a consistent 12-day temporal resolution is highly beneficial for overall data consistency. A 12-day repeat cycle is already sufficient for identifying long-term stationary targets; inserting additional detections from different modes or polarisations to create mixed sequences would disrupt this temporal consistency without adding significant practical benefit to our grounded iceberg identification. Clarifications have been added.

Author's Changes in Manuscript: We have added explicit clarifications to Section 3.1.3 at [Page 10, Line 222-225] noting that matching image pairs maintain strict mode and polarisation consistency, and explicitly stating the utilisation of continuous multi-temporal image sequences where $n \geq 3$.

Comment 7 & 8: Page 11, lines 222 and 225-227, page 15 lines 317-319, page 16 Table 2: thresholds and training parameters: Are they based on own experiments or taken from the literature? How strong does the sensitivity of detection performance depend on the selection of different thresholds and training parameters? Page 12, line 253: How many training sessions were necessary for this study?

Author's Response: We thank the reviewer for raising these important questions. As these parameters govern different stages of our pipeline, we address them categorically below:

1. **Deep Learning Training Parameters:** The training hyperparameters (such as learning rate, batch size, and weight decay) were determined through our own experiments using Bayesian optimisation to balance model convergence and segmentation accuracy. Approximately 50 independent training sessions were conducted during model development and hyperparameter optimisation.
2. **Grounded Iceberg Identification Algorithm Thresholds:** The matching thresholds (e.g., centroid distance of 10 pixels, area difference of 0.5, and Intersection over Union (IoU) limits) are empirical, derived through careful tuning and extensive visual inspection of our identification results. Rather than conducting a strictly quantifiable sensitivity analysis—which is challenging given the immense variability in iceberg sizes, grounding behaviours (e.g., completely stationary versus rotating in place), and high-density spatial clustering (where densely packed targets frequently merge during segmentation, as illustrated in Supplementary Figures S1 and S4)—we focused on establishing a geometrically sound balance. For instance, the centroid distance threshold required careful consideration: a value that is too large risks incorporating slow-drifting icebergs into stationary trajectories (increasing false positives), while a value that is too strict would prematurely break trajectories due to minor spatial shifts caused by iceberg tilting or rotation. Furthermore, in regions characterised by an exceptionally high density of small, mobile icebergs, such as the Bellingshausen Sea, we observed a specific type of false positive where small, “similarly shaped” drifting bergs coincidentally occupied overlapping positions between consecutive 12-day SAR passes. Because these coincidental overlaps satisfy the algorithm's temporal continuity and spatial constraints, no simple threshold adjustment can completely eliminate them without breaking the retrieval of genuinely grounded targets. A clarification has been added to Section 5.2 (Limitations).
3. **Physical Constraint Filters:** As briefly stated in Section 3.4, the post-processing filters are based on a combination of established literature and our empirical observations. The bathymetric limits (e.g., > 1000 m excluding grounding entirely) are grounded in the literature regarding maximum expected keel depths for Antarctic icebergs (e.g., Dowdeswell and Bamber, 2007). The combined thresholds of Area and Sea Ice Concentration (SIC) in shallower waters (< 600 m) were empirically determined by observing the radiometric behaviour of sea-ice rubble. This effectively filters out a significant volume of smaller sea-ice artefacts (e.g., removing over 3,500 targets in shallow waters) while safely retaining larger, deep-keeled icebergs.

Author's Changes in Manuscript: We have incorporated details on the 50 deep learning optimization training sessions in Section 3.2 at [Page 11, Line 251-254]. We have added an explicit discussion regarding empirical parameter thresholds, geometric trade-offs, and the coincidental overlap limitation of mobile icebergs in high-density regions within Section 5.2 at [Page 31, Line 656-662]. The physical and literature-based foundations for bathymetric filtering have been enhanced in Section 3.4 at [Page 16, Line 364-366].

Comment 9: Page 13, line 260: The term "trajectory reconstruction" irritated me at first sight because I associated drifting icebergs. I expected a formulation like "test whether the iceberg is stationary". One could provide a hint to section 3.3.2 where a detailed explanation is given.

Author's Response: We agree with the reviewer that the term "trajectory reconstruction" can incorrectly imply drifting motion. To prevent any reader confusion, we have changed this terminology to "stationary target identification". Furthermore, we have included a direct cross-reference to Section 3.3.2, where the detailed explanation of the stationarity test is provided.

Author's Changes in Manuscript: We have replaced the phrase "trajectory reconstruction" with "stationary target identification" throughout Section 3 and included direct cross-references to Section 3.3.2 at [Page 13, Line 283-291].

Comment 10: Page 16, section 3.4: The criteria for excluding icebergs dependent on SIC and water depth trigger the following comments: (a) It may be possible that the SIC data and/or seabed topography data are not accurate which could mean that an iceberg IS stationary although the application of the criteria excludes it with the given complementary data. Its exclusion means that I get a false negative (page 31 lines 631-634: here, the false positives are mentioned). (b) A hint of typical keel depths of Antarctic icebergs would be helpful.

Author's Response: We thank the reviewer for these insightful points. (a) We acknowledge that inaccuracies or coarse spatial resolutions in the underlying auxiliary datasets (such as IBCSO bathymetry or AMSR2 SIC) could theoretically lead to the erroneous exclusion of a truly stationary, grounded iceberg, thereby creating a false negative. We have explicitly noted this inherent limitation. (b) We appreciate this suggestion. As briefly noted in our Introduction, Antarctic icebergs retain substantial subsurface drafts typically spanning from tens of metres to depths exceeding 500 metres (Dowdeswell and Bamber, 2007). To ensure this context is readily available when discussing our filtering criteria, we have added a brief cross-reference in Section 3.4 reminding the reader of these typical keel depths. We have also added specific altimetry references: "As reported by Luckman et al. (2010), nine grounded icebergs in the western Weddell Sea have maximum grounding depths estimated from satellite altimetry ranging from 240 - 370 m."

Author's Changes in Manuscript: We have added sentences in Section 5.2 at [Page 32, Line 675-677] highlighting how auxiliary data resolution and accuracy introduce uncertainties and potential false negatives. Section 3.4 at [Page 17, Line 364-366] now features explicit cross-references and text detailing typical Antarctic iceberg keel depths, including the altimetry-derived measurements from Luckman et al. (2010).

Comment 11: Page 24, line 510: "Table 7" in parentheses?

Author's Response: We have corrected this formatting error.

Author's Changes in Manuscript: The parenthetical formatting around the table citation has been corrected at [Page 24, Line 538].

Comment 12: Page 24, Table 6: In the caption for the table, it could be made clear that here sequences of n images are analysed and "start" and "end" refer to the image number in the sequence at which an iceberg was first and last identified as stationary by the algorithm. Or point to section 4.1.2.

Author's Response: We thank the reviewer for the valuable suggestion. We have updated the caption of Table 6 to explicitly define the "start" and "end" terminology in the context of the analysed image sequences, and we have added a pointer to Section 4.1.2 for further clarity.

Author's Changes in Manuscript: The caption of Table 6 at [Page 25, Table 6 Caption, Line 511-513] has been expanded to define "start" and "end" phases relative to sequence image indexes, along with an explicit cross-reference pointing to Section 4.1.2.

Comment 13: Page 29, line 567: ice(Fraser. => missing space

Author's Response: We have corrected this typographical error in the revised manuscript.

Author's Changes in Manuscript: The missing space within the citation string has been corrected at [Page 28, Line 593].

Comment 14: Page 31, line 634: do you really mean areas with complex "fast-ice topography" and not "seabed topography"?

Author's Response: Yes, we specifically meant "fast-ice topography". Complex and heavily deformed fast-ice surfaces (such as pressure ridges and rubble fields) can produce strong radar backscatter signals that closely mimic the radiometric signatures of small icebergs.

Author's Changes in Manuscript: We have retained "fast-ice topography" and added a brief parenthetical clarification at [Page 32, Line 674-675] ("e.g., heavily deformed pressure ridges and sea-ice rubble fields") to prevent potential reading ambiguity.

Comment 15: Page 32, section "Future Work": (a) I am sceptical regarding the mentioned merits of altimetry. You get only a profile of surface elevation which may not be sufficient to conclude on volume and morphology of single icebergs. Altimeter data are of course a good complement for iceberg detection as such but problems will occur for smaller icebergs. (b) Multi-source image analysis includes different types of sensors (optical, thermal, radar) with different coverage and spatial resolution. For radar we have, e.g. different frequencies and polarizations. It is mentioned that cross-pol is preferable in iceberg detection, frequency may also have an impact. If you have already an idea what an optimal multi-source scenario could be, you should mention it. (c) Not clear to me: what is the "global contextual information"? How does the knowledge of distant conditions really help to improve detection of a single iceberg in its local environment?

Author's Response: We thank the reviewer for these constructive prompts to expand our Future Work section.

Altimetry: We agree with the reviewer's scepticism to some extent. The reviewer is correct that altimetry (such as ICESat-2) only provides narrow surface elevation profiles. Relying on these sparse profiles is insufficient to fully reconstruct the 3D morphology of individual icebergs, as the satellite ground tracks may only capture a narrow slice or miss tiny targets entirely. Our initial phrasing in Section 5.3 was admittedly overly optimistic regarding "accurate reconstruction of iceberg morphology" for single icebergs. For single-iceberg 3D reconstruction, field-based observations utilising digital elevation models alongside multibeam echosounders remain irreplaceable. However, combining SAR-derived 2D surface areas with altimetry-derived draft/freeboard estimations remains a valuable approach for statistical volume approximation (e.g., Luckman et al., 2010). Furthermore, these limitations are poised to be mitigated by next-generation missions like the Earth Dynamics Geodetic Explorer (EDGE) satellite, targeted for the early 2030s, which will employ swath mapping laser altimetry using 40 simultaneous beams to create contiguous, high-density coverage (Garvin et al., 2026). We have tempered our claims, clarifying that altimetry primarily facilitates critical draft and volume estimations when fused with 2D SAR boundaries, while highlighting future swath-mapping altimeters like EDGE.

Multi-source scenario: An optimal multi-source tracking and volume-estimation scenario would integrate high-resolution optical imagery (e.g., Sentinel-2) to cross-validate and compensate for SAR's signal degradation during the summer surface melt season, alongside multi-frequency SAR (e.g., combining our C-band Sentinel-1 baseline with L-band sensors, such as Rose-L) to maximise iceberg-to-sea-ice contrast under complex conditions. These specific sensor and frequency combinations have been explicitly included. **Global contextual information:** By "global contextual information", we refer to the broader spatial context beyond the standard 250 x 250 pixel sub-image tile input to the model. Incorporating this wider view serves two critical purposes: it allows the model to perceive the continuous geometry of giant icebergs that span across multiple tiles, thereby preventing fragmented segmentation; and it provides richer environmental constraints, allowing the model to better distinguish true targets from large-scale background interference (e.g., extensive patches of severe sea clutter or heavily deformed fast ice). We have rewritten this sentence to explicitly define these benefits.

Author's Changes in Manuscript: Section 5.3 at [Page 33, Line 709-738] has been thoroughly revised. We have adjusted the wording regarding altimetric capabilities, introduced the future EDGE mission concept, detailed the optimal multi-sensor fusion scenario (Sentinel-2 optical and Rose-L L-band SAR), and explicitly defined the spatial scale and operational advantages of "global contextual information".

Part 2: Responses to Anonymous Referee #2

General Comments

Reviewer: *This study presents the first high-resolution circum-Antarctic grounded iceberg dataset by integrating Sentinel-1 SAR imagery with a ResUNet deep learning model, multi-temporal trajectory matching, and physical constraints derived from bathymetry and sea ice concentration. This dataset provides an important basis for future quantitative analyses of the impacts of grounded icebergs on the Southern Ocean environment, the growth and maintenance of landfast ice, and the stability of near-coastal ice shelves. Overall, this study has considerable scientific value and application potential, and I recommend publication after minor revision.*

Author's Response: We sincerely thank the reviewer for the positive assessment and for recognising the scientific value and application potential of our dataset. We appreciate these constructive comments, which are helpful for improving the rigour of our manuscript and the usability of our dataset. Below, we address each comment point-by-point and outline the specific changes made in the revised manuscript.

Specific Comments

Comment 1: *Although the data publication page provides a general description of the main GeoPackage dataset, it still appears to lack detailed information on the individual attribute fields and their units. I suggest that the authors add a README file or a field-description table to the dataset, clearly listing the meaning, data type, and unit of each variable.*

Author's Response: Thank you for the helpful suggestion. To avoid redundancy, all individual attribute fields, including their names, descriptions, data types, and units, are comprehensively detailed in Table 1 of the revised manuscript. We have updated the Data Availability section to direct readers to Table 1 for full variable details.

Author's Changes in Manuscript: We have noted this addition in the Data Availability section at [Page 35, Line 763-764].

Comment 2: *Page 5, Line 123: The expression “0.016 km² (40 × 40 m)” is potentially ambiguous, because 40 × 40 m corresponds to 0.0016 km² rather than 0.016 km². According to the Methods section, connected components smaller than 10 pixels were removed as noise. Therefore, I suggest that the authors clarify here that the minimum retained iceberg area is 0.016 km², corresponding to 10 pixels at a 40 m spatial resolution.*

Author's Response: We thank the reviewer for pointing out this mathematical typo. In the revised manuscript, we have corrected this expression to eliminate any ambiguity regarding pixel metrics.

Author's Changes in Manuscript: The text in Section 2 at [Page 5, Line 126-127] has been amended to read: “icebergs ranging down to 0.016 km² (corresponding to 10 pixels at 40 m pixel spacing)”.

Comment 3: *Page 6, Lines 139-140: The authors state that EW imagery was primarily used, while IW imagery was used to fill gaps in EW coverage. However, it remains unclear whether the algorithm directly performs cross-mode matching when two consecutive images over the same region are acquired in EW and IW modes, respectively, or whether IW imagery is only used to supplement spatial coverage and trajectories are continued only within the same acquisition mode.*

I suggest adding a set of comparison figures to visualise cross-mode matching, including the EW and IW scenes, the predicted iceberg masks under both modes, and the differences in iceberg outlines.

Author's Response: We would like to clarify that our algorithm does not perform cross-mode matching during the spatiotemporal tracking process. Because our tracking framework analyses repeat-pass imagery along the same orbital track, the image pairs used for cross-temporal matching within any given trajectory exclusively consist of the same acquisition mode and polarisation (e.g., matching EW-HV to EW-HV, or IW-HH to IW-HH). IW imagery was utilised solely to supplement spatial coverage where EW data were missing, rather than filling temporal gaps within a single trajectory sequence. To prevent reader confusion, we have added an explicit clarification to Section 3.1.3 of the revised manuscript to emphasise that trajectory matching is restricted to consistent acquisition modes and polarisations.

Author's Changes in Manuscript: We have added explanatory text in Section 3.1.3 at [Page 10, Line 222-225] explicitly stating that tracking trajectories are processed strictly within homogeneous sensor acquisition modes and polarisations, avoiding cross-mode sequence stitching.

Comment 4: Page 10, Line 198: The manuscript states that repeat-pass images with temporal baselines of 12 days or multiples thereof were used for grounded iceberg identification for each orbital track. It is unclear whether all Antarctic coastal regions have at least two SAR images available for matching. I suggest that the authors clarify whether any regions had only a single valid SAR acquisition. If such regions exist, the authors should explain whether they were excluded from grounded iceberg identification.

Author's Response: We thank the reviewer for highlighting this detail. We clarify that all Antarctic coastal regions covered in our dataset have at least two consecutive valid SAR images available, with a temporal baseline of 12 days or multiples thereof. There were no regions with only a single valid SAR acquisition within our observation window. Therefore, no coastal sectors were excluded from the grounded iceberg identification process due to a lack of temporal sequences. We have added an explicit statement to Section 3.1.3 to ensure complete clarity.

Author's Changes in Manuscript: We have appended a clarifying sentence to Section 3.1.3 at [Page 10, Line 220-222], explicitly confirming that all processed coastal zones contained a minimum of two valid consecutive SAR repeat acquisitions, and no sector was excluded due to data sequence deficiency.

Comment 5: Page 16, Line 198: The authors use SIC and bathymetry data to remove iceberg candidates, but these auxiliary datasets have limitations in spatial resolution and accuracy. In particular, the AMSR2 SIC product has a spatial resolution of 10 km, which is much coarser than the scale of small icebergs. It may therefore fail to accurately represent the local sea ice conditions surrounding individual icebergs. As a result, some truly grounded icebergs may be incorrectly removed due to SIC uncertainty, leading to false negatives. I suggest that the authors further discuss this potential source of uncertainty.

Author's Response: We agree with the reviewer's insight. In the revised manuscript, we have added a dedicated paragraph to Section 5.2 (Limitations) discussing the uncertainties and potential false negatives associated with the use of auxiliary datasets of different resolutions (such as AMSR2 SIC and IBCSO bathymetry) in the filtering process.

Author's Changes in Manuscript: A comprehensive evaluation of auxiliary dataset constraints, highlighting spatial resolution mismatches (e.g., 10 km AMSR2 grid cell scales versus sub-kilometer iceberg scales) and their downstream impacts on false-negative metrics, has been detailed in Section 5.2 at [Page 32, Line 675-677].

Comment 6: Page 19-20, Lines 424-435: The authors state that the model shows strong cross-seasonal generalisation. However, the main validation area is a 50 × 50 km region in Adélie Land. Whether the validation results from this region are representative of the entire circum-Antarctic coastline requires further clarification.

Author's Response: We thank the reviewer for highlighting this point. As noted in our response to Reviewer 1 (Comment 5), the Adélie Land region was selected for the cross-seasonal, full-year validation primarily because it consistently presents a highly complex environmental background—specifically, a challenging mix of open water, landfast sea ice, and fragmented pack ice. This makes it an ideal, rigorous testbed for evaluating seasonal dynamics like summer surface melt.

To address the question of circum-Antarctic generalisation, we have provided comprehensive evidence in Section 3.5.1 and Table 5 (Extended Validation Dataset), where we evaluated single-frame detection performance across multiple other distinct coastal sectors (e.g., Princess Ragnhild Coast, Princess Martha Coast, and Enderby Land), spanning various combinations of acquisition modes and polarisations. These robust metrics demonstrate that our model maintains strong spatial generalisation capabilities across different Antarctic margins. Furthermore, our analysis indicates that cross-seasonal performance fluctuations are driven by specific environmental extremes (e.g., severe wind-driven sea clutter or heavily deformed sea ice rubble) rather than the geographic location itself. Because these driving physical factors are not unique to Adélie Land, we expect the model to maintain a similar level of year-round stability across the entire circum-Antarctic coastline.

Author's Changes in Manuscript: We have cross-referenced the single-frame spatial validation framework (Section 3.5.1 and Table 5) within the seasonal validation text of Section 3.5.2 at [Page 18, Line 393-397] to emphasize geographic diversity, and augmented the section with a physical discussion showing how localized clutter mechanics, rather than geographic coordinates, dictate performance stability across the circum-Antarctic coastline.

References

Dowdeswell, J. A., & Bamber, J. L. (2007). Keel depths of modern Antarctic icebergs and implications for sea-floor scouring in the geological record. *Marine Geology*, 243(1-4), 120-131. doi: <https://doi.org/10.1016/j.margeo.2007.04.008>

Fraser, A. D., Wongpan, P., Langhorne, P. J., Klekociuk, A. R., Kusahara, K., Lannuzel, D., ... & Wienecke, B. (2023). Antarctic landfast sea ice: A review of its physics, biogeochemistry and ecology. *Reviews of Geophysics*, 61(2), e2022RG000770. doi: <https://doi.org/10.1029/2022RG000770>

Garvin, J. B., McDonagh-Smith, P., Peters, R. M., Fricker, H. A., Armston, J. D., Blair, J. B., & Luthcke, S. B. (2026). Return on investment and economic impact of a NASA-funded swath mapping laser altimeter orbiter for Earth surface elevations situational awareness in 2031–2040. *New Space*, 21680256261434504. doi: <https://doi.org/10.1177/21680256261434504>

Luckman, A., Padman, L., & Jansen, D. (2010). Persistent iceberg groundings in the western Weddell Sea, Antarctica. *Remote Sensing of Environment*, 114(2), 385-391. doi: [10.1016/j.rse.2009.09.009](https://doi.org/10.1016/j.rse.2009.09.009)

Massom, R. A., Hill, K. L., Lytle, V. I., Worby, A. P., Paget, M. J., & Allison, I. (2001). Effects of regional fast-ice and iceberg distributions on the behaviour of the Mertz Glacier polynya, East Antarctica. *Annals of Glaciology*, 33, 391-398. doi:[10.3189/172756401781818518](https://doi.org/10.3189/172756401781818518)

Ohshima, K. I., Fukamachi, Y., Williams, G. D., Nihashi, S., Roquet, F., Kitade, Y., ... & Wakatsuchi, M. (2013). Antarctic Bottom Water production by intense sea-ice formation in the Cape Darnley polynya. *Nature Geoscience*, 6(3), 235-240. doi: <https://doi.org/10.1038/ngeo1738>

Grounded Icebergs around Antarctica: A High-Resolution Dataset Derived from Deep Learning and Sentinel-1 Synthetic Aperture Radar

Kaihong Jiao^{1,2}, Alexander D. Fraser^{1,2}, Johannes Lohse^{2,1,3}, Pat Wongpan^{1,2,4}, Caitlin Adams⁵, and Alexander C. Bradley⁵

¹Institute for Marine and Antarctic Studies (IMAS), University of Tasmania, nipaluna / Hobart, Australia

²Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, nipaluna / Hobart, Australia

³Department of Physics and Technology, UiT The Arctic University of Norway, Tromsø, Norway

⁴Australian Antarctic Division, Department of Climate Change, Energy, the Environment and Water, Kingston, Tasmania, Australia

⁵Geoscience Australia, Canberra, Australia

Correspondence: Kaihong Jiao (kaihongjiao0703@gmail.com)

Abstract. Icebergs frequently run aground on shoals on the continental shelf around Antarctica. Once rendered immobile, they can anchor landfast sea ice (fast ice) and their supply of limiting trace nutrients contribute to driving coastal marine productivity. They are also associated with seabed scouring, with implications for benthic marine ecosystems. Despite their importance, there is currently a lack of accurate, continent-wide automated mapping of grounded iceberg distribution and size and as a consequence, no large-scale, complete map of grounded icebergs exists. To address these gaps, this study implements an automated grounded iceberg detection framework based on Sentinel-1 Synthetic Aperture Radar (SAR) imagery, integrating a proposed ResUNet deep-learning network with a multi-temporal identification algorithm, incorporating strict physical constraints derived from bathymetry and sea ice concentration to mitigate environmental false positives. The method demonstrates strong robustness against interference from complex coastal conditions, achieving a detection F1 score exceeding 0.91 and successfully reducing the minimum detectable iceberg size to 0.016 km^2 . Since the presence of fast ice makes distinction between “truly grounded icebergs” and “those held motionless by fast ice”, we capitalise on the unprecedented low fast ice extent in early 2025 (late February to early April) to construct the first high-resolution, continent-wide dataset of grounded icebergs around Antarctica. A total of ~~38,905~~39,619 stationary icebergs were identified on the Antarctic continental shelf. We partition these stationary targets into “high-confidence grounded icebergs” and fast ~~ice-entrapped~~ice-entrapped candidates. Analysis shows that ~~70.5~~67.4 % of these targets are identified as high-confidence grounded. Across the entire dataset, tiny icebergs ($< 1 \text{ km}^2$) dominate numerically, accounting for ~~92.2~~92.6 % of the total. These high-density grounded iceberg clusters are primarily concentrated on shallow continental shelves and west of (i.e., downstream of) the actively disintegrating fronts of ice shelves, forming complex and discontinuous “grounded iceberg chains”. Our dataset reveals that these grounded icebergs cover a combined area of ~~13,719~~430 km^2 and are widely distributed along ~~56.7~~56.5 % of the Antarctic coastline, with just 14.2 % of the coastline containing 80 % of all grounded icebergs. Crucially, while typically overlooked, tiny icebergs ($< 1 \text{ km}^2$)

contribute ~~52.6~~53.9 % of this total grounded area. These dense clusters imply a potential “picket fence effect”, ~~providing a~~
acting as a series of physical anchors to stabilise fast ice. This provides a quantitative baseline for modelling fast ice stabil-
ity, assessing nutrient fluxes, and mapping benthic habitats. The grounded iceberg dataset (Jiao et al., 2026) is available at
<https://doi.org/10.25959/54sx-pt47>.

Antarctic icebergs originate from the calving of ice shelves and tidewater glaciers, a fundamental process that regulates the mass balance of the Antarctic Ice Sheet (Rignot et al., 2013; Greene et al., 2022). The discharge of ice into the Southern Ocean is not merely a passive decay but a critical component of ice-sheet dynamics; recent assessments indicate that calving accounts for approximately half of the continent's total mass loss, comparable to that of basal melting (Rignot et al., 2013; Greene et al., 2022). These calving events produce floating ice masses ranging from small fragments to massive tabular icebergs, which serve as proxies for ice-shelf break-up processes (Scambos et al., 2009). Importantly, these icebergs retain substantial subsurface drafts, typically spanning from tens of metres to depths exceeding 500 metres (Dowdeswell and Bamber, 2007). It is this substantial keel depth that predisposes them to interact with the seafloor on shallow continental shelves, transforming them from drifting bodies into stationary grounded features.

Once detached from their parent ice shelves, icebergs drift across the Southern Ocean, driven by the complex interplay of wind stress, ocean currents, the Coriolis force, and sea ice dynamics (Gladstone et al., 2001; Bigg et al., 1997). More recent models also account for interactions between icebergs, sea ice, and wave radiation stress (e.g. Merino et al., 2016; Rackow et al., 2017), which are increasingly recognised as essential components in Earth System Models (Vancoppenolle et al., 2025). When these deep-draft icebergs drift across the Antarctic continental shelves, where water depths are often less than 500 m (Dorschel et al., 2022), their keels can contact the seabed (Dowdeswell and Bamber, 2007; Olivé Abelló et al., 2025). This interaction anchors drifting ice, establishing them as “grounded icebergs”, stationary features that can remain in place for periods ranging from days to several decades (Woodworth-Lynas et al., 1991; Leane and Maddison, 2018).

Grounded icebergs play a critical role in the Antarctic sea ice system, serving as key anchor points for Antarctic landfast sea ice (fast ice) (Wright and Priestley, 1922) (“fast ice”; Wright and Priestley, 1922), and together with fast and pack ice, ice shelves and coastline, they form a complex coastal ice–ocean interaction network (Porter-Smith et al., 2021; Fraser et al., 2023; Vancoppenolle et al., 2025). Grounded icebergs not only fix and stabilise fast ice, prolonging its residence and enhancing the continuity and spatial cohesion of coastal ice zones (Massom et al., 2001; Li et al., 2020; Fraser et al., 2023), ~~but also, by obstructing sea ice export and altering ice distribution, promote~~. Furthermore, they act as topographic barriers that impede the upstream advection of sea ice; by blocking external pack ice from entering, they allow offshore winds to clear the downstream (leeward) side of sea ice (e.g., see Figure 11 in Fraser et al., 2023). This obstruction and alteration of ice distribution promotes polynya formation, the development of thick ice zones, and influence the intrusion of warm currents and ice shelf melting (Nakayama et al., 2014; Fraser et al., 2019; Bett et al., 2020; Olivé Abelló et al., 2025). Beyond these hydrodynamic effects, the deep keels of grounded icebergs physically scour the seafloor, creating extensive plough marks and resuspending sediments, which significantly reshapes local bathymetry and disturbs benthic environments (Woodworth-Lynas et al., 1991; Olivé Abelló et al., 2025; Kostov et al., 2026). Ecologically, grounded icebergs act as topographic barriers that impede sea ice drift, thereby facilitating the formation of latent heat polynyas (Ohshima et al., 2013), which are critical open-water areas that sustain high biological productivity, and they also serve as unique physical substrates for distinct microbial communities (Cefarelli et al., 2015; Bett et al., 2020). Their meltwater and nutrient release significantly enhance local productivity and alleviate nutrient

limitations for phytoplankton (Duprat et al., 2016; Lucas et al., 2025; Olivé Abelló et al., 2025). In addition, icebergs are an
60 important source of iron for polar oceans, and their heterogeneous release affects regional nutrient availability (Raiswell et al.,
2008; Herraiz-Borreguero et al., 2016; Hopwood et al., 2019). Therefore, monitoring the distribution and melting processes of
grounded icebergs is crucial for understanding polar marine ecology and climate feedbacks.

Traditionally, iceberg monitoring has relied heavily on manual analysis of satellite imagery. Operational agencies like the
U.S. National Ice Center track and name Antarctic icebergs with a major axis greater than 10 nautical miles (approximately
65 18.5 km), or recently, those with an area exceeding 20 square nautical miles (U.S. National Ice Center, 2024). This leaves a vast
number of smaller icebergs unmonitored. One of the main tools for large-scale iceberg monitoring and observations at fine spa-
tial resolution is spaceborne Synthetic Aperture Radar (SAR). Unlike optical remote sensing, SAR can penetrate clouds or fog
and does not rely on solar illumination, thus enabling year-round all-weather observations that are largely unaffected by atmo-
spheric conditions (Dierking, 2013; Lohse et al., 2020). By emitting microwave pulses and receiving the backscattered signals
70 from the surface, SAR generates high-resolution imagery that captures the texture, roughness, and backscatter characteristics of
sea ice and icebergs, thereby allowing discrimination between different ice types and open water (Willis et al., 1996; Dierking,
2013). In particular, the Sentinel-1 satellite constellation, launched by the European Space Agency, has provided continuous
and systematic observations of polar environments since 2014, with unprecedented Antarctic coverage (Torres et al., 2012).
Operating at C-band frequency with dual polarisation, its wide-swath acquisition modes achieve an effective balance between
75 spatial resolution and coverage, making it a core data source for current Antarctic sea ice and iceberg studies (Dierking and
Wesche, 2014).

To bridge the gap left by manual analysis and map the vast population of smaller icebergs, machine learning and deep learn-
ing technologies have been widely applied in iceberg detection tasks (Barbat et al., 2019; Braakmann-Folgmann et al., 2023;
Chen et al., 2026). Owing to their efficient data processing capabilities and adaptive learning characteristics, these technologies
80 have significantly enhanced the level of automation and detection accuracy. By fully utilising the high spatiotemporal resolution
and polarimetric characteristics of Sentinel-1 SAR data, machine learning algorithms can effectively extract iceberg texture,
shape, and backscatter features, thereby improving detection performance for small icebergs ($< 1 \text{ km}^2$; Barbat et al., 2021;
Koo et al., 2023; Chen et al., 2026). In particular, U-Net-based deep learning methods, through end-to-end feature extraction,
substantially enhance the ability to capture iceberg texture, shape, and contextual information under complex backgrounds,
85 showing greater robustness and detection accuracy (Ronneberger et al., 2015; He et al., 2016; Braakmann-Folgmann et al.,
2023).

Despite these technological advances, automated iceberg detection using SAR faces inherent seasonal limitations. Surface
meltwater during the summer months confounds the SAR backscatter signals, meaning most automated detection algorithms
perform reliably only during winter or early spring (Wesche and Dierking, 2012; Braakmann-Folgmann et al., 2023). Fur-
90 thermore, accurate identification of tiny icebergs (e.g., $< 0.4 \text{ km}^2$ or even $< 0.1 \text{ km}^2$) remains difficult year-round, as they
are often indistinguishable from speckle noise or complex multi-year ice textures in coastal zones (Barbat et al., 2021; Koo
et al., 2023; Evans et al., 2023; Færch et al., 2023). Beyond general detection, a fundamental challenge in isolating specifically
grounded icebergs is that stationarity in consecutive SAR images does not uniquely imply grounding. Floating icebergs are

frequently immobilised by fast ice (Fraser et al., 2023), making them indistinguishable from genuinely grounded targets based on motion analysis alone. This ambiguity is most pronounced during winter and early spring when fast ice extent is at its peak. This creates a conflicting seasonal requirement, which highlights a critical limitation in current iceberg products. For instance, while Chen et al. (2026) recently produced a high-resolution circum-Antarctic dataset of small icebergs ($> 0.04 \text{ km}^2$), their analysis was strictly limited to October of each year to ensure optimal general SAR detection performance. Given the extensive presence of fast ice during this season (Fraser et al., 2021), their snapshot contains multitudes of immobilised floating icebergs. Consequently, their dataset is unsuitable for capturing grounded iceberg dynamics and unable to reflect seasonal variations in their distribution.

Crucially, there is currently no dedicated automated product specifically for grounded icebergs. Although recent studies have mapped grounded icebergs to investigate their interaction with fast ice, these efforts largely rely on manual interpretation, which is labour-intensive and difficult to scale temporally (Li et al., 2020; Kostov et al., 2026). For example, Li et al. (2020) identified 24 specific zones containing groups of small grounded icebergs to demonstrate their significance in maintaining landfast ice. However, rather than extracting individual iceberg boundaries continent-wide, their approach was limited to demarcating broad regions and relying on detailed case studies for specific giant icebergs. Furthermore, such data record grounded icebergs merely as broad bounding zones or point locations, lacking essential geometric attributes such as precise shape and surface area, which are vital for quantifying their hydrodynamic drag and mechanical anchoring capacity within the fast ice matrix (Massom et al., 2001; Fraser et al., 2023).

To address the uncertainty between grounding and temporary entrapment by fast ice, this study deliberately restricts its temporal focus to the late summer and early autumn of 2025. While the Sentinel-1 catalogue provides a continuous data record spanning several years, applying automated detection across previous years, which consistently featured more extensive summer fast ice, would yield a dataset dominated by highly uncertain, fast-ice-induced stationary targets. The unprecedented, record low fast ice extent in early 2025 (Abram et al., 2025) – with less than $100,000 \text{ km}^2$ compared to the 2000-2018 baseline of $\sim 221,000 \text{ km}^2$ (Fraser et al., 2021) – offers a uniquely clear observational window to minimise “false positives” caused by fast ice immobilisation, and isolate truly grounded icebergs in open water. Furthermore, because macroscopic grounding patterns are predominantly dictated by relatively static physical constraints, such as regional bathymetry, ocean currents, and upstream calving sources (Massom et al., 2001; Tournadre et al., 2016; Kostov et al., 2026), the overall spatial distribution and density hotspots of grounded icebergs is not expected to change substantially from year to year (Fraser et al., 2012; Bett et al., 2020). Thus, although individual icebergs may remobilise, their aggregate spatial distribution likely remains stable. This makes the 2025 snapshot the most appropriate choice for a representative baseline for the Antarctic coastal system, intentionally prioritising data reliability over noisy multi-year temporal coverage.

Capitalising on these unique conditions, this study constructs the first circum-Antarctic grounded iceberg dataset to enable automated identification and temporal monitoring. Based on Sentinel-1 SAR imagery, a ResUNet deep learning framework was developed to achieve automatic detection and precise segmentation of icebergs around Antarctica, successfully identifying icebergs ranging down to 0.016 km^2 (~~40×40 m~~corresponding to 10 pixels at 40 m pixel spacing) and extracting their area, position, and border. Multi-temporal image analysis combined with feature-based matching algorithms is then applied to

identify grounded icebergs and monitor their temporal stability. The resulting dataset provides a robust foundation for understanding Antarctic coastal ice–ocean interactions and their impacts on polar climate and ecosystems, offering key data support for investigating the dynamic responses of grounded icebergs to fast ice stability, ice-shelf melting, and ecological processes.

2 Data

Analysis-ready Sentinel-1 SAR data were provided by Geoscience Australia under the Digital Earth Antarctica Program. 197 Sentinel-1 Ground Range Detected (~~GRD~~) ~~scenes~~ Medium Resolution (GRDM) scenes in the Extra Wide swath (EW) mode and 77 Single Look Complex (SLC) scenes in the Interferometric Wide swath (IW) mode were obtained from the Copernicus Data Space Ecosystem (CDSE) ~~were and~~ processed to a normalised, radiometrically terrain-corrected (RTC) gamma-nought backscatter product. EW GRDM data was processed using the pyroSAR (Truckenbrodt et al., 2019) and GAMMA (GAMMA Remote Sensing AG, n.d.) software packages. IW SLC data was processed using the Geoscience Australia sar-pipeline software that implements the NASA JPL RTC (OPERA Jet Propulsion Laboratory, 2026a) and ISCE3 (ISCE3 Development Team, 2026) software packages. Processing used the Copernicus 30 *m* Digital Elevation Model (DEM), with ellipsoidal heights applied for offshore data. The data were produced in decibel scaling at 40 *m* ~~spatial resolution for Extra-Wide-swath mode (EW)~~ ~~pixel spacing for EW~~ data and 20 *m* ~~spatial resolution for Interferometric Wide-swath mode (IW)~~ ~~pixel spacing for IW~~ data, and reprojected to polar stereographic coordinates (EPSG:3031) for analysis. ~~Default range and azimuth multilooking parameters defined by pyroSAR were used.~~ For EW data, thermal noise removal was applied and GRD border noise was removed from the input data using the pyroSAR method. The bicubic spline back-geocoding method was applied (Magnard et al., 2017). No additional multilooking was applied, maintaining the estimated number of looks (ENL) of the GRDM product of 10.7 (OPERA Project Team, 2024). For IW data, thermal noise removal, bistatic and static-tropospheric atmospheric delay corrections, radiometric terrain correction and geocoding was undertaken using the process described in the Product Specification Document for the OPERA Radiometric Terrain Corrected SAR Backscatter from Sentinel-1 (OPERA Jet Propulsion Laboratory, 2026b). To achieve the desired 20 *m* ground resolution, this workflow applies adaptive multi-looking. Unlike traditional fixed-window methods, this adaptive approach yields a spatially varying ENL across the scene based on local acquisition geometry and terrain (OPERA Project Team, 2024). Consequently, the scene-level ENL (approximately ~ 16 , estimated from resampling the nominal IW SLC $\sim 2.3 \times 14.1$ *m* pixel spacing to 20×20 *m*) should be understood as an approximate value rather than a fixed property of the product (OPERA Jet Propulsion Laboratory, 2026b)

In summary, the final gamma-nought intensity images obtained after pre-processing exhibit comparable visual characteristics for both EW and IW acquisition modes. However, IW images feature half the pixel spacing and slightly higher ENL compared to EW images. These differences stem from the intrinsic properties of the respective acquisition modes, where EW prioritises broader spatial coverage at the expense of spatial and/or radiometric resolution.

As the main aim of this study is to construct a circum-Antarctic grounded iceberg dataset, complete data coverage of the Antarctic coast is required. Therefore, EW mode (swath width approx. 410 *km*) data were primarily used where available, with

IW mode (swath width approx. 250 *km*) data used to fill gaps in the EW coverage. In addition, we found that in some SAR data, icebergs were more easily distinguished from sea ice in cross-polarisation (HV or VH) due to higher backscatter contrast, whereas in co-polarisation (HH) the backscatter from icebergs and sea ice can be similar, making it difficult for the model to differentiate them. Accordingly, this study primarily used cross-polarisation images for analysis where available. However, considering the limited spatial coverage of cross-polarisation data around Antarctica, co-polarisation images were used to fill the gaps in certain regions (Figure 1). To rigorously distinguish truly grounded icebergs from those immobilised by the fast sea ice matrix, we additionally employed a high-resolution map of the circum-Antarctic fast ice extent for early-mid March 2025 (i.e., the time of the climatological fast ice minimum; Abram et al., 2025) as a temporally static spatial mask.

In addition to Sentinel-1 SAR data, this study incorporated multiple supplementary datasets to support iceberg detection and validation. Topographic information was derived from the International Bathymetric Chart of the Southern Ocean (IBCSO; Dorschel et al., 2022) Version 2. Daily sea ice concentration (SIC) data were obtained from the European Organisation for the Exploitation of Meteorological Satellites Ocean and Sea Ice Satellite Application Facility (OSI SAF) Global Sea Ice Concentration product (OSI-408-a), derived from the Advanced Microwave Scanning Radiometer 2 (AMSR2) with a grid resolution of 10 *km* (EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF), 2025). Coastline boundaries were referenced from the high-resolution vector data of the Antarctic Digital Database (ADD, Version 7.11; Gerrish et al., 2025) maintained by the Scientific Committee on Antarctic Research (SCAR). Furthermore, optical imagery from the Sentinel-2 MultiSpectral Instrument (MSI; European Space Agency, 2015) was acquired to assist with manual annotation, ground-truth generation, and accuracy assessment. All supplementary datasets provided complete circum-Antarctic coverage, with the temporal window focused on late February to early April 2025, corresponding to the annual minimum of the Antarctic fast ice extent (Fraser et al., 2021), in the year of the lowest summertime fast ice extent on record (Abram et al., 2025).

3 Methods

This section outlines the novel approach for producing the grounded iceberg dataset, and is structured into five main sections: (3.1) data preparation, (3.2) iceberg detection, (3.3) grounded iceberg identification, (3.4) physical constraint filtering, and (3.5) performance validation. The workflow is illustrated in Figure 2. Boxes I through IV constitute the core production pipeline, progressing from data input to the generation of the final grounded iceberg dataset, while Box V details the accuracy assessment conducted on an independent validation set.

3.1 Data preparation and training dataset construction

3.1.1 Generation of training samples

For training U-Net-based models, iceberg regions were manually annotated to create the training dataset. The images were then divided into sub-images of approximately 250×250 pixels each. The exact tile dimensions were dynamically adjusted to ensure integer divisibility, thereby eliminating residual edge strips. Finally, sub-images were upsampled to 1536×1536 pixels




Figure 2. Technical workflow of the grounded iceberg detection and identification system. The pipeline processes Sentinel-1 imagery through deep learning segmentation, individual identification, and physical constraint filtering to generate the final circum-Antarctic grounded iceberg dataset.

using bilinear interpolation, as empirical tests indicated this resolution enhances boundary segmentation accuracy. From a total of 15 SAR images (yielding 4,220 sub-images), 702 sub-images containing icebergs under various environmental conditions were manually selected. The iceberg regions were then manually annotated using the Python LabelMe library. The accuracy of the manually drawn masks has been reported to be within 2 %–4 % of the iceberg area (Braakmann-Folgmann et al., 2022), and were verified by at least one study co-author. We generated binary mask images of the same size as the original images from the annotation files, defining pixels within the annotated regions as iceberg and all other pixels as background, as shown in Figure 3.

3.1.2 Model training dataset

For the model training dataset, nine coastal regions of Antarctica were selected to cover a diverse range of environmental conditions (Figure 1). The data were processed using the method described in Section 3.1.1 to construct the training dataset for iceberg detection. This training dataset covers data from February to March in 2021, 2023, and 2025, comprising a diverse mix of imagery from IW and EW swath modes, and spanning both HV and HH polarisations. These specific years were selected to capture a broad range of inter-annual variability in sea ice conditions and iceberg concentrations over a five-year period, ensuring the model’s robustness across different environmental regimes. In terms of target selection, the dataset primarily focuses on icebergs smaller than 100 km^2 (micro to medium-sized), maximising sensitivity for these challenging, under-




Figure 3. Cropped examples ($10 \times 10 \text{ km}$) from input SAR images acquired in Sentinel-1 EW mode with HV polarisation (top row) and corresponding binary masks generated from manual annotations (bottom row).

represented targets. To enable robust generalisation across these differing signal characteristics, a single ResUNet model was trained on this diverse dataset. This strategy allows the network to learn feature representations applicable to both polarisations, thereby mitigating the detection challenges associated with the lower contrast between icebergs and sea ice inherent in HH data. The final dataset was manually divided into a training set for model learning (618 sub-images, 88 %) and a test set (84 sub-images, 12 %) for assessing training performance. This internal assessment is distinct from the final performance validation, which is conducted on an independent validation dataset.

3.1.3 SAR Imagery for grounded iceberg dataset

To construct the circum-Antarctic grounded iceberg dataset, we acquired Sentinel-1 SAR imagery of the entire Antarctic continental shelf coastal margin primarily from March 2025, with the temporal window extended to late February and early April to ensure complete spatial coverage (Figure 1). This observation period was chosen to coincide with the timing of annual minimum of Antarctic fast ice extent (Fraser et al., 2021). Crucially, the record low fast ice extent observed in early 2025 minimises the risk of misidentifying temporarily immobilised floating icebergs as grounded (Abram et al., 2025). For each orbital track, repeat-pass images with temporal baselines of 12 days (or multiples due to data availability) were analysed. All Antarctic coastal regions included in the dataset are covered by at least two consecutive valid SAR acquisitions, ensuring that no regions were excluded from the stationarity evaluation due to a lack of repeat-pass data. Because our tracking framework analyses these repeat-pass images along the same track, the image pairs used for cross-temporal matching consist of the same acquisition mode and polarisation. Furthermore, continuous image sequences ($n \geq 2$) are routinely utilised to assess target stationarity. The dataset comprises a total of 274 Sentinel-1 SAR images, of which 197 were acquired in EW mode and 77

in IW mode. The total cumulative area of the acquired imagery is 9,983,860 km², corresponding to a unique footprint (union area) of 7,456,633 km² and an overlapping area of 2,527,226 km².

3.2 Iceberg detection

230 This study employs a modified U-Net model (Ronneberger et al., 2015), in which the encoder is replaced by a ResNet152 network (He et al., 2016), forming a ResUNet architecture (Diakogiannis et al., 2020) to enhance the extraction of iceberg features from SAR images. The model takes 1536×1536 single-channel backscatter sub-images as input (as described in Section 3.1.1 and shown in Figure 3), with corresponding manually-annotated masks of the same size indicating iceberg regions and background for training.

3.2.1 Model architecture

235 To enhance the model's ability to extract features from complex Antarctic coastal environmental backgrounds, this study employs ResNet152 (He et al., 2016) as the encoder and incorporates the Convolutional Block Attention Module (CBAM; Woo et al., 2018) to strengthen attention in both the channel dimension (identifying which features are meaningful) and the spatial dimension (localising where target objects are). The model consists of an encoder, decoder, skip connections, and embedded CBAM (Figure 4). The encoder uses an ImageNet-pretrained (Deng et al., 2009) ResNet152 to extract multi-scale
240 iceberg features. To accommodate single-channel input, the first convolutional layer is modified to accept single-channel data, and the first four convolutional blocks (convolution, batch normalisation, Rectified Linear Unit (ReLU), and max pooling) are frozen to preserve low-level feature stability, thereby enhancing the model's robustness in capturing iceberg edges and textures. CBAMs are placed at five levels within the encoder to filter the features. By enhancing the contrast between important targets and the environment, these modules help the model highlight icebergs while ignoring confusing background details like sea ice.
245 The decoder progressively restores spatial resolution through upsampling (nearest-neighbour interpolation) and convolutional operations, fusing features with the corresponding encoder levels. Finally, a 1×1 convolution maps the multi-channel feature representations onto a single-channel probability map, representing the likelihood of each pixel belonging to either iceberg or background, and a probability threshold of 0.5 is applied to binarise the map and generate the iceberg mask.

3.2.2 Model training and optimisation

250 To determine the optimal training parameters, Bayesian optimisation (Snoek et al., 2012) was employed to search hyperparameters, including learning rate, batch size, and weight decay. The selected training parameters were as follows: learning rate of 3×10^{-4} , batch size of 4, and weight decay coefficient of 1×10^{-4} , balancing model convergence speed and segmentation accuracy. In total, approximately 50 independent training runs were executed during the tuning phase and final training runs using the selected optimal configuration.




Figure 4. Schematic diagram of the proposed CBAM-ResUNet architecture. The network follows a standard encoder-decoder structure with a ResNet-152 backbone. The encoder path (left, blue nodes) extracts hierarchical features from the single-channel input. To enhance feature representation, Convolutional Block Attention Modules (CBAM, orange nodes) are integrated at the bottleneck and skip connections (dashed lines), applying both channel and spatial attention mechanisms to refine feature maps before fusion. The decoder path (right, green nodes) progressively recovers spatial resolution through upsampling and convolution operations to generate the final binary output mask. Acronyms: ResNet (Residual Neural Network); CBAM (Convolutional Block Attention Module); BN (Batch Normalization); ReLU (Rectified Linear Unit); Conv (Convolution); 1ch (1 Channel).

Iceberg pixels occupy only a small proportion of each image, and the distribution of positive and negative regions is highly imbalanced. To address this, the study employed a combined loss function (L_{Total}), which linearly integrates weighted Binary Cross-Entropy (L_{BCE}) and Tversky loss ($L_{Tversky}$) (Salehi et al., 2017). The weighted BCE loss is defined as:

$$L_{BCE} = -\frac{1}{N} \sum_{i=1}^N [w \cdot y_i \cdot \log(\sigma(x_i)) + (1 - y_i) \cdot \log(1 - \sigma(x_i))], \quad (1)$$

where N is the total number of pixels, y_i is the ground truth pixel value (0 for background, 1 for iceberg), $\sigma(x_i)$ is the sigmoid activation of the model output logits x_i , and w represents the positive sample weight. To further mitigate the impact of class imbalance and reduce false negatives, the Tversky loss is calculated based on the Tversky Index:

$$\text{Tversky Index} = \frac{TP + \epsilon}{TP + \alpha \cdot FP + \beta \cdot FN + \epsilon}, \quad (2)$$

$$L_{Tversky} = 1 - \text{Tversky Index}, \quad (3)$$

where TP , FP , and FN represent pixel-level true positive, false positive, and false negative counts, respectively. The parameters $\alpha = 0.6$ and $\beta = 0.4$ are set to impose a heavier penalty on false positives compared to false negatives, while ϵ is a small constant (set to 1×10^{-6}) added to ensure numerical stability and avoid division by zero. The final combined loss function is expressed as:

$$L_{Total} = 0.4 \cdot L_{BCE} + 0.6 \cdot L_{Tversky}, \quad (4)$$

where the weights 0.4 and 0.6 balance the contributions of each loss component to maintain segmentation accuracy while effectively reducing target omission.

Model training was conducted in a multi-GPU parallel environment using the Distributed Data Parallel (DDP) framework to improve training efficiency, with a gradient accumulation step of 2 to achieve a larger effective batch size. The AdamW optimiser (Loshchilov and Hutter, 2017) was used in combination with a cosine annealing learning rate scheduler, completing a full cycle every 20 epochs, with a minimum learning rate of 1×10^{-6} . To further stabilise training, automatic mixed precision (AMP; Micikevicius et al., 2017) and gradient scaling were applied to optimise computational efficiency. An early stopping mechanism was implemented, terminating training when the F1 score on the test set did not improve for 10 consecutive epochs, and the best-performing model was saved. In practice, each model typically required approximately 20–30 epochs to reach convergence.

In terms of computing resources, training was conducted on four NVIDIA Tesla V100 32GB GPUs within the Australian National Computational Infrastructure (NCI) Gadi supercomputer. Each training session took approximately half an hour.

3.3 Grounded iceberg identification

In this study, stationary icebergs are detected through a hierarchical filtering framework that integrates spatiotemporal identification with environmental constraints. We initially define a stationary iceberg candidate as a feature that persists at a fixed location for a minimum duration of 12 days. However, stationarity alone does not imply grounding, as icebergs may also be immobilised by fast ice. To operationalise this definition, we developed a multi-stage grounded iceberg identification pipeline, comprising three key phases: (1) Individual iceberg extraction ([Section 3.3.1](#)), which isolates distinct iceberg entities from segmentation masks; (2) ~~Trajectory-reconstruction~~ [Stationary target identification \(Section 3.3.2\)](#), which establishes temporal continuity based on geometric similarity and motion constraints defined by a maximum centroid displacement of 10 pixels; and (3) Quality control, which mitigates false positives using bathymetric, surrounding sea ice concentration criteria. Only candidates that survive this rigorous screening process are classified as stationary icebergs. We then use a March 2025 map of fast ice extent (Abram et al., 2025) to suggest which stationary icebergs are “high-confidence grounded” vs those possibly rendered stationary due to enclosure by fast ice.

3.3.1 Individual iceberg extraction

Since the binary masks predicted by the U-Net model provide only pixel-level classification without differentiating between individual iceberg instances, they cannot inherently distinguish individual icebergs. To address this, we implemented a block-based parallel processing method for connected component extraction. Given the wide coverage of SAR EW and IW modes and the density of small icebergs, masks are partitioned into multiple blocks to optimise processing. Each block undergoes independent connected component analysis using OpenCV (Bradski, 2000) to generate local labels and compute geometric properties (e.g., area, centroids, and bounding box), while regions smaller than 10 pixels are discarded as noise. Among these properties, the axis ratio is estimated through ellipse fitting to describe each iceberg's elongation:

$$\text{Axis Ratio} = \frac{a}{b}, \quad (5)$$

where a and b are the lengths of the major and minor axes, respectively. To mitigate pixel-level quantisation errors and improve the reliability of subsequent shape-based matching, the extracted contours are smoothed using a moving average convolution:

$$p'_i = \sum_{j=-k}^k p_{i+j} \cdot \frac{1}{2k+1}, \quad (6)$$

where p_i is the original coordinate (x_i or y_i) of the i -th contour point, p'_i is the resulting smoothed coordinate, and k is half of the kernel size used for the smoothing convolution. Simultaneously, to eliminate false positives arising from glacier features, we incorporated the SCAR ADD high-resolution Antarctic coastline polygons (Gerrish et al., 2025); any candidate region overlapping with these land masks is automatically excluded.

To resolve the splitting of connected regions across block boundaries, a Union-Find algorithm (Galler and Fisher, 1964) is applied to merge boundary labels. Subsequently, each connected region is mapped to a global label with a unique identifier, ensuring consistent representation across blocks. Ultimately, the workflow generates single-channel label masks for each individual iceberg, accompanied by a detailed attribute dataset (including area, centroid, and ID), providing the necessary data foundation for grounded iceberg identification and analysis.

3.3.2 Grounded iceberg identification algorithm

In this study we develop an algorithm for identifying grounded icebergs in SAR imagery, aimed at monitoring grounded icebergs using the multi-temporal mask data and region information obtained in Section 3.3.1. The method integrates block-based processing, feature extraction, cost matrix optimisation, and trajectory management, making it suitable for large-scale analysis of grounded icebergs in SAR data. The algorithm is structured into three main components: (1) data input, (2) processing workflow, and (3) output results.

In the data input stage, the algorithm primarily takes iceberg region information files and the corresponding single-channel mask files as inputs. The process begins with region loading and tile-based partitioning. Imagery and geographic information

data, including the coordinate reference system and affine transformation matrix, are read using the Rasterio Python library (Gillies, Sean and others, 2013), and the images are divided into tiles. Each region is then assigned to the corresponding tile based on its centroid location. The algorithm then extracts multiple features for each region, including the binary region and its properties from the mask (Section 3.3.1), contours and smoothed contours derived using OpenCV, and the major-to-minor axis ratio through ellipse fitting, providing a comprehensive description of each iceberg’s shape and characteristics.

The grounded iceberg spatiotemporal association component achieves trajectory continuation through cross-temporal region matching. The method employs the Hungarian algorithm (Kuhn, 1955) for cost matrix optimisation. The assignment problem is formulated to find the optimal matching that minimises the total cost:

$$\min \sum_{i,j} C_{ij} \cdot x_{ij} \quad (7)$$

subject to:

$$\sum_j x_{ij} = 1, \quad \sum_i x_{ij} = 1, \quad x_{ij} \in \{0, 1\}, \quad (8)$$

where C_{ij} is the element of the cost matrix representing the matching cost between region i and trajectory j , and x_{ij} is a binary variable indicating whether region i is assigned to trajectory j . For regions within the same tile across consecutive temporal phases, the cost matrix C_{ij} was constructed incorporating multiple metrics:

$$C_{ij} = w_1 \cdot \text{norm_dist}(c_i, c_j) + w_2 \cdot \text{norm_area}(M_i, M_j) + w_3 \cdot \text{norm_axis}(R_i, R_j, \theta_i, \theta_j) + w_4 \cdot \text{norm_contour}(S_i, S_j), \quad (9)$$

where w_1, w_2, w_3, w_4 are weights assigned to each metric ($w_1 = 0.05, w_2 = 0.30, w_3 = 0.60, w_4 = 0.05$). The components include the normalised centroid Euclidean distance, region-overlap cost, axis ratio, and contour-shape similarity based on Hu invariant moments (Hu, 1962). The region-overlap cost is calculated using the Intersection over Union (IoU):

$$\text{IoU}(M_1, M_2) = \frac{|M_1 \cap M_2|}{|M_1 \cup M_2|}, \quad (10)$$

where M_1 and M_2 are the binary mask regions at two consecutive timestamps, $|M_1 \cap M_2|$ is the number of overlapping pixels, and $|M_1 \cup M_2|$ is the number of pixels in the union. Hu moments, being invariant to translation, rotation, and scaling, enhance the robustness of the matching algorithm by quantifying shape similarity through the logarithmic differences of seven invariant moments.

To improve matching accuracy, the algorithm first excludes region pairs with excessive spatial separation using a centroid distance threshold (10 pixels), followed by filtering based on area difference (threshold 0.5) and IoU threshold (0.99, i.e., $\text{IoU} < 0.01$) to eliminate anomalous matches. If the total cost is below the threshold of 10^6 and represented the optimal

solution within the matchable range, the regions are deemed to correspond to the same iceberg across different times, and the trajectory information is updated; if no match is found, a new trajectory is initiated. If a grounded iceberg is missed in a single timestamp due to detection errors or merging with nearby icebergs but reappears in subsequent frames, the algorithm automatically reconnects its trajectory.

The identification results are exported as a tabular data file, with each trajectory record containing information such as timestamp, centroid, area, unique identifier, and matching cost. This data file, combined with multi-temporal mask data, forms the grounded iceberg dataset, the structure of which is detailed in Table 1.

Table 1. Structure and description of the circum-Antarctic grounded iceberg dataset attributes. UID stands for Unique ID, EW stands for Extra Wide swath mode, IW stands for Interferometric Wide swath mode, and WKT stands for Well-Known Text.

Field Name	Data Type	Unit	Description
Global_UID	String	-	Unique identifier (Orbit_TrackID, e.g., 045_113)
Orbit	String	-	Sentinel-1 relative orbit number (e.g., 45)
Timestamp	String	-	Polygon image acquisition time (ISO 8601, e.g., 2025-03-08T19:43:27Z)
Date_Range	String	-	Temporal window for verification (e.g., 2025-02-24T...Z - 2025-03-08T...Z)
Area_Mean_km2	Float	km^2	Mean surface area (e.g., 9.73)
Area_Range_km2	String	km^2	Min-Max area range observed (e.g., 9.39 - 10.07)
Area_Mean_px	Float	pixels	Mean area in pixels (e.g., 2450.5)
Area_Range_px	String	pixels	Min-Max pixel count range (e.g., 2350 - 2500)
Bed_Depth	Float	m	Seafloor depth from IBCSO v2 (e.g., -251.0)
Acquisition_Mode	String	-	Acquisition mode (EW/IW)
Latitude	Float	$^{\circ}$	Centroid latitude in WGS84 (e.g., -69.402)
Longitude	Float	$^{\circ}$	Centroid longitude in WGS84 (e.g., 14.584)
Northing_3031	Float	m	Projected Y coordinate in EPSG:3031 (e.g., 2188596.0)
Easting_3031	Float	m	Projected X coordinate in EPSG:3031 (e.g., 569429.4)
Fast_Ice_Overlap_Status	String	-	Spatial relationship with fast ice (Inside/Partial/Outside)
Geometry	Polygon	-	Vector polygon of iceberg boundary (WKT format)

3.4 Physical constraint filtering

Although the stationarity-based identification algorithm (Section 3.3.2) effectively distinguishes stationary from drifting targets, two main types of false positives can still occur in regions of high ice concentration or deep water: (1) floating icebergs temporarily immobilised by surrounding sea ice (especially fast ice, which can arrest iceberg advection; e.g., as shown in Figure 6 of (Fraser et al., 2010)), and (2) strong backscatter signals from dense sea ice rubble or ice deformation. To ensure the physical plausibility of the grounded iceberg dataset, we applied post-processing filters incorporating bathymetry and SIC data.

Identified iceberg trajectories were spatially matched with IBCSO bathymetry (Dorschel et al., 2022) and spatiotemporally aligned with daily AMSR2 SIC data (EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF), 2025). Based on Antarctic grounded iceberg characteristics (Massom et al., 2001; Dowdeswell and Bamber, 2007), (Massom et al., 2001; Dowdeswell and
 365 , as noted in Section 1), three hierarchical filtering criteria were defined (Table 2). Trajectories meeting any exclusion criterion at any time step were flagged as invalid and removed from the final dataset. This procedure markedly reduced false positives under complex sea ice conditions, ensuring high geographical and physical confidence in the resulting dataset. The quantitative impact of each filtering stage on the final grounded iceberg dataset is detailed in Section 4.2.1.

Table 2. Parameters for post-processing quality control filters. SIC: Sea Ice Concentration.

ID	Conditions	Thresholds	Physical rationale
1	Water Depth, SIC, & Area	Water Depth > 600 <i>m</i> SIC > 40 % Area < 1 <i>km</i> ²	Excludes floating icebergs temporarily immobilised by sea ice.
2	SIC & Area	SIC > 80 % Area < 0.1 <i>km</i> ²	Removes high backscatter features caused by sea ice rubble.
3	Water Depth	Water Depth > 1000 <i>m</i>	Eliminates targets in physically impossible ground- ing zones.

Preliminary visual inspection indicated that the dataset from orbit 54 (~146°E) exhibited anomalous textural characteristics, likely attributable to complex sea ice conditions. The presence of heavily deformed sea ice and surface roughness resulted in high backscatter signals that resembled the signature of icebergs. This environmental complexity led to widespread false positives, manifesting as small, fragmented artefacts. Furthermore, this region was not used in training the deep learning model. To mitigate this spatial heterogeneity, we implemented an ad-hoc quality control filtering strategy, applying a more stringent area threshold exclusively to this orbit: all tracked icebergs containing any individual observation smaller than 200 pixels
 375 (approximately 0.08 *km*²) were discarded. This targeted intervention effectively suppressed persistent noise artefacts in this orbit, and ensured that the reliability of the extracted trajectories in this region was consistent with that of the remainder of the global dataset.

To distinguish between truly grounded icebergs and floating icebergs immobilised by landfast sea ice, we performed a spatial overlay analysis using a high-resolution fast ice mask digitised from the same time period as the SAR data used in this
 380 study ((Abram et al., 2025))(Abram et al., 2025). Fast ice digitisation was manually achieved using a high-resolution cloud-free visible image composite, as described by (Fraser et al., 2009). Each stationary iceberg candidate was then classified based on its spatial relationship with this mask:

1. Inside: The stationary iceberg geometry is completely contained within the fast ice boundary, indicating that it may not be grounded.

- 385 2. Partial: The stationary iceberg geometry intersects the fast ice boundary but is not fully contained. This often occurs on the “downstream” side (typically to the west) of persistent fast ice features, and often indicates genuinely grounded icebergs which support fast ice growth and stability on the “upstream” side (typically to the east) (Fraser et al., 2012).
3. Outside: The stationary iceberg has no spatial overlap with the fast ice mask. Targets classified as ‘Outside’ are identified as high-confidence grounded icebergs, as no fast ice is present to inhibit their motion.

390 3.5 Validation

3.5.1 Iceberg detection model performance

To evaluate the performance of the iceberg detection model, we conducted an independent validation of the ResUNet model using a 50×50 km region in Adélie Land ($\sim 136^\circ\text{E}$) as the validation dataset. We manually interpreted and annotated 12 images, acquired in EW swath mode with cross-polarisation (HV) at monthly intervals from January 2021 to December 2021, 395 thereby establishing the ground-truth dataset for the icebergs. We chose to perform a year-round validation, even though the training dataset is only selected from the months of February and March. This allows us to assess the generalisation of our proposed method across different seasons and possible limitations caused by surface melt during the summer. The validation set underwent no data augmentation and was not used during model training, ensuring that the evaluation objectively reflects the model’s performance and generalisation ability. To ensure the reliability of these manual annotations, Sentinel-2 MSI optical 400 imagery was utilised as an auxiliary reference for visual verification (e.g., to allow manual separation of small icebergs from similarly-shaped small sea ice floes).

To ensure the model’s robustness and representativeness at the circum-Antarctic scale, we constructed an extended validation dataset comprising three additional independent observational scenes acquired between late February and mid-April 2025. These scenes were purposefully selected to cover all combinations of Sentinel-1 acquisition modes (EW and IW) and available 405 polarisations (HH and VH) used in this study, enabling an assessment of how differing radar configurations affect single-frame detection performance.

During validation, we computed standard pixel-wise performance metrics, including producer’s accuracy (recall), user’s accuracy (precision), and F1 score (the harmonic mean of precision and recall). Based on the comparison between the predicted and ground-truth masks, true positives (TP), true negatives (TN), false positives (FP), and false negatives (FN) were recorded. 410 These confusion categories are defined as follows: TP represents pixels correctly predicted as iceberg; TN represents pixels correctly predicted as background; FP represents pixels incorrectly predicted as iceberg; and FN represents pixels incorrectly predicted as background. These counts were then used to calculate the core performance metrics:

$$\text{Precision} = \frac{TP}{TP + FP + \epsilon}, \quad (11)$$

$$\text{Recall} = \frac{TP}{TP + FN + \epsilon}, \quad (12)$$

$$415 \quad F1 = \frac{2 \cdot \text{Precision} \cdot \text{Recall}}{\text{Precision} + \text{Recall} + \epsilon}, \text{ and} \quad (13)$$

$$\text{FAR} = \frac{FP}{FP + TN + \epsilon}, \quad (14)$$

where ϵ is a small constant (set to 1×10^{-6}) added to ensure numerical stability and avoid division by zero. Precision measures the proportion of predicted ‘iceberg pixels’ that are truly icebergs, reflecting the reliability of the model’s predictions; recall measures the proportion of all actual iceberg pixels that are successfully detected by the model, indicating its tendency
 420 to miss targets; and False Alarm Rate (FAR) measures the proportion of background pixels incorrectly classified as iceberg pixels, reflecting the model’s false detection rate.

3.5.2 Grounded iceberg identification

We also used the same $50 \times 50 \text{ km}$ area in Adélie Land, as described in Section 3.5.1, as the validation scene for the grounded iceberg identification algorithm. The evaluation employed a frame-based statistical approach, in which the identification results
 425 were manually classified. The specific evaluation metrics are defined as follows:

1. TP: the number of frames in which the algorithm successfully identified the grounded iceberg. That is, at a given time, the algorithm assigns the iceberg to the correct trajectory.
2. FN: the number of frames in which a grounded iceberg is actually present but the algorithm fails to generate a valid tracking result, also referred to as ‘missing frames’. This reflects instances where the algorithm loses the target in the
 430 time series.
3. FP: the number of frames in which the algorithm incorrectly generates a tracking result. This includes cases where background elements (such as sea ice or waves) are misidentified as grounded icebergs, or incorrectly classifies icebergs from different times as belonging to the same grounded iceberg.

Based on the above statistics, we selected Precision, Recall and F1 Score as the core evaluation metrics.

435 4 Results

Following the classification scheme of Wesche and Dierking (2015), the detected icebergs were divided into six categories based on their area: micro icebergs (A0, $< 0.1 \text{ km}^2$), tiny icebergs (A1, $0.1 - 1 \text{ km}^2$), small icebergs (A2, $1 - 10 \text{ km}^2$), medium icebergs (A3, $10 - 100 \text{ km}^2$), large icebergs (A4, $100 - 1,000 \text{ km}^2$), and giant icebergs (A5, $\geq 1,000 \text{ km}^2$). This classification enables effective differentiation of spatial distribution patterns across icebergs of varying sizes. However, it is important to note
 440 that no grounded icebergs belonging to the large (A4) or giant (A5) categories were identified in this dataset. This aligns with

our research focus on resolving the ‘missing’ population of smaller icebergs ($< 100 \text{ km}^2$), complementing the comprehensive records of large targets maintained by operational agencies (e.g., U.S. National Ice Center; U.S. National Ice Center, 2024). Consequently, our training strategy prioritised micro to medium icebergs (A0–A3) to ensure maximum sensitivity for these challenging, under-represented targets.

445 **4.1 System performance evaluation**

Using the independent validation set from Adélie Land described in Section 3.5, we conducted a systematic evaluation of the ResUNet model and the iceberg identification algorithm. The complex environmental characteristics of this region, which present substantial background clutter and variability, provide a suitable setting for assessing the robustness of the model in pixel-level detection and temporal tracking.

450 **4.1.1 Iceberg detection performance**

In terms of iceberg detection, the ResUNet model demonstrated strong and reliable performance on both the validation and test datasets. The model stopped training on the manually-partitioned test set with an F1 score of 0.8926, a precision of 0.8963, a recall of 0.9002, and a FAR of 0.0058. Its performance on the $50 \times 50 \text{ km}$ validation region in Adélie Land was similarly strong, achieving an average precision of 0.9145, a recall of 0.9131, an F1 score of 0.9121, and a false alarm rate of 0.0056
455 across twelve months. Table 3 presents the model’s performance metrics across selected images from different months.

Table 3. Monthly test metrics for the iceberg detection model. FPR: false positive rate.

Date	Precision	Recall	FPR	F1 Score
2021-01-09	0.9609	0.8652	0.0021	0.9106
2021-02-14	0.9065	0.8720	0.0044	0.8889
2021-03-10	0.8526	0.9368	0.0080	0.8927
2021-04-15	0.8763	0.9571	0.0089	0.9149
2021-04-27	0.9154	0.9426	0.0058	0.9288
2021-06-02	0.9058	0.9411	0.0065	0.9232
2021-07-03	0.9019	0.9507	0.0070	0.9256
2021-08-11	0.9377	0.9150	0.0043	0.9262
2021-09-18	0.9130	0.9447	0.0062	0.9286
2021-10-12	0.9171	0.9444	0.0059	0.9305
2021-11-17	0.9228	0.9325	0.0055	0.9276
2021-12-11	0.9646	0.7553	0.0014	0.8472

These metrics indicate that the model is capable of accurately and consistently detecting iceberg pixels, while maintaining a low rate of background pixels erroneously classified as icebergs. The statistical results presented in Table 3 further confirm the model's cross-seasonal generalisation. With the exception of December the model maintained a high level of detection stability throughout most of the year. The reduced model performance in December is likely caused by changes in backscatter resulting from surface melt and moisture in the snow cover, which is confirmed by visual inspection of example images from multiple regions. However, recall that December is not used for the generation of the final grounded iceberg dataset presented here. Except for these seasonal effects, the model handles changes in sea ice coverage and complex sea states effectively, demonstrating strong robustness in Antarctica's dynamic coastal environment. Figure 5 illustrates representative test samples, including the raw SAR imagery, Sentinel-2 optical imagery, manually annotated ground truth, and the model's predicted masks.

Visual inspection shows a high degree of agreement between the predicted iceberg boundaries and the ground truth. Notably, for icebergs with irregular edge morphology, the model produces smooth and well-fitted contours, demonstrating its effectiveness in preserving geometric detail.




Figure 5. Visual comparison of the iceberg ground truth mask and the predicted mask on SAR imagery ($\sim 136^\circ\text{E}$). (A) Sentinel-2 optical image acquired on 2021-04-06, shown as a false-colour composite using bands B08 (near-infrared), B04 (red), and B03 (green); (B) Sentinel-1 HV intensity image acquired on 2021-04-15; (C) iceberg ground truth mask obtained through manual annotation based on panel (B); (D) predicted mask generated by the deep learning model from the same Sentinel-1 image. In mask images (C) and (D), white regions represent the identified iceberg bodies, and black regions represent the background ocean or sea ice. Panels (B), (C), and (D) have exactly the same acquisition time (2021-04-15) and geographical extent, while panel (A) serves as a contemporaneous visible reference for validation.

Given the numerical dominance of tiny icebergs ($< 1 \text{ km}^2$) in the Antarctic coastal system, we further conducted a size-stratified performance evaluation. Using a threshold of 1 km^2 , we classified the ground truth targets into ‘small’ ($< 1 \text{ km}^2$) and ‘large’ ($\geq 1 \text{ km}^2$) categories. As detailed in Table 4, the model exhibits robust detection capabilities across both size classes. For large icebergs, the model achieved a high Recall of 0.9397, demonstrating its effectiveness in capturing the main bodies of extensive ice shelves and tabular icebergs. Crucially, for the challenging small iceberg category, which is often prone to sea clutter interference, the model maintained a Precision of 0.8652 and an F1 score of 0.8613. These results indicate that the model's high overall performance is not solely driven by large objects but reflects a genuine capability to distinguish sub-kilometer icebergs from complex backgrounds.

Table 4. Size-stratified detection performance (small ($<1 \text{ km}^2$) vs. large ($\geq 1 \text{ km}^2$) icebergs)

Category	Area (km^2)	Precision	Recall	F1 Score
Small	< 1	0.8652	0.8630	0.8613
Large	≥ 1	0.9404	0.9397	0.9387
Overall	All	0.9145	0.9131	0.9121

The extended validation across different combinations of acquisition modes and polarisations (Table 5) further elucidates the model’s sensitivity to radar mode and polarisation. While the model exhibits consistently high performance in cross-polarised imagery (e.g., F1 scores of 0.880 and 0.891 for IW mode, VH polarisation), its single-frame pixel-level accuracy fluctuates in co-polarised (HH) imagery. Specifically, the scenes acquired on 2025-03-02 using EW mode with HH polarisation and on
480 2025-03-11 using IW mode with HH polarisation show substantial drops in Precision (0.4908 and 0.4528, respectively) and overall F1 scores (~ 0.61).

Table 5. Extended pixel-level detection performance across different acquisition modes and polarisations.

Date	Location	Longitude	Mode	Polarisation	Precision	Recall	FPR	F1 Score
2025-02-27	Princess Ragnhild Coast	$\sim 25^\circ \text{E}$	IW	HH	0.6872	0.9251	0.0030	0.7886
2025-03-02	Princess Martha Coast	$\sim 15^\circ \text{W}$	EW	HH	0.4908	0.8006	0.0034	0.6085
2025-03-11	Princess Ragnhild Coast	$\sim 25^\circ \text{E}$	IW	HH	0.4528	0.9378	0.0060	0.6108
2025-03-23	Princess Ragnhild Coast	$\sim 25^\circ \text{E}$	IW	HH	0.8659	0.9335	0.0008	0.8984
2025-03-30	Enderby Land	$\sim 44^\circ \text{E}$	IW	VH	0.9097	0.8524	0.0025	0.8801
2025-04-04	Princess Ragnhild Coast	$\sim 25^\circ \text{E}$	IW	HH	0.8641	0.9205	0.0008	0.8914
2025-04-11	Enderby Land	$\sim 44^\circ \text{E}$	IW	VH	0.9041	0.8773	0.0022	0.8905
2025-04-19	Princess Martha Coast	$\sim 15^\circ \text{W}$	EW	HH	0.8779	0.8401	0.0006	0.8586

Confusion matrix analysis reveals that this performance drop stems not from missed targets, as Recall remains robust (e.g., 0.9378 for the 2025-03-11 scene), but from a marked spike in false positive. This is because HH polarisation often yields similar backscatter signatures for icebergs, deformed sea ice, and wind-driven clutter, making single-frame differentiation
485 inherently difficult. Crucially, these single-frame errors do not detract from the final dataset. Most False Positives in HH-polarised images represent transient features, such as drifting floes or transient surface roughness. Being highly mobile, they fail the strict temporal continuity constraints of our tracking algorithm (Section 3.3.2). Therefore, while varying polarisations cause unavoidable single-frame segmentation fluctuations, rigorous multi-temporal tracking acts as a robust spatiotemporal filter, neutralising transient noise to ensure a high-confidence final product.

Building on the validated single-frame iceberg detection performance, we further evaluated the grounded iceberg identification algorithm using the temporal SAR imagery of the Adélie Land validation set. The evaluation employed frame-based statistical metrics, with a focus on the algorithm’s ability to maintain target identity consistency across the temporal sequence.

Quantitative results indicate that the algorithm exhibits high stability over multiple consecutive frames. Frame-level evaluation metrics show a precision of 0.9590, a recall of 0.8660, and an F1 score of 0.9102. The high precision demonstrates that the algorithm effectively avoids misclassifying background noise, such as transient disturbances from drifting sea ice, as new iceberg trajectories. Moreover, few erroneous matches occur within the generated trajectories. The relatively lower recall is primarily attributed to object merging in densely-populated regions. Although the algorithm is generally able to maintain stable tracking, in areas of high iceberg density, adjacent icebergs are prone to being misidentified as a single connected component. Such segmentation errors hinder the generation and maintenance of independent trajectories for individual icebergs, leading to intermittent tracking interruptions or failures.

The robustness of the algorithm is largely attributed to its multi-feature matching strategy based on the Hungarian algorithm. In terms of shape invariance, although icebergs may undergo pixel-level morphological variations due to changes in imaging angle or minor tilting at different observation times, the incorporation of Hu invariant moments (Hu, 1962) plays a critical role. These moments consist of seven statistical descriptors derived from central moments that characterise the global shape distribution and remain theoretically constant under translation, scaling, and rotation. Regarding trajectory continuity, the results presented in Figure 6 confirm the robustness of the matching algorithm. By jointly utilising mask contours and centroid features, the algorithm achieves reliable object association over the temporal sequence, producing clear and continuous iceberg trajectories and validating the reliability of the matching results. Furthermore, the thresholding mechanism based on IoU and centroid distance effectively removes aberrant drift trajectories of non-grounded icebergs, ensuring that the final grounded iceberg dataset attains a high level of confidence.

To quantify the temporal uncertainty of grounded iceberg identification, we evaluated start and end dates of period where the algorithm identified each iceberg as stationary against manual expert assessments for 241 distinct algorithm-derived stationary trajectories (comprising 1,452 individual polygons) within the Adélie Land validation set (Table 6). As observed in visual assessments, high-density packing frequently causes adjacent icebergs to be segmented as a single merged polygon, while summer surface meltwater can lead to target fragmentation. Consequently, a single physical iceberg may be represented by multiple trajectories segments, or multiple icebergs by a single trajectory (Figure S4).

Overall, the algorithm successfully captured the full observable duration of stationarity for the vast majority of these identified trajectories (77.2%). When evaluating the physical targets corresponding to these trajectories, delayed identification (i.e., cases where the iceberg was assessed manually as stationary prior to the algorithm) occurred in 30 icebergs (12.4%), lagging by between 1 and 3 frames primarily due to sea-ice clutter obscuring initial target boundaries. Premature terminations (i.e., cases where the manually assigned period of stationarity extended longer than the algorithm’s assessment) affected 22 icebergs (9.1%), ending 1 to 2 frames early as summer surface meltwater degraded radar contrast. Complete identification failures were




Figure 6. Multi-temporal evolution of an iceberg grounding-ungrounding event. The gallery displays a sequence of Sentinel-1 SAR imagery patches showing the tracked iceberg throughout its grounding event (first two frames), its period of being grounded (middle two frames), and ungrounding (final two frames). The target iceberg is delineated by a red mask during its grounded phase. Images with blue titles indicate reference timestamps confirming that the iceberg was grounded only for the two middle frames.

exceptionally rare, occurring in only 3 instances (1.2%). While intermittent mid-trajectory misses were recorded (56 frame instances), our multi-temporal matching effectively bridged these gaps to maintain overall state continuity.

In summary, the validation results from Adélie Land demonstrate that the identification algorithm can maintain continuous identification and identity consistency of iceberg targets in complex environments, providing a reliable technical foundation for constructing a high-precision, circum-Antarctic grounded iceberg dataset.

4.2 Circum-Antarctic grounded iceberg dataset

Based on the previously validated detection and tracking framework, we applied our method to Sentinel-1 SAR imagery acquired from late February to early April 2025 around the Antarctic coastline (Figure 1), thereby constructing the first high-resolution circum-Antarctic grounded iceberg dataset. This dataset records the geographic locations, morphological contours, and geometric attributes of each grounded iceberg, revealing the macroscopic distribution patterns of grounded icebergs along the Antarctic coastline.

4.2.1 General Statistics

To quantify the effectiveness of the physical constraint filtering (described in Section 3.4), we tracked the number of stationary iceberg candidates removed at each stage. From the initial pool of ~~45,530~~ 50,448 tracked stationary targets, ~~6,625~~ 10,829 candidates were eliminated (Table 7).

After filtering, the final dataset consisted of ~~38,905~~ 39,619 stationary targets along the Antarctic continental shelf, with a combined area of ~~13,719~~ 430 km^2 . As shown in Figure 7, the frequency distribution of grounded iceberg areas is strongly skewed towards smaller sizes, with a mean area of ~~0.35~~ 0.34 km^2 and a median area of $0.13 km^2$. Notably, icebergs with an area of less than $1 km^2$ dominate numerically, accounting for as much as ~~92.2~~ 92.6 % of the total number. By area, tiny

Table 6. Correct versus incorrect grounded iceberg identification metrics. In this evaluation, temporal sequences of n images are analysed. The terms “Start” and “End” refer to the specific image frame number within the sequence at which an iceberg was first and last identified as stationary by the algorithm. For a detailed explanation of the evaluation methodology, refer to Section 4.1.2.

Category	Offset (Frames)	Count	Proportion (%)
Correct Identification Metrics			
Overall Successful Identification	—	238 Trajectories	99.1%
Captured Full Observable Duration	—	186 Trajectories	77.2%
Consistently Stationary (Zero Misses)	—	159 Trajectories	65.9%
Incorrect Identification & Categorisation Errors			
Delayed Identification (Start)	1	18 Icebergs	7.5%
Delayed Identification (Start)	2	6 Icebergs	2.5%
Delayed Identification (Start)	3	6 Icebergs	2.5%
Premature Termination (End)	1	17 Icebergs	7.1%
Premature Termination (End)	2	5 Icebergs	2.1%
Completely Missed Grounding	—	3 Icebergs	1.2%
Intermittent Misclassification	—	56 Frames	3.8%

Table 7. Detailed statistics of iceberg candidates removed by the physical constraint filters. Depth zones are divided into deep (> 600 m) and shallow (≤ 600 m) waters. The “Reason” corresponds to the filter IDs defined in Table 2 (C1: Depth > 600 m & SIC $> 40\%$ & Area < 1 km²; C2: SIC $> 80\%$ & Area < 0.1 km²; C3: Depth > 1000 m). SIC: Sea Ice Concentration.

Depth Zone	Reason	Count	% of Removed	Median Area (km ²)	Mean Max SIC (%)	Median Bed Depth (m)
Deep (> 600 m)	C1	2545	23.50	0.84	73.3	683.0
Deep (> 600 m)	C1+C2	1056	9.75	0.36	87.9	656.0
Deep (> 600 m)	C1+C3	602	5.56	0.29	69.3	2146.8
Deep (> 600 m)	C3	467	4.31	0.57	8.4	1164.0
Deep (> 600 m)	C1+C2+C3	200	1.85	0.26	86.8	1662.0
Shallow (≤ 600 m)	C2	5947	54.92	0.35	88.4	369.0
Shallow (≤ 600 m)	C1	9	0.08	2.26	71.9	597.0
Shallow (≤ 600 m)	C1+C2	3	0.03	0.24	87.3	598.5

icebergs (A0 and A1) also account for ~~52.58~~53.90%. It is important to note that this proportion is likely inflated, as the model’s limitation in fully segmenting large and giant icebergs leads to an underestimation of their total area.




Figure 7. Size distribution analysis of unique grounded icebergs. (A) Frequency distribution of the physical area (km^2) for all identified icebergs. The y-axis is presented on a logarithmic scale to account for the high frequency of smaller icebergs. Red and orange dashed lines indicate the mean and median physical areas, respectively. (B) Categorical distribution of icebergs across classes. Labels above the bars indicate the total count and the relative percentage of the total population for each category.

Of the ~~38,905~~39,619 stationary targets identified, spatial analysis reveals that 27,~~431~~(70.5~~634~~67.4 %) are not located within or adjacent to fast ice (i.e., classified as ‘~~Outside~~’“Outside”), confirming their status as high-confidence grounded. Meanwhile, ~~10,869~~11,436 targets are situated completely within fast ice, and ~~605~~626 exhibit partial overlap. While these ‘~~Inside~~’ and ‘~~Partial~~’“Inside” and “Partial” targets remain stationary, their grounding status is subject to higher uncertainty due to potential fastening by the surrounding fast ice.

4.2.2 Spatial Distribution

The spatial density of grounded icebergs around Antarctica exhibits heterogeneity. As shown in Figure 8, nearly all high-density hotspots are concentrated in shallow coastal regions. In these areas, shallow bathymetric features, such as continental banks and ridges, provide favourable anchoring points for iceberg grounding, especially downstream of regions where active ice-shelf calving supplies a plentiful source of icebergs. For example, in regions such as Thwaites Glacier ($\sim 106.75^\circ W$) and Fram Bank ($\sim 69.5^\circ E$), the maximum density reaches 41 icebergs per $25 km^2$, forming complex coastal “grounded iceberg chains”. In contrast, grounded icebergs are sparsely distributed in deep-water regions, such as Pine Island Trough ($\sim 101^\circ W$), where excessive water depths prevent long-term grounding. Similarly, low densities are observed in areas lacking productive

upstream sources; notably, while massive ice shelves such as the Ross Ice Shelf ($\sim 180^\circ$) produce giant tabular icebergs, they calve infrequently, resulting in fewer individual grounding events compared to smaller, frequently-calving ice shelves.




Figure 8. Spatial distribution of grounded icebergs. (A) Circum-Antarctic density heatmap (counts per $5 \times 5 \text{ km}$ grid) overlaid on International Bathymetric Chart of the Southern Ocean (IBCSO) bathymetry version 2 (Dorschel et al., 2022). Red boxes indicate the locations of the detail panels. (B, C) Close-up views of Thwaites Glacier and Western Getz Ice Shelf, respectively, displaying individual grounded iceberg polygons (orange) to illustrate their alignment along shallow continental banks. (D, E) Corresponding Sentinel-1 SAR imagery for the regions shown in (B) and (C). Panel (D) shows the Thwaites region (EW mode, HV polarisation) acquired on 17 March 2025, and Panel (E) shows the Western Getz region (IW mode, HH polarisation) acquired on 22 March 2025.

560 Despite regional heterogeneity, grounded icebergs remain a widely-encountered feature of the Antarctic margin. By projecting iceberg locations onto the main continental coastline and dividing the coast into 100 km intervals, we found that grounded icebergs are present along ~~56.71~~56.52 % of the entire coastline, further confirming their role as a widespread boundary condition for the coastal cryosphere. However, their spatial density is highly uneven. Cumulative distribution analysis based on 100 km coastal segments demonstrates extreme spatial clustering in both count and physical area. Specifically, 50 % of the

565 grounded icebergs (by number) are concentrated within only 4.9 % of the coastline, closely mirrored by the distribution of total grounded area (50 % within 3.8 %). Extending to the bulk of the population, 14.2 % of the coastline contains 80 % of

the total count, while 80 % of the total grounded area is confined to an even narrower ~~11.7–11.5~~ % extent (Figure 9). This indicates that, although these “micro-anchors” are widely distributed, ~~the “picket fence effect”~~ their influence is particularly pronounced in specific high-density hotspots. In these regions, the dense clustering creates a “picket fence effect”, acting as a series of physical anchors to stabilise fast ice and restrict its drift.

5 Discussion

5.1 Scientific Significance and Implications

The primary contribution of this study is the establishment of the first high-resolution, circum-Antarctic dataset of grounded icebergs around Antarctica, based on Sentinel-1 SAR imagery. By integrating deep learning-based detection with a multi-temporal identification algorithm, we successfully mapped the distribution of ~~38,905–39,619~~ stationary icebergs across the entire continental shelf. Crucially, we integrated high-resolution fast ice maps to partition these targets into high-confidence grounded icebergs (located in open water) and fast ice entrapped candidates. This novel classification partially resolves the ambiguity between grounding and fast ice immobilisation, revealing macroscopic distribution patterns previously obscured due to observational limitations. This achievement directly addresses a critical gap in current research on the Antarctic coastal cryosphere: the lack of automated methods for identifying grounded icebergs. Previous studies have largely relied on manual interpretation, which is restricted to specific regions or time periods and is unable to capture dynamic features at the circum-Antarctic scale (e.g., Li et al., 2020; Kostov et al., 2026).

Due to challenging background conditions caused by heterogeneous sea ice and textural similarities, icebergs smaller than 0.1 km^2 are easily missed by conventional SAR thresholding methods or earlier machine learning models (e.g., Wesche and Dierking, 2015; Barbat et al., 2021; Evans et al., 2023). For instance, while the unsupervised clustering approach by Evans et al., 2023 achieved automated detection, it faced challenges with sub-kilometre icebergs (Class A1, $< 1 \text{ km}^2$), reporting a mean Precision of 0.619 and an F1 score of 0.678 due to high false positive rates in complex environments. In contrast, our deep learning framework demonstrates superior feature extraction capabilities for these small targets, achieving a Precision of 0.8652 and an F1 score of 0.8613. Furthermore, our approach surpasses the detection limit of previous studies ($< 0.04 \text{ km}^2$, e.g., Chen et al., 2026), reducing the minimum detectable iceberg size to 0.016 km^2 . This improvement effectively mitigates interference from seasonal sea ice cover variations and complex sea conditions, maintaining high detection stability across most of the year and under a wide range of environmental conditions.

The dataset generated in this study provides a high-precision spatial foundation for analysing the interactions between grounded icebergs, fast ice, and ocean dynamics (Massom et al., 2001; Porter-Smith et al., 2021; Fraser et al., 2023). The spatial distribution confirms that high-density clusters of grounded icebergs are predominantly constrained by shallow bathymetry, typically accumulating on continental banks downstream of ice shelf calving fronts. From an ecological perspective, these dense iceberg clusters are not merely physical barriers but also hotspots of biogeochemical cycling (Duprat et al., 2016; Lucas et al., 2025) and freshwater input (Olivé Abelló et al., 2025). Grounded icebergs stabilise the surrounding sea ice (Fraser et al., 2023), which serves as an important habitats for plankton, sea ice algae, and benthic organisms (Arrigo, 2016; Meiners et al.,




Figure 9. Statistical distribution and spatial concentration of grounded icebergs along the Antarctic coastline. (A) Spatial distribution frequency. The histogram shows the number of grounded icebergs aggregated within 100 km segments along the primary Antarctic coastline. The inset map illustrates the reference coastline, with red markers indicating the cumulative distance (km) measured from 0° longitude, providing geographic context for the x-axis. (B) Cumulative distribution (Lorenz-style curve). The curve quantifies the degree of spatial clustering by plotting the cumulative percentage of grounded icebergs against the cumulative percentage of coastline segments, ordered by iceberg density. The dashed grey line represents a perfectly uniform distribution, for reference.

600 2018; Bett et al., 2020). At the same time, the release of iron from iceberg meltwater constitutes a key limiting factor for primary productivity along the Antarctic coast (Herraiz-Borreguero et al., 2016). In contrast to individual large icebergs, the numerous tiny icebergs identified in this study collectively contribute a larger total surface area (~~52.58~~ 53.90 % of the overall area) and exhibit higher perimeter-to-area ratios. These characteristics indicate that they may play a greater role than previously recognised in nutrient release and in enhancing local stratification (Duprat et al., 2016; Lucas et al., 2025).

605 Our study reveals the predominance of “tiny icebergs” ($< 1 \text{ km}^2$) within the grounded iceberg population, accounting for ~~92.59~~ 92.6 % of the total. Grounded icebergs are considered key anchoring points for Antarctic fast ice (Massom et al., 2001; Fraser et al., 2023). Previous studies have largely focused on the blocking effects of small to medium icebergs on fast ice (e.g. Li et al., 2020), but the high-density clusters of tiny icebergs revealed in this study suggest that such a “granular” distribution may create a more refined and stable “picket fence effect”. This indicates that, although individual small grounded icebergs cover
610 only a limited area, their large numbers and widespread presence in shallow waters allow them to play a notable anchoring role in the maintenance of fast ice.

Recent dynamical modelling supports the physical viability of these smaller anchors; Kostov et al. (2026) demonstrate that sediment resistance acts as the primary decelerator against atmospheric drag. This resistive force enables even smaller grounded features to withstand strong wind forcing and maintain their position for extended periods, thereby underpinning the stability
615 of this micro-scale “picket fence”. Collectively, these high-density clusters may effectively increase friction along the coastal boundary, restricting the drift and export of sea ice, while physically dissipating the kinetic energy of offshore floes and swell (Massom et al., 2001; Bett et al., 2020).

However, this stabilising mechanism is geographically highly heterogeneous. The finding that 50 % of grounded icebergs are concentrated along only 4.9 % of the coastline, while 80 % occupy just 14.2 %, implies that the Antarctic “picket fence”
620 functions less like a continuous wall and more like a series of discrete, high-intensity structural nodes. This extreme spatial inequality suggests that the stability of circum-Antarctic fast ice may disproportionately depend on these specific “hotspot” regions.

This stability mechanism, which relies on both the numerical dominance of tiny icebergs and their high local concentration, may indicate a unique sensitivity of the Antarctic coastal system to climate change. While their keel depths can be substantial due to potential overturning, tiny icebergs possess considerably lower total mass and thermal inertia compared to large
625 tabular icebergs. Consequently, they are more susceptible to rapid volume loss, disintegration, and subsequent “ungrounding” in response to ocean warming. This implies that even subtle perturbations in environmental conditions within these limited critical zones could lead to the rapid disintegration of this “micro-scale picket fence”, potentially triggering abrupt changes in the stability of fast ice through non-linear processes. In the context of the recent sharp decline in Antarctic fast ice extent
630 (Fraser et al., 2023; Abram et al., 2025; Doddridge et al., 2025), understanding this mechanism is particularly important. Our data suggest that subtle changes in the distribution of grounded icebergs may, through this non-linear collective effect, directly modulate the stability of fast ice and the formation potential of polynyas (Nakayama et al., 2014; Olivé Abelló et al., 2025).

5.2 Limitations

Although this study successfully constructed a circum-Antarctic grounded-iceberg dataset, several limitations remain in relation to data annotation, algorithmic adaptability, and environmental interference, all of which should be taken into account in subsequent applications and analyses.

The construction of a pan-Antarctic mosaic necessitates the integration of heterogeneous SAR data, introducing spatial and radiometric inconsistencies. To achieve complete coastal coverage, this study combines IW and EW modes. While this strategy maximises spatial coverage, the integration of different modes leads to spatially variable detection sensitivities. In our study, the EW mode demonstrated stronger detection capabilities compared to the IW mode, resulting in non-uniform performance baselines across different continental sectors. However, we note that this discrepancy is primarily driven by data availability and model training biases rather than the inherent physics of the SAR modes. Our deep learning model was trained on a dataset primarily composed of EW imagery. Furthermore, due to data availability, the mosaic utilises varying polarisation channels, primarily co-polarisation and cross-polarisation. Iceberg–ocean contrast differs between polarisations, and cross-polarisation typically provides superior contrast over co-polarisation. Consequently, the model’s segmentation sensitivity fluctuates across different orbital swaths, potentially introducing bias in regional iceberg density estimates.

In terms of data annotation and the detection of very small targets (< 25 pixels), both manual mask labelling and automated model prediction face substantial challenges. Extremely small icebergs (< 25 pixels) are often embedded within complex sea ice-dominated backgrounds, leading to ambiguous boundary definition. For targets smaller than 10 pixels in particular, local high backscatter caused by strong winds or wave breaking (sea clutter) is almost indistinguishable from real icebergs in terms of brightness (Marino et al., 2016). The close similarity in backscatter intensity introduces uncertainty into manual interpretation, which inevitably injects label noise into the training dataset. It also reduces the model’s discriminative power during inference, making it difficult to eliminate false positives under complex sea conditions. In addition, in extremely dense iceberg regions, the model occasionally merges closely adjacent icebergs into a single entity, potentially resulting in slight underestimation of the local iceberg count.

A further limitation of the spatiotemporal identification algorithm emerges in regions characterised by an extreme density of small, highly mobile icebergs, such as the Bellingshausen Sea. In these highly congested areas, coincidental spatial overlap introduces an unavoidable source of false positives. Even when the independent single-frame detection and multi-temporal tracking algorithms are functioning nominally, in some cases these may have been misidentified as grounded when different small drifting bergs sequentially occupied overlapping positions between the 12-day SAR passes. Because these transient targets coincidentally satisfy the geometric and spatial proximity thresholds required for trajectory continuation, the algorithm inherently interprets them as a single stationary entity.

A specific uncertainty arises from the reliance on bathymetric constraints for false positive removal. While the application of a 1,000 *m* depth threshold effectively eliminates spurious detections in deep open-ocean regions because grounding is physically impossible, it implies that similar detection artifacts likely persist in the shallower continental shelf regions. False positives caused by complex sea ice background texture, stationary fast ice ridges, or immobilised floating icebergs share iden-

tical radiometric characteristics in both deep and shallow waters. However, our filtering framework actively mitigates these across both regimes, with deep and shallow waters accounting for approximately 46% and 54% of the eliminated targets, respectively (Table 7). In the potential grounding zone ($< 600\text{ m}$), where depth criteria alone are insufficient, we utilise a combined area and SIC strategy to manage complex sea ice backgrounds. This approach is physically justified: in shallow waters, icebergs immobilised by sea ice are highly likely to be genuinely grounded if they are sufficiently large. Our area thresholds inherently retain these larger icebergs, which possess deeper keels capable of interacting with the seabed, while effectively filtering out smaller sea ice artifacts. Although a minor proportion of unquantifiable false positives persists, particularly in areas with ~~complex fast-ice topography~~ strong backscatter associated with large-scale deformed fast ice (especially multi-year fast ice), this dual approach improves the robustness of final dataset. However, we also acknowledge that inaccuracies or coarse spatial resolutions in these underlying auxiliary datasets (such as IBCSO bathymetry or AMSR2 SIC) could theoretically lead to the erroneous exclusion of truly grounded icebergs, thereby producing false negatives.

Regarding the distinction between grounding and entrapment (i.e., immobilisation by the fast ice itself), reliance on auxiliary datasets introduces specific uncertainties. In previous methodologies, distinguishing grounded icebergs from floating ones immobilised by landfast sea ice was a primary challenge, as both share identical radiometric characteristics. To address this, we integrated high-resolution fast ice maps to partition targets into three categories (Section 3.4), thereby effectively isolating high-confidence grounded icebergs in open water. However, this classification strategy implies that the reliability of the grounding status for ‘Inside’ and ‘Partial’ targets is inherently dependent on the accuracy and temporal synchronisation of the input fast ice mask. Any potential errors in the fast ice boundary definition could lead to misclassification of the grounding status for these specific candidates. Consequently, while the ‘Outside’ category represents a robust baseline of truly grounded targets, ‘the Inside’ and ‘Partial’ categories should be interpreted with consideration of the surrounding fast ice context.

Regarding the identification algorithm, although Hu invariant moments (Hu, 1962) are generally robust for shape-based matching, their stability decreases when applied to tiny icebergs (< 25 pixels). Owing to the limits of spatial resolution, shape descriptors for such small objects are easily affected by pixel-level quantisation errors, reducing the reliability of trajectory association. Similarly, in densely packed iceberg clusters, the frequent merging and separation of neighbouring icebergs can cause unstable variations in the segmentation results. These fluctuations can lead to abrupt changes in key object features (such as area and centroid), thereby disrupting the continuity of feature-based trajectory matching.

The temporal inconsistency of the data acquisition introduces environmental variability. As the dataset is a composite of acquisitions spanning late summer to early autumn, specifically February to April, environmental conditions are not uniform. Acquisitions from February may still be influenced by residual surface melt and wet snow. These conditions reduce the radar contrast between icebergs and the background, thereby lowering detection scores, as evidenced by the drop in F1 score for December to 0.8472 compared to the annual average of 0.9121 (Table 3). Conversely, images acquired in the colder transition period (March and April) may contain patches of residual high-backscatter multi-year fast ice. These features can be misclassified as icebergs due to their similar texture. Although we primarily targeted the March transition period to mitigate these extremes, this temporal spread implies that the background noise varies geographically across the dataset. Consequently, precision in specific regions may fluctuate, as indicated by the March precision of 0.8526 shown in Table 2.

Finally, due to the tile-processing strategy and the limited spatial context captured by the model, the ResUNet architecture still faces challenges when handling large and giant icebergs. Although it outperforms previous methods for detecting small to medium-sized icebergs, icebergs larger than the field of view of a single sub-image are difficult to segment completely. The lack of global contextual information prevents the model from perceiving the continuous geometry of an object that spans multiple tiles, which may lead to fragmented segmentation or missing boundaries in large-scale contours.

5.3 Future work

Future research will aim to fully exploit the scientific value of this dataset while addressing the limitations identified in the present study. First, multi-source remote sensing data fusion can be further advanced by integrating altimetry datasets (such as the Ice, Cloud, and land Elevation Satellite-2 – ICESat-2), extending the current two-dimensional monitoring to three-dimensional volume [estimation](#), and facilitating draft estimation (Markus et al., 2017). ~~This approach enables more accurate reconstruction of iceberg morphology and volume, while~~ [While fully reconstructing the 3D surface morphology of individual icebergs is challenging given the sparse nature of altimetry profiles, fusing these draft estimations with 2D SAR boundaries remains highly valuable for statistical volume approximations. This approach facilitates](#) quantifying freshwater flux from melting, thereby improving understanding of its impacts on local salinity, ocean circulation, and polar coastal ecosystems (Olivé Abelló et al., 2025). [Looking ahead, next-generation swath-mapping missions like the Earth Dynamics Geodetic Explorer \(EDGE\) promise to overcome these profiling limitations, providing contiguous coverage that will enable true 3D morphological reconstructions \(Garvin et al., 2026\).](#) In addition, multi-source data fusion may enhance monitoring across iceberg scales, from tiny to giant, providing a reliable basis for long-term dynamic databases and predicting climate-driven responses.

Second, a long-term dynamic monitoring system can be established using historical SAR archive data to reconstruct a decadal record of grounded icebergs across Antarctica, thereby revealing their long-term spatiotemporal evolution under a warming climate (Fraser et al., 2023; Doddridge et al., 2025). Building on this, process-based studies of the iceberg–sea ice–ocean system can be conducted by integrating high spatiotemporal resolution sea ice drift and ocean forcing data, quantifying the thermodynamic control of grounded iceberg distributions on fast ice stability (Massom et al., 2001; Fraser et al., 2023) and polynya formation (Nakayama et al., 2014), and exploring their potential cascading effects on polar coastal ecosystems (Arrigo, 2016; Meiners et al., 2018). However, accurate discrimination between stationary and truly grounded icebergs becomes more difficult prior to 2021, when summertime fast ice extent was far higher than in recent years (Abram et al., 2025).

Finally, to further improve model performance under complex conditions, future work could incorporate global contextual information [\(referring to the broader spatial context beyond the current \$250 \times 250\$ pixel sub-image input in this study\)](#) into the modelling framework. ~~Compared with features derived from single sub-images, integrating global~~ [This wider view serves two critical purposes. It allows the model to perceive the continuous geometry of larger icebergs that span across multiple tiles, thereby preventing fragmented segmentation. In addition, integrating a wider](#) spatial context provides richer environmental constraints, helping the model to ~~accurately~~ [recognise large-scale background interference \(such as extensive patches of severe sea clutter or heavily deformed fast ice\), allowing it to better](#) distinguish true targets from false signals (including those indis-

tinguishable by bathymetry) in cases where tiny icebergs are highly similar to background interference, such as sea clutter or residual fast ice.

6 Conclusion

740 In response to the current lack of automated monitoring of grounded icebergs across Antarctica, this study introduces an automated detection framework that integrates a proposed ResUNet deep-learning network with a multi-temporal identification algorithm. The method demonstrates strong robustness against signal interference in complex coastal environments, achieves stable detection performance ($F1 > 0.91$), and successfully reduces the minimum detectable iceberg size to 0.016 km^2 . Using Sentinel-1 SAR imagery, we construct the first high-resolution, continent-wide dataset of grounded icebergs in Antarctica.

745 Statistical analysis shows that almost ~~39~~40,000 stationary icebergs were identified along the Antarctic coastline in March 2025 (~~70.5~~67.4 % of which were classified as high-confidence grounded icebergs), with tiny icebergs ($< 1 \text{ km}^2$) dominating numerically and accounting for ~~92.2~~92.6 % of all grounded icebergs. Spatial patterns further reveal that these high-density iceberg clusters are concentrated mainly on shallow continental shelves and along the actively disintegrating fronts of ice shelves, forming complex and discontinuous “grounded iceberg chains”. Overall, these features are distributed along ~~56.71~~56.52 % of the continental coastline, with 80 % concentrated within just 14.2 % of the coastline. This finding builds on earlier
750 studies that predominantly examined medium and large icebergs, providing further evidence that aggregations of tiny icebergs may play a more important role than previously recognised in maintaining fast ice and regulating biogeochemical processes through a “picket fence effect”.

The dataset provides a new high-precision benchmark for quantifying the Antarctic ice–ocean interaction network, and it establishes a robust technical foundation for reconstructing long-term iceberg dynamics from historical imagery and projecting
755 their responses to climate change.

Author contributions. KJ: Conceptualisation, methodology, software, formal analysis, data curation, validation, visualisation, writing – original draft. ADF: Conceptualisation, supervision, methodological guidance, funding acquisition, writing – review & editing. JL: Conceptualisation, supervision, methodological guidance, writing – review & editing. PW: Supervision, methodological guidance, writing – review & editing. CA: Data curation, resources (Sentinel-1 processing). ACB: Data curation, resources (Sentinel-1 processing).

760 *Competing interests.* ADF is a member of the editorial board of ESSD. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declared.

7 Data availability

The grounded iceberg dataset is publicly available at <https://doi.org/10.25959/54sx-pt47> (Jiao et al., 2026). Detailed field descriptions are provided in Table 1.

765 8 Code availability

The code and datasets supporting this study are available on GitHub at <https://github.com/kj0703/Antarctic-Grounded-Iceberg-Detection> and are permanently archived on Zenodo at <https://doi.org/10.5281/zenodo.19228332>.

Acknowledgements. ADF is supported by the Australian Research Council, through grants FT230100234, LP170101090, LE220100103, and DP240100325. PW was supported under the Australian Research Council's Special Research Initiative for Antarctic Gateway Partnership (SR140300001). This project received grant funding from the Australian Government as part of the Antarctic Science Collaboration Initiative program. ADF and PW acknowledge the generous support of the Harris Charitable Trust through the Antarctic Science Foundation. This research was undertaken with the assistance of resources from the National Computational Infrastructure (NCI Australia), an NCRIS enabled capability supported by the Australian Government (Project jk72). CA and ACB publish with the permission of the CEO, Geoscience Australia. The authors acknowledge the use of Gemini 3 for assistance in optimising the data processing scripts. All AI-generated outputs
775 were rigorously reviewed and edited by the authors.

References

- Abram, N., Purich, A., England, M., McCormack, F., Strugnelli, J., Vance, T., Wienecke, B., and Fraser, A.: Emerging evidence of abrupt changes in the Antarctic environment, <https://doi.org/10.48361/Q2P7-G949>, 2025.
- Arrigo, K. R.: Sea ice as a habitat for primary producers, *Sea Ice*, p. 352–369, <https://doi.org/10.1002/9781118778371.ch14>, 2016.
- 780 Barbat, M. M., Wesche, C., Werhli, A. V., and Mata, M. M.: An adaptive machine learning approach to improve automatic iceberg detection from SAR images, *ISPRS Journal of Photogrammetry and Remote Sensing*, 156, 247–259, <https://doi.org/10.1016/j.isprsjprs.2019.08.015>, 2019.
- Barbat, M. M., Rackow, T., Wesche, C., Hellmer, H. H., and Mata, M. M.: Automated iceberg tracking with a machine learning approach applied to SAR imagery: A Weddell sea case study, *ISPRS Journal of Photogrammetry and Remote Sensing*, 172, 189–206, <https://doi.org/10.1016/j.isprsjprs.2020.12.006>, 2021.
- 785 Bett, D. T., Holland, P. R., Naveira Garabato, A. C., Jenkins, A., Dutrieux, P., Kimura, S., and Fleming, A.: The Impact of the Amundsen Sea Freshwater Balance on Ocean Melting of the West Antarctic Ice Sheet, *Journal of Geophysical Research: Oceans*, 125, <https://doi.org/10.1029/2020jc016305>, 2020.
- Bigg, G. R., Wadley, M. R., Stevens, D. P., and Johnson, J. A.: Modelling the dynamics and thermodynamics of icebergs, *Cold Regions Science and Technology*, 26, 113–135, [https://doi.org/10.1016/s0165-232x\(97\)00012-8](https://doi.org/10.1016/s0165-232x(97)00012-8), 1997.
- 790 Braakmann-Folgmann, A., Shepherd, A., Gerrish, L., Izzard, J., and Ridout, A.: Observing the disintegration of the A68A iceberg from space, *Remote Sensing of Environment*, 270, 112855, <https://doi.org/10.1016/j.rse.2021.112855>, 2022.
- Braakmann-Folgmann, A., Shepherd, A., Hogg, D., and Redmond, E.: Mapping the extent of giant Antarctic icebergs with deep learning, *The Cryosphere*, 17, 4675–4690, <https://doi.org/10.5194/tc-17-4675-2023>, 2023.
- 795 Bradski, G.: The opencv library., *Dr. Dobb's Journal: Software Tools for the Professional Programmer*, 25, 120–123, 2000.
- Cefarelli, A. O., Ferrario, M. E., and Vernet, M.: Diatoms (Bacillariophyceae) associated with free-drifting Antarctic icebergs: taxonomy and distribution, *Polar Biology*, 39, 443–459, <https://doi.org/10.1007/s00300-015-1791-z>, 2015.
- Chen, Z., Liu, X., Guan, Z., Li, T., Cheng, X., Li, T., Liu, Y., Liang, Q., Zheng, L., and Liu, J.: A six-year circum-Antarctic icebergs dataset (2018–2023), *Earth System Science Data*, 18, 147–166, <https://doi.org/10.5194/essd-18-147-2026>, 2026.
- 800 Deng, J., Dong, W., Socher, R., Li, L.-J., Li, K., and Fei-Fei, L.: ImageNet: A large-scale hierarchical image database, in: 2009 IEEE Conference on Computer Vision and Pattern Recognition, IEEE, <https://doi.org/10.1109/cvpr.2009.5206848>, 2009.
- Diakogiannis, F. I., Waldner, F., Caccetta, P., and Wu, C.: ResUNet-a: A deep learning framework for semantic segmentation of remotely sensed data, *ISPRS Journal of Photogrammetry and Remote Sensing*, 162, 94–114, <https://doi.org/10.1016/j.isprsjprs.2020.01.013>, 2020.
- Dierking, W.: Sea ice monitoring by synthetic aperture radar, *Oceanography*, 26, 100–111, 2013.
- 805 Dierking, W. and Wesche, C.: C-Band Radar Polarimetry—Useful for Detection of Icebergs in Sea Ice?, *IEEE Transactions on Geoscience and Remote Sensing*, 52, 25–37, <https://doi.org/10.1109/tgrs.2012.2234756>, 2014.
- Doddridge, E. W., Hobbs, W. R., Auger, M., Boyd, P. W., Chua, S. M. T., Cook, S., Corney, S., Emmerson, L., Fraser, A. D., Heil, P., Kelly, N., Lannuzel, D., Li, X., Liniger, G., Massom, R. A., Meyer, A., Reid, P., Southwell, C., Spence, P., Stekettee, A., Swadling, K. M., Teder, N., Wienecke, B., Wongpan, P., and Yamazaki, K.: Impacts of Antarctic summer sea-ice extremes, *PNAS Nexus*, 4, <https://doi.org/10.1093/pnasnexus/pgaf164>, 2025.
- 810 Dorschel, B., Hehemann, L., Viquerat, S., Warnke, F., Dreutter, S., Tenberge, Y. S., Accettella, D., An, L., Barrios, F., Bazhenova, E., Black, J., Bohoyo, F., Davey, C., De Santis, L., Dotti, C. E., Fremant, A. C., Fretwell, P. T., Gales, J. A., Gao, J., Gasperini, L., Greenbaum, J. S.,

- Jencks, J. H., Hogan, K., Hong, J. K., Jakobsson, M., Jensen, L., Kool, J., Larin, S., Larter, R. D., Leitchenkov, G., Loubrieu, B., Mackay, K., Mayer, L., Millan, R., Morlighem, M., Navidad, F., Nitsche, F. O., Nogi, Y., Pertuisot, C., Post, A. L., Pritchard, H. D., Purser, A.,
815 Rebesco, M., Rignot, E., Roberts, J. L., Rovere, M., Ryzhov, I., Sauli, C., Schmitt, T., Silvano, A., Smith, J., Snaith, H., Tate, A. J., Tinto, K., Vandenbossche, P., Weatherall, P., Wintersteller, P., Yang, C., Zhang, T., and Arndt, J. E.: The International Bathymetric Chart of the Southern Ocean Version 2, Scientific Data, 9, <https://doi.org/10.1038/s41597-022-01366-7>, 2022.
- Dowdeswell, J. and Bamber, J.: Keel depths of modern Antarctic icebergs and implications for sea-floor scouring in the geological record, *Marine Geology*, 243, 120–131, <https://doi.org/10.1016/j.margeo.2007.04.008>, 2007.
- 820 Duprat, L. P. A. M., Bigg, G. R., and Wilton, D. J.: Enhanced Southern Ocean marine productivity due to fertilization by giant icebergs, *Nature Geoscience*, 9, 219–221, <https://doi.org/10.1038/ngeo2633>, 2016.
- EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF): Global Sea Ice Concentration (AMSR-2) - OSI-408-a, <https://osi-saf.eumetsat.int/products/osi-408>, accessed: 2026-01-15, 2025.
- European Space Agency: Sentinel-2 User Handbook, ESA/ESTEC, Paris, France, issue 1, revision 2 edn., standard Document, 2015.
- 825 Evans, B., Faul, A., Fleming, A., Vaughan, D. G., and Hosking, J. S.: Unsupervised machine learning detection of iceberg populations within sea ice from dual-polarisation SAR imagery, *Remote Sensing of Environment*, 297, 113 780, <https://doi.org/10.1016/j.rse.2023.113780>, 2023.
- Fraser, A., Massom, R., and Michael, K.: A Method for Compositing Polar MODIS Satellite Images to Remove Cloud Cover for Landfast Sea-Ice Detection, *IEEE Transactions on Geoscience and Remote Sensing*, 47, 3272–3282, <https://doi.org/10.1109/tgrs.2009.2019726>,
830 2009.
- Fraser, A. D., Massom, R. A., and Michael, K. J.: Generation of high-resolution East Antarctic landfast sea-ice maps from cloud-free MODIS satellite composite imagery, *Remote Sensing of Environment*, 114, 2888–2896, <https://doi.org/10.1016/j.rse.2010.07.006>, 2010.
- Fraser, A. D., Massom, R. A., Michael, K. J., Galton-Fenzi, B. K., and Lieser, J. L.: East Antarctic Landfast Sea Ice Distribution and Variability, 2000–08, *Journal of Climate*, 25, 1137–1156, <https://doi.org/10.1175/jcli-d-10-05032.1>, 2012.
- 835 Fraser, A. D., Ohshima, K. I., Nishashi, S., Massom, R. A., Tamura, T., Nakata, K., Williams, G. D., Carpentier, S., and Willmes, S.: Landfast ice controls on sea-ice production in the Cape Darnley Polynya: A case study, *Remote Sensing of Environment*, 233, 111 315, <https://doi.org/10.1016/j.rse.2019.111315>, 2019.
- Fraser, A. D., Massom, R. A., Handcock, M. S., Reid, P., Ohshima, K. I., Raphael, M. N., Cartwright, J., Klekociuk, A. R., Wang, Z., and Porter-Smith, R.: Eighteen-year record of circum-Antarctic landfast-sea-ice distribution allows detailed baseline characterisation and
840 reveals trends and variability, *The Cryosphere*, 15, 5061–5077, <https://doi.org/10.5194/tc-15-5061-2021>, 2021.
- Fraser, A. D., Wongpan, P., Langhorne, P. J., Klekociuk, A. R., Kusahara, K., Lannuzel, D., Massom, R. A., Meiners, K. M., Swadling, K. M., Atwater, D. P., Brett, G. M., Corkill, M., Dalman, L. A., Fiddes, S., Granata, A., Guglielmo, L., Heil, P., Leonard, G. H., Mahoney, A. R., McMinn, A., van der Merwe, P., Weldrick, C. K., and Wienecke, B.: Antarctic Landfast Sea Ice: A Review of Its Physics, Biogeochemistry and Ecology, *Reviews of Geophysics*, 61, <https://doi.org/10.1029/2022rg000770>, 2023.
- 845 Færch, L., Dierking, W., Hughes, N., and Doulgeris, A. P.: A comparison of constant false alarm rate object detection algorithms for iceberg identification in L- and C-band SAR imagery of the Labrador Sea, *The Cryosphere*, 17, 5335–5355, <https://doi.org/10.5194/tc-17-5335-2023>, 2023.
- Galler, B. A. and Fisher, M. J.: An improved equivalence algorithm, *Communications of the ACM*, 7, 301–303, <https://doi.org/10.1145/364099.364331>, 1964.

- 850 GAMMA Remote Sensing AG: GAMMA Software, GAMMA Remote Sensing AG, Gümligen, Switzerland, <https://www.gamma-rs.ch>, accessed: 2026-01-28, n.d.
- Garvin, J. B., McDonagh-Smith, P., Peters, R. M., Fricker, H. A., Armston, J. D., Blair, J. B., and Luthcke, S. B.: Return on Investment and Economic Impact of a NASA-Funded Swath Mapping Laser Altimeter Orbiter for Earth Surface Elevations Situational Awareness in 2031–2040, *New Space*, <https://doi.org/10.1177/21680256261434504>, 2026.
- 855 Gerrish, L., Ireland, L., Fretwell, P., Cooper, P., and Skachkova, A.: High resolution vector polygons of the Antarctic coastline - VERSION 7.11, <https://doi.org/10.5285/7D6793C2-2513-4752-8F37-11DB5B791FB5>, 2025.
- Gillies, Sean and others: Rasterio: Access to geospatial raster data, <https://github.com/rasterio/rasterio>, gitHub repository, accessed: 2026-01-19, 2013.
- Gladstone, R. M., Bigg, G. R., and Nicholls, K. W.: Iceberg trajectory modeling and meltwater injection in the Southern Ocean, *Journal of Geophysical Research: Oceans*, 106, 19 903–19 915, <https://doi.org/10.1029/2000jc000347>, 2001.
- 860 Greene, C. A., Gardner, A. S., Schlegel, N.-J., and Fraser, A. D.: Antarctic calving loss rivals ice-shelf thinning, *Nature*, 609, 948–953, <https://doi.org/10.1038/s41586-022-05037-w>, 2022.
- He, K., Zhang, X., Ren, S., and Sun, J.: Deep Residual Learning for Image Recognition, in: 2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), p. 770–778, IEEE, <https://doi.org/10.1109/cvpr.2016.90>, 2016.
- 865 Herraiz-Borreguero, L., Lannuzel, D., van der Merwe, P., Treverrow, A., and Pedro, J. B.: Large flux of iron from the Amery Ice Shelf marine ice to Prydz Bay, East Antarctica, *Journal of Geophysical Research: Oceans*, 121, 6009–6020, <https://doi.org/10.1002/2016jc011687>, 2016.
- Hopwood, M. J., Carroll, D., Höfer, J., Achterberg, E. P., Meire, L., Le Moigne, F. A. C., Bach, L. T., Eich, C., Sutherland, D. A., and González, H. E.: Highly variable iron content modulates iceberg-ocean fertilisation and potential carbon export, *Nature Communications*, 10, <https://doi.org/10.1038/s41467-019-13231-0>, 2019.
- 870 Hu, M.-K.: Visual pattern recognition by moment invariants, *IEEE Transactions on Information Theory*, 8, 179–187, <https://doi.org/10.1109/tit.1962.1057692>, 1962.
- ISCE3 Development Team: ISCE3 Documentation, <https://isce-framework.github.io/isce3/>, accessed: 19 May 2026, 2026.
- Jiao, K., Fraser, A. D., Lohse, J., Wongpan, P., Adams, C., and Bradley, A. C.: Grounded Icebergs around Antarctica: A High-Resolution Dataset Derived from Deep Learning and Sentinel-1 Synthetic Aperture Radar, <https://doi.org/10.25959/54sx-pt47>, 2026.
- 875 Koo, Y., Xie, H., Mahmoud, H., Iqrah, J. M., and Ackley, S. F.: Automated detection and tracking of medium-large icebergs from Sentinel-1 imagery using Google Earth Engine, *Remote Sensing of Environment*, 296, 113 731, <https://doi.org/10.1016/j.rse.2023.113731>, 2023.
- Kostov, Y., Holland, P. R., Hogan, K. A., Smith, J. A., Jourdain, N. C., Mathiot, P., Olivé Abelló, A., Fleming, A. H., and Meijers, A. J. S.: Modelled dynamics of floating and grounded icebergs, with application to the Amundsen Sea, *The Cryosphere*, 20, 135–169, <https://doi.org/10.5194/tc-20-135-2026>, 2026.
- 880 Kuhn, H. W.: The Hungarian method for the assignment problem, *Naval Research Logistics Quarterly*, 2, 83–97, <https://doi.org/10.1002/nav.3800020109>, 1955.
- Leane, E. and Maddison, B.: A biography of iceberg B09B, *Australian Humanities Review*, 63, 99–115, 2018.
- Li, X., Shokr, M., Hui, F., Chi, Z., Heil, P., Chen, Z., Yu, Y., Zhai, M., and Cheng, X.: The spatio-temporal patterns of landfast ice in Antarctica during 2006–2011 and 2016–2017 using high-resolution SAR imagery, *Remote Sensing of Environment*, 242, 111 736, <https://doi.org/10.1016/j.rse.2020.111736>, 2020.
- 885

- Lohse, J., Doulgeris, A. P., and Dierking, W.: Mapping sea-ice types from Sentinel-1 considering the surface-type dependent effect of incidence angle, *Annals of Glaciology*, 61, 260–270, <https://doi.org/10.1017/aog.2020.45>, 2020.
- 890 Loshchilov, I. and Hutter, F.: Decoupled Weight Decay Regularization, arXiv preprint arXiv:1711.05101, <https://doi.org/10.48550/ARXIV.1711.05101>, 2017.
- Lucas, N. S., Brearley, J. A., Hendry, K. R., Spira, T., Braakmann-Folgmann, A., Abrahamsen, E. P., Meredith, M. P., and Tarling, G. A.: Giant iceberg meltwater increases upper-ocean stratification and vertical mixing, *Nature Geoscience*, 18, 305–312, <https://doi.org/10.1038/s41561-025-01659-7>, 2025.
- Magnard, C., Werner, C., and Wegmüller, U.: Gamma technical report: Interpolation and resampling, Tech. rep., Gamma Remote Sensing
895 Research and Consulting AG, Bern, Switzerland, 2017.
- Marino, A., Dierking, W., and Wesche, C.: A Depolarization Ratio Anomaly Detector to Identify Icebergs in Sea Ice Using Dual-Polarization SAR Images, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 5602–5615, <https://doi.org/10.1109/tgrs.2016.2569450>, 2016.
- Markus, T., Neumann, T., Martino, A., Abdalati, W., Brunt, K., Csatho, B., Farrell, S., Fricker, H., Gardner, A., Harding, D., et al.: The Ice, Cloud, and land Elevation Satellite-2 (ICESat-2): science requirements, concept, and implementation, *Remote sensing of environment*,
900 190, 260–273, <https://doi.org/https://doi.org/10.1016/j.rse.2016.12.029>, 2017.
- Massom, R. A., Hill, K. L., Lytle, V. I., Worby, A. P., Paget, M., and Allison, I.: Effects of regional fast-ice and iceberg distributions on the behaviour of the Mertz Glacier polynya, East Antarctica, *Annals of Glaciology*, 33, 391–398, <https://doi.org/10.3189/172756401781818518>, 2001.
- Meiners, K. M., Vancoppenolle, M., Carnat, G., Castellani, G., Delille, B., Delille, D., Dieckmann, G. S., Flores, H., Fripiat, F., Grotti, M.,
905 Lange, B. A., Lannuzel, D., Martin, A., McMinn, A., Nomura, D., Peeken, I., Rivaro, P., Ryan, K. G., Stefels, J., Swadling, K. M., Thomas, D. N., Tison, J., van der Merwe, P., van Leeuwe, M. A., Weldrick, C., and Yang, E. J.: Chlorophyll-a in Antarctic Landfast Sea Ice: A First Synthesis of Historical Ice Core Data, *Journal of Geophysical Research: Oceans*, 123, 8444–8459, <https://doi.org/10.1029/2018jc014245>, 2018.
- Merino, N., Le Sommer, J., Durand, G., Jourdain, N. C., Madec, G., Mathiot, P., and Tournadre, J.: Antarctic icebergs melt over the Southern
910 Ocean: Climatology and impact on sea ice, *Ocean Modelling*, 104, 99–110, <https://doi.org/10.1016/j.ocemod.2016.05.001>, 2016.
- Micikevicius, P., Narang, S., Alben, J., Damos, G., Elsen, E., Garcia, D., Ginsburg, B., Houston, M., Kuchaiev, O., Venkatesh, G., and Wu, H.: Mixed Precision Training, <https://doi.org/10.48550/ARXIV.1710.03740>, 2017.
- Nakayama, Y., Timmermann, R., Schröder, M., and Hellmer, H.: On the difficulty of modeling Circumpolar Deep Water intrusions onto the Amundsen Sea continental shelf, *Ocean Modelling*, 84, 26–34, <https://doi.org/10.1016/j.ocemod.2014.09.007>, 2014.
- 915 Ohshima, K. I., Fukamachi, Y., Williams, G. D., Nihashi, S., Roquet, F., Kitade, Y., Tamura, T., Hirano, D., Herraiz-Borreguero, L., Field, I., Hindell, M., Aoki, S., and Wakatsuchi, M.: Antarctic Bottom Water production by intense sea-ice formation in the Cape Darnley polynya, *Nature Geoscience*, 6, 235–240, <https://doi.org/10.1038/ngeo1738>, 2013.
- Olivé Abelló, A., Mathiot, P., Jourdain, N. C., Kostov, Y., Holland, P. R., Gascoin, S., and Rousset, C.: Iceberg Grounding Enhances the Release of Freshwater on the Antarctic Continental Shelf, *Journal of Geophysical Research: Oceans*, 130,
920 <https://doi.org/10.1029/2025jc022857>, 2025.
- OPERA Jet Propulsion Laboratory: OPERA Radiometric Terrain-Corrected (RTC) Product, <https://github.com/opera-adt/RTC>, accessed: 19 May 2026, 2026a.
- OPERA Jet Propulsion Laboratory: Product Specification Document for the OPERA Radiometric Terrain Corrected SAR Backscatter from Sentinel-1, https://d2pn8kiwq2w21t.cloudfront.net/documents/ProductSpec_RTC-S1.pdf, accessed: 19 May 2026, 2026b.

- 925 OPERA Project Team: Algorithm Theoretical Basis Document (ATBD) for the OPERA Radiometric Terrain Corrected (RTC) Product from Sentinel-1, Technical Report D-107393, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA, <https://www.jpl.nasa.gov/>, initial Release, Version 1.0, 2024.
- Porter-Smith, R., McKinlay, J., Fraser, A. D., and Massom, R. A.: Coastal complexity of the Antarctic continent, *Earth System Science Data*, 13, 3103–3114, <https://doi.org/10.5194/essd-13-3103-2021>, 2021.
- 930 Rackow, T., Wesche, C., Timmermann, R., Hellmer, H. H., Juricke, S., and Jung, T.: A simulation of small to giant Antarctic iceberg evolution: Differential impact on climatology estimates, *Journal of Geophysical Research: Oceans*, 122, 3170–3190, <https://doi.org/10.1002/2016jc012513>, 2017.
- Raiswell, R., Benning, L. G., Tranter, M., and Tulaczyk, S.: Bioavailable iron in the Southern Ocean: the significance of the iceberg conveyor belt, *Geochemical Transactions*, 9, <https://doi.org/10.1186/1467-4866-9-7>, 2008.
- 935 Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341, 266–270, <https://doi.org/10.1126/science.1235798>, 2013.
- Ronneberger, O., Fischer, P., and Brox, T.: U-Net: Convolutional Networks for Biomedical Image Segmentation, p. 234–241, Springer International Publishing, ISBN 9783319245744, https://doi.org/10.1007/978-3-319-24574-4_28, 2015.
- Salehi, S. S. M., Erdogmus, D., and Gholipour, A.: Tversky Loss Function for Image Segmentation Using 3D Fully Convolutional Deep Networks, p. 379–387, Springer International Publishing, ISBN 9783319673899, https://doi.org/10.1007/978-3-319-67389-9_44, 2017.
- 940 Scambos, T., Fricker, H. A., Liu, C.-C., Bohlander, J., Fastook, J., Sargent, A., Massom, R., and Wu, A.-M.: Ice shelf disintegration by plate bending and hydro-fracture: Satellite observations and model results of the 2008 Wilkins ice shelf break-ups, *Earth and Planetary Science Letters*, 280, 51–60, <https://doi.org/10.1016/j.epsl.2008.12.027>, 2009.
- Snoek, J., Larochelle, H., and Adams, R. P.: Practical bayesian optimization of machine learning algorithms, *Advances in neural information processing systems*, 25, 2012.
- 945 Torres, R., Snoeij, P., Geudtner, D., Bibby, D., Davidson, M., Attema, E., Potin, P., Rommen, B., Floury, N., Brown, M., Traver, I. N., Deghaye, P., Duesmann, B., Rosich, B., Miranda, N., Bruno, C., L’Abbate, M., Croci, R., Pietropaolo, A., Huchler, M., and Rostan, F.: GMES Sentinel-1 mission, *Remote Sensing of Environment*, 120, 9–24, <https://doi.org/10.1016/j.rse.2011.05.028>, 2012.
- Tournadre, J., Bouhier, N., Girard-Ardhuin, F., and Rémy, F.: Antarctic icebergs distributions 1992–2014, *Journal of Geophysical Research: Oceans*, 121, 327–349, <https://doi.org/10.1002/2015jc011178>, 2016.
- 950 Truckenbrodt, J., Cremer, F., Baris, I., and Eberle, J.: pyroSAR: A framework for large-scale SAR satellite data processing, in: *Proceedings of the 2019 Conference on Big Data from Space (BiDS’19)*, pp. 197–200, Publications Office of the European Union, Munich, Germany, <https://doi.org/10.2760/848593>, 2019.
- U.S. National Ice Center: Antarctic Iceberg Naming and Tracking Information, accessed: 2025-12-03, 2024.
- 955 Vancoppenolle, M., Jourdain, N. C., Fichefet, T., Fraser, A. D., Massonnet, F., Mathiot, P., Madec, G., Mehlmann, C., Abelló, A. O., Atwater, D. P., Billy, C., Brearley, A., Delfieu, J., Kostov, Y., Lemaire, E., Marson, J. M., Petit, J., Pirlet, N., Wongpan, P., Burgard, C., de Lavergne, C., Holland, P. R., Hutchinson, K., Noël, M., Ortega, É., Piedagnel, E., Rampal, P., Ridley, J., Rousset, C., Sallée, J.-B., and Thomas, M.: Why Icebergs and their Interactions with Sea Ice Should Be Included in Earth System Models?, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/bams-d-25-0322.1>, 2025.
- 960 Wesche, C. and Dierking, W.: Iceberg signatures and detection in SAR images in two test regions of the Weddell Sea, Antarctica, *Journal of Glaciology*, 58, 325–339, <https://doi.org/10.3189/2012j0g11j020>, 2012.

- Wesche, C. and Dierking, W.: Near-coastal circum-Antarctic iceberg size distributions determined from Synthetic Aperture Radar images, *Remote Sensing of Environment*, 156, 561–569, <https://doi.org/10.1016/j.rse.2014.10.025>, 2015.
- Willis, C. J., Macklin, J. T., Partington, K. C., Teleki, K. A., Rees, W. G., and Williams, R. G.: Iceberg detection using ERS-1 synthetic aperture radar, *International Journal of Remote Sensing*, 17, 1777–1795, <https://doi.org/10.1080/01431169608948739>, 1996.
- Woo, S., Park, J., Lee, J.-Y., and Kweon, I. S.: CBAM: Convolutional Block Attention Module, p. 3–19, Springer International Publishing, ISBN 9783030012342, https://doi.org/10.1007/978-3-030-01234-2_1, 2018.
- Woodworth-Lynas, C., Josenhans, H., Barrie, J., Lewis, C., and Parrott, D.: The physical processes of seabed disturbance during iceberg grounding and scouring, *Continental Shelf Research*, 11, 939–961, [https://doi.org/10.1016/0278-4343\(91\)90086-1](https://doi.org/10.1016/0278-4343(91)90086-1), 1991.
- Wright, C. S. and Priestley, R. E.: *British (Terra Nova) Antarctic Expedition 1910–1913: Glaciology*, Harrison and Sons, Ltd., London, 1922.