

# Weather station data from the Mount Everest region, Nepal: 3,810-8,810 m above sea level

Arbindra Khadka<sup>1,2</sup>, L. Baker Perry<sup>3,4</sup>, Tom Matthews<sup>5</sup>, Tenzing Gyalzen Sherpa<sup>6</sup>, Chitra Bahadur Shrestha<sup>1</sup>, Dibas Shrestha<sup>1</sup>, Deepak Aryal<sup>1</sup>, Subash Tuladhar<sup>7</sup>, Niraj Pradhananga<sup>7</sup>, Dinkar Kayastha<sup>7</sup>, Brian Raichle<sup>4</sup>, Peter Athans<sup>8</sup>, Dawa Yangzum Sherpa<sup>8</sup>, Keith Garrett<sup>9</sup>, Garrett Wheeler<sup>10</sup>, Tom Young<sup>11</sup>, Aurora Elmore<sup>12</sup>

- 1 Central Department of Hydrology and Meteorology, Tribhuvan University, Kirtipur, Nepal
- 2 Université Grenoble Alpes, CNRS, IRD, IGE, Grenoble, France
- 3 Department of Geography (Climatology), University of Nevada, Reno, Nevada, USA
- 4 Research Institute for Environment, Energy, and Economics, Department of Sustainable Technology & the Built Environment, Appalachian State University, Boone, North Carolina
- 5 Department of Geography, King's College London, London, UK
- 6 Khumbu Climbing Centre, Phortse, Nepal
- 7 Department of Hydrology and Meteorology, Nepal
- 8 The North Face, Alameda, California
- 9 Mount Washington Observatory, North Conway, New Hampshire, USA
- 10 Campbell Scientific, Logan, Utah, USA
- 11 RM Young, Traverse City, Michigan, USA
- 12 National Oceanic and Atmospheric Administration, Maryland, USA

Corresponding Author: Arbindra Khadka [arbindra.khadka@cdhm.tu.edu.np](mailto:arbindra.khadka@cdhm.tu.edu.np)

## Abstract

~~Between 2019 and 2025, as part of the The~~ National Geographic ~~and~~ Rolex Perpetual Planet Everest expeditions, ~~a team of National Geographic Explorers, scientists, and elite climbing Sherpas~~ <sup>2019-2022</sup> installed a network of Automatic Weather Stations (AWSs) to improve understanding of the climate at high altitudes in the Nepal Himalaya. This knowledge is critical in the Mount Everest (Khumbu) region, due to its extreme altitude, popularity amongst trekkers and mountaineers, and its importance as a source of freshwater for downstream communities. Here we present quality controlled (QC) meteorological data from six AWSs ~~extending~~ from Phortse (3,810 m above sea level, m asl) to Bishop Rock (8,810 m asl) in the Everest region, including the seasonal climatology ~~focused on the three lower elevation AWSs~~, and a comparison with ERA5 reanalysis data from the South Col AWS. The data is accessible from <https://doi.org/10.5281/zenodo.18849098> (Khadka, 2026).

## 1. Introduction

Field research in High Mountain Asia (HMA), which ~~contains the largest glacierized~~ ~~has the most glacier~~ area outside of the polar regions (Brun et al., 2017; Pfeffer et al., 2014), is extremely difficult. The region's high ~~altitude result~~ ~~elevations result~~ in low barometric pressure (and hence relatively low oxygen availability: Matthews et al., 2020a), and extreme weather, challenging ~~both~~ personnel and equipment ~~during the assembling of alike in the task of assembling~~ high-quality, long-term observational datasets (Salerno et al., 2015; Shea et al., 2015; Khadka et al., 2022; Matthews et al., 2020a). However, there are strong incentives to overcome these difficulties. First, the region is an important freshwater source for ~~a vast number of~~ communities downstream (Immerzeel et al., 2010). Second, the extreme weather, combined with steep terrain, places the region at a relatively high risk of hydrometeorological hazards (Miner et al., 2020). Establishing meteorological observational networks both helps ~~to~~ improve understanding of water resource availability ~~in under~~ a changing climate, and ~~supports the reduction of~~ ~~to reduce~~ hydrometeorological risks.

Within HMA, the Khumbu region has attracted significant ~~cryospheric~~ research interest (Salerno et al., 2015; Perry et al., 2020; Wagnon et al., 2021; King et al., 2020; Thakuri et al., 2014, 2019; Sherpa et al., 2023; Potocki et al., 2022). Its extreme elevation, comprising the highest (Mount Everest) and fourth-highest (Lhotse) mountains on earth, as well as the highest-elevation glacier combined with its thriving tourism and mountaineering industry, enable altitudinal gradients in meteorological quantities to be assessed across the entire elevational range of HMA glaciers; drive high demand for accurate weather observations and forecasts (due to the high exposure of trekkers and mountaineers); and provide relatively strong logistical support for fieldwork.

Compared to elsewhere in HMA, the Khumbu region has already been relatively well instrumented with ~~a~~ Automatic ~~w~~Weather ~~s~~Stations (AWSs). For instance, an AWS was deployed at Pyramid (5,035 m asl) by EvK2CNR in 1994, and their network was later extended up to South Col (in 2008) (Salerno et al., 2015, 2025). ~~In 2010, the French National Research Institute for Sustainable Development ) (Salerno et al., 2015). Since 2010, The Institute for Sustainable Development~~ (IRD France; GLACIOCLIM) and Nepalese partners began long-term ~~weather~~ measurements at Mera glacier up to 6,352 m asl and up to 5,450 m asl along the Changri Nup glacier nearby Khumbu Glacier (Figure 1; Khadka et al., 2022). On Everest's northern slopes, AWSs were installed at the North Col (7,028 m asl) and Ruopula Pass (6,560 m asl) (Yang et al., 2011) though these ceased operation by 2008 and 2010, respectively. More recently, Chinese scientists reportedly installed eight AWSs up to near the summit but these data remain unpublished (The Himalayan Times, 2022) and the state of the network remains unknown.

Despite significant investments by numerous research groups, critical gaps persist in understanding HMA cryospheric processes, climate change signals, and data dependability. Weather forecasts for peaks above 6,000 m asl, ~~which can have—of life-or-and-death consequences for relevance for the region’s many~~ mountaineers ~~(Matthews et al., 2020a)~~, also remain highly uncertain ~~due to the~~because of lack of ground-~~validated-based~~ observations for validation (Matthews et al., 2020a). Furthermore, a scarcity of high-quality, elevation-dependent datasets hinders efforts to refine regional climate models and downscale reanalysis products, which is necessary to understand longer-term, larger-scale climate change in this region (Khadka et al., 2022).

During the 2019–2025 period, our team of researchers installed and maintained AWSs in six different locations along Everest’s southern slopes as part of the National Geographic and Rolex Perpetual Planet Everest Expeditions. These stations have since generated novel insights, including: seasonal oxygen availability at high elevations and climate-influenced changes in Everest’s physiological summit height (Matthews et al., 2020b); sublimation dynamics at South Col Glacier (Sherpa et al., 2023); seasonal variability in cloud cover and visibility (Grey et al., 2022); extreme solar irradiance patterns (Matthews et al., 2020a); energy balance studies (Matthews et al., 2020a; Potocki et al., 2022; Brun et al., 2023); and monsoon behaviour at extreme altitudes (Perry et al., 2020; Khadka et al., 2021).

In this paper, we focus on describing the latest available open-access data (<https://doi.org/10.5281/zenodo.18849098>; Khadka, 2026). We first provide a summary of the network (Section 2), before outlining quality control (QC, Section 3). We ~~conclude close (Section 4)~~ by summarizing the seasonal climatology at stations with the most complete records (Section 4), the correspondence between AWS-South Col AWS observations and ERA5 ~~at the South Col~~(Section 5), and by sharing new insights from the highest stations with more intermittent records.

## 2. Observation area and ~~d~~Data

The geographical locations of the different AWSs are shown in Figure 1. The five stations were initially installed in the primary Khumbu valley from Phortse (3,810 m asl; hereafter ~~referred as~~-PH), Base Camp (5,315 m asl; hereafter ~~referred as~~-BC), Camp II (6,464 m asl; hereafter ~~referred as~~-CII), South Col (7,945 m asl; hereafter ~~referred as~~-SC), and Balcony (8,430 m asl; hereafter ~~referred as~~-BA) during the 2019 expedition (Matthews et al., 2020a). The BA station was toppled by extreme winds and anchor failure in January 2020, so in May 2022 a team of researchers and Sherpas returned to install a new AWS at Bishop Rock (8,810 m asl; hereafter ~~referred as~~-BR), remove the BA station, and upgrade the ~~station at the~~-SC AWS (Matthews et al., 2022).

Data have been recorded using Campbell ~~S~~scientific CR1000X dataloggers, at intervals ranging from 10 minutes to daily. However, here we present quality-controlled hourly data-sets in ~~University~~-Coordinated Universal Time (UTC). The respective variables and sensors include air temperature (T), relative humidity (RH), wind speed (WS), radiation variables (~~I~~incoming (in) and outgoing (out) shortwave (SW) and long wave (LW) radiation), pressure (pPa), and precipitation (P) (Table 1). We also include data from experimental pitot wind sensors installed at the SC and BR in 2022.



Figure 1. Geographical location of different automatic weather stations installed during the 2019-2022 National Geographic (NGS) and Rolex Perpetual Planet Everest expeditions, together with operational EvK2CNR and GLACIOCLIM stations in the Khumbu region of Nepal. Black symbols indicate the National Geographic Society AWSs, while the other symbols indicate EvK2CNR and GLACIOCLIM stations. Plus signs indicate important locations and camps, including Everest Base Camp. Changri Nup indicates the area where meteorological and glaciological observations have been conducted since 2010. The red line shows the standard climbing route to Mount Everest from the southern, Nepali side, in the Khumbu region of Nepal. The plus symbol represents camps in the valley including Everest Base Camp and the Changri Nup point is the area where Meteorological glaciological observation has been monitored since 2010 and the red lines represents the normal climbing route to Mount Everest from South side.

## 2.1 AWS details and measured variables

Two types of tripods were used for the AWSs: the Campbell Scientific CM106B, which has an adjustable height of between 2.1 and 3 m, was used for the lower three (PH, BC, and CII) AWSs. The tripod was affixed with a datalogger box, solar panel, various sensors and a grounding rod and lightning-protection system (Table 1, Figures S1 and S2). An antenna for grounding to protect against lightning (Table 1, see Supplementary Information for photographs). An OTT HydroMet Pluvio<sup>2</sup> with a double-A-shield was installed separately at the PH and BC AWSs to measure precipitation. For the upper three (SC, BA and BR) AWSs, a 7-kg custom aluminum tripod designed by Campbell Scientific specifically for Everest was used (Matthews et al., 2020a). The PH and BC AWSs were secured in the soil with stakes pushed 0.5–1 m into the ground, while the other higher elevation AWSs were bolted on to rocks and also guyed to additional anchor points. For the lower elevation AWSs (PH, BC, and CII; < 6,500 m asl), the batteries were installed inside housed within the data-logger box enclosure mounted on the and on a tripod. For the upper elevation AWSs (SC, BA, BR; > 6,500 m asl), the batteries were installed in a separate insulated box to provide protection protect it from very low temperatures, and both Both the data-logger and battery boxes were

separately bolted to the rock to reduce wind drag on the tripod. The overall weight of the lower elevation AWSs is approximately 60 kg and the upper elevation AWSs weigh 52 kg, including the additional pitot tube (~2 kg) and other sensors added in 2022 (Matthews et al., 2020a, 2022). The BA AWS was found to have been toppled during the 2020 maintenance visit, possibly due to extreme wind gusts; however, the other tripods remain in position at the time of writing. Below, we detail provide measurement details for the respective meteorological variables. This information is also summarised in Table 1.

#### **2.1.1 Air Temperature and Relative Humidity**

Air temperature (°C) and relative humidity (%) ~~are were~~ measured using a Vaisala HMP155A-L5-PT sensor installed in a naturally ventilated METSPEC 14-plate solar radiation shield. Additionally, a CS109 temperature probe, ventilated by a METSPEC 6-plate solar radiation shield, was installed for air temperature measurement at each station in 2019. While both shield types reduce radiative heating, the absence of active aspiration, particularly for the less protected CS109 with its ~~6six~~ 6-plate shield may introduce a positive diurnal temperature bias. In 2022, however, only dual Vaisala HMP-155A-L5-PT sensors with METSPEC 14-plate solar radiation shields were installed in the SC and BR AWSs. ~~The sensors are installed 2 m from the ground for the PH, BC and CII AWSs, and 1.5 m for the higher AWSs. The sensors were installed 2 m from the ground for the lower elevation AWSs (PH, BC, and CII), and 1.5 m for the higher elevation AWSs. We chose a lower height for the high-altitude stations to protect them from strong winds in two ways. First, wind speeds are generally lower closer to the ground. Second, a shorter pole creates less leverage (torque) on the tripod base, which reduces the risk of tripod failure during severe wind gusts.~~

#### **2.1.2 Wind Speed and Direction**

Initially, in 2019, the R. M. Young 05108-45 ~~anemometers were was~~ ~~in-at~~ used at all AWSs for measuring wind speed and direction. The SC and BA ~~AWSs were each equipped with two had two of these~~ sensors for redundancy. However, all four R. M. Young 05108-45s failed in the extreme conditions encountered at the BA and SC. We therefore replaced the ~~two double~~ 05108-45s at the SC with a new polycarbonate version of that sensor; we also added a Richards C5C anemometer and Pitot tube, custom made by the Mount Washington Observatory. This same sensor array for measuring wind speed was used at the BR station ~~following the failure after the demise~~ of the BA AWS (Matthews et al., 2022). All wind sensors were installed 2.3 m above the ground surface for lower AWSs (PH, BC, and CII) and ~2 m for higher elevation AWSs (SC, BA, and BR).

#### **2.1.3 Air Pressure**

The ~~measurement of~~ barometric air pressure (hPa) at PH ~~was measured using a Vaisala PTB110, whereas the Vaisala PTB210 was used at the other AWSs because of the lower barometric air pressure at higher elevations. At all stations, the pressure sensors were housed inside the datalogger enclosure. At the three lower-elevation AWSs (PH, BC, and CII), the enclosure was mounted approximately 1–1.5 m above the ground. is achieved using the Vaisala PTB-110, whilst the Vaisala PTB210 is employed in all other AWSs as higher AWSs weather is comparatively harsh with lower pressure. The sensors are installed within the data logger box of all AWSs. However, the data logger box of the three lower AWSs (PH, BR, and CII) is located on the tripod mast, at a distance of 1–1.5 m above the ground surface.~~

#### **2.1.4 Radiation**

~~The four radiation components were measured using paired Hukseflux SR30 pyranometers for incoming and outgoing shortwave radiation and paired Hukseflux IR01 pyrgeometers for incoming and outgoing longwave radiation at PH. An Apogee SN-500-SS net radiometer was used at CII, while Hukseflux NR01 net radiometers were used at the other AWSs. The instruments were mounted on crossarms approximately 1.5 m above the surface (Figures S1 and S2). The measurement of the four components of radiation is facilitated by the Hukseflux SR30 (up/down shortwave radiation); Hukseflux IR20 (up/down longwave radiation) at PH; Apogee SN-500-SS at CII; and Hukseflux NR01 at other AWSs using the cross arm (Figure~~

S1, S2). The radiation sensors are mounted on crossarms at approximately 1.5 m above the surface at all locations.

### 2.1.5 Precipitation

At the PH and BC AWSs, the weighing OTT Pluvio<sup>2</sup> precipitation gauges equipped with double-A-tilt wind shields, with a height of 1.5 m (from the surface at the mouth of Pluvio) were installed at the PH and BC AWSs. At both sites the OTT Pluvio<sup>2</sup> is/were partially filled with an environmentally friendly antifreeze agent/solution, which is replaced regularly when the reservoir is emptied during regular maintenance visits. An optical disdrometer (OTT Parsivel<sup>2</sup>) and a snow-level sonic sensor (SR50A) were installed to complement the precipitation measurements. These datasets are included in the data archive but are not analysed in the present study because they require specialised quality-control procedures, including the removal of artefacts caused by wind and blowing snow (Table 1; Löffler-Mang and Joss, 2000; Ryan et al., 2008; Pérez-Tello et al., 2026). These datasets will be analysed in future work.

### 2.1.6 Time Lapse Photo

In addition to the meteorological measurements at the BC AWS, a Campbell Scientific CCFC fField Ccamera was installed looking-facing eastward towards the summit of Mount Everest. The camera captures photographs two times a day; at 0937 Nepal time (NPT) (0352 UTC) and at 1437 NPT (0852 UTC), respectively (Grey et al., 2022).

## 2.2 Real-time data transfer

At the beginning of the project in 2019, (2019) all the stations (except PH) were configured to transmit real-time data using the Thuraya satellite network, whereas, the PH station used Inmarsat telemetry. Data transmission from PH ceased following a battery failure in September 2024. Separately, the failure of the Thuraya 3 satellite in April 2024 caused increasingly intermittent connectivity at the stations using Thuraya, followed by a complete loss of transmission. After the failure of the Thuraya 3 satellite in April 2024, connectivity became intermittent, and then stopped altogether. In 2025, we therefore moved BC and CII to the Iridium Short Burst Data (SBD) service, as used by other remote networks (e.g., on the Greenland Ice Sheet: Fausto et al., 2021). The SC station was not converted to Iridium SBD in 2024 due to logistical, financial, and time constraints and therefore limited data are only currently available via data via direct download. The PH station was originally configured to transfer data using the network but was switched to the Nepal Telecom 4G network in 2025<sup>4</sup>. Though only limited variables are transmitted through SBD, the new data transmitted from SBD and BGAN-INMARSAT are not included in the current Zenodo <https://doi.org/10.5281/zenodo.18849098> archive (Khadka, 2026). Additional data will be added to the repository following retrieval, quality control, and validation. The new data will be added after the next fieldwork probably in spring 2026.

203 Table 1. ~~AWSS~~ Locations of ~~AWSS~~, data availability, sensors and their height from the surface during the installation (updated  
204 from ~~Matthews~~ et al. (2020a)).

|                                                 | Phortse (PH)                                                                                     | Base Camp (BC)                                                | Camp II (CIH)                                        | South Col (SC) 2019                             | South Col (SC) 2022                                                    | Balcony (BA)                                     | Bishop Rock (BR)                                                       |
|-------------------------------------------------|--------------------------------------------------------------------------------------------------|---------------------------------------------------------------|------------------------------------------------------|-------------------------------------------------|------------------------------------------------------------------------|--------------------------------------------------|------------------------------------------------------------------------|
| <b>Lat, Lon, and Elevation</b>                  | 27.8456°N<br>86.7472°E<br>3,810 m                                                                | 27.9952°N<br>86.8406°E<br>5,315 m                             | 27.9810°N<br>86.9023°E<br>6,464 m                    | 27.9719°N<br>86.9295°E<br>7,945 m               | 27.9719°N<br>86.9295°E<br>7,945 m                                      | 27.9826°N<br>86.9292°E<br>8,430 m                | 27.98805°N<br>86.92521°E<br>8,810 m                                    |
| <b>Data availability</b>                        | April 2019 to <del>April 2025</del> <sup>present</sup>                                           | October 2019 to <del>April 2025</del> <sup>present</sup>      | May 2019 to <del>April 2025</del> <sup>present</sup> | June 2019 to July 2021                          | May 2022 to <del>April 2025</del> <sup>present</sup>                   | May 2019 to January 2020                         | May 2022 to July 2023                                                  |
| <b>Sensor level above ground (m)</b>            | 2 m for temperature and humidity, 2.3 m for wind                                                 | 2 m                                                           | 2 m                                                  | 1.5 m for temperature and humidity, 2m for wind | 1.5 m for temperature and humidity, 2m for wind                        | 1.5 m for temperature and humidity, 2 m for wind | 1.5 m for temperature and humidity, 2 m for wind                       |
| <b>Air temperature sensor</b>                   | Vaisala HMP155A-L5-PT; CS109                                                                     | Vaisala HMP155A-L5-PT; CS109                                  | Vaisala HMP155A-L5-PT; CS109                         | Vaisala HMP155A-L5-PT; CS109                    | 2 × Vaisala HMP155A-L5-PT                                              | Vaisala HMP155A-L5-PT; CS109                     | 2 × Vaisala HMP155A-L5                                                 |
| <b>Relative humidity sensor</b>                 | Vaisala HMP155A-L5-PT                                                                            | Vaisala HMP155A-L5-PT                                         | Vaisala HMP155A-L5-PT                                | Vaisala HMP155A-L5-PT                           | 2 × Vaisala HMP155A-L5-PT                                              | Vaisala HMP155A-L5-PT                            | 2 × Vaisala HMP155A-L5-PT                                              |
| <b>Wind speed and direction sensor</b>          | R.M. Young 05108-45                                                                              | R.M. Young 05108-45                                           | R.M. Young 05108-45                                  | 2 × R.M. Young 05108-45                         | Polycarbonate R.M. Young 05108-45, Richards C5C anemometer, Pitot tube | 2 × R.M. Young 05108-45                          | Polycarbonate R.M. Young 05108-45, Richards C5C anemometer, Pitot tube |
| <b>Air pPressure sensor</b>                     | Vaisala PTB 110                                                                                  | Vaisala PTB210                                                | Vaisala PTB210                                       | Vaisala PTB210                                  | Vaisala PTB210                                                         | Vaisala PTB210                                   | Vaisala PTB210                                                         |
| <b>Radiation sensor</b>                         | 2 × Hukseflux SR30 (up/down shortwave radiation); 2 × Hukseflux IR20 (up/down thermal radiation) | Hukseflux NR01                                                | Apogee SN-500-SS                                     | Hukseflux NR01                                  | Hukseflux NR01                                                         | Hukseflux NR01                                   | Hukseflux NR01                                                         |
| <b>Precipitation sensor</b>                     | OTT Pluvio <sup>2</sup> and double- <del>A</del> alter shield                                    | OTT Pluvio <sup>2</sup> and double- <del>A</del> alter shield | -                                                    | -                                               | -                                                                      | -                                                | -                                                                      |
| <b>Time Lapse pPhoto</b>                        | -                                                                                                | Campbell Scientific Canada CCFC Field Camera                  | -                                                    | -                                               | -                                                                      | -                                                | -                                                                      |
| <b>Present weather sensor</b>                   | OTT Parsivel <sup>2</sup>                                                                        | OTT Parsivel <sup>2</sup>                                     | -                                                    | -                                               | -                                                                      | -                                                | -                                                                      |
| <b>Relative surface elevation change sensor</b> | Campbell Scientific SR50A                                                                        | Campbell Scientific SR50A                                     | Campbell Scientific SR50A                            | -                                               | -                                                                      | -                                                | -                                                                      |
| <b>Data Logger</b>                              | Campbell Scientific CR1000X                                                                      | Campbell Scientific CR1000X                                   | Campbell Scientific CR1000X                          | Campbell Scientific CR1000X                     | Campbell Scientific CR1000X                                            | Campbell Scientific CR1000X                      | Campbell Scientific CR1000X                                            |
| <b>Logger enclosure</b>                         | Standard Campbell Scientific                                                                     | Standard Campbell Scientific                                  | Standard Campbell Scientific                         | Pelican case with military-spec quick-connects  | Pelican case with military-spec quick-connects                         | Pelican case with military-spec quick-connects   | Pelican case with military-spec quick-connects                         |

|                           |                                                                                                                                                       |                                                                                                                  |                                                                                                            |                                                                                                                                                              |                                                                                                                                                               |                                                                                                                                                              |                                                                                                                                                               |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Batteries</b>          | 24 Ah                                                                                                                                                 | 3 × 8Ah                                                                                                          | 3 × 8Ah                                                                                                    | 3 × 8Ah                                                                                                                                                      | 3 × 8Ah                                                                                                                                                       | 3 × 8Ah                                                                                                                                                      | 3 × 8Ah                                                                                                                                                       |
| <b>Charging</b>           | 2 × 20_W solar panels                                                                                                                                 | 2 × 20_W solar panels                                                                                            | 2 × 20_W solar panels                                                                                      | 2 × 10_W solar panels                                                                                                                                        | 3 × 10_W solar panels                                                                                                                                         | 2 × 10_W solar panels                                                                                                                                        | 3 × 10_W solar panels                                                                                                                                         |
| <b>Telemetry</b>          | Inmarsat, Sierra wireless with NTC <del>simSIM</del>                                                                                                  | Thuraya; 400 MHz radio, Iridium                                                                                  | Thuraya; 400 MHz radio, Iridium                                                                            | Thuraya; 400 MHz radio                                                                                                                                       | Thuraya; 400 MHz radio                                                                                                                                        | Thuraya; 400 MHz radio                                                                                                                                       | Thuraya; 400 MHz radio                                                                                                                                        |
| <b>Sampling interval</b>  | 3 s (wind); 60 s (radiation, air pressure, temperature, relative humidity, precipitation) 3600 s (present weather, relative surface elevation change) | 5 s (wind); 60 s (temperature, relative humidity, precipitation); 600 s (air pressure); 3600 s (present weather) | 60 s (temperature, relative humidity, wind); 600 s (radiation); 3600 s (relative surface elevation change) | 5 s (wind); 60 s (temperature, relative humidity, radiation); 600 s (pressure)                                                                               | 5 s (wind); 60 s (temperature, relative humidity); 600 s (pressure)                                                                                           | 5 s (wind); 60 s (temperature, relative humidity); 600 s (pressure)                                                                                          | 5 s (wind); 60 s (temperature, relative humidity); 600 s (pressure)                                                                                           |
| <b>Tripod</b>             | Campbell Scientific CM106B                                                                                                                            | Campbell Scientific CM106B                                                                                       | Campbell Scientific CM106B                                                                                 | Custom Aluminum                                                                                                                                              | Custom Aluminum                                                                                                                                               | Custom Aluminum                                                                                                                                              | Custom Aluminum                                                                                                                                               |
| <b>Approximate Weight</b> | <del>Not available_</del>                                                                                                                             | <del>Not available_</del>                                                                                        | <del>Not available_</del>                                                                                  | Total: 50 kg<br>Pelican case with logger: 8 kg<br>Pelican case with batteries: 16 kg<br>Tripod: 7 kg<br>Crossarms, mounts, and bolts: 11 kg<br>Sensors: 8 kg | Total: 52 kg<br>Pelican case with logger: 8 kg<br>Pelican case with batteries: 16 kg<br>Tripod: 7 kg<br>Crossarms, mounts, and bolts: 11 kg<br>Sensors: 10 kg | Total: 50 kg<br>Pelican case with logger: 8 kg<br>Pelican case with batteries: 16 kg<br>Tripod: 7 kg<br>Crossarms, mounts, and bolts: 11 kg<br>Sensors: 8 kg | Total: 52 kg<br>Pelican case with logger: 8 kg<br>Pelican case with batteries: 16 kg<br>Tripod: 7 kg<br>Crossarms, mounts, and bolts: 11 kg<br>Sensors: 10 kg |

205

### 206 3. Data treatment and quality control procedure

207 Data gaps, sensor failures, and erroneous readings were common due to factors such as riming (icing of  
208 sensors), heavy snowfall partially or completely burying the upper stations, battery depletion, extreme  
209 temperature fluctuations, and communication failures. Acknowledging these challenges, we implemented,  
210 a multi-stage quality control (QC) procedure to identify and flag spurious data points ([Figure 2](#)).

211 Our QC protocol takes a tiered approach, progressing from simple flagging to more complex, variable-  
212 specific corrections. The procedures applied to the raw dataset are as follows:

- 213 • **Data flagging:** Where available, manufacturer-provided diagnostic data (Table S-1) and internal  
214 sensor flags were used to identify and filter out physically implausible or erroneous observations.
- 215 • **Relative humidity correction:** Physically, RH cannot exceed 100% under stable conditions. Any  
216 instances in which RH was greater than 100% were assumed to be caused by sensor uncertainty or  
217 rapid cooling events, and were capped at 100%. Artificially low ~~values were also present in the raw~~  
218 ~~measurements~~ ~~RH in our raw measurements were also present~~ due to the logger program calculating  
219 RH with respect to water rather than ice when air temperatures were below 0 °C. We therefore  
220 corrected these instances by computing vapour pressure (*e*) from measured RH, and saturation

vapour pressure ( $e_s$ ) with respect to ice using the air temperature (Buck, 1981, 1996). Corrected RH was hence  $100 \times (e/e_s)$ .

- **Wind sensor freezing detection:** Periods of zero wind speed coupled with zero wind direction variability are a strong indicator of sensor freezing (e.g., encased in rime ice). These data points were identified and removed from the dataset rather than being treated as valid calm conditions.
- **Shortwave radiation corrections:**
  - **Night time values:** All incoming shortwave radiation below  $7 \text{ W m}^{-2}$  (a common threshold for sensor uncertainty during night time) were set to zero (Shea et al., 2015; Khadka et al., 2022).
  - **Albedo-based correction:** Fresh snowfall or deposition can accumulate on the upward-facing pyranometer, temporarily capping or shading the sensor. This obstruction leads to a substantial underestimation of incoming shortwave radiation (SWin), while the downward-facing sensor continues to accurately measure the outgoing shortwave radiation (SWout) of the fresh snow surface. Consequently, the calculated surface albedo ( $\alpha = \text{SWout}/\text{SWin}$ ) can artifactually exceed realistic threshold values for snow (typically  $\alpha > 0.95$ ). To address this instrumental artifact and ensure strict quality control, we adopted the correction protocol developed by Shea et al. (2015) to adjust the compromised SWin values during these specific cases. Specifically, for periods Fresh snowfall can cause the surface albedo ( $\alpha = \text{SWout} / \text{SWin}$ ) to exceed a realistic value for snow (typically  $\alpha \leq 0.95$ ). For periods where the calculated albedo exceeded 0.95, it was assumed that SWin was underestimated (e.g., due to sensor riming or a tilted sensor). SWin was recalculated as  $\text{SWout}/0.95$  to provide a more realistic estimate (Shea et al., 2015; Khadka et al., 2022)
- **Precipitation:** The Pluvio<sup>2</sup> weighing precipitation gauges were equipped with double-A-shield shields to minimize wind-induced under-catch (Figure S1). Only positive accumulation records were considered. No correction for wind under-catch was applied in this dataset; users are advised to apply such corrections using established methods (e.g., Kochendorfer et al., 2022; Graves et al., 2026)(e.g.,).




Figure 2. Periods of data availability for the principal sensors at each AWS from installation until May 2025. Solid horizontal lines indicate periods for which quality-controlled observations are available. Time periods when stations/sensors were operational since the beginning of installation to the last field visit in May 2025. The solid lines represent the period of available data.

## 4. Climatology of the region

Figure 3 presents the annual cycles of shows the monthly precipitation and temperature at PH and BC for December 2019–November 2022, together with monthly mean, minimum, and maximum temperatures based on all available observations at CII, SC, BA, and BR (from December 2019 to November 2022; Figure 2) for PH (3,840 m) and BC (5,315 m), the two AWSs in the network that include measurements of precipitation. Whilst this is a very short period for understanding climate variability, there is a clear seasonal pattern in precipitation and temperature at both sites. Across the network, the lowest temperatures generally occur during winter (December–February). Temperatures then rise progressively during the pre-monsoon season (March–May), coinciding with increasing incoming shortwave radiation and longer daylight duration (Figure S4). The annual precipitation is 790 and 629 mm, with 72 and 77% of this falling during the monsoon at the PH and BC AWSs, respectively. The average mean annual temperature at PH is was 4.1 °C at PH, BC is -3.1 °C at BC, and CII is -10.2 °C at CII. The with mean monthly temperature amplitude of temperature is was 6.7 °C (max 8.0 °C in March, min 4.6 °C in July) at PH, 6.5 °C (max 7.9 °C in March, min 4.9 °C in December) at BC and 6.4 °C (max 7.8 °C in March, min 3.6 °C in December) at CII, respectively. At SC, July was the warmest month, with a mean temperature of -12.2 °C and a mean diurnal range of 10.2 °C. February was the coldest month, with a mean temperature of -29.7 °C and a mean diurnal range of 4.4 °C. Precipitation at PH and BC also clearly delineates the seasonal cycle: winter is predominantly dry, precipitation gradually increases during the pre-monsoon, and the largest amounts occur during the June–September monsoon season, which accounts for 72% and 77% of the annual precipitation at PH and BC, respectively. Precipitation subsequently declines during the post-monsoon period as cooler and drier winter conditions become established.




Figure 3. Mean annual cycle of monthly precipitation (snow and total precipitation in mm liquid water equivalent, histograms) and monthly mean, minimum and maximum air temperature ( $T$ , lines) during December 2019–November 2022 at PH, BC, and CII AWSs

(a-c) and monthly mean, minimum and maximum temperature for all the available period for SC, BA and BR AWSs (d-f). The error bars represent the monthly standard deviation for all different variables and BA error bars are omitted because of the limited number of observations.




Figure 3. Mean annual cycle of monthly precipitation (snow and total precipitation in mm liquid water equivalent, histograms) and mean monthly mean, minimum and maximum air temperature ( $T$ , lines) during December 2019–November 2022 at PH, BC and CII AWSs. The bar represents the standard deviation for all different variables.

Seasonal mean values, and elevation gradients between the PH and BC AWSs (separated by  $\sim 1,500$  m) also reveal some interesting characteristics (Table 2). While these linear elevation gradients provide a useful baseline, it is important to note that the precipitation gradient across this elevational range is fundamentally non-linear, with the rate of change varying significantly between specific altitudes and locations. When calculated as a single linear gradient across this elevational range, the mean annual precipitation gradient was  $-107 \text{ mm km}^{-1}$ , but it declines by almost 50 % during the monsoon to  $-56 \text{ mm km}^{-1}$ . This contrast indicates that while the monsoon brings the highest total volume of water, non-monsoonal seasons (winter, pre-, and post-monsoon) feature much more dramatic relative precipitation drops with increasing elevation. The precipitation gradients average  $-107 \text{ mm km}^{-1}$  ( $-17 \% \text{ km}^{-1}$ ) average over three years, but it declines by almost 50 % during the monsoon (to  $-56 \text{ mm km}^{-1}/12 \% \text{ km}^{-1}$ ). Overall, the amount of precipitation measured at the both (PH and BC) observing sites (PH and BC) is/were higher than those reported e amount measured at the Pheriche (4,200 m asl; 540 mm during November 2016–December 2020) and Pyramid (5,050 m asl; 449 mm during 1994–2012 and 591 mm during November 2016–December 2020) by IRD stations- (Khadka et al., 2022; Salerno et al., 2015)(Khadka et al., 2022). Similarly, between PH and BC, the mean temperature gradient (TG) is  $-4.8 \text{ }^{\circ}\text{C km}^{-1}$  with least negative in winter ( $-4.5 \text{ }^{\circ}\text{C km}^{-1}$ ) and most negative during the pre-monsoon season ( $-5.4 \text{ }^{\circ}\text{C km}^{-1}$ ). When data from four stations with minimal data gaps (PH, BC, CII, and SC) are used to calculate linear and non-linear TGs, the resulting gradients are more strongly negative than those calculated using only the PH and BC AWSs. Both the seasonal and annual TGs derived from all four stations (Table 2) are more negative (pre-monsoon, monsoon, annual:  $-6.5$ ,  $-5.7$ , and  $-6.0 \text{ }^{\circ}\text{C km}^{-1}$ , respectively) than the values previously reported by Salerno et al. (2015) and Graves et al. (2026), calculated using several stations in the Koshi basin between 1,000–8,000 m asl and Dudh Koshi basin between 2,600–8,400 m asl. However, the TG calculated from all four AWSs (Table 2) is comparable at both seasonal and annual scales to the TG reported by Salerno et al. (2015). Crucially, we note that reported non-linearity in the temperature-altitude relationship in the Khumbu region (Khadka et al., 2021) limits the comparability of lapse rates calculated across different elevational ranges. Consequently, we follow Graves et al. (2026) and provide a non-linear equation for the temperature gradients between PH and SC (Table 2; Figure S4). Similarly, between the PH and BC, the mean temperature gradient is  $-4.8 \text{ }^{\circ}\text{C km}^{-1}$  with minimum  $-4.5 \text{ }^{\circ}\text{C km}^{-1}$  during the winter and maximum  $-5.4 \text{ }^{\circ}\text{C km}^{-1}$  during the Pre-monsoon. Both

seasonal and annual temperature gradients (Table 2) are also less negative than reported previously (pre-monsoon, monsoon, annual:  $-6.5$ ,  $-5.7$ ,  $-6.0$   $^{\circ}\text{C km}^{-1}$ , respectively) by Salerno et al. (2015), calculated using several stations on Koshi basin between  $\sim 1,000$ – $8,000$  m asl. However, we note that reported non-linearity in the temperature-altitude relationship in the Khumbu region (Khadka et al., 2021) limits the comparability of lapse rates calculated across different elevational ranges.

Both incoming longwave (LWin) and shortwave radiation (SWin) exhibit strong seasonality and clear variation with altitude (Figure S4). Both LWin and SWin reach their lowest values during winter when temperatures are coldest. As the pre-monsoon begins, rising air temperatures drive a steady increase in LWin, which remains high throughout the monsoon season. Notably, the maximum daily LWin at PH during the monsoon exceeds  $350 \text{ W m}^{-2}$ , while at SC it remains below  $\sim 250 \text{ W m}^{-2}$ . This contrast likely reflects both the lower atmospheric temperature at SC and differences in effective atmospheric emissivity, potentially associated with lower water-vapour content and reduced cloud cover at higher elevations.

SWin is lowest during mid-winter and gradually increases toward the pre-monsoon period. However, the influence of monsoonal cloud cover is weaker at higher-elevation sites (CII and SC) compared to lower-elevation stations (PH and BC), indicating reduced atmospheric moisture at higher altitudes and supporting the presence of a strong negative precipitation gradient with elevation (Graves et al., 2026).

Table 2. Mean seasonal and annual temperature ( $^{\circ}\text{C}$ ) and precipitation (P, mm) at PH and BC AWSs and the calculated linear Temperature and Precipitation gradients (TG and PG respectively), and the non-linear TG equation derived across the stations. \*z represents the altitude in kilometres (km) used to calculate the non-linear TG.

| Season                 | Base Camp (BC)                     |                    | Phortse (PH)                       |                    | TG from PH and BC AWSs ( $^{\circ}\text{C km}^{-1}$ ) | TG from PH, BC, CII and SC AWSs ( $^{\circ}\text{C km}^{-1}$ ) | TG non-linear equation from PH, BC, CII and SC AWSs ( $^{\circ}\text{C km}^{-1}$ ) | PG from PH and BC AWSs (mm $\text{km}^{-1}$ / % $\text{km}^{-1}$ ) |
|------------------------|------------------------------------|--------------------|------------------------------------|--------------------|-------------------------------------------------------|----------------------------------------------------------------|------------------------------------------------------------------------------------|--------------------------------------------------------------------|
|                        | Temperature ( $^{\circ}\text{C}$ ) | Precipitation (mm) | Temperature ( $^{\circ}\text{C}$ ) | Precipitation (mm) |                                                       |                                                                |                                                                                    |                                                                    |
| Winter (Dec-Feb)       | -8.8                               | 27                 | -2.1                               | 56                 | -4.5                                                  | -6.5                                                           | $-0.75z^2 + 2.4z - 0.7$                                                            | -20/-73                                                            |
| Pre-Monsoon (Mar-May)  | -4.5                               | 87                 | 3.6                                | 115                | -5.4                                                  | -6.7                                                           | $-0.5z^2 - 0.75z + 13.8$                                                           | -19/-22                                                            |
| Monsoon (Jun-Sept)     | 2.4                                | 484                | 9.7                                | 567                | -4.8                                                  | -5.6                                                           | $-0.26z^2 - 2.53z + 23.2$                                                          | -56/-12                                                            |
| Post-Monsoon (Oct-Nov) | -3.1                               | 31                 | 3.9                                | 51                 | -4.6                                                  | -6.6                                                           | $-0.78z^2 + 2.56z + 5.4$                                                           | -13/-44                                                            |
| Annual                 | -3.1                               | 629                | 4.1                                | 790                | -4.8                                                  | -6.3                                                           | $-0.57z^2 + 0.42z + 10.4$                                                          | -107/-17                                                           |

Table 2. Mean seasonal and annual Temperature ( $^{\circ}\text{C}$ ) and precipitation (P, mm) at PH and BC AWSs and the calculated Temperature and precipitation gradients (TG and PG) between these AWSs.

| Season       | Base Camp (BC)                     |                    | Phortse (PH)                       |                    | TG ( $^{\circ}\text{C km}^{-1}$ ) | PG ( $\text{mm km}^{-1}$<br>$\pm\% \text{ km}^{-1}$ ) |
|--------------|------------------------------------|--------------------|------------------------------------|--------------------|-----------------------------------|-------------------------------------------------------|
|              | Temperature ( $^{\circ}\text{C}$ ) | Precipitation (mm) | Temperature ( $^{\circ}\text{C}$ ) | Precipitation (mm) |                                   |                                                       |
| Winter       | -8.8                               | 27                 | -2.1                               | 56                 | -4.5                              | -20/ 73                                               |
| Pre-Monsoon  | -4.5                               | 87                 | 3.6                                | 115                | -5.4                              | -19/ 22                                               |
| Monsoon      | 2.4                                | 484                | 9.7                                | 567                | -4.8                              | -56/ 12                                               |
| Post-Monsoon | -3.1                               | 31                 | 3.9                                | 51                 | -4.6                              | -13/ 44                                               |
| Annual       | -3.1                               | 629                | 4.1                                | 790                | -4.8                              | -107/ 17                                              |

## 5. Comparison with ERA5

Monitoring the meteorology at the highest altitudes in the Himalaya at elevations where glaciers are accumulating mass, and where mountaineers are most at risk from extreme weather provides the greatest test to observational efforts. Accordingly, there is considerable interest in using reanalysis data to extend brief in-situ records. However, to facilitate such extensions, it is necessary to establish the extent to which such reanalysis products can reproduce the observed meteorology (Khadka et al., 2022). ERA5, the fifth-generation global reanalysis produced by ECMWF, provides hourly atmospheric variables from 1940 onward at multiple pressure levels and at relatively high spatial resolution and is widely used as a reference dataset (Hersbach et al., 2020). While adjacent higher AWSs (BR and BA) suffer from highly limited data availability, SC AWS has longer observational record and remains operational. This is also the location of the highest-altitude camp on the main southern (Nepalese) Everest climbing route, where critical resources (e.g., tents and oxygen) are stored, and where climbers can spend considerable time. Consequently, we focus on comparisons between temperature, relative humidity and wind speed between the SC AWS and ERA5 reanalysis (Hersbach et al., 2020) data extracted from the nearest grid point at the 350 hPa pressure level (most representative of the SC: mean air pressure 2019-2021 = 377 hPa). Our comparison is limited to air temperature—relevant for cold stress and with relatively high data completeness at the BA and BR sites. Here, we therefore compare temperature, relative humidity and wind speed between the SC AWS (the longest record from above Camp II, where extreme weather poses the greatest risk for mountaineers) and ERA5 reanalysis (Hersbach et al., 2020) data from the nearest grid point at the 350 hPa pressure level (most representative of the SC: mean air pressure 2019-2021 = 377 hPa).





Figure 4. Hourly air temperature, relative humidity (with respect to water) and wind speed from the SC AWS and ERA5 350 hPa pressure level data (from the nearest grid point to the SC AWS) for the available data period of 2019 (a,b,c) and 2022 (d,e,f). The solid black and red lines represent the 10-day running means of relative humidity and the blue dots represent the observed daily maximum wind speed.



**Figure 4.** Hourly air temperature, relative humidity (with respect to water) and wind speed from the SC AWS and ERA5 350 hPa pressure level data (from the nearest grid point to the SC AWS) for the available data period of 2019 (a,b,c) and 2022 (d,e,f). The solid dark black and red relative humidity line represents the 10-days running mean values.

During both 2019 and 2022, air temperature exhibited reasonably close co-variation between the SC observations and ERA5, with an  $R^2 > 0.6$  (Figure 4 and Table 3). ERA5 systematically underestimates air temperature relative to ~~measurement~~the observations, although some of that reflects the higher altitude (lower air pressure) at 350 hPa relative to the SC data. Diurnal variability is also considerably greater in the SC dataset (Figures S35 and S64), likely reflecting the influences of diurnal fluctuations in the measured surface heat fluxes at SC, which are missing in ERA5 as this (higher altitude) pressure level is representative of the free atmosphere. To further assess the representativeness of ERA5 at high elevations, we compared ERA5 350 hPa temperatures with shorter observational records from BA (mean air pressure = 354 hPa) and BR (mean air pressure = 340.5 hPa). In 2019, the coefficient of determination ( $R^2$ ) between ERA5 and BA found 0.85, while in 2022 the  $R^2$  between ERA5 and BR was 0.56. Despite the shorter datasets, these results indicate that ERA5 temperatures at 350 hPa track observed variability in atmospheric conditions at these very high elevation sites.

The  $R^2$  values for AWS and ERA5 relative humidity were 0.49 and 0.57 in 2019 and 2022; however, the MAE and RMSE for the relative humidity are  $>20\%$ . The 10-day running means of relative humidity from both the AWS observations and ERA5 clearly capture the seasonal transitions, with monsoon onset at SC occurring on 1 July 2019 and 14 June 2022 (Figure 4). The  $R^2$  (0.49 and 0.57) between the AWS and ERA5 relative humidity were adequate for both 2019 and 2022, however the MAE and RMSE for the relative humidity are  $>20\%$ . From the 10 days running mean both AWS and ERA5 relative humidity clearly shows the seasonal transitions. In 2019 and 2022 the monsoon onset was 1<sup>st</sup> July and 14<sup>th</sup> June at SC.

In both years, the wind speed exhibited the highest correlation between SC observations and ERA5 ( $R^2 = 0.70$ ). However, ERA5 consistently overestimates mean wind speeds at the SC, (with MAE = values of 4.8 and 5.43  $\text{m s}^{-1}$  in 2019 and 2022, respectively (Table 3 and Figure 4). This is again consistent with the ERA5 grid point being representative of the higher-altitude free atmosphere; frictional drag at the SC reduces the wind speed. We also observe faster winds in the post-monsoon in 2022 compared to 2019. Based on the generally good agreement between ERA5 reanalysis data and observations at the SC, we anticipate that the reanalysis could be used to gap-fill and extend intermittent meteorological records from Mount Everest's upper slopes—correcting the time-varying biases using empirical-statistical, or machine-learning based approaches (Van Wyk de Vries et al., 2024; Potocki et al., 2022).

Table 3. Values of each statistical metrics ( $R^2$ , MAE, RMSE) obtained by the comparing the hourly SC AWS measured observations and with ERA5 350 hPa temperature (T), relative humidity (RH) and wind speed (WS) at hourly scale for 2019 and 2022.

| Year | Variables                | $R^2$ | MAE  | RMSE |
|------|--------------------------|-------|------|------|
| 2019 | T ( $^{\circ}\text{C}$ ) | 0.67  | 3.7  | 4.1  |
|      | RH (%)                   | 0.49  | 24.6 | 31.4 |
|      | WS ( $\text{m s}^{-1}$ ) | 0.70  | 4.8  | 5.6  |
| 2022 | T ( $^{\circ}\text{C}$ ) | 0.62  | 3.1  | 3.6  |
|      | RH (%)                   | 0.57  | 23.4 | 30.4 |
|      | WS ( $\text{m s}^{-1}$ ) | 0.70  | 5.3  | 6.1  |

## 6. Research insight and importance of the data

Observations from the SC (Figure 4) highlight the potential for mean winds exceeding  $30 \text{ m s}^{-1}$  and gusts above  $60 \text{ m s}^{-1}$  speeds that can be extremely dangerous for mountaineers: at high by increasing the risk of being blown over, and elevated the probability of cold injury (Sherpa et al., 2023; Van Wyk de Vries et al., 2024). Elsewhere, atmospheric pressure data from the AWS network described here have shown that oxygen availability at the summit is highly variable on synoptic timescales, as it depends on the partial pressure of atmosphere. Consequently, the “apparent elevation” of Mt. Everest (i.e., how high the mountain would feel without supplemental oxygen) can vary by almost 750 m, and during the most challenging winter conditions an ascent without supplemental oxygen may be impossible data from the network of AWSs described here has shown oxygen availability at the summit to be highly variable at synoptic timescales, meaning that the “apparent elevation” of Mt. Everest (how high the mountain would feel for those not breathing supplemental oxygen) can vary by almost 750 m, and that during the most challenging winter conditions an ascent without the aid of supplemental oxygen might be impossible (Matthews et al., 2020b). Accurately forecasting wind speeds and oxygen variability for the upper slopes of Mount Everest is therefore of high importance for increasing safety on the mountain. The data from the AWSs provided-presented here provide the foundation to achieve such improvements. At the a minimum, they can be used to assess uncertainty in the freely available forecasts widely used by expeditions (Matthews et al., 2020b); but they can also be used to train model output statistics including novel applications of machine learning to provide improved, hyper-local forecasts for identifying optimal climbing windows (Van Wyk de Vries et al., 2024).

The AWS data shared here can also be extremely useful for understanding cryosphere-climate interactions at the highest elevations within HMA. For instance, the archive has already been used to estimate the surface energy balances at the summit of Mt. Everest (Matthews et al., 2020a), and for the South Col Glacier (at  $\sim 8,000 \text{ m asl}$ ), revealing a high-altitude system that might be acutely sensitive to changes in effective precipitation due to extremely high levels of insolation and thus highly responsive to albedo variations (Matthews et al., 2020a; Potocki et al., 2022; Brun et al., 2023). However, significant uncertainties still remain, mainly because the observational record is short. These could be improved by extending the analysis with downscaled reanalysis products like ERA5 Yet significant uncertainties remain, not least because the

observational record presented here is brief. Understanding here could be improved through extending these observations with downscaled reanalyses products (such as ERA5).

The data here are useful not only for assessing cryosphere-climate interactions at extreme elevations but could be used to improve understanding of surface energy and mass balances within the ablation areas of the region's glaciers, and at lower elevations below. The precipitation measurements, for instance, are particularly high quality, ~~as they were collected behind double-Alter-taken-behind double-Alter-shields shields~~ to reduce under-catch (which likely accounts for the greater accumulations relative to nearby stations without such shields: Pheriche and Pyramid), and hence offer a valuable opportunity to constrain mass inputs. More generally, when used in combination with the generally lower-altitude observations from the GLACIOCLIM and EvK2CNR networks, the data shared here can improve quantification of elevational gradients in key meteorological variables needed in distributed (e.g., glacier or hydrological) modelling assessments across up to approximately five vertical kilometres, and hence the entire glacierised elevational range of the Khumbu region.

Data across such a large elevational range could also provide valuable insight into regional-scale atmospheric dynamics. For instance, the timing and character of the monsoon's arrival—critical ~~in-for~~ determining the annual precipitation totals (Perry et al., 2020), and a key control on melt energy (Khadka et al., 2021; Matthews et al., 2020a) can be assessed right up to the upper accumulation area of the Khumbu ~~G~~glacier. Important further work might improve understanding of the relevant physical drivers behind elevational differences in the monsoon's arrival, how they interact with local anabatic and potential katabatic forcing, and how they might change with further global warming (Salerno et al., 2023).

Ultimately, though, these data might be used in ways far beyond the illustrations above. The upper slopes of ~~Mount-~~ Everest represented the last frontier for human exploration, reached successfully by humans 42 years after the South Pole was attained. Meteorological measurements from the highest reaches only began in 2022 (at the BR). We are still, therefore, at the very beginnings of exploring the weather of this extreme environment, and the list of questions that these data might help answer from disciplines beyond the climate sciences (e.g., biology: Dragone et al., 2023) might conceivably grow rapidly.

## Acknowledgements

This research was supported by the National Geographic and Rolex Perpetual Planet Everest Expeditions and conducted in partnership with Tribhuvan University, with approval from all relevant agencies of the Government of Nepal. We acknowledge Sujatha Bagal and the entire National Geographic Society team for all their support~~conducted in partnership with National Geographic Society (NGS), Rolex, and Tribhuvan University, with approval from all relevant agencies of the Government of Nepal. We acknowledge Sujatha Bagal and all the NGS team for all their support.~~ We also ~~wish to~~ thank the communities of the Khumbu ~~r~~Region, Shangri-La Nepal Treks, Seven Summit Treks, and we further recognize the extraordinary efforts of our entire Sherpa climbing teams since 2019, whose physical and mental prowess at altitude made installation and maintenance of the AWSs possible.

## Authors ~~c~~Contributions

BP, TM, and A~~C~~E planned the initial data collection strategy. BP and TM managed the funding and collaboration with the National Geographic Society and Rolex for the 2019-2025 period. AK, BP, TM, and TGS installed and maintained the AWSs with the Sherpa supports. BP and TM worked closely with Campbell Scientific to design and program the AWSs. AK, BP, and TM contributed to the data analysis, manuscript designing. AK prepared the paper with contributions from all the co-authors.

## Data availability

The quality controlled dataset is available in CSV format from Zenodo (<https://doi.org/10.5281/zenodo.18849098>; Khadka, 2026). Selected data can also be downloaded from the National Geographic Society weather-data portal (<https://www.nationalgeographic.org/projects/perpetual-planet/everest/weather-data/>). The raw disdrometer and all available wind speed data are available at Zenodo. Additionally, the data processing code is available at [https://github.com/arbindrakhadka/AWS\\_QC](https://github.com/arbindrakhadka/AWS_QC) available at: <https://doi.org/10.5281/zenodo.18849098> (Khadka, 2026) moreover the data in csv format and can be downloaded from <https://www.nationalgeographic.org/projects/perpetual-planet/everest/weather-data/>.

## References

- Brun, F., Berthier, E., Wagnon, P., Kääb, A., and Treichler, D.: A spatially resolved estimate of High Mountain Asia glacier mass balances from 2000 to 2016, *Nat. Geosci.*, 10, 668–673, <https://doi.org/10.1038/ngeo2999>, 2017.
- Brun, F., King, O., Réveillet, M., Amory, C., Planchot, A., Berthier, E., Dehecq, A., Bolch, T., Fourteau, K., Brondex, J., Dumont, M., and Mayer, C.: Everest South Col Glacier did not thin during the period 1984 – 2017, *Cryosph.*, 17, 3251–3268, <https://doi.org/10.5194/tc-17-3251-2023>, 2023.
- Buck, A. L.: New Equations for Computing Vapor Pressure and Enhancement Factor, *J. Appl. Meteorol. Climatol.*, 20, 1527–1532, [https://doi.org/10.1175/1520-0450\(1981\)020<1527:NEFCVP>2.0.CO;2](https://doi.org/10.1175/1520-0450(1981)020<1527:NEFCVP>2.0.CO;2), 1981.
- Buck, A. L.: Buck research cr-1a user's manual, Buck Res. Instruments Boulder, CO, USA, 1996.
- Dragone, N. B., Perry, L. B., Solon, A. J., Seimon, A., Seimon, T. A., and Schmidt, S. K.: Genetic analysis of the frozen microbiome at 7900 m a.s.l., on the South Col of Sagarmatha (Mount Everest), *Arctic, Antarct. Alp. Res.*, 55, 2164999, <https://doi.org/10.1080/15230430.2023.2164999>, 2023.
- Fausto, R. S., Van As, D., Mankoff, K. D., Vandecrux, B., Citterio, M., Ahlstrøm, A. P., Andersen, S. B., Colgan, W., Karlsson, N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., Nielsen, S., Pedersen, A., Shields, C. L., Solgaard, A. M., and Box, J. E.: Programme for Monitoring of the Greenland Ice Sheet (PROMICE) automatic weather station data, *Earth Syst. Sci. Data*, 13, 3819–3845, <https://doi.org/10.5194/ESSD-13-3819-2021>, 2021.
- Graves, B., Matthews, T., Perry, B. L., Edwards, T., Taylor, R., Brun, F., Wagnon, P., Khadka, A., and Shrestha, D.: Modelling Accumulation of a High-Altitude Himalayan Glacier, *J. Glaciol.*, 72, e72, <https://doi.org/10.1017/jog.2026.10147>, 2026.
- Grey, L., Johnson, A. V., Matthews, T., Perry, L. B., Elmore, A. C., Khadka, A., Shrestha, D., Tuladhar, S., Baidya, S. K., Aryal, D., and Gajurel, A. P.: Mount Everest's photogenic weather during the post-monsoon, *Weather*, 77, 156–160, <https://doi.org/10.1002/WEA.4184>, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

517 Immerzeel, W. W., Van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the Asian water  
518 towers, *Science* ~~(80-)~~, 328, 1382–1385, <https://doi.org/10.1126/science.1183188>, 2010.

519 Khadka, A.: NGS Mount Everest AWS Network, Zenodo, <https://doi.org/10.5281/zenodo.18849098>,  
520 2026.

521 Khadka, A., Matthews, T., Perry, L. B., Koch, I., Wagnon, P., Shrestha, D., Sherpa, T. G., Aryal, D., Tait,  
522 A., Sherpa, T. C., Tuladhar, S., Baidya, S. K., Elvin, S., Elmore, A. C., Gajurel, A., and Mayewski, P. A.:  
523 Weather on Mount Everest during the 2019 summer monsoon, *Weather*, 76, 205–207,  
524 <https://doi.org/10.1002/wea.3931>, 2021.

525 Khadka, A., Wagnon, P., Brun, F., Shrestha, D., Lejeune, Y., and Arnaud, Y.: Evaluation of ERA5-Land  
526 and HARv2 reanalysis data at high elevation in the upper Dudh Koshi basin (Everest region, Nepal), *J.*  
527 *Appl. Meteorol. Climatol.*, 61, 931–954, <https://doi.org/10.1175/jamc-d-21-0091.1>, 2022.

528 King, O., Bhattacharya, A., Ghuffar, S., Tait, A., Guilford, S., Elmore, A. C., and Bolch, T.: Six Decades  
529 of Glacier Mass Changes around Mt. Everest Are Revealed by Historical and Contemporary Images, *One*  
530 *Earth*, 3, 608–620, <https://doi.org/10.1016/J.ONEEAR.2020.10.019>, 2020.

531 Kochendorfer, J., Earle, M., Rasmussen, R., Smith, C., Yang, D., Morin, S., Mekis, E., Buisan, S., Roulet,  
532 Y.-A., Landolt, S., Wolff, M., Hoover, J., Thériault, J. M., Lee, G., Baker, B., Nitu, R., Lanza, L., Colli,  
533 M., and Meyers, T.: How Well Are We Measuring Snow Post-SPICE?, *Bull. Am. Meteorol. Soc.*, 103,  
534 E370–E388, <https://doi.org/10.1175/BAMS-D-20-0228.1>, 2022.

535 Löffler-Mang, M. and Joss, J.: An optical disdrometer for measuring size and velocity of hydrometeors., *J.*  
536 *Atmos. Ocean. Technol.*, 17, 130–139, 2000.

537 Matthews, T., Perry, L. B., Koch, I., Aryal, D., Khadka, A., Shrestha, D., Abernathy, K., Elmore, A. C.,  
538 Seimon, A., Tait, A., Elvin, S., Tuladhar, S., Baidya, S. K., Potocki, M., Birkel, S. D., Kang, S., Sherpa, T.  
539 C., Gajurel, A., and Mayewski, P. A.: Going to Extremes: Installing the World’s Highest Weather Stations  
540 on Mount Everest, *Bull. Am. Meteorol. Soc.*, 101, E1870–E1890, [https://doi.org/10.1175/BAMS-D-19-](https://doi.org/10.1175/BAMS-D-19-0198.1)  
541 [0198.1](https://doi.org/10.1175/BAMS-D-19-0198.1), 2020a.

542 Matthews, T., Perry, L. B., Lane, T. P., Elmore, A. C., Khadka, A., Aryal, D., Shrestha, D., Tuladhar, S.,  
543 Baidya, S. K., Gajurel, A., Potocki, M., and Mayewski, P. A.: Into Thick(er) Air? Oxygen Availability at  
544 Humans’ Physiological Frontier on Mount Everest, *iScience*, 23, 101718,  
545 <https://doi.org/10.1016/j.isci.2020.101718>, 2020b.

546 Matthews, T., Perry, L. B., Khadka, A., Sherpa, T. G., Shrestha, D., Aryal, D., Tuladhar, S., Thapa, N.,  
547 Pradhananga, N., Athans, P., Sherpa, D. Y., Guy, H., Seimon, A., Elmore, A. C., Li, K., and Alexiev, N.:  
548 Weather observations reach the summit of Mount Everest, *Bull. Am. Meteorol. Soc.*, 1,  
549 <https://doi.org/10.1175/BAMS-D-22-0120.1>, 2022.

550 Miner, K. R., Mayewski, P. A., Baidya, S. K., Broad, K., Clifford, H., Elmore, A., Gajurel, A. P., Giri, B.,  
551 Guilford, S., Hubbard, M., Jaskolski, C., Koldewey, H., Li, W., Matthews, T., Napper, I., Perry, L. B.,  
552 Potocki, M., Priscu, J. C., Tait, A., Thompson, R., and Tuladhar, S.: An overview of physical risks in the  
553 Mt. Everest region, *One Earth*, 3, 547–550, <https://doi.org/10.1016/j.oneear.2020.10.008>, 2020.

554 Pérez-Tello, M. A., Platero, I., Valdivia, J. M., Martinez-Castro, D., Villalobos-Puma, E. E., and Silva, F.  
555 Y.: Evaluation and Correction of Precipitation Types Measured by a PARSIVEL2 Disdrometer in a  
556 Tropical Glacier Environment, *EGUsphere*, 1–36, <https://doi.org/10.5194/EGUSPHERE-2026-2705>,

2026.

Perry, L. B., Matthews, T., Guy, H., Koch, I., Khadka, A., Elmore, A. C., Shrestha, D., Tuladhar, S., Baidya, S. K., Maharjan, S., Wagnon, P., Aryal, D., Seimon, A., Gajurel, A., and Mayewski, P. A.: Precipitation Characteristics and Moisture Source Regions on Mt. Everest in the Khumbu, Nepal, *One Earth*, 3, 594–607, <https://doi.org/10.1016/j.oneear.2020.10.011>, 2020.

Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Graham, J., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Mölg, M., Paul, F., Radic', V., Radic', R., Rastner, P., Raup, B. H., Rich, J., Sharp, M. J., and Consortium, R.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*, 60, 537–552, <https://doi.org/10.3189/2014JoG13J176>, 2014.

Potocki, M., Mayewski, P. A., Matthews, T., Perry, L. B., Schwikowski, M., Tait, A. M., Korotkikh, E., Clifford, H., Kang, S., Sherpa, T. C., Singh, P. K., Koch, I., and Birkel, S.: Mt. Everest's highest glacier is a sentinel for accelerating ice loss, *npj Clim. Atmos. Sci.*, 5, 1–8, <https://doi.org/10.1038/s41612-022-00230-0>, 2022.

Ryan, W. A., Doesken, N. J., and Fassnacht, S. R.: Ultrasonic snow depth sensors for national weather service. Snow measurements in the US: evaluation of operational readiness, *Hydrol. Process.*, 22, 2748–2757, <https://doi.org/10.1002/HYP.7065>, 2008.

Salerno, F., Guyennon, N., Thakuri, S., Viviano, G., Romano, E., Vuillermoz, E., Cristofanelli, P., Stocchi, P., Agrillo, G., Ma, Y., and Tartari, G.: Weak precipitation, warm winters and springs impact glaciers of south slopes of Mt. Everest (central Himalaya) in the last 2 decades (1994-2013), *Cryosph.*, 9, 1229–1247, <https://doi.org/10.5194/tc-9-1229-2015>, 2015.

Salerno, F., Guyennon, N., Yang, K., Shaw, T. E., Lin, C., Colombo, N., Romano, E., Gruber, S., Bolch, T., Alessandri, A., Cristofanelli, P., Putero, D., Diolaiuti, G., Tartari, G., Verza, G., Thakuri, S., Balsamo, G., Miles, E. S., and Pellicciotti, F.: Local cooling and drying induced by Himalayan glaciers under global warming, *Nat. Geosci.*, 16, 1120–1127, <https://doi.org/10.1038/s41561-023-01331-y>, 2023.

Salerno, F., Guyennon, N., Colombo, N., Melis, M. T., Dessì, F. G., Verza, G., Bista, K., Sheharyar, A., and Tartari, G.: What is climate change doing in the Himalaya? Thirty years of the Pyramid Meteorological Network (Nepal), *Earth Syst. Sci. Data*, 17, 4293–4304, <https://doi.org/10.5194/ESSD-17-4293-2025>, 2025.

Shea, J. M., Wagnon, P., Immerzeel, W. W., Biron, R., Brun, F., and Pellicciotti, F.: A comparative high-altitude meteorological analysis from three catchments in the Nepalese Himalaya, *Int. J. Water Resour. Dev.*, 31, 174–200, <https://doi.org/10.1080/07900627.2015.1020417>, 2015.

Sherpa, T. C., Matthews, T., Perry, L. B., Thapa, A., Singh, P. K., Khadka, A., Koch, I., Pelto, M., Panday, P., Aryal, D., Shrestha, D., Kang, S., and Mayewski, P. A.: Insights from the first winter weather observations near Mount Everest's summit, *Weather*, 78, 344–348, <https://doi.org/10.1002/wea.4374>, 2023.

Thakuri, S., Salerno, F., Smiraglia, C., Bolch, T., D'Agata, C., Viviano, G., and Tartari, G.: Tracing glacier changes since the 1960s on the south slope of Mt. Everest (central Southern Himalaya) using optical satellite imagery, *Cryosph.*, 8, 1297–1315, <https://doi.org/10.5194/tc-8-1297-2014>, 2014.

Thakuri, S., Dahal, S., Shrestha, D., Guyennon, N., Romano, E., and Colombo, N.: Elevation-dependent

597 warming of maximum air temperature in Nepal during 1976–2015, *Atmos. Res.*, 228, 261–269,  
598 <https://doi.org/https://doi.org/10.1016/j.atmosres.2019.06.006>, 2019.

599 The Himalayan Times: NatGeo scientists instal highest weather station on Mt Everest - The Himalayan  
600 Times, 2022.

601 Wagnon, P., Brun, F., Khadka, A., Berthier, E., Shrestha, D., Vincent, C., Arnaud, Y., Six, D., Dehecq,  
602 A., Ménégoz, M., and Jomelli, V.: Reanalysing the 2007-19 glaciological mass-balance series of Mera  
603 Glacier, Nepal, Central Himalaya, using geodetic mass balance, *J. Glaciol.*, 67, 117–125,  
604 <https://doi.org/10.1017/jog.2020.88>, 2021.

605 Van Wyk de Vries, M., Matthews, T., Perry, L. B., Thapa, N., and Wilby, R.: At-scale Model Output  
606 Statistics in mountain environments (AtsMOS v1.0), *Geosci. Model Dev.*, 17, 7629–7643,  
607 <https://doi.org/10.5194/gmd-17-7629-2024>, 2024.

608 Yang, W., Guo, X., Yao, T., Yang, K., Zhao, L., Li, S., and Zhu, M.: Summertime surface energy budget  
609 and ablation modeling in the ablation zone of a maritime Tibetan glacier, *J. Geophys. Res. Atmos.*, 116,  
610 14116, <https://doi.org/10.1029/2010JD015183>, 2011.

611

612