

# C-PEAT’s Global Peatland Carbon Database (v.2025)

Julie Loisel<sup>1,2</sup>, Angela Gallego-Sala<sup>3</sup>, Daniela Ransby<sup>4</sup>, Emily Rabel<sup>1</sup>, Cornelia Behrens<sup>5</sup>, Stefanie Schumacher<sup>4</sup>, Kathe Todd-Brown<sup>6</sup>, Vaasuki Marupaka<sup>6</sup>, C-PEAT Working Group Members<sup>^</sup>

<sup>1</sup> Dept. of Geography, Texas A&M University, College Station, USA.

<sup>2</sup> Dept. of Geography, University of Nevada, Reno, USA.

<sup>3</sup> Dept. of Geography, University of Exeter, Exeter, UK.

<sup>4</sup> PANGAEA, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany.

<sup>5</sup> PANGAEA, MARUM – Center for Marine Environmental Sciences, University of Bremen, Germany.

<sup>6</sup> Dept. of Environmental Engineering Sciences, University of Florida, Gainesville, USA.

<sup>^</sup> A full list of data contributors appears at the end of the paper.

Correspondence to: Julie Loisel (jloisel@unr.edu)


**Abstract.** Field-based measurements are foundational to the study of short- and long-term peatland carbon dynamics. For decades, the scientific community has amassed hundreds of valuable empirical datasets in the form of peat core records from around the world. Those records typically include peat depth, basal age, peat organic matter content, peat dry bulk density, peat organic density, and/or carbon and nitrogen content. Once combined with chronological constraints and models, peat core time series can be used to estimate changes in peat-carbon accumulation rates through time. Consolidating these peat records can help improve global peat-carbon stock estimates and quantifications of past, present, and future greenhouse gas exchanges between peatlands and the atmosphere. Large-scale synthesis can also shed light on the sensitivity of peat-carbon accumulation processes to climate change and provide context for current and future global environmental change. We can also use spatial and temporal peat data to inform, validate, and benchmark existing models that include peatland representations. This paper presents the first formal version of PAGES’ C-PEAT Global Peatland Carbon Database (GD), which is available for download in the PANGAEA and International Soil Carbon Network (ISCN) data repositories. The C-PEAT GD contains 267 independently catalogued peat cores and a large number of observations from those cores, including: peat depths, organic matter content values, dry bulk density values, organic density values, as well as carbon and nitrogen content values. Raw and calibrated chronological data are included for each individual dataset when available. The metadata fields are easily searchable and interoperable, as per PANGAEA’s standards. The main objective of this article is to describe the structure and content of the database, itself aimed at increasing the use, assimilation, and interoperability of peat-core data across disciplines. The C-PEAT GD can be accessed at the PANGAEA data repository (<https://doi.org/10.1594/PANGAEA.986891>; Loisel et al., 2025).

## 1 Introduction

Peatlands have stored a vast quantity of carbon (C) in their thick organic soils over the past ~10,000 years (Gorham 1991; Yu et al. 2010). The effect of this long-term **carbon** sink on the global climate system has been that of a net cooling (Smith et al. 2004; Frolking et al. 2006; Yu 2012), despite large methane (CH<sub>4</sub>) emissions to the atmosphere (Spahni et al. 2013; Packalen et al. 2014; Treat et al. 2021). While peatlands only occupy 3% of the global land area, their current **carbon** store has been estimated at ~ 500-1000 gigatonnes (Gt) (Yu 2012; Nichols & Peteet 2019), with an apparent long-term averaged **carbon** accumulation rate (aCAR) of 23 g C m<sup>-2</sup> yr<sup>-1</sup> (Loisel et al. 2014). We note that peat soil **carbon** density and aCAR are known to vary over space and time, as they depend on a number of internal factors that include local vegetation communities, hydrological setting, and topography, as well as a suite of external factors such as air temperature, moisture regime, and climate seasonality (Jones & Yu 2010; Charman et al. 2013; Cobb et al. 2017; Hoyt et al. 2019). Other key

agents capable of altering a peatland's **carbon** store include wildfire, permafrost dynamics, sea-level change, atmospheric pollution and dust deposition, and anthropogenic land-use change (Dommain et al. 2011; Gibson et al. 2018; Hooijer et al. 2012; Hoyt et al. 2020; Turetsky et al. 2020).

Over the past ~ 35 years, the literature has become rich with empirical- and model-based studies aimed at quantifying peatland **carbon** storage from around the world as well as their past, present, and future CARs (Fig. 1a). While the majority of these studies have focused on northern high-latitude peatlands (> 45° N; Gorham 1991; Turunen et al. 2002; Yu 2012), there has been an increase in the number of studies based in the temperate, tropical, and southern high-latitude regions (Kurnianto et al. 2015; Kock et al. 2020; Ruwaimana et al. 2020), as well as global assessments (Gallego-Sala et al. 2018; Qiu et al. 2020; Loisel et al. 2021; Müller & Joos 2021). In particular, studies of tropical peatlands have been on the rise since 2010 (Fig. 1b).




**Figure 1.** Number of peer-reviewed publications that relate to peat carbon, per year (1991-2024), according to Web of Science. **(a)** Global peatlands (search terms: peat\* AND carbon AND (stock\* OR stor\*) AND rate\*). **(b)** Tropical peatlands (search terms: peat\* AND carbon AND (stock\* OR stor\*) AND rate\* AND tropic\*).

The 2010s have also brought along many large-scale peat-data compilations (Fig. 2), perhaps in part thanks to the broad goals of increasing scientific availability/reproducibility (Ellison 2010; Bond-Lamberty et al. 2016). The need for transparency in methods, code, and data **has** also been fueling an increase in collaborative, open science (Adams 2012; Wolkovitch et al. 2012), which helps fulfill a growing need for integrated knowledge and actionable science, in addition to bringing an additional element of trust to the body of science (Grand et al. 2012). In particular, synthesis work has been contributing to initiating new collaborations and building working groups across the scientific community (Baron et al. 2017). In the case of ecological data, syntheses and meta-analyses increase visibility of, and accessibility to, individual case studies (Markowitz et al. 2015). These efforts may help the safeguarding, rescuing, and disseminating of ecological data that would otherwise be lost or remain inaccessible, particularly if unpublished (Dietze et al. 2014; Vines et al. 2014; Harden et al. 2017). Indeed, access to raw ecological data is generally thought to range between 1 and 10% (Reichman et al. 2011; Wolkovitch et al. 2012), making data curation, data accessibility, and data compatibility essential parts of a scientist's work today. This is particularly the case for field ecological observations and measurements that cannot be reproduced in a context of climate change (Wolkovitch et al. 2012).

It has been argued that *synthesis work* provides critical knowledge to solving environmental problems and informing policy (Carpenter 2009; Hampton & Parker 2011; Dicks et al. 2014). By integrating multiple lines of evidence, the power of synthesis work lies in its ability to develop new knowledge that broadens perspectives. The synthesis of existing research provides **a** means to assess individual case studies and explore generalities among datasets; it limits emphasis on specificities and instead focuses on distilling information to improve scientific understanding. In the case of peat core data, the combination of hundreds of peat records can help improve global peat-carbon stock estimates and quantifications of past,

present, and future greenhouse gas exchanges between peatlands and the atmosphere (e.g., MacDonald et al. 2006; Yu 2012; Yu et al. 2013). Large-scale synthesis can also shed light on the sensitivity of peat-carbon accumulation processes to climate change (e.g., Charman et al. 2013; Loisel et al. 2014; Treat et al. 2019) and provide context for current and future global environmental change (e.g., Gallego-Sala et al. 2018; Hugelius et al. 2020; Qiu et al. 2020). We can also use spatial and temporal peat data to inform, validate, and benchmark existing Earth System Models that include peatland representations (Kleinen et al. 2012; Spahni et al. 2013; Qiu et al. 2018; Alexandrov et al. 2020; Chaudhary et al. 2020; Müller & Joos 2021a, 2021b; Chadburn et al. 2022).

Here we present the first formal version of PAGES' C-PEAT Global Peatland Carbon Database (GD), which is now available for download on the PANGAEA data repository. The C-PEAT GD contains 267 previously published independent peat cores and a large number of observations from those cores: 267 peat core depths, 43,467 peat dry bulk density values, 31,473 peat organic matter content values, 4,016 peat organic matter density values, 42,566 carbon content values, and 10,771 nitrogen content values. The main objective of this article is to describe the structure and content of the database, itself aimed at increasing the use, assimilation, and interoperability of peat-core data across disciplines. The database is made of four synthesis papers that were published as part of international community efforts deployed by the C-PEAT working group over the past decade. Those four synthesis papers are introduced below, and in Fig. 2, with the latter also acknowledging a number of additional synthesis papers on long-term peatland dynamics and carbon stock. C-PEAT is a group of over 350 scientists dedicated to synthesizing data and knowledge on the evolution of peatlands through Earth's recent history via collaborations between international researchers. C-PEAT operates under the umbrella of the parent organizations Past Global Changes (PAGES) and Future Earth, and has received further support from the International Union of Quaternary Research (INQUA) and UNESCO's International Geoscience Programme (IGCP) to organize community workshops and training sessions.



**Figure 2.** Timeline of some key synthesis papers on peatland carbon. Blue: northern hemisphere; Orange: global.

**Commented [JL1]:** this is the revised figure.


Yu et al. (2010) compiled **carbonC** accumulation records from 76 sites, which were used to reconstruct changes in the peat-  
**carbonC** pool since the Last Glacial Maximum for the northern, tropical, and southern peatland regions. Charman et al.  
(2013) then focused on 24 well-dated peat profiles from the northern high-latitude regions to examine changes in C  
accumulation over the past millennium. Loisel et al. (2014) augmented the Yu et al.'s northern dataset by using peat core  
data from over 200 sites to report Holocene peat properties (i.e., bulk density, organic matter, **carbonC**, and nitrogen (N)  
content) and aCARs for 127 of those sites. Gallego-Sala et al. (2018) further extended the last millennium work from  
Charman et al. (2013) by combining over 250 millennium-aged peat-core records from around the globe with a modeling  
approach to quantify the future global peatland carbon sink function. We note that the C-PEAT group has worked on other  
peatland databases worth mentioning here. For instance, Treat et al. (2015) looked at peatland plant macrofossil records  
from 280 sites across the northern permafrost zone and reported differences in peat properties as well as aCARs among the  
different vegetation types and environmental classes. Yu et al. (2013) and Loisel et al. (2017) combined 2808 basal peat ages  
from previously published papers to reconstruct spatial and temporal patterns of Holocene peatland area change. Lastly,  
Treat et al. (2019) used over 1000 stratigraphic records of buried peatlands from around the world to reconstruct peat-

forming wetland dynamics over the past 130,000 years. The datasets from those last few papers are also available on PANGAEA, but the peat cores have not yet been cataloged individually. Recently, the compilations by Widyastuti et al. (2025) and Skye et al. (2026) have also contributed global peat depth datasets.

2. Methods

2.1. Definitions and potential uses for the C-PEAT GD

The C-PEAT GD includes high-resolution peat-core geochemical and chronological data that can be used to calculate a number of soil carbonC indices (Fig. 3). At the peat-core level, the information can be used to quantify peatland carbonC storage (units vary, but typically shown as  $g\ C\ cm^{-2}$ ,  $kg\ C\ m^{-2}$ , or  $Mg\ C\ ha^{-1}$ ) and C density ( $kgC/m^3$ ). These two indices are commonly used to estimate C stocks across biomes and the terrestrial biosphere (e.g., Armentano & Menges 1986; Gorham 1991; Petrokofsky et al. 2012). Apparent Ccarbon accumulation rates (aCAR;  $g\ C\ m^{-2}\ yr^{-1}$ ) can be obtained by dividing a core's carbonC density by the basal age of that core (Tolonen & Turunen 1996); multiple ages and carbonC measurements along a peat profile can also be used to estimate temporal changes in aCAR (e.g., Yu et al. 2010), though this technique has methodological limits; see belowaCAR should not be misinterpreted as net carbon balance (Young et al. 2019, 2021). In previous synthesis work, aCAR was calculated for each peat layer (Fig. 3); the layer data were then combined into bins (e.g., 50, 100, 500, 1000 years) to provide time-weighted aCAR (Yu et al. 2010; Jones & Yu 2011; Loisel et al. 2014; Fig. 3). The latter makes it easier to compare apparent accumulation rates across sites and build regional assessments. It is important to note that aCAR is an imperfect measure of past accumulation rates because peatlands may be subject to secondary peat loss - from fire, erosion, a lasting drought, or management - that takes place long after initial peat deposition and early-stage diagenesis (Yu 2012; Froelking et al 2014; Young et al. 2019, 2021). Thus, a fundamental problem with aCAR is that it cannot rule out whether secondary peat loss has taken place. Aggregating a large number of core site data for which multiple age determinations and carbonC measurements are available can help alleviate the issue, if one assumes that; indeed, secondary decomposition can be seen as site-specific “noise” that would be subdued in a synthesis product, though large-scale changes could affect several sites at once. Recent aCAR calculations (past ~150 years; Figure 3) are technically possible using our database. But here again, the use of these datasets comes with important limitations. For instance, we emphasize that near-surface records tend to represent incompletely decomposed, young peat, such that recent aCAR should not be directly compared against longer-term aCAR (Piilo et al. 2019; Young et al. 2019). In fact, the past ~150 years of peat accumulation are often omitted from synthesis work (e.g., Charman et al. 2013; Gallego-Sala et al. 2018). Lastly, it should be mentioned that the C-PEAT GD database also includes nitrogenN content; nitrogenN stock, soil nitrogenN density, and aNAR can thus be calculated.



**Figure 3.** Datasets available from the C-PEAT Global Peatland Carbon Database. The \* denotes soil carbon indices that can be derived from the database. For additional details, refer to Tables 2-3-4. Photo credit: Julie Loisel.

**Commented [JL2]:** this is a revised version of Fig 3.

160 **2.2. Peat carbon database architecture**  
 Here we present the first version (coined v.2025) of the C-PEAT GD. The following paragraphs provide the general database structure. We start with a brief introduction of the collaborative process that gave birth to the database. We then describe data aggregation and formatting, list the types of data/metadata that are available, and we finish by presenting data selection criteria as well as data coverage. The C-PEAT GD can be accessed via PANGAEA at: [https://www.pangaea.de/?q=project:label:PAGES\\_C-PEAT](https://www.pangaea.de/?q=project:label:PAGES_C-PEAT). The latter landing page contains links to download individual records; the entire database can also be mass-downloaded using R or Python (Ransby et al. 20252026). Metadata and data can be obtained from the PANGAEA repository with scripts using the ‘pangaeR’ R package (Chamberlain et al. 2021) and the ‘pangaeapy’ Python library (Huber et al. 2025). More information on submission of new or revised datasets can be found in sect. 4.1.

170 This database is the product of approximately 10 years of collaborative work by the C-PEAT working group, which is endorsed by PAGES. This synthesis effort focuses on peatland-C records; it is a contribution to C-PEAT’s second developmental phase and fulfills the objective of making the community’s peat-C data widely available (Loisel & Gallego-Sala 2018). The C-PEAT GD is the culmination of multiple synthesis efforts that have taken place over the past decade (see section 1 and Fig. 2). For each one of these previous studies, calls for participation were widely distributed. Data inclusion criteria differed between papers to match their different scopes; in each case, decisions were made by a group of lead authors in consultation with the data contributors (Table 1). The workload for assembling the data and metadata that make up C-PEAT’s GD was coordinated at Texas A&M University. All data from the four compilation papers are included in the C-PEAT GD.


180 **Table 1.** Criteria for inclusion used by the four peat-data synthesis articles that form the C-PEAT Global Peatland Carbon Database (v.2025).

| References               | Chronological constraints                                             | Geochemical constraints                                  | Geographic constraints |
|--------------------------|-----------------------------------------------------------------------|----------------------------------------------------------|------------------------|
| Yu et al. 2010           | multiple calibrated ages                                              | some bulk density measurements                           | none                   |
| Charman et al. 2013      | at least 3 evenly spaced dates spanning the last 1000 years           | contiguous bulk density measurements at <5 cm resolution | latitude: >45N         |
| Loisel et al. 2014       | at least 1 date per thousand years (on average for the entire record) | some bulk density or organic matter density measurements | latitude: >45N         |
| Gallego-Sala et al. 2018 | at least 2 radiocarbon dates from the past 1000 years                 | contiguous bulk density measurements at <2 cm resolution | none                   |

185 **2.3. Data aggregation, formatting, and availability in PANGAEA**  
 As a data publisher for Earth & Environmental Sciences, PANGAEA (Felden et al. 2023) has an almost 40-year history as an open-access library for archiving, publishing, and disseminating georeferenced data from the Earth, environmental, and biodiversity sciences. Originally evolving from a database for sediment cores, it is operated as a joint facility of the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, and the Center for Marine Environmental Sciences at the University of Bremen. PANGAEA has been certified with the Core Trust Seal since 2019. Each published dataset can be cited with a specific data citation and an associated unique Digital Object Identifier (DOI) to cross-link to related resources

such as manuscripts and data. A broad spectrum of contextual information ('metadata') describing the where, how, when, and why of a measurement is given. Additionally, the methods and devices used for sampling and analysis as well as references of related literature and external resources are provided. All data and metadata are compiled in close collaboration between the scientists and trained field experts acting as data editors. Both data and metadata are checked for completeness and plausibility, ensuring high quality standards according to the FAIR data principles. Semantic interoperability during data curation is ensured through strict application and dynamic evolution of terminologies according to international protocols and standards, such as Environment ontology (EnvO), the Integrated Taxonomic Information System (ITIS), and Quantities, Units, Dimensions and DataTypes Ontology (QUDT). This harmonization and standardization promote not only readability and further processability by machines, but also a high degree of reusability of the data stock. In addition to the classic access to data via the website, an integrative use of data in the form of a DataWarehouse and a set of tools for programmatic data processing are available for this purpose. The two applications, 'pangaeapy' (Huber et al. 2025) and 'pangaear' (Chamberlain et al. 2021), respectively, are written for the scripting languages Python and R, and make use of the well-developed interoperability framework of PANGAEA.

Each peat core record that is a part of our GD was either directly contributed by the original data owner or obtained from the published literature by the different synthesis author teams (Fig. 4). In most cases, records that were identified in the literature were then obtained from the corresponding author via email. In a few cases, datasets were digitized from publication tables, appendices, and supplementary materials. The synthesis team leads provided templates to the data contributors to facilitate aggregation. In the case of data acquired from publications, the synthesis teams entered the data into the templates themselves. At this stage, data quality was manually assessed by said teams (Table 1). The data were then compiled and used for analysis. During the second stage –(i.e., in preparation for the C-PEAT GD)– the data and metadata were reviewed and formatted to harmonize and standardize them with the PANGAEA database. To date, PANGAEA hosts 758 C-PEAT datasets obtained from 267 distinct cores (267 geochemistry datasets, 264 age determination datasets, and 227 calibrated ages datasets). These datasets were authored by 81 individuals (or groups of individuals). The datasets refer to 84 journal articles as well as 4 theses, dissertations, and conference papers. Each dataset is associated with an "Event", which is a metadata descriptor used by PANGAEA for characterizing the location site / station where a physical sample or measurement took place. For the C-PEAT database, "Event" is a synonym for a "peat core". While the database contains individual peatlands that harbor several cores, each "core" is represented individually. There is no direct way to differentiate by "peatland site", besides by looking at the "event" names: cores collected from a single site typically have the same first letters in their name (e.g., event: Mariana\_coreMF03-1 vs. event: Mariana\_coreMF03-2). Mandatory metadata for each Event comprises a unique "Label" as well as "Latitude" and "Longitude" information (in decimal degrees). Further optional metadata include an additional / alternative core Label, Sampling method, Elevation (meters above sea level) and a Comment (free text) (Fig. 4). Duplicate cores across the 4 synthesis papers were reconciled to avoid duplications in the database. Cores from the most recent synthesis were selected over data from the same core site that were previously published (e.g., we would include a core presented by Charman et al. (2013) and leave its older duplicate from Yu et al. (2010)). This selection allowed to integrate the most updated values into the C-PEAT database, should the core data were updated over time.



**Figure 4.** Workflow to integrate C-PEAT data into PANGAEA. Each peat core (called an Event in PANGAEA) is linked to Location and Methods metadata. Each peat layer includes geochemical data, raw age determinations, and calibrated ages (referred to as Data in PANGAEA). **Three** datasets per core are created, each with a distinct DOI. Those data are linked to Methods and References from the original publication (if published).

The geochemistry datasets contain the key peat geochemical properties along with the ancillary data that may complement, or help interpret, the data (Table 2). These ancillary data include entries such as the main peat type. Age determination datasets provide raw **and/or calibrated** age control data **from different chronological sources**, such as radiocarbon ( $^{14}\text{C}$ ),  $^{210}\text{Pb}$ , or tephra chronostratigraphy, together with information on the type of dated material (bulk, wood, pollen etc.) (Table 3). **Since different authors opt for different nomenclatures and notation systems when presenting their raw dates and calibrated ages, PANGAEA uses a relatively long list of parameters aimed at maintaining the original meaning of the original data entries.** For example, the parameters “AGE [ ka BP ]” and “Age model [ ka ]” are used to store the calibrated age for each sample along a peat core, and the data are presented in thousands of calibrated years Before Present (BP). In publications where age is instead presented in calendar years, the PANGAEA parameters “AGE [ a ]”, “AGE [ a, AD/CE ]”, or “AGE [ a, years ago ]” are used. The same goes with archiving age uncertainties: some authors calculate their own uncertainty range (PANGAEA parameter: “Age, dated, uncertainty [  $\pm$  ]”) while others opt to present one standard deviation (PANGAEA parameter: “Age, dated standard deviation [  $\pm$  ]”). Age uncertainties can also be presented in ka or in years, and different parameters are used accordingly. Table 3 presents the complete list of parameters (and associated short descriptions) that are used to archived the C-PEAT GD data entries pertaining to dates and ages. Age-depth relationships of individual peat cores, **including their upper and lower limits**, are presented in the calibrated ages datasets (Table 4). **Note**



that ~~These~~ ages were obtained from the original publications published papers and are thus based on different age-depth modeling techniques and calibration curves.

255 Geochemical metadata were obtained from the contributing authors (or extracted from the peer-reviewed articles) and they  
relate to study core site and author information, as well as analytical and sampling methods (Table 2). Chronological  
260 metadata relate to the different types of dating methods, laboratories where the analysis were performed, and they also  
include additional sampling information (when available from the original articles) (Table 3).

**Table 2.** Geochemical parameters included in the C-PEAT Global Peatland Carbon Database, and-with associated  
descriptions and metadata.

| Parameter [unit]                                 | Description                                                                                                                                             | Parameter short name |
|--------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|
| DEPTH, sediment/rock [m]                         | <u>Ssample midpoint depth, measured from the surface</u>                                                                                                | Depth sed            |
| AGE [ka BP]                                      | <u>calibrated age (in thousands of years Before Present, based on the original age-depth curve)</u>                                                     | Age                  |
| Sample thickness [cm]                            | <u>distance (cm) between top and bottom of a sample</u>                                                                                                 | Samp thick           |
| Sample volume [cm^3]                             | <u>amount of material</u>                                                                                                                               | Samp vol             |
| Water content, wet mass [%]                      | <u>weight-%</u>                                                                                                                                         | Water wm             |
| Peat type []                                     | <u>category</u>                                                                                                                                         | Peat type            |
| Sedimentation rate per year [cm/a]               | <u>-amount of peat accumulated per year</u>                                                                                                             | SR                   |
| Peat carbon accumulation rate per year [g/m^2/a] | <u>weight (g) of carbon accumulated per unit of surface area (m^2) per year</u>                                                                         | PCAR                 |
| Organic matter [%] *                             | weight-%                                                                                                                                                | OM                   |
| Carbon [%]                                       | <del>converted</del> calculated from other data such as LOI                                                                                             | C                    |
| Carbon, inorganic, total [%]                     | weight-%                                                                                                                                                | TIC                  |
| Carbon, organic, total [%]                       | weight-%                                                                                                                                                | TOC                  |
| Carbon, total [%]                                | weight-%                                                                                                                                                | TC                   |
| Comment []                                       | <u>free text</u>                                                                                                                                        | Comment              |
| Cumulative mass [g/cm^2]                         | <u>total accumulated peat dry mass up to the sample's depth, where CM = 0 is the deepest sample of the core and CM = max at the surface of the core</u> | Cum mass             |
| Density, dry bulk [g/cm^3]                       | <u>ratio of dry mass to peat bulk volume</u>                                                                                                            | DBD                  |

|                                  |                                                               |           |
|----------------------------------|---------------------------------------------------------------|-----------|
| Density, organic carbon [g/cm^3] | <a href="#">ratio of organic carbon to peat bulk volume -</a> | Corg dens |
| Density, organic matter [g/cm^3] | <a href="#">ratio of organic matter to peat bulk volume -</a> | OM dens   |
| Nitrogen, total [%]              | weight-%                                                      | TN        |

\* In some datasets, the parameter is named Loss on ignition [%] — both were merged for the synthesis.

**Table 3.** Age determination parameters included in the C-PEAT Global Peatland Carbon Database, ~~and with~~ associated [descriptions and](#) metadata.

| Parameter [unit]                  | Description                                                      | Parameter short name |
|-----------------------------------|------------------------------------------------------------------|----------------------|
| DEPTH, sediment/rock [m]*         | <a href="#">sample midpoint depth, measured from the surface</a> | Depth sed            |
| Depth, top/min [m]                | <a href="#">sample top depth, measured from the surface</a>      | Depth top            |
| Depth, bottom/max [m]             | <a href="#">sample bottom depth, measured from the surface</a>   | Depth bot            |
| Sample code/label []              | <a href="#">sample name</a>                                      | Sample label         |
| Sample thickness []               | <a href="#">distance (cm) between top and bottom of a sample</a> | Samp thick           |
| Laboratory code/label []          | <a href="#">sample code (provided by a dating laboratory)</a>    | Lab label            |
| Method comment []                 | <a href="#">free text</a>                                        | Method comm          |
| Fraction modern carbon []         | <a href="#">radiocarbon concentration</a>                        | F14C                 |
| Fraction modern carbon, error [±] | <a href="#">laboratory error associated with F14C</a>            | F14C e               |
| Δ14C [‰]                          | <a href="#">radiocarbon concentration</a>                        | Δ14C                 |
| Δ14C, standard deviation [±]      | <a href="#">laboratory error associated with Δ14C</a>            | Δ14C std dev         |

|                                                                             |                                                                                                              |                   |
|-----------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|-------------------|
| Activity of radiocarbon in percent of modern carbon [pMC]                   | <a href="#">percent of modern carbon in a sample</a>                                                         | Activity          |
| Activity of radiocarbon in percent of modern carbon, standard deviation [±] | <a href="#">laboratory error associated with 14C</a><br><a href="#">Activity</a>                             | Activity std dev  |
| Age, dated [ka]                                                             | <a href="#">calibrated age (in thousands of years Before Present, based on the original age-depth curve)</a> | Age dated         |
| Age, dated, uncertainty [±]                                                 | <a href="#">age uncertainty (in ka) as reported in the original publication</a>                              | Age dated unc     |
| Age, dated standard deviation [±]                                           | <a href="#">age uncertainty if reported as 1 standard deviation in the original publication</a>              | Age dated std dev |
| Age, dated material [ ]**                                                   | <a href="#">type of material dated (e.g., leaf, bulk sediment, charcoal, etc.)</a>                           | Dated material    |
| Age, comment [ ]                                                            | <a href="#">free text</a>                                                                                    | Comm              |
| Age model [ka]                                                              | <a href="#">calibrated age (in thousands of years Before Present, based on the original age-depth curve)</a> | Age model         |
| Age, error [±]                                                              | <a href="#">age error as reported in the original publication</a>                                            | Age e             |
| Age [a]                                                                     | <a href="#">calibrated age (in calendar years, based on the original age-depth curve)</a>                    | Age               |
| Age, uncertainty [±]                                                        | <a href="#">age uncertainty (in calendar years) as reported in the original publication</a>                  | Age unc           |

|                                      |                                                                                                            |                 |
|--------------------------------------|------------------------------------------------------------------------------------------------------------|-----------------|
| Age, standard deviation [±]          | <u>age uncertainty (in calendar years) if reported as 1 standard deviation in the original publication</u> | Age std dev     |
| Age [a AD/CE]                        | <u>calibrated age (in calendar years, based on the original age-depth curve)</u>                           | Age             |
| Age, maximum/old [a AD/CE]           | <u>oldest calibrated age (in calendar years, based on the original age-depth curve)</u>                    | Age max         |
| Age, minimum/young [a AD/CE]         | <u>youngest calibrated age (in calendar years, based on the original age-depth curve)</u>                  | Age min         |
| <u>Age [a years ago]</u>             | <u>calibrated age (in calendar years, based on the original age-depth curve)</u>                           | <u>Age</u>      |
| Calendar age [ka BP]                 | <u>calibrated age (in thousands of calendar years, based on the original age-depth curve)</u>              | Cal age         |
| Calendar age, standard deviation [±] | <u>age uncertainty (in calendar years) as reported in the original publication</u>                         | Cal age std dev |
| Comment []                           | <u>free text</u>                                                                                           | Comment         |

\*sample midpoint depth — \*\*type of material dated (e.g., bulk, pollen, charecoal, etc.)

**Table 4.** Calibrated age parameters included in the C-PEAT Global Peatland Carbon Database, and-with associated descriptions and metadata.

| Parameter [unit]                    | Parameter short name | Description                                    |
|-------------------------------------|----------------------|------------------------------------------------|
| DEPTH, sediment/rock [m]            | Depth sed            | Sample midpoint depth                          |
| Calendar age [ka BP]                | Cal age              | Contains a method comment (e.g., OxCal, Bacon) |
| Calendar age, maximum/old [ka BP]   | Cal age max          | Contains a method comment (e.g., OxCal, Bacon) |
| Calendar age, minimum/young [ka BP] | Cal age min          | Contains a method comment (e.g., OxCal, Bacon) |

275

#### 2.4. Data coverage, resolution, duration, and chronological constraints

280 Data coverage ~~and representativity~~ of the database ~~are is~~ dependent upon data availability, which is mostly out of the control of the data analysts. In the case of the C-PEAT GD, all available records that fulfilled our criteria for inclusion (Table 1) were accepted. As for the resolution and duration of the peat records included in the C-PEAT GD, they vary greatly. For example, the minimal geochemical data resolution was set at every 2 cm by Gallego-Sala et al. (2018) whereas it was set at 5 cm by Charman et al. (2013). This is because those previous synthesis activities had different research objectives (Table 1).

#### 2.5. Analysis performed on the C-PEAT GD

285 To ~~determine-illustrate our~~ overall data coverage ~~and the representativity of our database~~, we used a climate envelope and climate region analysis (Köppen) approach. For each ~~coresite~~, mean annual temperature (MAT) and total annual precipitation (MAP) ~~were~~ extracted from a gridded climate product (Fick & Hijmans 2017). We also investigated the peat geochemical properties of our peat samples. For each variable, central tendencies (mean, median and mode) and dispersions (minimum, maximum, variance, and standard deviation) were calculated. ~~Lastly, we compared our peat soil data against a mineral soil database (USDA-NRCS-NCSS), which is currently harmonized with ISCN v.4 (Todd-Brown et al. In prep). The National Cooperative Soil Survey (NCSS) is a comprehensive soil characterization dataset that contains 15,819 pedons spanning the world (as of July 2024). In addition to comparing central tendencies between these two datasets, we also contrast changes in soil organic C density along depth for peat vs. mineral soils.~~


295

### 3. Results and Discussion

#### 3.1. Data coverage over space and time


300 Spatial coverage of the dataset is global, with the majority of ~~sites-cores~~ (220/267) located in the northern extratropical region (Fig. 5). With the exception of southern South America (Patagonia; 11 cores), data availability is very sparse across the southern extratropical region. As for the tropical belt (around 20° N to 20° S), the C-PEAT dataset includes 34 peat core records (Fig. 5). Future efforts should prioritize the synthesis of tropical core records.

305 Temporal coverage of the peat cores varies over time, ~~with the number of basal ages steadily increasing throughout the Holocene~~ (Fig. 6a). To our knowledge, all peat records found in this database are continuous and their surface is assumed to be of modern age (~~Fig. 6a~~). We note that the oldest age available along a peat core sets the record's duration; in most cases, peat cores extend back to peatland initiation, though many cores do not. In total, 24 peat core records are younger than 1000 years (of which 11 core chronologies were solely constrained by <sup>210</sup>Pb ages), 110 records are at least 5000 years old, and 16 cores initiated prior to 10,000 years ago. The oldest records are found in southern Patagonia and Alaska, presumably because of the earlier deglacial history of those regions. The oldest tropical ~~site-core~~ available in this database is approximately 11,600 years old, though we note that most of the tropical cores included in the dataset are less than 5000 years old (median = 2940 years).




315 **Figure 5.** Spatial distribution of peat cores included in the C-PEAT Global Peatland Carbon Database. Orange circles  
 | represent individual peat cores. The global peatland area (in **blackgreen**) is from UNEP (2022). Map source: Esri, TomTom,  
 | FAO, USGS | Powered by Esri.

320 The temporal resolution of the peat core records varies considerably. For instance, the number of chronological constraints  
 | along a single peat core varies from 1 to 41 (median = 6; Fig. 6b). Dividing the maximum age of a record by the number of  
 | dates available for said record yields an average of 0.76 dates / 1000 years, with 64 records characterized by an averaged  
 | temporal resolution that is greater than 1 date / 1000 years. In general, there are more chronological constraints for the late  
 | Holocene than the mid- and early-Holocene time periods, in part because the data from Gallego-Sala et al. (2018) focused on  
 | the past millennium. Peat accumulation rates also vary widely between **sitescores**. In general, the peatland archives therein  
 | offer a multi-decadal resolution, with an average peat accumulation rate of  $0.057\text{cm}\cdot\text{yr}^{-1}$ , ranging from  $0.005$  to  $0.5\text{cm}\cdot\text{yr}^{-1}$   
 325 (Fig. 6c).



330 **Figure 6.** Temporal resolution of peat cores included in the C-PEAT Global Peatland Carbon Database. **(a)** Cumulative  
 | number of peat core records over time. **(b)** Number of chronological constraints per core. **(c)** Mean apparent rate of peat  
 | accumulation for each core (in  $\text{cm}\cdot\text{yr}^{-1}$ ). On panels b) and c), the 25th-75th percentile ranges (IQR – **interquartile range**) are  
 | displayed as the boxes, with the medians (50th percentile) marked by horizontal black lines inside. The whiskers represent  
 |  $1.5 \times \text{IQR}$  and the circles represent outliers.

335 **3.2. Data representativcoveragety**  
The climate envelope and climate region analyses confirm that peatland **sites-cores** within this database are found across a broad range of hydroclimatic conditions and climate classes, from the tropical rainforest and the tropical savannah to the polar tundra, and many other classes in between. It is clear that cold climate **sites-cores** dominate the dataset (222/267 sites) and that more cores from the warmer bioclimatic regions would help balance representation in the database (Fig. 7).



345 **Figure 7.** Distribution of **core** sites included in the C-PEAT Global Peatland Carbon Database in the context of the  
Köppen-Geiger climate classification. Peatland **sites-cores** (orange circles on the map) are the same as on Fig. 6. The  
climate map is from Beck et al. (2023) and the global peatland area (in black) is from UNEP (2022). The biplot presents  
those same peatland **sites-cores** (color-coded using the Köppen climate classes) in a climate space. The legend reads as  
follows (from Beck et al. 2023): 1 Af (Tropical, rainforest), 2 Am (Tropical, monsoon), 3 Aw (Tropical, savannah), 4 BWh  
350 (Arid, desert, hot), 5 BWk (Arid, desert, cold), 6 BSh (Arid, steppe, hot), 7 BSk (Arid, steppe, cold), 8 Csa (Temperate, dry  
summer, hot summer), 9 Csb (Temperate, dry summer, warm summer), 10 Csc (Temperate, dry summer, cold summer), 11  
Cwa (Temperate, dry winter, hot summer), 12 Cwb (Temperate, dry winter, warm summer), 13 Cwc (Temperate, dry winter,  
cold summer), 14 Cfa (Temperate, no dry season, hot summer), 15 Cfb (Temperate, no dry season, warm summer), 16 Cfc  
355 (Temperate, no dry season, cold summer), 17 Dsa (Cold, dry summer, hot summer), 18 Dsb (Cold, dry summer, warm  
summer), 19 Dsc (Cold, dry summer, cold summer), 20 Dsd (Cold, dry summer, very cold winter), 21 Dwa (Cold, dry  
winter, hot summer), 22 Dwb (Cold, dry winter, warm summer), 23 Dwc (Cold, dry winter, cold summer), 24 Dwd Cold, dry  
winter, very cold winter), 25 Dfa (Cold, no dry season, hot summer), 26 Dfb (Cold, no dry season, warm summer), 27 Dfc  
(Cold, no dry season, cold summer), 28 Dfd (Cold, no dry season, very cold winter), 29 ET (Polar, tundra), 30 EF (Polar,  
frost). Map source: Esri, TomTom, FAO, USGS | Powered by Esri.

360 **3.3. Key peat properties and characteristics**  
When combined, the peat records yield a total of 267 peat **core** depths, 43,467 dry bulk density values, 31,473 organic matter  
content values, 4,016 peat organic matter density values, 42,566 carbon content values (**total-carbon-and-carbon**  
**eomined%C, %TC, %TOC combined**), and 10,771 nitrogen content values. Descriptive statistics of the dataset are  
presented in the paragraphs below as well as in Fig. 8 and Table 5.

365 Peat core depth is provided for all 267 cores and ranges from 0.30 to 10.95 m, with a mean value of  $2.24 \pm 1.9$  m (1 standard  
deviation (SD); Table 5). We note that 'peat core depth' refers to the deepest sample provided **for-along a sitecore**; while

most peat cores in the dataset extend back to peatland initiation, many cores may not. When in doubt, it is recommended to refer to the original publication to confirm whether the record starts at peat inception.


Peat dry bulk density values ( $n = 43,467$ ) are available for 263 [sites-cores](#) and range from 0.01 to  $3.380 \text{ g } \text{cm}^{-3}$ , with a mean value of  $0.140 \pm 0.178 \text{ g } \text{cm}^{-3}$  (1 SD; Fig. 8a; Table 5). Organic matter content data ( $n = 31,473$ ) are available for 174 [sites-cores](#) and range from 0 to 100% (though there are some data points > 100%), with a mean value of  $88.5 \pm 20.1\%$  (1 SD; Fig. 8b; Table 5). Organic matter density values (OMD;  $n = 4016$ ) are available for 20 [sites-cores](#) and range from 0.008 to  $0.399 \text{ g OM } \text{cm}^{-3}$ , with a mean value of  $0.117 \pm 0.046 \text{ g OM } \text{cm}^{-3}$  (1 SD; Table 5).

In the database, carbon content is presented as three distinct variables: C%, TC%, and TOC%. While C% refers to estimated [carbon](#) content based on [organic matter](#) measurements ([authors often multiply their OM% by 50% to estimate C% and data users are prompted to refer to the original publications to know more about the conversion factors that were used](#)), reports of TC (total carbon) and TOC (total organic carbon) data refer to values that were measured directly, typically through elemental analysis. Here we report C% data from 151 [sites-cores](#) ( $n = 26,842$ ), TC% from 114 [sites-cores](#) ( $n = 14,430$ ), and TOC% data from 11 [sites-cores](#) ( $n = 1,532$ ), for a total of 42,804 measurements. We note that 16 [sites-cores](#) report both C% and TC% values, that 7 [sites-cores](#) report C% and TOC% values, that 1 [site-cores](#) reports TC% and TOC% values, and that 1 [site-core](#) reports C%, TC%, and TOC% values. Carbon content (C%) estimates range from 0.0 to 71.7%, with a mean value of  $46.7 \pm 10.5\%$  (1 SD; Fig. 8c; Table 5). Total carbon content (TC%) values range from 0.0 to 100%, with a mean value of  $45.4 \pm 16.4\%$  (1 SD; Fig. 8c; Table 5). Total organic carbon content (TOC%) values range from 0.0 to 77.0%, with a mean value of  $37.7 \pm 19.6\%$  (1 SD; Fig. 8c; Table 5). Lastly, organic carbon density values (OCD;  $n = 3,952$ ) are available for 28 [sites-cores](#) and range from 0.005 to  $0.195 \text{ g C/cm}^3$ , with a mean value of  $0.058 \pm 0.024 \text{ g C } \text{cm}^{-3}$  (1 SD; Table 5). We acknowledge that a few individual data points are above the expected data ranges (e.g., %C > 60%); we did not exclude said data points from the database. Data users can easily filter outliers as they see fit.

Total nitrogen content (TN) data are reported for 87 [sites-cores](#) ( $n = 10,770$ ) and range from 0 to 6.3% (there is an outlier at 56.6% that was omitted from Fig. 8d), with a mean value of  $1.2 \pm 0.7\%$  (1 SD; Fig. 8d; Table 5).

~~Overall, the central tendencies and data ranges presented therein are in line with previously published datasets (e.g., Loisel et al. 2014).~~ While the averaged data (Table 5) may be used as reference [values](#), we advise data users to further dig into the dataset, as the database structure allows for much finer analysis. For instance, it is possible (and simple) to query the dataset for a particular region of the world and/or time period, with the goal to obtain more specific peat characteristics.





**Figure 8.** Descriptive statistics for the C-PEAT Global Peatland Carbon Database. **(a)** Dry bulk density; **(b)** Organic matter content; **(c)** Carbon content, where the 3 datasets are displayed (see sect. 3.3 for details), and **(d)** Nitrogen content. For each variable, the left inner panel displays the **interquartile range (IQR)** (represented as the box, with the median shown as a horizontal black line) while the right inner panel presents the entire data range (including the outliers, denoted as circles). Whiskers present 1.5 x **IQR**.


**Table 5.** Central tendencies and dispersions for the C-PEAT Global Peatland Carbon Database (267 cores).

| Variable name and sample count |        | Central tendencies |        | Dispersions |         |       |
|--------------------------------|--------|--------------------|--------|-------------|---------|-------|
|                                | n      | mean               | median | minimum     | maximum | SD    |
| total peat depth (m)           | 267    | 2.24               | 1.61   | 0.30        | 10.95   | 1.90  |
| DBD (g/cm <sup>3</sup> )       | 43,467 | 0.140              | 0.098  | 0.001       | 3.380   | 0.178 |
| OM (%)                         | 31,473 | 88.5               | 96.5   | 0.0         | 100.0   | 20.1  |
| OMD (g OM/cm <sup>3</sup> )    | 4016   | 0.117              | 0.109  | 0.008       | 0.399   | 0.046 |

|                            |        |       |       |       |       |       |
|----------------------------|--------|-------|-------|-------|-------|-------|
| C (%)                      | 26,842 | 46.7  | 50.8  | 0.0   | 71.7  | 10.5  |
| TC (%)                     | 14,430 | 45.4  | 48.4  | 0.0   | 100.0 | 16.4  |
| TOC (%)                    | 1532   | 37.7  | 46.0  | 0.0   | 77.0  | 19.6  |
| OCD (g C/cm <sup>3</sup> ) | 3952   | 0.058 | 0.055 | 0.005 | 0.195 | 0.024 |
| TN (%)                     | 10,771 | 1.2   | 1.0   | 0.0   | 56.6  | 0.7   |

3.4. Comparing peat vs. mineral soil data

The peat vs. mineral soil data comparisons provided a way to quantify key carbon stock differences between these two soil types. On average, soil organic carbon content for the top 100 cm was 16 times greater in peat soils (17.53 g/cm<sup>2</sup> area mass) than in mineral soils (1.08 g/cm<sup>2</sup>). Most mineral soils in the NCSS database exhibit very low SOC stocks (0–1.33 g/cm<sup>2</sup>), with a skewed distribution for higher stocks (max = 55 g/cm<sup>2</sup>) (Table 6). The distribution of soil organic carbon over depth (top 100 cm) also highlighted key differences, with mineral soil data exhibiting a clear decreasing trend in organic carbon with depth that is absent from the peat soil data (Fig. 9a). Deeper records, which we limit to 600 cm because that is the deepest mineral soil data, show a similar trend (Fig. 9b). Overall, peat soils are rich in organic matter (Table 5) and therefore have higher concentrations of organic carbon compared to mineral soils (Table 6). This result is supported by the order-of-magnitude lower bulk density values for peat soils (mean: 0.14 g/cm<sup>3</sup>) compared to those of mineral soils (mean: 1.54 g/cm<sup>3</sup>).



**Figure 9.** Comparison of soil organic carbon data between the C-PEAT Global Peatland Carbon Database (267 cores) and the NCSS soil pedons (15,819 pedons). (a) Depth distribution of soil organic carbon stocks for the top 100 cm soil profile for C-PEAT samples (top) vs. NCSS soils (bottom). (b) Depth distribution of soil organic carbon stocks down to 600 cm for C-PEAT samples (top) vs. NCSS soils (bottom).

**Table 6.** Central tendencies and dispersions for the NCSS soil pedons (15,819 pedons).

| Variable name and sample count | Central tendencies | Dispersions |
|--------------------------------|--------------------|-------------|
|--------------------------------|--------------------|-------------|

|                                           | n      | mean   | median  | minimum | maximum | SD     |
|-------------------------------------------|--------|--------|---------|---------|---------|--------|
| Bulk density (oven-dry)                   | 78,551 | 1.5452 | 1.58    | -0.03   | 4.47    | 0.263  |
| Soil Organic Carbon (g/cm <sup>3</sup> )  | 78,551 | 0.0123 | -0.0063 | -0.3746 | -0.6142 | 0.019  |
| estimated_organic_carbon                  | 30,634 | 1.3655 | 0.5     | -3.9    | -62.4   | 3.419  |
| Organic C walkley-black (%)               | 48,018 | 0.9304 | 0.4     | -0.15   | 54      | 1.895  |
| Total coarse fraction (wt_pet_gt_2_mm_ws) | 78,551 | -8.96  | -1      | 0       | 4229    | 28.767 |

4. Usage notes

4.1. Database availability, updates, and submission of new datasets

The C-PEAT GD v.2025 presented therein can be accessed via PANGAEA, under the project label C-PEAT. There also exist R and Python scripts that allow for mass-downloads of the entire dataset (Ransby et al. 2026<sup>5</sup>). The database is expected to keep evolving over time, as new datasets are integrated, existing datasets are extended (or corrected), and new sharing standards are developed. Formal database updates will be tentatively initiated every two to three years; said updates will be described online, directly within the data repository. Future updates will be reflected in the database identifier, which will include the update year (e.g., v.2027, v.2030, etc.). Between updates, database users are asked to flag any existing issues by contacting the PANGAEA editors through the contact form <https://www.pangaea.de/contact/>; a new version of the erroneous data can then be published. The inclusion of new datasets or the expansion of existing ones are performed manually. Data contributors interested in sharing one or a handful of datasets are asked to follow the instructions and use the templates provided by PANGAEA ([https://wiki.pangaea.de/wiki/Best\\_practice\\_manuals\\_and\\_templates](https://wiki.pangaea.de/wiki/Best_practice_manuals_and_templates)). Contributors who wish to share a larger quantity of datasets (> 10) should contact a PANGAEA editor through the contact form linked above.

Whether they used the database in its entirety or only a subset, users are asked to cite the C-PEAT GD v.2025. When possible, and particularly when a subsequent product makes use of a few specific records, we request that the original publication(s) from which the primary data were extracted be cited as well.

4.2. Outstanding usage notes

Lastly, it is important to recognize the limitations of this database. Most importantly, the dataset is incomplete, in that there remain ~~many numerous~~ peat core sites that have not yet been integrated into this dataset. As a reminder, the C-PEAT GD v.2025 only includes ~~sites~~ ~~cores~~ that were already integrated to previously published data synthesis products. Second, and as mentioned in the text, not all peat core records are complete, meaning that some ~~of them~~ ~~cores~~ may not have reached the bottom of the peatland. ~~Along those lines~~ ~~Similarly~~, we remind the users that peat cores are not necessarily representative of an entire peat basin; rather, their record should be interpreted as a snapshot of an ecosystem’s history. Additionally, individual research teams might have had different ~~research~~ goals than the data users when they selected coring ~~sites~~ ~~locations~~ and collected their samples, such that some cores therein could represent the deepest portion of a peatland basin, but others might have been purposefully collected near ~~basin margins~~ ~~the edge~~. It is always advised to read the primary data source to gain further knowledge about the study sites ~~and core locations~~.

Appendices  
n/a

**Code and data availability**

495 The database is publicly available via the PANGAEA data repository (<https://doi.org/10.1594/PANGAEA.986891>; Loisel et al. 2025). The datasets can be explored and queried at [https://www.pangaea.de/?q=project:label:PAGES\\_C-PEAT](https://www.pangaea.de/?q=project:label:PAGES_C-PEAT). Code for accessing the data using Python and R is available on Zenodo (<https://doi.org/10.5281/zenodo.20734314><https://doi.org/10.5281/zenodo.17465602>; Ransby et al. 20265).

**Supplement link**

500 n/a

**Team list**

^C-PEAT Working Group Members and data contributors:  
A Korhola, A Lamarre, B Fiałkiewicz-Kozieł, B O'Reilly, B van Geel, C Bochicchio, C Tarnocai, D Anderson, D Hammarlund, D Large, D Mauquoy, D Peteet, DH Vitt, DJ Charman, DW Beilman, E Karofeld, E Klein, E-S Tuittila, F Chambers, F De Vleeschouwer, FJG Mitchell, G Le Roux, G Magnan, G Mallon, GM MacDonald, GT Swindles, I Bauer, IC Prentice, J Alm, J Bunbury, J Holmquist, J McCarroll, J Nichols, JA Christen, K Barber, L Ukonmaanaho, LR Belyea, M Amesbury, M Blaauw, M Galka, M Garneau, M Lamentowicz, M Lavoie, M Mäkilä, M Packalen, M Rundgren, M Turetsky, M Välranta, M van der Linden, MC Jones, N Phadtare, N Rausch, P Camill, P Kuhry, P Mathijssen, P Oksanen, PDM Hughes, PJH Richard, R Moschen, RK Booth, S Andersson, S Brewer, S Finkelstein, S Harrison, S Robinson, S van Bellen, ST Jackson, T Nieminen, T Ronkainen, T Thom, TR Moore, U Kokfelt, Ü Sillasoo, V Hohl, W Hinchcliffe, Y Corish, Y Huang, Z Yu, A Martínez Cortizas, ABK Sannel, C Courtney-Mustaphi, C Jones, C Massa, D Carless, E Cressey, E Githumbi, F Mitchell, G Wang, H Mackay, J Hribljan, J Uglow, J Zaragoza-Castells, K Marcisz, L Orme, M Clifford, N Sanderson, N Silva-Sánchez, N Steinberg, P Friedlingstein, P Moss, R Chimner, R de Jong, R Marchant, S Björck, S Moreton, SE Page, T Blyakharchuk, T Lacourse, T Mighall, T Roland, TE Turner, W Zhou, Y Zhao.

515 **Author contributions**

JL & AGS built and managed the C-PEAT GD, and coordinated the broader C-PEAT network; JL, AGS, SU & ER assembled or entered datasets and/or metadata into the database; DR, CB & SS performed quality control, term standardization, database cleaning, and/or QC certification; JL, KTB & VM analyzed data and generated figures for this paper; JL wrote [and revised](#) the manuscript; AGS, DR, SU, ER, CB, SS, KTB & VM reviewed and improved [a](#)-preliminary versions of this manuscript.

**Competing interests**

n/a

**Disclaimer**

525 Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to

include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

**Acknowledgements**

530 PAGES received support from the Swiss Academy of Sciences, the US National Science Foundation, and the Chinese Academy of Sciences at the time of database preparation and redaction of this manuscript.

**Financial support**

535 Support for this work includes NSF-2142177 and NSF-1802838 to JL. The C-PEAT working group acknowledges financial support from its parent organizations Past Global Changes (PAGES) and Future Earth, and support from the International Union of Quaternary Research (INQUA) and UNESCO’s International Geoscience Programme (IGCP) for workshops and training events.

**Review statement**

n/a

540 **References**

Adams J. 2012. The rise of research networks. *Nature*. 490(7420):335-6.

Alexandrov GA, Brovkin VA, Kleinen T, Yu Z. 2020. The capacity of northern peatlands for long-term carbon sequestration. *Biogeosciences*. 17(1):47-54.

545 Armentano TV, Menges ES. 1986. Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *The Journal of Ecology*. 755-74.

Baron JS, Specht A, Garnier E, Bishop P, Campbell CA, Davis FW, Fady B, Field D, Gross LJ, Guru SM, Halpern BS. 550 2017. Synthesis centers as critical research infrastructure. *BioScience*. 67(8):750-9.

Beck HE, McVicar TR, Vergopolan N, Berg A, Lutsko NJ, Dufour A, Zeng Z, Jiang X, van Dijk AI, Miralles DG. 2023. High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Scientific data*. 10(1):724.

555 Bond-Lamberty B, Smith AP, Bailey V. 2016. Running an open experiment: transparency and reproducibility in soil and ecosystem science. *Environmental Research Letters*. 11(8):084004.

Carpenter SR, Armbrust EV, Arzberger PW, Chapin III FS, Elser JJ, Hackett EJ, Ives AR, Kareiva PM, Leibold MA, 560 Lundberg P, Mangel M. 2009. Accelerate synthesis in ecology and environmental sciences. *BioScience*. 59(8):699-701.

Chadburn SE, Burke EJ, Gallego-Sala AV, Smith ND, Bret-Harte MS, Charman DJ, Drewer J, Edgar CW, Euskirchen ES, Fortuniak K, Gao Y. 2022. A new approach to simulate peat accumulation, degradation and stability in a global land surface scheme (JULES vn5. 8\_accumulate\_soil) for northern and temperate peatlands. *Geoscientific Model Development*. 15(4):1633-57.

Chamberlain S, Woo K, MacDonald A, Zimmerman N, Simpson G. 2021. pangaeac: Client for the 'Pangaea' Database. R package version 1.1.0, <https://CRAN.R-project.org/package=pangaeac>.

Charman DJ, Beilman DW, Blaauw M, Booth RK, Brewer S, Chambers FM, Christen JA, Gallego-Sala A, Harrison SP, Hughes PD, Jackson ST. 2013. Climate-related changes in peatland carbon accumulation during the last millennium. *Biogeosciences*. 10(2):929-44.

Chaudhary N, Westermann S, Lamba S, Shurpali N, Sannel AB, Schurgers G, Miller PA, Smith B. 2020. Modelling past and future peatland carbon dynamics across the pan-Arctic. *Global Change Biology*. 26(7):4119-33.

Cobb AR, Hoyt AM, Gandois L, Eri J, Dommain R, Abu Salim K, Kai FM, Haji Su'ut NS, Harvey CF. 2017. How temporal patterns in rainfall determine the geomorphology and carbon fluxes of tropical peatlands. *Proceedings of the National Academy of Sciences*. 114(26):E5187-96.

Dicks LV, Hodge I, Randall NP, Scharlemann JP, Siriwardena GM, Smith HG, Smith RK, Sutherland WJ. 2014. A transparent process for "evidence-informed" policy making. *Conservation Letters*. 7(2):119-25.

Dietze MC, Serbin SP, Davidson C, Desai AR, Feng X, Kelly R, Kooper R, LeBauer D, Mantooth J, McHenry K, Wang D. 2014. A quantitative assessment of a terrestrial biosphere model's data needs across North American biomes. *Journal of Geophysical Research: Biogeosciences*. 119(3):286-300.

Dommain R, Couwenberg J, Joosten H. 2011. Development and carbon sequestration of tropical peat domes in south-east Asia: links to post-glacial sea-level changes and Holocene climate variability. *Quaternary Science Reviews*. 30(7-8):999-1010.

Ellison AM. 2010. Repeatability and transparency in ecological research. *Ecology*. 91(9):2536-9.

Felden J, Möller L, Schindler U, Huber R, Schumacher S, Koppe R, Diepenbroek M, Glöckner FO. 2023. PANGAEA-Data publisher for earth & environmental science. *Scientific Data*. 10(1):347.

Fick, S.E. and R.J. Hijmans, 2017. WorldClim 2: new 1km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*. 37(12):4302-4315.

Frolking S, Talbot J, Subin ZM. 2014. Exploring the relationship between peatland net carbon balance and apparent carbon accumulation rate at century to millennial time scales. *The Holocene*, 24(9):1167-73.

Frolking S, Roulet N, Fuglestad J. 2006. How northern peatlands influence the Earth's radiative budget: Sustained methane emission versus sustained carbon sequestration. *Journal of Geophysical Research: Biogeosciences*, 111(G1).

Gallego-Sala AV, Charman DJ, Brewer S, Page SE, Prentice IC, Friedlingstein P, Moreton S, Amesbury MJ, Beilman DW, Björck S, Blyakharchuk T. 2018. Latitudinal limits to the predicted increase of the peatland carbon sink with warming. *Nature climate change*. 8(10):907-13.

- 610 Gibson CM, Chasmer LE, Thompson DK, Quinton WL, Flannigan MD, Olefeldt D. 2018. Wildfire as a major driver of recent permafrost thaw in boreal peatlands. *Nature communications*. 9(1):3041.
- Gorham E. 1991. Northern peatlands: role in the carbon cycle and probable responses to climatic warming. *Ecological applications*. 1(2):182-95.
- 615 Grand A, Wilkinson C, Bultitude K, Winfield AF. 2012. Open science: a new “trust technology”? *Science Communication*. 34(5):679-89.
- Hampton SE, Parker JN. 2011. Collaboration and productivity in scientific synthesis. *BioScience*. 61(11):900-10.
- 620 Harden JW, Hugelius G, Ahlström A, Blankinship JC, Bond-Lamberty B, Lawrence CR, Loisel J, Malhotra A, Jackson RB, Ogle S, Phillips C. 2017. Networking our science to characterize the state, vulnerabilities, and management opportunities of soil organic matter. *Global Change Biology*. 24(2):e705-18.
- 625 Hooijer A, Page S, Jauhiainen J, Lee WA, Lu XX, Idris A, Anshari G. 2012. Subsidence and carbon loss in drained tropical peatlands. *Biogeosciences*. 9(3):1053-71.
- Hoyt AM, Gandois L, Eri J, Kai FM, Harvey CF, Cobb AR. 2019. CO<sub>2</sub> emissions from an undrained tropical peatland: Interacting influences of temperature, shading and water table depth. *Global Change Biology*. 25(9):2885-99.
- 630 Hoyt AM, Chaussard E, Seppäläinen SS, Harvey CF. 2020. Widespread subsidence and carbon emissions across Southeast Asian peatlands. *Nature Geoscience*. 13(6):435-40.
- 635 Huber, R; Röttenbacher, J; Selke, N; CodeShredder; Spreckelen, F; Balamurugan, A; Lowe, S; Stocker, M; Kempf, D; Iris-Hinrichs; Schindler, U & qays abou housien. (2025). *pangaea-data-publisher/pangaeapy: v1.1.0 (v1.1.0)*. Zenodo. <https://doi.org/10.5281/zenodo.15749619>
- Hugelius G, Loisel J, Chadburn S, Jackson RB, Jones M, MacDonald G, Marushchak M, Olefeldt D, Packalen M, Siewert MB, Treat C. 2020. Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. *Proceedings of the National Academy of Sciences*. 117(34):20438-46.
- 640 Jones MC, Yu Z. 2010. Rapid deglacial and early Holocene expansion of peatlands in Alaska. *Proceedings of the National Academy of Sciences*. 107(16):7347-52.
- 645 Kleinen T, Brovkin V, Schuldt RJ. 2012. A dynamic model of wetland extent and peat accumulation: results for the Holocene. *Biogeosciences*. 9(1):235-48.
- Kock ST, Schitteck K, Mächtle B, Maldonado A, Vos H, Lupo LC, Kulemeyer JJ, Wissel H, Schäbitz F, Lücke A. 2020. Multi-centennial-scale variations of South American summer monsoon intensity in the southern central Andes (24–27 S) during the late Holocene. *Geophysical research letters*. 47(4):e2019GL084157.
- 650 Kurnianto S, Warren M, Talbot J, Kauffman B, Murdiyarso D, Frohling S. 2015. Carbon accumulation of tropical peatlands over millennia: a modeling approach. *Global Change Biology*. 21(1):431-44.
- 655 Loisel J, Gallego-Sala A. 2018. New research directions for the PAGES C-PEAT working group. *Past Global Changes Magazine*, 26(2): 91.

- Loisel J, Yu Z, Beilman DW, Camill P, Alm J, Amesbury MJ, Anderson D, Andersson S, Bochicchio C, Barber K, Belyea LR. 2014. A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation. *The Holocene*. 24(9):1028-42.
- Loisel J, van Bellen S, Pelletier L, Talbot J, Hugelius G, Karran D, Yu Z, Nichols J, Holmquist J. 2017. Insights and issues with estimating northern peatland carbon stocks and fluxes since the Last Glacial Maximum. *Earth-Science Reviews*. 165:59-80.
- Loisel J, Gallego-Sala AV, Amesbury MJ, Magnan G, Anshari G, Beilman DW, Benavides JC, Blewett J, Camill P, Charman DJ, Chawchai S. 2021. Expert assessment of future vulnerability of the global peatland carbon sink. *Nature Climate Change*. 11(1):70-7.
- Loisel J, Ransby D, Gallego-Sala AV (eds.) 2025. C-PEAT global database (v. 2025) [dataset editorial publication]. PANGAEA. <https://doi.org/10.1594/PANGAEA.986891>
- MacDonald GM, Beilman DW, Kremenetski KV, Sheng Y, Smith LC, Velichko AA. 2006. Rapid early development of circumarctic peatlands and atmospheric CH<sub>4</sub> and CO<sub>2</sub> variations. *Science*. 314(5797):285-8.
- Markowitz VM, Chen IM, Chu K, Pati A, Ivanova NN, Kyrpides NC. 2015. Ten years of maintaining and expanding a microbial genome and metagenome analysis system. *Trends in microbiology*. 23(11):730-41.
- Müller J, Joos F. 2021a. Global peatlands under future climate—seamless model projections from the last glacial maximum. *Biogeosciences*. 18:3657-87.
- Müller J, Joos F. 2021b. Committed and projected future changes in global peatlands—continued transient model simulations since the Last Glacial Maximum. *Biogeosciences*. 18(12):3657-87.
- Nichols JE, Peteet DM. 2019. Rapid expansion of northern peatlands and doubled estimate of carbon storage. *Nature Geoscience*. 12(11):917-21.
- Packalen MS, Finkelstein SA, McLaughlin JW. 2014. Carbon storage and potential methane production in the Hudson Bay Lowlands since mid-Holocene peat initiation. *Nature Communications*. 5(1):4078.
- Petrokofsky G, Kanamaru H, Achard F, Goetz SJ, Joosten H, Holmgren P, Lehtonen A, Menton MC, Pullin AS, Wattenbach M. 2012. Comparison of methods for measuring and assessing carbon stocks and carbon stock changes in terrestrial carbon pools. How do the accuracy and precision of current methods compare? A systematic review protocol. *Environmental Evidence*. 1:1-21.
- Piilo SR, Zhang H, Garneau M, Gallego-Sala A, Amesbury MJ, Väiranta MM. 2019. Recent peat and carbon accumulation following the Little Ice Age in northwestern Québec, Canada. *Environmental Research Letters*. 14(7):075002.
- Qiu C, Zhu D, Ciais P, Guenet B, Krinner G, Peng S, Aurela M, Bernhofer C, Brümmer C, Bret-Harte S, Chu H. 2018. ORCHIDEE-PEAT (revision 4596), a model for northern peatland CO<sub>2</sub>, water, and energy fluxes on daily to annual scales. *Geoscientific Model Development*. 11(2):497-519.
- Qiu C, Zhu D, Ciais P, Guenet B, Peng S. 2020. The role of northern peatlands in the global carbon cycle for the 21st century. *Global Ecology and Biogeography*, 29(5):956-73.
- Ransby, D; Riemann-Campe, K & Sanderson, N (2026<sup>65</sup>). C-PEAT data retrieval from PANGAEA (R & Python (2.1)). Zenodo. <https://doi.org/10.5281/zenodo.17465602> <https://doi.org/10.5281/zenodo.20734314>



- Reichman OJ, Jones MB, Schildhauer MP. 2011. Challenges and opportunities of open data in ecology. *Science*. 331(6018):703-5.
- Ruwaimana M, Anshari GZ, Silva LC, Gavin DG. 2020. The oldest extant tropical peatland in the world: a major carbon reservoir for at least 47 000 years. *Environmental Research Letters*. 15(11):114027.
- Smith LC, MacDonald GM, Velichko AA, Beilman DW, Borisova OK, Frey KE, Kremenetski KV, Sheng Y. 2004. Siberian peatlands a net carbon sink and global methane source since the early Holocene. *Science*. 303(5656):353-6.
- Spahni R, Joos F, Stocker BD, Steinacher M, Yu Z. 2013. Transient simulations of the carbon and nitrogen dynamics in northern peatlands: From the Last Glacial Maximum to the 21st century. *Climate of the Past*. 9(3):1287-308.
- Tolonen K, Turunen J. 1996. Accumulation rates of carbon in mires in Finland and implications for climate change. *The Holocene*. 6(2):171-8.
- Treat CC, Jones MC, Camill P, Gallego-Sala A, Garneau M, Harden JW, Hugelius G, Klein ES, Kokfelt U, Kuhry P, Loisel J. 2015. Effects of permafrost aggradation on peat properties as determined from a pan-Arctic synthesis of plant macrofossils. *Journal of Geophysical Research: Biogeosciences*. 121(1):78-94.
- Treat CC, Kleinen T, Broothaerts N, Dalton AS, Dommain R, Douglas TA, Drexler JZ, Finkelstein SA, Grosse G, Hope G, Hutchings J. 2019. Widespread global peatland establishment and persistence over the last 130,000 y. *Proceedings of the National Academy of Sciences*. 116(11):4822-7.
- Treat CC, Jones MC, Brosius L, Grosse G, Anthony KW, Frolking S. 2021. The role of wetland expansion and successional processes in methane emissions from northern wetlands during the Holocene. *Quaternary Science Reviews*. 257:106864.
- Turetsky MR, Abbott BW, Jones MC, Anthony KW, Olefeldt D, Schuur EA, Grosse G, Kuhry P, Hugelius G, Koven C, Lawrence DM. 2020. Carbon release through abrupt permafrost thaw. *Nature Geoscience*. 13(2):138-43.
- Turunen J, Tomppo E, Tolonen K, Reinikainen A. 2002. Estimating carbon accumulation rates of undrained mires in Finland—application to boreal and subarctic regions. *The Holocene*. 12(1):69-80.
- UNEP. 2022. Global Peatlands Assessment – The State of the World’s Peatlands: Evidence for action toward the conservation, restoration, and sustainable management of peatlands. Main Report. Global Peatlands Initiative. United Nations Environment Programme, Nairobi.
- Vines TH, Albert AY, Andrew RL, Débarre F, Bock DG, Franklin MT, Gilbert KJ, Moore JS, Renaut S, Rennison DJ. 2014. The availability of research data declines rapidly with article age. *Current Biology*. 24(1):94-7.
- Wolkovich EM, Regetz J, O'Connor MI. 2012. Advances in global change research require open science by individual researchers. *Global Change Biology*. 18(7):2102-10.
- Young DM, Baird AJ, Charman DJ, Evans CD, Gallego-Sala AV, Gill PJ, Hughes PD, Morris PJ, Swindles GT. 2019. Misinterpreting carbon accumulation rates in records from near-surface peat. *Scientific reports*. 9(1):17939.
- Young DM, Baird AJ, Gallego-Sala AV, Loisel J. 2021. A cautionary tale about using the apparent carbon accumulation rate (aCAR) obtained from peat cores. *Scientific Reports*. 11(1):9547.
- Yu Z. 2012. Northern peatland carbon stocks and dynamics: a review. *Biogeosciences*. 9(10):4071-85.

760 Yu Z, Loisel J, Brosseau DP, Beilman DW, Hunt SJ. 2010. Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters*. 37(13).

765 Yu Z, Loisel J, Turetsky MR, Cai S, Zhao Y, Frolking S, MacDonald GM, Bubier JL. 2013. Evidence for elevated emissions from high-latitude wetlands contributing to high atmospheric CH<sub>4</sub> concentration in the early Holocene. *Global Biogeochemical Cycles*. 27(1):131-40.