

1 Daily Snow Depth in the Southern Andes (2010–2024): A

2 Quality-Controlled Dataset from Chile and Argentina

3 Alexis Caro¹, Javier Medina^{2,3}, Freddy Saavedra^{2,3}, Fernando Gimeno^{4,5}, Jorge Huenante⁶, Mariano
4 Masiokas⁷, Cristian Orrego⁸, Ana Hernández^{2,3}, James McPhee^{9,10}, Pierre Pitte⁷, Sebastián Krogh¹¹,
5 David Farias-Barahona¹, Nevenka Bulovic¹², Iñigo Irarrazaval¹³, Shelley MacDonell^{8,14}, Mauricio
6 Zambrano-Bigiarini^{5,15}, Carlos Romero²

7

8 ¹Department of Geography, Universidad de Concepción, Concepción, Chile

9 ²Laboratorio de Teledetección Ambiental, Facultad de Ciencias Naturales y Exactas. Universidad de Playa Ancha,
10 Valparaíso, Chile

11 ³HUB Ambiental, Universidad de Playa Ancha, Valparaíso, Chile

12 ⁴Doctorado en Ciencias de Recursos Naturales, Universidad de la Frontera, Temuco, Chile

13 ⁵Center for Climate and Resilience Research, Universidad de Chile, Santiago, Chile

14 ⁶Dirección General de Aguas, Ministerio de Obras Públicas, Chile

15 ⁷Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA), CCT CONICET Mendoza, Argentina

16 ⁸Centro de Estudios Avanzados en Zonas Áridas (CEAZA), La Serena, Chile

17 ⁹Department of Civil Engineering, University of Chile, Santiago, Chile

18 ¹⁰Advanced Mining Technology Center, University of Chile, Santiago, Chile

19 ¹¹Departamento de Recursos Hídricos, Universidad de Concepción, Campus Chillán, Chillán, Chile

20 ¹²Sustainable Minerals Institute, The University of Queensland, Brisbane, Australia

21 ¹³Centro de Investigación en Ecosistemas de la Patagonia, Coyhaique, Chile

22 ¹⁴Waterways Centre, University of Canterbury, Christchurch, New Zealand

23 ¹⁵Department of Civil Engineering, Universidad de la Frontera, Temuco, Chile

24

25 Correspondence to: Alexis Caro (alexis.caro.paredes@gmail.com), and Freddy Saavedra (freddy.saavedra@upla.cl)

26

27

28

29

30

31 Abstract.

32 Seasonal snow is a critical component of the water cycle in the Southern Andes of Chile and Argentina. Quantitative
33 assessments of snow accumulation remain constrained by the scarcity, heterogeneity, and inconsistency of *in situ*
34 observations, resulting in substantial uncertainties in mountain hydrological modeling. To address this gap, we compiled and
35 quality-controlled snow depth observations to produce a consistent daily dataset from 81 stations between 21°S and 54°S
36 covering 2010–2024. Our quality-control procedure involved data compilation and preprocessing, followed by
37 harmonization of snow depth observations, primarily by correcting the ground reference level relative to the soil surface
38 during snow-free periods and removing anomalous spikes and observations outside physically plausible ranges. This process
39 improved data reliability, increasing the Physical Consistency Index (PCI) from 87 % to 95 % at selected stations in the
40 Maipo River Basin, while reducing the median number of days with data across stations by 23% (from 1,392 to 1,074 days).
41 The snow depth data availability increased from one station in 2010 to 57 stations in 2024, largely driven by the expansion
42 of the monitoring network operated by the General Directorate of Water (DGA), Chile. However, this expansion remains
43 spatially uneven across the Andean zones. The Mediterranean Andes have the largest number of stations (39), as well as the
44 largest number of stations with highly complete records with 17 stations reaching 80–100 % data coverage, compared with
45 only nine stations each in the Arid and Wet Andes, highlighting persistent spatial and temporal gaps. Using this newly
46 quality-controlled dataset, we find that snow depth generally increases from the Arid to the Wet Andes in association with
47 increasing precipitation, whereas its relationship with elevation is not consistently positive. The snow depth–elevation
48 relationship is nonlinear in the Arid and Mediterranean Andes, with maximum observed snow depths at 4,300 m a.s.l. in the
49 Elqui River Basin and 3,300 m a.s.l. in the Maipo River Basin. In contrast, the available observations indicate a positive
50 snow depth–elevation relationship in the Wet Andes (Maule–Itata River Basin). However, these relationships should be
51 interpreted in the context of the specific spatial and temporal conditions represented by the available observations. This
52 open-access, quality-controlled snow depth dataset (Medina and Caro, 2026, <https://zenodo.org/records/21577171>)
53 represents the largest and most complete collection of continuous snow depth observations for the Southern Andes and
54 provides a basis for hydrological applications, reanalysis evaluation, and seasonal streamflow forecasting.

55 1 Introduction

56 Seasonal snowpacks are a key component of mountain hydrology, storing winter precipitation and releasing water during the
57 dry season, thereby sustaining downstream water resources (Stewart et al., 2009). Although snow accumulation generally
58 occurs during winter, in tropical regions it can also occur during summer, in response to the seasonal precipitation regime
59 (Caro et al., 2021). However, quantifying snow storage and its spatial variability remains challenging due to the limited
60 availability and uneven distribution of *in situ* observations (Masiokas et al., 2020). This limitation is particularly pronounced
61 in the Southern Andes (20° S–55° S), where long-term, quality-controlled observations of snow depth (SD) and snow water
62 equivalent (SWE) remain scarce and fragmented across monitoring networks. In the Arid and Mediterranean Andes, the

63 seasonal snowpack is a fundamental hydrological resource, providing water for human consumption and supporting
64 ecosystems, agriculture, hydropower generation, and industrial activities including mining (SERNAGEOMIN, 2021;
65 Masiokas et al., 2020).

66 The hydrological role of the seasonal snowpack varies across Andean climatic zones. In the Mediterranean Andes (31°
67 S–36° S), seasonal and interannual streamflow variability of the main mountain rivers is primarily controlled by snowmelt,
68 with secondary contributions from glacier melt and minimal influence from rainfall. As a result, river discharge exhibits a
69 relatively simple unimodal regime, with peak flows occurring in late spring and early summer (Masiokas et al., 2016, 2019;
70 Burger et al., 2019; Ayala et al., 2020). In contrast, in the Wet Andes (36° S–55° S), snowmelt accounts for approximately 26
71 % of the annual runoff (Krogh et al., 2015), and mountain rivers exhibit more complex seasonal patterns influenced by
72 precipitation events throughout the year. Since 2010, the Arid and Mediterranean Andes have experienced a prolonged
73 megadrought characterized by precipitation deficit and reduced snow accumulation (Garreaud et al., 2017; 2020). This has
74 led to reduced river discharge, decreasing water availability and increased conflicts over water rights as demand increasingly
75 exceeds supply (Rivera et al., 2016; Álvarez–Garretón et al., 2021). Although glacier melt has partially buffered streamflow
76 reductions during the ongoing megadrought, sustaining runoff despite significant precipitation deficits, this compensatory
77 effect is associated with accelerated ice loss. Its contribution is therefore expected to diminish in the future, potentially
78 exacerbating water scarcity during prolonged dry conditions (Ayala et al., 2025; Caro et al., 2025).

79 SD and SWE are key variables for quantifying the hydrological contribution of the seasonal snowpack. While SWE is more
80 directly related to water storage than SD, it is considerably more difficult to measure. SWE can be measured directly or
81 estimated by combining SD with measured or modeled snow density values (Ntokas et al., 2021). Specifically, the
82 end-of-winter spatial distribution of SD and SWE are crucial for forecasting spring and summer streamflow in
83 high-mountain regions (Shaw et al., 2020a). However, because *in situ* snow observations are typically sparse and mostly
84 located at low elevations, snow cover, SD, and SWE are commonly estimated using spatial extrapolation supported by
85 satellite imagery and modeling approaches (Cornwell et al., 2016; Cortés and Margulis, 2017; Shaw et al., 2020a; Bulovic et
86 al., 2025; Saavedra et al., 2026). In Chile, these limitations are further exacerbated for SWE, as observations are substantially
87 more restricted than SD data in terms of spatial density and temporal coverage, with large data gaps. In Argentina, the
88 number of stations reporting SWE data is also low, with most sites concentrated in the Dry Andes where snowmelt
89 represents a crucial water resource for the adjacent lowlands. Regular daily SWE observations began in the early twenty-first
90 century, although manual winter measurements from a limited number of snow courses date back to the early 1950s.

91 SD monitoring networks in the Southern Andes have expanded in recent years, increasing spatial coverage across a wider
92 range of latitudes and elevations. In Chile, the growing demand for spatially distributed snow information has driven
93 government agencies and private entities to expand these monitoring networks. The General Directorate of Water (Dirección
94 General de Aguas, DGA) has consolidated national SD observations since 2010, while research institutions and universities,

95 including the Centro de Estudios Avanzados en Zonas Áridas (CEAZA) and the Department of Civil Engineering at the
96 Universidad de Chile (UdeChile), operate stations in the Arid and Mediterranean Andes. In the Wet Andes, monitoring
97 efforts are led by the Centro de Estudios Científicos (CECs) and the Centro de Investigación en Ecosistemas de la Patagonia
98 (CIEP). However, SWE observations from CEAZA, UdeChile, and CIEP are either unavailable or limited to short and
99 discontinuous periods and, therefore, SWE records remain largely restricted to a small number of DGA stations. In
100 Argentina, several institutions have expanded snow monitoring networks across the Andes, including the Instituto Argentino
101 de Nivología, Glaciología y Ciencias Ambientales (IANIGLA), which has monitored SD variations near selected glaciers
102 since 2014. Other institutions, typically responsible for water management at the provincial and regional levels, conduct
103 SWE observations to support seasonal streamflow forecasts using automatic sensors (snow pillows, snow scales) and
104 sporadic manual measurements at selected sites.

105 Publicly available SD data face challenges related to data acquisition, heterogeneous formats, and limited quality control,
106 reducing their suitability for technical and scientific analyses. Existing regional studies of the Southern Andes (Cornwell et
107 al., 2016; Cortés and Margulis, 2017; Bulovic et al., 2025), *in situ* observations are limited to only 12–26 sites. Considering
108 all stations included in these studies, the interquartile range spans 32.4° S–34.8° S in latitude and approximately 2,500–3,600
109 m a.s.l. in elevation, indicating a strong concentration of observations at mid-elevations in central Chile–Argentina Andes.
110 However, these datasets have been used to extrapolate SD and SWE conditions across a much broader region (29° S–44° S)
111 of the Andes, encompassing climatically contrasting mountain basins and elevations well beyond those represented by the
112 observation network (e.g., Bulovic et al., 2025), including high-elevation areas where seasonal snow accumulation is critical
113 for sustaining downstream water resources during the dry season (Masiokas et al., 2020).

114 This limited spatial and elevational coverage introduces substantial uncertainty in characterizing how snow depth varies with
115 elevation. While some studies in the Southern Andes report positive SD–elevation gradients from *in situ* observations
116 (Cornwell et al., 2016; Cortés and Margulis, 2017), evidence from other mountain regions suggests more complex,
117 non-linear relationships, with SD increasing toward a peak and decreasing at the highest elevations due to precipitation
118 patterns and snow redistribution processes such as wind transport, sloughing, and avalanching (Grünwald et al., 2014).
119 These contrasting findings highlight the lack of observational constraints at high elevations and limit our ability to robustly
120 assess snow storage across elevation gradients in the Southern Andes.

121 Despite valuable progress in SD and SWE observations across the Southern Andes, estimating SWE from basin to regional
122 scales remains challenging due to a restricted observational record. Latitudinal and elevational sampling biases limit the
123 representativeness of current datasets, directly impacting the reliability of regional simulations. These challenges are
124 exacerbated by the lack of a coordinated regional framework to date; no initiative has produced a homogenized
125 quality-controlled SD dataset for the Southern Andes. This hinders our understanding of mountain hydrological processes
126 and their variability across elevations and basins in the Southern Andes. It also limits hydrological model calibration and

validation and increases uncertainty in climate change impact projections by hampering the generation of robust hydrological simulations, particularly for extreme years.

In this study, we provide the first daily, homogenized, quality-controlled SD dataset for the Southern Andes. The dataset covers 2010–2024 and integrates observations from automatic stations operated by government agencies, research centers, and universities across Chile and Argentina. Quality control was performed using a three-step Python-based algorithm. To illustrate its scientific utility, we further analyze snow depth–elevation relationships across river basins and climatic zones, addressing a key but still uncertain aspect of snow accumulation patterns in the Andes. This comprehensive and consistent dataset provides an observational basis for improving model calibration and seasonal streamflow forecasting, and for strengthening short- and long-term projections of water availability at local and regional scales under climate change scenarios.

2 Study area

The Southern Andes can be divided into three major climatic zones: the Arid, Mediterranean, and Wet Andes (Saavedra et al., 2017; Sarricolea et al., 2017; Caro et al., 2021). These zones are shown in Figure 1 across the study area (21° S–54° S), together with the distribution of the SD stations analyzed in this study. In the northern Arid Andes, precipitation predominantly occurs during autumn (MAM) and winter (JJA). However, in its northernmost sector, significant precipitation also occurs during the austral summer (DJF), associated with the Bolivian Winter. At high elevations, air temperatures typically can remain below freezing for several months, reaching minimum values of -29 °C at Guanaco Glacier (5,324 m a.s.l., 2008–2011) (MacDonell et al., 2013). The Mediterranean Andes encompass the most densely populated areas of Chile (65 % of the country's total population from the Valparaíso to Ñuble regions, INE, 2024), as well as 74 % of the country's agricultural land, and also support the country's main electricity generation (INE, 2022). Further south in the Wet Andes, the mountain range transitions to a wetter climate influenced by Pacific frontal systems. Although mean monthly precipitation also peaks in autumn–winter, it remains substantial through spring–summer (SON–DJF), exceeding 150 mm month⁻¹ at the Lago Vargas station (1977–1991) (Dussaillant et al., 2012).

Across Chile and Argentina, persistent snow cover, defined as areas where seasonal snow remains on the ground for several months each year, covers approximately 34,370 km² between 29° S and 36° S (Saavedra et al., 2018). Mountain river basins in Chile generally drain westward to the Pacific Ocean, whereas those in Argentina drain eastward to the Atlantic Ocean. Hydrological regimes also vary latitudinally. Southern basins are dominated by winter precipitation, with secondary contributions from spring snowmelt, producing a pluvio–nival regime. In contrast, northern basins rely mainly on spring and early summer snowmelt and, particularly on the Chilean side, to a lesser extent on winter precipitation, resulting in predominantly nival to nivo–pluvial regimes (Masiokas et al., 2019). These spatial patterns are governed by large-scale atmospheric circulation, as well as by latitudinal gradients, elevation, and local physiographic controls.




Figure 1. Spatial distribution of the 81 snow depth stations retained after quality control across 26 river basins and three Andean climatic zones (Arid, Mediterranean, and Wet). Station locations are identified according to the contributing institutional networks (DGA, IANIGLA, CEAZA, UdeChile, and CIEP), together with the 37 stations excluded during the quality-control procedure. The Elqui-Jáchal, Maipo-Mendoza, and Maule-Itata river basins, located in each climate zone, are analyzed in greater detail in subsequent sections. Basemap imagery: Esri World Imagery © Esri, Maxar, Earthstar Geographics, and the GIS User Community | Powered by Esri.

158 **3 Quality-control procedure**

159 The complete data-processing and quality-control workflow is summarized in Figure 2. This procedure comprises three
160 sequential steps, ranging from data compilation (Step 1) to harmonization of observations (Step 3). The resulting dataset was
161 subsequently evaluated using an independent cross-variable validation based on concurrent snow depth, precipitation, and
162 air temperature observations. Finally, SD-elevation relationships were analyzed at the river-basin scale.




Figure 2. Workflow of the data processing and quality-control procedure applied to the 2010–2024 snow. The logos at the top indicate the institutional networks that contributed the snow depth observations used to develop the final quality-control dataset.

163 **3.1 Step 1 – Compilation of snow depth observations**

164 This study compiled SD observations from 118 stations distributed across the Southern Andes (17° S–55° S), including both
165 hourly and daily records, although hourly observations were not available at all stations. In addition, precipitation (Pr) and
166 air temperature (AT) data were compiled to assess the physical consistency of the SD records at stations located in the Elqui
167 and Maipo river basins (Table 3). Data were obtained from DGA, CEAZA, UdeChile, and CIEP in Chile, and from
168 IANIGLA in Argentina.

169 SD observations were obtained either through direct requests or by downloading publicly available datasets. The original
170 data for Chile are available from <https://snia.mop.gob.cl> (DGA), <https://www.ceazamet.cl> (CEAZA),
171 http://190.121.23.217/index_cdataaysen.php (CIEP) and Huerta et al., (2019). The data available for Argentina were
172 provided by IANIGLA and is partially available from <https://observatorioandino.com/estaciones/>. Given the heterogeneous
173 structure of these sources, the data were reorganized into a tabular format, with each column representing a monitoring
174 station and each row containing the daily mean values for SD. This initial data compilation was consolidated into a single
175 raw data file.

176 **3.2 Step 2 – Preprocessing of snow depth observations**

177 An initial visual screening was conducted to identify and exclude stations with records unsuitable for subsequent quality
178 control and analysis based on the following criteria: (i) insufficient monthly data availability (less than 50 % of daily
179 observations), and (ii) insufficient annual coverage (fewer than three months with available data during the snow season,
180 from May to October). In addition, several stations were excluded because their records only became available in late 2025
181 or 2026. Following this procedure, 81 of the 118 stations with SD observations were retained, providing coverage across the
182 Andes between 21° and 54° S. The selected stations span elevations from 30 to 5,804 m a.s.l., with records covering the
183 period 2010–2024. Table 1 summarizes the main characteristics of the compiled SD dataset aggregated by river basin, while
184 metadata for the selected and excluded stations are provided in Tables A1 and A2, respectively.

Table 1. Characteristics of snow depth stations by river basin across the Southern Andes (21° S–54° S). CL and AR refer to Chile and Argentina, respectively.					
Andean zone	River Basin (Country)	N° Stations	Latitude (median, °S)	Station elevation (median, m a.s.l.)	Sources
Arid Andes	Salar Michincha (CL)	1	21.2	5,087	DGA
	Salar Atacama (CL)	1	23.0	5,625	DGA
	Tres Cruces (CL)	2	27.1	5,503	DGA
	Copiapó (CL)	1	28.5	3,985	DGA
	Jáchal (AR)	1	30.2	4,754	IANIGLA
	Elqui (CL)	7	30.2	3,660	CEAZA, DGA
	Limarí (CL)	3	30.9	3,529	DGA, CEAZA
	Choapa (CL)	2	31.7	3,432	CEAZA, DGA

Mediterranean Andes	Petorca (CL)	1	32.2	3,143	DGA
	Aconcagua (CL)	2	32.9	3,691	DGA
	Mendoza (AR)	2	33.2	3,544	IANIGLA
	Maipo (CL)	21	33.3	3,340	DGA, UdeChile
	Rapel (CL)	7	34.7	2,665	DGA
	Mataquito (CL)	1	35.2	2,602	DGA
	Tunuyán (AR)	1	35.3	2,453	IANIGLA
Wet Andes	Maule (CL)	4	36.0	1,993	DGA
	Itata (CL)	5	36.9	1,809	UdeChile, DGA
	Biobio (CL)	5	37.6	1,784	DGA
	Valdivia (CL)	2	39.7	1,683	DGA
	Bueno (CL)	1	40.7	920	DGA
	Yelcho (CL)	1	43.3	575	DGA
	Aysén (CL)	2	45.6	1,310	CIEP, DGA
	Costeras e Islas R. Aisén R. Baker (CL)	1	46.7	1,187	CIEP, DGA
	Baker (CL)	2	46.8	555	DGA
	Santa Cruz (AR)	4	49.3	984	IANIGLA
	Islas sur Estrecho de Magallanes (CL)	1	53.7	70	DGA

185 A series of automated algorithms implemented in Python using Pandas library (The pandas development team, 2025) were
186 applied to perform three quality-control procedures: adjustment of the ground reference level (Step 2.1, Zero-level
187 correction), corresponding to the soil surface established after sensor installation or maintenance; removal of anomalous
188 spikes (Step 2.2, Spike removal); and identification and removal of implausible or physically unrealistic values (Step 2.3,
189 Physical range check). For these automated procedures, parameters and thresholds were calibrated at the river-basin scale to
190 account for regional SD variability and ensure physical consistency across stations sharing similar climatic environments. In
191 addition to this basin-level processing, a supplementary manual filtering step was applied on a station-by-station basis.

192 **Step 2.1. Zero-level correction:** This step realigns the ground reference level to account for instrumental drift during
193 snow-free periods. The algorithm detects stable periods using a moving-window range (the difference between maximum
194 and minimum SD) with a 0.5 cm stability threshold. Window lengths were calibrated by river basin (3–5 days; median: 5
195 days) to balance sensitivity to instrumental drift against short-term SD variability. From these stable windows, we defined a
196 ‘validated offset’ as the calculated displacement from the ground reference level. To protect actual shallow snow records,
197 windows with median values exceeding a 5 cm snow threshold are preserved and not used for correction. Stable windows
198 with median SD >5 cm were excluded from offset estimation to avoid misclassifying shallow, persistent snow as
199 instrumental drift. The 5 cm threshold was selected conservatively to exceed both nominal sensor noise (± 1 cm) and typical
200 surface-roughness effects (2 cm). If the validated offset is very small (within 0.3 cm), it is treated as noise and set to zero.
201 Finally, to maintain data continuity, this offset is subtracted from subsequent observations only if the drift exceeds a 0.3 cm
202 tolerance. Although sensor accuracy is ± 1 cm (<https://www.campbellsci.com/sr50a>), our use of daily averages from
203 high-frequency (hourly) observations over multi-day windows dampens random noise. This reduction in variance justifies

the stricter 0.3 cm threshold, enabling the detection of subtle systematic shifts that would otherwise be obscured by instrumental error.

In addition, the SR50A sensors, which have been widely used across stations in Chile and Argentina, have recently begun to be replaced at some stations by a newer generation of sensors with higher measurement precision. The sensor used at each station is indicated in Table A1.

Step 2.2. Spike removal: To identify and remove isolated spikes in the daily mean SD time series, the median absolute deviation (MAD, Rousseeuw and Croux, 1993) was applied using a centered moving window. The window size (ω) was adjusted for each basin within a range of 3 to 15 days to account for local SD variability. Shorter windows increase sensitivity to local anomalies in stable snowpacks, whereas longer windows provide a more robust baseline under highly variable conditions. For the Arid Andes, window sizes typically ranged from 3 to 9 days (median: 3), while the Mediterranean and Wet Andes required longer windows (ranging from 6 to 15 days and 9 to 15 days, respectively) to obtain a robust moving median under greater SD variability. For a given day t , the MAD is calculated as:

$$MAD_t = median_w \left(\left| SD_i - \widetilde{SD}_i \right| \right) \quad \text{Equation 1}$$

where SD_i represents the daily observation within the sliding window ω centered at day t , and $\widetilde{SD}_i$ is the median of that window.

A daily observation was flagged as a spike and removed if it met either of two independent criteria. The first criterion evaluated deviations from the window median using a scaling factor, k . Conceptually, k acts as a tolerance threshold that defines the acceptable range of natural variability around the median. Because snow accumulation is inherently irregular, k was adjusted to balance the removal of instrumental errors with the preservation of true snowfall events. This adjustment was determined through expert judgment, integrating visual inspection of time series with local knowledge of station-specific signal-to-noise ratios and SD variability. In this context, a lower k value provides a stricter filter for noise removal, while a higher k value is more permissive to accommodate natural variability and ensure that extreme physical signals are not erroneously discarded. Across the study area, k values ranged from 4 to 7, with median values calculated across basins being 4 in the Arid and Mediterranean Andes and 6 in the Wet Andes. To avoid inadvertently removing true extreme snowfall events, this criterion required that any suspected spike exhibit clear discontinuities ($> k \cdot MAD$) relative to both adjacent days. Additionally, the absolute difference between the two adjacent days had to remain small ($< k \cdot MAD$), confirming that the event represented an isolated anomaly rather than a sustained change in SD. The second criterion acted as an absolute physical filter. Regardless of MAD, an observation was removed if the absolute difference relative to both adjacent days

231 exceeded a basin-specific threshold representing an implausibly large day-to-day change in SD. These thresholds were
232 established through expert judgment based on visual inspection of station records, physical plausibility, and inter-station
233 consistency within each basin. Given the steep latitudinal precipitation gradient across the Andes, threshold values ranged
234 from 20 to 150 cm (median: 50 cm) in the Arid Andes, 100 to 200 cm (median: 100 cm) in the Mediterranean Andes, and 60
235 to 350 cm (median: 140 cm) in the Wet Andes.

236 **Step 2.3. Physical range check:** A physical range filter was applied to the SD time series at each station.

237 Lower and upper plausible thresholds were defined for each river basin based on station elevation, regional climatology, and
238 expert knowledge. Observations below the ground reference level (defined in Step 2.1) were set to zero, whereas values
239 exceeding the upper physical thresholds were considered implausible and removed. The median physical limits for basins
240 were 0–140 cm in the Arid Andes, 0–360 cm in the Mediterranean Andes, and 0–300 cm in the Wet Andes. In some stations,
241 the lower threshold in the Mediterranean and Wet Andes was set to 2 cm to account for sensor noise (± 1 cm) and local
242 surface roughness (e.g., low vegetation or rocky terrain). Setting these near-zero values to zero prevents instrumental and
243 surface-related noise from being misinterpreted as shallow snow accumulation.

244 3.3 Step 3 – Harmonization of snow depth observations

245 When overlapping daily SD observations exist for the same station due to data being compiled from multiple sources, a
246 single unified time series is constructed. For example, the DGA dataset for some stations may include multiple versions of
247 the same SD observation, arising either from data updates or from the incorporation of post-processed information obtained
248 from external sources for specific stations (e.g., universities or other monitoring agencies). This step is particularly important
249 for any future updates of the quality-controlled SD dataset, as revised data releases from the original data providers may
250 contain different observations for the same station and date than those included in earlier versions. Such differences may
251 arise from technical decisions by data providers or from sensor replacements affecting SD observations.

252 The output of this step is a cleaned SD file that preserves the structure of the original raw SD dataset while containing
253 quality-controlled values for each monitoring station.

254 3.4 Cross-variable validation of snow depth observations

255 A multivariable validation procedure was applied to assess the physical consistency of both raw and quality-controlled SD
256 observations. The analysis was restricted to stations in the Elqui and Maipo River Basins where daily observations of SD, Pr,
257 and AT were simultaneously available at the same station during the May–October period, corresponding to the snow
258 accumulation season in the Southern Andes (Masiokas et al., 2020; Bulovic et al., 2025). The approach is based on the joint
259 physical consistency among these variables. Snow accumulation, defined as a positive day-to-day change in SD (positive
260 Δ SD), is expected to occur during precipitation events under sufficiently cold air temperature conditions. In contrast, snow

261 ablation (negative ΔSD) may result from multiple processes (e.g., melt, sublimation, compaction, or wind redistribution),
262 which are not directly observed at most weather stations. Accordingly, the validation was restricted to snow accumulation
263 events.

264 We developed and implemented a new Physical Consistency Index (PCI) to classify each day as physically consistent ($\phi_i = 1$) or inconsistent ($\phi_i = 0$), following a multivariable framework inspired by the Perkins Skill Score (PSS, Perkins et al. 2008) and the Automated Quality assurance of daily surface observations procedures (AQ, Durre et al., 2010). This
267 procedure filters the record to identify the robust snowfall signals that collectively constitute the index denominator (N).
268 First, to mitigate measurement noise, a station-specific threshold (τ) was established as the 5th percentile of the positive
269 ΔSD distribution, with a minimum of 1 cm to align with sensor accuracy. A day was retained as a potential snow
270 accumulation event only when $\Delta SD > \tau$; consequently, all other observations, including negligible shifts ($\Delta SD \leq \tau$), stable
271 snowpacks ($\Delta SD = 0$), and ablation periods (negative ΔSD), were excluded from the denominator to ensure the index targets
272 active snowfall signals. Second, to ensure physical consistency during snow accumulation events, we retained only days with
273 measurable precipitation ($Pr > 1$ mm), thereby reducing uncertainties associated with gauge undercatch and wind-driven
274 snow redistribution. For these events, an air temperature threshold (AT_{rain}) was defined as the station-specific 95th percentile
275 of AT to account for the large climatic variability across the Andes. A fixed usual threshold (e.g., 0–2 °C) was not adopted
276 because the air temperature associated with snowfall varies substantially with local conditions. Using a station-specific
277 percentile therefore provides a locally adapted threshold that better represents the upper temperature limit at which snowfall
278 is still observed at each station. For example, the median air temperature threshold ranged from -3.2 to 2.2 °C (median: -2.2
279 °C) in the Elqui River Basin and from -3.8 to 3.8 °C (median: -1.5 °C) in the Maipo River Basin. Accumulation days with
280 AT exceeding this threshold were classified as physically inconsistent ($\phi_i = 0$), whereas those with AT below the threshold
281 were considered physically consistent and retained as valid snow accumulation signals ($\phi_i = 1$). The threshold is therefore
282 intended to identify potential outliers in the observed snow accumulation signal rather than to represent a physically derived
283 rain-snow transition temperature.

284 The PCI for each station was calculated as the ratio between the number of physically consistent snow accumulation days ($\phi_i = 1$) and the total number of identified snow accumulation days with $Pr > 1$ mm. This multivariable equation specifically
286 targets the consistency of precipitation driven accumulation. By excluding days without precipitation from the denominator
287 we minimize uncertainties associated with wind driven snow redistribution and instrumental noise, which do not represent
288 true snowfall signals. This index is defined in Equation 2, where ϕ_i is a binary consistency indicator for each day i , taking a
289 value of 1 for physically consistent conditions and 0 otherwise.

$$PCI = \frac{1}{N} \sum_{i=1}^N \phi_i \quad , \quad \text{Equation 2}$$

where ϕ_i is the consistency indicator for each i daily snow accumulation day, defined as:

$$\phi_i = \begin{cases} 1, & \text{if } (\Delta SD_i > \tau) \wedge (Pr_i > 1 \text{ mm}) \wedge (AT_i \leq AT_{rain}) \\ 0, & \text{otherwise} \end{cases}$$

While the PCI provides a relative measure of the physical consistency of snow accumulation events, its interpretation in isolation can be misleading. As a ratio, highly noisy raw SD observations may artificially inflate PCI values, in some cases making them comparable to or even higher than those derived from the quality-controlled SD dataset. To complement the PCI, two additional metrics were developed in this study and computed using the same event definition and filtering criteria described in Equation 2. First, the Net Inconsistency Reduction (ΔE) is defined in Equation 3 as the difference between the number of physically inconsistent days detected in the raw (N_{raw}) and quality-controlled SD datasets (N_{QC}), with positive values indicating a reduction in inconsistencies. Second, the Noise Ratio (NR) is defined in Equation 4 as the proportion of the total number of apparent accumulation events detected in the raw dataset ($N_{event, raw}$) to the number of events retained after quality-control ($N_{event, QC}$). The interpretation of this metric allows for three scenarios: (1) $NR > 1$ indicates that the quality-control procedures up to Step 3 effectively identified and removed spurious positive SD spikes, revealing the presence of noise or measurement artifacts in the raw SD dataset; (2) $NR = 1$ suggests a high-quality raw record where no events were removed or recovered; and (3) $NR < 1$ indicates signal recovery, where the cleaning algorithm or manual quality control allowed the identification of valid accumulation events that were previously obscured or incorrectly recorded in the raw data. For example, a station with $NR = 2.0$ has had half of its raw accumulation signals removed as noise, whereas a station with $NR = 0.5$ has doubled its count of valid events through the correction of systematic errors.

$$\Delta E = N_{raw} - N_{QC} , \quad \text{Equation 3}$$

$$NR = \frac{N_{event, raw}}{N_{event, QC}} \quad \text{Equation 4}$$

The validation was restricted to the Elqui and Maipo River Basins because these basins provide suitable conditions for evaluating the physical consistency of snow accumulation using concurrent SD, Pr, and AT observations. In particular, they contain stations with overlapping observations of the three variables and sufficient altitudinal coverage to evaluate snow accumulation under contrasting meteorological conditions. These characteristics are essential for applying the proposed multivariable validation consistently across stations and for reducing the influence of spatial sampling limitations. The

validation could not be extended to other regions, particularly the Wet Andes, because the availability of concurrent SD, Pr, and AT observations is more limited and unevenly distributed along the elevation gradient. This limitation is particularly evident for SD in the Maule–Itata basins, as shown later in the study.

3.5 Snow depth and elevation analysis by river basin

Due to the limited temporal continuity of SD observations across stations located at different elevations and in different river basins for the same dates, the observations were grouped into three precipitation year types. Based on precipitation records from the General Directorate of Water (DGA) and the Dirección Meteorológica de Chile (DMC), we identified, for three basins, years representative of different ranges of total annual precipitation: dry [0th to 30th percentile], normal (30th to 70th percentile], and wet (70th to 100th percentile] (Table A3). For this classification, precipitation records were obtained from relatively high–elevation stations within each basin to better represent mountain precipitation conditions. Specifically, we used the La Laguna Embalse station (3,160 m a.s.l.) in the Elqui River Basin, 19 stations in the Maipo River Basin, and the Diguillín (670 m a.s.l.), Río Diguillín en San Lorenzo–Atacalco (727 m a.s.l.), and Las Trancas stations (1,242 m a.s.l.) in the Maule–Itata River Basin. These basins were selected because they contain a sufficient number of SD observations spanning a broad elevation range.

A positive correlation between elevation and precipitation is generally expected, reflecting the influence of orographic enhancement of precipitation (Scaff et al., 2017). In contrast, mean air temperature is expected to exhibit a negative correlation with elevation, with lower mean temperatures typically observed at higher elevations (Voordendag et al., 2021; Aguayo et al., 2024; Caro et al., 2024). To quantify the combined influence of these elevation–dependent processes, the SD–elevation relationship was analyzed using the 25th, 50th, and 75th percentiles, as well as the cumulative distribution function (CDF), across three precipitation year types, providing a robust basis for comparing SD across elevations and contrasting climate conditions in the Andean zones.

To analyze the SD–elevation relationship, only stations with at least 70 % of daily SD data availability per month during the accumulation period (1 May to 31 October) were retained (Alvial Vásquez et al., 2020; Cordero et al., 2024), and elevation was derived from the 30 m SRTM digital elevation model (Far et al., 2007). This approach enabled consistent comparison of SD observations across elevations within each precipitation year type. Additionally, the median July snow line elevation, together with its 25th percentile, was estimated for these three river basins using satellite-derived snow persistence for the period 2000–2024 to provide a reference for evaluating the elevational distribution of the SD observations. July corresponds to the month with the lowest snow line elevation in the year.

Monthly snow persistence was derived from daily MODIS Terra and Aqua snow products (MOD10A1/MYD10A1, Collection 6.1; 500 m spatial resolution), processed in Google Earth Engine and reported by Saavedra et al. (2026). Snow presence (1) and absence (0) were defined using a normalized difference snow index (NDSI) threshold of 0.4. Elevation was

341 extracted from the same digital elevation model described above. Following Saavedra et al. (2026), the snow line elevation
342 was defined as the 5 % snow persistence contour. Its elevation was then estimated by intersecting this contour with the DEM
343 and calculating the median elevation of the intersecting pixels.

344 4 Results

345 4.1 Spatial and temporal distribution of snow depth observations across the Southern Andes

346 SD observations are available from 81 stations distributed across 26 river basins between 21° S and 54° S, for the period
347 2010–2024. The Mediterranean Andes contain the largest number of stations (37 stations in 7 basins), followed by the Wet
348 Andes (28 stations in 11 basins) and the Arid Andes (16 stations in 8 basins). The number of stations with available SD
349 observations increased markedly over the study period. From 2015 onwards, the number of stations reporting SD data
350 increased substantially, with a relatively steady growth until 2022, followed by relative stabilization through 2024. Over the
351 full study period, the DGA contributed the largest number of stations (52 stations), followed by the UdeChile (12 stations),
352 IANIGLA (eight stations), CEAZA (seven stations), and the CIEP (two stations). These institutions exhibit distinct spatial
353 coverage patterns. The DGA and IANIGLA provide the broadest geographical coverage across Chile and Argentina,
354 respectively. In contrast, CEAZA is concentrated in the Elqui, Limarí, and Choapa River Basins within the Arid Andes; the
355 UdeChile contributed stations mainly in the Mapocho and Itata River Basins across the Mediterranean and Wet Andes; and
356 the CIEP is focused on the Aysén and Baker River Basins in the Wet Andes.

357 Figure 3 presents the years with SD observations across the 81 stations of our final quality-controlled snow depth dataset,
358 considering records from May to October over the period 2010–2024. Prior to 2015, SD observations were sparse and almost
359 exclusively provided by the DGA, with only eight stations reporting data, predominantly located in the Mediterranean Andes
360 (five stations). Between 2015 and 2022, the network expanded considerably from 14 to 52 stations with SD observations.
361 This increase is still concentrated in the Mediterranean Andes (24 stations), largely driven by DGA, but also reflects the
362 incorporation of additional data sources, including UdeChile (eight stations) and IANIGLA (three stations). In parallel, the
363 Arid Andes showed a notable increase in coverage, from zero stations prior to 2015 to 12 stations, mainly operated by DGA
364 (six stations) and CEAZA (five stations). In contrast, the Wet Andes remain comparatively underrepresented throughout this
365 period, with the stations reporting SD observations increasing from seven in 2015 to 16 in 2022, despite their substantially
366 larger spatial extent relative to the other Andean zones.

367 During 2023–2024, the SD observational network reached its maximum spatial extent, with up to 57 stations reporting SD
368 observations in 2024 across the Andes. This expansion is primarily driven by a substantial increase in stations operated by
369 DGA in both the Arid and Wet Andes. In 2024, 25 stations report SD observations in the Mediterranean Andes, of which 22
370 are operated by DGA, while two stations correspond to IANIGLA and one to CEAZA. In the Arid Andes, 13 stations report

SD observations, of which nine are operated by DGA. In the Wet Andes, 19 stations report SD data, with 16 operated by DGA. By 2024, the DGA played a dominant role in the recent expansion of the SD monitoring network, including the incorporation of SD observations at the northernmost and southernmost limits of the Southern Andes.

In addition to the expansion of the SD observational network in 2024, further observational potential arises from stations recently installed or upgraded during 2025 and 2026, as well as from stations currently affected by measurement inconsistencies that are expected to be resolved in the near future (Table A2). [This indicates that the availability of SD observations is likely to continue increasing beyond the period analysed here.](#) Conversely, a subset of stations corresponds to short-term deployments associated with specific research projects. These stations typically provide observations over limited time windows and contribute to characterizing SD conditions during particular years only. As such, they fall outside the core long-term monitoring network, which is primarily sustained by the DGA and IANIGLA.

In terms of SD data availability, Figure 3 also shows the percentage of days with data from 0 to 100 % in the period May to October. Across the 81 stations of our final SD dataset, the year-by-year analysis of data availability reveals marked differences among the Arid, Mediterranean, and Wet Andes in terms of both data completeness and temporal continuity. In the Arid Andes, only six stations achieve high data availability, with SD observations covering 80–100 % of days per year for at least five years. Among them, Quebrada Larga (DGA) and La Laguna (CEAZA) exhibit the highest data availability, followed by Capayán (IANIGLA). The number of stations achieving 80–100 % annual data availability increases markedly to 17 in the Mediterranean Andes. The highest data availability corresponds to DGA stations, with continuous 11 year records at Laguna Negra and Olivares Gamma, followed by Portillo station. These stations also exhibit the longest temporal coverage, in addition to the Laguna Los Cristales and Termas del Flaco stations. Further south, in the Wet Andes, only six stations achieve 80–100 % annual data availability for at least five years. Among them, Lo Aguirre (DGA) stands out as the station with the longest record in the Southern Andes, beginning in 2010, followed by Nevado de Longaví (DGA) in 2011 and Aonikenk (IANIGLA) in 2014. Notably, continuous records of at least nine years are observed at Volcán Chillán and Alto Mallines stations.




Figure 3. Availability of quality-controlled daily snow depth observations for the 81 selected stations in Chile and Argentina (21–54° S) from 2010 to 2024. Stations are ordered latitudinally from north to south. Cell colors represent the annual availability of daily observations during the May–October snow season, classified from no data to high data availability. Colored circles beside each station indicate the institutional network from which the observations were obtained.

394 Figure 4 presents the daily SD observations in selected stations along the Southern Andes for the 2010–2024 period. SD
395 observations show a consistent increase from the arid to the wet zones of Chile, in agreement with the corresponding
396 gradient in total precipitation (Sarricolea et al., 2017). In the Arid Andes, north of 27° S, the high-elevation stations of Ojos
397 del Salado and Cerro Chajnantor (both above 5,600 m a.s.l.) exhibit limited seasonal snow depth, with most accumulation
398 occurring during the austral summer. In contrast, Tapado station (30° S, 4,300 m a.s.l.), located further south, shows larger
399 snow accumulation predominantly during the winter season. This contrast highlights a marked difference in the seasonal
400 timing of snow accumulation within the same Andean zone. In the Mediterranean Andes, stations located in the Aconcagua
401 and Maipo River Basins exceed 200 cm of SD, with Las Melosas station (33° S, 3,300 m a.s.l.) reaching values above 300
402 cm. Despite the high precipitation that occurs in the Wet Andes (Sarricolea et al., 2017), the possible predominance of
403 rainfall over snowfall, combined with warmer mountain temperatures compared to those observed further north during the
404 winter months, limits snow accumulation. As a result, SD values during recent years are close to 200 cm at the Volcán
405 Chillán station (37° S, 2,000 m a.s.l.), and reach around 150 cm east of the Southern Patagonian Icefield at the Aonikenk
406 station (49° S, 1,200 m a.s.l.).

407 A clear west–east gradient is observed across the Arid and Mediterranean Andes, with higher SD values on the western side
408 compared to the eastern side. This pattern is consistent with the stronger influence of Pacific moisture on the western slopes.
409 (Viale and Nuñez, 2011). In the Arid and Mediterranean Andes, at latitudes where both Chilean and Argentine SD stations
410 are available, stations located on the western side of the Andes show higher SD values than those on the eastern side. For
411 example, stations in the Elqui River Basin present SD values exceeding 50 cm between 2018 and 2021, whereas stations in
412 the Jáchal River Basin rarely exceed this threshold, even for stations located above 4,000 m a.s.l. (Tapado TPF and
413 Capayán). A similar pattern is observed in the Aconcagua and Maipo river basins compared to the Mendoza River Basin.
414 However, this contrast in snow accumulation is not evident in the Wet Andes, where San Rafael station located on the
415 western side of the Andes and Cordón Divisadero and Aonikenk stations located on the eastern side show similar SD values
416 during 2022–2024, at latitudes where the occurrence of rainfall strongly influences snow accumulation.

417 The megadrought signal in the Chilean Central Andes between 2010 and 2021 (Garreaud et al., 2025) is evident in the SD
418 observations, including a marked recovery of SD since 2023. Particularly, extremely dry conditions in 2019 and 2021 are
419 evident, with SD stations located in the Aconcagua, Maipo, and Mendoza river basins above 3,000 m a.s.l. showing reduced
420 SD values. This signal is also identifiable in stations within the Maule and Itata river basins.




Figure 4. Daily snow depth time series for selected stations across the Southern Andes of Chile (CL) and Argentina (AR), spanning 21° S–54° S during 2010–2024. Station locations and institutional networks are shown on the map. Time series illustrate seasonal and interannual SD variability and periods of missing data. Note that the vertical scale differs among panels. Basemap: Esri World Imagery © Esri, Maxar, Earthstar Geographics, and the GIS User Community.

4.2 Quality assessment of snow depth observations: comparison of raw and quality-controlled datasets

This section evaluates the quality of daily SD observations in both the raw and quality-controlled datasets over the 2010–2024 period. Table 2 summarizes the number of days with available SD observations and the relative data availability, defined as the proportion of days with data between the first and last available records, for both raw and quality-controlled

425 datasets across 26 river basins spanning from 21° S to 54° S. Details by station are shown in Table A4. Median values are
426 reported throughout this section because the distribution of data availability across stations is strongly skewed by a few
427 stations with exceptionally long records, making the median a more representative measure of the typical station than the
428 mean.

429 Overall, a strong positive correspondence is observed between raw and quality-controlled effective days with data ($r = 0.9$),
430 indicating that the quality-control procedure preserves the relative ranking of stations in terms of data availability. No clear
431 latitudinal dependence is identified in the reduction of data availability after cleaning. However, substantial basin-scale
432 differences emerge. The largest median reductions in data availability ($> 50\%$) are concentrated in basins of the Wet Andes,
433 including Valdivia (73%), Biobío (63%), Baker (58%), and the Costeras e Islas R. Aisén R. Baker River Basins (55%). These
434 basins also exhibit relatively high initial data availability, suggesting that the quality-control procedure removes observations
435 identified as inconsistent or exhibiting unusually high variability based on statistical and manual criteria.

436 The Vn. Mocho Choshuenco station largely explains the substantial number of SD observations filtered after applying the
437 quality-control procedure in the Valdivia River Basin, as SD observations for the 2019–2022 period exhibit high dispersion
438 that prevents both automated procedures and manual inspection from clearly identifying the seasonal snow accumulation
439 cycle. In particular, the absence of clearly identifiable snow-free periods prevents reliable determination of the ground
440 reference level the detection of low SD variability, leading to a large proportion of data being classified as unreliable and
441 subsequently removed. Similar challenges were observed in the Biobío River Basin, particularly at Chenquenco and Liucura.
442 At Chenquenco, several months in 2023 were discarded because SD remained below 2 cm and could not be reliably
443 distinguished from sensor noise or surface roughness; consequently, only observations from 2024 were retained. At Liucura,
444 only the 2022 record was preserved because the 2023–2024 period showed irregular autumn–winter accumulation,
445 anomalously high summer SD values in 2024, and no stable ground reference level. A comparable situation is found in the
446 Baker River Basin and Costeras e Islas R. Aisén R. Baker, where the large reduction in SD observations is mainly explained
447 by two out of three stations. At Steffen station, although data are available for 2020–2024, the very limited variability (< 5
448 cm) prevents the identification of a clear seasonal cycle, with only a potential accumulation signal in 2022 reaching
449 approximately 250 cm, whereas at G. San Rafael station, despite data availability from 2015 to 2024, the 2015–2019 period
450 was excluded because accumulation and ground reference level months could not be distinguished, as illustrated by years
451 such as 2016 and 2019, when higher SD values occur during spring (SON) and summer (DJF), which is inconsistent with the
452 expected seasonal cycle in this zone.

453 Intermediate reductions of days with data after cleaning (20–50 %) are observed in several basins across all climatic zones,
454 notably Rapel River Basin (48 %) in the Mediterranean Andes and Limarí River Basin (48 %) in the Arid Andes, indicating
455 that data loss is not exclusively controlled by latitude but is instead strongly influenced by station-specific issues. In the
456 Rapel River Basin, the large proportion of SD observations failing the quality-control procedure is primarily associated with

three stations: Laguna El Yeso, Laguna Los Cristales, and G. Universidad. Different sources of inconsistency are identified. At Laguna El Yeso, data from 2013 and 2015 were excluded due to the absence of snow accumulation during expected winter months, the lack of a consistent ground reference level, and highly incomplete records limited to isolated spring–summer periods between 2019 and 2021. At Laguna Los Cristales, SD observations from 2021 to 2024 were removed due to the absence of variability in the SD time series. However, at the G. Universidad station data from 2017 to 2021 were excluded due to high short-term variability in SD observations over consecutive days; nevertheless, a more detailed, station-specific analysis could potentially recover a substantial number of valid observations. A comparable situation is observed in the Limarí River Basin, where data exclusion is restricted to the Cerro Vega Negra station (2015 and 2021–2023), driven by abrupt SD variations from near 0 to approximately 200 cm within 1–2 days, likely reflecting inverted or mis-scaled observations (e.g., meters instead of centimeters), which, despite being partially captured by the quality-control procedure, can still be misinterpreted as valid snow accumulation periods and are therefore removed.

In contrast, minimal reductions in the number of days with data after the quality-control procedure (< 1%) are observed in basins with only one station, such as Yelcho, Bueno, and Tunuyán river basins. Notably, the Elqui and Maipo river basins, despite having the highest number of stations (7 and 21 stations, respectively), also exhibit very low median reductions (< 2%). This indicates a high potential for SD analyses in these basins, supported by the combination of dense observational networks and consistently high-quality records relative to other basins across the Southern Andes.

Out of the 81 SD stations of our final quality-controlled dataset, 45 stations distributed between 21° S and 49° S show reductions in data availability below 10%, with a median of 1,236 days with valid observations, after the quality-control procedure. Within this group, the Elqui River Basin includes five stations, Glaciar Tapado en Los Corrales, Tapado (TPF), Los Corrales, La Laguna, and El Jote, while the Maipo River Basin comprises 11 stations: Glaciar Juncal Sur, Glaciar Olivares Gamma, Glaciar Olivares Alfa, AMTC10, Piuquenes7, La Parva, Farellones, AMTC12, Laguna Negra, Las Melosas, and Las Hualtatas. The relatively high data availability and long observational records at these stations provide a robust basis for characterizing temporal variations in SD within these basins.

Table 2. Effective days of snow depth observations for raw and quality-controlled datasets by river basin, considering all available records over the 2010–2024 period.							
Basin (country)	N° Stations	Raw data		Clean data		Reduction of days with data after cleaning (%)	Total basin area (km²)
		Effective days with data (median)	Relative data availability (%)	Effective days with data (median)	Relative data availability (%)		
Salar Michincha (CL)	1	1,085	96	1,062	94	2	2,674
Salar Atacama (CL)	1	733	71	622	60	15	5,307
Tres Cruces (CL)	2	592	81	537	74	9	15,618
Copiapo (CL)	1	988	100	975	99	1	18,703
Jáchal (AR)	1	2,933	100	2,892	99	1	32,318
Elqui (CL)	7	1,769	84	1,742	83	2	9,825
Limarí (CL)	3	2,940	82	1,536	47	48	11,696
Choapa (CL)	2	2,720	97	2,596	92	5	7,653

Petorca (CL)	1	2,831	97	2,638	90	7	1,988
Aconcagua (CL)	2	2,825	95	2,297	77	19	7,334
Mendoza (AR)	2	1,866	100	1,809	99	3	21,830
Maipo (CL)	21	1,261	51	1,236	50	2	15,273
Rapel (CL)	7	2,461	57	1,282	53	48	13,766
Mataquito (CL)	1	2,910	90	2,145	66	26	6,332
Tunuyán (AR)	1	2,560	89	2,529	89	1	22,195
Maule (CL)	4	3,636	83	2,736	63	25	21,052
Itata (CL)	5	1,016	100	977	96	4	11,326
Biobío (CL)	5	1,129	99	419	38	63	24,369
Valdivia (CL)	2	1,081	75	296	62	73	10,244
Bueno (CL)	1	598	95	592	94	1	15,366
Yelcho (CL)	1	248	99	247	99	0	4,084
Aysén (CL)	2	1,112	93	762	63	32	11,456
Costeras e Islas R. Aisén R. Baker (CL)	1	2,567	75	1,149	83	55	35,153
Baker (CL)	2	1,354	99	573	96	58	20,945
Santa Cruz (AR)	4	1,490	61	1,313	57	12	29,938
Islas sur Estrecho de Magallanes (CL)	1	414	100	285	99	31	27,931

480 4.3 Assessing the physical consistency of snow depth with precipitation and air temperature by station

481 Table 3 summarizes the results of the PCI metric applied to stations located in the Elqui and Maipo River Basins, which are
482 the two basins with the largest number of SD stations and the most comprehensive elevation distribution among the 26 river
483 basins analyzed in this study. Two and five SD stations met the conditions required for this validation in Elqui and Maipo
484 River Basins, respectively, including overlapping daily SD, Pr, and AT data during the accumulation period (May–October)
485 over the period 2017 to 2024, and satisfied the three criteria related to air temperature threshold, snow accumulation, and
486 precipitation events (see in Section 3.4). These strict data availability requirements substantially reduced the number of days
487 available for PCI analysis. In the Elqui River Basin, the median number of days with overlapping quality–controlled SD,
488 precipitation, and air temperature observations decreased from 360 to 19 days after applying all selection criteria. In the
489 Maipo River Basin, the corresponding median decreased from 965 to 79 days. This reduction is primarily explained as the
490 filter explicitly isolates precipitation–driven accumulation days, discarding the majority of the winter season characterized by
491 zero precipitation, stable snowpacks, or ablation periods. Complete results for both the raw and quality–controlled datasets
492 are provided in Tables A5 and A6.

493 Across the analyzed stations, the quality–control procedure improves the stability of physical consistency metrics. In the
494 Elqui River Basin, the PCI in the quality–controlled dataset ranges between 90.0 % and 92.9 %, compared to 90.5 % and
495 93.9 % in the raw dataset, indicating only minor changes after cleaning. In contrast, in the Maipo River Basin, the PCI range
496 narrows substantially after quality–control, converging to 94.4%–94.9 % from a wider range of 87.3 %–95.1 % in the raw
497 dataset. This reduction in dispersion reflects a more consistent representation of physically plausible conditions following the
498 application of the quality–control procedure, while no clear relationship is observed between PCI and station elevation in the

499 river basins. While the clean PCI is relatively stable, relying on it in isolation can be misleading. Because PCI is a ratio,
500 similar values can arise from datasets with substantially different numbers of retained accumulation events. An example of
501 this is observed at the Cerro Olivares station (Elqui River Basin), where despite a high Noise Ratio ($NR = 2.1$) indicating
502 that more than half of the raw SD days were spurious values, the raw PCI (90.5 %) is similar to the clean PCI (90.0 %).
503 Complementary, a similar behavior is observed at the Termas del Plomo station (Maipo River Basin), where a higher ΔE
504 value (11) indicates a greater frequency of physically inconsistent accumulation signals, such as rain-to-snow transitions,
505 which are effectively filtered by the algorithm. Despite this substantial removal of inconsistent events, the clean PCI (94.9
506 %) remains close to the raw PCI (95.1 %), reinforcing that similar PCI values can mask important improvements in the
507 physical consistency of the dataset.

508 The quality-control procedure evaluation in these stations can be categorized into three groups. The first group comprises
509 stations with higher absolute error removal ($\Delta E \geq 2$), exhibiting an increase in PCI from 87.3% in the raw data to 94.9% after
510 quality-control procedure. This group includes the Termas del Plomo, Glaciar Juncal Sur, and Glaciar Olivares Alfa stations.
511 The improvement in PCI is associated with the removal of a substantial number of physically inconsistent days (NR ranging
512 from 2 to 11). Notably, NR does not show a direct relationship with the PCI. The second group comprises stations with low
513 error correction ($\Delta E \leq 1$) and varying proportions of discarded noise (NR ranging from 1.0 to 2.1), which exhibit a slight
514 decrease from raw PCI to quality-control PCI. This behavior is observed at Las Melosas, Glaciar Tapado Corrales and Cerro
515 Olivares stations in the Elqui River Basin. This marginal decline does not indicate a deterioration in data quality. Rather, the
516 quality-control procedure removes spurious noise, thereby reducing the total sample size (as reflected by NR), which
517 increases the relative influence of the few remaining natural inconsistencies on the final PCI percentage. The third group
518 exhibits a distinct behavior, represented by the Glaciar Olivares Gamma station, which stands out as a unique case of signal
519 recovery. Unlike the previous groups, this pattern is explained by its metrics. Although no errors were detected by the
520 automated filters ($\Delta E = 0$), manual correction of the record (executed in Step 2) increased the PCI from 89.8% to 94.4%.
521 This improvement did not result from the automated noise filters, but from the correction of specific issues in the raw data
522 associated with height-unit continuity errors caused by sensor handling or configuration issues. These errors could not be
523 corrected automatically and therefore required manual intervention. This process nearly doubled the total number of valid
524 accumulation events ($NR = 0.6$), diluting the few remaining inconsistencies and increasing the final PCI. The increase in the
525 number of valid accumulation events reduced the proportional contribution of the remaining inconsistent events, thereby
526 increasing PCI. Although this station behaves as an outlier, it includes more than 100 analyzed days over six years, enabling
527 this outcome, in contrast to Cerro Olivares, which comprises only 10 analyzed days within a single year of observations.

Table 3. Assessment of snow depth, precipitation, and air temperature data before and after the quality-control procedure in the Elqui and Maipo river basins for the accumulation period May–October, 2017–2024.							
Station name	Basin	Elevation (m a.s.l.)	N° of overlapping days in QC dataset	ΔE	NR	Raw PCI (%)	Clean PCI (%)

			With negative and positive Δ SD and Pr > 0 mm	With SD > 1 cm	With Pr > 1 mm	Start–End period				
Cerro Olivares	Elqui	3,566	182	11	10	2024–2024	1	2.1	90.5	90.0
Glaciar Tapado Corrales		4,065	538	29	28	2020–2023	0	1.2	93.9	92.9
Termas Del Plomo	Maipo	3,027	710	77	59	2017–2022	11	1.9	87.3	94.9
Las Melosas		3,317	543	91	79	2022–2024	1	1.0	95.1	94.9
Glaciar Olivares Gamma		3,628	1050	156	107	2019–2024	0	0.6	89.8	94.4
Glaciar Juncal Sur		4,035	1041	113	110	2019–2024	4	1.4	93.4	94.6
Glaciar Olivares Alfa		4,230	965	78	72	2019–2024	2	1.4	93.9	94.4

528 **4.4 Deriving snow depth–elevation patterns from *in situ* measurements**

529 We selected three river basins representing the Arid (Elqui), Mediterranean (Maipo), and Wet Andes (Maule–Itata) because
530 they provide relatively consistent SD observations distributed across broad elevation gradients over the 2010–2024 period,
531 despite the heterogeneous spatial coverage and temporal gaps that characterize the complete dataset. Despite this selection,
532 none of these basins contain continuous observations for all years and elevations, which limits the analysis of SD variability
533 on a year–by–year basis. To address this limitation, we adopt a representative–year approach based on three precipitation
534 conditions, under dry, normal, and wet years (Table A3).

535 Figure 5 presents the cumulative distribution function (CDF) of daily SD observations at each station during the seasonal
536 snow accumulation period (1 May–31 October) for the representative dry, normal, and wet years. For each station and
537 precipitation–year type, the median SD (CDF = 0.5) was used as a representative measure of snow accumulation, providing a
538 consistent basis for comparing SD across elevations and climatic conditions. Across the three basins, a coherent large–scale
539 pattern emerges: SD increases with elevation and from dry to wet years, reflecting the combined control of orographic
540 precipitation and zonal climatic gradients. In dry years, SD distributions are strongly compressed toward low values,
541 considering elevation from 2,000 to 4,500 m a.s.l., with several stations exhibiting less than 50 cm of median SD. In contrast,
542 wet years show a pronounced increase of SD with higher variability of SD over CDF 0.5. This transition is especially
543 marked in the Wet Andes basin (Maule–Itata), where median SD exceeds 200 cm at all stations during wet conditions.
544 Despite this general behavior, the CDF–based analysis indicates that the relationship between SD and elevation is not strictly
545 positive within basins: (i) a positive SD–elevation relationship in the Arid Andes, (ii) a non–positive SD–elevation pattern in
546 the Mediterranean Andes, characterized by mid–elevation SD maxima, and (iii) a consistently high and positive
547 SD–elevation relationship in the Wet Andes.

548 In the Elqui River Basin (Arid Andes), SD exhibits a strong dependence on elevation and precipitation–year type. During the
549 dry year (2021), [median daily SD values](#) are close to zero at most stations below 3,600 m a.s.l., with only the highest
550 elevation station, Tapado TPF (4,306 m a.s.l.), reaching 16 cm. Under normal conditions (year 2018), SD increases
551 substantially at high elevations (38 cm at Tapado TPF station), while remaining limited at mid elevations (4 cm at Llano de
552 Liebres station). The wet year (2022) reveals a marked increase of SD across all elevations, with station coverage broadly
553 comparable to that available for the dry year, with a clear altitudinal gradient: median SD reaches 132 cm at 4,306 m a.s.l.
554 and remains relatively high (e.g., 55 cm at 3,660 m a.s.l. at El Jote station), although lower elevation stations still exhibit
555 reduced accumulation (< 5 cm at 3,200 m a.s.l. at La Laguna station). Overall, the observations suggest increasing SD
556 toward the highest sampled elevations, although the relationship is not monotonic in all years.

557 In the Maipo River Basin (Mediterranean Andes), SD exhibits a complex and spatially heterogeneous distribution along the
558 elevation gradient. During the dry year (2021), median SD remains close to zero at the highest elevations (e.g., Tupungatito
559 Bajo, 4,425 m a.s.l.), while intermediate elevations (3,000–3,400 m a.s.l.) show substantially higher values (e.g. 53 cm at Las
560 Melosas and 27 cm at Termas del Plomo), indicating a non-monotonic SD–elevation relationship. In the normal year (2018),
561 SD increases markedly across a broad elevation range, with several stations between 3,300 and 3,600 m a.s.l. exceeding 100
562 cm (e.g., Piuquenes14: 102 cm; Las Melosas: 129 cm), whereas lower elevations exhibit reduced values and the highest
563 elevations lack observations. During the wet year (2024), median SD reaches its maximum, particularly at mid elevations
564 (e.g. 260 cm at Las Melosas and 130 cm at Laguna Negra), while remaining comparatively low at elevations above 4,000 m
565 a.s.l. (close to 5 cm at G. Tupungatito Bajo and G. Olivares Alfa). [The persistence of this mid–elevation maximum suggests](#)
566 [the combined influence of several processes. First, mean winter precipitation in the subtropical Andes has been shown to](#)
567 [peak on the windward slopes below the crest of the Andes, rather than at the highest elevations, as a result of orographic](#)
568 [enhancement and upstream flow blocking \(Viale and Nuñez, 2011\). Second, wind–driven snow redistribution and enhanced](#)
569 [sublimation at higher elevations, where drier atmospheric conditions prevail, may further reduce snow accumulation, thereby](#)
570 [preventing the development of a simple positive SD–elevation relationship. Although winter–focused studies addressing](#)
571 [these processes remain scarce in this region, Ayala et al. \(2017\) highlighted the significant role of sublimation on](#)
572 [wind–exposed surfaces of the Juncal Norte Glacier near 4,500 m a.s.l.](#)

573 In the Maule–Itata River Basin (Wet Andes), SD exhibits a more consistent and coherent increase with both elevation and
574 climatic wetness. Even during the dry year (2019), [median daily SD values](#) remain relatively high compared to northern
575 basins (e.g., 77 cm at 1,989 m a.s.l. at Lo Aguirre and 46 cm at 1,977 m a.s.l. at Nevado Longaví). In the normal year (2017),
576 SD increases substantially, reaching 350 cm at 2,439 m a.s.l. (Los Cóndores), with all stations exceeding 50 cm. During the
577 wet year (2024), SD shows a strong amplification across all elevations, with median values exceeding 200 cm at all stations
578 and reaching up to 380 cm at Los Cóndores. Unlike the Elqui and Maipo river basins, the SD–elevation relationship remains
579 largely positive and continuous, reflecting more humid atmospheric conditions and a reduced influence of sublimation at

580 high elevations, which in turn favors more consistent snow accumulation across the elevation gradient in the Wet Andes
581 (Schaefer et al., 2020).




Figure 5. Cumulative distribution functions of snow depth by elevation for specific years with dry, normal, and wet precipitation conditions across three river basins corresponding to each Andean zone. The dry (2021) and normal (2018) years are shared between the Arid and Mediterranean Andes, whereas wet years correspond to 2022 in the Arid Andes and 2024 in the Mediterranean Andes; in contrast, the Wet Andes are represented by a different set of years, with dry (2019), normal (2017), and wet (2024) precipitation conditions. The analysis is based on daily snow depth observations from 1 May to 31 October in the Elqui, Maipo, and Maule–Itata river basins.

582 In addition, an additional assessment of SD observations was conducted by examining the SD–elevation relationship
583 together with the hypsometry of the three selected river basins.

584 Figure 6A shows that observed SD generally increases from the Arid to the Wet Andes, consistent with the regional
585 precipitation gradient (Sarricolea et al., 2017). Furthermore, similar SD–elevation patterns, analysed through CDFs, were
586 observed in the three river basins. Although the available stations do not fully represent the spatial variability within each
587 basin, they constitute the most comprehensive in situ SD observations currently available and provide observational

588 constraints on SD–elevation relationships. In the Elqui River Basin (29° S), the available observations indicate a positive
589 relationship between SD and elevation during the dry year (2021), based on five stations spanning elevations from 3,209 to
590 4,306 m a.s.l. A similar positive relationship is observed during the normal year (2018), although this pattern is poorly
591 constrained because only two stations are available, spanning elevations up to 3,500 m a.s.l. However, during the wet year
592 (2022), based on six stations spanning elevations from 3,209 to 4,306 m a.s.l., this pattern is interrupted: SD increases up to
593 approximately 3,600 m a.s.l., then declines toward 4,000 m a.s.l., before increasing again up to the highest elevation with
594 available SD observations (4,300 m a.s.l.). These features are also reflected in the 25th and 75th percentile SD values. In the
595 Maipo River Basin (33° S), the available observations indicate that the SD–elevation relationship remains positive up to
596 approximately 2,800–3,300 m a.s.l. across all precipitation years, based on at least 10 stations spanning elevations from
597 2,020 to 4,425 m a.s.l. The available observations further suggest that above 3,300 m a.s.l., SD begins to decline. In contrast,
598 the available SD observations in the Maule–Itata River Basin (37° S), based on at least three stations during the dry year and
599 six stations during the wet year, indicate a consistently positive SD–elevation relationship.

600 Although such SD–elevation patterns have not been previously documented in Chile at the basin scale over a wide elevation
601 range, existing studies restricted to small microcatchments with short observational periods provide only partial insights.
602 Mendoza et al. (2020) examined microcatchments in the Arid and Mediterranean Andes within a relatively narrow elevation
603 range, finding that SD increased between 3,600 and 3,900 m a.s.l. in 2018, followed by a decrease at higher elevations, and
604 that SD values declined on slopes steeper than 35°. Similarly, Shaw et al. (2020a,b) showed in a microcatchment of the
605 Mediterranean Andes that SWE increased from approximately 3,000 to 3,800 m a.s.l., then declined up to 5,400 m a.s.l., in
606 the period 2017–2018.

607 The elevational distribution of SD stations under these precipitation regimes supports the observed SD patterns across the
608 three river basins. Figure 6B presents the distribution of SD stations by elevation alongside the hypsometric curve for each
609 basin. The lower bound of the hypsometric curve represents the minimum elevation of July snow persistence over the
610 2000–2024 period, while the elevation corresponding to the 25th percentile of snow persistence is indicated by a thick line.

611 Despite the generally well–distributed elevation range of stations, the Elqui River Basin includes five SD stations that do not
612 provide observations during the analyzed normal precipitation year, highlighting the importance of CEAZA stations in
613 characterizing the SD–elevation relationship in this basin. A comparable spatial distribution of SD stations across the three
614 precipitation regimes is observed in the Maipo River Basin; however, of the 18 stations, nine lack observations for the
615 selected normal year, particularly at the lowest and highest elevations. Although the SD network is largely composed of
616 DGA stations, 13 spanning a broad elevation range (2,020–4,230 m a.s.l.), stations operated by UdeChile provide important
617 complementary coverage in the Mapocho catchment. In contrast, the Maule–Itata River Basin comprises only DGA stations,
618 with three stations lacking data during dry precipitation years.

As a result, when considering only these three specific precipitation conditions, just six stations capture SD variability across the Maule–Itata River Basin, which encompasses a snow–covered area of 30,690 km² as defined by the snow line elevation estimated here. This contrasts with the smaller snow–covered areas of the Elqui (6,460 km²) and Maipo (7,730 km²) River Basins, where a denser SD station network is available.




Figure 6. Measured relationships between snow depth and elevation for three selected precipitation years, classified as dry (D), normal (N), and wet (W), across three river basins representing the Arid (Elqui, 29° S), Mediterranean (Maipo, 33° S), and Wet (Maule–Itata, 37° S) Andes in Chile. Only SD observations from stations with at least 70% of daily observations per month during the accumulation period (1 May to 31 October) are included. In each upper panel (A), bars represent the 25th, 50th, and 75th percentile SD values, derived from all available observations for the corresponding selected precipitation year. The lower panel (B) compares station elevations with the basin hypsometric curves. The thin line indicates the minimum elevation of July snow persistence over the 2000–2024 period, while the thick line marks the elevation at which snow persistence reaches the 25th percentile (SP25). Circles denote the elevation of SD stations and are labeled according to the selected precipitation year (D, N, or W). Stations without labels correspond to sites with SD observations available for all three selected precipitation years.

623 5 Data availability

624 The quality-controlled dataset is openly available through Zenodo at <https://zenodo.org/records/21577171> (Medina and
625 Caro, 2026). The repository provides daily snow depth observations from 69 automatic stations across the Southern Andes
626 for 2010–2024, together with station metadata and a dataset overview figure. An interactive snow depth exploration tool is
627 available at <https://javiermedinamen-art.github.io/hidromet>, developed by the Laboratorio de Teledetección Ambiental. A
628 concise file manifest describing all distributed files and their contents is provided in Table A1 and Table A4. The same
629 information is available in the Zenodo repository in the file Station metadata for snow depth observations.

630 Input data were obtained from public repositories and direct institutional contributions, including the DGA, CEAZA (or
631 CEAZAMET), IANIGLA, the University of Chile, and CIEP. DGA data are distributed through the Chilean open-data
632 framework, which allows data reuse and redistribution with proper attribution. CEAZA data usage policies explicitly allow
633 their use in scientific publications with acknowledgment of the original source. Datasets from IANIGLA, the University of
634 Chile, and CIEP were provided directly by these institutions for scientific research purposes, and co-authors actively
635 affiliated with these institutions facilitated access to the corresponding datasets. The original observations were reprocessed
636 and integrated into the quality-controlled dataset distributed with this study, while maintaining attribution to the original data
637 providers.

638 6 Code availability

639 The full data-processing and quality-control workflow comprises three sequential processing steps, followed by an
640 independent cross-variable validation, as described in Section 3: (1) compilation of multi-institutional snow depth records;
641 (2) automated and manual processing, including zero-level correction (Step 2.1), spike removal (Step 2.2), and
642 physical-range filtering (Step 2.3), complemented by station-level expert screening; and (3) harmonization of overlapping
643 series. The resulting dataset was subsequently evaluated through cross-variable validation using the Physical Consistency
644 Index (PCI).

645 An executable demonstration of Step 2.1 (zero-level/ground-reference correction; function `detect_flat_offsets`) is available
646 on GitHub at https://github.com/javiermedinamen-art/hidromet/tree/main/sd_cleaning_sample. The sample is applied to the
647 Valle Olivares station (05706003; 33° S) and includes the Python module, a Jupyter notebook, an example input CSV (date
648 and snow depth in centimetres), dependency specifications, and a basin-level parameter table for Steps 2.1–2.3
649 (`QC_parameters_by_river_basin`). Default parameters in the sample correspond to the Maipo River Basin settings used for
650 Valle Olivares (window = 5 days; var_thr = 0.5 cm; tol = 0.3 cm; drift_thr = 0.3 cm; snow_thr = 5 cm).

Steps 2.2, 2.3, and 3 are fully specified in Section 3, including the algorithms, decision rules, and thresholds used in the processing workflow. The complete source code implementing these steps is not publicly available but can be obtained from the authors upon request. Expected input and output formats for the automated cleaning stage are daily tabular files with one column per station (raw to quality-controlled snow depth in centimetres). The open dataset provides quality-controlled daily snow depth for 69 stations; institutional restrictions prevent redistribution of the remaining series included in the 81-station analysis.

7 Conclusions

This open-access, quality-controlled dataset provides the largest and most complete collection of continuous daily snow depth observations currently available for the Southern Andes, comprising 81 stations across 21°–54° S over 2010–2024. Of the 118 candidate snow depth stations initially compiled across Chile and Argentina, 81 were retained after quality control. Records from 69 stations are openly distributed through Zenodo, while data-use restrictions prevent redistribution of the remaining 12 records. The quality-control procedure improved data reliability, increasing the newly proposed Physical Consistency Index from 90.5 % in the raw dataset to 92.9 % in the quality-controlled dataset for stations in the Elqui River Basin, and from 87.3 % to 94.9 % for stations in the Maipo River Basin. These results support the effectiveness of the quality-control procedure at the stations where independent cross-variable was possible, while resulting in an average 21% reduction in data availability across the 26 river basins analysed. The main conclusions are as follows.

- No clear latitudinal pattern is evident in station data quality across the 26 river basins. Nevertheless, several of the largest reductions in data availability after quality control occurred in Wet Andes basins, particularly Valdivia, Biobío, and Baker (>50 %). In contrast, in the Arid and Mediterranean Andes, the Elqui and Maipo River Basins containing 7 and 21 retained stations, respectively, showed only minor reductions after quality control. The combination of relatively large station numbers and limited data loss makes these basins particularly suitable for basin-scale SD analyses.
- Only three river basins provide sufficient elevational coverage to examine basin-scale quality-controlled SD-elevation relationship: Elqui, Maipo, and Maule-Itata. Nevertheless, the available stations do not fully represent the spatial variability of snow conditions within these basins. The available observations indicate non-linear SD-elevation relationships in two of these basins, with maximum observed SD at approximately 3,300 m a.s.l. in the Maipo River Basin (Mediterranean Andes) and 4,300 m a.s.l. in the Elqui River Basin (Arid Andes). Elqui River Basin shows increasing SD toward higher elevations but with non-monotonic behavior under wet conditions. In contrast, SD observations in the Maule-Itata River Basin (Wet Andes) indicate a predominantly positive SD-elevation relationship.

681 - The spatial distribution of SD monitoring stations does not reflect the relative spatial extent of snow-covered areas
682 in the study domain. The Maule–Itata River Basin has the largest snow-covered area (30,690 km²) yet a
683 comparatively lower station density than the smaller Elqui (6,460 km²) and Maipo (7,730 km²) River Basins. In
684 these basins, the DGA network is strongly complemented by stations operated by CEAZA and Universidad de
685 Chile, which provide important complementary elevational and spatial coverage for improving the characterization
686 of SD–elevation patterns.

687 This research underscores the need for sustained efforts to improve the availability and spatial representativeness of *in situ*
688 snow depth observations through informed monitoring strategies and policy decisions.

689 Author contributions

690 AC conceived the study and defined its scope. AC and JM compiled the station dataset, designed the quality-control
691 procedure, performed the analyses, and prepared the tables and figures. JM developed the quality-control algorithm. AC
692 drafted the first version of the manuscript. All co-authors reviewed and revised the manuscript.

693 Competing interests

694 The contact author has declared that none of the authors has any competing interests.

695 Acknowledgements

696 The authors gratefully acknowledge DGA, CEAZA, the Department of Civil Engineering at the University of Chile, Marlene
697 Huerta, CIEP, CECS, and IANIGLA for providing the snow depth data used in this study. We also thank Valentina Contreras
698 and Antonia Flores for their contributions to data compilation. Finally, we thank the reviewers for their constructive
699 comments, which helped improve the quality of this article.

700 Financial support

701 This research has been supported by the following grants:
702 ANID/FONDEF/N.IT24I0092; ANID/FONDECYT/POSTDOCTORADO/N.3261164.

703 References

704 Aguayo R., León-Muñoz J., Aguayo M., Baez-Villanueva O., Fernandez A., Zambrano-Bigiarini M., Fernández, A., and
705 Jacques-Coper, M.: PatagoniaMet: A multi-source hydrometeorological dataset for Western Patagonia. Sci. Data. 11, 6,
706 10.1038/s41597-023-02828-2, 2024.

708 Alvarez-Garretón, C., Boisier, J. P., Garreaud, R., Seibert, J., and Vis, M.: Progressive water deficits during multiyear
709 droughts in basins with long hydrological memory in Chile, *Hydrol. Earth Syst. Sci.*, 25, 429–446,
710 doi:10.5194/hess-25-429-2021, 2021.

711

712 Alvial Vásquez F-J, Abarca-del-Río R and Ávila AI.: High-Resolution Precipitation Gridded Dataset on the South-Central
713 Zone (34° S–41° S) of Chile, *Front. Earth Sci.* 8:519975. doi: 10.3389/feart.2020.519975, 2020.

714

715 Ayala, Á., Pellicciotti, F., Peleg, N., and Burlando, P.: Melt and surface sublimation across a glacier in a dry environment:
716 distributed energy-balance modelling of Juncal Norte Glacier, Chile, *J. Glaciol.*, 63, 803–822, doi:10.1017/jog.2017.46,
717 2017.

718

719 Ayala, Á., Farías-Barahona, D., Huss, M., Pellicciotti, F., McPhee, J., and Farinotti, D.: Glacier runoff variations since 1955
720 in the Maipo River basin, in the semiarid Andes of central Chile, *Cryosphere*, 14, 2005–2027, doi:10.5194/tc-14-2005-2020,
721 2020.

722

723 Ayala, Á., Muñoz-Castro, E., Farinotti, D., Farías-Barahona, D., Mendoza, P. A., MacDonell, S., McPhee, J., Vargas, X., and
724 Pellicciotti, F.: Less water from glaciers during future megadroughts in the Southern Andes, *Commun Earth Environ* 6, 860,
725 doi:10.1038/s43247-025-02845-6, 2025.

726

727 Bulovic, N., Johnson, F., Lievens, H., Shaw, T. E., McPhee, J., Gascoin, S., Demuzere, M., and McIntyre, N.: Evaluating the
728 performance of Sentinel-1 SAR derived snow depth retrievals over the extratropical Andes Cordillera, *Water Resour. Res.*,
729 61, doi:10.1029/2024WR037766, 2025.

730

731 Burger, F., Ayala, Á., Farías, D., Shaw, T. E., MacDonell, S., Brock, B., McPhee, J., and Pellicciotti, F.: Interannual
732 variability in glacier contribution to runoff from a high-elevation Andean catchment: Understanding the role of debris cover
733 in glacier hydrology, *Hydrol. Process.*, 33, 214–229, doi:10.1002/hyp.13354, 2019.

734

735 Caro, A., Condom, T., and Rabatel, A.: Climatic and morphometric explanatory variables of glacier changes in the Andes
736 (8–55° S): New insights from machine learning approaches, *Front. Earth Sci.*, 9, 713011, doi:10.3389/feart.2021.713011,
737 2021.

738

739 Caro, A., Condom, T., Rabatel, A., Champollion, N., García, N., and Saavedra, F.: Hydrological response of Andean
740 catchments to recent glacier mass loss, *Cryosphere*, 18, 2487–2507, doi:10.5194/tc-18-2487-2024, 2024.

741

742 Caro, A., Condom, T., Rabatel, A., Aguayo, R., and Champollion, N.: Future glacio–hydrological changes in the Andes: a
743 focus on near-future projections up to 2050, *Sci Rep* 15, 10991, doi:10.1038/s41598-025-88069-2, 2025
744

745 Cordero, R. R., Feron, S., Damiani, A., MacDonell, S., Carrasco, J., Pizarro, J., Karas, C., Jorquera, J., Sepulveda, E.,
746 Cabello, F., Fernandoy, F., Wang, C., Khan, A. L., and Casassa, G.: Rapid decline in extratropical Andean snow cover driven
747 by the poleward migration of the Southern Hemisphere westerlies, *Sci Rep* 14, 26365, doi:10.1038/s41598-024-78014-0,
748 2024.
749

750 Cornwell, E., Molotch, N. P., and McPhee, J.: Spatio-temporal variability of snow water equivalent in the extra-tropical
751 Andes Cordillera from distributed energy balance modeling and remotely sensed snow cover, *Hydrol. Earth Syst. Sci.*, 20,
752 411–430, doi:10.5194/hess-20-411-2016, 2016.
753

754 Cortés, G. and Margulis, S.: Impacts of El Niño and La Niña on interannual snow accumulation in the Andes: Results from a
755 high-resolution 31-year reanalysis, *Geophys. Res. Lett.*, 44, 6859–6867, doi:10.1002/2017GL073826, 2017.
756

757 Durre, I., Menne, M. J., Gleason, B. E., Houston, T. G., and Vose, R. S.: Comprehensive automated quality assurance of
758 daily surface observations, *J. Appl. Meteorol. Clim.*, 49, 1615–1633, doi:10.1175/2010JAMC2375.1, 2010.
759

760 Dussailant, A., Buytaert, W., Meier, C., and Espinoza, F.: Hydrological regime of remote catchments with extreme gradients
761 under accelerated change: the Baker basin in Patagonia, *Hydrolog. Sci. J.*, 57, 1530–1542,
762 doi:10.1080/02626667.2012.726993, 2012.
763

764 Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., Roth, L., Seal,
765 D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M., Burbank, D., Alsdorf, D.: The Shuttle Radar Topography
766 Mission. *Rev. Geophys.* 45, RG2004, doi:10.1029/2005RG000183, 2007.
767

768 Garreaud, R. D., Alvarez-Garretón, C., Barichivich, J., Boisier, J. P., Christie, D., Galleguillos, M., LeQuesne, C., McPhee,
769 J., and Zambrano-Bigiarini, M.: The 2010–2015 megadrought in central Chile: Impacts on regional hydroclimate and
770 vegetation, *Hydrol. Earth Syst. Sci.*, 21, 6307–6327, doi:10.5194/hess-21-6307-2017, 2017.
771

772 Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D.: The central Chile
773 mega drought (2010–2018): A climate dynamics perspective, *Int. J. Climatol.*, 40, 421–439, doi:10.1002/joc.6219, 2020.
774

775 Garreaud, R., Boisier, J. P., Alvarez-Garretón, C., Christie, D. A., Carrasco-Escaff, T., Vergara, I., Chávez, R. O., Aldunce,
776 P., Camus, P., Suazo-Álvarez, M., Masiokas, M., Castro, G., Muñoz, A., Zambrano-Bigiarini, M., Fuster, R., and Godoy, L.:

777 Hyperdroughts in central Chile: drivers, impacts, and projections, *Hydrol. Earth Syst. Sci.*, 29, 5347–5369,
778 doi:10.5194/hess-29-5347-2025, 2025.

779

780 Grünewald, T., Bühler, Y., and Lehning, M.: Elevation dependency of mountain snow depth, *Cryosphere*, 8, 2381–2394,
781 doi:10.5194/tc-8-2381-2014, 2014.

782

783 Huerta, M.L., Molotch N.P., McPhee J.: Snowfall interception in a deciduous *Nothofagus* forest and implications for spatial
784 snowpack distribution, *Hydrological Processes*, 33, 1818–1834, doi:10.1002/hyp.13439, 2019.

785

786 Instituto Nacional de Estadísticas (INE): Resultados Censo 2024, available at: <https://censo2024.ine.gob.cl/resultados/>
787 (last access: 10 December 2025), 2024.

788

789 Instituto Nacional de Estadísticas (INE): Gráficas nacionales – Resultados del VIII Censo Nacional Agropecuario y Forestal,
790 available at: <https://www.ine.gob.cl/censoagropecuario/resultados-finales/graficas-nacionales>
791 (last access: 10 December 2025), 2022.

792

793 Krogh, S. A., Pomeroy, J. W., and McPhee, J.: Physically based mountain hydrological modeling using reanalysis data in
794 Patagonia, *J. Hydrometeorol.*, 16, 172–193, doi:10.1175/JHM-D-13-0178.1, 2015.

795

796 MacDonell, S., Kinnard, C., Mölg, T., Nicholson, L., and Abermann, J.: Meteorological drivers of ablation processes on a
797 cold glacier in the semi-arid Andes of Chile, *Cryosphere*, 7, 1513–1526, doi:10.5194/tc-7-1513-2013, 2013.

798

799 Masiokas, M. H., Rabatel, A., Rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, A., Bown, F., Berthier, E.,
800 Dussailant, I., and MacDonell, S.: A review of the current state and recent changes of the Andean cryosphere, *Front. Earth*
801 *Sci.*, 8, 99, doi:10.3389/feart.2020.00099, 2020.

802

803 Masiokas, M. H., Cara, L., Villalba, R., Pitte, P., Luckman, B. H., Toum, E., Christie, D. A., Le Quesne, C., and Mauget, S.:
804 Streamflow variations across the Andes (18°–55°S) during the instrumental era, *Sci Rep*, 9, 17879,
805 doi:10.1038/s41598-019-53981-x, 2019.

806

807 Masiokas, M. H., Christie, D. A., Le Quesne, C., Pitte, P., Ruiz, L., Villalba, R., Luckman, B. H., Berthier, E., Nussbaumer,
808 S. U., González-Reyes, Á., McPhee, J., and Barcaza, G.: Reconstructing the annual mass balance of the Echaurren Norte
809 glacier (Central Andes, 33.5° S) using local and regional hydroclimatic data, *The Cryosphere*, 10, 927–940,
810 doi:10.5194/tc-10-927-2016, 2016.

811

812 Medina, J., and Caro, A.: The Southern Andes Daily Snow Depth Dataset (2010–2024): quality-controlled dataset from
813 Chile and Argentina, Zenodo [data set], <https://zenodo.org/records/21577171>, 2026.

814

815 Mendoza, P. A., Shaw, T. E., McPhee, J., Musselman, K. N., Revuelto, J., and MacDonell, S.: Spatial distribution and scaling
816 properties of lidar-derived snow depth in the extratropical Andes. *Water Resources Research*, 56, e2020WR028480,
817 doi:10.1029/2020WR028480, 2020.

818

819 Ntokas, K. F. F., Odry, J., Boucher, M. A., and Garnaud, C.: Investigating ANN architectures and training to estimate snow
820 water equivalent from snow depth, *Hydrol. Earth Syst. Sci.*, 25, 3017–3040, doi:10.5194/hess-25-3017-2021, 2021.

821

822 Pandas development team: pandas-dev/pandas: Pandas (version 2.3.0), Zenodo, doi:10.5281/zenodo.15597513, 2025.

823

824 Perkins, R. M. and Jones, J. A.: Climate variability, snow, and physiographic controls on storm hydrographs in small forested
825 basins, western Cascades, Oregon, *Hydrol. Process.*, 22, 4949–4964, doi:10.1002/hyp.7117, 2008.

826

827 Rivera, D., Godoy-Faúndez, A., Lillo, M., Alvez, A., Delgado, V., Gonzalo-Martín, C., Menasalvas, E., Costumero, R., and
828 García-Pedrero, Á.: Legal disputes as a proxy for regional conflicts over water rights in Chile, *Journal of Hydrology*, 535,
829 36–45, doi:10.1016/j.jhydrol.2016.01.057, 2016.

830

831 Rousseeuw, P. J., and Croux, C.: Alternatives to the Median Absolute Deviation. *Journal of the American Statistical*
832 *Association*, 88(424), 1273–1283, doi:10.1080/01621459.1993.10476408, 1993.

833

834 Saavedra, F. A., Kampf, S. K., Fassnacht, S. R., and Sibold, J. S.: A snow climatology of the Andes Mountains from MODIS
835 snow cover data, *Int. J. Climatol.*, 37, 1526–1539, doi:10.1002/joc.4795, 2017.

836

837 Saavedra, F. A., Kampf, S. K., Fassnacht, S. R., and Sibold, J. S.: Changes in Andes snow cover from MODIS data,
838 2000–2016, *The Cryosphere*, 12, 1027–1046, doi:10.5194/tc-12-1027-2018, 2018.

839

840 Saavedra F., Hernández-Duarte A., Caro A., Rabatel A., Fassnacht S., Condom T., Kampf S., Masiokas M., Romero C.,
841 Aguirre Y., González D., Contreras V., Medina J. and Arancibia P.: Spatial and temporal variability of snow in the Andes
842 using MODIS snow product 2000–2025. *Front. Earth Sci.* 14:1564035, doi: 10.3389/feart.2026.1564035, 2026.

843

844 Sarricolea, P., Herrera-Ossandon, M., and Meseguer-Ruiz, Ó.: Climatic regionalisation of continental Chile, *J. Maps*, 13,
845 66–73, doi:10.1080/17445647.2016.1259592, 2017.

846

847 Scaff, L., Rutllant, J. A., Rahn, D., Gascoin, S., and Rondanelli, R.: Meteorological interpretation of orographic precipitation
848 gradients along an andes west slope basin at 30° S (Elqui valley, Chile). *J. Hydrometeorol.* 18 (3), 713–727,
849 doi:10.1175/JHM-D-16-0073.1, 2017.

850

851 Schaefer, M., Fonseca-Gallardo, D., Farías-Barahona, D., and Casassa, G.: Surface energy fluxes on Chilean glaciers:
852 measurements and models, *The Cryosphere*, 14, 2545–2565, doi:10.5194/tc-14-2545-2020, 2020.

853

854 Servicio Nacional de Geología y Minería (SERNAGEOMIN). Anuario de la Minería de Chile (Tech. Rep.). Servicio
855 Nacional de Geología y Minería, 2021.

856

857 Shaw, T. E., Caro, A., Mendoza, P., Ayala, Á., Pellicciotti, F., Gascoin, S., and McPhee, J.: The utility of optical satellite
858 winter snow depths for initializing a glacio-hydrological model of a high-elevation Andean catchment, *Water Resour. Res.*,
859 56, e2020WR027188, doi:10.1029/2020WR027188, 2020a.

860

861 Shaw, T. E., Gascoin, S., Mendoza, P. A., Pellicciotti, F., and McPhee, J.: Snow Depth Patterns in a High Mountain Andean
862 Catchment from Satellite Optical Tristereoscopic Remote Sensing, *Water Resour. Res.*, 56, doi:10.1029/2019wr024880,
863 2020b.

864

865 Stewart, I. T.: Changes in snowpack and snowmelt runoff for key mountain regions, *Hydrological Processes*, 23, 78–94,
866 doi:10.1002/hyp.7128, 2009.

867

868 Viale, M. and Nuñez, M. N.: Climatology of Winter Orographic Precipitation over the Subtropical Central Andes and
869 Associated Synoptic and Regional Characteristics, *J. Hydrometeorol.*, 12, 481–507, doi:10.1175/2010JHM1284.1, 2011.

870

871 Voordendag, A., Réveillet, M., MacDonell, S., and Lhermitte, S.: Snow model comparison to simulate snow depth evolution
872 and sublimation at point scale in the semi-arid Andes of Chile, *Cryosphere*, 15, 4241–4259, doi:10.5194/tc-15-4241-2021,
873 2021.

Table A1. Characteristics of 81 selected stations with snow depth observations in Chile and Argentina across 21–54° S

Name	Latitude (°)	Longitude (°)	Elevation (m a.s.l.)	Basin	Source	Code	Snow depth sensor
Volcán Auncanquilcha	-21.1906	-68.4528	5,087	Fronterizas Salar Michincha	DGA	02000002	Campbell/SR50A was replaced by Lufft/SHM31
Cerro Chajnantor Cumbre	-22.9872	-67.7422	5,625	Endorreica entre Front. y S. Atacama	DGA	02400002	Campbell/SR50A was replaced by Lufft/SHM31
Volcán Ojos del Salado bajo Refugio Atacama	-27.0519	-68.5350	5,201	Endorreicas entre Front. y Vertiente	DGA	03031002	Campbell/SR50A was replaced by Lufft/SHM31
Volcán Ojos del Salado en Refugio Tejos	-27.0869	-68.5378	5,804	Endorreicas entre Front. y Vertiente	DGA	03031003	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Maranceles	-28.4572	-69.7236	3,985	Río Copiapo	DGA	03413001	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Tapado en Los Corrales	-30.1554	-69.8829	4,065	Río Elqui	DGA	04300004	Campbell/SR50A was replaced by Lufft/SHM31
Tapado (TPF)	-30.1584	-69.9084	4,306	Río Elqui	CEAZA	0430004	Lufft/SHM31
Los Corrales	-30.1615	-69.8759	3,982	Río Elqui	CEAZA	0430014	Campbell/SR50A
Capayán	-30.1734	-69.8038	4,754	Río Jáchal	IANIGLA	Capayán	Campbell/SR50A
La Laguna	-30.2032	-70.0371	3,209	Río Elqui	CEAZA	0430009	Campbell/SR50A
Llano de Las Liebres	-30.2542	-69.9345	3,565	Río Elqui	CEAZA	0430012	Campbell/SR50A
Cerro Olivares	-30.2567	-69.9355	3,566	Río Elqui	DGA	04300001	Campbell/SR50A was replaced by Lufft/SHM31
El Jote	-30.4053	-70.2795	3,660	Río Elqui	CEAZA	0430003	Campbell/SR50A
Quebrada Larga Cota 3500	-30.7252	-70.2918	3,550	Río Limarí	DGA	04520006	Campbell/SR50A was replaced by Lufft/SHM31
Cerro Vega Negra	-30.9059	-70.5165	3,529	Río Limarí	DGA	04511004	Campbell/SR50A was replaced by Lufft/SHM31
Tascadero	-31.2628	-70.5399	3,427	Río Limarí	CEAZA	0450001	Campbell/SR50A
Casa del Canto	-31.4173	-70.5890	3,570	Río Choapa	CEAZA	0470001	Campbell/SR50A
El Soldado	-32.0066	-70.3213	3,293	Río Choapa	DGA	04700002	Campbell/SR50A was replaced by Lufft/SHM31
Nacimiento Del Sobrante	-32.1889	-70.4829	3,143	Río Petorca	DGA	05100003	Campbell/SR50A was replaced by Lufft/SHM31
Las Cuevas	-32.8130	-70.0530	3,197	Río Mendoza	IANIGLA	Cuevas	Campbell/SR50A
Portillo	-32.8424	-70.0970	3,032	Río Aconcagua	DGA	05401007	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar León Negro	-32.9936	-70.0283	4,350	Río Aconcagua	DGA	05400005	Campbell/SR50A was replaced by Lufft/SHM31
Portillo Argentino	-33.6219	-69.6078	4,336	Río Mendoza	IANIGLA	Portillo	Campbell/SR50A

Table A1. Characteristics of 81 selected stations with snow depth observations in Chile and Argentina across 21–54° S

Name	Latitude (°)	Longitude (°)	Elevation (m a.s.l.)	Basin	Source	Code	Snow depth sensor
Glaciar Juncal Sur	-33.1175	-70.1119	4,035	Río Maipo	DGA	05706004	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Olivares Gamma	-33.1564	-70.1528	3,628	Río Maipo	DGA	05706002	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Olivares Alfa	-33.1786	-70.2181	4,230	Río Maipo	DGA	05706005	Campbell/SR50A was replaced by Lufft/SHM31
Valle Olivares	-33.1910	-70.1141	2,786	Río Maipo	DGA	05706003	Campbell/SR50A was replaced by Lufft/SHM31
Yerba Loca Alto	-33.2092	-70.2769	3,340	Río Maipo	UdeChile	AMTC10	Campbell/SR50A
Yerba Loca Medio	-33.2730	-70.3000	2,520	Río Maipo	UdeChile	AMTC8	Campbell/SR50A
Molina en Cepo	-33.2910	-70.2173	3,380	Río Maipo	UdeChile	AMTC13	Campbell/SR50A
Piuquenes 10	-33.3126	-70.2481	3,570	Río Maipo	UdeChile	Piuquenes 10	MaxBotix/MB7389 HRXL-MaxSonar-WRMT
Piuquenes 7	-33.3133	-70.2561	3,483	Río Maipo	UdeChile	Piuquenes 7	MaxBotix/MB7389 HRXL-MaxSonar-WRMT
Piuquenes 6	-33.3165	-70.2611	3,556	Río Maipo	UdeChile	Piuquenes 6	MaxBotix/MB7389 HRXL-MaxSonar-WRMT
Piuquenes 14	-33.3176	-70.2516	3,380	Río Maipo	UdeChile	Piuquenes 14	MaxBotix/MB7389 HRXL-MaxSonar-WRMT
La Parva	-33.3306	-70.2972	2,703	Río Maipo	DGA	05721019	Campbell/SR50A was replaced by Lufft/SHM31
Farellones	-33.3517	-70.3128	2,452	Río Maipo	DGA	05720005	Campbell/SR50A was replaced by Lufft/SHM31
Molina Medio	-33.3753	-70.2413	2,200	Río Maipo	UdeChile	AMTC12	Campbell/SR50A
Glaciar Tupungatito Bajo	-33.3858	-69.8381	4,425	Río Maipo	DGA	05705004	Campbell/SR50A was replaced by Lufft/SHM31
Portezuelo Glaciar Echaurren Norte	-33.5764	-70.1289	3,847	Río Maipo	DGA	05703013	Campbell/SR50A was replaced by Lufft/SHM31
Valle Echaurren Norte	-33.5939	-70.1103	2,950	Río Maipo	DGA	05703012	Campbell/SR50A was replaced by Lufft/SHM31
Termas Del Plomo	-33.6133	-69.9058	3,027	Río Maipo	DGA	05703014	Campbell/SR50A was replaced by Lufft/SHM31
Laguna Negra	-33.6659	-70.1077	2,785	Río Maipo	DGA	05703009	Campbell/SR50A was replaced by Lufft/SHM31
Las Melosas Ruta de Nieve	-33.8636	-70.2727	3,317	Río Maipo	DGA	05701010	Campbell/SR50A was replaced by Lufft/SHM31
Las Hualtatas	-34.0677	-70.1089	2,020	Río Maipo	DGA	05701011	Campbell/SR50A was replaced by Lufft/SHM31
Codelco Cerro Teniente	-34.2047	-70.2692	2,950	Río Rapel	DGA	06005005	Campbell/SR50A was replaced by Lufft/SHM31

Table A1. Characteristics of 81 selected stations with snow depth observations in Chile and Argentina across 21–54° S

Name	Latitude (°)	Longitude (°)	Elevation (m a.s.l.)	Basin	Source	Code	Snow depth sensor
Laguna El Yeso	-34.4118	-70.2084	2,140	Río Rapel	DGA	06000004	Campbell/SR50A was replaced by Lufft/SHM31
Laguna Los Cristales	-34.5611	-70.5097	2,357	Río Rapel	DGA	06013008	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Cortaderal	-34.6608	-70.2561	3,156	Río Rapel	DGA	06001000	Campbell/SR50A was replaced by Lufft/SHM31
Glaciar Universidad en Río San Andrés	-34.7186	-70.3592	2,436	Río Rapel	DGA	06023000	Campbell/SR50A was replaced by Lufft/SHM31
Cerro Los Guzmanes	-34.8861	-70.4636	3,125	Río Rapel	DGA	06021002	Campbell/SR50A was replaced by Lufft/SHM31
Termas Del Flaco	-34.8930	-70.3300	2,665	Río Rapel	DGA	06020001	Campbell/SR50A was replaced by Lufft/SHM31
Paso Vergara	-35.1948	-70.5114	2,602	Río Mataquito	DGA	07100006	Campbell/SR50A was replaced by Lufft/SHM31
Toqui Lautaro	-35.2928	-70.5163	2,453	Río Tunuyán	IANIGLA	Toqui	Campbell/SR50A
Lo Aguirre	-35.9713	-70.5721	1,989	Río Maule	DGA	07301000	Campbell/SR50A was replaced by Lufft/SHM31
NUEVA LO AGUIRRE	-35.9723	-70.5718	1,997	Río Maule	DGA	07301002	Campbell/SR50A was replaced by Lufft/SHM31
Los Condores	-36.0097	-70.5478	2,439	Río Maule	DGA	07301001	Campbell/SR50A was replaced by Lufft/SHM31
Nevado Longavi	-36.2344	-71.1435	1,977	Río Maule	DGA	07350008	Campbell/SR50A was replaced by Lufft/SHM31
Nevado Chillan	-36.8473	-71.4160	2,546	Río Itata	DGA	08130009	Campbell/SR50A was replaced by Lufft/SHM31
Volcan Chillan	-36.8939	-71.4116	2,078	Río Biobio	DGA	08374001	Campbell/SR50A was replaced by Lufft/SHM31
Node3	-36.9103	-71.3989	1,938	Río Itata	UdeChile	081003	Campbell/SR50A
Node2	-36.9104	-71.4089	1,809	Río Itata	UdeChile	081002	Campbell/SR50A
Node1	-36.9148	-71.4166	1,633	Río Itata	UdeChile	081001	Campbell/SR50A
Node4	-36.9205	-71.4180	1,695	Río Itata	UdeChile	081004	Campbell/SR50A
Alto Mallines	-37.1578	-71.2430	1,784	Río Biobio	DGA	08372001	Campbell/SR50A was replaced by Lufft/SHM31
Los Corralitos	-37.5772	-71.1575	1,785	Río Biobio	DGA	08370010	Campbell/SR50A was replaced by Lufft/SHM31
Chenqueco	-38.0647	-71.3547	778	Río Biobio	DGA	08308003	Campbell/SR50A was replaced by Lufft/SHM31
Liucura	-38.6449	-71.0907	1,034	Río Biobio	DGA	08301001	Campbell/SR50A was replaced by Lufft/SHM31
Volcán Villarrica Pichillancahue	-39.4350	-71.8758	1,800	Río Valdivia	DGA	10105003	Campbell/SR50A was replaced by Lufft/SHM31

Name	Latitude (°)	Longitude (°)	Elevation (m a.s.l.)	Basin	Source	Code	Snow depth sensor
Complejo Volcánico Mocho Choshuenco	-39.9514	-72.0586	1,565	Río Valdivia	DGA	10110003	Campbell/SR50A was replaced by Lufft/SHM31
Cardenal Samore	-40.6831	-71.9864	920	Río Bueno	DGA	10321001	Campbell/SR50A was replaced by Lufft/SHM31
Cuesta Moraga	-43.3360	-72.3990	575	Río Yelcho	DGA	10710004	Campbell/SR50A was replaced by Lufft/SHM31
Cerro Divisadero	-45.6165	-72.0134	1,430	Río Aysén	CIEP-CECs	1130001	Campbell/SR50A
Cerro Divisadero en el Fraile	-45.6270	-72.0040	1,190	Río Aysén	DGA	11314002	Campbell/SR50A was replaced by Lufft/SHM31
Portezuelo Ibañez	-46.0778	-72.0510	1,080	Río Baker	DGA	11503002	Campbell/SR50A was replaced by Lufft/SHM31
Hielo Norte en Glaciar San Rafael	-46.7053	-73.5936	1,187	Cost. e Islas R. Aisén R. Baker	DGA	11440001	Campbell/SR50A was replaced by Lufft/SHM31
Steffen	-47.5555	-73.6958	30	Río Baker	CIEP	1150003	Campbell/SR50A
Epulef	-49.0523	-72.9505	898	Río Santa Cruz	IANIGLA	Epulef	Campbell/SR50A
Aonikenk	-49.2774	-72.9975	1,233	Río Santa Cruz	IANIGLA	Aonikenk	Campbell/SR50A
Chalten	-49.3381	-72.8844	400	Río Santa Cruz	IANIGLA	Chalten	Campbell/SR50A
Pliegue Tumbado	-49.3793	-72.9368	1,070	Río Santa Cruz	IANIGLA	Tumbado	Campbell/SR50A
Glaciar Snoring	-53.7230	-72.6080	70	Islas al sur Est. de Magallanes	DGA	12730001	Campbell/SR50A was replaced by Lufft/SHM31

Name	Lat (°)	Long (°)	Elevation (m a.s.l.)	Basin	Source	Issue
Volcán Tacora	-17.6940	-69.7730	5,047	Río Lluta Alto	DGA	Poor data quality from 2022 to 2025
Volcán Quimsachata	-19.7510	-68.7870	5,076	Quebrada de Tarapacá	DGA	Poor data quality in 2022
Paso Jama	-22.9220	-67.7010	4,680	Salar de Atacama	DGA	Poor data quality in 2025
Toconao Pueblo	-23.1861	-68.0056	2,491	Salar de Atacama	DGA	Poor data quality from 2017 to 2026; no ground reference height
Paso Sico	-23.8225	-67.4388	4,295	Salar de Atacama	DGA	Poor data quality between 2017 and 2026
Volcán LLullaillaco	-24.6760	-68.5700	4,800	Endorreicas Salar Atacama-Vertiente Pacífico	DGA	Data since 2026
El Tapado 2	-30.1540	-69.8960	4,250	Río Elqui	CEAZA	Poor data quality
Cuesta Blanca	-31.0970	-70.4170	3,284	Río Limari	DGA	Data available only during summer 2025–2026
Los Azules	-31.2300	-70.5360	3,408	Río Limari	DGA	Data available only during summer 2025–2026
Nido de Cóndores	-32.6400	-70.0300	5,563	Mendoza	IANIGLA	Poor data quality

Table A2. Characteristics of 37 excluded stations with snow depth observations in Chile and Argentina across 17–55° S.						
Name	Lat (°)	Long (°)	Elevation (m a.s.l.)	Basin	Source	Issue
Plaza de Mulas	-32.6539	-70.0651	4,320	Mendoza	IANIGLA	Poor data quality between 2018 to 2025
Morenas Coloradas	-32.9601	-69.3700	3,442	Río Mendoza	IANIGLA	Poor data quality
G. Tupungatito Alto	-33.3939	-69.8125	5,500	Río Maipo	DGA	Poor data quality and noisy measurements from 2021 to 2024
Glaciar San Francisco	-33.8040	-70.0710	2,238	Río Maipo	DGA	Poor data quality from 2012 to 2026
Volcán Azufre	-35.3111	-70.5842	3,154	Río Mataquito	DGA	Constant values between 2022–2023; poor data quality
Sierra Velluda	-37.4725	-71.4358	2,702	Río Biobio	DGA	Poor data quality between 2022 and 2025
Cauñicu	-37.7130	-71.4840	621	Río Biobio	DGA	Poor data quality; anomalous peaks during periods without snowfall
Vn. Puyehue	-40.5930	-72.1140	2,120	Río Bueno	DGA	Data since 2026
Monte Tronador	-41.0760	-71.8770	430	Río Petrohue	DGA	No reference height available, but good data series from 2019 to 2025
Vn. Michimahuida Sur	-42.8520	-72.4420	820	Río Yelcho	DGA	Data since 2026
Vc. Yanteles	-43.4464	-72.7914	1,326	Río Corcovado	DGA	Poor data quality during five months in 2024
Teniente Vidal	-45.5970	-72.1140	308	Río Aysen	DGA	Date shift from 2022 to 2025
G. Calluqueo Alto	-47.6000	-72.3990	1,858	Río Baker	DGA	Available data just some months in 2025
Glaciar Lucia	-48.3510	-73.3010	100	Río Pascua	DGA	No data available in 2025
Estero Pantoja	-49.0200	-73.0261	748	Río Pascua	DGA	No data available in 2025
Glaciar Chico	-49.1560	-73.1390	1,611	Río Pascua	DGA	Poor data quality; continuous data between 2022 and 2025
Fiordo Stage	-49.7197	-74.7253	52	Isla Wellington	DGA	Poor data quality from 2023 to 2025
Torre Paine	-51.1750	-72.9540	25	Río Serrano	DGA	Poor data quality from 2023 to 2025
Estación Dorotea	-51.6040	-72.3340	470	Río Hollemberg	DGA	Poor data quality from 2021 to 2025; no ground reference height, but potentially recoverable
Casas Viejas	-51.7000	-72.3260	230	Río Hollemberg	DGA	Poor data quality from 2021 to 2025; no ground reference height
C. Vidal	-52.2410	-72.2030	1,016	Vertiente del Atlántico	DGA	Data since 2026
Villa Tehuelche	-52.4280	-71.4160	190	Rio Penitente	DGA	Poor data quality during several months in 2025
Gran Campo Nevado	-52.8320	-73.2430	166	Costeras e Islas entre R Hollemberg, Golfo Alte. Laguna Blanca	DGA	Data since 2026
Cerro Mirador	-53.1460	-71.0040	580	Costeras e Islas Orientales de la P Brunswick	DGA	Poor data quality from 2022 to 2025; no ground reference height, but potentially recoverable
Pampa Huanaco	-54.0510	-68.8030	150	Río Grande	DGA	No data available for seven months in 2025
Lago deseado	-54.3410	-68.8260	170	Río Grande	DGA	Available data just some months in 2025
Robalo Alto	-54.9780	-67.6740	360	Islas Navarino y Gable	DGA	Available data just some months in 2025

Table A3. Precipitation–year types by river basin identified from percentiles of total precipitation: dry (P0–30), normal (P30–70), and wet (P70–100), based on available data for 2013–2024.			
Year	Elqui	Maipo	Maule-Itata
2013	Normal	-	Normal
2014	Dry	Normal	Normal
2015	Wet	Normal	Normal
2016	Wet	Wet	Dry
2017	Normal	Normal	Normal
2018	Normal	Normal	Normal
2019	Dry	Dry	Dry
2020	Dry	Dry	Dry
2021	Dry	Dry	Dry
2022	Wet	Normal	Normal
2023	Dry	Wet	Wet
2024	Wet	Wet	Normal

877

Table A4. Potential and effective total days by station with snow depth data									
Station name	Latitude (°)	Raw data				Clean data			
		year start	year end	Potential n° days with data	Effective n° days with data (%)	year start	year end	Potential n° days with data	Effective n° days with data (%)
Vn. Auncanquilcha	-21.2	2021-12-01	2024-12-31	1,127	1,085 (96%)	2021-12-01	2024-12-31	1,127	1,062 (94%)
Cerro Chajnantor	-23.0	2022-03-01	2024-12-31	1,037	733 (71%)	2022-03-01	2024-12-31	1,037	622 (60%)
Vn. Ojos del Salado A.	-27.1	2023-01-03	2024-12-31	729	518 (71%)	2023-01-03	2024-12-31	729	481 (66%)
Vn. Ojos del Salado T.	-27.1	2023-01-03	2024-12-31	729	666 (91%)	2023-01-03	2024-12-31	729	592 (81%)
Glaciar Maranceles	-28.5	2022-04-19	2024-12-31	988	988 (100%)	2022-04-19	2024-12-31	988	975 (99%)
G. Tapado Corrales	-30.2	2020-02-27	2024-12-31	1,770	1,769 (100%)	2020-02-27	2024-12-31	1,770	1,742 (98%)
Tapado (TPF)	-30.2	2018-03-21	2023-10-25	2,045	1,853 (91%)	2018-03-21	2023-10-25	2,045	1,844 (90%)
Los Corrales	-30.2	2022-02-16	2023-07-25	525	343 (65%)	2022-02-16	2023-01-30	349	336 (96%)
Capayan	-30.2	2016-12-20	2024-12-31	2,934	2,933 (100%)	2016-12-20	2024-12-31	2,934	2,892 (99%)
La Laguna	-30.2	2014-06-22	2024-12-31	3,846	2,667 (69%)	2015-06-01	2024-12-31	3,502	2,579 (74%)
Llano de Las Liebres	-30.3	2017-05-16	2024-12-31	2,787	1,295 (46%)	2017-05-16	2021-07-31	1,538	1,098 (71%)
Cerro Olivares	-30.3	2020-12-23	2024-12-31	1,470	1,439 (98%)	2020-12-23	2024-12-31	1,470	735 (50%)
El Jote	-30.4	2019-04-02	2024-12-31	2,101	2,098 (100%)	2019-04-02	2024-12-31	2,101	2,057 (98%)
Quebrada Larga Cota	-30.7	2015-02-20	2024-12-31	3,603	3,428 (95%)	2015-03-07	2024-12-31	3,588	3,373 (94%)
Cerro Vega Negra	-30.9	2015-03-06	2024-10-30	3,527	2,940 (83%)	2016-01-01	2024-10-30	3,226	1,536 (48%)
Tascadero	-31.3	2020-02-19	2024-12-31	1,778	1,299 (73%)	2020-02-19	2024-12-31	1,778	1,282 (72%)
Casa del Canto	-31.4	2019-03-14	2024-12-31	2,120	2,120 (100%)	2019-03-14	2024-12-31	2,120	2,093 (99%)
El Soldado	-32.0	2015-06-08	2024-12-31	3,495	3,319 (95%)	2015-06-08	2024-12-31	3,495	3,099 (89%)
Nacimiento Sobrante	-32.2	2017-01-01	2024-12-31	2,922	2,831 (97%)	2017-01-01	2024-12-31	2,922	2,638 (90%)
Las Cuevas	-32.8	2018-01-03	2024-12-31	2,555	2,553 (100%)	2018-01-03	2024-12-31	2,555	2,543 (100%)
Portillo	-32.8	2013-04-24	2024-12-31	4,270	4,258 (100%)	2013-04-24	2024-12-31	4,270	3,748 (88%)
Glaciar León Negro	-33.0	2020-06-10	2024-12-31	1,666	1,392 (84%)	2020-06-10	2024-12-31	1,666	846 (51%)
Portillo Argentino	-33.6	2020-01-10	2023-04-01	1,178	1,178 (100%)	2020-01-10	2022-12-31	1,087	1,074 (99%)
Glaciar Juncal Sur	-33.1	2019-01-18	2024-12-31	2,175	2,175 (100%)	2019-01-18	2024-12-31	2,175	2,115 (97%)
G. Olivares Gamma	-33.2	2014-05-18	2024-12-31	3,881	3,881 (100%)	2014-05-18	2024-12-31	3,881	3,597 (93%)

Table A4. Potential and effective total days by station with snow depth data

Station name	Latitude (°)	Raw data				Clean data			
		year start	year end	Potential n° days with data	Effective n° days with data (%)	year start	year end	Potential n° days with data	Effective n° days with data (%)
Glaciar Olivares Alfa	-33.2	2019-03-26	2024-12-31	2,108	2,008 (95%)	2019-03-26	2024-12-31	2,108	1,954 (93%)
Valle Olivares	-33.2	2014-05-19	2024-12-31	3,880	3,557 (92%)	2015-01-02	2024-12-31	3,652	2,794 (77%)
Yerba Loca Alto	-33.2	2017-10-10	2019-11-22	774	177 (23%)	2017-10-10	2019-11-22	774	176 (23%)
Yerba Loca Medio	-33.3	2016-09-02	2017-10-19	413	364 (88%)	2017-01-01	2017-10-19	292	254 (87%)
Molina en Cepo	-33.3	2017-04-04	2019-11-22	963	118 (12%)	2017-04-04	2017-06-01	59	59 (100%)
Piuquenes10	-33.3	2018-03-11	2020-01-20	681	639 (94%)	2018-05-21	2020-01-20	610	544 (89%)
Piuquenes7	-33.3	2018-03-11	2020-04-24	776	715 (92%)	2018-03-11	2020-04-24	776	702 (90%)
Piuquenes6	-33.3	2018-03-10	2021-12-06	1,368	820 (60%)	2018-03-10	2019-12-31	662	653 (99%)
Piuquenes14	-33.3	2018-03-11	2024-01-09	2,131	2,113 (99%)	2018-05-11	2024-01-09	2,070	1,558 (75%)
La Parva	-33.3	2021-07-20	2024-12-31	1,261	1,261 (100%)	2021-07-20	2024-12-30	1,260	1,236 (98%)
Farellones	-33.4	2023-11-07	2024-12-31	421	420 (100%)	2023-11-07	2024-12-31	421	416 (99%)
Molina Medio	-33.4	2017-04-04	2017-06-01	59	59 (100%)	2017-04-04	2017-06-01	59	58 (98%)
G. Tupungatito Bajo	-33.4	2020-09-24	2024-12-31	1,560	1,560 (100%)	2020-09-24	2024-12-31	1,560	1,342 (86%)
P. G. Echaurren Norte	-33.6	2023-06-07	2024-12-31	574	563 (98%)	2023-06-07	2024-12-31	574	464 (81%)
Valle Echaurren Norte	-33.6	2022-01-12	2024-11-14	1,038	1,028 (99%)	2023-01-01	2024-11-14	684	676 (99%)
Termas Del Plomo	-33.6	2016-01-12	2024-07-04	3,097	2,578 (83%)	2017-03-06	2022-12-31	2,127	1,792 (84%)
Laguna Negra	-33.7	2013-04-10	2024-12-31	4,284	3,896 (91%)	2013-04-10	2024-12-31	4,284	3,844 (90%)
Las Melosas	-33.9	2017-01-01	2024-12-31	2,922	2,551 (87%)	2017-01-01	2024-12-31	2,922	2,509 (86%)
Las Hualtatas	-34.1	2021-05-20	2024-12-31	1,322	1,322 (100%)	2021-05-20	2024-12-31	1,322	1,311 (99%)
Codelco Cerro Teniente	-34.2	2024-05-31	2024-12-31	215	215 (100%)	2024-05-31	2024-12-31	215	210 (98%)
Laguna El Yeso	-34.4	2013-04-11	2024-12-31	4,283	2,750 (64%)	2016-01-01	2024-12-31	3,288	1,978 (60%)
Laguna Los Cristales	-34.6	2013-03-09	2024-12-31	4,316	2,724 (63%)	2013-03-09	2018-08-31	2,002	1,282 (64%)
Glaciar Cortaderal	-34.7	2018-05-31	2024-10-13	2,328	2,163 (93%)	2018-05-31	2023-08-31	1,919	1,829 (95%)
G. Universidad	-34.7	2011-06-06	2024-12-31	4,958	2,461 (50%)	2022-05-17	2024-12-31	960	707 (74%)
Cerro Los Guzmanes	-34.9	2024-04-19	2024-12-31	257	253 (98%)	2024-04-19	2024-12-31	257	249 (97%)
Termas Del Flaco	-34.9	2013-04-09	2024-12-31	4,285	3,855 (90%)	2013-04-09	2024-12-31	4,285	3,410 (80%)
Paso Vergara	-35.2	2016-02-18	2024-12-31	3,240	2,910 (90%)	2016-02-18	2024-12-31	3,240	2,145 (66%)
Toqui Lautaro	-35.3	2016-12-06	2024-10-23	2,879	2,560 (89%)	2017-01-01	2024-10-23	2,853	2,529 (89%)
Lo Aguirre	-36.0	2010-06-11	2024-12-31	5,318	5,011 (94%)	2010-06-11	2024-12-31	5,318	3,885 (73%)
Nueva Lo Aguirre	-36.0	2021-06-23	2024-12-31	1,288	1,288 (100%)	2021-06-23	2024-12-31	1,288	1,285 (100%)
Los Condores	-36.0	2014-11-01	2024-12-31	3,714	3,360 (90%)	2015-01-02	2024-12-31	3,652	2,538 (70%)
Nevado Longaví	-36.2	2011-03-01	2024-12-31	5,055	3,911 (77%)	2011-03-01	2024-12-31	5,055	2,934 (58%)
Nevado Chillán	-36.8	2022-05-02	2024-12-31	975	975 (100%)	2022-05-02	2024-12-30	974	491 (50%)
Vn. Chillán	-36.9	2015-03-20	2024-12-31	3,575	3,390 (95%)	2016-03-04	2024-12-31	3,225	3,106 (96%)
Node3	-36.9	2015-01-01	2017-10-12	1,016	1,016 (100%)	2015-01-01	2017-10-12	1,016	1,015 (100%)
Node2	-36.9	2015-01-01	2017-10-12	1,016	1,016 (100%)	2015-01-01	2017-09-05	979	977 (100%)
Node1	-36.9	2015-01-01	2017-10-12	1,016	1,016 (100%)	2015-01-01	2017-10-12	1,016	1,014 (100%)
Node4	-36.9	2015-01-01	2017-10-12	1,016	1,016 (100%)	2015-01-01	2017-08-07	950	948 (100%)
Alto Mallines	-37.2	2015-12-16	2024-12-31	3,304	3,221 (97%)	2016-01-02	2024-12-31	3,287	3,191 (97%)
Los Corralitos	-37.6	2023-03-17	2024-12-31	656	420 (64%)	2023-03-17	2024-12-31	656	419 (64%)
Chenqueco	-38.1	2023-05-31	2024-12-31	581	562 (97%)	2024-01-02	2024-12-31	365	354 (97%)
Liucura	-38.6	2021-11-23	2024-12-31	1,135	1,129 (99%)	2022-01-02	2022-12-31	364	357 (98%)
Vn. Villarrica	-39.4	2021-12-13	2023-11-10	698	478 (68%)	2021-12-13	2023-11-10	698	385 (55%)
Vn.Mocho Choshuenco	-40.0	2019-01-14	2024-12-31	2,179	1,683 (77%)	2024-04-25	2024-12-31	251	207 (82%)
Cardenal Samore	-40.7	2023-04-14	2024-12-31	628	598 (95%)	2023-04-14	2024-12-31	628	592 (94%)
Cuesta Moraga	-43.3	2024-04-26	2024-12-31	250	248 (99%)	2024-04-26	2024-12-31	250	247 (99%)
Cerro Divisadero	-45.6	2023-06-08	2024-12-31	573	481 (84%)	2023-06-08	2024-12-31	573	467 (82%)

Table A4. Potential and effective total days by station with snow depth data									
Station name	Latitude (°)	Raw data				Clean data			
		year start	year end	Potential n° days with data	Effective n° days with data (%)	year start	year end	Potential n° days with data	Effective n° days with data (%)
Cerro Divisadero Fraile	-45.6	2019-12-31	2024-12-31	1,828	1,743 (95%)	2019-12-31	2024-12-31	1,828	1,056 (58%)
Portezuelo Ibañez	-46.1	2022-07-07	2024-12-31	909	909 (100%)	2022-07-07	2024-12-31	909	906 (100%)
Glaciar San Rafael	-46.7	2015-09-01	2024-12-31	3,410	2,567 (75%)	2021-03-13	2024-12-31	1,390	1,149 (83%)
Steffen	-47.6	2019-12-19	2024-12-31	1,840	1,799 (98%)	2022-01-01	2022-10-09	282	240 (85%)
Epulef	-49.1	2016-01-31	2024-11-05	3,202	1,891 (59%)	2016-01-31	2023-12-31	2,892	1,573 (54%)
Aonikenk	-49.3	2014-03-30	2024-11-10	3,879	3,092 (80%)	2014-03-30	2024-11-10	3,879	3,072 (79%)
Chalten	-49.3	2021-12-09	2024-12-31	1,119	1,089 (97%)	2022-01-01	2024-12-31	1,096	1,053 (96%)
Pliegue Tumbado	-49.4	2020-03-27	2023-04-04	1,104	1,062 (96%)	2020-03-27	2022-12-31	1,010	954 (94%)
Glaciar Snoring	-53.7	2023-08-28	2024-10-14	414	414 (100%)	2024-01-01	2024-10-14	288	285 (99%)

878

Table A5. Summary of raw data reduction of successive quality–control filters for PCI construction.				
Station	Total days with SD, Pr and AT simultaneous variables	Season filter (May to October)	Snow accumulation filter (ASD > 10 mm)	Precipitation filter (Pr > 1 mm)
Glaciar Tapado Corrales	1,085	546	34	33
Cerro Olivares	705	420	24	21
Glaciar Olivares Alfa	1,992	1,008	114	99
Glaciar Juncal Sur	2,112	1,082	162	151
Glaciar Olivares Gamma	2,095	1,100	68	59
Las Melosas	1,015	548	95	82
Termas Del Plomo	2,245	1,065	182	110

879

Table A6. Summary of clean data reduction of successive quality–control filters for PCI construction.					
Station	Total days with SD, Pr and AT simultaneous variables	Season filter (May to October)	Snow accumulation filter (ASD > 10 mm)	Precipitation filter (Pr > 1 mm)	AT _{rain}
Glaciar Tapado Corrales	1,066	538	29	28	-2.2
Cerro Olivares	363	182	11	10	2.2
Glaciar Olivares Alfa	1,907	965	78	72	-3.8
Glaciar Juncal Sur	2,063	1,041	113	110	-3.7
Glaciar Olivares Gamma	1,839	1,050	156	107	-0.9
Las Melosas	983	543	91	79	-1.5
Termas Del Plomo	1,469	710	77	59	3.8

880