

# A Global Database of Soil Methane uptake (SMUD)

Jiawei Jiang<sup>1</sup>, Zhifeng Yan<sup>1,2\*</sup>, Jinshi Jian<sup>3,4\*</sup>, Jesper Riis Christiansen<sup>5</sup>, Xinchu Wang<sup>1,2</sup>, Ren-Min Yang<sup>1</sup>, Zhen Zhang<sup>6</sup>, Philippe Ciais<sup>7</sup>

<sup>1</sup>Institute of Surface-Earth System Science, School of Earth System Science, Tianjin University, Tianjin 300072, China.

<sup>2</sup>Tianjin Bohai Rim Coastal Earth Critical Zone National Observation and Research Station, Tianjin University, Tianjin 300072, China.

<sup>3</sup>State Key Laboratory of Soil and Water Conservation and Desertification Control, College of Soil and Water Conservation Science and Engineering, Yangling District, Xianyang 712100, China.

<sup>4</sup>Institute of Soil and Water Conservation, Chinese Academy of Sciences & Ministry of Water Resources, 26 Xinong Road, Yangling District, Xianyang 712100, China.

<sup>5</sup>Department of Geoscience and Natural Resource Management, University of Copenhagen, Frederiksberg, DK 1958, Denmark

<sup>6</sup>National Tibetan Plateau Data Center (TPDC), State Key Laboratory of Tibetan Plateau Earth System, Environment and Resource (TPESER), Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, 100101, China

<sup>7</sup>Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, 91190, Gif sur Yvette, France.

Corresponding to: [Zhifeng Yan \(yanzf17@tju.edu.cn\)](mailto:yanzf17@tju.edu.cn) and [Jinshi Jian \(jinshi@nwfufu.edu.cn\)](mailto:jinshi@nwfufu.edu.cn)

## Abstract

Soil methane uptake (SMU), the important biological sink for atmospheric methane (CH<sub>4</sub>) in terrestrial ecosystems, represents an important component of the global CH<sub>4</sub> budget. Although field SMU observations have been conducted since the 1980s, they have not yet been systematically compiled into a unified and openly available database. Here, we present the global Soil Methane Uptake Database (SMUD), which includes 2,427 site-level records from 920 peer-reviewed publications spanning 1986 to 2025. SMUD contains SMU measurements at multiple temporal scales, including daily (n = 1,425), monthly (n = 2,001), seasonal (n = 1,720), and annual (n = 1,098), together with associated soil moisture and soil temperature data at daily and monthly scales where available. The collected datasets cover all major climatic zones and ecosystem types and reveal pronounced spatial and temporal variability in SMU. Our analyses show that seasonal- and biome-specific patterns are important for understanding intra-annual variability in SMU. In particular, quantitative classification showed that the relationships of SMU with soil temperature and moisture differed among biomes, making ecosystems to be grouped into four response types: temperature-dominated, moisture-dominated, jointly regulated by temperature and moisture, and no clear dominant control. Overall, SMUD provides a comprehensive basis for characterizing global SMU patterns.

设置了格式: 字体: (中文) + 中文正文 (等线)

设置了格式: 默认段落字体, 字体: (默认) + 西文正文 (等线), (中文) + 中文正文 (等线), 五号

设置了格式: 字体: (中文) + 中文正文 (等线)

设置了格式: 字体: (中文) + 中文正文 (等线), 11 磅

设置了格式: 默认段落字体, 字体: (默认) + 西文正文 (等线), (中文) + 中文正文 (等线), 五号

设置了格式: 字体: (中文) + 中文正文 (等线), 11 磅

删除了: the second-largest and represents a biologically mediated sink for atmospheric methane (CH<sub>4</sub>) in terrestrial ecosystems

删除了: plays a non-negligible role in regulating the global CH<sub>4</sub> budget.

删除了:

删除了: Field SMU observations have been conducted since the 1980s, but have not yet been systematically compiled within any unified and openly available framework.

删除了: along with soil moisture and temperature data at daily and monthly scales.

删除了: The collected datasets cover all major climatic zones and ecosystem types, across which SMU exhibits pronounced spatial and temporal variability

删除了: We found that seasonal- and biome-specific variations were key to understanding intra-annual changes in SMU.

删除了: In particular, the effects of soil temperature and moisture on SMU are biome-specific: observations from Temperate Grassland are primarily associated with temperature changes, whereas those from Tropical Rainforest and Desert are mainly controlled by moisture, and other biomes show a mix of controlling factors.

investigating underlying mechanisms, and improving estimates of the global SMU budget. The database, associated data, and code for reproducing the results are available at <https://doi.org/10.5281/zenodo.18299391> (Jiang et al., 2026). We aim to maintain SMUD as a collaborative, community-supported, and regularly updated resource for the research community.

删除了: Overall, SMUD provides a comprehensive basis for understanding the global patterns of SMU, unraveling the underlying mechanisms, and estimating the global SMU budget.

删除了: to reproduce

设置了格式: 字体: (中文) 等线

## 1 Introduction

### 1.1 The critical role of soil methane uptake (SMU) in the global methane cycle and its climatic significance.

删除了: its broader impacts

Among the major long-lived greenhouse gases, methane ( $\text{CH}_4$ ) has contributed approximately  $0.5^\circ\text{C}$  to global warming and accounts for about 30% of observed global warming (Masson-Delmotte et al., 2021; Shindell et al., 2012). Per unit mass,  $\text{CH}_4$  exerts a stronger warming effect than carbon dioxide ( $\text{CO}_2$ ) over short timescales due to its higher radiative efficiency (Saunois et al., 2025). Since the Industrial Revolution, the persistent increase in atmospheric  $\text{CH}_4$  has reflected a sustained imbalance between its emissions and sinks (Gauci et al., 2024).  $\text{CH}_4$  mitigation has thus become a central focus of international climate policy, as highlighted by the Global Methane Pledge, which aims to reduce anthropogenic  $\text{CH}_4$  emissions by at least 30% by 2030 relative to 2020 levels (Saunois et al., 2025). In this context, a comprehensive understanding of both anthropogenic sources and natural sinks is essential to accurately assess the net  $\text{CH}_4$  budget and mitigation potential.

删除了: three

删除了: On a per-unit basis

删除了:  $\text{CH}_4$

Soils are key environments for  $\text{CH}_4$  cycling, where  $\text{CH}_4$  is consumed by aerobic methanotrophs and generated by anaerobic methanogens. The resulting net  $\text{CH}_4$  flux is therefore determined by the balance between  $\text{CH}_4$  oxidation and production, which is regulated by multiple environmental and biological factors influencing methanotrophic and methanogenic activities (Ni and Groffman, 2018; Bridgman et al., 2013). Soil methane uptake (SMU), driven by the microbial oxidation of atmospheric  $\text{CH}_4$ , represents the second-largest atmospheric  $\text{CH}_4$  sink after atmospheric oxidation and the major biologically mediated oxidative sink in terrestrial ecosystems (Song et al., 2024; Saunois et al., 2025). However, human land-use change has substantially weakened this sink (Smith et al., 2000). For example, without the widespread conversion of natural ecosystems to croplands since the mid-19th century, SMU in temperate regions would likely have been up to three times higher, reaching  $27\text{ Tg CH}_4\text{ yr}^{-1}$  instead of  $8\text{ Tg CH}_4\text{ yr}^{-1}$  (Ojima et al., 1993; Tate, 2015). Given that the global atmospheric  $\text{CH}_4$  burden has been growing at approximately  $19.0\text{--}36.0\text{ Tg CH}_4\text{ yr}^{-1}$  during this century, strengthening SMU could provide a complementary opportunity to partially offset this increase (Saunois et al., 2025). Moreover, SMU represents a potentially manageable terrestrial  $\text{CH}_4$  sink, as it can be directly influenced by human activities through changes in land use, soil properties, and management practices (De Bernardi et al., 2022).

删除了: In this context, a comprehensive understanding of both anthropogenic sources and natural sinks is required to assess the net

设置了格式: 字体: 非加粗

删除了:  $\text{CH}_4$

删除了: For example, without the widespread conversion of natural ecosystems to croplands since the mid-19th century, SMU in temperate regions would likely be higher by up to a factor of three, at

设置了格式: 字体: 非加粗

删除了: Given that the global atmospheric  $\text{CH}_4$  burden is growing at approximately

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

删除了: in this century, enhancing SMU thus represents a potential strategy to partially offset or mitigate this increase

删除了: remains highly uncertain

Global SMU estimates remain highly uncertain, with bottom-up estimates ranging from 11 to  $49\text{ Tg CH}_4\text{ yr}^{-1}$  (Saunois et al., 2025). This uncertainty arises in part from the substantial spatiotemporal variability of SMU (Jiang et al., 2025). Temporally, continuous measurements

have revealed large fluctuations, with interannual variability reaching up to 163.2% (Luo et al., 2012). Intra-annual variability is also pronounced, with distinct seasonal patterns observed across diverse climate zones and ecosystems (Luo et al., 2013). Spatially, SMU exhibits considerable heterogeneity across latitudes (Murguia-Flores et al., 2018) and among biomes (Jiang et al., 2025). These variations are shaped by environmental conditions and management practices, both of which introduce substantial uncertainty into global SMU estimation (Song et al., 2024). Current global SMU estimates generally rely on data-driven or process-based approaches. Data-driven methods aggregate observations across ecosystems or climate zones but are limited by observational bias, limited spatial coverage and coarse spatial representation. Process-based models simulate grid-scale SMU by representing CH<sub>4</sub> oxidation and diffusion processes, but they commonly rely on spatially invariant or weakly constrained parameters, which may be inconsistent with observations and lead to divergent estimates (Jiang et al., 2025). Together, these limitations indicate that a major source of uncertainty in current global SMU estimates arises from model parameterization and the limited observational basis used to constrain model parameters. Addressing this issue requires a more comprehensive, standardized, and robust global database to better support model development, parameter optimization, and large-scale SMU assessment.

## 1.2 The need for a unified global SMU database

Many field measurements have been conducted since the 1980s to measure SMU across a wide range of climatic conditions and ecosystems (see Supplementary file “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410”, Fig. 1). However, most existing SMU estimates rely on small-scale and site-specific observations, which remain difficult to synthesize because of fragmented reporting, inconsistent measurement units, and the lack of standardized approaches across research groups. Therefore, standardizing and integrating existing SMU measurements into a coherent data framework is essential for supporting synthesis studies and advancing global CH<sub>4</sub> cycle research.

Since 2000, several databases based on in situ measurements have been developed to support global SMU estimates (Jiang et al., 2025). However, these databases have limited representation across ecosystems and climatic zones and are therefore insufficient to fully capture the spatiotemporal variability of SMU. For example, Grosso et al. (2000) focused on observations from Europe and North America rather than providing a global assessment. Dutaur et al. (2007) presented a global estimate, but their dataset was still dominated by observations from Europe and North America. Although Murguia-Flores et al. (2018) reported global SMU estimates, their analysis was based on a relatively limited database comprising approximately 100 observations from about 30 studies.

Although the number of SMU measurements has increased over time, existing databases remain inadequate for robustly estimating global SMU. Most global SMU estimates have been derived from data with low temporal resolution, often relying on annual averages that overlook important intra-annual variability (Li et al., 2025). In addition, many previous estimates used simplified ecosystem classifications (Li et al., 2025) or focused on a single ecosystem type (Ni and Groffman, 2018), limiting their usefulness for comprehensive syntheses of global SMU estimates.

删除了: These spatial differences are largely driven by variations in environmental conditions and management practices, which introduce substantial uncertainty in estimating global SMU

删除了: Global SMU estimates rely on data-driven and process-based approaches

删除了: Data-driven methods aggregate observations by ecosystem or climate but are limited by data bias and coarse spatial representation.

删除了: Process-based models simulate grid-scale SMU using CH<sub>4</sub> oxidation and diffusion processes, yet commonly assume invariant parameters inconsistent with observations, resulting in divergent estimates

删除了: field experiments

删除了: CH<sub>4</sub>

删除了: CH<sub>4</sub>

删除了: However, most existing SMU estimates rely on small-scale, site-specific observations, which are fragmented and unsystematic due to diverse measurement units and the absence of standardized approaches across research groups

删除了: past SMU measurements

删除了: CH<sub>4</sub>

删除了: Several databases have been developed based on in situ measurements to support global SMU estimates since 2000

删除了: However, these databases lack comprehensive representation of different ecosystems and climatic zones, and are insufficient to capture spatiotemporal variability due to data scarcity.

删除了: For example, Grosso et al. (2000) focused exclusively on Europe and North America observations rather than providing a global assessment, whereas Dutaur et al. (2007) presented a global estimate that was nevertheless dominated by data from Europe and North America; similarly, although Murguia-Flores et al. (2018) reported

删除了: Although the count of measurements has grown over time, existing databases are still inadequate to robustly

删除了: Most global SMU estimates have been derived from low temporal resolution data and primarily annual averages

Consequently, compiling comprehensive and standardized global SMU measurements, together with associated soil physicochemical properties where available, is essential for generalizing and upscaling existing observations and improving the reliability of SMU estimates. Such a database would also facilitate research linking SMU to key soil biogeochemical processes (Aronson and Helliher, 2010) and plant-associated processes (Voigt et al., 2023; Chai et al., 2020) that regulate SMU, as well as modeling SMU responses to changing environmental conditions (Song et al., 2024; Murguía-Flores et al., 2018).

Our primary aim is to harmonize disparate data sources, standardize measurement units, compile associated metadata, and integrate global SMU observations into a comprehensive database across multiple temporal and spatial scales. We also aim to provide an open and robust platform that facilitates synthesis, benchmarking, and modeling of SMU dynamics under varying climatic and environmental conditions. Furthermore, we aim to foster collaboration and knowledge exchange within the scientific community, thereby supporting evidence-based assessments relevant to CH<sub>4</sub> mitigation planning and climate change research.

## 2 Methods

### 2.1 Database compilation

Relevant peer-reviewed publications reporting SMU were identified through a systematic search of two literature databases: Web of Science (<https://clarivate.com.cn/solutions/web-of-science/>) and the China National Knowledge Infrastructure (CNKI, <http://www.cnki.net/>). The search covered peer-reviewed articles published between 1986 and 2025, and the search strategy used Boolean logic with the following items: TI = (CH<sub>4</sub> OR methane OR greenhouse gas) AND TS = (soil AND (\*flux OR oxidat\* OR uptake OR consumption OR exchange OR sink)). The title, abstract, figures, and tables of each study were screened to determine eligibility. In total, approximately 10,800 papers were retrieved.

Studies were considered eligible if they met the following criteria: (i) they reported in situ field measurements of SMU; laboratory incubation experiments and purely modeling-based studies without field observations were excluded; (ii) when multiple experimental treatments were reported, the unamended control treatment was selected; (iii) they reported at least one type of temporally aggregated SMU data, including daily, weekly, monthly, seasonal, or annual values. Measurements made at hourly resolution within a single day were excluded. The flux data in SMUD were organized into four categories based on temporal aggregation: Daily (see Supplementary file “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410”), Monthly, Seasonal, and Annual (see Supplementary file “Studies and Fluxes”). In total, 920 papers met these criteria, with about 84% (773) from the Web of Science and the rest from CNKI (Fig. 1). In this study, “site-level records” refer to unique database entries compiled from individual studies and sites, whereas “measurements” refer to CH<sub>4</sub> flux values available at specific temporal resolutions. A single site-level record may contribute measurements at daily, monthly, seasonal, and annual resolutions. SMU data were provided by the authors or extracted by digitizing graphs in the papers using the GetData Graph Digitizer (version 2.26, <http://www.getdata-graph-digitizer.com/>) when data were presented in the graphs. Seasonal and annual values were

删除了: Consequently, obtaining comprehensive and accurate global SMU measurements, coupled with associated soil physicochemical properties that can be used to generalize and upscale existing observations, is essential for improving the reliability of SMU estimates

删除了: Additionally, such a database would facilitate further research into linking and modeling key soil biogeochemical

删除了: and plant biological processes

删除了: as well as modelling how SMU responds to dynamic and changing environmental conditions

删除了: Our primary aim is to harmonize disparate data sources, standardize measurement units, ensure complete metadata, and integrate global SMU observations to construct a comprehensive database across different temporal and spatial scales

删除了: We also aim to provide a robust and open platform that facilitates advanced research and enhances the modeling and prediction of CH<sub>4</sub> dynamics under varying climatic and environmental conditions.

删除了: Furthermore, we hope to foster collaboration and knowledge exchange within the scientific community, ultimately informing policy decisions and mitigation strategies to address increasing atmospheric CH<sub>4</sub> concentrations and their implications for climate change.

删除了: C

删除了: All available studies reporting SMU were compiled from peer-reviewed publications through a systematic search in databases: Web of Science

删除了: CH<sub>4</sub>

删除了: were used to decide whether to include it.

删除了: In total, almost 10040 papers were retrieved.

删除了: the study reported in situ field measurements of SMU

删除了: the control group without amendment was selected when multiple treatments were applied

删除了: at least one of the following types of SMU data was reported: mean annual, seasonal, monthly, weekly, or daily values; measurements at the hourly scale within a single day

删除了: three

aggregated from shorter timescale fluxes when not provided in the original publication. Additional site information, including location latitude, longitude, elevation, country, region, climatic characteristics, and ecosystem type (e.g., forest, grassland, and agriculture) was also collected (Table 1, Supplementary file “Studies and Fluxes”). Mean annual temperature (MAT) and mean annual precipitation (MAP) for each site were extracted from ERA5-Land Monthly Aggregated data (Fig. S1, doi:10.24381/cds.68d2bb30). When latitude, longitude, or elevation was missing, the missing information was supplemented using Google Earth (https://earth.google.com/) based on the original site description.

The data were categorized according to the Whittaker biome classification (Fig. 2), including tundra (Tun), boreal forest (BF), temperate seasonal forest (TSF), temperate rainforest (TRF), temperate grassland (TG), desert (Des), tropical rainforest (TRRF), tropical seasonal forest (TRSF), savanna (Sav), subtropical desert (SDes), woodland (Wood), and shrubland (Shrub). Observations outside the Whittaker biome framework were reclassified into additional categories, including alpine (Alp), agricultural land (Agri), bare land (Bare), urban areas (Urban), and others. The “others” category included special habitats or sources not represented in the Whittaker framework, such as saline-alkaline soils, blanket peats, landfills, wetlands, riparian zones, floodplains, ant nests, and atmosphere–soil–cave systems.

## 2.2 Database structure

The “Studies and Fluxes” file contains publication information, site-specific metadata, including location, climatic characteristics, and ecosystem type, and experimental details, including sampling time, measurement duration, sampling frequency, and measurement methods; field descriptions are provided in Table 1. This file also includes monthly, seasonal, and annual SMU fluxes, either as reported by the original authors or derived from shorter-timescale measurements. The second file, named “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410”, contains the compiled daily and monthly SMU measurements (Table 2). Each record includes information on the sampling year, day of year (DOY), and corresponding SMU fluxes. A unique identifier (Paper–Study ID) links corresponding entries between the two files.

In the “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410” the original SMU units reported in the source publications were retained and described using four components: gas species, mass basis, area basis, and time basis. For cross-study comparison, all SMU values were converted to a standardized unit of  $\mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ . When CH<sub>4</sub> fluxes were expressed as CO<sub>2</sub>-equivalents, a 100-year global warming potential factor of 27.2 was used for CH<sub>4</sub> relative to CO<sub>2</sub> (i.e., Global Warming Potential of CH<sub>4</sub> relative to CO<sub>2</sub> over a 100-year period) (Masson-Delmotte et al., 2021). The “Studies and Fluxes” file contains 66 fields (Table 1), whereas the “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410” file contains 6 fields (Table 2).

Table 1. Categories of database fields and corresponding descriptions for the Supplementary file “Studies and Fluxes”. SMU values in this file have been standardized to unified units of  $\mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$  for comparison across studies.

- 删除了: Additional site information, including location (latitude, longitude, elevation, country, and region), climate
- 删除了: see
- 删除了: and
- 删除了: were acquired
- 删除了: ,
- 删除了: When latitude, longitude, and elevation were
- 设置了格式
- 删除了: The data were categorized according to biomes in
- 删除了: Observations outside the Whittaker biome
- 删除了: SMUD consists of two separate Supplementary file
- 删除了: This file also includes monthly, seasonal, and annual
- 删除了: CH<sub>4</sub>
- 删除了: CH<sub>4</sub>
- 删除了: acquired daily and monthly SMU measurements
- 删除了: An index number (paper-study ID) maps entries
- 删除了: SMU units in the
- 删除了: CH<sub>4</sub>
- 删除了: CH<sub>4</sub>
- 删除了: file follow the original units reported in the source
- 删除了: All SMU values were converted to
- 删除了: CH<sub>4</sub>
- 删除了: <sup>2</sup>
- 删除了: yr
- 删除了: ·
- 设置了格式: 字体颜色: 自动设置, 降低量 2 磅
- 设置了格式: 字体: 非加粗
- 设置了格式: 下标
- 删除了: The calculation of CO<sub>2</sub>-equivalents of CH<sub>4</sub> used a
- 设置了格式: 下标
- 删除了: CH<sub>4</sub>
- 删除了: CH<sub>4</sub>
- 删除了: CH<sub>4</sub>
- 删除了: Both files are provided in comma-separated values
- 带格式的: 缩进: 首行缩进: 1 字符

| Num                                          | Field name                   | Unit                                                        | Description                                                                                                 |
|----------------------------------------------|------------------------------|-------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|
| Metadata                                     |                              |                                                             |                                                                                                             |
| 1                                            | Paper_number                 | -                                                           | Paper number                                                                                                |
| 2                                            | Study_number                 | -                                                           | Study number                                                                                                |
| 3                                            | Paper_title                  | -                                                           | Paper Title                                                                                                 |
| 4                                            | Paper_author                 | -                                                           | The name of the first author                                                                                |
| 5                                            | Paper_year                   | -                                                           | The year the paper was published                                                                            |
| 6                                            | Paper_journal                | -                                                           | Journal of Publication                                                                                      |
| 7                                            | Paper_language               | -                                                           | English or Chinese                                                                                          |
| 8                                            | Entry_date                   | -                                                           | Entry date                                                                                                  |
| 9                                            | Collector                    | -                                                           | Data collector                                                                                              |
| 10                                           | Checker                      | -                                                           | Data checker                                                                                                |
| Site and condition                           |                              |                                                             |                                                                                                             |
| 11                                           | Continent                    | -                                                           | Continent                                                                                                   |
| 12                                           | Country                      | -                                                           | Country                                                                                                     |
| 13                                           | Region                       | -                                                           | State/province/region                                                                                       |
| 14                                           | Site_name                    | -                                                           | Name of study site                                                                                          |
| 15                                           | Study_duration               | month                                                       | Duration of measurement                                                                                     |
| 16                                           | Study_frequence              | per month                                                   | frequency of measurement                                                                                    |
| 17                                           | Flux_method                  | -                                                           | Flux measurement method:<br>Static chamber; Eddy covariance<br><u>CH<sub>4</sub></u> concentration analysis |
| 18                                           | <u>CH<sub>4</sub></u> method | -                                                           | technique: Gas chroma<br>Laser spectroscopy                                                                 |
| 19                                           | Latitude                     | -                                                           | Latitude, decimal; positive =<br>North, negative = South                                                    |
| 20                                           | Longitude                    | -                                                           | Longitude, decimal; positive =<br>East, negative = West                                                     |
| 21                                           | Elevation                    | m                                                           | Elevation                                                                                                   |
| 22                                           | MAT                          | °C                                                          | Reported mean annual<br>temperature                                                                         |
| 23                                           | MAP                          | mm                                                          | Reported mean annual<br>precipitation                                                                       |
| 24                                           | Climate                      | -                                                           | Climate zones of sites                                                                                      |
| 25                                           | Ecosystem                    | -                                                           | Ecosystems of sites                                                                                         |
| 26                                           | Biome                        | -                                                           | Biomes are categorized according<br>to the Whittaker biome diagram<br>(see Methods)                         |
| Annual, seasonal, and monthly SMU            |                              |                                                             |                                                                                                             |
| 27-37                                        | SMU Jan to Dec               | μg <u>CH<sub>4</sub></u> -C m <sup>-2</sup> h <sup>-1</sup> | Monthly SMU flux                                                                                            |
| 38-41                                        | SMU Spr to Win               | μg <u>CH<sub>4</sub></u> -C m <sup>-2</sup> h <sup>-1</sup> | Seasonal SMU flux                                                                                           |
| 42                                           | annual                       | μg <u>CH<sub>4</sub></u> -C m <sup>-2</sup> h <sup>-1</sup> | Annual SMU flux                                                                                             |
| Soil moisture (SM) and soil temperature (ST) |                              |                                                             |                                                                                                             |
| 43-55                                        | SM Jan to Dec                | WFPS (%)                                                    | Monthly mean soil moisture at<br>topsoil                                                                    |

|       |               |    |                                          |
|-------|---------------|----|------------------------------------------|
| 56-66 | ST Jan to Dec | °C | Monthly mean soil temperature at topsoil |
|-------|---------------|----|------------------------------------------|

Table 2. Categories of database fields and corresponding descriptions for the Supplementary file “CH<sub>4</sub> uptake data CH<sub>4</sub> FLUX 1 1410”. Original flux values and units are preserved as reported in the corresponding literature records and have not been standardized in this table.

| Num | Field name   | Unit                         | Description                                                                                                                                                                                                   |
|-----|--------------|------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1   | Paper_number | -                            | Paper number                                                                                                                                                                                                  |
| 2   | Study_number | -                            | Study number                                                                                                                                                                                                  |
| 3   | Meas_year    | -                            | The year the study was performed<br>This field was used to separate different sub-records (e.g., layers or treatments) reported within the same paper, study, and year; these sub-records were then averaged. |
| 4   | Meas_mark    | -                            | A single number is used to denote the day of year                                                                                                                                                             |
| 5   | DOY          | -                            |                                                                                                                                                                                                               |
| 6   | Fluxes       | Given by the original papers | Corresponding observation fluxes                                                                                                                                                                              |

## 2.3 Quality control

A quality-control (QC) procedure was applied to the database, and quality flags (Q01–Q07) were used to identify records with known issues, potential uncertainties, or specific data characteristics. When multiple sources of the same SMU value were available, values directly reported by the authors were prioritized over values calculated from other reported data or digitized from figures. Laboratory incubation data (Q01) and duplicate records (Q02) were excluded from subsequent analyses. The specific sampling time was recorded to improve documentation of SMU variability at fine temporal scales, which can influence monthly, seasonal, and annual estimates. When the exact sampling time was not reported, the corresponding records were flagged as Q03 and assigned to the “Other” category for sampling time in the “Studies and Fluxes” file. Because incorrect units may result in unrealistic flux values, confirmed and suspected unit errors were identified based on unit-conversion checks and information from figures, tables, and relevant textual descriptions; these records were flagged as Q04. Records with confirmed erroneous flux units in the original publications were excluded from the analysis. Site coordinates and locations were checked using Google Earth (<https://earth.google.com/>) to identify and correct inaccurately reported site locations; corrected records were flagged as Q05. For records lacking coordinate information, geographic coordinates were supplemented based on the descriptions of study sites or plots. Records with unresolved uncertainties or other known data issues that could not be assigned to Q01–Q05 were flagged as Q06 to alert users to potential problems. In addition, daily-scale records were identified using Q07; this flag indicates temporal resolution rather than a data-quality problem.

带格式的: 缩进: 首行缩进: 0 字符

删除了: CH4

删除了: CH4

删除了: A basic quality-control procedure was applied to the database, and quality flags (Q01-Q07) were used to identify entries with known issues or potential problems.

删除了: We gave preference to data directly reported by the authors, under the assumption that such data were of higher quality than those calculated or extracted from figures or tables.

删除了: Incubation data (Q01) and duplicate records (Q02) were excluded.

删除了: ensure accurate documentation

删除了: When the exact sampling time was not reported, the corresponding records were flagged in Q03 and classified into the “Other” category (Supplementary file “Studies and Fluxes”)

删除了: Incorrect units often result in outliers; therefore, both confirmed and suspected unit errors were identified based on unit conversion results, as well as information from figures, tables, and surrounding text (Q04).

删除了: Data with erroneous flux units in the original publication were excluded from the analysis

删除了: Coordinate projections in Google Earth

删除了: were used to identify and correct inaccurately reported site locations (Q05)

删除了: supplied

删除了: local plots

删除了: Despite these efforts, some errors persisted; therefore, records annotated as “other known problem with data” were systematically flagged to alert users to potential issues (Q06).

删除了: In addition, we identified and marked the daily-scale data (Q07).

## 2.4 Data analysis

The consistency between MAT and MAP reported in the original studies and those derived from ERA5-Land was assessed by extracting ERA5-Land values from the 0.1° grid cells corresponding to site coordinates and fitting linear regressions between ERA5-Land-derived and study-reported values. Regression slope, intercept,  $R^2$ , and  $N$  were reported, with the 1:1 line shown as a reference (Fig. S1). ERA5-Land-derived MAT and MAP showed strong agreement with study-reported values (MAT:  $R^2 = 0.86$ , slope = 0.898; MAP:  $R^2 = 0.79$ , slope = 0.781). Locally measured or site-reported MAT and MAP values were prioritized when available, particularly for sites located in mountainous areas with strong elevation gradients. SMU values were obtained either directly from the literature or derived from shorter-timescale measurements. Monthly SMU fluxes from January to December were calculated as arithmetic means of daily fluxes within the corresponding DOY ranges for each site or study record. Seasons were defined as follows: spring (March–May in the Northern Hemisphere and September–November in the Southern Hemisphere), summer (June–August in the Northern Hemisphere and December–February of the following year in the Southern Hemisphere), autumn (September–November in the Northern Hemisphere and March–May in the Southern Hemisphere), and winter (December–February of the following year in the Northern Hemisphere and June–August in the Southern Hemisphere). Seasonal fluxes were calculated only if data were available for all months within the corresponding season. Annual values were calculated when field measurements covered a full 12-month period and included values for all four seasons, or when the original study reported paired seasonal values representing the full annual cycle, such as growing versus dormant seasons or wet versus dry seasons. All SMU values were expressed in or converted to units of  $\mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ .  $\text{CH}_4$  uptake was defined as positive, and only uptake observations were used in subsequent analyses. Monthly soil moisture and soil temperature were also calculated from daily observations using the same DOY ranges. Soil moisture was expressed as water-filled pore space (WFPS). If WFPS was not reported, its value was calculated based on the reported volumetric ( $\theta_v$ ,  $\text{m}^3 \text{m}^{-3}$ ) or gravimetric ( $\theta_g$ ,  $\text{g g}^{-1}$ ) soil moisture as follows (Borken et al., 2006):

$$\text{WFPS} = \theta_v / (1 - \text{BD} / 2.65) = \theta_g \times \text{BD} / (1 - \text{BD} / 2.65) \quad (1)$$

where  $\text{BD}$  is bulk density ( $\text{g cm}^{-3}$ ); when  $\text{BD}$  was not reported, it was retrieved from SoilGrids250m 2.0 (Poggio et al., 2021), and  $2.65 \text{ g cm}^{-3}$  is the assumed particle density. All data were initially organized in Microsoft Excel and archived as CSV files in the supplementary files “Studies and Fluxes” and “ $\text{CH}_4$  uptake data  $\text{CH}_4$  FLUX 1 1410”. The coefficient of variation ( $CV$ ) was used to quantify the relative magnitude of intra-annual SMU variability:

$$\sigma = \sqrt{\sum_{i=1}^N (x_i - \mu)^2 / N} \quad (2)$$

$$CV = (\sigma / \mu) \cdot 100\% \quad (3)$$

where  $\sigma$  is the standard deviation,  $x_i$  is the  $i$ -th observation,  $\mu$  is the sample mean, and  $N$  is the sample size.  $CV$  was classified as low ( $CV < 10\%$ ), medium ( $10\% \leq CV < 20\%$ ), high ( $20\% \leq CV < 30\%$ ), and very high ( $CV \geq 30\%$ ) (Vaz et al., 2017; Mashalaba et al., 2020).

To support the classification of temperature-dominated, moisture-dominated, jointly

设置了格式: 非突出显示

删除了: In this study, the consistency between study-reported MAT/MAP and ERA5-Land-derived MAT/MAP was assessed by sampling ERA5-Land at site coordinates (0.1°) and fitting linear regressions (ERA5 vs. original)

删除了: are

删除了: SMU values were obtained in two ways: directly from the literature or by computation.

删除了: Monthly SMU fluxes (Jan–Dec) were derived as

删除了: -

删除了: -

删除了: -

删除了: -

删除了: -

删除了: -

删除了: Annual values were calculated when field

删除了:  $\text{CH}_4$

删除了:  $^{-2}$

删除了:  $^{-1}$ .

删除了:  $\text{CH}_4$

设置了格式: 上标

设置了格式: 上标

删除了: Monthly soil moisture and soil temperature were a

设置了格式: 字体: (中文) + 中文正文 (等线)

删除了:  $^{-3}$ ;

设置了格式: 上标

设置了格式: 上标

删除了: All the data were organized and archived in

设置了格式: 字体: 倾斜

删除了: We used the coefficient of variation ( $CV$ ) to expres

删除了:

删除了:

删除了:

删除了:

删除了:

删除了:

controlled, and no-clear-dominance systems, we conducted ecosystem-specific regression analyses of SMU as a function of soil temperature and soil moisture using standardized variables (Schielzeth, 2010). For each ecosystem group, we fitted multiple linear regression models with standardized variables, using SMU as the response variable and soil temperature and soil moisture as predictors. The relative importance of temperature and moisture was evaluated based on the significance and absolute magnitude of the standardized regression coefficients. Ecosystems were classified as temperature-dominated when the temperature coefficient was significant and larger than the moisture coefficient, moisture-dominated when the moisture coefficient was significant and larger than the temperature coefficient, jointly controlled when both predictors were significant and showed comparable standardized effects. Ecosystems that did not meet these criteria were classified as having no clear dominant control. These analyses were intended to support descriptive classification of empirical patterns rather than establish causal mechanisms.

Data processing and calculations were performed using MATLAB R2021a (MATLAB code files 1–3). Figures were generated using R version 4.5.0 (R Core Team, 2025; R code files 1–11). The major R packages used for plotting included ggplot2, dplyr, tidyr, readxl, and patchwork.

### 3 Results and discussion

#### 3.1 Overview of the SMUD




Figure 1. Temporal trends in peer-reviewed publications on soil methane uptake (SMU), from the Web of Science and CNKI databases. The green line indicates the globally averaged atmospheric  $\text{CH}_4$  concentration (adapted from NOAA Global Monitoring Laboratory records).

Before the final data screening and filtering procedures, all 3,050 initially compiled records were evaluated using the Q01–Q07 QC system to identify potential issues and improve data

删除了: (Schielzeth, 2010)

设置了格式: 字体: (默认) Times New Roman, 11 磅

设置了格式: 字体: (默认) Times New Roman, 11 磅

删除了:

删除了: Peer-reviewed publications on soil methane uptake over time

删除了:  $\text{CH}_4$

删除了: adapted from [https://gml.noaa.gov/ccgg/trends\\_ch4/](https://gml.noaa.gov/ccgg/trends_ch4/)

transparency. The proportions of records flagged as Q01–Q07 were 0.56% (n = 17), 0.03% (n = 1), 17.44% (n = 532), 2.69% (n = 82), 4.13% (n = 126), 0.23% (n = 7), and 47.70% (n = 1,455), respectively. These QC flags were primarily used to annotate potential data limitations or specific data characteristics prior to subsequent filtering and harmonization procedures. Because individual records may contain multiple QC flags, the reported percentages are not mutually exclusive. The current database includes 2,427 site-level records from 1,155 sites extracted from 920 peer-reviewed publications, spanning publication years from 1986 to 2025 and covering measurement years from April 1983 to November 2023. The majority of publications (approximately 84%) are in English, with the remainder published in Chinese. The dataset comprises 1,425 daily-scale measurements from 681 sites, 2,001 monthly-scale measurements from 973 sites, 1,720 seasonal-scale measurements from 795 sites, and 1,098 annual-scale measurements from 535 sites.

The number of SMU studies has gradually increased since the 1980s, with a marked increase over the past two decades. The expansion of SMU studies may be partly associated with advances in CH<sub>4</sub> measurement techniques. Measurement approaches have evolved from laboratory wet-chemical analyses to gas chromatography, automated chamber systems, laser spectroscopy, and broader atmospheric CH<sub>4</sub> monitoring, enabling observations across multiple temporal and spatial scales (Khalil et al., 1989; Werle et al., 1993; Baldocchi, 2003; Crosson, 2008; Jacob et al., 2016). The increase in SMU publications also appears to coincide with shifts in atmospheric CH<sub>4</sub> over the last 40 years; after a relatively stable period, atmospheric CH<sub>4</sub> concentrations began to rise again around 2007 (Reay et al., 2018). One possible interpretation is that the renewed increase in atmospheric CH<sub>4</sub> may have contributed to increased scientific attention to SMU.

### 3.2 Spatial distribution of data

The database provides broad spatial coverage and encompasses most key regions where SMU studies have been conducted (Fig. 2). The most frequently represented countries include China (n = 761 records, hereafter “n =” was omitted), the United States (320), Germany (139), Japan (131), Canada (121), Finland (90), Australia (80), Brazil (71), Indonesia (52) and the United Kingdom (51). Across continents, Asia contributed 43.0% of all records (1,023), followed by Europe (22.7%, n = 540, hereafter “n =” was omitted), North and Central America (21.0%, 499), South America (5.1%, 122), Oceania (4.1%, 97), Africa (3.5%, 84), and Antarctica (0.7%, 16). These results indicate a strong sampling bias toward observations from the Northern Hemisphere. Records are primarily concentrated in eastern Asia, North America, and Europe. In contrast, large areas of Oceania, Africa, and the polar regions remain sparsely represented (Fig. 2). This uneven geographical distribution reflects the current availability of field observations but may introduce additional uncertainty in global extrapolation and large-scale modeling applications, especially in regions with sparse observational coverage.

删除了:

删除了: 914

设置了格式: 非突出显示

删除了: 1048

删除了: of

设置了格式: 非突出显示

设置了格式: 非突出显示

设置了格式: 非突出显示

设置了格式: 非突出显示

删除了: The number of studies has gradually increased since the 1980s, with a pronounced surge over the past two decades.

删除了: The expansion of SMU studies is closely linked to advances in CH<sub>4</sub> measurement techniques: detection methods have evolved from laboratory wet-chemical analysis to automated chromatography, laser spectroscopy, and further to satellite remote sensing and intelligent monitoring, forming a multi-scale and comprehensive monitoring system (Baldocchi, 2003; Jacob et al., 2016; Werle et al., 1993; Crosson, 2008; Khalil et al., 1989).

删除了: Moreover, the pattern appears to align with shifts in atmospheric CH<sub>4</sub> over the last 40 years: following a relatively steady period, CH<sub>4</sub> levels began to rise again around 2007 (Reay et al., 2018).

删除了: One possible interpretation is that the renewed increase in atmospheric CH<sub>4</sub> has helped draw more attention to SMU....

删除了: has

删除了: have substantially fewer measurements




Figure 2. Geographic distribution of global SMU measurement sites across biomes at monthly, seasonal, and annual scales.

722 The database covers a wide range of climatic zones and ecosystem types across both natural  
 723 and human-modified ecosystems.  
 724 In terms of climatic classification, most records are located d in temperate zones, accounting  
 725 for 60.3% (n = 1,446, hereafter “n =” was omitted) of the dataset, followed by boreal zone 18.5%  
 726 (443), tropical zone 15.7% (377), and alpine zone 5.5% (131). Relative to their global land area,  
 727 temperate regions are overrepresented: they account for approximately 31% of the global land  
 728 surface but contribute about 60% of the total records, reflecting a sampling bias toward mid-  
 729 latitude regions. By contrast, although alpine regions occupy only approximately 3% of the  
 730 global land area, they contribute 5.5% of the records in SMUD, likely because this database  
 731 includes a relatively large number of SMU studies from the Qinghai–Tibet Plateau in China, a  
 732 region that was underrepresented in previous global databases. These findings further highlight  
 733 the need to increase future SMU observations in tropical regions, which account for  
 734 approximately 24% of the global land area yet contain only 15.7% of the records. This  
 735 imbalance may weaken model calibration and evaluation in tropical regions, thereby  
 736 contributing to uncertainty in global SMU budget estimates.  
 737 Ecosystem representation also varied substantially across the database. Forests represent the  
 738 largest share of the dataset, contributing 48.1% (n = 1,153, hereafter “n =” was omitted) of  
 739 records. Within the forest category, rainforests account for 1.4% (33). Agricultural lands  
 740 account for 22.4% (538), possibly reflecting increasing research interest in the effects of land  
 741 use and management on SMU. Grasslands account for 16.7% (399). Although spatially  
 742 clustered in certain regions, these records collectively cover a wide range of terrestrial  
 743 ecosystems and climatic zones.  
 744 Despite spatial clustering in certain regions, these records collectively provide broad  
 745 ecosystem and climatic coverage, (Fig. 3). The best represented biomes include temperate  
 746 seasonal forests, which account for 27.6% (n = 662, hereafter “n =” was omitted) of the total  
 747 dataset, followed by agricultural lands (22.4%, 538), temperate grasslands (11.2%, 269), and  
 748 boreal forests (10.4%, 248). Less represented biomes include tropical seasonal forests (9.3%,  
 749 223), alpine regions (5.5%, 131), and tundra (3.7%, 89).  
 750

删除了: natural and anthropogenic biomes

删除了: d

删除了: s

删除了: Considering land area, temperate regions account for ~31% of the global land surface but contribute about 60% of the total records, reflecting a sampling bias toward mid-latitude regions.

删除了: Interestingly, although alpine regions occupy only ~3% of the global land area, they contribute 5.5% of the records in the database, likely due to the inclusion of a relatively large number of SMU studies from the Qinghai–Tibet Plateau in China, which were underrepresented in previous global databases.

删除了: These findings further highlight the need to enhance future investigations into SMU in tropical regions, which remain underrepresented in current datasets.

设置了格式: 字体: (中文) 等线, 11 磅

删除了: The data span a broad range of terrestrial ecosystems, with varying levels of representation.

删除了: Forests dominate the dataset, contributing

删除了: ,

删除了: while rainforests, as a specific forest subtype, represent...

删除了: reflecting the increasing research focus in recent years on the impacts of land use and management on SMU.

删除了: According to the Whittaker biome classification, the global SMU records were categorized into distinct biome types...

删除了: most represented




Figure 3. Distribution of global SMU records (gray markers) within the Whittaker biome framework, which represents global climate space based on mean annual temperature (MAT) and mean annual precipitation (MAP). The inset shows the distribution of additional biome categories (Alpine, Agriculture, Bare, Urban, and Others) that fall outside the Whittaker classification.

删除了: figure

删除了: additional biomes

### 3.3 Observed annual and seasonal fluxes




Figure 4. Comparison of annual and seasonal SMU observations across biomes: Tundra (Tun), Boreal Forest (BF), Temperate Seasonal Forest (TSF), Temperate Rainforest (TRF), Tropical Rainforest (TRRF), Tropical Seasonal Forest (TRSF), Savanna (Sav), Subtropical Desert (SDes), Temperate Grassland (TG), Desert (Des), Woodland (Wood), Shrubland (Shrub), Alpine (Alp), Agriculture (Agri), Bare (Bare), and Urban. In the annual panel, biomes are ordered by the median SMU values. Black lines indicate the median SMU, and the crosses (x) denote the mean for each biome. In violin plots, the width represents the kernel density of observations at a given value. The number of flux values is indicated after each x-axis label. Plot types were selected based on sample size: scatter plots for  $n < 10$ , box plots for  $10 \leq n < 20$ , and violin plots for  $n \geq 20$ .

At the annual scale, forests generally exhibited higher SMU than grasslands and cultivated soils in the database, consistent with previous studies (Fig. 4; Dutaur and Verchot, 2007; Reay et al., 2018; Tate, 2015). Among forest biomes, the high median fluxes observed in tropical rainforests ( $46.5 \mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ ) and temperate rainforests ( $42.3 \mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ ), differ from some earlier estimates (Dutaur and Verchot, 2007). This discrepancy may partly reflect seasonal sampling bias in earlier tropical rainforest studies, many of which focused on the hot and wet season, potentially leading to underestimated annual averages. Moreover, many earlier studies grouped rainforests together with other forest types, potentially masking their relatively high SMU (Li et al., 2025). The high SMU observed in shrublands (Fig. S2) are consistent with results from the Terrestrial Ecosystem Model (TEM) (Zhuang et al., 2013). Alpine regions showed the highest SMU among climatic zones (Fig. S3); however, their mean SMU was slightly lower than the value reported for alpine grasslands (mean  $54.2 \mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ ), likely because the Alpine category in this study includes multiple ecosystem types, such as forests and grasslands (Wang et al., 2025). Savanna, agricultural, and desert soils exhibited relatively low SMU, especially in subtropical deserts, probably due to moisture limitation, land-

删除了:  $\text{CH}_4$

删除了: uptake

删除了: scatter plots for  $<10$ , box plots for  $10-20$ , and violin plots for  $\geq 20$ .

删除了: On an annual scale (Fig. 4), as commonly recognized in previous studies, forests exhibit the highest  $\text{CH}_4$  sink rates compared to grasslands and cultivated soils in the database (Dutaur and Verchot, 2007; Reay et al., 2018; Tate, 2015)

删除了: Among forest biomes, the notably high median flux observed in rainforests ( $46.5 \mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$ )

设置了格式: 上标

设置了格式: 上标

删除了: differs from some earlier estimates (Dutaur and Verchot, 2007),

删除了: likely because many previous studies on tropical rainforests focused on the hot and wet season (i.e., April-October)

删除了: which could have led to an underestimation of annual averages.

删除了: fluxes

删除了: Alpine regions exhibited the highest SMU across climatic zones (Fig. S3); however, the mean SMU was slightly lower than that reported for alpine grasslands

删除了:  $\text{CH}_4$

删除了:  $\text{m}^{-2} \text{h}^{-1}$

删除了: likely because the Alpine category aggregates multiple ecosystem types (e.g., forests and grasslands) (Wang et al., 2025).

management practices, nitrogen inputs, sparse vegetation and harsh environmental conditions, and the suppressive effects of biological soil crusts (Hu et al., 2022; Jia et al., 2018).

The observed global mean annual SMU ranged from approximately 0 to 409.0 ( $\mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ , hereafter the same units) with a median of 21.2 (interquartile range: 9.3–39.2). The large variability in SMU, together with its divergent responses to soil moisture, likely reflects differences in local environmental conditions, including climate (Dutaur and Verchot, 2007), ecosystem type (Wang et al., 2014), and soil properties (Kou et al., 2017). Within our database, the highest  $\text{CH}_4$  consumption rate was recorded in a tropical forest in Hainan, China, whereas the lowest detectable flux was observed in a subtropical desert in Ningxia Hui Autonomous Region, China.

Natural biomes account for the majority of global SMU (Zhuang et al., 2004), with particularly high SMU observed in temperate and tropical regions. In the database, natural biomes showed relatively high SMU, including woodlands (median  $21.0 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ , hereafter the same units), temperate grasslands (24.8), tropical seasonal forests (25.0), alpine regions (25.3), temperate seasonal forests (29.8), shrublands (36.9), temperate rainforests (42.3) and tropical rainforests (46.5) all exhibit relatively high SMU. These patterns are consistent with previous estimates for temperate and tropical regions (Murguia-Flores et al., 2018; Zhou et al., 2024). In contrast, boreal ecosystems, such as boreal forests (11.9), arid regions, including savannas (6.7), subtropical deserts (13.5), and deserts (13.7), and human-impacted areas, such as agricultural lands (11.3) and bare lands (12.1), generally showed lower annual mean SMU. Notably, rainforests show exceptionally high annual mean SMU, exceeding  $40 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ .

SMU exhibits high seasonal variability across biomes (Figs. 4 and S4). Thirteen biomes showed pronounced seasonal variability,  $CV \geq 20\%$ . Thirteen biomes showed pronounced seasonal variability, with CV values  $\geq 20\%$ : tundra (92.3%), temperate rainforests (86.3%), woodlands (59.3%), boreal forests (55.2%), shrublands (48.2%), temperate grasslands (38.0%), bare lands (34.6%), deserts (30.6%), savannas (28.9%), temperate seasonal forests (24.8%), alpine regions (24.2%), agricultural lands (22.4%), and tropical rainforests (20.9%). This variability may reflect intra-annual environmental changes as well as differences in measurement timing and sampling frequency. These findings emphasize the importance of accounting for seasonal variability when quantifying SMU across biomes. In addition, comparisons between seasonal fluxes and annual mean values in certain ecosystems revealed systematic differences, indicating that estimates that do not account for intra-annual variation may overestimate or underestimate annual SMU budgets.

To evaluate the potential influence of excluding  $\text{CH}_4$  emission records, we quantified the number and proportion of negative flux values. Negative values accounted for 10.72% of the annual records. Seasonally, the proportion of negative records ranged from 9.79% in autumn to 12.83% in winter and was also relatively high in summer (12.43%). The influence of this filtering varied among biomes. Agricultural soils had the largest absolute number of excluded annual emission records, which accounted for 17.84% of valid annual observations in this biome. These results indicate that  $\text{CH}_4$  emissions occurred in a subset of observations. Therefore, excluding these records may influence estimates of mean  $\text{CH}_4$  uptake and seasonal variability, particularly during periods when soils temporarily shift from  $\text{CH}_4$  sinks to  $\text{CH}_4$  sources.

删除了: Savanna, agricultural, and desert soils exhibit relatively low  $\text{CH}_4$  sink rates, especially in subtropical deserts, likely due to moisture limitation, management practices, nitrogen inputs, barren environmental conditions, and suppressive effects of biological soil crusts (Hu et al., 2022; Jia et al., 2018).

删除了: The observed global annual  $\text{CH}_4$  uptake rates ranged from ~0

删除了: .

删除了:  $\text{CH}_4$

删除了:  $\text{m}^{-2} \text{ h}^{-1}$

删除了: The large variability in SMU and its divergent responses to moisture reflect discrepancies in local environments such as climate

删除了: The highest consumption rate was measured in a  
设置了格式: 字体: (中文) + 中文正文 (等线)

删除了:  $\text{CH}_4$

删除了: uptake

删除了: fluxes

删除了: natural biomes such as woodlands

删除了:  $\text{CH}_4$

删除了:  $\text{m}^{-2} \text{ h}^{-1}$

删除了:  $\text{CH}_4$  uptake rates

删除了: result

删除了: In contrast, boreal ecosystems such as boreal forests

删除了:  $\text{CH}_4$

删除了: uptake rates

删除了:  $\text{CH}_4$

删除了:  $\text{m}^{-2} \text{ h}^{-1}$

删除了: Thirteen biomes show pronounced seasonal

删除了: ,

删除了: including tundra (92.3%), temperate rainforests

删除了: This emphasizes the critical role of seasonal

删除了: In addition, for certain ecosystems, comparisons

设置了格式: 字体: (默认) Times New Roman

设置了格式: 字体: (默认) Times New Roman

3.4 Monthly variations of each biome





Figure 5. Intra-annual variations of SMU, soil moisture (expressed as WFPS, water-filled pore space), and soil temperature (T) across biomes in the Northern Hemisphere. The black lines indicate the median SMU for each biome. The count of flux values in each month is labeled beneath the x-axis ticks. The biomes were grouped into four categories: (A) temperature-dominated, (B) moisture-dominated, (C) jointly controlled, (D) no clear dominant control, based on the covariation of SMU with soil moisture and soil temperature.

删除了: 分节符(下一页)

带格式的: 居中

带格式的: 正文, 段落间距段前: 0 磅, 段后: 0 磅, 行距: 单倍行距



删除了:

删除了: CH<sub>4</sub>

删除了: uptake

设置了格式: 非突出显示

删除了: five

设置了格式: 非突出显示

删除了: moisture and temperature

删除了: promot

删除了: moisture and temperature jointly constrained, and (E)

删除了: no obvious pattern

设置了格式: 非突出显示

设置了格式: 非突出显示

删除了: based on the interactions between moisture and temperature

删除了:

设置了格式: 非突出显示

At the global scale, monthly SMU anomalies showed weak temporal covariation with soil moisture and soil temperature when data were pooled across all sites (Fig. S5). In contrast, ecosystem-specific mean monthly SMU patterns showed stronger covariation with soil moisture and soil temperature (Figs. 5 and S5). The large variability in SMU and its divergent sensitivities to soil moisture and temperature likely reflect heterogeneity in local environmental conditions. In general, low soil moisture can inhibit CH<sub>4</sub> oxidation by suppressing microbial metabolism (Yue et al., 2022), whereas high soil moisture can constrain SMU by reducing CH<sub>4</sub> diffusion into soil pores (Shukla et al., 2013). Across ecosystems, SMU showed either positive or negative sensitivity to soil moisture, depending on ecosystem type and background moisture conditions. Soil temperature generally followed air-temperature seasonality, peaking in warm seasons and reaching minima in cold seasons. The combined effects of soil temperature and soil moisture may therefore lead to distinct seasonal patterns and magnitudes of SMU across ecosystems.

Based on the covariation between SMU dynamics and soil hydrothermal variables, biomes were classified into four groups (Fig. 5). In temperature-dominated (e.g., Temperate Grassland; Fig. 5A), where SMU generally increased from winter to summer, broadly following the seasonal increase in soil temperature, whereas the seasonal pattern of soil moisture was less consistently aligned with SMU. In moisture-dominated (e.g., Boreal Forest; Fig. 5B), where seasonal variation in SMU corresponded more closely to changes in soil moisture, suggesting that moisture availability and associated diffusion constraints may be more important controls on SMU. In jointly controlled (e.g., Tropical Rainforest; Fig. 5C), characterized by both soil temperature and soil moisture showing significant associations with SMU, which indicates that seasonal SMU in tropical rainforest cannot be attributed primarily to a single driver; rather, temperature and moisture likely act together to shape observed SMU variability. In contrast, some ecosystems were classified as having no clear dominant control (e.g., Desert; Fig. 5D), because neither soil temperature nor soil moisture showed a clear dominant association with SMU. These results should be interpreted as a descriptive classification of empirical patterns rather than as evidence of direct causal mechanisms. The regression analyses identify whether SMU covaries more strongly with soil temperature, soil moisture, or both within each ecosystem, but they do not exclude the influence of other environmental and biogeochemical factors such as soil texture, pH, nitrogen availability, vegetation type, substrate limitation, or microbial community structure. In the Southern Hemisphere, no consistent SMU seasonality was evident across most biomes (Fig. S7), which likely reflects sparse observational coverage, uneven monthly sampling, and the smaller land area represented in the database. Therefore, the biome-level classifications are most robust for ecosystems with adequate sample size and clear seasonal coverage, while patterns in data-limited regions should be interpreted with caution.

Monthly SMU exhibited pronounced heterogeneity across both temporal and spatial scales within and among sites, ranging from approximately 0 to 792.9 μg CH<sub>4</sub>-C m<sup>-2</sup> h<sup>-1</sup> (see Supplementary file “Studies and Fluxes”). The variability of monthly mean soil CH<sub>4</sub> fluxes in each ecosystem was quantified using the CV (see Methods, Fig. S8). In the Northern Hemisphere, CVs for 16 biomes exceeded 10%, indicating moderate to high intra-annual variability and pronounced differences in seasonal SMU amplitude among biomes. The highest median monthly SMU occurred in August in shrublands, whereas the lowest occurred in November in subtropical deserts.

删除了: At the global scale, monthly SMU anomalies show weak temporal covariation with soil moisture and soil

删除了: In contrast, ecosystem-specific mean monthly SMU

删除了: The large variability in SMU and its divergent

删除了: In general, low soil moisture inhibits CH<sub>4</sub> oxidation

删除了: exhibits

删除了: Soil temperature largely follows air-temperature

删除了: The combined effects of soil temperature and

删除了: Based on the interactive effects of soil moisture and

删除了: five

删除了: ):

删除了:

删除了: and

删除了: Tropical Rainforest

删除了: where the monthly variability of SMU aligns more

删除了: ;

删除了: promot

删除了: Boreal Forest

设置了格式: 字体: (默认) Times New Roman, 11 磅

设置了格式: 字体: (默认) Times New Roman, 11 磅

删除了: clear seasonal cycles in both drivers and a

删除了: jointly constrained (e.g., Tropical Seasonal Forest

设置了格式: 字体: (默认) Times New Roman, 11 磅

设置了格式: 字体: (默认) Times New Roman, 11 磅

删除了: and no obvious pattern (e.g., Savanna; Fig. 5E),

设置了格式: 字体: (默认) Times New Roman, 11 磅

设置了格式: 字体: (默认) Times New Roman, 11 磅

删除了: In the Southern Hemisphere, soil temperature

删除了: exhibits

带格式的: 缩进: 首行缩进: 1 字符

删除了: ~

删除了: CH<sub>4</sub>

删除了: CH<sub>4</sub>

删除了: exceed

删除了: The largest median monthly SMU occurred in

## 4 Future development and access to SMUD

### 4.1 Limitations of the current SMUD

The current literature search may not have captured all studies related to SMU, particularly those conducted in environments where SMU is intermittent or temporally variable, such as riparian zones, wetlands, and seasonally wet agricultural regions. Measurements reported in publications written in languages other than English or Chinese were not included in the database. Furthermore, the existing data structure could be further streamlined and automated to facilitate data access, reuse, and analysis.

The original SMUD database retains both positive CH<sub>4</sub> fluxes, representing uptake, and negative CH<sub>4</sub> fluxes, representing emissions. For the analyses presented in this manuscript, emission fluxes were excluded to focus specifically on SMU patterns. We note that this filtering affects the representation of net soil-atmosphere CH<sub>4</sub> exchange; therefore, users should interpret the analyses presented here primarily as reflecting SMU capacity rather than comprehensive net CH<sub>4</sub> exchange, likely overestimating the total SMU. In addition, although soil temperature and soil moisture are key drivers of SMU and are included in the current version of SMUD, other environmental factors, such as soil pH, C/N ratio, clay content, and other soil texture- or nutrient-related properties, are also known to influence CH<sub>4</sub> oxidation. These factors may regulate methanotrophic activity, gas diffusion, and substrate availability, and their effects on SMU may vary with spatial scale, biome type, and seasonal conditions. Therefore, the current version of SMUD is better suited for analyses of SMU patterns and their relationships with climate, seasonality, and soil hydrothermal conditions, but is less suitable for independently resolving the effects of intrinsic soil physicochemical controls. Future applications could address this limitation by linking SMUD with external soil datasets, thereby supporting more comprehensive assessments of climatic, hydrothermal, and edaphic controls on soil CH<sub>4</sub> uptake.

As new SMU studies continue to be published, additional data are continually emerging. We intend to maintain and periodically update a dynamic version of SMUD. Over time, we envision these updates evolving into a collaborative effort involving the broader research community, ultimately transforming SMUD into a shared resource that can be integrated with other related initiatives.

### 4.2 Future development and applications

SMUD has broad applications across multiple disciplines, including biogeochemistry, ecology, and climate science. A global SMU database provides harmonized measurements of CH<sub>4</sub> fluxes and associated environmental drivers, together with standardized soil parameters, such as soil moisture and soil temperature, thereby supporting more robust bottom-up estimates of SMU. The database can also help process-based models represent SMU more mechanistically by providing observational constraints on CH<sub>4</sub> oxidation and its environmental controls, thereby reducing uncertainty in global SMU estimates, helping to reconcile discrepancies with top-down atmospheric inversions, and improving predictions of CH<sub>4</sub>–

删除了: Weaknesses

删除了: The current literature search was insufficient to capture all studies related to SMU, particularly those conducted in environments with intermittent CH<sub>4</sub> uptake, such as riparian zones, wetlands, and seasonally wet agricultural regions

删除了: Measurements reported in publications written in languages other than English or Chinese are not included in the database.

删除了: Furthermore, the existing data structure could be streamlined and automated to facilitate easier access and analysis.

删除了:

删除了: s

删除了: A global SMU database provides harmonized measurements of CH<sub>4</sub> fluxes and environmental drivers and incorporates standardized soil parameters (e.g., soil moisture and soil temperature), thereby supporting more accurate bottom-up estimates of the soil CH<sub>4</sub> sink.

climate feedbacks under global change (Luo et al., 2013; Murguia-Flores et al., 2018; Ni and Groffman, 2018; Saunois et al., 2025; Smith et al., 2000). By unifying flux measurements across biomes, including tundra, boreal forests, savannas, and agricultural areas, the database helps reveal the spatiotemporal patterns of SMU and supports assessments of SMU responses to climate, vegetation, and land use (Dutaur and Verchot, 2007). Integrating point-based SMU observations with Earth observation products enables the upscaling of SMU estimates from local sites to regional and global scales (Oertel et al., 2016; Tian et al., 2016). The database further provides essential ground-truth observations for calibrating and validating satellite-based and process-based estimates of global SMU.

In Earth system science, shared databases supported by modern computational tools can expand the user community, strengthen analytical capacity, improve reproducibility, and enhance public trust. Moreover, version-control technologies developed within the open-source community enable “living” databases that can be continually updated, tracked, and improved by their users (Bond-Lamberty and Thomson, 2010). Thus, our goal is to build a global database of published SMU observations and provide it to researchers both as a citable static archive and as a dynamic, community-driven resource.

#### 4.3 Database access and updates

All data and code used to reproduce the results in this study are available at <https://doi.org/10.5281/zenodo.18299391> (Jiang et al., 2026). The data are licensed under the Creative Commons Attribution 4.0 International license (CC BY 4.0, <https://creativecommons.org/licenses/by/4.0/>), and the accompanying code is licensed under the open-source MIT license (<https://opensource.org/license/mit>). A detailed description of the data-processing workflow is provided in (Figure S9).

#### 5 Conclusions

SMUD provides a comprehensive global dataset of SMU across both temporal and spatial dimensions. The database includes observations at four temporal scales—daily, monthly, seasonal, and annual—and spans a wide range of climatic zones and ecosystem types. Seasonal variation is essential for understanding intra-annual changes in SMU and elucidating its spatiotemporal patterns. The data reveal distinct SMU capacities among biomes. The overlooks of biome diversity within ecosystem types contribute to the uncertainty in SMU estimates due to failing to reliably capture the intra-ecosystem differences of SMU.

SMUD is intended to compile and make accessible the broad body of SMU observations, mainly published over the past three decades. In addition, the database aims to harness open-source software technologies to foster a dynamic, collaborative, and continually improving data platform for researchers and other users in the biological, environmental, and Earth sciences.

#### Acknowledgements

This research was financially supported by the National Natural Science Foundation of China (grant nos. 42477367 and 42221001). We sincerely thank Professor Ben Bond-Lamberty from

删除了: This database also enables process-based models to represent SMU more mechanistically, reducing uncertainty in global SMU estimates, helping reconcile discrepancies with top-down atmospheric inversions, and improving predictions of CH<sub>4</sub>-climate feedbacks under global change (Luo et al., 2013; Murguia-Flores et al., 2018; Ni and Groffman, 2018; Saunois et al., 2025; Smith et al., 2000).

删除了: By unifying flux measurements across biomes (e.g., tundra, boreal forests, savanna, and agricultural areas), the database enables to reveal the spatiotemporal patterns of SMU and to examine the response of SMU to climate, vegetation, and land use (Dutaur and Verchot, 2007).

删除了: Integrating point-based SMU data with Earth observation products enables upscaling of CH<sub>4</sub> uptake estimates from local sites to regional and global scales (Oertel et al., 2016; Tian et al., 2016).

删除了: The database further offers essential ground truth for calibrating and validating satellite-based and process-based estimates of global SMU.

删除了: In Earth system science, shared databases supported by modern computational tools can expand the user community, strengthen analytical capacity, and enhance public trust.

删除了: Moreover, version-control technologies from the open-source community enable “living” databases that can be continually updated and improved by their users (Bond-

删除了: Thus, our goal is to build a global database of published SMU data and provide it to researchers both as a

设置了格式: 默认段落字体, 字体: (默认) + 西文正文 (等线), (中文) + 中文正文 (等线), 五号

设置了格式: 字体: (中文) + 中文正文 (等线)

删除了: .

删除了: data

删除了: Seasonal variation is shown to be essential for understanding intra-annual changes in SMU and elucidating

删除了: across

删除了: studies

删除了: aspires to

删除了: This research was financially supported by the National Natural Science Foundation of China [42477367,

Pacific Northwest National Laboratory for his valuable comments and careful editing of the manuscript. We also thank all SMUD contributors for their efforts in collecting and compiling the database. ChatGPT ~~was~~ used only for English language polishing (grammar, spelling, and clarity) and for assistance with code editing and debugging.

删除了: were

## Author contributions

Z.Y. and J.W.J. conceived this study; Z.Y., J.S.J and J.W.J. designed the primary analysis; J.W.J. built the database, developed the model ~~codes~~, and wrote the first draft. All authors contributed to data interpretation and preparation of manuscript text.

删除了: code

## References

- Aronson, E. and Helliker, B.: Methane flux in non-wetland soils in response to nitrogen addition: a meta-analysis, *Ecology*, 91, 3242-3251, 2010.
- Baldocchi, D. D.: Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future, *Global Change Biol.*, 9, 479-492, 2003.
- Bond-Lamberty, B. and Thomson, A.: A global database of soil respiration data, *Biogeosciences*, 7, 1915-1926, 2010.
- Bridgman, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales, *Global Change Biol.*, 19, 1325-1346, 2013.
- Chai, L. L., Hernandez-Ramirez, G., Hik, D. S., Barrio, I. C., Frost, C. M., Soto, C. C., and Esquivel-Hernandez, G.: A methane sink in the Central American high elevation paramo: Topographic, soil moisture and vegetation effects, *Geoderma*, 362, ARTN 114092 10.1016/j.geoderma.2019.114092, 2020.
- Crosson, E.: A cavity ring-down analyzer for measuring atmospheric levels of methane, carbon dioxide, and water vapor, *Applied Physics B*, 92, 403-408, 2008.
- De Bernardi, M., Priano, M. E., Fernández, M. E., Gyenge, J., and Juliarena, M. P.: Impact of land use change on soil methane fluxes and diffusivity in Pampean plains, Argentina, *Agric., Ecosyst. Environ.*, 329, 107866, 2022.
- Dutaur, L. and Verchot, L. V.: A global inventory of the soil ~~CH<sub>4</sub>~~ sink, *Global Biogeochem. Cycles*, 21, 10.1029/2006gb002734, 2007.
- Gauci, V., Pangala, S. R., Shenkin, A., Barba, J., Bastviken, D., Figueiredo, V., Gomez, C., Enrich-Prast, A., Sayer, E., and Stauffer, T.: Global atmospheric methane uptake by upland tree woody surfaces, *Nature*, 631, 796-800, 2024.
- Grosso, S. D., Parton, W., Mosier, A., Ojima, D., Potter, C., Borken, W., Brumme, R., Butterbach-Bahl, K., Crill, P., and Dobbie, K.: General ~~CH<sub>4</sub>~~ oxidation model and comparisons of ~~CH<sub>4</sub>~~ oxidation in natural and managed systems, *Global Biogeochem. Cycles*, 14, 999-1019, 2000.
- Hu, Y., Xu, B., Wang, Y., He, Z., Zhang, P., and Wang, G.: Reference for different sensitivities of greenhouse gases effluxes to warming climate among types of desert biological soil crust, *Sci. Total Environ.*, 830, 154805, 2022.
- Jacob, D. J., Turner, A. J., Maasakkers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever,

删除了: CH<sub>4</sub>

删除了: CH<sub>4</sub>

删除了: CH<sub>4</sub>

1287 J., and Frankenberg, C.: Satellite observations of atmospheric methane and their value for quantifying  
1288 methane emissions, *Atmos. Chem. Phys.*, 16, 14371-14396, 2016.

1289 Jia, R., Teng, J., Chen, M., Zhao, Y., and Gao, Y.: The differential effects of sand burial on CO<sub>2</sub>, CH<sub>4</sub>,  
1290 and N<sub>2</sub>O fluxes from desert biocrust-covered soils in the Tengger Desert, China, *Catena*, 160, 252-260,  
1291 2018.

1292 Jiang, J., Yan, Z., and Jian, J.: Global Soil Methane Uptake Database (SMUD) v1.0.0, Zenodo [data set],  
1293 <https://doi.org/10.5281/zenodo.18299391>, 2026.

1294 Jiang, J., Yan, Z., Jian, J., Peng, S., Tian, H., Morris, K. A., Ellam, R. M., Christiansen, J. R., Chen, H.,  
1295 and Dong, J.: Global Soil Methane Uptake Estimated by Scaling Up Local Measurements, *Global*  
1296 *Change Biol.*, 31, e70194, 2025.

1297 Khalil, M. A. K., Rasmussen, R., and Shearer, M.: Trends of atmospheric methane during the 1960s and  
1298 1970s, *Journal of Geophysical Research: Atmospheres*, 94, 18279-18288, 1989.

1299 Kou, Y., Li, J., Wang, Y., Li, C., Tu, B., Yao, M., and Li, X.: Scale-dependent key drivers controlling  
1300 methane oxidation potential in Chinese grassland soils, *Soil Biol. Biochem.*, 111, 104-114, 2017.

1301 Li, Z., Shang, Y., Wang, C., Wang, J., Liu, X., Li, F., Chen, G., Cheng, H., Zou, J., and Liu, S.: Generally  
1302 Reduced Sink Capacity of Upland Soils for Atmospheric Methane Over the Past Three Decades (1993–  
1303 2022), *Global Change Biol.*, 31, e70248, 2025.

1304 Luo, G., Kiese, R., Wolf, B., and Butterbach-Bahl, K.: Effects of soil temperature and moisture on  
1305 methane uptake and nitrous oxide emissions across three different ecosystem types, *Biogeosciences*, 10,  
1306 3205-3219, 2013.

1307 Luo, G. J., Brüggemann, N., Wolf, B., Gasche, R., Grote, R., and Butterbach-Bahl, K.: Decadal  
1308 variability of soil CO<sub>2</sub>, NO, N<sub>2</sub>O, and CH<sub>4</sub> fluxes at the Högwald Forest, Germany, *Biogeosciences*, 9,  
1309 1741–1763, <https://doi.org/10.5194/bg-9-1741-2012>, 2012.

1310 Mashalaba, L., Galleguillos, M., Seguel, O., and Poblete-Olivares, J.: Predicting spatial variability of  
1311 selected soil properties using digital soil mapping in a rainfed vineyard of central Chile, *Geoderma*  
1312 *Regional*, 22, e00289, 2020.

1313 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y.,  
1314 Goldfarb, L., and Gomis, M.: Climate change 2021: the physical science basis, Contribution of working  
1315 group I to the sixth assessment report of the intergovernmental panel on climate change, 2, 2021.

1316 Murguía-Flores, F., Arndt, S., Ganesan, A. L., Murray-Tortarolo, G., and Hornibrook, E. R. C.: Soil  
1317 Methanotrophy Model (MeMo v1.0): a process-based model to quantify global uptake of atmospheric  
1318 methane by soil, *Geosci. Model Dev.*, 11, 2009-2032, 10.5194/gmd-11-2009-2018, 2018.

1319 Ni, X. and Groffman, P. M.: Declines in methane uptake in forest soils, *Proc. Natl Acad. Sci. USA*, 115,  
1320 8587-8590, 10.1073/pnas.1807377115, 2018.

1321 Oertel, C., Matschullat, J., Zurba, K., Zimmermann, F., and Erasmi, S.: Greenhouse gas emissions from  
1322 soils—A review, *Geochemistry*, 76, 327-352, 10.1016/j.chemer.2016.04.002, 2016.

1323 Ojima, D., Valentine, D., Mosier, A., Parton, W., and Schimel, D.: Effect of land use change on methane  
1324 oxidation in temperate forest and grassland soils, *Chemosphere*, 26, 675-685, 1993.

1325 Poggio, L., De Sousa, L. M., Batjes, N. H., Heuvelink, G. B., Kempen, B., Ribeiro, E., and Rossiter, D.:  
1326 SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty, *Soil*, 7, 217-  
1327 240, 2021.

1328 Reay, D. S., Smith, P., Christensen, T. R., James, R. H., and Clark, H.: Methane and Global  
1329 Environmental Change, *Annual Review of Environment and Resources*, 43, 165-192, 10.1146/annurev-  
1330 environ-102017-030154, 2018.

删除了: CH<sub>4</sub>

删除了: CH<sub>4</sub>

Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguía-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., van der Werf, G. R., Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, *Earth Syst. Sci. Data*, 17, 1873–1958, 10.5194/essd-17-1873-2025, 2025.

Shindell, D., Kuylenstierna, J. C., Vignati, E., Van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S. C., Müller, N., Janssens-Maenhout, G., and Raes, F.: Simultaneously mitigating near-term climate change and improving human health and food security, *Science*, 335, 183–189, 2012.

Schielzeth, H.: [Simple means to improve the interpretability of regression coefficients, \*Methods in Ecology and Evolution\*, 1, 103–113, 2010.](#)

Shukla, P. N., Pandey, K. D., and Mishra, V. K.: Environmental Determinants of Soil Methane Oxidation and Methanotrophs, *Crit. Rev. Environ. Sci. Technol.*, 43, 1945–2011, 10.1080/10643389.2012.672053, 2013.

Smith, K., Dobbie, K., Ball, B., Bakken, L., Sitaula, B., Hansen, S., Brumme, R., Borken, W., Christensen, S., and Priemé, A.: Oxidation of atmospheric methane in Northern European soils, comparison with other ecosystems, and uncertainties in the global terrestrial sink, *Global Change Biol.*, 6, 791–803, 2000.

Song, H., Peng, C., Zhu, Q., Chen, Z., Blanchet, J.-P., Liu, Q., Li, T., Li, P., and Liu, Z.: Quantification and uncertainty of global upland soil methane sinks: Processes, controls, model limitations, and improvements, *Earth-Science Reviews*, 104758, 2024.

Tate, K. R.: Soil methane oxidation and land-use change – from process to mitigation, *Soil Biol. Biochem.*, 80, 260–272, 10.1016/j.soilbio.2014.10.010, 2015.

Tian, H., Lu, C., Ciais, P., Michalak, A. M., Canadell, J. G., Saikawa, E., Huntzinger, D. N., Gurney, K. R., Sitch, S., and Zhang, B.: The terrestrial biosphere as a net source of greenhouse gases to the atmosphere, *Nature*, 531, 225–228, 2016.

Vaz, M. A. B., Pacheco, P. S., Seidel, E. J., and Ansuji, A. P.: Classification of the coefficient of variation to variables in beef cattle experiments, *Cienc. Rural*, 47, e20160946, 2017.

Voigt, C., Virkkala, A. M., Hould Gosselin, G., Bennett, K. A., Black, T. A., Detto, M., Chevrier-Dion, C., Guggenberger, G., Hashmi, W., Kohl, L., Kou, D., Marquis, C., Marsh, P., Marushchak, M. E., Nesic, Z., Nykanen, H., Saarela, T., Sauheitl, L., Walker, B., Weiss, N., Wilcox, E. J., and Sonnentag, O.: Arctic soil methane sink increases with drier conditions and higher ecosystem respiration, *Nat Clim Chang*, 13, 1095–1104, 10.1038/s41558-023-01785-3, 2023.

Wang, Chen, H., Zhu, Q., Peng, C., Wu, N., Yang, G., Zhu, D., Tian, J., Tian, L., Kang, X., He, Y., Gao, Y., and Zhao, X.: Soil methane uptake by grasslands and forests in China, *Soil Biol. Biochem.*, 74, 70–81, 10.1016/j.soilbio.2014.02.023, 2014.

Wang, Wang, J., Wang, S., D'Imperio, L., Elberling, B., Ambus, P., Zhang, Z., Ito, A., Li, Y., Pan, J., Song, L., Liu, N., Zhang, R., Chen, W., and Niu, S.: Soil Moisture Threshold of Methane Uptake in Alpine Grassland Ecosystems, *Glob Chang Biol*, 31, e70062, 10.1111/gcb.70062, 2025.

Werle, P., Mücke, R., and Slemr, F.: The limits of signal averaging in atmospheric trace-gas monitoring by tunable diode-laser absorption spectroscopy (TDLAS), *Applied Physics B*, 57, 131–139, 1993.

1377 Yue, P., Zuo, X., Li, K., Li, X., Wang, S., and Misselbrook, T.: Precipitation changes regulate the annual  
1378 methane uptake in a temperate desert steppe, *Sci. Total Environ.*, 804, 150172, 2022.  
1379 Zhou, X., Xiao, W., Cheng, L., Smaill, S. J., and Peng, S.: Unveiling the impact of soil methane sink on  
1380 atmospheric methane concentrations in 2020, *Global Change Biol.*, 30, e17381, 2024.  
1381 Zhuang, Q., Melillo, J. M., Kicklighter, D. W., Prinn, R. G., McGuire, A. D., Steudler, P. A., Felzer, B.  
1382 S., and Hu, S.: Methane fluxes between terrestrial ecosystems and the atmosphere at northern high  
1383 latitudes during the past century: A retrospective analysis with a process-based biogeochemistry model,  
1384 *Global Biogeochem. Cycles*, 18, 2004.  
1385 Zhuang, Q., Chen, M., Xu, K., Tang, J., Saikawa, E., Lu, Y., Melillo, J. M., Prinn, R. G., and McGuire,  
1386 A. D.: Response of global soil consumption of atmospheric methane to changes in atmospheric climate  
1387 and nitrogen deposition, *Global Biogeochem. Cycles*, 27, 650-663, 10.1002/gbc.20057, 2013.▼

删除了:

Schielzeth, H.: Simple means to improve the interpretability of regression coefficients, *Methods in Ecology and Evolution*, 1, 103-113, 2010.