



Spatial patterns of global land management intensity are influenced by socioeconomic, biophysical and behavioural factors

Ankita Saxena^{1*}, Calum Brown¹, Karina Winkler¹, Almut Arneith^{1,2} and Mark Rounsevell^{1,2,3}

¹ Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Karlsruhe Institute of Technology, Kreuzackbahnstraße 19, 82467, Garmisch-Partenkirchen, Germany

² Institute of Geography and Geo-ecology, Karlsruhe Institute of Technology, Kaiserstraße 12, Building 10.50, 76131, Karlsruhe, Germany

³ School of Geosciences, University of Edinburgh, Drummond Street, Edinburgh, EH8 9XP, UK

Correspondence to: Ankita Saxena (ankita.saxena@kit.edu)

Abstract. Land systems are increasingly influenced not only by land-use change but also by land management intensity. However, there limitations exist in data and in systematic understanding of management intensity and how it is shaped by socioeconomic, biophysical and human behaviour. We develop a global dataset of land management intensity for 2020 at 0.01 ° spatial resolution, distinguishing unmanaged, very extensive, extensive, and intensive management across cropland, pasture, and forest systems. Intensive management occupies about 22 % of global managed land, while extensive and very extensive management dominate (78 %). Intensive cropland management accounts for around 44 % of cropland area, but intensive pasture is limited to 10 % of pasture area, and intensive forest systems to only 5 % of forest area, revealing distinct sectoral contrasts. Management intensity is highly heterogeneous, with intensive cropland concentrated in North America, Europe, and South and East Asia, while extensive management dominates in Africa and Latin America. Five countries account for nearly half of global intensive cropland. Income, market access, population density, and aridity influences cropland management intensity, whereas pastures and forests show more complex relationships. Comparison with global land decision-making types shows spatial consistency (68 %), suggesting that land management intensity is influenced by land-user behaviour.

Keywords: Global land-use and land cover, Land management intensity, human-behaviour

1 Introduction

Human use of land is a major driving force in the functioning of the Earth system, influencing food production, biodiversity, biogeochemical cycles, and climate regulation (IPCC, 2022). Agriculture and forestry together occupy nearly 70 % of the planet's ice-free surface, contribute roughly one quarter of global greenhouse-gas emissions, and account for more than 70 % of freshwater withdrawals (Foley et al., 2011; FAO, 2020). Over recent decades, research on land-use and land-cover change has substantially improved understanding of processes such as agricultural expansion, deforestation, and urban growth (e.g. Winkler et al., 2021, 2025). However, much less attention has been given to land management intensity, despite its major



influence on productivity, resource use and environmental impacts (Kehoe et al., 2015, Matez et al., 2025). Between 2000 and 2020, global cropland area expanded by less than 10 %, while agricultural output increased by more than 45 %, indicating a widespread shift from land expansion toward intensification of production (FAOSTAT, 2022). Similarly, the human appropriation of net primary production (HANPP) has more than doubled since 1900 and now exceeds one quarter of potential terrestrial productivity (Kastner et al., 2022). These dynamics indicate that analysing land-cover and land-use transitions alone is insufficient for a comprehensive understanding of land system change and its implications for humans and the environment (Arneth et al., 2025; Bayer et al., 2023; Zabel et al., 2019).

Land management intensity reflects the level and combination of inputs applied to land, including irrigation, fertilisers, and other management interventions, which together determine resource use, and influence productivity and the environment (Erb et al., 2017; Haberl et al., 2014). Existing global studies have typically focussed on agronomic or climatic outcomes of management intensity without characterising intensity itself (Alexander et al., 2015). Widely used state of the art land-use and land-cover datasets such as HILDA+ (Winkler et al., 2021), HANPP (Kastner et al., 2022), HYDE (Goldewijk et al., 2017), EarthStat (<http://www.earthstat.org>) and Global Pasture Watch (<https://developers.google.com/earth-engine/datasets/publisher/global-pasture-watch>) provide valuable insights into land-use or land-cover change, biomass appropriation, crop production, and specific management components, but lack a harmonised representation of land management intensity across cropland, pasture and forest systems. However, recent years have seen substantial progress in developing spatially-explicit global datasets that captured specific components of land management. These include high-resolution maps of fertiliser application (e.g., Tian et al., 2022; Adalibieke et al., 2023), irrigation extent and intensity in cropland systems (Liu et al., 2021; Nagaraj et al., 2021; Mehta et al., 2024; Kebede et al., 2024), and livestock densities and grazing pressure in pasture systems (e.g., Global Pasture Watch). Notwithstanding their value, these datasets are largely based on sector-specific indicators, making it difficult to assess land management intensity consistently across land systems and world regions.

Beyond management practices themselves, land management intensity is inherently linked to socioeconomic and behavioural processes (Rounsevell and Arneth, 2011). Economic growth, infrastructure, market access, population density and inequality determine the incentives and abilities of land users to intensify or extensify land management (Stehfest et al., 2019; Ma et al., 2024). At the same time, land management decisions are mediated by human behaviours such as values, aspirations, risk perceptions, and local constraints (Lambin and Meyfroidt, 2011; Rounsevell et al., 2021; Vortkamp and Hilker, 2023). Previous studies incorporating human decision-making into global land-use analyses highlighted regional diversity in land-use strategies (Malek et al., 2019; Malek & Verburg, 2020). However, these behavioural and socioeconomic dimensions have not been examined in relation to spatial patterns of land management intensity using spatially-explicit and globally consistent datasets.

Overall, two key knowledge gaps remain. First, there is no harmonised global representation of land management intensity that consistently distinguishes management intensity levels across cropland, pasture and forest systems. Second, the extent to which observed patterns of management intensity align with underlying socioeconomic drivers and land-use decision-



65 making behaviour remain largely unexplored. To address these gaps, we make three key contributions. First, we develop a
harmonised, spatially-explicit global dataset of land management with distinct levels of management intensity across
cropland, pasture, and forest systems. Second, we integrate multiple variables in a unified framework, enabling a consistent
comparison of management intensity across land systems and regions. Third, we link spatial patterns of land management
intensity with socioeconomic and biophysical variables and with land-use decision making types. We then develop new
70 insights into the behavioural and socioeconomic dimensions of land systems which have not previously been analysed using
a globally consistent and spatially-explicit framework.

To analyse global land management intensity and its association with explanatory variables we pose the following research
questions;

1. How is land management intensity spatially distributed across cropland, pasture, and forest systems globally, and how
75 does this distribution vary between regions, income groups, and countries?
2. To what extent are observed patterns of land management intensity associated with socioeconomic drivers such as
economic development, inequality, market access, and population density, and biophysical variables (aridity index and
elevation), and do these relationships differ across land-use sectors?
3. Do spatial patterns of land management intensity align with land-use decision-making types?

80 We address these questions by developing a harmonised, spatially-explicit global dataset of land management intensity for
the year 2020 at 0.01° spatial resolution. The dataset captures cropland, pasture, and forest systems, each classified into
management intensity categories of unmanaged, very extensive, extensive, and intensive, derived from multiple biophysical
and management indicators. Using this dataset, we (i) analyse the global spatial distribution of land management intensity
across regions, income groups and countries; (ii) quantify relationships between management intensity and important
85 socioeconomic drivers including GDP per capita, income inequality (GINI), market access, population density, aridity index
and elevation using multinomial logistic regression; and (iii) assess the correspondence between management intensity
patterns and independently derived global land-use decision-making types (Malek and Verburg, 2020). These analyses
provide an understanding of the drivers and behavioural influences of land management intensity at the global scale and
provide a basis for integrating management intensity into a global land system analysis.

90 **2 Methods**

We develop a global gridded land management intensity dataset at 0.01 ° spatial resolution for 2020 (Saxena et al., 2026).
Agricultural crops, forests, and pasture land are usually managed through species-selection, the application of fertilisers (and
agro-chemicals more broadly) and irrigation, among other inputs. The level of management intensity reflects these inputs,
which influence yields and the production of various commodities. Here we define four levels of management intensity
95 (intensive, extensive, very extensive and unmanaged) in agricultural, pastoral, and forestry systems, and combine the spatial
datasets with urban, agroforestry, photovoltaics, unmanaged (sparse/ no vegetation/natural unmanaged), and water classes. In



100 this study, intensive management refers to a system with high input use and strong human intervention (high fertiliser application rates, irrigation, and active management practices). Extensive management represents practices of moderate input use with a greater reliance on natural processes such as rainfall. Very extensive management refers to a low-input system
105 relying on minimal intervention and natural conditions. An additional ‘unmanaged’ category was used to represent forests, shrublands and pastures with no signs of active human intervention. Furthermore, we categorised cropland based on its primary purpose (e.g., food, feed, or bioenergy), the data were spatially distributed by country using statistics from the Food and Agriculture Organization of the United Nations (2024). We then analysed the spatial distribution of land management intensity globally and by world income groups, and its correlation with socioeconomic and biophysical indicators (see
110 below). We finally assessed the correlation between land management intensities and the land decision-making types from Malek and Verburg (2020). To harmonise different datasets of varying spatial characteristics, we performed all the necessary preprocessing to bring the data to a consistent level that we later integrated. All input layers were aligned to a common grid of 0.01 ° resolution (Fig. 1). The datasets available at coarser resolution (e.g., 5-arcminutes) were resampled using a nearest-neighbour approach which is standard in large-scale studies.

110 The individual datasets and the methods used are discussed in detail in the following sub-sections.

2.1 Data used

The identities and spatial extents of land-use and land-cover classes were drawn from the global HILDA+ dataset (Winkler et al., 2021) at 0.01 ° (Table 1). The cropland areas were refined into specific crop categories by integrating the CROPGRIDS v1.08 dataset (Tang et al., 2024), which provided primary crop data for maize, wheat, soy, rice, and starchy
115 roots at 5-arcminutes resolution for the year 2020. To determine management intensity for these crops we utilised a crop-specific N fertiliser dataset (Adalibieke et al., 2023). Pasture land management intensity was derived from total N fertilizer datasets (Tg N yr-1) (Tian et al., 2022) at 5-arcminutes. The HILDA+ forest management (managed/unmanaged) dataset (Winkler et al., 2021) was combined with a 100 m resolution forest management dataset (Lesiv et al., 2022) to determine
120 forest management intensity. To account for large-scale photovoltaic (PV) deployment worldwide, we integrated a global PV panels dataset at 20 m spatial resolution (Li et al., 2025), ensuring that solar energy installations were spatially represented in the land-use dataset. We utilised some input datasets corresponding to different reference years than 2020, particularly irrigation and forest management data from 2015. These datasets were considered as they provide the most consistent and currently available spatially-explicit information. To harmonise these inputs, we assumed that large-scale spatial patterns of irrigation and forest management remained relatively stable over short periods of time at the global scale. Thus, the final
125 dataset represented an approximation for the year 2020.

Country-wise FAO crop use data were applied to categorise land-use into food-feed-bioenergy uses using random distribution ensuring spatially-explicit classification at the national level. To characterise socioeconomic contexts, we used the global gridded socio-economic dataset of Perkins et al. (2025) at 1 km resolution. Some of the input variables used to construct the land management intensity dataset included socioeconomic and accessibility related variables in their



130 underlying methodologies, which however differed from the variables used in this study in terms of spatial resolution,
 temporal reference and role. The passive variables of the inputs of the management intensity dataset relied largely on
 country-level or coarse-resolution inputs and earlier reference years, whereas the analysis presented here utilised the
 spatially-explicit and harmonised socioeconomic datasets specifically for the year 2020 (Perkins et al., 2025). These
 variables were explicitly used as explanatory factors in a separate regression framework, and were not used as direct inputs
 135 into the construction of the management intensity dataset. We utilised four globally available socioeconomic indicators
 representing complementary dimensions of development and accessibility and two biophysical variables – aridity index and
 elevation. Gross domestic product (GDP) per capita was used as a proxy for financial capital, reflecting economic capacity.
 Population density captures aspects of human capital and demographic pressure on land systems, influencing labour
 availability and land management intensity. Market accessibility represents infrastructural and manufactured capital,
 140 approximating access to markets and services that facilitate intensive land management. Income inequality, measured using
 the GINI coefficient, was included as an indicator of social structure and distributional inequality. GINI may influence
 decisions related to the adoption of land management practices as it is determined by the ability to access to the land, capital,
 and agricultural inputs Together, these variables provided a parsimonious representation of socioeconomic conditions. In
 addition, we included aridity index and elevation to account for important biophysical constraints on land management. The
 145 aridity index represented long-term moisture availability, where lower values indicate more arid conditions and higher values
 indicate more humid environments. This is important for understanding the feasibility of intensive and irrigated land
 management. Elevation was included as a proxy for terrain-related constraints such as climate limitations, which influence
 the intensity of land management practices. Finally, we compared the land management intensity dataset with a land-use
 decision-making dataset (Malek and Verburg 2020) to assess the correlation with the management practices and land-use
 150 decision-making types. More details on the datasets used are provided in Table 1.

Table 1: Salient details on the data used in the creation of land management intensity dataset.

Data	Categories	Spatial resolution	Year	source
Land-use and land cover	<ul style="list-style-type: none"> – urban, – annual crops, – tree crops, – agroforestry, – pasture/rangeland, – forest (unknown/other), – forest (evergreen, – needle leaf), – forest (evergreen, broad leaf), – forest (deciduous, needle leaf), – forest (deciduous, broad leaf), – forest (mixed), – unmanaged grass/shrubland, – sparse/no vegetation, 	0.01 °	2020	HILDA+ (Winkler et al., 2020, 2021)



	– water			
Crops	– wheat, – maize, – rice, – soy, – starchy-roots	5-arcminutes	2020	CROPGRIDS v1.08 (Tang et al., 2024)
Irrigation	– Irrigation/non-irrigation	5-arcminutes	2015	Mehta et al., 2024
N-fertiliser by crops	– N-fertilizer rates in crops	5-arcminutes	2020	Adalibieke et al., 2023
N-fertiliser in pasture	– N-fertilizer in pasture <ul style="list-style-type: none"> • No3 fertilizer • NH4 fertilizer • N manure application • N manure deposition 	5-arcminutes	2020	Tian et al., 2022
Forest managed/unmanaged	– Forest managed/unmanaged map	0.01 °	2020	HILDA+ (Winkler et al., 2021)
Forest management	– Forest management map	100 m	2015	Lesiv et al., 2022
Ground-mounted PV areas	– PV areas	20 m	2020	Li et al., 2025
Socio-economic dataset	– GDP/capita – Gini index – Market access – Working age population	1 km	2020	Perkins et al., 2025
Biophysical dataset	– Aridity index – Digital Elevation Model (DEM)	30 arc-seconds 1 arc-second	1970-2000	Zomer et al., 2022 ASTER GDEM
Land use decision making	– Professional intensifier – Professional commercialist – Market-oriented smallholder – Subsistence-oriented smallholder – Eco-agriculturalist – The survivalist	10 km	2020	Malek and Verburg 2020
Crop use distribution	FAO statistics for crops used for food-feed-bioenergy	Country-wise	2020	Our World in Data

2.2 Mapping crop types and management intensity onto agriculture land

To map the primary crop types – wheat, maize, rice, oil-N-fixing crops (soy), and starchy-roots (potatoes and tubers) – we used the overall cropland extent from the HILDA+ dataset and then integrated crop specific datasets from CROPGRIDS v1.08 (Tang et al., 2024) to spatially allocate each crop type. The crop-specific harvested area layers were extracted and



harmonised to a common spatial grid, with strict enforcement of identical resolution, extent, and coordinate and projection systems across all crop types. Cropland cells with non-zero harvested area for at least one crop were treated as cropland grid cells. A binary presence-absence map was created for each crop type, indicating whether a specific crop was present in a given grid cell. These binary maps were then aggregated into a single dataset to represent the distribution of all crop types. Missing or negative values were assigned as zero. To address overlapping crop types within a grid cell, we allocated the crop type with the highest harvested area as the dominant crop, ensuring one-cell one-crop representation. To ensure consistency across the dataset, we reprojected the data onto the 0.01 ° resolution by aggregating neighbouring cells and assigning the majority crop type within each aggregation unit.

To assess the management intensity associated with each crop type, we utilised a crop-specific N fertiliser, manuring, and irrigation dataset (Table 1, Adalibieke et al., 2023, Mehta et al., 2024). Fertiliser application rates (kg N/ha per year) were derived for wheat, maize, rice, soybean, and starchy roots. For each crop, we considered various N fertiliser types including Urea, nitrate fertilizers (ammonium nitrate (AN) and calcium ammonium nitrate (CAN)), compound fertilizers (ammonium phosphate (AP), N K compounds (NK), N P K compounds (NPK), other NP (ONP)), AA&NS (anhydrous ammonia (AA) and N solutions (NS)), other synthetic fertilizers (other N straight (ONS) and ammonium sulphate (AS)) and MA&CR (manure (MA) and crop residues (CR)). These fertiliser components were aggregated at the grid cell level to obtain total N application rates for each crop type. Prior to aggregation, values were screened to remove implausible negatives and extreme outliers. The total crop-wise N fertiliser rates (originally at 5-arcmin resolution) were resampled to the 0.01 ° grid using the nearest-neighbour resampling approach, and thus all finer resolution cells within each coarse grid cell inherit the averaged value. To determine the management intensity, we classified the crop-specific N fertiliser rates dataset based on absolute N fertiliser thresholds applied to the dominant crop in each cell. Three intensity classes were defined: very extensive (< 50 kg N ha⁻¹ yr⁻¹), extensive (50-100 kg N ha⁻¹ yr⁻¹), and intensive (≥ 100 kg N ha⁻¹ yr⁻¹) (Kleijn et., 2009, Overmars et al., 2014, Dou et al., 2021). Missing fertiliser values were conservatively treated as zero application and classified as very extensive management. The absolute thresholds were applied consistently across all regions and crop types to ensure comparability of management intensity classes at the global scale. We did not include other chemical inputs such as pesticides because pesticide use is often associated with agricultural intensification and tends to co-vary with fertiliser application rates (Pergner et al., 2023). Including both variables would therefore introduce redundancy. The irrigation dataset at 0.05° resolution (Mehta et al., 2024) was also used to distinguish intensive management with and without irrigation.

2.3 Mapping Pasture Management Intensity

The spatial extent of pasture land was derived from the HILDA+ dataset, ensuring consistency with the broader land-use and land cover classification (though note that the underlying data is lower-resolution for pasture than for other land use types). To create the management intensity dataset, we combined data for synthetic (NO₃⁻, NH₄⁺) fertiliser and manure (organic N), giving total N fertiliser for pastures (Table 1). To convert the total N data into application rates, the fraction of managed pasture within each grid cell was calculated by spatially aggregating the managed pasture mask.




Figure 1: Methodology for land use management dataset creation



190

Grid-cell areas were computed as a function of latitude and total N fertilisers were converted to application rates per unit pasture area ($\text{kg N ha}^{-1} \text{ yr}^{-1}$). Cells without pasture were excluded from the calculation. Pasture management intensity was classified into three intensity categories based on the N application rates: very extensive ($< 50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), extensive ($50\text{-}100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), and intensive ($\geq 100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). The absolute thresholds were applied consistently across all regions and pastures to ensure comparability of management intensity classes at the global scale. Similar to cropland, the pasture fertiliser inputs (5-arcmin) were resampled to map onto a 0.01° grid.

195

2.4 Mapping Forest types and management intensity

The forest categories considered in this study included needle leaf evergreen, broad leaf evergreen, needle leaf deciduous, broad leaf deciduous, and mixed forest. Forest management intensity was determined by combining the HILDA+ forest management dataset with additional information from Lesiv et al.'s (2022) forest management dataset, both of which were projected at 0.01° resolution using a majority method. Forest areas were classified into intensive, extensive and unmanaged based on the level of human intervention and management practices. Planted forests were considered as intensively-managed, naturally regenerating forests with signs of management, such as selective logging, or other forestry interventions, were considered as extensively-managed (Lesiv et al., 2022) and forests with no substantial signs of active human intervention (i.e., no or minimal direct management inputs) were considered as natural forests with minimal or no active management.

200

205

2.5 Photovoltaic (PV) Farm Mapping

Spatial information on PV farms for the year 2020 was aggregated from high resolution raster data (20 m) to the harmonised land management intensity dataset at 0.01° resolution (Li et al., 2025). Individual PV tiles were processed sequentially and reprojected onto the 0.01° grid using average resampling. The PV fraction of underlying high-resolution cells covered by PV installations, resulted in a continuous PV coverage fraction between 0 and 1. We calculated the total PV land area as 8432 km^2 (Li et al., 2025). PV cell allocation was restricted to non-water, non-urban, and non-forest areas and allocated by ranking candidate cells by descending PV fraction and allocating PV cells until the total PV area was achieved using latitude-adjusted cell areas.

210

215

Finally, the land management intensity dataset for year 2020 was produced at 0.01° global grid using a deterministic, rule-based mapping scheme designed to preserve broad land-use and land cover classes from HILDA+ while adding finer details on the management intensity. The land management intensity dataset relies on some low-resolution input datasets (e.g., crop and pasture fertiliser and irrigation), while some of the input datasets were at finer resolutions (e.g., forest management, forest types, etc.). Therefore, the final dataset represented a harmonised integration of varying underlying input data. Water and urban were mapped directly from HILDA+, while cropland, pasture and forest classes were assigned with the management intensity levels derived from respective land management layers. Conservative management intensity was applied where management information was missing or invalid. The cropland cells were assigned to an 'other crops' class unless defined crop types and management intensity levels were available. Managed pastures were assigned as very extensive management unless classified otherwise. Forest cells were assigned to extensive management for the respective forest type, when the management intensity information was not present.

220

225

2.6 Sensitivity analysis of a range of thresholds for the management intensity classification

We performed a systematic sensitivity analysis by varying the thresholds used to define the management intensity classes (baseline case: $50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ and $100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) for a range of possible threshold combinations i.e., by varying the lower threshold between $40 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ and $60 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ and the upper threshold between $80 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ and $120 \text{ kg N ha}^{-1} \text{ yr}^{-1}$.



230 N ha⁻¹ yr⁻¹ resulting in nine combined scenarios. For each combination, the spatial distribution of management intensity was
quantified (i) the global share of cropland under each management class and (ii) spatial agreement with the baseline
thresholds (50 kg N ha⁻¹ yr⁻¹ and 100 kg N ha⁻¹ yr⁻¹). The spatial agreement was computed as the percentage of grid cells for
which assigned management intensity class remained unchanged relative to the baseline classification. The sensitivity
analysis showed that the management intensity classification remained broadly stable across alternative threshold
235 combinations (Fig. A4; Tables S1-S2). For cropland systems, spatial agreement with the baseline classification ranged
between 89.7 % and 96.5 %, while pasture systems showed even higher agreement ranging from 97.7 % to 99 %. Although
changes in thresholds altered the relative shares of very extensive, extensive, and intensive classes, the overall spatial
patterns remained largely consistent.

2.7 Relationship between land management intensity and socioeconomic and biophysical indicators

240 To examine the association between socioeconomic and biophysical conditions and land management intensity, we
employed multinomial logistic regression model at the grid cell level. To evaluate that the socioeconomic variables do not
inculcate multicollinearity, a pairwise correlation coefficient and the variance inflation factor (VIF) were computed for all
the variables and the land-use types. Separate models were developed for cropland, pasture and forest systems, with
management intensity (very extensive, extensive, and intensive) as the dependent variables. Very extensive management was
245 used as the reference category in the model. Independent variables comprised GDP per capita, Gini index (income
inequality), market accessibility and working age population, giving indicators of the primary socioeconomic capital groups
(financial, social, manufactured and human; Perkins et al., 2025). All these datasets were harmonised to the same spatial
resolution of 0.01 ° and for the year 2020 as the land management intensity dataset. These predictors were standardised prior
to the use in the regression model to allow a direct comparison across the variables. The individual regression models were
250 developed to analyse the variations both globally and in the World Bank income groups (lower-middle income, upper-
middle income, high-income and low-income). We also computed the average marginal effects, which quantify the change
in the predicted probability of each management intensity category associated with a unit change in each variable. Marginal
effects were computed for all explanatory variables and management intensity classes across the land systems and income
groups. In addition, to assess where socioeconomic drivers explain land management intensity well, and where they do not,
255 we computed spatial residuals between observed land management intensity and predicted class probabilities derived from
the multinomial logistic regression models. For each land sector and management intensity class, residuals were calculated
as the difference between the observed binary management class membership and the model-predicted probability at each
grid cell. Positive residuals indicate locations where a management intensity class occurs more frequently than predicted by
socioeconomic variables alone, while negative residuals indicate under prediction.

260 We performed a robustness check by using a spatial thinning method to evaluate whether spatial clustering has a role in the
regression relationships. We performed an analysis using progressively coarser resolution samples such as 0.10 °, 0.25 °, and
0.50 ° to compute the regression for an individual set-up. The direction and magnitude of marginal effects were compared
against the original resolution model (0.01 °). The outcome indicated that marginal-effect relationships retained their
direction across progressively coarser spatial samples (Fig. A8–A10; Table A6). In particular, GDP, market access, aridity
265 index, and elevation effects remained largely stable across cropland, pasture, and forest systems, although effect magnitudes
varied under coarser thinning. Mixed responses occurred mainly for selected pasture and forest relationships, indicating that
some associations are more sensitive to spatial aggregation. However, such influences were very limited. Overall, the results
suggest that the main regression patterns are not solely driven by spatial clustering of neighbouring grid cells.



2.8 Relationship between land management intensity and the land decision-making types

270 To assess the consistency between land management intensity and land-use decision-making behaviour, we harmonised
 management intensity classes with decision-making types derived by Malek and Verburg (2020). The mapping was based on
 the conceptual correspondence between decision-making strategies, which reflected actors' objectives, capacities, and
 constraints, and observable land management outcomes. Decision-making types associated with market orientation, capital
 access, and production optimisation including professional intensifiers and professional commercialists were mapped to
 275 intensive or extensive management classes (Table 2). These decision-making types corresponded to active engagement in
 land-use markets and adaptive management strategies, which were inferred as either high-input intensive systems or input-
 efficient extensive systems depending on biophysical and economic conditions. Decision-making types emphasising
 environmental concerns or multifunctionality, such as eco-agriculturalists, were mapped primarily to extensive management,
 reflecting lower input use and greater reliance on ecosystem processes. In contrast, decision-making types constrained by
 280 limited resources or risk tolerance such as subsistence-oriented smallholders and survivalists were mapped to very extensive
 management (Table 2). Several decision-making categories represented co-presence or hybrid strategies, where multiple
 behavioural types coexist within the same spatial unit. For these cases, we applied multiple management intensity classes
 (e.g. intensive, extensive, and very extensive). Decision-making classes with low classification confidence or insufficient
 information were excluded from the comparison. Spatial comparisons were then conducted at the grid-cell level. A match
 285 was defined when the observed land management intensity at a given grid cell fell within the set of possible management
 intensity classes associated with the corresponding decision-making type. This flexible matching approach allows for one-to-
 many relationships between behaviour and management outcomes, reflecting the fact that similar decision-making strategies
 may lead to different management intensities depending on local environmental, and institutional conditions. To evaluate the
 spatial consistency between the land decision-making types and the land management intensity, pixel-wise comparison
 290 metrics were computed including total matched (hit) cells, mismatches, and overall accuracy. Additionally, confusion
 metrics summaries were computed. It is to note that the decision-making dataset is derived using proxy variables such as
 GDP, irrigation and accessibility from earlier datasets and at a coarser spatial resolution. Similar variables were used in this
 study to perform the regression analysis which differed in terms of their role, spatial representation, and time coverage. The
 comparison between the newly developed land management intensity dataset and the decision-making types should therefore
 295 be interpreted as broad consistency between behavioural patterns and management intensity.

Table 2. Mapping of land-use decision making types mapping onto land management intensities classes based on Malek and Verburg (2020)

Decision-making type	Mapped management intensity	Rationale
Professional intensifier	intensive or extensive	high-input intensive systems and input-efficient extensive systems depending on biophysical conditions
Professional commercialist	intensive or extensive	profit-oriented with flexibility in input intensity based on biophysical conditions
Co-presence of commercialist and professional intensifier	intensive or extensive	Mixed professional land-use within the same spatial unit depending upon the market situation and biophysical conditions
Copresence of professional commercialist, professional intensifier and eco-agriculturalist	intensive or extensive or very extensive	Behavioural heterogeneity spanning production-oriented and environmentally oriented
Eco-agriculturalist	extensive	Emphasis on environmental stewardship and reduced input use
Subsistence oriented smallholder	very extensive	Small-scale production with limited input use



Survivalist	very extensive	Minimal intervention and resource constrained
Co- presence of professional intensifier and eco-agriculturalist	intensive or extensive or very extensive	Hybrid decision-making strategies reflecting both production and conservation
Market-Oriented Smallholder	intensive or extensive	Market oriented small-scale production with limited input use
Co-presence of professional commercialist and eco-agriculturalist	intensive or extensive or very extensive	Coexistence of commercial and environmental land-use strategies
Low Confidence / Unclassified	ignored in analysis	

3 Results

3.1 Global distribution of land management intensity


300 The global land management intensity dataset (Fig. 2) revealed spatial heterogeneity in cropland, pasture and forest management intensity at 0.01 ° resolution, enabling a spatially consistent global assessment of land management patterns (figures below are expressed as percentages of the entire global terrestrial surface, unless otherwise indicated) (Saxena et al., 2026). Approximately 79 % of the Earth’s land area was found to be managed to some degree, ranging from very extensive to intensive management across cropland, pasture and forest systems. The remaining 21 % of the total land area is

305 predominantly unmanaged and consists largely of sparsely- and non-vegetated areas, and ice/snow covered areas with minimal human intervention. Intensive management occupies about 22 % of the global managed land area (i.e., cropland, pasture, and forestry), while extensive and very extensive management together accounted for 78 % (Table A3). Despite perceptions of widespread intensive management, most managed land globally remains under extensive or very extensive management (Table A3). Although, management intensity varied substantially across land-use and land-cover sectors.

310 Croplands accounted for around 10.5 % of the global terrestrial surface, yet contained the highest share of intensive management among all land-use and land-cover sectors (Fig. 3). Of global cropland, 44 % was managed intensively comprising around 30 % irrigated management and 14% rainfed management (Fig. A1). Intensive and extensive croplands were concentrated in densely populated and highly industrialised regions, where infrastructure, fertiliser use, and irrigation were widely available (Fig. A2).




Cropland




Very extensive Extensive rainfed Extensive irrigated Intensive rainfed Intensive irrigated

Pasture



Unmanaged pasture Very extensive Extensive Intensive

Forest



Unmanaged forest Extensive Intensive

320 Figure 3: Spatial distribution of land management intensity in cropland, pasture, and forest, categorised into very extensive, extensive, intensive, unmanaged forest, and pasture.



The results revealed a sectoral contrast, where intensive management remained concentrated in cropland systems, while pasture and forest systems remained predominantly extensive or very extensive despite their large spatial extent. At the aggregate level, pasture and forest land dominated managed land areas, and together
325 accounted for more than two-thirds of the terrestrial surface. Pasture land occupied around 36 % of the global terrestrial surface, but was predominantly managed at low intensity, with only 10 % under intensive management, 44 % under extensive management, and 46 % under very extensive or minimal management (Fig. A1 & A5). Forests covered around 31.3 % of the global land surface, of which only around 5 % was managed intensively, 40 % managed extensively and 55 % remained unmanaged (mainly in boreal and tropical regions)
330 (Fig. A1). The mapped global distributions of land management intensity aligned closely with independent reference datasets (Fig. A5, Table A3), with deviations (on an area-basis) generally within ± 3 percentage points of FAO and IPCC estimates, supporting the robustness of the mapped dataset.

3.2 Economic, regional and country-level patterns in land management intensity


Land management intensity revealed distinct and non-uniform relationships with world income groups, world
335 regions, and countries across land-use sectors (Fig. 4–5 & Fig. A1).

3.2.1 Land management in different income groups

Our analysis indicated that economic development does not uniformly align with intensive management, but has relationship that vary with land-use type and regional characteristics. In cropland systems, economic development and intensive management are strongly linked, but more complex effects are found in pasture and
340 forest systems. In all income groups, management intensity varied substantially across land-use sectors (Fig. 4), indicating that the influence of economic development may vary between cropland, pasture, and forest systems. In cropland, intensive management (rainfed and irrigated combined) accounted for nearly half (49 % and 48 %) of cropland area in high-income and upper-middle income countries respectively, compared to smaller shares in lower-middle income (36 %) and low-income countries (28 %). Conversely, very extensive cropland dominated
345 in low- and lower-middle income countries, where it accounted for approximately 55 % and 50 % of cropland, compared to 35 % globally. Pasture systems were predominantly managed at low intensity across all income groups. Globally, unmanaged and very extensive pastures together accounted for approximately 90 % of pasture area, with intensive pasture representing only about 10 % (Fig. 4). Intensive pasture management was most prevalent in upper-middle and lower-middle income countries (17 % and 22 % respectively) and least common
350 in high-income countries (3 %). Forest management revealed a contrasting pattern, with unmanaged forests dominating globally (55 %) and particularly in high- and low-income countries (61 % and 52 % respectively). Intensive forest management was comparatively rare but occurred disproportionately in upper-middle income (7 %) and high-income countries (5 %).

3.2.2 Land management in different world regions

355 Regional patterns indicate that land management intensity may not only be driven by economic development (i.e., income), but also by regional conditions, and institutional contexts, leading to diverse management intensities across the globe (Fig. A1). Intensive cropland management was concentrated in North America (56% of the cropland area) and East Asia & Pacific (55 %), followed by Europe & Central Asia (47 %) and South Asia (41 %).



360

Figure 4: Proportion of cropland, pasture, and forest area under different management intensities globally and by world economic groups. Cropland management categories included very extensive, rainfed extensive, irrigated extensive, rainfed intensive, irrigated intensive. Pasture management included unmanaged, very extensive, extensive and intensive. Forest management included unmanaged forest, extensive and intensive forest.



365 In contrast, very extensive cropland dominated in Latin America & Caribbean (62 %), Sub-Saharan Africa (56
%) and the Middle East & North Africa (53 %), revealing an uneven distribution of management intensity across
regions. Pasture systems were largely unmanaged or very extensively managed across all regions, but showed
greater variability in Latin America & Caribbean (i.e., very extensive – 42 %, intensive – 18 %), and South Asia
(i.e., very extensive – 46 %, intensive – 23 %). Forest management also exhibited regional differences: intensive
370 forest management was most pronounced in East Asia & Pacific (13 %), followed by Europe & Central Asia (6
%) and North America (4 %). Unmanaged forests dominated in Europe & Central Asia (63 %), while extensive
forest management was most prevalent in South Asia (80 %), the Middle East & North Africa (76 %), and Sub-
Saharan Africa (57 %).

3.2.3 Concentration of intensive management at the country level

375 Global patterns of land management intensity were found to be disproportionately driven by a small number of
countries, perhaps indicating a role of national-level policies, and land-use practices in determining management
intensity decisions. The United States, China, India, Canada and Ukraine together accounted for nearly half of
global intensive cropland area (49.8 %), with individual contributions of 13.1 %, 12.4 %, 8.7 %, 7.7 % and 6.7 %
respectively. A similar concentration pattern was observed in pasture systems, where China (21.4 %), Brazil (8.2
380 %), Madagascar (4.4 %), Mozambique (3.3 %), and the United States (2.8 %), together accounted for around 40
% of global intensive pasture area (Fig. 5). Intensive forest management was even more concentrated, with China
(26.5 %), the United States (13.4 %), Russia (8.6 %), Sweden (6.6 %), and Japan (5 %) together representing
around 60 % of intensively managed forest worldwide (Fig. 5).



3.3 Socioeconomic and biophysical drivers of land management intensity

Multinomial logistic regression models were used to quantify the relationship of socioeconomic (economic development i.e., GDP per capita, inequality i.e., GINI, infrastructure accessibility i.e., market access, and population density) and biophysical variables (elevation and aridity index – higher values are more humid and lower values more arid) with the probability of different land management intensity classes across cropland, pasture and forest systems. A pairwise correlation and the variance inflation factor (VIF) for all the variables and land-use types were computed to check for multicollinearity. A weak to moderate correlation (-0.09 to +0.37) and low VIF (+1.1 to +3.0) values were found to indicate the absence of multicollinearity (Fig. A6). Non-linear relationships were identified between socioeconomic variables and land management intensity, with differences in direction and magnitude across land-use and land-cover classes and world income groups (Table A4, Fig. A7–A9). Marginal effects were computed to quantify the changes in predicted class probabilities associated with a one-standard deviation increase in each socioeconomic variable (Fig. 6).

3.3.1 Relationship between socioeconomic and biophysical variables, and land management intensity

Socioeconomic and biophysical variables were correlated with land management intensity with the direction and magnitude of these relationships varying across land-use sectors and income groups (Fig. 6, Fig. A7-S9). Cropland management intensity showed a positive association with economic development and market connectivity (Fig. A7 & Fig. 6). Higher GDP per capita and improved market access were correlated with higher probabilities of intensive (with irrigated) cropland management, while extensive and very extensive cropland systems declined. For example, a one standard deviation increase in GDP was associated with increases of up to 17 percentage points in the probability of intensive cropland management. Population density was also positively associated with intensive cropland management, especially in upper- and lower-middle income regions. In contrast, higher income inequality (GINI) and increasing aridity index were correlated with greater probabilities of extensive or very extensive cropland management. Elevation showed more heterogeneous relationships across regions (Fig. A7). The details related to marginal effects are summarised in Table A5.

Pasture management showed associations with market access and climate conditions than with economic development. Increased market accessibility and higher aridity index were correlated with greater probabilities of intensive pasture management across several regions, with aridity index showing a higher association (up to 22 percentage points in some regions) (Fig. A8 & Fig. 6). Population density and inequality showed weaker but positive relationships with intensive pasture management, whereas GDP per capita exhibited relatively limited and region-specific correlations.

Forest management intensity was found the most heterogeneous among the three sectors. Market accessibility was correlated with higher probabilities of intensive and extensive forest management, accompanied by substantial declines in unmanaged forest probability across regions (Fig. A9). In high- and upper-middle-income regions, greater inequality was associated with increased probabilities of extensive forest management and lower probabilities of unmanaged forest.




Figure 6: Marginal effects of socioeconomic variables on land management intensities. Individual plot shows an average marginal effect of socioeconomic variables on the predicted probability of land management intensity (very extensive, extensive, and intensive) for cropland, pasture and forest across global and world economic group. Bars represent average marginal effects calculated from multinomial logistic regression method with 95% confidence interval. Positive (negative) values indicate an increase (decrease) in the probability of a given management intensity class associated with a one unit change in the corresponding socioeconomic variable.


420



425 Aridity was associated with higher probabilities of unmanaged forest, while GDP per capita showed positive associations with intensive forest management primarily in low-income regions. Population density exhibited mixed relationships, with denser populations associated with more intensive forest management in high-income regions but with greater probabilities of extensive or unmanaged forests in several middle- and low-income regions. Detailed regional marginal effects are reported in Table A5.

430 **3.4 Comparison between land management intensity and land decision-making types**

A comparison between the land management intensity dataset presented in this study and the decision-making types of Malek & Verburg (2020) (Table A3) provided insights into the behavioural dimensions of land management. The decision-making categories represented diverse farmer and land manager behaviours, ranging from survivalist and subsistence smallholders to professional intensifiers and eco-agriculturalists (Malek & Verburg 2020). Fig. 7 shows the global spatial distribution of consistency with one or more plausible management intensity classes and mismatch between land management intensity and land-use decision-making types (more details in Methods). Across all land systems, a total of 319,045 valid grid cells were compared of which, 215,263 grid-cells (68 %) showed spatial consistency with one or more plausible management intensity classes, while 32 % showed divergence. This level of consistency indicated that, despite conceptual differences, land management intensity captures important aspects of underlying land-use behaviour at the global scale. Large regions of consistency were observed in intensively managed cropland systems, including parts of North America, Europe, East Asia, and South Asia, where high-input management aligns with decision-making types characterised by commercialisation and intensive management. Consistency was also evident in regions dominated by very extensive or extensive land systems, such as parts of boreal Eurasia, central Africa, and arid and semi-arid regions where low-input management corresponds to decision-making types characterised by limited intervention. In contrast, spatial mismatches were concentrated in heterogenous landscapes, where land-use decisions and management intensity vary over small spatial extents. These included regions with mixed land-use systems, transitional management regimes, or overlapping decision-making strategies. In such cases, similar decision-making types may correspond to multiple management intensities or management outcomes may diverge from dominant behavioural classifications.



450 **Figure 7: Comparing land management intensities and land-use decision-making types (given by Malek and Verburg (2020)) showing spatial consistency between the decision-making types and management intensities.**

4 Discussion

This study integrates land-use and land-cover thematic information with land management intensity using multiple datasets, providing a harmonised and spatially-explicit representation of global land management at 0.01-degrees. The relationships of management intensity with socioeconomic and biophysical conditions, and with land-decision making types, deepens



455 understanding of land-system management globally, reflecting its varied dependencies and manifestations. The results reveal
that management intensity is unevenly represented across land-use and land-cover categories and across world regions,
indicating that land-use and land-cover datasets alone are insufficient to characterise levels of intensity (Erb et al., 2017;
Kuemmerle et al., 2013; Meyfroidt et al., 2022). Globally, intensive management accounted for around 22 % of land areas
across cropland, pasture and forest systems, while extensive and very extensive together dominated, covering about 33 %
460 and 45 % respectively. Intensive management represented around 44 % of global cropland area, but only about 10 % of
pasture and 5 % of forest area. Despite widespread narratives of intensive management, most land, especially pasture and
forest, remains under extensive to very extensive management (see also Erb et al., 2016; Turner et al., 2022).

Heterogenous management patterns occur even within broadly similar landscapes, showing strong regional differences in
biophysical and socioeconomic conditions. Intensive pasture management is relatively limited and appears to be determined
465 more by biophysical conditions and accessibility than by economic development. Similar relationships were found by
Herrero et al. (2013 and Jayne et al. (2019). Forest management intensity is greater in Northern and Central Europe and parts
of East Asia, while unmanaged forests dominate large areas of the Amazon basin, Central Africa and Northern Eurasia.

A relatively small number of countries were accounted for a large share of globally intensive land management. We found
that the United States, China, India, Canada, and Ukraine together account for nearly half (49.8 %) of intensive cropland
470 globally. Similarly, China, Brazil, Madagascar, Mozambique, and United States account for 40.2 % of intensive pasture,
while China, the United States, Russia, Sweden, and Japan together represent around 60 % of the intensive forest
management. These patterns are not wholly attributable to economics; high-income regions tend to exhibit higher shares of
intensive management in cropland and forest sectors, but extensive and very extensive management remain prevalent,
particularly in pasture. Conversely, some low and lower-middle income regions show pockets of intensive, often irrigated
475 cropland, indicating that management intensity emerges from interactions between biophysical and socioeconomic
conditions rather than from the economy alone.

The relationships between management intensity and socioeconomic and biophysical variables are sector dependent and also
non-linear. Higher GDP and improved market access are generally correlated with increased probabilities of intensive
(especially irrigated) cropland management. This pattern reflects the capital and infrastructure intensive nature of high-input
480 cropping (Foley et al., 2011). Similarly, population density correlates with the management intensity indicating a higher land
pressure, which appears to provide incentives to intensify production (Zabel et al., 2019). In contrast, income inequality
appears to play a comparatively minor and often dampening role suggesting that uneven access to resources can limit
transitions toward high-input management even where aggregate economic conditions are favourable (Ostrom, 2009;
Lambin & Meyfroidt, 2011).

485 Pasture management follows a slightly different pattern compared to cropland. Pasture is dominated by very extensive and
extensive management with intensive systems occupying only a small area across income groups. Socioeconomic factors
show weaker relationships, with market access primarily associated with a transition from very extensive to extensive
management rather a shift towards intensive pasture management. This response reflects that grazing systems are often



490 constrained by biophysical conditions (e.g., cold or aridity), and relatively low returns on capital intensive inputs. Unlike
cropland, pasture management is therefore less directly linked to economic growth and more to market accessibility among
other factors. Pasture (and rangeland) is also a difficult land use class to identify, especially at lower management intensities,
making firm conclusions difficult to draw at global extent (Harrison et al., 2025). Forest management intensity shows a
heterogeneous response to socioeconomic variables. Increased market access is associated with a decline in less-intensively
495 managed forests and a corresponding increase in predominantly extensive forest management, suggesting that accessibility
facilitates forest management. However, this does not relate to widespread intensive forest management, probably because of
constraints from government regulation and environment policies (Rudel et al., 2010). Economic development is therefore
insufficient to explain intensive forest management, and in some contexts higher income coincides with reduced intensive
management due to conservation policies or land-use zoning.

The comparison between management intensity and land-use decision making types reveals conceptual consistency at the
500 global scale (Malek & Verburg, 2020). Approximately two-thirds of valid grid cells show consistency between the two
datasets, despite their different conceptual foundations. The consistency indicates that management intensity aligns with core
aspects of land-use decision-making and the behaviours that underpin it. The regions with greatest consistency have clearly
defined land-use strategies, such as intensively managed cropland systems in major agricultural regions and very extensive
and unmanaged land systems in remote or environmentally constrained areas. Areas of mismatch, in contrast, tend to occur
505 in heterogeneous landscapes with multiple decision-making strategies or where similar strategies lead to different
management outcomes. These mismatches imply either mixed land management practices or environmental constraints that
were not assessed in this study, and which may reduce the influence of behavioural drivers. Linking management intensity
with a behavioral dimension provided a broader understanding of land management decisions which helps to understand
these decisions in addition to economics.

510 **5 Limitations and uncertainties**

The dataset developed here represents a spatially harmonised approximation of global land management conditions around
2020 and may not fully capture all locally relevant management practices. For example, the absolute thresholds used to
classify management intensity may not capture regional differences in agricultural practices. Using this approach does,
however, enable a consistent global comparison of management intensity patterns, essential for assessing patterns and
515 relationships. The dataset integrates multiple sourced inputs with different spatial resolutions, temporal coverage, and
methodological assumptions. Several crop and pasture management indicators originally available at coarse resolutions were
resampled to the 0.01° grid. This approach preserves finer land-use classes, but inevitably propagates underlying
uncertainties associated with the coarse data. These are particularly pertinent to pasture classes, where data are not only of
coarse spatial and temporal resolution but also more uncertain than equivalent datasets for other land cover classes (Harrison
520 et al., 2025). Some datasets were derived from slightly earlier time periods than 2020 and may therefore not fully represent



525 recent changes. Missing fertiliser values were conservatively treated as absence of fertiliser input and therefore assigned to the very extensive management class. This approach avoids artificially inflating intensive management for relatively small number of grid-cells, although it may lead to some underestimation of management intensity where fertiliser information was unavailable. However, this limitation reflects a broader challenge in global land system science, where key processes cannot be directly measured but instead need to be approximated using best-available data.

530 In addition, some of the input datasets used to derive newly developed land management intensity and decision-making types incorporate socioeconomic and accessibility-related variables in their construction. Although these differ in scale, time period, and purpose from the predictors used in this study, some level of partial dependence cannot be fully excluded. The relationships have therefore been interpreted as correlations. Since the mapping of decision-making types onto management intensity was purposefully based on a one-to-many relationship between land-decision-making types and management intensity classes to consider behavioural heterogeneity and adaptive responses to local conditions, a perfect spatial consistency would neither be expected nor conceptually appropriate.

535 These considerations mean that our findings should be interpreted carefully, as broad global indications of patterns rather than specific explanations of them. Going further towards explanatory analysis would require significantly improved data sources describing land system management. Nevertheless, the results of this study show that such advances would likely be highly informative and reveal genuine relationships between biophysical conditions, human behaviour, socio-economics and land management. The dataset developed in this study already provides a robust baseline of current global land management patterns. It can support the initialisation, parameterisation and validation of land-use models, enabling a better representation of spatial heterogeneity in management practices. In addition, the study can be useful in the assessment of policy-oriented analyses for example land-based climate mitigation strategies – assessing trade-offs between ecosystem service provision and environmental impacts.

Appendix A

545





Figure A1: Proportion of cropland, pasture, and forest area under different management intensities by World Bank regions in 2020. Cropland management categories included very extensive, extensive rainfed, extensive irrigated, intensive rainfed, intensive irrigated. Pasture management included unmanaged, very extensive, extensive and intensive pasture. Forest management included unmanaged, extensive and intensive forest. Regions include: EAP – East Asia & Pacific, ECA – Europe & Central Asia, LAC – Latin America & Caribbean, MENA – Middle East & North Africa, NA – North America, SA – South Asia, SSA – Sub-Saharan Africa.

550



Global cropland management intensity



555 **Figure A2: Global cropland management intensity. Cropland management categories included very extensive (Ve), extensive**
rainfed (Erf), extensive irrigated (Eir), intensive rainfed (Irf), intensive irrigated (Iir). Roots include tubers and potatoes, and ‘O’
refers to ‘other crops’.

560


565

570

575



Global land-use and land cover distribution



580

Figure A3: Global Land-use and land cover distribution. The map shows broader land-use and land cover classes.




Figure A4: Sensitivity analysis of cropland and pasture management intensity classifications to different thresholds values. The x-axis represents the lower and upper thresholds used for the management intensity classes, while the y-axis indicates changes in the spatial agreement with respect to the baseline in percentage points.

585

590




Table A1: Sensitivity analysis of cropland management intensity classification for different thresholds. The table includes shares of very extensive, extensive, and intensive land and spatial agreement with the baseline threshold.

Scenario	Very extensive threshold	Intensive threshold	Very extensive managed crops (%)	Extensive managed crops (%)	Intensive managed crops (%)	Agreement with baseline (%)
40_80	40	80	46.62	13.55	39.83	92.5
40_100	40	100	46.62	17.58	35.8	96.53
40_120	40	120	46.62	24.07	29.31	90.04
50_80	50	80	50.09	10.08	39.83	95.97
50_100	50	100	50.09	14.11	35.8	100
50_120	50	120	50.09	20.6	29.31	93.51
60_80	60	80	53.91	6.26	39.83	92.15
60_100	60	100	53.91	10.29	35.8	96.18
60_120	60	120	53.91	16.78	29.31	89.68

595 **Table A2: Sensitivity analysis of pasture management intensity classification for different thresholds. The table includes shares of very extensive, extensive, and intensive land and spatial agreement with the baseline threshold.**

Scenario	Very extensive threshold	Intensive threshold	Very extensive managed pasture (%)	Extensive managed pasture (%)	Intensive managed pasture (%)	Agreement with baseline (% managed pasture)
40_80	40	80	92.12	3.81	4.07	97.72
40_100	40	100	92.12	4.78	3.10	98.68
40_120	40	120	92.12	5.45	2.43	98.01
50_80	50	80	93.44	2.49	4.07	99.04
50_100	50	100	93.44	3.46	3.10	100.00
50_120	50	120	93.44	4.13	2.43	99.33
60_80	60	80	94.47	1.46	4.07	98.01
60_100	60	100	94.47	2.43	3.10	98.97
60_120	60	120	94.47	3.10	2.43	98.30



600 **Figure A5: Land-use and land cover distribution and a comparison of modelled and reference global land-use shares. Land-use distribution by (b) groups, (c) types, and (d) management intensities. Land-use groups are the broader land-use and cover sectors, Land-use types are finer land-use types, while Land-use management intensity is further classification of the land-use types in various management intensities. Blue bars denote modelled estimates from this study, and red bars indicate reference data.**



605

Table A3: Global land-use, land cover, and management intensity comparison. Modelled values are derived from this study (2020 baseline). Reference values are derived from global datasets and various studies, where sources do not explicitly report global percentages, values are interpreted following the published methods.


Land-use / land cover	Modelled (%)	Reference (%)	Sources	Interpretation of reference value
Cropland	10.5	12–14	Taylor and Rising (2023), FAO UN (2024), HYDE 3.3 (Goldewijk et al.)	Derived from cropland area (ca.1.55–1.6 billion ha) divided by global ice-free land area
Forest	31.4	30–32	FAO FRA (2020), Kay et al., 2022	Directly reported global forest share around 31.1% of land area
Agroforestry	0.7	0.3–1.0	IPCC SRCCL (2019)	IPCC SRCCL highlights agroforestry as widespread but poorly mapped
Pasture	36	25–35	FAO (2022); Kay et al., 2022	Range synthesised from global pasture and rangeland estimates
Unmanaged land	18.6	20–25	IPCC SRCCL (2019)	Inferred from land cover classes and IPCC definitions of unmanaged or minimally managed ecosystems
Urban	0.74	0.5–0.8	FAO UN 2025	Urban land share inferred by combining urban population data with global urban land expansion estimates
Photovoltaics (PV)	0.01	<0.05	IRENA (2023); Fraunhofer ISE	Estimated by converting installed global PV capacity to land area using reported values
Water	1.8	2.0–2.5	FAO AQUASTAT	Directly reported global inland water surface fraction
Land-use/ land-cover subtype	Modelled (%)	Reference (%)	Sources	Interpretation of reference value
C3 cereals	4.48	4–6	FAO UN 2025	Derived from global harvested area fractions reported by FAO UN 2025 i.e., 731.9 Mha for cereals
C4 cereals	2.77	2–2.5	FAO UN 2025	Derived from crop-specific global area estimates i.e., 275 Mha for C4 cereals
Rice	1.3	~1.1%	FAO UN 2025	Derived from global harvested area fractions reported by FAO UN 2025 164 Mha for rice
Oil & N-fix crops (soy)	1.4	1–1.5	FAO UN 2025	Synthesised from soybean and oil/nitrogen-fixing crop area statistics i.e., 135 Mha
Starchy roots	0.18	0.1–0.3	FAO UN 2025	Derived from FAO crop category totals i.e., 17 M ha
Other crops	0.3	-	-	Residual crop category other than major crop types
Broadleaf	4.96	4–6	HILDA+ (Winkler et al.,	Derived from the percentage share of the



deciduous forest			2021), Ma et al., 2023	forest cover
Broadleaf evergreen forest	8.56	8-14	HILDA+ (Winkler et al., 2021), Ma et al., 2023	Derived from the percentage share of the forest cover
Needleleaf deciduous forest	4.42	3-5	HILDA+ (Winkler et al., 2021), Ma et al., 2023	Derived from the percentage share of the forest cover
Needleleaf evergreen forest	5.67	6-8	HILDA+ (Winkler et al., 2021), Ma et al., 2023	Derived from the percentage share of the forest cover
Land management intensity	Modelled (%)	Reference (%)	Sources	Interpretation of reference value
Very extensive crops	3.8	1-3	IPCC SRCCL (2019)	Inferred from low-input cropland systems described in IPCC SRCCL
Extensive crops	1.9	1-2	IPCC SRCCL (2019); FAO	Synthesised from sources
Intensive crops	4.3	3-5	IPCC SRCCL (2019)	Derived from high input intensity and irrigated cropping
Very extensive pasture	16.9	18-20	Bahar et al., 2020	Derived from the mentioned source
Extensive pasture	0.01	-	Bahar et al., 2020	Based on extensive grazing systems and low stocking densities
Intensive pasture	4.3	5-8	Bahar et al., 2020	Derived from the mentioned source
Intensive forest	1.58	2-3	FAO FRA (2020), Forest Extent Indicator (2024)	Derived from roughly 290-300 million hectares
Extensive forest	13.5	20-22%	FAO FRA (2020), Forest Extent Indicator (2024)	Derived from the mentioned source
Unmanaged forest	16.2	14-18	FAO FRA (2020), Forest Extent Indicator (2024)	Primary and minimally disturbed forest areas
Unmanaged pasture	14.8	14-18	FAO (2022); IPCC SRCCL	Low-input grazing and semi-natural rangelands
Intensive	8%	The percentages are with respect to the total managed land area excluding snow/ice, water and natural unmanaged land		
Extensive	19%			
Very extensive	69%			
Other land-use	4%			
Intensive	22%	The percentages are with respect to the total managed land covering cropland, pasture, and forestry only		
Extensive	33%			
Very extensive	45%			



Pearson correlation for the socioeconomic variables



610 **Figure A6: Multicollinearity check using pair-wise regression analysis and the variance inflation factor (VIF) methods for the**
 615 **socioeconomic variables**

615

620

625

630

635






Figure A7: Predicted relationships between crop management intensity and socioeconomic drivers across global income groups. The x-axes of each plot show the standardised predictor value (z-score) for various variables (GDP, GINI, market access, and population), and the y-axis indicates the predicted probability of each management intensity (very extensive, extensive, and intensive).



640 **Figure A8: Predicted relationships between pasture management intensity and socioeconomic drivers across global income groups. The x-axes of each plot show the standardised predictor value (z-score) for various variables (GDP, GINI, market access, and population), and the y-axis indicates the predicted probability of each management intensity (very extensive, extensive, and intensive).**



645 **Figure A9: Predicted relationships between forest management intensity and socioeconomic drivers across global income groups. The x-axes of each plot show the standardised predictor value (z-score) for various variables (GDP, GINI, market access, and population), and the y-axis indicates the predicted probability of each management intensity (very extensive, extensive, and intensive)**



650

Table A4: Multinomial logistic regression. Multinomial logistic regression results showing the association between socioeconomic variables and land-use management intensity levels (Extensive, Intensive) for cropland, pasture, and forest land-use types. Coefficients reported are alongside 95% confidence intervals. All models use the very extensive (VExt or unmanaged) category as the reference level. The asterisks (*) indicate significance levels: * p < 0.001.**

Land type	Region	Intensity	Variable	Coefficient	95% CI	p-value	Signif.
Crops	Global	Extensive	GDP	0.502	[0.500, 0.504]	<0.001	***
		Extensive	GINI	0.202	[0.201, 0.204]	<0.001	***
		Extensive	MarketAccess	0.181	[0.179, 0.183]	<0.001	***
		Extensive	Population	0.646	[0.641, 0.650]	<0.001	***
		Extensive	const	1.095	[1.093, 1.097]	<0.001	***
		Intensive	GDP	0.968	[0.966, 0.970]	<0.001	***
		Intensive	GINI	0.381	[0.380, 0.383]	<0.001	***
		Intensive	MarketAccess	0.615	[0.613, 0.617]	<0.001	***
		Intensive	Population	0.834	[0.830, 0.839]	<0.001	***
		Intensive	const	0.896	[0.895, 0.898]	<0.001	***
	High Income	Extensive	GDP	0.454	[0.452, 0.457]	<0.001	***
		Extensive	GINI	0.175	[0.172, 0.177]	<0.001	***
		Extensive	MarketAccess	0.273	[0.270, 0.276]	<0.001	***
		Extensive	Population	-0.048	[-0.050, -0.046]	<0.001	***
		Extensive	const	1.243	[1.240, 1.246]	<0.001	***
		Intensive	GDP	0.763	[0.760, 0.766]	<0.001	***
		Intensive	GINI	0.271	[0.268, 0.274]	<0.001	***
		Intensive	MarketAccess	0.615	[0.611, 0.618]	<0.001	***
		Intensive	Population	-0.214	[-0.217, -0.210]	<0.001	***
		Intensive	const	1.312	[1.309, 1.315]	<0.001	***
	Low Income	Extensive	GDP	0.022	[0.016, 0.028]	<0.001	***
		Extensive	GINI	0.088	[0.083, 0.093]	<0.001	***
		Extensive	MarketAccess	0.127	[0.120, 0.133]	<0.001	***
		Extensive	Population	0.431	[0.419, 0.444]	<0.001	***
		Extensive	const	0.506	[0.501, 0.511]	<0.001	***
		Intensive	GDP	-0.083	[-0.091, -0.076]	<0.001	***
		Intensive	GINI	0.12	[0.113, 0.128]	<0.001	***
Intensive		MarketAccess	0.607	[0.599, 0.615]	<0.001	***	



		Intensive	Population	0.482	[0.469, 0.496]	<0.001	***
		Intensive	const	-0.828	[-0.835, -0.821]	<0.001	***
	Lower Middle Income	Extensive	GDP	0.149	[0.146, 0.153]	<0.001	***
		Extensive	GINI	0.006	[0.003, 0.009]	<0.001	***
		Extensive	MarketAccess	0.153	[0.149, 0.157]	<0.001	***
		Extensive	Population	1.485	[1.474, 1.497]	<0.001	***
		Extensive	const	1.297	[1.292, 1.302]	<0.001	***
		Intensive	GDP	-0.166	[-0.170, -0.163]	<0.001	***
		Intensive	GINI	0.086	[0.082, 0.090]	<0.001	***
		Intensive	MarketAccess	0.971	[0.966, 0.976]	<0.001	***
		Intensive	Population	1.708	[1.697, 1.720]	<0.001	***
		Intensive	const	0.666	[0.661, 0.671]	<0.001	***
	Upper Middle Income	Extensive	GDP	0.107	[0.103, 0.110]	<0.001	***
		Extensive	GINI	0.53	[0.528, 0.533]	<0.001	***
		Extensive	MarketAccess	0.025	[0.022, 0.028]	<0.001	***
		Extensive	Population	1.158	[1.145, 1.172]	<0.001	***
		Extensive	const	1.143	[1.139, 1.147]	<0.001	***
		Intensive	GDP	0.483	[0.480, 0.487]	<0.001	***
		Intensive	GINI	0.794	[0.791, 0.798]	<0.001	***
		Intensive	MarketAccess	0.527	[0.523, 0.531]	<0.001	***
		Intensive	Population	1.562	[1.549, 1.576]	<0.001	***
		Intensive	const	1.001	[0.996, 1.005]	<0.001	***
Forest	Global	Extensive	GDP	0.044	[0.044, 0.045]	<0.001	***
		Extensive	GINI	0.509	[0.509, 0.510]	<0.001	***
		Extensive	MarketAccess	1.052	[1.051, 1.053]	<0.001	***
		Extensive	Population	-0.418	[-0.419, -0.417]	<0.001	***
		Extensive	const	-0.41	[-0.411, -0.410]	<0.001	***
		Intensive	GDP	0.18	[0.179, 0.182]	<0.001	***
		Intensive	GINI	0.432	[0.430, 0.433]	<0.001	***
		Intensive	MarketAccess	1.636	[1.635, 1.638]	<0.001	***
		Intensive	Population	-0.409	[-0.411, -0.407]	<0.001	***
		Intensive	const	-2.932	[-2.934, -2.930]	<0.001	***



	High Income	Extensive	GDP	-0.135	[-0.136, -0.134]	<0.001	***
		Extensive	GINI	0.977	[0.976, 0.978]	<0.001	***
		Extensive	MarketAccess	1.41	[1.408, 1.412]	<0.001	***
		Extensive	Population	-0.343	[-0.345, -0.341]	<0.001	***
		Extensive	const	-0.505	[-0.506, -0.503]	<0.001	***
		Intensive	GDP	-0.284	[-0.287, -0.282]	<0.001	***
		Intensive	GINI	1.001	[0.999, 1.004]	<0.001	***
		Intensive	MarketAccess	2.017	[2.015, 2.020]	<0.001	***
		Intensive	Population	-0.436	[-0.438, -0.434]	<0.001	***
		Intensive	const	-2.996	[-2.999, -2.993]	<0.001	***
	Low Income	Extensive	GDP	0.644	[0.640, 0.648]	<0.001	***
		Extensive	GINI	0.174	[0.171, 0.176]	<0.001	***
		Extensive	MarketAccess	0.207	[0.204, 0.211]	<0.001	***
		Extensive	Population	-0.377	[-0.383, -0.372]	<0.001	***
		Extensive	const	-0.25	[-0.252, -0.247]	<0.001	***
		Intensive	GDP	-0.068	[-0.123, -0.011]	0.018	*
		Intensive	GINI	-0.109	[-0.150, -0.068]	<0.001	***
		Intensive	MarketAccess	0.922	[0.896, 0.947]	<0.001	***
		Intensive	Population	-0.273	[-0.310, -0.236]	<0.001	***
		Intensive	const	-6.456	[-6.502, -6.410]	<0.001	***
	Lower Middle Income	Extensive	GDP	-0.337	[-0.339, -0.335]	<0.001	***
		Extensive	GINI	0.039	[0.037, 0.041]	<0.001	***
		Extensive	MarketAccess	0.298	[0.295, 0.300]	<0.001	***
		Extensive	Population	-0.236	[-0.239, -0.233]	<0.001	***
		Extensive	const	0.168	[0.166, 0.170]	<0.001	***
		Intensive	GDP	-0.154	[-0.162, -0.146]	<0.001	***
		Intensive	GINI	0.091	[0.080, 0.102]	<0.001	***
		Intensive	MarketAccess	1.585	[1.576, 1.595]	<0.001	***
		Intensive	Population	-1.141	[-1.162, -1.119]	<0.001	***
		Intensive	const	-4.678	[-4.693, -4.662]	<0.001	***
Upper Middle	Extensive	GDP	0.164	[0.162, 0.165]	<0.001	***	
	Extensive	GINI	0.418	[0.416, 0.419]	<0.001	***	



	Income	Extensive	MarketAccess	0.941	[0.939, 0.943]	<0.001	***
		Extensive	Population	-0.329	[-0.331, -0.326]	<0.001	***
		Extensive	const	-0.321	[-0.322, -0.320]	<0.001	***
		Intensive	GDP	0.238	[0.236, 0.241]	<0.001	***
		Intensive	GINI	0.015	[0.012, 0.017]	<0.001	***
		Intensive	MarketAccess	1.513	[1.510, 1.515]	<0.001	***
		Intensive	Population	-0.112	[-0.114, -0.110]	<0.001	***
		Intensive	const	-2.337	[-2.340, -2.334]	<0.001	***
Pasture	Global	Extensive	GDP	-0.396	[-0.398, -0.394]	<0.001	***
		Extensive	GINI	0.295	[0.294, 0.297]	<0.001	***
		Extensive	MarketAccess	0.91	[0.909, 0.912]	<0.001	***
		Extensive	Population	0.061	[0.060, 0.062]	<0.001	***
		Extensive	const	-2.546	[-2.547, -2.544]	<0.001	***
		Intensive	GDP	-0.009	[-0.012, -0.007]	<0.001	***
		Intensive	GINI	0.392	[0.390, 0.394]	<0.001	***
		Intensive	MarketAccess	1.075	[1.073, 1.076]	<0.001	***
		Intensive	Population	0.071	[0.070, 0.072]	<0.001	***
		Intensive	const	-3.595	[-3.598, -3.592]	<0.001	***
	High Income	Extensive	GDP	-0.476	[-0.479, -0.472]	<0.001	***
		Extensive	GINI	0.444	[0.441, 0.447]	<0.001	***
		Extensive	MarketAccess	1.245	[1.242, 1.247]	<0.001	***
		Extensive	Population	-0.036	[-0.038, -0.034]	<0.001	***
		Extensive	const	-3.711	[-3.716, -3.706]	<0.001	***
		Intensive	GDP	-0.405	[-0.409, -0.400]	<0.001	***
		Intensive	GINI	0.414	[0.410, 0.418]	<0.001	***
		Intensive	MarketAccess	1.38	[1.377, 1.383]	<0.001	***
		Intensive	Population	-0.078	[-0.081, -0.075]	<0.001	***
		Intensive	const	-4.391	[-4.398, -4.384]	<0.001	***
	Low Income	Extensive	GDP	-0.4	[-0.406, -0.395]	<0.001	***
		Extensive	GINI	0.275	[0.272, 0.278]	<0.001	***
		Extensive	MarketAccess	0.211	[0.208, 0.215]	<0.001	***
		Extensive	Population	0.171	[0.165, 0.176]	<0.001	***




		Extensive	const	-2.194	[-2.198, -2.190]	<0.001	***
		Intensive	GDP	-0.331	[-0.343, -0.318]	<0.001	***
		Intensive	GINI	0.407	[0.400, 0.414]	<0.001	***
		Intensive	MarketAccess	0.28	[0.272, 0.287]	<0.001	***
		Intensive	Population	0.207	[0.201, 0.213]	<0.001	***
		Intensive	const	-4.101	[-4.110, -4.092]	<0.001	***
	Lower Middle Income	Extensive	GDP	-0.217	[-0.220, -0.213]	<0.001	***
		Extensive	GINI	0.189	[0.185, 0.192]	<0.001	***
		Extensive	MarketAccess	0.536	[0.533, 0.539]	<0.001	***
		Extensive	Population	0.136	[0.132, 0.140]	<0.001	***
		Extensive	const	-1.781	[-1.785, -1.778]	<0.001	***
		Intensive	GDP	-0.234	[-0.241, -0.227]	<0.001	***
		Intensive	GINI	0.221	[0.215, 0.228]	<0.001	***
		Intensive	MarketAccess	0.643	[0.638, 0.649]	<0.001	***
		Intensive	Population	0.158	[0.153, 0.163]	<0.001	***
		Intensive	const	-3.431	[-3.439, -3.423]	<0.001	***
	Upper Middle Income	Extensive	GDP	-0.246	[-0.248, -0.243]	<0.001	***
		Extensive	GINI	0.354	[0.352, 0.356]	<0.001	***
		Extensive	MarketAccess	0.915	[0.913, 0.917]	<0.001	***
		Extensive	Population	0.232	[0.229, 0.235]	<0.001	***
		Extensive	const	-2.25	[-2.252, -2.248]	<0.001	***
		Intensive	GDP	-0.057	[-0.060, -0.053]	<0.001	***
		Intensive	GINI	0.432	[0.429, 0.435]	<0.001	***
		Intensive	MarketAccess	1.05	[1.047, 1.052]	<0.001	***
		Intensive	Population	0.249	[0.246, 0.252]	<0.001	***
		Intensive	const	-3.217	[-3.221, -3.214]	<0.001	***




Table A5: Summary of marginal-effect relationships between socioeconomic/biophysical variables and land management intensity across land-use sectors and income groups


Land sector	Variable	Main relationship	Dominant management response	Direction	Marginal effect range (percentage points)	Strongest regions/groups
Cropland	GDP per capita	Positive	Intensive irrigated cropland	↑	+10 to +17	High-income, upper-middle-income
Cropland	Market access	Positive	Intensive cropland	↑	+8 to +14 / -18	Most regions
			Very extensive	↓		
Cropland	Population density	Positive	Intensive cropland	↑	+5 to +10	Upper-/lower-middle income
Cropland	GINI	Negative for intensive	Extensive cropland	↑	-3 to -8	Several regions
Cropland	Aridity index	Positive for low-intensity systems	Very extensive cropland	↑	+7 to +14	Multiple regions
Cropland	Elevation	Region dependent	Mixed responses	↑↓	-10 to +15	Region specific
Pasture	Market access	Positive	Intensive pasture	↑	+1 to +6	High-, upper-, lower-middle income
Pasture	Aridity index	Strong positive	Intensive pasture	↑	up to +22	Most regions
Pasture	Population density	Slight positive	Intensive pasture	↑	+2 to +6	Multiple regions
Pasture	GDP per capita	Weak positive	Intensive pasture	↑	-2 to +8	Low-income
Forest	Market access	Positive	Intensive/extensive forest	↑	up to +40	Multiple regions
Forest	GINI	Positive for extensive	Extensive forest	↑	+5 to +22	High-, upper-middle income
Forest	Aridity index	Positive for unmanaged	Unmanaged forest	↑	variable	Multiple regions
Forest	Population density	Mixed	Region-dependent	-	variable	Region specific



660 **Figure A10: Spatial thinning robustness analysis of marginal effects for cropland management intensity classes.** Marginal effects were recalculated using progressively coarser spatial samples (0.10°, 0.25°, and 0.50°). Lines show marginal effects for each thinning level relative to the full-resolution model (0.01°), while shaded areas indicate confidence intervals.



665 **Figure A11: Spatial thinning robustness analysis of marginal effects for pasture management intensity classes. Marginal effects were recalculated using progressively coarser spatial samples (0.10°, 0.25°, and 0.50°). Lines show marginal effects for each thinning level relative to the full-resolution model (0.01°), while shaded areas indicate confidence intervals.**



670 **Figure A12: Spatial thinning robustness analysis of marginal effects for cropland management intensity classes.** Marginal effects were recalculated using progressively coarser spatial samples (0.10°, 0.25°, and 0.50°). Lines show marginal effects for each thinning level relative to the full-resolution model (0.01°), while shaded areas indicate confidence intervals.



Table A6: spatial thinning based robustness check for various coarser spatial samples such as 0.10°, 0.25°, and 0.50°

Land	MgmtClass	Variable	full	thin_0p10deg	thin_0p25deg	thin_0p50deg	Direction_stability
crops	Ext_IRR	Aridity_index	-	-	-	-	stable
crops	Ext_IRR	Elevation	-	-	-	-	stable
crops	Ext_IRR	GDP	-	-	-	-	stable
crops	Ext_IRR	GINI	+	+	+	+	stable
crops	Ext_IRR	MarketAccess	-	-	+	+	mixed
crops	Ext_IRR	Population	-	-	-	-	stable
crops	Ext_RF	Aridity_index	-	-	-	-	stable
crops	Ext_RF	Elevation	-	-	-	-	stable
crops	Ext_RF	GDP	-	-	-	-	stable
crops	Ext_RF	GINI	+	+	+	-	mixed
crops	Ext_RF	MarketAccess	-	-	-	-	stable
crops	Ext_RF	Population	-	-	-	-	stable
crops	Int_IRR	Aridity_index	-	-	-	-	stable
crops	Int_IRR	Elevation	+	+	+	+	stable
crops	Int_IRR	GDP	+	+	+	+	stable
crops	Int_IRR	GINI	+	+	+	+	stable
crops	Int_IRR	MarketAccess	+	+	+	+	stable
crops	Int_IRR	Population	+	+	+	+	stable
crops	Int_RF	Aridity_index	+	+	+	+	stable
crops	Int_RF	Elevation	+	+	+	+	stable
crops	Int_RF	GDP	-	+	+	+	mixed
crops	Int_RF	GINI	-	-	-	-	stable
crops	Int_RF	MarketAccess	-	-	-	-	stable
crops	Int_RF	Population	-	-	-	-	stable
crops	VExt	Aridity_index	+	+	+	+	stable
crops	VExt	Elevation	-	-	-	-	stable
crops	VExt	GDP	-	-	-	-	stable
crops	VExt	GINI	+	+	+	+	stable
crops	VExt	MarketAccess	-	-	-	-	stable
crops	VExt	Population	+	+	+	+	stable
forest	Ext	Aridity_index	-	-	-	-	stable
forest	Ext	Elevation	-	-	-	-	stable
forest	Ext	GDP	-	-	-	-	stable
forest	Ext	GINI	+	+	+	+	stable
forest	Ext	MarketAccess	+	+	+	+	stable
forest	Ext	Population	-	-	-	-	stable
forest	Int	Aridity_index	-	+	+	+	mixed



forest	Int	Elevation	-	-	-	-	stable
forest	Int	GDP	+	+	-	+	mixed
forest	Int	GINI	+	+	+	+	stable
forest	Int	MarketAccess	+	+	+	+	stable
forest	Int	Population	-	-	-	-	stable
forest	UNM	Aridity_index	+	+	+	+	stable
forest	UNM	Elevation	+	+	+	+	stable
forest	UNM	GDP	+	+	+	+	stable
forest	UNM	GINI	-	-	-	-	stable
forest	UNM	MarketAccess	-	-	-	-	stable
forest	UNM	Population	+	+	+	+	stable
pasture	Ext	Aridity_index	+	+	+	+	stable
pasture	Ext	Elevation	+	+	+	+	stable
pasture	Ext	GDP	-	-	-	-	stable
pasture	Ext	GINI	-	+	-	+	mixed
pasture	Ext	MarketAccess	+	+	+	+	stable
pasture	Ext	Population	+	-	-	-	mixed
pasture	Int	Aridity_index	+	+	+	+	stable
pasture	Int	Elevation	+	-	-	-	mixed
pasture	Int	GDP	-	-	-	-	stable
pasture	Int	GINI	+	+	+	+	stable
pasture	Int	MarketAccess	+	+	+	+	stable
pasture	Int	Population	+	+	-	-	mixed
pasture	VExt	Aridity_index	-	-	-	-	stable
pasture	VExt	Elevation	-	+	+	+	mixed
pasture	VExt	GDP	+	+	+	+	stable
pasture	VExt	GINI	-	-	-	-	stable
pasture	VExt	MarketAccess	-	-	-	-	stable
pasture	VExt	Population	-	-	+	+	mixed




Figure A13: Distribution of predicted probabilities for land management intensity. The distributions for predicted probabilities are shown for cropland, pasture and forest systems across global and world income groups. Violins represent full distribution across grid cells, while embedded boxplots indicate the interquartile range (25th- 75th percentiles), median, and dispersion within each group.

680



Data availability

We provide a global land management intensity dataset explorer (<https://ee-ankitasaxena03as.projects.earthengine.app/view/global-land-use-management>), and these data are available at <https://doi.org/10.5281/zenodo.18249970> (Saxena et al., 2026) for research purposes. Input data for socioeconomic variables (GDP, GINI, market access and working age population) can be accessed through <https://doi.org/10.5281/zenodo.17580780> (Perkins et al., 2025) and world region boundaries can be found at <https://datacatalog.worldbank.org/>. Any other data that support the findings of this study are included within the article, online methods and supplementary materials.

Conclusions

Overall, the study shows that management intensity cannot be treated as a homogeneous response to socioeconomic or biophysical change, since it emerges from complex, sector-specific and non-linear interactions between socioeconomic, biophysical, and behavioural factors. These findings challenge the simplified assumptions that are commonly embedded in global land-use and integrated assessment models, where economic growth is often assumed to drive land-use change uniformly (e.g., Popp et al., 2017; Stehfest et al., 2019; Ma et al., 2024). Instead, the results show distinct relationships with socioeconomic variables across cropland, pasture and forest management. At the same time, broad consistency between land management intensity classes and land-use decision-making types indicates that management intensity can serve as a useful proxy if partial for land-use behaviour at large scales (Brown et al., 2017; Rounsevell et al., 2021). While the observed mismatches further indicate that decision-making strategies do not translate into a single management outcome, particularly in heterogenous landscapes.

Author contributions

AS: Conceptualization, Data curation, Methodology, Validation, Investigation, Formal analysis, Writing – original draft, Writing – review & editing. CB: Conceptualization, Methodology, Writing – review & editing. KW: Writing – review & editing, AA: Methodology, Writing – Writing – review & editing, MR: Conceptualization, Methodology, Writing – review & editing.

Competing interests

Authors declare no competing interests.



Acknowledgements

Authors acknowledge the KIT Future Fields Programme and Helmholtz Association, Germany for supporting this work. The authors also acknowledge the use of AI-based generative tools (GPT-5.5) to improve the language and readability in some parts of the manuscript, with all content carefully reviewed and assessed by the authors.

710 References

- Adalibieke, W., Cui, X., Cai, H., You, L. and Zhou, F., 2023. Global crop-specific nitrogen fertilization dataset in 1961–2020. *Scientific data*, 10(1), p.617.
- Arneith, A., Alexander, P., Fuchs, R., Shin, Y., Claudet, J., Coll, M., Leadley, P., Rondinini, C. & Rounsevell, M.D.A. (2025). The biodiversity-climate-food nexus: illustrating challenges and solutions using the Green Shoots framework. *One Earth*, <https://doi.org/10.1016/j.oneear.2025.101455>
- 715 Alexander, P., Rounsevell, M.D., Dislich, C., Dodson, J.R., Engström, K. and Moran, D., 2015. Drivers for global agricultural land use change: The nexus of diet, population, yield and bioenergy. *Global Environmental Change*, 35, pp.138–147.
- Bahar, N.H., Lo, M., Sanjaya, M., Van Vianen, J., Alexander, P., Ickowitz, A. and Sunderland, T., 2020. Meeting the food
720 security challenge for nine billion people in 2050: What impact on forests?. *Global Environmental Change*, 62, p.102056.
- Bayer, A.D., Lautenbach, S. and Arneith, A., 2023. Benefits and trade-offs of optimizing global land use for food, water, and carbon. *Proceedings of the National Academy of Sciences*, 120(42), p.e2220371120.
- Brown, C., Alexander, P., Holzhauser, S. and Rounsevell, M.D., 2017. Behavioral models of climate change adaptation and mitigation in land-based sectors. *Wiley Interdisciplinary reviews: climate change*, 8(2), p.e448.
- 725 Dou, Y., Cosentino, F., Malek, Z., Maiorano, L., Thuiller, W. and Verburg, P.H., 2021. A new European land systems representation accounting for landscape characteristics. *Landscape Ecology*, 36(8), pp.2215–2234.
- Erb, K.H., Luysaert, S., Meyfroidt, P., Pongratz, J., Don, A., Kloster, S., Kuemmerle, T., Fetzel, T., Fuchs, R., Herold, M. and Haberl, H., 2017. Land management: data availability and process understanding for global change studies. *Global change biology*, 23(2), pp.512–533.
- 730 Erb, K.H., Lauk, C., Kastner, T., Mayer, A., Theurl, M.C. and Haberl, H., 2016. Exploring the biophysical option space for feeding the world without deforestation. *Nature communications*, 7(1), p.11382.
- FAO (2022). FAOSTAT Statistical Database. Food and Agriculture Organization of the United Nations. <http://www.fao.org/faostat>
- Food and Agriculture Organization of the United Nations (2024) – with major processing by Our World in Data.
735 <https://ourworldindata.org/forest-area>
- FAO FRA (2020). Global Forest Resources Assessment 2020: Main report. Food and Agriculture Organization of the United Nations. <https://www.fao.org/forest-resources-assessment/en/>



- Fraunhofer ISE (2023). Photovoltaics Report. Fraunhofer Institute for Solar Energy Systems ISE.
<https://www.ise.fraunhofer.de>
- 740 Foley, J.A., Ramankutty, N., Brauman, K.A., Cassidy, E.S., Gerber, J.S., Johnston, M., Mueller, N.D., O'Connell, C., Ray, D.K., West, P.C. and Balzer, C., 2011. Solutions for a cultivated planet. *Nature*, 478(7369), pp.337-342.
- “Forest Extent Indicator.” Global Forest Review, updated April 4, 2024. Washington, DC: World Resources Institute. Available online at <https://research.wri.org/gfr/forest-extent-indicators/forest-extent>.
- Griscom, B.W., Adams, J., Ellis, P.W., Houghton, R.A., Lomax, G., Miteva, D.A., Schlesinger, W.H., Shoch, D., Siikamäki, J.V., Smith, P. and Woodbury, P., 2017. Natural climate solutions. *Proceedings of the National Academy of Sciences*, 114(44), pp.11645-11650.
- Haberl, H., Erb, K.H. and Krausmann, F., 2014. Human appropriation of net primary production: patterns, trends, and planetary boundaries. *Annual Review of Environment and Resources*, 39, pp.363-391.
- Harrison, G.R., Rigge, M., Assal, T.J., Applestein, C., James, D.K. and McCord, S.E., 2025. An accuracy assessment of satellite-derived rangeland fractional cover. *Ecological Indicators*, 172, p.113267.
- 750 Herrero, M., Havlík, P., Valin, H., Notenbaert, A., Rufino, M.C., Thornton, P.K., Blümmel, M., Weiss, F., Grace, D. and Obersteiner, M., 2013. Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proceedings of the National Academy of Sciences*, 110(52), pp.20888-20893.
- IPCC (2022). Special Report on Climate Change and Land (SRCCL). Intergovernmental Panel on Climate Change.
755 <https://www.ipcc.ch/srccl/>
- IPCC AR6 (2022). Sixth Assessment Report: Working Group III – Climate Change Mitigation. Intergovernmental Panel on Climate Change. <https://www.ipcc.ch/report/ar6/wg3/>
- IRENA (2023). Renewable Capacity Statistics 2023. International Renewable Energy Agency.
<https://www.irena.org/Statistics>
- 760 Jayne, T.S., Snapp, S., Place, F. and Sitko, N., 2019. Sustainable agricultural intensification in an era of rural transformation in Africa. *Global Food Security*, 20, pp.105-113.
- Kastner, T., Matej, S., Forrest, M., Gingrich, S., Haberl, H., Hickler, T., Krausmann, F., Lasslop, G., Niedertscheider, M., Plutzer, C. and Schwarzmüller, F., 2022. Land use intensification increasingly drives the spatiotemporal patterns of the global human appropriation of net primary production in the last century. *Global Change Biology*, 28(1), pp.307-322.
- 765 Kay, M., Bunning, S., Burke, J., Boerger, V., Bojic, D., Bosc, P.M., Clark, M., Dale, D., England, M., Hoogeveen, J. and Koo-Oshima, S., 2022. The state of the world's land and water resources for food and agriculture 2021. Systems at breaking point.
- Kebede, E.A., Oluoch, K.O.A., Siebert, S., Mehta, P., Hartman, S., Jägermeyr, J., Ray, D., Ali, T., Brauman, K.A., Deng, Q. and Xie, W., 2025. A global open-source dataset of monthly irrigated and rainfed cropped areas (MIRCA-OS) for the 21st
770 century. *Scientific Data*, 12(1), p.208.



- Kehoe, L., Kuemmerle, T., Meyer, C., Levers, C., Václavík, T. and Kreft, H., 2015. Global patterns of agricultural land-use intensity and vertebrate diversity. *Diversity and Distributions*, 21(11), pp.1308-1318.
- Klein Goldewijk, K., Beusen, A., Doelman, J. and Stehfest, E., 2017. Anthropogenic land use estimates for the Holocene–HYDE 3.2. *Earth System Science Data*, 9(2), pp.927-953.
- 775 Kleijn, D., Kohler, F., Báldi, A., Batáry, P., Concepción, E.D., Clough, Y., Díaz, M., Gabriel, D., Holzschuh, A., Knop, E. and Kovács, A., 2009. On the relationship between farmland biodiversity and land-use intensity in Europe. *Proceedings of the royal society B: biological sciences*, 276(1658), pp.903-909.
- Lambin, E.F. and Meyfroidt, P., 2011. Global land use change, economic globalization, and the looming land scarcity. *Proceedings of the national academy of sciences*, 108(9), pp.3465-3472.
- 780 Ma, H., Crowther, T.W., Mo, L., Maynard, D.S., Renner, S.S., Van den Hoogen, J., Zou, Y., Liang, J., De-Miguel, S., Nabuurs, G.J. and Reich, P.B., 2023. The global biogeography of tree leaf form and habit. *Nature Plants*, 9(11), pp.1795-1809.
- Matej, S., Weidinger, F., Kaufmann, L., Roux, N., Gingrich, S., Haberl, H., Krausmann, F. and Erb, K.H., 2025. A global land-use data cube 1992–2020 based on the Human Appropriation of Net Primary Production. *Scientific Data*, 12(1), p.511.
- 785 Meyfroidt, P., Chowdhury, R.R., de Bremond, A., Ellis, E.C., Erb, K.H., Filatova, T., Garrett, R.D., Grove, J.M., Heinimann, A., Kuemmerle, T. and Kull, C.A., 2018. Middle-range theories of land system change. *Global environmental change*, 53, pp.52-67.
- Meyfroidt, P., De Bremond, A., Ryan, C.M., Archer, E., Aspinall, R., Chhabra, A., Camara, G., Corbera, E., DeFries, R., Díaz, S. and Dong, J., 2022. Ten facts about land systems for sustainability. *Proceedings of the National Academy of Sciences*, 119(7), p.e2109217118.
- 790 Lambin, E.F. and Meyfroidt, P., 2011. Global land use change, economic globalization, and the looming land scarcity. *Proceedings of the national academy of sciences*, 108(9), pp.3465-3472.
- Lesiv, M., Schepaschenko, D., Buchhorn, M., See, L., Dürauer, M., Georgieva, I., Jung, M., Hofhansl, F., Schulze, K., Bilous, A. and Blyshchyk, V., 2022. Global forest management data for 2015 at a 100 m resolution. *Scientific Data*, 9(1), p.199.
- 795 Li, A., Liu, L., Li, S., Cui, X., Chen, X. and Cao, X., 2025. Global photovoltaic solar panel dataset from 2019 to 2022. *Scientific Data*, 12(1), p.637.
- Liu, X., Zheng, J., Yu, L., Hao, P., Chen, B., Xin, Q., Fu, H. and Gong, P., 2021. Annual dynamic dataset of global cropping intensity from 2001 to 2019. *Scientific data*, 8(1), p.283.
- 800 Ma, J., Tian, L., Zhang, Y., Yang, X., Li, Y., Liu, Z., Zhou, L., Wang, Z. and Ouyang, W., 2024. Global property rights and land use efficiency. *Nature Communications*, 15(1), p.8525.
- Malek, Ž., Douw, B., Van Vliet, J., Van Der Zanden, E.H. and Verburg, P.H., 2019. Local land-use decision-making in a global context. *Environmental Research Letters*, 14(8), p.083006.



- Malek, Ž. and Verburg, P.H., 2020. Mapping global patterns of land use decision-making. *Global Environmental Change*, 65, p.102170.
- 805
- Mehta, P., Siebert, S., Kummu, M., Deng, Q., Ali, T., Marston, L., Xie, W. and Davis, K.F., 2024. Half of twenty-first century global irrigation expansion has been in water-stressed regions. *Nature Water*, 2(3), pp.254-261.
- Nagaraj, D., Proust, E., Todeschini, A., Rulli, M.C. and D'Odorico, P., 2021. A new dataset of global irrigation areas from 2001 to 2015. *Advances in Water Resources*, 152, p.103910.
- 810
- Ostrom, E., 2009. A general framework for analyzing sustainability of social-ecological systems. *Science*, 325(5939), pp.419-422.
- Overmars, K.P., Schulp, C.J., Alkemade, R., Verburg, P.H., Temme, A.J., Omtzigt, N. and Schaminée, J.H., 2014. Developing a methodology for a species-based and spatially explicit indicator for biodiversity on agricultural land in the EU. *Ecological Indicators*, 37, pp.186-198.
- 815
- Parente, L., Sloat, L., Mesquita, V., Consoli, D., Stanimirova, R., Hengl, T., Bonannella, C., Teles, N., Wheeler, I., Hunter, M. and Ehrmann, S., 2024. Annual 30-m maps of global grassland class and extent (2000–2022) based on spatiotemporal Machine Learning. *Scientific data*, 11(1), p.1303.
- Pergner, I. and Lippert, C., 2023. On the effects that motivate pesticide use in perspective of designing a cropping system without pesticides but with mineral fertilizer—a review. *Agronomy for Sustainable Development*, 43(2), p.24.
- 820
- Perkins, O., Saxena, A., Seo, B., Millington, J., Brown, C., & Rounsevell, M. (2025). Gridded socio-economic capitals for the SSPs [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.17580780>
- Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L., Dietrich, J.P., Doelmann, J.C., Gusti, M. and Hasegawa, T., 2017. Land-use futures in the shared socio-economic pathways. *Global Environmental Change*, 42, pp.331-345.
- 825
- Rounsevell, M.D. and Arneth, A., 2011. Representing human behaviour and decisional processes in land system models as an integral component of the earth system. *Global Environmental Change*, 21(3), pp.840-843.
- Rounsevell, M.D., Arneth, A., Brown, C., Cheung, W.W., Gimenez, O., Holman, I., Leadley, P., Luján, C., Mahevas, S., Maréchaux, I. and Péliissier, R., 2021. Identifying uncertainties in scenarios and models of socio-ecological systems in support of decision-making. *One Earth*, 4(7), pp.967-985.
- 830
- Rudel, T.K., Schneider, L. and Uriarte, M., 2010. Forest transitions: An introduction. *Land use policy*, 27(2), pp.95-97.
- Saxena, A., Brown, C., Winkler, K., Arneth, A. and Rounsevell, M., 2026. Global Gridded Land Management Dataset at 0.01-degrees. <https://doi.org/10.5281/zenodo.18249970>
- Stehfest, E., van Zeist, W.J., Valin, H., Havlik, P., Popp, A., Kyle, P., Tabeau, A., Mason-D'Croz, D., Hasegawa, T., Bodirsky, B.L. and Calvin, K., 2019. Key determinants of global land-use projections. *Nature communications*, 10(1),
- 835
- p.2166.
- Tang, F.H., Nguyen, T.H., Conchedda, G., Casse, L., Tubiello, F.N. and Maggi, F., 2024. CROPGRIDS: a global geo-referenced dataset of 173 crops. *Scientific Data*, 11(1), p.413.



- 840 Tian, H., Bian, Z., Shi, H., Qin, X., Pan, N., Lu, C., Pan, S., Tubiello, F.N., Chang, J., Conchedda, G. and Liu, J., 2022. History of anthropogenic Nitrogen inputs (HaNi) to the terrestrial biosphere: A 5-arcmin resolution annual dataset from 1860 to 2019. *Earth System Science Data Discussions*, 2022, pp.1-32.
- Turner, B., Devisscher, T., Chabaneix, N., Woroniecki, S., Messier, C. and Seddon, N., 2022. The role of nature-based solutions in supporting social-ecological resilience for climate change adaptation. *Annual Review of Environment and Resources*, 47(1), pp.123-148.
- UN DESA (2018). World Urbanization Prospects: The 2018 Revision. United Nations Department of Economic and Social Affairs, Population Division. <https://population.un.org/wup/>
- 845 Utrecht University/PBL Netherlands Environmental Assessment Agency - History Database of the Global Environment (HYDE v 3.3, 2023). Klein Goldewijk, C.G.M., Beusen, A., Doelman, J., Stehfest, E., 2017, Anthropogenic land use estimates for the Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927–953
- Vortkamp, I. and Hilker, F.M., 2023. Farmers' land-use decision-making: A dynamical modelling approach that integrates qualitative knowledge about social norms into a quantitative model. *People and Nature*, 5(4), pp.1147-1159.
- 850 Winkler, K., Fuchs, R., Rounsevell, M. and Herold, M., 2020. *HILDA+ Global land use change between 1960 and 2019 [Dataset]*. PANGAEA [online]
- Winkler, K., Fuchs, R., Rounsevell, M. and Herold, M., 2021. Global land use changes are four times greater than previously estimated. *Nature communications*, 12(1), p.2501.
- 855 Winkler, K., Fuchs, R., Rounsevell, M. and Herold, M., 2025. Six decades of global crop yield increase and cropland expansion from 1960 to 2020. *Environmental Research Communications*, 7(5), p.055013.
- Zabel, F., Delzeit, R., Schneider, J.M., Seppelt, R., Mauser, W. and Václavík, T., 2019. Global impacts of future cropland expansion and intensification on agricultural markets and biodiversity. *Nature communications*, 10(1), p.2844.
- 860 Zomer, R.J., Xu, J. and Trabucco, A., 2022. Version 3 of the global aridity index and potential evapotranspiration database. *Scientific Data*, 9(1), p.409.