



A multi-decadal, depth-resolved Area-of-Habitat dataset for global marine species

Yixuan Li^{1, #}, Shijun Zheng^{1, #}, Le Yu^{1,2, *}, Zhenrong Du³, Fanghua Xu¹, Mingjie Li⁴,
Zhibing Jiang⁵, Yibo Liao⁵, Yining Chen⁵, Bangyi Tao⁶

¹ Department of Earth System Science, Ministry of Education Key Laboratory for Earth System Modeling, Institute for Global Change Studies, Tsinghua University, Beijing 100084, China

² Ministry of Education Ecological Field Station for East Asian Migratory Birds, Beijing 100084, China

³ School of Information and Communication Engineering, Dalian University of Technology, Dalian 116024, China

⁴ Technology Innovation Center for South China Sea Remote Sensing, Surveying and Mapping Collaborative Application, South China Sea Development Research Institute, Ministry of Natural Resources, Guangzhou 510300, China

⁵ Key Laboratory of Marine Ecosystem Dynamics, Second Institute of Oceanography, Ministry of Natural Resources, Hangzhou 310012, China

⁶ State Key Laboratory of Satellite Ocean Environment Dynamics, Second Institute of Oceanography, Ministry of Natural Resources, Hangzhou 310012, China

These authors contributed equally.

* Corresponding author: Le Yu (leyu@tsinghua.edu.cn)

Abstract

Marine biodiversity is being rapidly redistributed in response to ocean change, yet globally consistent datasets that resolve suitable habitat across both depth and time remain scarce. Here, we present a multi-decadal, depth-resolved Area-of-Habitat (AOH) dataset for a globally distributed subset of marine species assessed by the IUCN Red List. We integrated approximately 45.5 million occurrence records with IUCN expert range maps and physical and biogeochemical ocean reanalysis data spanning 1993–2025. Species-specific MaxEnt models were successfully fitted for 5,619 species (primarily fishes (4,815 species), together with reef-forming corals, molluscs, lobsters, marine mammals, marine reptiles, mangroves and seagrasses. The fitted models were



28 projected across 50 standard depth layers for 1995, 2000, 2005, 2010, 2015, 2020 and
29 2025. Phytoplankton, zooplankton and macroalgae were not included. Model
30 predictions were constrained to the original IUCN ranges to generate continuous, range-
31 constrained estimates of environmental suitability, operationally defined here as marine
32 AOH. The resulting dataset comprises species-level layers and aggregated products
33 resolved by depth and year. Internal consistency assessments showed that predicted
34 suitability was generally higher at species occurrence locations across the evaluated
35 species and depth layers. Independent comparisons with global coral-reef distributions
36 and FishGlob scientific survey observations further supported the ecological
37 plausibility and spatial realism of the resulting AOH estimates.

38 Aggregated analyses revealed pronounced vertical differentiation in marine AOH,
39 with shallow and deeper waters exhibiting distinct spatial concentrations and temporal
40 trajectories. By integrating geographic space, ocean depth and multi-decadal
41 environmental change within a consistent modelling framework, this dataset provides
42 a standardized resource for investigating marine habitat redistribution, identifying
43 persistent areas of high suitability and potential climate refugia, and evaluating the
44 correspondence between suitable habitats and marine protected areas in a changing
45 ocean. The dataset is publicly available at <https://doi.org/10.5281/zenodo.21933262>
46 (Yu and Li, 2026).

47 **1. Introduction**

48 Marine biodiversity is declining under the combined pressures of climate change,
49 overexploitation, pollution and habitat degradation, with broad consequences for
50 ecosystem functioning and the benefits that marine ecosystems provide to society
51 (IPBES, 2019; Pörtner et al., 2023). In response, the Kunming–Montreal Global
52 Biodiversity Framework calls for the effective conservation and management of at least
53 30% of coastal and marine areas by 2030 (CBD, 2022). Achieving this target requires
54 spatially explicit, repeatable and comparable information on species and their habitats,
55 enabling biodiversity patterns, conservation priorities and protected-area coverage to



56 be assessed consistently across space and time. Area of Habitat (AOH) mapping
 57 provides a standardized framework for delineating environmentally suitable habitat
 58 within species' geographic ranges and for aggregating such information across species
 59 to support biodiversity assessment (Brooks et al., 2019; Lumbierres et al., 2022). When
 60 generated consistently across multiple time points, AOH maps can further characterize
 61 changes in the extent and spatial distribution of suitable habitat, thereby supporting
 62 biodiversity monitoring and conservation planning (Yang et al., 2025; Li et al., 2025).
 63 Extending this framework to marine systems therefore offers an important basis for
 64 assessing large-scale patterns of marine habitat suitability and tracking their changes
 65 through time.

66 Substantial progress has been made in developing standardized AOH datasets for
 67 terrestrial biodiversity, including multispecies products for mammals, birds and
 68 amphibians at regional to global scales (Rondinini et al., 2011; Ficetola et al., 2015;
 69 Palacio et al., 2021; Lumbierres et al., 2022). More recent datasets have further
 70 incorporated historical time series or future scenario projections, enabling dynamic
 71 assessments of habitat change (Yang et al., 2025; Li et al., 2025). The need for dynamic
 72 assessment also extends to marine systems, including deeper waters where species
 73 adapted to long-term environmental stability are nevertheless expected to face
 74 increasing climate velocities even under ambitious mitigation pathways (Brito-Morales
 75 et al., 2020; UNEP, 2026). Comparable global marine mapping has advanced primarily
 76 through habitat-suitability and species-distribution frameworks such as AquaMaps and
 77 AquaX, which combine occurrence records, geographic constraints and environmental
 78 information to estimate suitable areas for thousands of marine species (Kaschner et al.,
 79 2006; Ready et al., 2010; Reygondeau et al., 2026). However, existing global marine
 80 products remain limited in two important respects. First, most represent either
 81 climatological baselines or a small number of widely separated future periods, rather
 82 than a consistent multi-decadal sequence suitable for examining historical changes in
 83 marine habitat. Such limited temporal coverage constrains the ability to distinguish



84 persistent redistribution from short-term variability and to identify when suitable
 85 habitats have expanded, contracted or shifted. Second, existing products generally
 86 represent marine distributions in two-dimensional geographic space or distinguish only
 87 broadly between surface and bottom environments. Yet the ocean is strongly vertically
 88 stratified: temperature, oxygen, salinity, light, productivity and pressure vary
 89 substantially through the water column, and both species composition and habitat
 90 suitability can differ markedly among depth layers (Duffy and Chown, 2017; Woolley
 91 et al., 2016; Woods et al., 2023; Duhamet et al., 2023). Models that associate
 92 observations collected at depth with surface conditions, or that collapse the water
 93 column into a small number of broad depth categories, may therefore mischaracterize
 94 species' environmental niches and obscure vertically structured habitat redistribution
 95 (Duffy and Chown, 2017; Robinson et al., 2017; Klaassen et al., 2025). Together,
 96 limited temporal coverage and coarse vertical representation constrain comprehensive
 97 assessments of marine habitat change across space, depth and time.

98 To address these limitations, we developed a multi-decadal, depth-resolved AOH
 99 dataset for global marine species. We integrated approximately 45.5 million occurrence
 100 records from the Global Marine Biodiversity Dataset (Dou, 2024) with physical and
 101 biogeochemical ocean reanalysis data spanning 1993–2025 from two Copernicus
 102 Marine Service products (Copernicus Marine Service, 2024a, 2024b). Environmental
 103 conditions were harmonized across 50 standard depth layers and used to fit species-
 104 specific MaxEnt models for 5,619 marine species. The fitted models were projected for
 105 seven target years (1995, 2000, 2005, 2010, 2015, 2020 and 2025), selected at regular
 106 five-year intervals to provide temporally consistent snapshots and to include 2020 as a
 107 policy-relevant reference point marking the endpoint of the Aichi Biodiversity Targets
 108 and the transition to the post-2020 global biodiversity framework. These projections
 109 generated species-level estimates of range-constrained environmental suitability across
 110 geographic space, depth and time. We assessed the resulting dataset using
 111 complementary internal consistency checks at the species and depth-layer levels



112 together with independent comparisons against global coral-reef distributions and
 113 scientific bottom-trawl observations. By integrating long-term temporal coverage with
 114 explicit vertical resolution, this dataset provides a standardized resource for
 115 investigating marine habitat redistribution, examining depth-dependent changes in
 116 habitat suitability, identifying persistent areas of high suitability and potential climate
 117 refugia, and evaluating the correspondence between suitable habitats and marine
 118 protected areas in a changing ocean.


119 2. Methods

120 2.1 Overall workflow

121 We developed an integrated workflow to generate a multi-decadal, depth-resolved
 122 Area-of-Habitat (AOH) dataset for global marine species (Fig. 1). The workflow
 123 combined species occurrence records, IUCN expert range maps, and depth-resolved
 124 physical and biogeochemical ocean variables to estimate species-specific
 125 environmental suitability across geographic space, depth and time. It consisted of three
 126 main stages: preparation and harmonization of biological and environmental inputs;
 127 species-specific distribution modelling and spatiotemporal projection; and construction,
 128 aggregation and evaluation of the resulting AOH products. First, species occurrence
 129 records, IUCN range maps and depth-resolved environmental variables were compiled,
 130 screened and harmonized to a common spatial, temporal and vertical framework.
 131 Second, independent MaxEnt models were fitted for individual species using
 132 occurrence records together with depth-specific, sampling-bias-adjusted background
 133 points, and the fitted models were projected across seven target years and the depth
 134 layers represented by each species. Third, the continuous suitability predictions were
 135 constrained to the original IUCN ranges to generate species-level marine AOH layers.
 136 These layers were subsequently aggregated using area-weighted metrics to characterize
 137 spatial, vertical and temporal patterns, and their plausibility was assessed through
 138 internal consistency checks and independent external comparisons.



139



140

141 **Fig. 1. Workflow for generating the multi-decadal, depth-resolved marine Area-of-Habitat dataset.**

142 2.2 Species and environmental data

143 We first compiled accepted scientific names and species-specific depth limits for
144 marine species. Because the IUCN Red List covers marine species that have undergone
145 formal conservation assessment, the resulting species pool represents a taxonomically
146 incomplete subset of described marine biodiversity rather than a comprehensive
147 inventory of all marine species. In this study, “global marine species” therefore refers



148 to the worldwide spatial coverage of the IUCN-assessed marine taxa included in the
149 analysis, subject to the availability of usable depth, occurrence and range information.
150 Upper and lower depth limits were available for 16,445 species.

151 We then obtained approximately 140 million raw marine species occurrence
152 records from the Global Marine Biodiversity Dataset (Dou, 2024), released by the
153 Oceanographic Data Center, Chinese Academy of Sciences. This point-based dataset
154 integrates occurrence records originally sourced from the Ocean Biodiversity
155 Information System (OBIS) and the Global Biodiversity Information Facility (GBIF),
156 provides environmental attributes matched to the coordinates, depths and collection
157 times of individual records, and includes species names matched to the accepted names
158 provided by IUCN. Records were screened for valid scientific names, geographic
159 coordinates, depths and collection years. Where an observed depth was unavailable, it
160 was assigned using the median of the species' IUCN depth range; records lacking both
161 observed and IUCN-derived depth information were excluded. Following this initial
162 screening and matching with the IUCN species list, approximately 45.5 million
163 occurrence records representing 15,128 species remained, while 1,317 IUCN species
164 without usable occurrence records were excluded. To reduce the risk of unstable models
165 under our depth-stratified, density-weighted background-sampling design, we required
166 a minimum of 15 valid occurrence records per species for species-level MaxEnt model
167 fitting, consistent with the threshold adopted in comparable marine MaxEnt
168 applications (La Manna et al., 2024). Applying this criterion retained 5,673 species for
169 model training (Section 2.3).

170 Pairwise correlations among the 11 candidate predictors were evaluated using the
171 harmonized, spatially continuous CMEMS environmental grids (Copernicus Marine
172 Service, 2024a, 2024b). These predictors represent environmental variables commonly
173 used to characterize marine species distributions, including temperature, salinity, ocean
174 currents, oxygen conditions, nutrient availability, acidity and primary productivity
175 (Tyberghein et al., 2012; Assis et al., 2018; Pickens et al., 2021). The 11 variables



176 included seawater temperature, salinity, eastward current velocity, northward current
 177 velocity, chlorophyll-a concentration, nitrate concentration, phosphate concentration,
 178 silicate concentration, dissolved oxygen concentration, pH and net primary production.
 179 Pairwise Pearson correlations were calculated to assess collinearity, with $|r| > 0.8$
 180 considered indicative of strong correlation. Based on the observed correlation structure
 181 across representative depth layers, together with ecological relevance, we retained
 182 seawater temperature, salinity, chlorophyll-a concentration, nitrate concentration and
 183 dissolved oxygen concentration. Although temperature and dissolved oxygen were
 184 strongly correlated in some upper-ocean layers, both variables were retained because
 185 they represent distinct thermal and aerobic constraints on marine species distributions,
 186 and their correlation varied substantially with depth. Their selection was also supported
 187 by their established relevance to marine species distributions. Temperature is a major
 188 environmental driver shaping the distribution and diversity patterns of marine
 189 organisms (Tittensor et al., 2010), while dissolved oxygen constrains their aerobic
 190 habitats (Deutsch et al., 2020). Salinity and nitrate reflect hydrographic and resource
 191 gradients and have been identified as important predictors of marine species
 192 distributions, although their relative importance varies among taxonomic groups and
 193 depth zones (Egorova et al., 2025). Chlorophyll-a provides an indicator of
 194 phytoplankton biomass and associated resource availability and has contributed
 195 substantially to explaining the spatiotemporal distributions of pelagic fishes (Wang et
 196 al., 2018).

197 The selected predictor layers were obtained from the Copernicus Marine Service
 198 (CMEMS) for model calibration and spatiotemporal projection (Copernicus Marine
 199 Service, 2024a, 2024b). Monthly data covering 1993–2025 were aggregated into annual
 200 means. The physical variables were provided at a native horizontal resolution of
 201 approximately 0.083° ($1/12^\circ$), whereas the biogeochemical variables were provided at
 202 0.25° . To establish a common horizontal grid, the biogeochemical variables were
 203 resampled to the 0.083° physical grid using nearest-neighbour interpolation. Seawater



204 temperature and salinity were provided across 50 standard depth layers, whereas
 205 chlorophyll-a concentration, nitrate concentration and dissolved oxygen concentration
 206 were originally provided across 75 depth layers. The biogeochemical variables were
 207 also aligned with the 50 physical depth layers using nearest-neighbour interpolation.
 208 The processed environmental layers were used to extract environmental values for
 209 occurrence and background records and to generate the temporal and depth-resolved
 210 model projections described in Section 2.3. Species range polygons were obtained from
 211 the IUCN Red List spatial database and used to screen occurrence records and constrain
 212 the spatial extent of the species-level AOH estimates. We retained polygons with
 213 presence classified as Extant, Probably Extant, or Possibly Extant and origin classified
 214 as Native, Reintroduced, or Introduced. Multiple polygons belonging to the same
 215 species were merged by scientific name. To exclude geographically distant occurrence
 216 records while retaining observations near mapped range boundaries, each species' range
 217 was buffered outward by approximately 500 km. Occurrence records were intersected
 218 with the corresponding buffered ranges, and only records located within these areas
 219 were retained for model fitting.

220 The labels L1–L50 denote the 50 physical-product depth coordinates, from
 221 approximately 0.5 to 5,727.9 m, rather than 50 equal-thickness depth intervals. This
 222 retains the native vertical framework of the CMEMS physical reanalysis (Copernicus
 223 Marine Service, 2024b). For Figures 2–4, eight illustrative levels were used: L1, L2,
 224 L5, L10, L15, L20, L30 and L40 (0.5, 1.5, 5.1, 13.5, 29.4, 65.8, 380.2 and 1,941.9 m).
 225 This subset provides a compact view of densely resolved shallow waters and
 226 progressively deeper levels; it is a visualization subset, not a separate ecological
 227 zonation or a statistically optimized selection. Modelling retained the full 50-level
 228 framework, subject to species-level data availability.

229 The approximately 500-km range buffer was a pragmatic tolerance for screening
 230 occurrences near expert-mapped range boundaries, while excluding geographically
 231 remote records. A 500-km buffering scale has also been used in marine SDMs, for



example around occurrence-derived domains in clownfishes (García-Jiménez et al., 2025). That application differs from buffering IUCN polygons and provides a methodological precedent rather than evidence that 500 km is optimal here. The buffer does not represent a common dispersal distance across taxa, and it does not expand the final AOH maps, which remain restricted to the original IUCN polygons. Alternative buffer distances were not evaluated in this study.

2.3 Species distribution modelling and temporal projection

We fitted an independent MaxEnt model for each species using the maxnet package in R (Phillips et al., 2017). MaxEnt estimates environmental suitability by contrasting the environmental conditions associated with species occurrences against those available at a set of background locations, making it suitable for presence-only occurrence data for which systematically collected absence records are unavailable (Phillips et al., 2006). Model calibration was based on the five environmental predictors retained in Section 2.2. Separate models were fitted for individual species so that each model characterized the environmental conditions associated with that species rather than pooling environmental responses across taxa. Feature classes were selected using the default sample-size-dependent settings of maxnet, and models were fitted using the default regularization multiplier (regmult = 1) (Phillips, 2021).

To reduce the influence of spatial sampling bias, we adapted the target-group background approach to depth-resolved marine occurrence data. Spatially biased occurrence records combined with uniformly sampled background points can cause presence-background models to reproduce geographic variation in sampling intensity rather than the environmental conditions associated with species occurrences. Drawing background points according to the sampling distribution of the broader target group can reduce this effect (Phillips et al., 2009; Barber et al., 2022). We therefore pooled the occurrence records of all included species and constructed a kernel-density surface for each of the 50 standard depth layers using a bandwidth of 50 km and a spatial resolution of 0.083°. These depth-specific surfaces represented the spatial distribution



260 of marine sampling within each layer and were used as probability weights for
261 background selection. For each species, background points were drawn only from the
262 depth layers represented by its occurrence records, and the total number of background
263 points was distributed among these layers in proportion to the number of occurrence
264 records in each layer. Locations were then sampled using the corresponding kernel-
265 density values as weights. A maximum of 20,000 background points was used when
266 more than 40,000 valid candidate cells were available; otherwise, half of the valid
267 candidate cells were sampled. Candidate points lacking complete values for any of the
268 five predictors were excluded.

269 The 50-km bandwidth is a fixed smoothing scale for the pooled sampling-effort
270 proxy, rather than a species-specific movement or habitat scale. It was used to reduce
271 highly localized sampling peaks while retaining broad sampling gradients and was not
272 optimized separately for each taxon or depth. The density surfaces were generated on
273 the same 0.083° grid used for the harmonized environmental predictors.

274 Occurrence and background records were matched to the environmental data
275 according to their geographic coordinates, collection year and assigned depth. The
276 nearest available annual environmental layer and standard depth layer were identified
277 for each record, from which values of the five retained predictors were extracted.
278 Occurrence locations were designated as presences and the density-weighted sampled
279 locations as background records, forming a separate calibration dataset for each species.
280 Records with incomplete predictor values were removed before model fitting. Of the
281 5,673 species meeting the minimum occurrence threshold, independent MaxEnt models
282 were fitted successfully for 5,619 species; species for which model fitting failed were
283 excluded from subsequent processing. Predictions were generated using the cloglog
284 output format implemented in maxnet, producing continuous suitability values ranging
285 from 0 to 1 (Phillips et al., 2017). These values represent relative environmental
286 suitability under the fitted model and should not be interpreted directly as occurrence
287 probability or observed occupancy.



Each fitted model was projected onto environmental conditions for seven target years: 1995, 2000, 2005, 2010, 2015, 2020 and 2025. Predictions were generated separately for each standard depth layer represented by the species, thereby preserving variation in environmental suitability across both time and depth. Predictor values outside the ranges encountered during model calibration were clamped to the corresponding calibration ranges using the default prediction settings of maxnet, thereby reducing unsupported extrapolation into environmental conditions not represented by the training data (Stohlgren et al., 2011). For each species, the environmental grids and resulting suitability estimates were spatially constrained using the original, unbuffered IUCN range polygons; predictions outside the mapped range were excluded. The resulting continuous suitability estimates were stored separately for each species, target year and depth layer and were subsequently used to construct the marine AOH dataset described in Section 2.4.

2.4 Construction and aggregation of the marine AOH dataset

Conventional terrestrial AOH mapping commonly refines a species' range using habitat and elevation associations. Because the present analysis did not apply a separate marine habitat-type layer, we operationally defined marine AOH as the environmentally suitable portion of the original IUCN range. The resulting values should therefore be interpreted as range-constrained environmental suitability rather than as direct estimates produced by conventional terrestrial habitat and depth filtering.

$$AOH_{s,i,l,t} = R_{s,i} S_{s,i,l,t} \quad (1)$$

Here, $S_{s,i,l,t}$ is the predicted suitability of species s in grid cell i , depth layer l and year t , and $R_{s,i}$ is the species-specific IUCN range mask, taking a value of 1 inside the original IUCN range and 0 outside it. Applying this mask retained the continuous suitability estimates within the species' range and excluded predictions outside it. Buffered ranges were used only to screen occurrence records and define the modelling domain; the final suitability estimates were constrained using the original IUCN polygons.



Species-level AOH layers were stored separately for each target year and depth layer. To summarize the joint distribution of modelled species, continuous AOH values were summed cell by cell:

$$SS_{i,l,t} = \sum_{s \in S_l} AOH_{s,i,l,t} \quad (2)$$

where $SS_{i,l,t}$ is the summed suitability in grid cell i , depth layer l and year t , and S_l is the set of modelled species represented in that layer. Continuous values were retained during aggregation to avoid introducing an arbitrary binary suitability threshold (Calabrese et al., 2014; Grenié et al., 2020). Accordingly, summed suitability represents the cumulative modelled suitability of the included species and should not be interpreted as directly observed species richness.

Because grid-cell area varies with latitude, layer-level totals were calculated using cell-area weights:

$$T_{l,t} = \sum_{i \in \Omega_{l,t}} A_i SS_{i,l,t} \quad (3)$$

where $T_{l,t}$ is the area-weighted summed suitability for depth layer l and year t , A_i is the area of grid cell i , and $\Omega_{l,t}$ is the set of cells with valid values in that layer and year. We subsequently summarized spatial and temporal variation using the percentage change relative to 2020, the proportion of the layer-level total occurring in tropical waters, the contribution of the highest-valued cells occupying 5% of valid area, the total valid area and the maximum cell value. The temporal trajectories in Fig. 4 use area-weighted means over valid cells, expressed relative to 2020.

Temporal change was evaluated across the seven target years. For each depth layer, an ordinary least-squares regression was fitted to the area-weighted totals. Cell-level trends were estimated only for grid cells with complete values across all target years, and the area-weighted proportion of cells with positive slopes was calculated for each depth layer. The outcomes of these analyses and the assessments of dataset plausibility are reported in the Results section. For interactive exploration, the aggregated layers



339 can be visualized by target year and depth layer through the Marine AOH Viewer
 340 provided in the Data availability section.

341 **2.5 Internal consistency and external validation**

342 Because globally consistent independent observations are scarce for a dynamic,
 343 depth-resolved marine AOH dataset, we assessed the plausibility of the 2020
 344 predictions through two complementary internal consistency checks and two
 345 comparisons with independent external datasets. The internal checks followed the
 346 point-prevalence-versus-model-prevalence framework proposed by Dahal et al. (2022),
 347 which evaluates whether predicted suitable conditions are more strongly concentrated
 348 at species occurrence locations than across the corresponding AOH rasters. Because the
 349 occurrence records used in these assessments were also used during model calibration,
 350 these analyses were interpreted as internal consistency checks rather than independent
 351 tests of predictive performance.

352 Framework A was conducted at the species level. For each species, occurrence
 353 points were pooled across all represented depth layers, and the corresponding cloglog
 354 suitability values were extracted from the 2020 predictions. To characterize suitability
 355 across the modelled range, we randomly sampled up to 1,000 valid AOH pixels from
 356 each represented depth layer of the species' IUCN-range-masked rasters. Framework B
 357 was conducted at the depth-layer level. For each of the 50 standard depth layers,
 358 suitability values were pooled across the occurrence points of all species represented in
 359 that layer, together with up to 10,000 valid AOH pixels sampled from each contributing
 360 species' raster. In both frameworks, AOH pixels were restricted to non-missing cells
 361 within the original IUCN range of the corresponding species.

362 We evaluated five fixed suitability thresholds, τ , ranging from 0.1 to 0.5 at
 363 intervals of 0.1. At each threshold, point prevalence was calculated as

$$P_{point}(\tau) = \frac{n(\text{occurrence points}, \hat{S} \geq \tau)}{n(\text{occurrence points})} \quad (4)$$



364 where n (occurrence points, $\hat{S} \geq \tau$) is the number of occurrence points with
 365 predicted suitability $\hat{S}$ at or above threshold τ , and $n(\text{occurrence points})$ is the total
 366 number of occurrence points evaluated. Model prevalence was calculated as

$$P_{\text{model}}(\tau) = \frac{n(\text{AOH pixels}, \hat{S} \geq \tau)}{n(\text{AOH pixels})} \quad (5)$$

367 where $n(\text{AOH pixels}, \hat{S} \geq \tau)$ is the number of sampled valid AOH pixels with
 368 predicted suitability at or above the threshold, and $n(\text{AOH pixels})$ is the total number
 369 of sampled valid AOH pixels evaluated within the species' original IUCN range. A
 370 species or depth layer was considered to pass the threshold-based assessment when
 371 $P_{\text{point}}(\tau) > P_{\text{model}}(\tau)$, indicating that occurrence locations exceeded the threshold
 372 more frequently than expected from the corresponding proportion of the modelled AOH.
 373 We additionally compared mean suitability at occurrence points with mean suitability
 374 across sampled AOH pixels to evaluate the same relationship without converting the
 375 continuous predictions into binary suitable and unsuitable areas. Extracted values were
 376 retained so that all thresholds and continuous summaries could be calculated from the
 377 same occurrence-point and AOH-pixel samples.

378 The two external datasets were selected to evaluate complementary aspects of
 379 model plausibility. FishGlob provides species-level, station-based scientific bottom-
 380 trawl observations with associated year and depth information (Maureaud et al., 2024),
 381 enabling comparison of predicted suitability between detections and non-detections. In
 382 contrast, the UNEP-WCMC global coral-reef dataset provides an independently
 383 mapped distribution of reef habitat (UNEP-WCMC et al., 2021), allowing broad-scale
 384 spatial correspondence of shallow-water predictions to be assessed.

385 The first external comparison evaluated whether predicted suitability for reef-
 386 building corals corresponded with independently mapped coral reef locations. Of 372
 387 species identified from the IUCN REEF_FORMING_CORALS group, 272 had 2020
 388 predictions for all five shallow layers (L1–L5; representative depths 0.5, 1.5, 2.6, 3.8
 389 and 5.1 m). Continuous predictions were summed across these species within each layer



390 and then averaged equally across the five layers, without applying a species-level binary
 391 suitability threshold. The common analysis domain retained ocean cells with consistent
 392 valid-prediction support across layers for each species; cells outside all species'
 393 prediction support retained the original aggregate-zero background convention. Grid
 394 cells intersecting UNEP-WCMC reef polygons (UNEP-WCMC et al., 2021) were
 395 classified as reef cells. Reef and non-reef values were compared using a one-sided
 396 Mann–Whitney U test, sampling at most 200,000 cells per group (seed 42); descriptive
 397 metrics used all eligible cells. Spatial agreement was summarized using the inside-to-
 398 outside mean ratio and reef enrichment among cells at or above the 75th percentile of
 399 positive mean summed suitability, relative to the reef-cell fraction in the corresponding
 400 domain. These comparisons were repeated within the Great Barrier Reef, Coral
 401 Triangle and Caribbean, with percentile cutoffs calculated separately for each domain.
 402 The analysis evaluated horizontal spatial correspondence without matching reef
 403 locations to actual bathymetric depths.

404 The second external comparison used FishGlob data (Maureaud et al., 2024),
 405 which integrates 29 scientific bottom-trawl surveys comprising 216,548 stations across
 406 the continental shelves and slopes of North America and Europe. Species captured at a
 407 station were treated as presences, whereas surveyed stations at which a species was not
 408 captured were treated as non-detections. Matching FishGlob data with the 5,619
 409 successfully modelled species yielded 1,053 candidate species. For each candidate
 410 species, stations within its surveyed range were assigned to presence and non-detection
 411 groups. When more than 3,000 non-detection stations were available, 3,000 stations
 412 were randomly selected. Station coordinates, survey years and sampling depths were
 413 matched to the nearest target year and standard prediction depth layer. Stations outside
 414 the original IUCN range or lacking a valid prediction were excluded, and only species
 415 with at least 10 valid stations in both groups were retained. For each retained species,
 416 we compared mean continuous suitability between presence and non-detection stations.
 417 A comparison was considered directionally consistent when mean predicted suitability



418 was higher at presence stations than at non-detection stations. Non-detections can
419 reflect imperfect catchability as well as unsuitable conditions.

420 **2.6 Assumptions and constraints**

421 First, species–environment relationships fitted to the pooled calibration records
422 are assumed to be transferable among the seven target years and represented depth
423 levels. Annual means, nearest-year/depth matching and imputed depths simplify
424 seasonal variability, vertical movements and observation uncertainty. Predictions
425 therefore describe environmental suitability under these assumptions, rather than
426 observed occupancy or abundance.

427 Second, static IUCN polygons delimit all final predictions, so suitable conditions
428 outside these boundaries cannot be recovered by this product. Density-weighted
429 backgrounds approximate sampling effort but cannot fully correct taxonomic,
430 geographic or temporal biases. The fixed bandwidth and buffer settings were not
431 optimized across taxa. Environmental predictors do not explicitly represent substrate or
432 biotic interactions.


433 Third, comparisons among depths depend on the species and valid cells
434 represented in each layer. Summed suitability is not observed species richness, and an
435 area-weighted mean differs from an area-integrated total when the valid area changes.
436 Internal checks reuse calibration occurrences; reef comparisons assess horizontal
437 habitat correspondence; and trawl non-detections do not establish biological absence.
438 External data provenance alone does not establish that every validation record was
439 independent of the OBIS/GBIF-derived calibration pool because no record-level
440 overlap audit was performed. These assumptions and constraints should be considered
441 when interpreting the results and validation outcomes.



442 **3. Results**

443 **3.1 Global depth-stratified patterns of AOH**

444 Aggregated AOH showed pronounced vertical differentiation across the eight
445 representative depth layers examined for 2020 (selected from L1–L40, approximately
446 0.5–1,942 m; Fig. 2). Shallow-water layers were characterized by strong spatial
447 concentration and a predominance of AOH in tropical waters, whereas deeper layers
448 exhibited progressively more diffuse spatial patterns. From L2 to L15 (approximately
449 1.5–29 m), more than half of the area-weighted global AOH total occurred within the
450 tropical band (23.5°S–23.5°N), and the highest-valued grid cells occupying 5% of valid
451 area accounted for well over half of the corresponding layer-level total. From L20
452 downward (approximately 66–1,942 m), both tropical dominance and spatial
453 concentration declined markedly and subsequently remained comparatively stable. The
454 area-weighted total AOH also decreased substantially with depth, declining by
455 approximately 79% from L1 to L40. In contrast, the proportion of grid cells retaining
456 valid predictions decreased more gradually, from just over 70% at L1 to approximately
457 59% at L40. The shallowest layer, L1, differed from the immediately deeper shallow
458 layers by exhibiting lower tropical dominance and weaker spatial concentration.
459 Together, these patterns reveal a clear transition from highly concentrated shallow-
460 water AOH to more spatially diffuse distributions at greater depths.



461

462 **Fig. 2. Global depth-resolved patterns of aggregated marine AOH in 2020.** The accompanying
 463 latitudinal profiles show the mean $\pm$ SD of aggregated AOH across longitudes at each latitude. Aggregated
 464 AOH represents summed modelled environmental suitability across species. (a–h) Maps and latitude profiles
 465 for the eight representative layers. (i) Area-integrated summed suitability. (j) Contributions of the tropical
 466 band and the highest-valued cells occupying 5% of valid area to the corresponding layer total.




3.2 Temporal changes in AOH during 1995-2025

Temporal trends in aggregated AOH varied markedly among depth layers between 1995 and 2025 (Fig. 3). Pixel-wise linear trends were predominantly negative from L1 to L30, with the strongest overall decline occurring at L5 and a comparatively weaker negative trend at L20. L40 was the only representative depth layer showing a positive net trend, with increasing AOH over more than half of its valid area. At L30, negative trends were concentrated primarily in the mid-latitudes of both hemispheres, particularly in the Southern Hemisphere, whereas trends were close to zero within approximately 23.5° of the equator and became positive poleward of 50°. Prediction coverage at L50 was insufficient to support reliable trend estimation.


Layer-level time series showed similarly depth-dependent temporal trajectories (Fig. 4). The five shallowest representative layers (L1, L2, L5, L10 and L15; approximately 0.5–29 m) generally reached their highest area-weighted mean AOH around 2000 and subsequently declined toward 2020, although the magnitude of change varied among layers. L20 fluctuated within a comparatively narrow range throughout the study period. In contrast, L30 exhibited a persistent decline across the seven target years and remained below its earlier levels in 2025, whereas L40 increased consistently from 1995 to 2025. These contrasting trajectories indicate that temporal changes in marine AOH were strongly depth dependent, with shallow and intermediate waters generally declining while deeper layers showed distinct or opposing patterns.

The increasing trajectory at L40 represents a modelled change in environmental suitability at approximately 1,941.9 m, rather than a measured increase in deep-sea abundance or direct evidence of downward migration. The post-2020 comparison contains only the 2020 and 2025 target years and therefore represents a five-year endpoint difference rather than a continuously observed annual trend.





493 **Fig. 3. Pixel-wise linear trends in aggregated marine AOH across eight representative depth layers during**
494 **1995–2025.** Linear trends were estimated for each grid cell using aggregated AOH values from seven target
495 years (1995, 2000, 2005, 2010, 2015, 2020 and 2025), and only grid cells with valid predictions for all seven
496 years were included. The accompanying latitudinal profiles show the mean $\pm$ SD of pixel-wise slopes across
497 longitudes at each latitude. Panels (a–h) show the maps and profiles; panel (i) shows the area-weighted
498 proportions of grid cells with decreasing, zero or increasing trends across the eight depth layers, with labels
499 indicating the proportion with positive slopes. Slope direction does not imply statistical significance.



500



501 **Fig. 4. Temporal trajectories of area-weighted mean marine AOH across eight representative depth**
502 **layers during 1995–2025.** For each depth layer, continuous species-level AOH values were summed across
503 species and then averaged over valid grid cells using grid-cell-area weights to obtain the layer-level mean for
504 each of the seven target years (1995, 2000, 2005, 2010, 2015, 2020 and 2025). In panel a, values are expressed
505 as percentage change relative to the corresponding 2020 mean, allowing temporal trajectories to be compared
506 among depth layers despite differences in absolute AOH magnitude. (b) Percentage change over each
507 consecutive five-year interval, relative to the earlier value. (c) Net percentage change from 1995 to 2025,
508 relative to 1995.
509



510 3.3 Validation

511 We evaluated the 2020 predictions at complementary biological and spatial scales.
 512 Following the point-prevalence-versus-model-prevalence framework of Dahal et al.
 513 (2022), internal consistency was examined first at the species level and then across
 514 depth layers. The predictions were subsequently compared with two independent
 515 external datasets representing reef-associated spatial patterns and station-level species
 516 observations.

517 3.3.1 Species-level internal consistency

518 Among the 4,996 species evaluated for 2020, occurrence-point prevalence
 519 exceeded model prevalence for the majority of species across all five suitability
 520 thresholds ($\tau = 0.1$ – 0.5 ; Fig. 5). The species-level pass rate declined progressively as
 521 the threshold became more restrictive, from 94.70% (4,731/4,996) at $\tau = 0.1$ to 94.38%,
 522 92.55%, 90.27% and 85.69% (4,281/4,996) at $\tau = 0.2, 0.3, 0.4$ and 0.5 , respectively. At
 523 the fixed threshold of $\tau = 0.5$, most species lay above the 1:1 line, indicating that
 524 suitability values exceeding the threshold occurred proportionally more often at known
 525 occurrence locations than across sampled pixels within the corresponding range-
 526 constrained AOH. The threshold-free comparison of continuous suitability showed the
 527 same overall pattern, with mean predicted suitability generally higher at occurrence
 528 points than across sampled AOH pixels. Only 132 species (2.6%) failed the prevalence
 529 comparison at all five thresholds, indicating that the observed relationship was robust
 530 to the choice of suitability threshold. Together, these results demonstrate strong internal
 531 consistency between species occurrences and the corresponding modelled suitability
 532 patterns. However, because the occurrence records used in these comparisons also
 533 contributed to model calibration, the analysis should be interpreted as an internal
 534 consistency assessment rather than as an independent evaluation of predictive
 535 performance.




Fig. 5. Depth-resolved internal consistency of marine AOH predictions in 2020. Mean predicted suitability (cloglog; mean $\pm$ SD, panel a) is shown for occurrence points and sampled AOH pixels across the 47 standard depth layers with sufficient data for evaluation. Occurrence-point suitability was extracted from the 2020 species-level predictions, whereas AOH-pixel suitability was sampled from valid cells within the corresponding original IUCN ranges. Panel b shows the species pass rate for each depth layer, defined as the proportion of constituent species for which occurrence-point prevalence exceeded model prevalence at the fixed suitability threshold of 0.5. Because the occurrence records used in this assessment also contributed to model calibration, the comparison represents an internal consistency check rather than an independent validation of predictive performance. Panel c shows the numbers of occurrence records and species represented at each depth.

3.3.2 Depth-layer-level internal consistency

Of the 50 standard depth layers, 47 contained sufficient data for evaluation. Across all 47 layers, mean predicted suitability at occurrence points was higher than that across




sampled AOH pixels, and this pattern persisted from shallow waters to the deepest
 evaluated layers (Fig. 5). Although both the magnitude and variability of predicted
 suitability differed among depths, the layer-specific species pass rate at the fixed
 threshold of $\tau = 0.5$ remained consistently high, averaging $91.95\% \pm 3.28\%$ and ranging
 from 86.78% to 100.00%. The lowest pass rate occurred at approximately 9.6 m,
 whereas the two deepest evaluated layers reached 100%. These results indicate that the
 concentration of higher predicted suitability at known occurrence locations was
 maintained across the water column and was not restricted to species-rich surface layers.
 As with the species-level assessment, however, the occurrence records used here also
 contributed to model calibration. The analysis should therefore be interpreted as a
 depth-resolved internal consistency assessment rather than as an independent validation
 of predictive performance.

3.3.3 External spatial validation using the UNEP-WCMC global coral reef distribution

For the 272 reef-building coral species evaluated using the 2020 L1–L5 average
 (representative depths 0.5–5.1 m), aggregated predicted suitability showed strong
 spatial correspondence with the independent UNEP-WCMC global coral reef
 distribution (Fig. 6). Mean shallow-layer summed suitability was 42.17 within mapped
 reef boundaries and 1.98 outside them, corresponding to a 21.33-fold difference. Areas
 of high predicted suitability were also disproportionately associated with mapped reefs:
 9.60% of grid cells at or above the 75th percentile of positive mean summed suitability
 overlapped reef boundaries, compared with a global spatial baseline of 0.3927%,
 representing a 24.45-fold enrichment. The same directional pattern was observed across
 three major coral-reef regions. Mean summed suitability was consistently higher within
 than outside mapped reef boundaries in the Great Barrier Reef, Coral Triangle and
 Caribbean, with inside-to-outside ratios of 3.12, 1.79 and 1.86, respectively. The
 agreement observed at both global and regional scales indicates that the aggregated



577 shallow-layer predictions recovered the spatial distribution of a well-established marine
578 habitat independently of the data used for model calibration.



579




580 **Fig. 6. External spatial validation of shallow-layer coral suitability against mapped reefs. Continuous**
 581 **2020 predictions for 272 reef-building coral species were summed within layers and averaged equally across**
 582 **L1–L5 (representative depths 0.5–5.1 m), without species-level binarization. (a) Global mean summed**
 583 **suitability and UNEP-WCMC reef cells. (b–d) Great Barrier Reef, Coral Triangle and Caribbean, using the**
 584 **same colour scale. Reef symbols aid visibility without enlarging the quantitative mask. (e) Cumulative**
 585 **suitability distributions inside and outside reef cells, including aggregate-zero ocean background. (f)**
 586 **Corresponding mean suitability. (g) Reef enrichment among cells at or above the 75th percentile of positive**
 587 **suitability, relative to each domain's reef-cell baseline. The common global domain comprised 24,114 reef**
 588 **and 6,115,775 non-reef cells. Grid cells are spatially dependent; this comparison does not validate actual**
 589 **vertical occupancy.**

590 3.3.4 External station-level validation using FishGlob trawl observations

591 FishGlob provided an independent station-level evaluation of species-specific
 592 AOH predictions using scientific bottom-trawl observations from continental shelf and
 593 slope ecosystems. Of the 1,053 FishGlob species shared with the modelled dataset, 459
 594 met the requirement of at least 10 valid presence stations and 10 valid non-detection
 595 stations after matching observations to the corresponding prediction year, depth layer
 596 and original IUCN range (Fig. 7). Species not retained generally lacked sufficient
 597 matched stations or valid predictions within the required spatiotemporal and geographic
 598 constraints. Across the 459 evaluated species, mean predicted suitability was higher at
 599 presence stations (0.315) than at non-detection stations (0.245). A total of 324 species
 600 (70.6%) lay above the 1:1 line, indicating higher mean predicted suitability at presence
 601 than at non-detection stations, whereas 135 species showed equal or lower suitability
 602 at presence stations. This predominance of species above the 1:1 line indicates that the
 603 model predictions generally discriminated observed presences from non-detections in
 604 an independent survey dataset. These species-level station contrasts cannot be ranked
 605 directly against reef-level enrichment, because the two comparisons differ in response
 606 variable, aggregation, spatial domain and observation process. Nevertheless, the overall
 607 directional consistency provides independent support for the ability of the species-level



608 predictions to distinguish between locations where species were observed and surveyed
 609 locations where they were not detected.



610

611 **Fig. 7. External validation of species-level AOH predictions using FishGlob scientific bottom-trawl**
 612 **observations.** The comparison included 459 species for which at least 10 valid presence stations and 10 valid
 613 non-detection stations could be matched to the corresponding prediction year, depth layer and original IUCN
 614 range. In panel a, for each species, the x-axis shows mean predicted suitability at non-detection stations and
 615 the y-axis shows mean predicted suitability at presence stations. The diagonal 1:1 line indicates equal mean
 616 suitability between the two groups; points above the line therefore represent species for which predicted
 617 suitability was higher at presence than at non-detection stations. A total of 324 species (70.6%) occurred
 618 above the 1:1 line. Point colour represents the difference between mean suitability at presence and non-
 619 detection stations, and point area increases with the logarithm of the total number of stations included for
 620 each species. (b) Distribution of species-level presence-minus-non-detection mean differences. (c) Mean
 621 differences in relation to the number of presence records per species. Non-detection does not establish
 622 biological absence.



623 4. Discussion

624 The principal contribution of this dataset is its representation of marine AOH
 625 across horizontal space, depth and multiple decades. By integrating species occurrence
 626 records, expert range constraints and depth-resolved environmental variables, this
 627 framework extends AOH-based biodiversity assessment into the marine environment
 628 and provides a consistent basis for examining habitat redistribution through three
 629 dimensions. The observed patterns highlight that marine AOH is strongly structured
 630 along the vertical axis, with different depth layers showing distinct spatial
 631 concentrations and temporal trajectories. Such depth-dependent patterns are consistent
 632 with the importance of explicitly representing depth in marine species-distribution
 633 modelling (Duffy and Chown, 2017), the pronounced bathymetric structuring and
 634 turnover documented for marine fishes (Duhamet et al., 2023), and evidence that
 635 climate velocities—and hence the expected pressures for species redistribution—differ
 636 substantially among ocean depth zones (Brito-Morales et al., 2020). However, these
 637 results represent changes in modelled environmental suitability rather than direct
 638 changes in species richness, occupancy, abundance or movement; therefore, changes in
 639 AOH should not be interpreted as direct evidence of population redistribution.

640 The validation analyses provide complementary evidence that the dataset captures
 641 meaningful ecological patterns. Internal assessments showed that predicted suitability
 642 was generally higher at occurrence locations than across the corresponding range-
 643 constrained prediction areas, although these evaluations represent consistency checks
 644 because the same occurrence records were used for model calibration. External
 645 comparisons further supported the spatial realism of the predictions: predicted
 646 suitability for reef-building corals was strongly concentrated within mapped reef areas,
 647 and most FishGlob-tested species showed higher mean suitability at observed-presence
 648 stations than at observed-absence stations. These two external assessments evaluate
 649 different properties of the predictions: the coral-reef comparison examines broad-scale
 650 spatial correspondence with an independently mapped habitat, whereas the FishGlob



651 comparison tests discrimination between observed-presence and observed-absence
 652 stations. Their results are therefore not expected to be directly equivalent, because
 653 different evaluation approaches may target different properties of model performance
 654 and differ in their degree of independence from the calibration data (Araújo et al., 2019).
 655 The FishGlob comparison may additionally be influenced by variation among surveys
 656 in sampling design, spatiotemporal coverage, and gear- and species-specific
 657 catchability because the integrated dataset combines multiple scientific bottom-trawl
 658 surveys and primarily samples fish communities associated with the seafloor
 659 (Maureaud et al., 2024). Together, these assessments support the comparative use of
 660 the dataset while highlighting uncertainty associated with taxa, depth layers and
 661 observation methods.

662 Reef-building corals are sessile after settlement, so mapped reef habitat provides
 663 a relatively persistent spatial reference compared with mobile fish detections.
 664 Nevertheless, reef extent and condition can change, and mapped reef habitat does not
 665 demonstrate the presence of each coral species.

666 The dataset should be interpreted according to its operational definition of marine
 667 AOH. Conventional terrestrial AOH is defined as suitable habitat within a species’
 668 geographic range and is commonly mapped by refining IUCN range maps using habitat
 669 associations and elevational limits (Brooks et al., 2019; Lumbierres et al., 2022). In this
 670 dataset, this concept is operationally extended to the marine environment by defining
 671 species-level marine AOH as continuous environmental suitability constrained by the
 672 original IUCN range. Species-level layers therefore represent range-constrained
 673 suitability estimates rather than binary occupied areas, and aggregated summed
 674 suitability should not be interpreted as equivalent to observed species richness.
 675 Taxonomic coverage was restricted to marine taxa assessed by the IUCN Red List and
 676 was further limited by the availability of usable depth, occurrence and range data and
 677 successful model fitting. The resulting dataset should therefore not be interpreted as a
 678 taxonomically complete representation of global marine biodiversity. Uneven



679 occurrence sampling and the omission of taxon-specific ecological predictors are
 680 widely recognized sources of uncertainty in marine SDMs (Robinson et al., 2017;
 681 Pickens et al., 2021; Klaassen et al., 2025). Additional dataset-specific uncertainty
 682 arises from assigning missing observation depths using available species-level depth
 683 information, applying static expert range polygons, using annually averaged
 684 environmental conditions and excluding some ecological constraints, such as substrate
 685 characteristics, prey availability and biotic interactions. Depth-specific target-group
 686 background sampling can reduce the influence of spatially uneven sampling effort
 687 (Phillips et al., 2009; Barber et al., 2022), while geographic range constraints and
 688 prediction clamping limit geographically implausible projections and unsupported
 689 environmental extrapolation (Elith et al., 2010; Stohlgren et al., 2011). Nevertheless,
 690 these procedures cannot completely eliminate uncertainty associated with the
 691 occurrence data, environmental predictors and modelling assumptions.

692 Within these limitations, this dataset provides a standardized framework for
 693 comparing marine habitat suitability across species, depth layers and time periods.
 694 Species-level predictions can support assessments of changes in suitable environmental
 695 conditions within mapped ranges, while aggregated products enable analyses of
 696 vertically differentiated habitat patterns, persistent suitability hotspots and potential
 697 climate refugia. The dataset may also support spatial comparisons between suitable
 698 marine habitats, potential climate refugia and existing marine protected areas (Brito-
 699 Morales et al., 2020; Zhuang et al., 2025). Its depth-resolved structure further provides
 700 an opportunity to evaluate whether protected areas encompass suitable environmental
 701 conditions across the water column rather than only at the surface. Future applications
 702 should integrate species-specific ecological knowledge, local observations and explicit
 703 uncertainty assessments to improve the interpretation of marine biodiversity change.

704 **5. Conclusion**

705 We developed a multi-decadal, depth-resolved marine Area-of-Habitat (AOH)
 706 dataset for 5,619 species across 50 standard depth layers. Using environmental



707 reanalysis data spanning 1993–2025, we generated AOH for seven discrete target years
 708 (1995, 2000, 2005, 2010, 2015, 2020, and 2025) to capture temporal variation at 5-year
 709 intervals. By integrating species occurrence records, IUCN expert range maps, and
 710 vertically resolved physical and biogeochemical ocean variables within a consistent
 711 modelling framework, the dataset provides range-constrained estimates of
 712 environmental suitability across geographic space, ocean depth, and time. Its principal
 713 advance lies in explicitly incorporating the vertical dimension into long-term, global
 714 marine habitat mapping, thereby extending existing approaches beyond predominantly
 715 static or surface-oriented representations. The contrasting spatial patterns and temporal
 716 trajectories observed among depth layers further demonstrate the importance of
 717 accounting for depth when assessing large-scale changes in marine habitat suitability.
 718 Complementary internal consistency assessments and independent external
 719 comparisons support the ecological plausibility of the resulting predictions.

720 The dataset provides a standardized basis for comparing suitable environmental
 721 conditions among species, regions, depth zones, and time periods. Species-level layers
 722 enable changes in habitat suitability to be examined for individual species across years
 723 and depths, whereas aggregated products facilitate analyses of broader spatial patterns,
 724 vertically structured suitability hotspots, and potential climate refugia. The dataset may
 725 also support assessments of how suitable marine habitats overlap with existing marine
 726 protected areas across the water column. As improved occurrence records, ocean
 727 reanalysis products, and species-specific ecological information become available, the
 728 framework can be updated and refined accordingly. The dataset should nevertheless be
 729 interpreted as continuous, range-constrained environmental suitability rather than as
 730 direct observations of species occupancy, abundance, or richness.

731 6. Acknowledgements

732 This work was supported by the National Key R&D Program of China
 733 (2024YFF1307600), the National Natural Science Foundation of China (42401314),
 734 and the Technology Innovation Center for South China Sea Remote Sensing, Surveying



735 and Mapping Collaborative Application, Ministry of Natural Resources (RSSMCA-
736 2025-B006).

737 **7. Data availability**

738 The data for this manuscript are available at
739 <https://doi.org/10.5281/zenodo.21933262> (Yu and Li, 2026). An interactive
740 visualization of the aggregated marine AOH layers is available through the Marine
741 AOH Viewer at <https://zzssjj.projects.earthengine.app/view/marine-aoh-viewer>, where
742 users can explore the dataset by target year and standard depth layer. The application
743 provides a visualization interface, whereas the complete species-level and aggregated
744 data files are archived on Zenodo.

745

746



References

- Araújo, M. B., Anderson, R. P., Márcia Barbosa, A., Beale, C. M., Dormann, C. F., Early, R., Garcia, R. A., Guisan, A., Maiorano, L., Naimi, B., O'Hara, R. B., Zimmermann, N. E., and Rahbek, C.: Standards for distribution models in biodiversity assessments, *Science Advances*, 5, <https://doi.org/10.1126/sciadv.aat4858>, 2019.
- Assis, J., Tyberghein, L., Bosch, S., Verbruggen, H., Serrão, E. A., and De Clerck, O.: Bio-ORACLE v2.0: Extending marine data layers for bioclimatic modelling, *Global Ecology and Biogeography*, 27, 277–284, <https://doi.org/10.1111/geb.12693>, 2018.
- Barber, R. A., Ball, S. G., Morris, R. K. A., and Gilbert, F.: Target-group backgrounds prove effective at correcting sampling bias in Maxent models, *Diversity and Distributions*, 28, 128–141, <https://doi.org/10.1111/ddi.13442>, 2022.
- Brito-Morales, I., Schoeman, D. S., Molinos, J. G., Burrows, M. T., Klein, C. J., Arafah-Dalmau, N., Kaschner, K., Garilao, C., Kesner-Reyes, K., and Richardson, A. J.: Climate velocity reveals increasing exposure of deep-ocean biodiversity to future warming, *Nature Climate Change*, 10, 576–581, <https://doi.org/10.1038/s41558-020-0773-5>, 2020.
- Brooks, T. M., Pimm, S. L., Akçakaya, H. R., Buchanan, G. M., Butchart, S. H. M., Foden, W., Hilton-Taylor, C., Hoffmann, M., Jenkins, C. N., Joppa, L., Li, B. V., Menon, V., Ocampo-Peñuela, N., and Rondinini, C.: Measuring terrestrial area of habitat (AOH) and its utility for the IUCN red list, *Trends in Ecology & Evolution*, 34, 977–986, <https://doi.org/10.1016/j.tree.2019.06.009>, 2019.
- Calabrese, J. M., Certain, G., Kraan, C., and Dormann, C. F.: Stacking species distribution models and adjusting bias by linking them to macroecological models, *Global Ecology and Biogeography*, 23, 99–112, <https://doi.org/10.1111/geb.12102>, 2014.
- Comte, L. and Olden, J. D.: Climatic vulnerability of the world's freshwater and marine fishes, *Nature Climate Change*, 7, 718–722, <https://doi.org/10.1038/nclimate3382>, 2017.
- Convention on Biological Diversity: Kunming-Montreal Global Biodiversity Framework, 2022.
- Copernicus Marine Service: Global Ocean Biogeochemistry Hindcast, https://data.marine.copernicus.eu/product/GLOBAL_MULTIYEAR_BGC_001_029/description, last access: 21 August 2026, 2024a.
- Copernicus Marine Service: Global Ocean Physics Reanalysis, https://data.marine.copernicus.eu/product/GLOBAL_MULTIYEAR_PHY_001_030/description, last access: 21 August 2026, 2024b.
- Dahal, P. R., Lumbierres, M., Butchart, S. H. M., Donald, P. F., and Rondinini, C.: A validation standard for area of habitat maps for terrestrial birds and mammals,



- 787 Geoscientific Model Development, 15, 5093–5105, [https://doi.org/10.5194/gmd-](https://doi.org/10.5194/gmd-15-5093-2022)
 788 15-5093-2022, 2022.
- 789 Deutsch, C., Penn, J. L., and Seibel, B.: Metabolic trait diversity shapes marine
 790 biogeography, *Nature*, 585, 557–562, <https://doi.org/10.1038/s41586-020-2721-y>,
 791 2020.
- 792 Dou, F.: Global Marine Biodiversity Dataset with Environmental Elements,
 793 Oceanographic Data Center, Chinese Academy of Sciences [data set],
 794 [https://msdc.qdio.ac.cn/data/metadata-special-](https://msdc.qdio.ac.cn/data/metadata-special-detail?id=1745022461423996930&otherId=1745022461554020354)
 795 detail?id=1745022461423996930&otherId=1745022461554020354 (last access:
 796 21 August 2026), 2024.
- 797 Duffy, G. and Chown, S.: Explicitly integrating a third dimension in marine species
 798 distribution modelling, *Marine Ecology Progress Series*, 564, 1–8,
 799 <https://doi.org/10.3354/meps12011>, 2017.
- 800 Duhamet, A., Albouy, C., Marques, V., Manel, S., and Mouillot, D.: The global depth
 801 range of marine fishes and their genetic coverage for environmental DNA
 802 metabarcoding, *Ecology and Evolution*, 13, <https://doi.org/10.1002/ece3.9672>,
 803 2023.
- 804 Durán, A. P., Green, J. M. H., West, C. D., Visconti, P., Burgess, N. D., Virah-Sawmy,
 805 M., and Balmford, A.: A practical approach to measuring the biodiversity impacts
 806 of land conversion, *Methods in Ecology and Evolution*, 11, 910–921,
 807 <https://doi.org/10.1111/2041-210x.13427>, 2020.
- 808 Egorova, Y., Reygondeau, G., Cheung, W. W. L., and Pakhomov, E. A.: Species
 809 Distribution Models for Mesopelagic Mesozooplankton Community, *Journal of*
 810 *Biogeography*, 52, 13–26, <https://doi.org/10.1111/jbi.15011>, 2025.
- 811 Elith, J., Kearney, M., and Phillips, S.: The art of modelling range-shifting species,
 812 *Methods in Ecology and Evolution*, 1, 330–342, [https://doi.org/10.1111/j.2041-](https://doi.org/10.1111/j.2041-210x.2010.00036.x)
 813 210x.2010.00036.x, 2010.
- 814 Ficetola, G. F., Rondinini, C., Bonardi, A., Baisero, D., and Padoa-Schioppa, E.:
 815 Habitat availability for amphibians and extinction threat: a global analysis,
 816 *Diversity and Distributions*, 21, 302–311, <https://doi.org/10.1111/ddi.12296>, 2015.
- 817 García-Jiménez, A., Talbi, M., Fitzgerald, L. M., Heim, A., Marcionetti, A., Schmid,
 818 S., Bertrand, J., Shaughnessy, A., Santiago, C., Rangseethampanya, P.,
 819 Ruttanachuchote, P., Aunkhongthong, W., Pengsakun, S., Sutthacheep, M.,
 820 Malinsky, M., Frédérich, B., Cortesi, F., Kochzius, M., Yeemin, T., Gaboriau, T.,
 821 and Salamin, N.: Habitat Specialisation Impacts Clownfish Demographic
 822 Resilience to Pleistocene Sea-Level Fluctuations, *Molecular Ecology*, 34, e70134,
 823 <https://doi.org/10.1111/mec.70134>, 2025.
- 824 Grenié, M., Violle, C., and Munoz, F.: Is prediction of species richness from stacked
 825 species distribution models biased by habitat saturation?, *Ecological Indicators*,
 826 111, 105970, <https://doi.org/10.1016/j.ecolind.2019.105970>, 2020.



- 827 IPBES: Global Assessment Report on Biodiversity and Ecosystem Services of the
 828 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem
 829 Services, IPBES, <https://doi.org/10.5281/zenodo.6417333>, 2019.
- 830 The IUCN Red List of Threatened Species: <https://www.iucnredlist.org>, last access: 21
 831 August 2026.
- 832 Kaschner, K., Watson, R., Trites, A., and Pauly, D.: Mapping world-wide distributions
 833 of marine mammal species using a relative environmental suitability (RES) model,
 834 Marine Ecology Progress Series, 316, 285–310,
 835 <https://doi.org/10.3354/meps316285>, 2006.
- 836 Klaassen, M., Marques, T. A., Alves, F., and Fernandez, M.: Trends in marine species
 837 distribution models: a review of methodological advances and future challenges,
 838 Ecography, <https://doi.org/10.1002/ecog.07702>, 2025.
- 839 La Manna, G., Ronchetti, F., Moro Merella, M., Vargiu, R., Perretti, F., and Ceccherelli,
 840 G.: Using spatial distribution modeling of commercial species to inform
 841 management of small-scale fisheries in a Mediterranean marine protected area,
 842 Ecology and Evolution, 14, <https://doi.org/10.1002/ece3.70169>, 2024.
- 843 Li, B., Cheng, C., Zhang, T., Mu, N., Li, Z., Yang, S., and Wu, X.: Global 1-km habitat
 844 distribution for endangered species and its spatial changes under future warming
 845 scenarios, Scientific Data, 12, <https://doi.org/10.1038/s41597-025-05898-6>, 2025.
- 846 Lumbierres, M., Dahal, P. R., Soria, C. D., Di Marco, M., Butchart, S. H. M., Donald,
 847 P. F., and Rondinini, C.: Area of habitat maps for the World’s terrestrial birds and
 848 mammals, Scientific Data, 9, <https://doi.org/10.1038/s41597-022-01838-w>, 2022.
- 849 Maureaud, A. A., Palacios-Abrantes, J., Kitchel, Z., Mannocci, L., Pinsky, M. L.,
 850 Fredston, A., Beukhof, E., Forrest, D. L., Frelat, R., Palomares, M. L. D., Pecuchet,
 851 L., Thorson, J. T., van Denderen, P. D., and Mériçot, B.: FISHGLOB_data: an
 852 integrated dataset of fish biodiversity sampled with scientific bottom-trawl
 853 surveys, Scientific Data, 11, 24, <https://doi.org/10.1038/s41597-023-02866-w>,
 854 2024.
- 855 Palacio, R. D., Negret, P. J., Velásquez-Tibatá, J., and Jacobson, A. P.: A data-driven
 856 geospatial workflow to map species distributions for conservation assessments,
 857 Diversity and Distributions, 27, 2559–2570, <https://doi.org/10.1111/ddi.13424>,
 858 2021.
- 859 Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I.-C.,
 860 Clark, T. D., Colwell, R. K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S.,
 861 Frusher, S., Garcia, R. A., Griffiths, R. B., Hobday, A. J., Janion-Scheepers, C.,
 862 Jarzyna, M. A., Jennings, S., Lenoir, J., Linnetved, H. I., Martin, V. Y.,
 863 McCormack, P. C., McDonald, J., Mitchell, N. J., Mustonen, T., Pandolfi, J. M.,
 864 Pettorelli, N., Popova, E., Robinson, S. A., Scheffers, B. R., Shaw, J. D., Sorte, C.
 865 J. B., Strugnell, J. M., Sunday, J. M., Tuanmu, M.-N., Vergés, A., Villanueva, C.,
 866 Wernberg, T., Wapstra, E., and Williams, S. E.: Biodiversity Redistribution under



- 867 Climate change: Impacts on Ecosystems and Human well-being, *Science*, 355,
 868 eaai9214, <https://doi.org/10.1126/science.aai9214>, 2017.
- 869 Phillips, S. J., Anderson, R. P., and Schapire, R. E.: Maximum entropy modeling of
 870 species geographic distributions, *Ecological Modelling*, 190, 231–259,
 871 <https://doi.org/10.1016/j.ecolmodel.2005.03.026>, 2006.
- 872 Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J., and
 873 Ferrier, S.: Sample selection bias and presence-only distribution models:
 874 implications for background and pseudo-absence data, *Ecological Applications*,
 875 19, 181–197, <https://doi.org/10.1890/07-2153.1>, 2009.
- 876 Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., and Blair, M. E.: Opening
 877 the black box: an open-source release of Maxent, *Ecography*, 40, 887–893,
 878 <https://doi.org/10.1111/ecog.03049>, 2017.
- 879 Phillips, S.: maxnet: Fitting ‘Maxent’ Species Distribution Models with ‘glmnet’, R
 880 package version 0.1.4, <https://CRAN.R-project.org/package=maxnet>, 2021.
- 881 Pickens, B. A., Carroll, R., Schirripa, M. J., Forrestal, F. C., Friedland, K. D., and Taylor,
 882 J. C.: A systematic review of spatial habitat associations and modeling of marine
 883 fish distribution: A guide to predictors, methods, and knowledge gaps, *PLOS ONE*,
 884 16, e0251818–e0251818, <https://doi.org/10.1371/journal.pone.0251818>, 2021.
- 885 Pörtner, H.-O., Scholes, R. J., Arneeth, A., Barnes, D. K. A., Burrows, M. T., Diamond,
 886 S. E., Duarte, C. M., Kiessling, W., Leadley, P., Managi, S., McElwee, P., Midgley,
 887 G., Ngo, H. T., Obura, D., Pascual, U., Sankaran, M., Shin, Y. J., and Val, A. L.:
 888 Overcoming the coupled climate and biodiversity crises and their societal impacts,
 889 *Science*, 380, 2023.
- 890 Ready, J., Kaschner, K., South, A. B., Eastwood, P. D., Rees, T., Rius, J., Agbayani,
 891 E., Kullander, S., and Froese, R.: Predicting the distributions of marine organisms
 892 at the global scale, *Ecological Modelling*, 221, 467–478,
 893 <https://doi.org/10.1016/j.ecolmodel.2009.10.025>, 2010.
- 894 Reygondeau, G., Egorova, Y., Boerder, K., Tittensor, D. P., Kaschner, K., Kesner-
 895 Reyes, K., Bailly, N., and Cheung, W. W. L.: AquaX: An enhanced and revised
 896 AquaMaps framework to model marine species distributions and biodiversity,
 897 *PLOS One*, 21, e0335823, <https://doi.org/10.1371/journal.pone.0335823>, 2026.
- 898 Robinson, N. M., Nelson, W. A., Costello, M. J., Sutherland, J. E., and Lundquist, C.
 899 J.: A Systematic Review of Marine-Based Species Distribution Models (SDMs)
 900 with Recommendations for Best Practice, *Frontiers in Marine Science*, 4,
 901 <https://doi.org/10.3389/fmars.2017.00421>, 2017.
- 902 Rondinini, C., Di Marco, M., Chiozza, F., Santulli, G., Baisero, D., Visconti, P.,
 903 Hoffmann, M., Schipper, J., Stuart, S. N., Tognelli, M. F., Amori, G., Falcucci, A.,
 904 Maiorano, L., and Boitani, L.: Global habitat suitability models of terrestrial
 905 mammals, *Philosophical Transactions of the Royal Society B: Biological Sciences*,
 906 366, 2633–2641, <https://doi.org/10.1098/rstb.2011.0113>, 2011.



- 907 Stohlgren, T. J., Jarnevich, C. S., Esaias, W. E., and Morisette, J. T.: Bounding species
 908 distribution models, *Current Zoology*, 57, 642–647,
 909 <https://doi.org/10.1093/czoolo/57.5.642>, 2011.
- 910 Tittensor, D. P., Mora, C., Jetz, W., Lotze, H. K., Ricard, D., Berghe, E. V., and Worm,
 911 B.: Global patterns and predictors of marine biodiversity across taxa, *Nature*, 466,
 912 1098–1101, <https://doi.org/10.1038/nature09329>, 2010.
- 913 Tyberghein, L., Verbruggen, H., Pauly, K., Troupin, C., Mineur, F., and De Clerck, O.:
 914 Bio-ORACLE: a global environmental dataset for marine species distribution
 915 modelling, *Global Ecology and Biogeography*, 21, 272–281,
 916 <https://doi.org/10.1111/j.1466-8238.2011.00656.x>, 2012.
- 917 UNEP-WCMC, WorldFish Centre, WRI, and TNC: Global distribution of coral reefs,
 918 Version 4.1, <https://doi.org/10.34892/t2wk-5t34>, 2021.
- 919 UNEP: Limiting Overshoot - Navigating exceedance of 1.5°C and pathways towards
 920 return, UNEP - UN Environment Programme, 2026.
- 921 Wang, L., Kerr, L. A., Record, N. R., Bridger, E., Tupper, B., Mills, K. E., Armstrong,
 922 E. M., and Pershing, A. J.: Modeling marine pelagic fish species spatiotemporal
 923 distributions utilizing a maximum entropy approach, *Fisheries Oceanography*, 27,
 924 571–586, <https://doi.org/10.1111/fog.12279>, 2018.
- 925 Woods, B. L., Van, A. P., Hindell, M. A., Raymond, B., Saunders, R. A., Walters, A.,
 926 and Rowan Trebilco: Species distribution models describe spatial variability in
 927 mesopelagic fish abundance in the Southern Ocean, *Frontiers in Marine Science*,
 928 9, <https://doi.org/10.3389/fmars.2022.981434>, 2023.
- 929 Woolley, S. N. C., Tittensor, D. P., Dunstan, P. K., Guillera-Arroita, G., Lahoz-Monfort,
 930 J. J., Wintle, B. A., Worm, B., and O’Hara, T. D.: Deep-sea diversity patterns are
 931 shaped by energy availability, *Nature*, 533, 393–396,
 932 <https://doi.org/10.1038/nature17937>, 2016.
- 933 Yang, Y., Yang, Y., Li, X., Yu, T., Pei, K., Zhong, Y., and Xie, Y.: Area of Habitat
 934 maps for national key protected wildlife in China, *Scientific Data*, 12,
 935 <https://doi.org/10.1038/s41597-025-05489-5>, 2025.
- 936 Yu, Le and Li, Y.: A dynamic, depth-resolved Area-of-Habitat dataset for global marine
 937 species, *Zenodo*, <https://doi.org/10.5281/zenodo.21933262>, 2026.
- 938 Zhuang, H., Zhao, L., Wang, Z., Zhang, Z., and Yang, L.: Identifying global marine
 939 climate refugia through a conservative approach to ocean biodiversity preservation,
 940 *Nature Communications*, 16, <https://doi.org/10.1038/s41467-025-65791-z>, 2025.