



SYSU TWSA v1.0: Global High-Resolution Terrestrial Water Storage Anomalies via Satellite Gravimetry

Yuhao Xiong^{1,2}, Wei Feng^{1,2*}, Jun Huang^{1,2}, Hongbing Bai^{1,2}, Guangyu Jian^{1,2}, Min Zhong^{1,2}

¹School of Geospatial Engineering and Science, Sun Yat-Sen University, Zhuhai 519082, China.

5 ²Key Laboratory of Comprehensive Observation of Polar Environment (Sun Yat-sen University), Ministry of Education, Zhuhai 519082, China.

Correspondence to: Wei Feng (fengwei@sysu.edu.cn)

Abstract. Publicly available global high-resolution terrestrial water storage anomaly (TWSA) datasets derived from satellite gravimetry remain scarce. Many existing global downscaling products rely heavily on hydrological models. Consequently, their performance can degrade in regions where key mass variations observed by the Gravity Recovery and Climate Experiment (GRACE) and its successor mission GRACE Follow-On (GFO) are poorly represented in the models, notably those associated with mountain glaciers and large lakes. Here we provide SYSU TWSA, a global monthly 0.5° TWSA dataset spanning April 2002 to December 2022, generated using a joint-inversion spatial downscaling framework that integrates large-scale constraints from GRACE/GFO, high-resolution spatial patterns from the WaterGAP Global Hydrological Model (WGHM), and additional mascon groups that explicitly represent mountain glaciers and selected large or rapidly changing lakes. The dataset helps alleviate the current shortage of global high-resolution products and explicitly strengthens the representation of glacier- and lake-related signals. We assess SYSU TWSA through four complementary evaluations: (1) basin-wise consistency with raw GRACE/GFO estimates, (2) a basin water-balance consistency check, (3) independent evaluation against *in situ* groundwater well observations, and (4) comparisons with representative downscaled products in both the spectral and spatial domains. SYSU TWSA shows strong agreement with GRACE/GFO at the basin scale, with coefficients of determination (R^2) exceeding 0.85 across basin-size classes. In small basins, consistency with terrestrial water fluxes derived from the basin water-balance equation improves substantially, with NSE increasing by 17.1% relative to raw GRACE/GFO across 1,200 basins. Agreement with groundwater wells also improves, with correlations increasing at 67.7% of 28,248 wells. Comparisons with representative assimilation-based and deep-learning downscaled products further indicate that SYSU TWSA achieves competitive overall accuracy while strengthening the representation of glacier- and lake-related signals.

1 Introduction

Water is a critical resource that sustains Earth's ecosystems and supports human societies, and its spatiotemporal distribution and availability are largely governed by the global water cycle. As an essential component of the climate system, the global water cycle redistributes water resources across space and time, thereby shaping regional water sustainability, the frequency



and intensity of hydrological extremes (e.g., floods and droughts), and influencing global sea level change (Bierkens, 2015; Rodell et al., 2018; Reager et al., 2016). In this context, terrestrial water storage integrates changes across multiple water compartments, including surface water, soil moisture, groundwater, snow, and ice, and serves as a key indicator of regional water balance as well as the hydrological response to climate change and human activities. Consequently, high-accuracy, continuous monitoring of water storage dynamics is essential for advancing our understanding of the water cycle, supporting water-security management, and addressing global change.

With the launch of the Gravity Recovery and Climate Experiment (GRACE) mission and its successor, GRACE Follow-On (GFO), satellite gravimetry has provided continuous observations of time-variable gravity fields, enabling the tracking of global surface mass redistribution (Wahr et al., 1998; Tapley et al., 2019; Feng et al., 2023; Rodell and Reager, 2023). Among these mass transport processes, terrestrial water storage anomaly (TWSA) constitutes one of the dominant contributors to time-variable gravity signals (Rodell and Reager, 2023). However, constraints imposed by satellite orbit design, payload instrument precision, and background model errors limit the effective spatial resolution of monthly GRACE/GFO solutions to ~330 km (Vishwakarma et al., 2018; Tapley et al., 2019), which substantially restricts their utility for small basin-scale hydrological research and applications (Pail et al., 2015; Chen et al., 2022; Wiese et al., 2022). Improving the effective spatial resolution of GRACE/GFO-derived TWSA is therefore critical for extending its hydrological applicability.

One of the main challenges in GRACE/GFO spatial downscaling is the lack of independent, high-precision measurements of TWSA, which makes it difficult to both develop downscaling frameworks and rigorously validate their accuracy. Existing studies generally follow two main pathways. The first is assimilation-based downscaling, which incorporates the large-scale observational constraints from GRACE/GFO into a high-resolution model to improve spatial resolution (Eicker et al., 2014; Li, B et al., 2019; Gerdener et al., 2023; Gou and Soja, 2024). However, such approaches typically rely heavily on hydrological models. These models still exhibit large uncertainties in time-varying signals from interannual to long-term scales (Scanlon et al., 2018) and often have deficiencies or missing modules for key processes such as groundwater dynamics and mountain glacier mass changes. Because GRACE/GFO observations contain signals associated with these processes, downscaling performance is limited in regions where the models do not simulate them reliably (Gou and Soja, 2024). The second pathway is statistical downscaling. It establishes statistical relationships between GRACE/GFO observations and finer-scale variables (e.g., precipitation, evapotranspiration, and temperature) to distribute large-scale signals onto finer spatial scales (Ning et al., 2014; Yin et al., 2018; Vishwakarma et al., 2021). Recent advances in machine learning have further improved the ability to capture nonlinear relationships (Seyoum and Milewski, 2017; Zuo et al., 2021; Zhang et al., 2021; Yin et al., 2022). Nevertheless, global statistical downscaling often faces challenges related to the generalizability of empirical statistical relationships under spatial heterogeneity, overfitting in machine-learning training, and the reliability of transferring relationships from GRACE/GFO-effective scales to finer local scales. When the training samples are drawn directly from gridded GRACE/GFO TWSA products, such as official mascon products provided at 0.5° or 0.25° (Loomis et al., 2019; Save et al., 2016; Wiese et al., 2016), although distributed on fine grids, their information



65 content is substantially coarser than the nominal resolution (Watkins et al., 2015; Vishwakarma et al., 2021). As a result, relationships learned from such targets may be valid only at large scales, that is, within the effective GRACE/GFO resolution, while their applicability at finer local resolutions, such as those of the high-resolution predictor variables, remains insufficiently demonstrated.

Against this backdrop, publicly available global high-resolution TWSA products remain scarce. Existing assimilation-based products, despite their strengths, often inherit model structural limitations that can leave glacier mass changes poorly constrained in glacierized regions. Meanwhile, large lakes that are effectively detectable by GRACE/GFO, such as the North American Great Lakes and Lake Victoria in Africa, are overlooked in some downscaled products, thereby weakening their ability to monitor lake storage variations. To address these issues, we recently developed a joint-inversion spatial downscaling framework and validated it across China (Xiong et al., 2025a). The framework does not require training a predictive model or calibrating hydrological model parameters, making it more flexible and computationally efficient in practice. It further improves performance in regions where hydrological models are less reliable by allowing mascon groups to be defined and incorporated flexibly, thereby supplementing glacier and large-lake signals that are not well represented in hydrological models. Building on this framework, we extend the approach to the global domain and develop a high-resolution SYSU TWSA product to alleviate the current scarcity of global high-resolution datasets and to better represent glacierized regions and large lakes. We evaluate its performance at the basin scale, that is, in basins comparable to the effective GRACE/GFO resolution, and conduct independent checks using a water-balance constraint and *in situ* groundwater well observations to assess its potential advantages. Finally, we compare the SYSU product with existing representative global downscaled products.

2 Data

85 2.1 GRACE/GFO Data and Processing

In this study, we used the monthly GRACE/GFO Level 2 gravity field products from April 2002 to December 2022, provided in the form of spherical harmonic coefficients (up to degree and order 90). The post-processing of these coefficients included the following steps: (1) replacing the degree-1 term with monthly geocenter estimates computed by Landerer (2019), and substituting the C_{20} and C_{30} coefficients with satellite laser ranging (SLR) estimates (Sun et al., 2016; Loomis et al., 2020); (2) removing the mean gravity field over January 2004 to December 2009; (3) applying a GIA correction using the ICE-6G model (Peltier et al., 2018); (4) filtering stripe noise using the DDK3 filter (Kusche et al., 2009); (5) reducing the topography effects (Yang et al., 2022); and (6) synthesizing the spherical harmonics to derive surface mass changes on a 0.5° grid (Wahr et al., 1998). Notably, although the surface mass changes were represented on a 0.5° grid after synthesis, their effective spatial resolution remained approximately 330 km, following $20,000/l$, where l denotes the maximum spherical harmonic degree.



2.2 WaterGAP Global Hydrological Model Outputs

The WaterGAP Global Hydrology Model (WGHM; Müller Schmied et al., 2021) outputs used in this study covered 2002–2022 and were forced by the GSWP3-W5E5 climate dataset. WGHM provides gridded monthly water storage components at 0.5° resolution, simulated within a vertical and lateral water-balance framework, and represents many storage compartments, including lakes, wetlands, rivers, reservoirs, groundwater, soil moisture, and snow water equivalent. Notably, WGHM does not simulate mountain glacier mass changes and does not explicitly represent several large lakes, such as the North American Great Lakes, Lake Victoria in Africa, and rapidly expanding lakes in the Tibetan Plateau.

2.3 Mountain Glacier Catalogue

The Randolph Glacier Inventory (RGI) version 6.0 is a global glacier outline dataset designed to provide a static snapshot of glacier extents worldwide (Pfeffer et al., 2014). It provides polygonal boundaries for individual glaciers, compiled through the Global Land Ice Measurements from Space (GLIMS) initiative. In this study, we used RGI v6.0 to delineate global mountain glacier mascon groups.

2.4 Hydrometeorological Data

In this study, monthly precipitation (P), evapotranspiration (E), and runoff (R) data were adopted from the ERA5-Land dataset. These variables were used in the water-balance equation to evaluate the potential advantages of the SYSU product relative to the raw GRACE/GFO observations at the global level-4 basins (Lehner and Grill, 2013). ERA5-Land is a high-resolution land reanalysis produced by the ECMWF ERA5 system, and the data are available from the Copernicus Climate Data Store (Service, 2019).


2.5 *In situ* ground wells Data

The *in situ* groundwater well observations used in this study were obtained from the Global Groundwater Monitoring Network (GGMN), coordinated by the International Groundwater Resources Assessment Centre (IGRAC), and were used as independent data to evaluate the performance of the SYSU TWSA product. It should be noted that GGMN does not provide a comprehensive global coverage of groundwater observations; its spatial distribution and temporal continuity depend on the degree to which national and sub-national agencies openly share their data, resulting in substantial heterogeneity across regions. To ensure comparability, we pre-processed the groundwater level time series by aggregating the original measurements to monthly means and retained only wells with at least 60 months (5 years) of valid records during April 2002 to December 2022. This screening yielded 28,248 wells for the validation analysis.



3 Joint Inversion Downscaling Framework

125 The key strategy of the joint inversion downscaling was to integrate the large-scale information from GRACE/GFO with high-resolution spatial information from the model and self-defined mascon groups (Xiong et al., 2025a). In this framework, TWSA data were regarded as a linear combination of spatial basis functions with temporal evolution coefficients. The high-resolution spatial basis functions were comprised of (i) spatial patterns extracted from WGHM-simulated TWSA compartments (Sect. 3.1) and (ii) supplemental basis functions constructed for signals not explicitly simulated in WGHM by defining global mountain glaciers and selected lakes mascon groups (Sect. 3.2-3.3). For the temporal evolution, we first
130 ensured consistency with the effective GRACE/GFO resolution by truncating all basis functions to the same maximum spherical harmonic degree and order as the GRACE/GFO fields and applying the DDK3 filter. This scale matching is essential for the subsequent least-squares fit, because the temporal coefficients can be robustly estimated only when the observations and basis functions share a consistent effective spatial resolution. We then fitted GRACE/GFO observations using these filtered spatial basis functions via a least-squares adjustment to estimate the temporal coefficients associated with
135 each basis function. These adjusted coefficients replaced the potentially unreliable temporal evolution simulated by the model, and provided temporal information for the self-defined mascon groups. Finally, the downscaled TWSA fields were reconstructed by combining the adjusted temporal evolution with the unfiltered (high-resolution) spatial basis functions. The overall workflow was summarized in Fig. 1.



140 **Figure 1: Flowchart of the joint inversion downscaling method.** SWES denotes snow water equivalent storage, SMS is soil moisture storage, GWS represents groundwater storage, and RESS refers to reservoir storage.

3.1 Extracting WGHM-simulated spatial patterns

WGHM provides gridded monthly mass change for multiple vertically integrated TWSA compartments, including snow water equivalent, reservoir storage, soil moisture, groundwater, wets, and rivers. We applied independent component analysis (ICA) to these compartments to extract spatial patterns, which were then used as spatial basis functions in the inversion. During the ICA decomposition, we retained the spatial patterns associated with the dominant trend, seasonal, and interannual variation. These patterns typically correspond to larger singular values. Retaining the dominant patterns acts as the effect of regularization, helping mitigate solution instabilities in the global inversion that can arise from an over-parameterized model.

150 3.2 Defining spatial basis functions for glacier mascon groups

The glacier mascon groups are user-defined regions of the Earth’s surface designed to encompass concentrated glacierized areas exceeding 100 km² (Ciraci et al., 2020). Each mascon group is represented by multiple small blocks, whose centroids



are defined on the 0.5° grid. These mascon groups do not necessarily coincide with the exact glacier outlines, but are designed to capture glacier mass changes detectable by GRACE/GFO (Jacob et al., 2012).

155 Building on the global glacier mascon groups of Jacob et al. (2012) and Ciraci et al. (2020), we refined the mascon groups using the glacier information provided by RGI v6.0. Global mountain glaciers were divided into 14 major regions: (1) Alaska; (2) Western Canada and the United States; (3) Canadian Archipelago; (4) Iceland; (5) Svalbard; (6) Scandinavia; (7) Russian Arctic; (8) North Asia; (9) Central Europe; (10) Caucasus and Middle East; (11) High Mountain Asia; (12) Low Latitudes; (13) Southern Andes; and (14) New Zealand. Each major region was further subdivided into several subregions,
160 shown in different colors (Fig. A1). Within each mascon unit, the glacier-covered areas were coded as 1 and the non-glacier areas as 0. We then incorporated these self-defined glacier mascon groups as spatial basis functions for global mountain glaciers, thereby supplementing the missing glacier component in WGHM within the joint-inversion framework.

3.3 Building spatial basis functions for lake mascon groups

We defined three lake mascon groups representing (1) the Great Lakes (North American), (2) Victoria Lake (Africa), and (3)
165 50 representative lakes (Tibetan Plateau), as shown in Fig. A2. The lake dataset in the Tibetan Plateau was taken from Li, X. et al. (2019). Because these lakes are relatively small, we applied ICA to the gridded lake storage variations and retained the leading spatial patterns that capture the long-term trend and the seasonal cycle. These patterns were then used as spatial basis functions for the lake group in the Tibetan Plateau. In contrast, the Great Lakes and Victoria Lake were defined as individual mascon units, assigned a value of 1 inside the lake and 0 outside. We then incorporated these lake mascon groups as spatial
170 basis functions to represent lake storage variations observed by GRACE/GFO yet not explicitly simulated in WGHM.

4 Validation

In this section, we evaluate the performance of the SYSU downscaled product from four perspectives: (1) comparison with the raw GRACE/GFO estimates at the basin scale; (2) assessment of its potential advantages using a water-balance equation; (3) independent evaluation against *in situ* groundwater well observations; and (4) comparison with other downscaled
175 products in both the spectral and spatial domains.


4.1 Compared with GRACE/GFO at the basin scale

Given the lack of independent high-precision TWSA measurements at the global scale, a direct validation of downscaled products at the grid scale (0.5°) is challenging. The only globally available TWSA observations are provided by GRACE/GFO. However, GRACE/GFO does not truly resolve variability at the 0.5° grid scale and therefore cannot serve as
180 a reliable reference at this scale. By contrast, GRACE/GFO observations are generally reliable at the basin scale when basin sizes are comparable to the effective GRACE/GFO resolution (Boergens et al., 2022), which enables an indirect evaluation. It should be noted, however, that a direct basin-scale comparison between a downscaled product and GRACE/GFO is still



185 inappropriate. GRACE/GFO measurements can suffer from signal leakage in some basins (Landerer and Swenson, 2012). Although a range of signal restoration approaches has been proposed (Vishwakarma et al., 2017; Xiong et al., 2025b; Long et al., 2015), their accuracy is difficult to assess across individual basins in the absence of ground truth. Under these circumstances, it is more appropriate to forward-process the downscaled product by expanding it into spherical harmonics to the same maximum degree and order as GRACE/GFO and applying the same smoothing filter before evaluation. This ensures that the two datasets have comparable spectral content, that is, an equivalent effective resolution.

190 In this study, we selected 89 basins of different sizes from the GRDC dataset (GRDC, 2020). Based on basin area, we grouped them into large basins (40 basins; $> 500,000 \text{ km}^2$), medium basins (36 basins; $100,000\text{-}500,000 \text{ km}^2$), and small basins (13 basins; $40,000\text{-}100,000 \text{ km}^2$), as shown in Fig. 2. In addition, global hydroclimatic conditions were stratified into arid (A: $0.03 < \text{AI} < 0.2$), semi-arid (SA: $0.2 < \text{AI} < 0.5$), sub-humid (SH: $0.5 < \text{AI} < 0.65$), and humid (H: $\text{AI} > 0.65$) classes based on the aridity index (Trabucco and Zomer, 2019).



1 Glomaa	2 Weser	3 Gota	4 Po	5 Rukwa	6 Kizilirmak	7 Doce	8 Comoe
9 San Joaquin	10 Cuyuni	11 Daugava Dvina	12 Panuco	13 Rhone	14 Loire	15 Santiago	16 Elbe
17 Melrhir	18 Ural	19 Ogooue	20 YongDing He	21 Olenek	22 Liao He	23 Har US Nuur	24 Fraser
25 Yana	26 Thelon	27 Thorgay	28 Krishna	29 Magdalena	30 Uruguay	31 Salween	32 Neva
33 Khatanga	34 Churchill	35 Severnaya Dvina	36 Godavari	37 Pechora	38 Parnaiba	39 Indigirka	40 Hamun
41 Colorado	42 Zhu	43 Limpopo	44 Volta	45 Lake Balkhash	46 Irrawaddy	47 Rio De La Plata	48 Don
49 Senegal	50 Dnieper	51 Brahmaputra	52 Sao Francisco	53 Kolyma	54 Colorado	55 Columbia	56 Bravo
57 Okavango	58 Tocantins	59 Mekong	60 Danube	61 Jubba	62 Yukon	63 Indus	64 Shatt Al Arab
65 Orinoco	66 Yellow	67 Orange	68 Ganges	69 Saint Lawrence	70 Murray	71 Nelson	72 Lake Eyre
73 Aral Sea	74 Zambezi	75 Volga	76 Tarim He	77 Yangtze	78 Mackenzie	79 Niger	80 Amur
81 Lena	82 Chad	83 Yenisey	84 Parana	85 Ob	86 Mississippi	87 Nile	88 Congo
89 Amazon							

195 **Figure 2: Global distribution of 89 river basins ranked by ascending area. Humidity is characterized by the Aridity Index (AI). Basin boundaries are based on GRDC (2020) datasets (available at: <https://mrb.grdc.bafg.de/>).**

We expanded the SYSU product into spherical harmonics up to the same maximum degree and order as GRACE/GFO and applied the same spatial smoothing filter, yielding a filtered SYSU product with an effective spatial resolution consistent with GRACE/GFO. To assess performance across basin scales, we averaged (latitude-weighted) both GRACE/GFO and the filtered SYSU TWSA for large, medium, and small basins to derive basin-wise time series. We then concatenated the time series within each basin-size class into column vectors and generated scatter-density plots with GRACE/GFO on the X-axis and the filtered SYSU product on the Y-axis, from which the coefficient of determination (R^2) was calculated. As shown in Fig. 3, after matching the effective spatial resolution, the filtered SYSU product agrees closely with GRACE/GFO in basin-



mean TWSA, with R^2 exceeding 0.85 for all basin-size classes. Specifically, R^2 reached 0.95 for large basins, 0.87 for
205 medium basins, and 0.86 for small basins. These results indicate that the SYSU product is highly reliable and support the
effectiveness of the joint inversion downscaling framework in preserving the basin-scale accuracy of GRACE/GFO signals.
Notably, performance is relatively lower for small basins. This is expected because GRACE/GFO observations are generally
more reliable for large basins (Vishwakarma et al., 2018; Boergens et al., 2022), whereas small basins may perform
relatively poorly due to their inherently low signal strength, which is more significantly affected by the measurement
210 accuracy of GRACE/GFO (~2 cm) and leakage errors.




Figure 3: Scatter density plots of the basin-wise average TWSA in different scale basins estimated from the filtered SYSU products (Y-axis) versus GRACE/GFO measurements (X-axis).

As shown in Fig. 4, we compared the TWSA time series for different scale basins. For large basins, both SYSU and filtered
215 SYSU products agree well with GRACE/GFO observations. The median root-mean-square error (RMSE) between the
filtered SYSU product and GRACE/GFO is 1.27 cm, which falls within the GRACE/GFO measurement uncertainty
(Vishwakarma et al., 2018; Tapley et al., 2019). The corresponding median Nash-Sutcliffe efficiency (NSE) is
approximately 0.95, and the median correlation coefficient (CC) is 0.98. For medium basins, performance decreases slightly
relative to large basins, but the majority of basins still show strong agreement in their time series, with an RMSE of 2.36 cm,
220 an NSE of 0.83, and a CC of 0.95. For small basins, the consistency decreases further, with an RMSE of 3.75 cm, an NSE of
0.81, and a CC of 0.95.

Although GRACE/GFO observations are generally affected by signal attenuation and leakage, these effects are relatively
small for large basins (Xiong et al., 2025b). Spherical harmonic truncation and spatial smoothing can cause signal spreading
beyond basin boundaries, but most of the signal remains concentrated within the target region. As the joint-inversion
225 downscaling framework preserves the large-scale accuracy of GRACE/GFO measurements, all three products show a high
level of consistency in large basins. However, as the basin area decreases, the effects of attenuation and leakage typically
become more pronounced due to the limited effective spatial resolution of GRACE/GFO. This tends to dampen basin-wise
TWSA amplitudes and underestimate long-term trends (Scanlon et al., 2012; Long et al., 2015; Vishwakarma et al., 2016).
Improving spatial resolution can alleviate these issues. Consequently, in basins where attenuation and leakage are severe, the




230 basin-average TWSA derived from the SYSU product may exhibit larger amplitudes and trends than GRACE/GFO, as
observed for the Cuyuni River basin (ID: 10), the Daugava Dvina basin (ID: 11), the Gota basin (ID: 3), and the Glomma
basin (ID: 1). Consistently, when SYSU is degraded to the same effective resolution as GRACE/GFO (i.e., the filtered
SYSU product), the differences in basin-average TWSA are substantially reduced. In addition, for basins in arid regions,
such as the Hamun basin (ID: 40) and Lake Balkhash (ID: 45), the filtered SYSU product still shows discrepancies relative
235 to GRACE/GFO. This may reflect the weak hydrological signals in these arid environments, for which both GRACE/GFO
and WGHM have limited sensitivity to detect or simulate the variations accurately (Gou and Soja, 2024; Xiong et al., 2025a).



Figure 4: Comparison of time series in different basin scales: (a) large basins, (b) medium basins, and (c) small basins.



To verify the performance of the SYSU product in preserving long-term trends and the amplitudes of the annual and semi-annual signals, we compared the consistency of signal estimates among the three TWSA products in large, medium, and small basins, i.e., the SYSU, the filtered SYSU, and GRACE/GFO, as shown in Fig. 5. For large basins, the long-term trend and the annual and semi-annual amplitudes estimated from SYSU closely match those from GRACE/GFO, with R^2 of 0.90, 0.99, and 0.96, respectively, and most points clustering near the red dashed line. Notably, the filtered SYSU product does not show a clear improvement over the SYSU in its agreement with GRACE/GFO, with R^2 of 0.94, 0.99, and 0.92, respectively. This indicates that GRACE/GFO signals are reliable for large-basin scales, where leakage effects are relatively small, and that our downscaling framework effectively inherits their accuracy. For medium basins, the agreement between SYSU and GRACE/GFO decreases, with R^2 of 0.69, 0.94, and 0.83 for the long-term trend, the annual and semi-annual amplitudes, respectively. The filtered SYSU product also shows no substantial improvement, with R^2 of 0.72, 0.97, and 0.80. In contrast, for small basins, SYSU shows relatively poor agreement with GRACE/GFO observations, whereas the consistency between the filtered SYSU product and GRACE/GFO increases markedly, with R^2 of 0.70, 0.90, and 0.95. This contrast indirectly suggests that GRACE/GFO performs poorly at small basins, where severe leakage errors can prevent the native observations from capturing the underlying signals. By increasing the spatial resolution, the SYSU product can substantially alleviate this problem. Therefore, the apparent mismatch between SYSU and GRACE/GFO, together with the closer agreement between filtered SYSU and GRACE/GFO, mainly reflects differences in effective spatial resolution. Figures 5b and 5c compare the basin-wise secular trends and annual amplitudes derived from the SYSU product and GRACE/GFO. Overall, the two datasets agree closely, with RMSEs of 0.24 cm yr^{-1} for the trend and 1.93 cm for the annual amplitude. The largest trend difference is found in the Daugava Dvina basin (ID: 11), reaching -0.67 cm yr^{-1} , whereas the largest discrepancy in annual amplitude occurs in the Cuyuni basin (ID: 10), at 15.23 cm . These differences are consistent with leakage effects (Fig. 4).



260

Figure 5: Comparison of TWSA signals across different basin scales. (a) Scatter plots comparing long-term trends, annual amplitudes, and semi-annual amplitudes estimated by the SYSU and filtered SYSU products (Y-axis) against GRACE/GFO observations (X-axis). (b) Spatial distribution map illustrating basin-scale trend of the SYSU product and GRACE/FO observations. (c) Spatial distribution map showing basin-wise annual amplitude of the SYSU product and GRACE/FO observations.

265


4.2 Evaluation Based on Water Balance Equation in Sub-Basins

To evaluate the potential advantages of the downscaled product, we used a water balance equation ($TWF = P - E - R$) to assess whether the SYSU product improves performance relative to the raw GRACE/GFO observations. We selected 1,200 level-4 basins globally (Lehner and Grill, 2013) as the evaluation domain and used NSE as the performance metric. Considering data gaps in the GRACE/GFO record and the degraded data quality during the single-accelerometer period, we restricted the analysis to January 2003-December 2010. For the SYSU and GRACE/GFO, terrestrial water flux (TWF) was

270



estimated from monthly TWSA using a centered difference approach ($TWF = TWSA(t + 1) - TWSA(t - 1) / 2\Delta t$, Long et al., 2014). For ERA5-Land, TWF was computed from P, E, and R, and an adjacent-month smoothing scheme was applied to reduce potential high-frequency noise (Landerer et al., 2010).



275

Figure 6: Agreement assessment of TWF estimates in 1200 level-4 basins. Comparison between SYSU product, GRACE/GFO observations, and water balance-derived TWF quantified via NSE. (a) Basin-wise NSE of SYSU product. (b) Basin-wise NSE of GRACE/GFO observations. (c) NSE differences between SYSU and GRACE/GFO across watersheds sorted by ascending drainage area; positive values indicate superior performance of the SYSU product. The red dashed line marks the GRACE/GFO effective resolution limit of 63,000 km² (Vishwakarma et al., 2018).

280

Figure 6 summarizes the agreement between TWF estimates derived from the SYSU product and GRACE/GFO, and those computed from an independent water balance equation. The SYSU product shows a strong ability to close the basin-wise water balance across the 1,200 basins, with 71.7% of basins yielding positive NSE values. By comparison, 59.2% of basins show positive NSE when TWF is derived from the raw GRACE/GFO. As shown in Fig. 6a-b, most negative NSE values occur in arid regions where hydrological signals are weak, such as the Sahara in Africa and the Arabian Peninsula in the Middle East. For basins larger than the GRACE/GFO effective-resolution limit (63,000 km²), SYSU provides little improvement over GRACE/GFO, and the two products yield nearly identical NSE values. In contrast, for basins smaller than this limit, the NSE increases by 17.1%, indicating a clear advantage of SYSU at finer spatial scales. The most evident improvements relative to GRACE/GFO are found in small basins in regions such as the western European coasts and the

285



290 eastern Chinese coasts (Fig. 6a-b). These results suggest that, provided P, E, and R estimates are sufficiently accurate, improving the spatial resolution of TWSA products can yield TWF estimates that are closer to those derived by the basin water balance equation. It is worth noting that this comparison should be interpreted as a consistency check rather than a strict independent validation, because uncertainties in P, E, and R and unaccounted human interventions (e.g., inter-basin transfers) can affect basin water-balance closure.

295 **4.3 Comparison with *In situ* Wells data**

To evaluate the extent to which the SYSU TWSA product improves spatial detail relative to the coarse resolution GRACE/GFO observations, we used 28,248 *in situ* groundwater wells worldwide as an independent reference. Because well records measure groundwater level changes whereas satellite gravimetry provides water storage changes, a strict magnitude comparison would require converting water levels to storage using parameters such as specific yield. Such parameters are difficult to obtain reliably at the global scale. We therefore used the correlation coefficient as a consistency metric to quantify whether SYSU better matches well observations than the raw GRACE/GFO product. Specifically, for each well location, we interpolated the SYSU and GRACE/GFO TWSA fields to the site to obtain corresponding time series, and then computed their correlations with the observed groundwater level time series. By comparing correlations before and after downscaling, we directly tested whether the added spatial detail yields a more accurate and physically consistent representation of local hydrological variations. As shown in Fig. 7, downscaling leads to a clear improvement: compared with raw GRACE/GFO, correlations between SYSU TWSA and groundwater level time series increase at 67.7% of wells. Regional statistics indicate particularly pronounced improvements in the United States, India, and Australia, where correlations increase at 77.4%, 76.2%, and 77.7% of wells, respectively, whereas the improvement is less evident in Europe. Overall, the results indicate that the downscaled SYSU TWSA is more consistent with observed groundwater level variations than the raw GRACE/GFO observations, highlighting the value of downscaling.




Figure 7: Difference in correlation coefficients between the SYSU product and *in situ* groundwater well observations versus raw GRACE/GFO and well observations, calculated across 28,248 groundwater wells. Positive values indicate stronger agreement between the SYSU product and well data.

315 4.4 Comparison with other products

We further compared the SYSU product with GRACE/GFO spherical harmonic and mascon solutions, as well as with WGHM and several representative downscaled products, to evaluate the performance of each dataset. The downscaled products considered here were derived from WGHM through data assimilation. One uses a conventional Ensemble Kalman Filter (EnKF) approach (GLWS 02, hereafter GLWS), and the other is based on a self-supervised deep learning approach
320 (Gou and Soja, 2024). The evaluation was conducted from two perspectives, in the spectral and spatial domains.



4.4.1 Spectral domain

In this section, we compare each downscaled product with GRACE/GFO and WGHM in the spherical harmonic domain. To ensure a consistent evaluation, we converted land TWSA from all downscaled products, the WGHM outputs, and GRACE/GFO (including both spherical harmonic and mascon solutions) into spherical harmonic coefficients. Notably, the expansion was performed over the global land domain and excluded the Greenland and Antarctic ice sheets to avoid contamination of the assessment by polar mass variations.




Figure 8: RMS of spherical harmonic coefficients derived by different TWSA products (2003-2019). RMS is expressed in terms of geoid change here, in mm-units.

Figure 8 shows the RMS of each spherical harmonic coefficient's contribution to geoid variability (up to degree and order 90) for the different products. The results indicate that all downscaled products effectively remove much of the high-frequency content inherited from WGHM, yet their RMS values remain slightly higher than those of GRACE/GFO. In addition, compared with the EnKF-based GLWS product, both the deep learning downscaling product of Gou and Soja (2024) and the joint-inversion SYSU product better preserve the low-degree coefficients observed by GRACE/GFO. However, beyond roughly degree 30, SYSU retains substantially more variability than Gou and Soja's product.



Figure 9 shows the degree variance curves for October 2005, and the overall behavior is consistent with the patterns in Fig. 8. A clear offset is evident between WGHM and GRACE/GFO. This may reflect stronger signals in WGHM over Siberia and North America (Gerdener et al., 2023), or may arise because the WGHM fields were not demeaned. Over degrees 0-25, all downscaled products remain in close agreement with GRACE/GFO. Beyond about degree 30, however, the curves begin to diverge. Because GRACE/GFO solutions are filtered and regularized, their high-degree power is suppressed. In contrast, the SYSU and GLWS products retain more high-frequency variance than the product of Gou and Soja (2024), indicating that they preserve more small-scale signals but also carry a higher noise level.




Figure 9: Degree variance for different products in October 2010 expressed by geoid heights (mm).

Given that individual spherical harmonic coefficients can be used to assess how a downscaling method corrects the hydrological signal from WGHM (Gerdener et al., 2023), we selected C_{20} as a representative example (Fig. 10). The GLWS product is closer to GRACE/GFO than WGHM, but it still tends to overestimate GRACE/GFO and exhibits larger temporal variability. This suggests that the EnKF-based assimilation inherits characteristics from both GRACE/GFO and WGHM and improves the hydrological model to some extent, yet remains constrained by its underlying error-model assumptions (Gerdener et al., 2023). By contrast, the downscaled product of Gou and Soja (2024) and the SYSU product agree closely with GRACE/GFO, with C_{20} values lying between the GRACE/GFO spherical harmonic and mascon solutions. These results indicate that deep learning-based and joint-inversion downscaling can offer advantages over conventional EnKF assimilation.




Figure 10: Time series of C_{20} spherical harmonic coefficient derived by different products.

355 4.4.2 Spatial domain

In this section, we focus on spatial comparisons. Figure 11a shows the long-term trends estimated from each product from January 2003 to December 2019. Consistent with previous studies (Döll et al., 2014; Scanlon et al., 2018), the WGHM-simulated TWSA trends are substantially smaller in magnitude than those derived from GRACE/GFO. In addition, except for the polar regions, WGHM does not effectively capture the mountain glacier melt signals evident in GRACE/GFO observations. Despite these discrepancies, the two datasets show good agreement in some regions, particularly in areas with intensive groundwater depletion (e.g., the Northwest India Plain, the North China Plain, and California's Central Valley) and in regions characterized by pronounced wetting-drying variability (e.g., the Amazon and La Plata basins and the African Great Lakes region). Notably, relative to GRACE/GFO, WGHM exhibits richer spatial detail in the trend patterns.

All three downscaled products largely retain the spatial resolution of the WGHM. Their long-term trends exhibit much finer spatial detail than GRACE/GFO, revealing more localized positive and negative trend patterns. For example, the river-channel structure within the Amazon Basin is clearly resolved. However, in some regions, such as the northern Great Plains of the United States, GRACE/GFO observations indicate a recovery in groundwater storage (Rateb et al., 2020; Scanlon et al., 2023), whereas the downscaled product of Gou and Soja (2024) does not correct the WGHM simulation and still suggests continued groundwater depletion. By contrast, the SYSU and GLWS products successfully adjust the negative WGHM trend and align with the GRACE/GFO observations, underscoring the critical contribution of GRACE/GFO constraints within both the joint-inversion framework and the EnKF approach. More importantly, relative to the other two downscaled products, SYSU incorporates mascon groups for mountain glaciers and selected lakes during the inversion, enabling it to capture glacier-melt signals and storage increases in large lakes. Compared with GRACE/GFO, SYSU provides substantially enhanced spatial resolution, yielding a more localized depiction of the spatial details of mountain glaciers and large lakes.




Figure 11: (a) Long-term trends in TWSA estimated from different products for the period January 2003 to December 2019. (b) Same as (a), but for annual amplitudes. (c) Same as (a), but for phases, indicating the month of peak amplitude (e.g., 1 represents January).



380 Figures 11b and 11c compare the annual amplitude and its phase (defined as the month of maximum amplitude) among the
different products. Because seasonal water storage variation in most basins is dominated by the annual signals, the products
show broadly consistent annual amplitudes. Moreover, the downscaled products not only achieve a spatial resolution
comparable to that of WGHM, but also produce amplitudes more consistent with GRACE/GFO than WGHM. This indicates
that these downscaling approaches benefit from incorporating GRACE/GFO constraints to improve hydrological-model
385 representations of seasonal variation. For the phase (Fig. 11c), WGHM exhibits a 1-2 month offset relative to GRACE/GFO
in multiple regions. GLWS reduces this discrepancy to some extent but still deviates from GRACE/GFO, whereas the SYSU
and Gou and Soja (2024) products show phases that are generally closer to GRACE/GFO. Locally, however, phase advances
or delays are still apparent, which may reflect genuine regional hydrological differences. A notable discrepancy appears in
390 northern Africa, where the phase from SYSU and from Gou and Soja (2024) differs substantially. SYSU is close to the
GRACE/GFO spherical harmonic solution, whereas Gou and Soja (2024) is close to the mascon solution. This discrepancy
likely reflects the extensive desert cover in northern Africa, where hydrological signals are weak and the uncertainty of
GRACE/GFO observations is relatively large (Boergens et al., 2022). Moreover, given the limited effective spatial resolution
of GRACE/GFO, it remains difficult to determine which downscaled product provides a more accurate phase estimate in this
region.

395 The qualitative comparisons of long-term trends, annual amplitudes, and phases suggest that EnKF, deep learning, and our
joint-inversion downscaling approaches all have considerable potential to improve WGHM-simulated TWSA estimates.
Given that GRACE/GFO is highly accurate at its effective spatial resolution (Vishwakarma et al., 2018; Boergens et al.,
2022), we quantify the improvement of each of the three downscaled products over WGHM in estimating TWSA.
Specifically, we assess which product matches GRACE/GFO most closely when evaluated at the same effective resolution.

400 To this end, we expanded the WGHM outputs and the three downscaled products into spherical harmonics to the same
maximum degree and order as GRACE/GFO (degree/order 90) and applied the same DDK3 filtering to match the effective
spatial resolution of GRACE/GFO. We then derived basin-average time series for 288 global level-3 basins (Lehner and
Grill, 2013) over January 2003 to December 2019 and evaluated their agreement with GRACE/GFO using RMSE, NSE, CC,
and R^2 (Table 1).

405 **Table 1. Evaluation of different products in 288 basins. Metrics include the median values of RMSE (in cm), NSE, CC, and R^2 . For each row, the best-performing value is shown in bold, and the second-best is underlined.**

Product type	RMSE	NSE	CC	R^2
SYSU	<u>2.24</u>	0.71	0.90	0.92
Gou and Soja (2024)	1.87	<u>0.68</u>	<u>0.88</u>	<u>0.83</u>
GLDAS CLSM DA v2.2	2.61	0.43	0.75	0.66
GLWS	3.40	0.14	0.68	0.45
WGHM	3.27	0.14	0.65	0.26



In terms of RMSE, all three downscaled products outperform WGHM (Table 1 and Fig. 12). Specifically, the median RMSE between WGHM and GRACE/GFO is 3.27 cm (mean: 4.41 cm). The GLWS product performs similarly to WGHM overall, with a median RMSE of 3.40 cm (mean: 4.04 cm). By contrast, the SYSU product and the [Gou and Soja \(2024\)](#) product achieve low RMSEs and both outperform GLWS. SYSU yields a median RMSE of 2.24 cm (mean: 2.44 cm), whereas [Gou and Soja \(2024\)](#) yields a median RMSE of 1.87 cm (mean: 2.37 cm). These values support the overall quality of the latter two downscaled products, given that GRACE/GFO uncertainties are on the order of 2-3 cm ([Wahr et al., 2006](#); [Tapley et al., 2019](#)). Notably, in glacierized regions, SYSU performs markedly better than the [Gou and Soja \(2024\)](#) product (e.g., along the Gulf of Alaska coast). This advantage likely reflects that SYSU not only leverages information from WGHM but also explicitly incorporates mountain-glacier mascon groups. Finally, the largest RMSE values for SYSU (11.66 cm) and for [Gou and Soja \(2024\)](#) (17.06 cm) both occur in the glacierized Canadian Arctic Archipelago, which may be influenced by leakage from Greenland.




Figure 12: Basin-wise RMSE comparison for three downscaled products and the WGHM output compared to GRACE/GFO observations.

In terms of NSE (Table 1 and Fig. 13), the SYSU and [Gou and Soja \(2024\)](#) products perform particularly well, with median values of 0.71 and 0.68, respectively. By comparison, WGHM performs poorly, with a median NSE of only 0.14, and GLWS shows little improvement over WGHM, with the same median NSE of 0.14. The CC shows a consistent pattern (Table 1 and Fig. 14). SYSU and [Gou and Soja \(2024\)](#) achieve median CC values of 0.90 and 0.88, respectively, followed by



425 GLWS (0.68), whereas WGHM is lowest (0.65). Low NSE and CC values are concentrated in arid regions, such as northern Africa and Mongolia, likely reflecting weak hydrological signals and the limited sensitivity of both GRACE/GFO and WGHM in such environments (Gou and Soja, 2024; Xiong et al., 2025a).






Figure 13: Same as Fig. 12, but for NSE.



430

Figure 14: Same as Fig. 12, but for CC.

To further examine overall agreement, we concatenated the basin-average time series from the 288 basins into column vectors and constructed scatter-density plots for each product, using GRACE/GFO as the reference on the X-axis and the corresponding basin-average estimates derived from each product on the Y-axis. We then computed the R^2 for each product. As shown in Fig. 15, SYSU achieves the highest R^2 (0.92), followed by Gou and Soja (2024) (0.83), GLWS (0.45), and WGHM (0.26). Together with the results in Figs. 12-14, these comparisons indicate that all three downscaled products improve WGHM-simulated TWSA estimates to varying degrees, yielding basin-wise signals that more closely match GRACE/GFO observations, with SYSU showing the strong overall performance.




440 **Figure 15: Comparison of three downscaled products and the WGHM output with GRACE/GFO observations at the basin scale. The color density of the scatter points represents the frequency of data point distribution. The R^2 characterizes the goodness of fit between each product and the observations.**

In addition, another widely used assimilation-based product is the GLDAS CLSM DA v2.2 daily dataset at 0.25° resolution, which assimilates CSR mascon observations into the Catchment Land Surface Model (CLSM) using an EnKF method (Li, 445 Bailing et al., 2019). Figure 16 compares GLDAS CLSM DA v2.2 with GRACE/GFO in 288 global level-3 basins. The results show that, although GLDAS CLSM DA v2.2 and GLWS are both derived through EnKF assimilation, GLDAS CLSM DA v2.2 exhibits close agreement with GRACE/GFO across multiple metrics (RMSE=2.61 cm, NSE=0.43, CC=0.75, and $R^2=0.66$), whereas the agreement is weak for GLWS (RMSE=3.4 cm, NSE=0.14, CC=0.68, and $R^2=0.45$). Nevertheless, the SYSU product (RMSE=2.24 cm, NSE=0.71, CC=0.90, and $R^2=0.92$) and the Gou and Soja (2024) product (RMSE=1.87 450 cm, NSE=0.68, CC=0.88, and $R^2=0.83$) show closer agreement with GRACE/GFO than GLDAS CLSM DA v2.2 in the 288



basins. Notably, GLDAS CLSM DA v2.2 shows degraded performance in arid regions and in glacierized areas (e.g., Alaska), likely because the CLSM framework lacks an explicit module for mountain-glacier processes.



455 **Figure 16: Basin-wise comparison of GLDAS CLSM DA v2.2 product with GRACE/GFO observations: (a) RMSE, (b) NSE, (c) CC, and (d) scatter plot validation (R^2 indicates the goodness of fit).**

5 Conclusion

We generate SYSU TWSA, a global monthly terrestrial water storage anomaly dataset at 0.5° spatial resolution spanning April 2002 to December 2022, developed to address two persistent limitations of current global gravimetry-derived downscaled products. First, publicly available global high-resolution TWSA datasets remain scarce. Second, many existing
460 downscaling approaches rely heavily on hydrological models and therefore tend to perform poorly in regions where key GRACE/GFO observable mass variations are not adequately simulated or explicitly represented, particularly mountain glaciers and large or rapidly changing lakes. The SYSU TWSA is generated using a joint-inversion spatial downscaling framework that integrates (i) the large-scale observational constraint from GRACE/GFO, (ii) high-resolution spatial patterns from the WaterGAP Global Hydrological Model (WGHM), and (iii) self-defined mascon groups to explicitly represent mass
465 variations associated with global mountain glaciers and selected large or rapidly changing lakes that are observable by GRACE/GFO but are not adequately represented in WGHM. By design, the framework constrains the temporal evolution of



each spatial basis function using GRACE/GFO observations, while retaining high-resolution spatial detail in the final reconstruction, thereby reducing reliance on model-simulated temporal variation.


470 Comprehensive evaluations demonstrate that SYSU TWSA preserves the large-scale fidelity of GRACE/GFO and provides finer spatial details. After matching effective spatial resolution, SYSU TWSA shows strong basin-wise consistency with GRACE/GFO across basin-size classes ($R^2 > 0.85$). In basins below the GRACE/GFO effective-resolution limit, SYSU-derived terrestrial water flux estimates exhibit improved consistency with water-balance closure relative to raw GRACE/GFO (NSE increase of 17.1% across 1,200 level-4 basins). Independent comparisons with 28,248 *in situ* groundwater wells further indicate that the enhanced spatial detail is physically meaningful, with correlations increasing at 475 67.7% of sites relative to raw GRACE/GFO. Comparisons against representative assimilation-based and deep-learning downscaled products in both the spectral and spatial domains show that SYSU TWSA achieves competitive overall performance while providing improved representation of glacier- and lake-related signals.

480 SYSU TWSA is intended for applications that require a globally consistent, gravimetry-constrained TWSA dataset with enhanced spatial detail, including basin-scale water-budget studies, groundwater and cryosphere-related analyses, and intercomparison of global water storage products. Users should note that, although the framework strengthens the representation of glacier- and lake-related signals and enhances spatial detail, uncertainties remain due to GRACE/GFO measurement noise. In addition, the downscaled fields inherit spatial information from WGHM. Thus, any structural biases in WGHM spatial patterns may propagate into the product. In addition, water balance-based evaluations may be affected by uncertainties in precipitation, evapotranspiration, and runoff estimates and by human interventions that are not fully 485 represented in the basin closure framework.

The SYSU TWSA dataset is openly available through <https://doi.org/10.11888/Terre.tpd.303322>. We anticipate that SYSU TWSA will facilitate global and regional investigations of terrestrial water storage variability by providing a high-resolution product that remains consistently constrained by satellite gravimetry observations.

6 Appendix A: The self-defined mascon groups of mountain glaciers and lakes

490 This appendix provides the spatial configuration of the self-defined mascon groups used as supplemental spatial basis functions in the joint-inversion framework (Sects. 3.2-3.3). The figures are included for reference and visualization of the group definitions.



495 **Figure A1: Global distribution of mountain glacier mascon groups.** A total of 14 regions are outlined with dashed lines, each consisting of subregions represented by different colors.




Figure A2: Spatial distribution of lake Mascon groups.



7 Data availability

The SYSU TWSA v1.0 dataset is openly available through <https://doi.org/10.11888/Terre.tpd.303322> (Xiong et al., 2026).
500 The GRACE/GFO Level-2 data (downscaled from <https://icgem.gfz-potsdam.de/sl/temporal>) and mascon solution (available at http://grace.jpl.nasa.gov/data/get-data/jpl_global_mascons) are given by ICGEM and JPL, respectively (GRACE-FO, 2023; Wiese et al., 2024). Monthly WGHM outputs can be accessed via: <https://doi.org/10.1594/PANGAEA.948461> (Müller Schmied et al., 2020). GLWS data can be publicly obtained from <https://doi.org/10.1594/PANGAEA.954742> (Gerdener et al., 2023). The downscaling product of Gou and Soja (2024) is available at <https://doi.org/10.3929/ethz-b-000648738> (Gou
505 and Soja, 2024). The GLDAS CLSM DA 2.2 product are available at https://hydro1.gesdisc.eosdis.nasa.gov/data/GLDAS/GLDAS_CLSM025_DA1_D.2.2/ (Li, B et al., 2019). The ERA5-Land products are available at <https://cds.climate.copernicus.eu> (Service, 2019). The level-3 and level-4 basins boundary data are available at <https://www.hydrosheds.org/products/hydrobasins> (Lehner and Grill, 2013). GRDC (2020) datasets is available at <https://mrp.grdc.bafg.de/> (GRDC, 2020). The *in situ* groundwater well observations provided by the IGRAC can be
510 downloaded at <https://un-igrac.org/data/dataset/> (IGRAC, 2024).

Author contributions

Conceptualization: Yuhao Xiong, Wei Feng, Min Zhong; Methodology: Yuhao Xiong; Investigation: all; Visualization: Yuhao Xiong; Supervision: Wei Feng, Min Zhong; Funding acquisition: Yuhao Xiong, Wei Feng, Min Zhong; Validation: all; Writing-original draft: Yuhao Xiong; Writing-review & editing: all.

515 Competing interests

The contact author has declared that none of the authors has any competing interests.

Disclaimer

Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes
520 every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements

We acknowledge the supercomputing resources supported by school of Geospatial Engineering and Science, Sun Yat-Sen University.



Financial support

525 This work is jointly supported by the National Natural Science Foundation of China (Grant No. 42574068), and National Gravitation Laboratory, Huazhong University of Science and Technology (Grant No. NGL-2025-025). This research was also supported by the International Space Science Institute (ISSI) in Bern, through ISSI International Team project #496 (Time-Variable Gravity Field Modelling and Simulation from Present and Future Gravity Satellite Missions).

References

- Bierkens, M. F.: Global hydrology 2015: State, trends, and directions, *Water Resources Research*, 51, 4923-4947, <https://doi.org/10.1002/2015WR017173>, 2015.
- 535 Boergens, E., Kvas, A., Eicker, A., Dobsław, H., Schawohl, L., Dahle, C., Murböck, M., and Flechtner, F.: Uncertainties of GRACE-Based Terrestrial Water Storage Anomalies for Arbitrary Averaging Regions, *Journal of Geophysical Research: Solid Earth*, 127, e2021JB022081, <https://doi.org/10.1029/2021JB022081>, 2022.
- Chen, J., Cazenave, A., Dahle, C., Llovel, W., Panet, I., Pfeffer, J., and Moreira, L.: Applications and Challenges of GRACE and GRACE Follow-On Satellite Gravimetry, *Surveys in Geophysics*, 43, 305-345, <https://doi.org/10.1007/s10712-021-09685-x>, 2022.
- 540 Ciraci, E., Velicogna, I., and Swenson, S.: Continuity of the Mass Loss of the World's Glaciers and Ice Caps From the GRACE and GRACE Follow-On Missions, *Geophysical Research Letters*, 47, e2019GL086926, <https://doi.org/10.1029/2019GL086926>, 2020.
- Döll, P., Müller Schmied, H., Schuh, C., Portmann, F. T., and Eicker, A.: Global-scale assessment of groundwater depletion and related groundwater abstractions: Combining hydrological modeling with information from well observations and GRACE satellites, *Water Resources Research*, 50, 5698-5720, <https://doi.org/10.1002/2014WR015595>, 2014.
- 545 Eicker, A., Schumacher, M., Kusche, J., Döll, P., and Schmied, H. M.: Calibration/Data Assimilation Approach for Integrating GRACE Data into the WaterGAP Global Hydrology Model (WGHM) Using an Ensemble Kalman Filter: First Results, *Surveys in Geophysics*, 35, 1285-1309, <https://doi.org/10.1007/s10712-014-9309-8>, 2014.
- 550 Feng, W., Xiong, Y., Yi, S., Zhong, B., Chen, X., Zhong, Y., Pan, Y., Liu, L., Wang, W., and Zhong, M.: Recent Progress on Hydrogeodesy in China, *Journal of Geodesy and Geoinformation Science*, 6, 124-134, <https://doi.org/10.11947/j.JGGS.2023.0312>, 2023.
- Gerdener, H., Kusche, J., Schulze, K., Döll, P., and Klos, A.: The global land water storage data set release 2 (GLWS2.0) derived via assimilating GRACE and GRACE-FO data into a global hydrological model, *Journal of Geodesy*, 97, 73, <https://doi.org/10.1007/s00190-023-01763-9>, 2023.
- 555 Gou, J. and Soja, B.: Global high-resolution total water storage anomalies from self-supervised data assimilation using deep learning algorithms, *Nature Water*, 2, 139-150, <https://doi.org/10.1038/s44221-024-00194-w>, 2024.
- GRACE-FO: GRACE-FO Level-2 Monthly Geopotential Spherical Harmonics CSR Release 6.2 (RL06.2), NASA Physical Oceanography Distributed Active Archive Center [dataset], <https://doi.org/10.5067/GFL20-MC062>, 2023.
- GRDC: GRDC Major River Basins, 2020.
- 560 IGRAC: The Global Groundwater Monitoring Network (GGMN), <https://doi.org/10.58154/6Z0Y-DA34>, 2024.
- Jacob, T., Wahr, J., Pfeffer, W. T., and Swenson, S.: Recent contributions of glaciers and ice caps to sea level rise, *Nature*, 482, 514-518, <https://doi.org/10.1038/nature10847>, 2012.



- 565 Kusche, J., Schmidt, R., Petrovic, S., and Rietbroek, R.: Decorrelated GRACE time-variable gravity solutions by GFZ, and their validation using a hydrological model, *Journal of geodesy*, 83, 903-913, <https://doi.org/10.1007/s00190-009-0308-3>, 2009.
- Landerer, F.: Monthly estimates of degree-1 (geocenter) gravity coefficients, generated from GRACE (04-2002–06/2017) and GRACE-FO (06/2018 onward) RL06.1 solutions, GRACE Technical Note 13, The GRACE Project, NASA Jet Propulsion Laboratory, https://doi.org/https://podaac.jpl.nasa.gov/gravity/grace-documentation#Level2Documentation/TN-13_GEOC_JPL_RL0601.txt), 2019.
- 570 Landerer, F. W. and Swenson, S. C.: Accuracy of scaled GRACE terrestrial water storage estimates, *Water Resources Research*, 48, <https://doi.org/10.1029/2011WR011453>, 2012.
- Landerer, F. W., Dickey, J. O., and Güntner, A.: Terrestrial water budget of the Eurasian pan-Arctic from GRACE satellite measurements during 2003–2009, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014584>, 2010.
- 575 Lehner, B. and Grill, G.: Global river hydrography and network routing: baseline data and new approaches to study the world's large river systems, *Hydrological Processes*, 27, 2171-2186, <https://doi.org/10.1002/hyp.9740>, 2013.
- Li, B., Rodell, M., Kumar, S., Beaudoin, H. K., Getirana, A., Zaitchik, B. F., de Goncalves, L. G., Cossetin, C., Bhanja, S., Mukherjee, A., Tian, S., Tangdamrongsub, N., Long, D., Nanteza, J., Lee, J., Policelli, F., Goni, I. B., Daira, D., Bila, M., de Lannoy, G., Mocko, D., Steele-Dunne, S. C., Save, H., and Bettadpur, S.: Global GRACE Data Assimilation for Groundwater and Drought Monitoring: Advances and Challenges, *Water Resources Research*, 55, 7564-7586, <https://doi.org/10.1029/2018WR024618>, 2019.
- 580 Li, X., Long, D., Huang, Q., Han, P., Zhao, F., and Wada, Y.: High-temporal-resolution water level and storage change data sets for lakes on the Tibetan Plateau during 2000–2017 using multiple altimetric missions and Landsat-derived lake shoreline positions, *Earth Syst. Sci. Data*, 11, 1603-1627, <https://doi.org/10.5194/essd-11-1603-2019>, 2019.
- 585 Long, D., Longuevergne, L., and Scanlon, B. R.: Uncertainty in evapotranspiration from land surface modeling, remote sensing, and GRACE satellites, *Water Resources Research*, 50, 1131-1151, <https://doi.org/10.1002/2013WR014581>, 2014.
- Long, D., Longuevergne, L., and Scanlon, B. R.: Global analysis of approaches for deriving total water storage changes from GRACE satellites, *Water Resources Research*, 51, 2574-2594, <https://doi.org/10.1002/2014WR016853>, 2015.
- 590 Loomis, B. D., Luthcke, S. B., and Sabaka, T. J.: Regularization and error characterization of GRACE mascons, *Journal of Geodesy*, 93, 1381-1398, <https://doi.org/10.1007/s00190-019-01252-y>, 2019.
- Loomis, B. D., Rachlin, K. E., Wiese, D. N., Landerer, F. W., and Luthcke, S. B.: Replacing GRACE/GRACE-FO With Satellite Laser Ranging: Impacts on Antarctic Ice Sheet Mass Change, *Geophysical Research Letters*, 47, e2019GL085488, <https://doi.org/10.1029/2019GL085488>, 2020.
- 595 Müller Schmied, H., Cáceres, D., Eisner, S., Flörke, M., Herbert, C., Niemann, C., Peiris, T. A., Popat, E., Portmann, F. T., and Reinecke, R.: The global water resources and use model WaterGAP v2. 2d: Model description and evaluation, *Geoscientific Model Development*, 14, 1037-1079, <https://doi.org/10.5194/gmd-14-1037-2021>, 2021.
- Müller Schmied, H., Cáceres, D., Eisner, S., Flörke, M., Herbert, C., Niemann, C., Peiris, T. A., Popat, E., Portmann, F. T., Reinecke, R., Shadkam, S., Trautmann, T., and Döll, P.: The global water resources and use model WaterGAP v2.2d - Standard model output, PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.918447>, 2020.
- 600 Ning, S., Ishidaira, H., and Wang, J.: Statistical downscaling of GRACE-derived terrestrial water storage using satellite and GLDAS products, *Journal of Japan Society of Civil Engineers, Ser. B1 (Hydraulic Engineering)*, 70, I_133-I_138, https://doi.org/10.2208/jscejhe.70.I_133, 2014.
- 605 Pail, R., Bingham, R., Braitenberg, C., Dobslaw, H., Eicker, A., Güntner, A., Horwath, M., Ivins, E., Longuevergne, L., Panet, I., Wouters, B., and Panel, I. E.: Science and User Needs for Observing Global Mass Transport to Understand Global Change and to Benefit Society, *Surveys in Geophysics*, 36, 743-772, <https://doi.org/10.1007/s10712-015-9348-9>, 2015.
- Peltier, W. R., Argus, D. F., and Drummond, R.: Comment on “An Assessment of the ICE-6G_C (VM5a) Glacial Isostatic Adjustment Model” by Purcell et al, *Journal of Geophysical Research: Solid Earth*, 123, 2019-2028, <https://doi.org/10.1002/2016JB013844>, 2018.
- 610 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., and Sharp, M. J.: The



- Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of Glaciology*, 60, 537-552, <https://doi.org/10.3189/2014JG13J176>, 2014.
- 615 Rateb, A., Scanlon, B. R., Pool, D. R., Sun, A., Zhang, Z., Chen, J., Clark, B., Faunt, C. C., Haugh, C. J., Hill, M., Hobza, C., McGuire, V. L., Reitz, M., Müller Schmied, H., Sutanudjaja, E. H., Swenson, S., Wiese, D., Xia, Y., and Zell, W.: Comparison of Groundwater Storage Changes From GRACE Satellites With Monitoring and Modeling of Major U.S. Aquifers, *Water Resources Research*, 56, e2020WR027556, <https://doi.org/10.1029/2020WR027556>, 2020.
- 620 Reager, J. T., Gardner, A. S., Famiglietti, J. S., Wiese, D. N., Eicker, A., and Lo, M.-H.: A decade of sea level rise slowed by climate-driven hydrology, *Science*, 351, 699-703, <https://doi.org/10.1126/science.aad8386>, 2016.
- Rodell, M. and Reager, J. T.: Water cycle science enabled by the GRACE and GRACE-FO satellite missions, *Nature Water*, 1, 47-59, <https://doi.org/10.1038/s44221-022-00005-0>, 2023.
- Rodell, M., Famiglietti, J. S., Wiese, D. N., Reager, J. T., Beaudoin, H. K., Landerer, F. W., and Lo, M. H.: Emerging trends in global freshwater availability, *Nature*, 557, 651-659, <https://doi.org/10.1038/s41586-018-0123-1>, 2018.
- 625 Save, H., Bettadpur, S., and Tapley, B. D.: High-resolution CSR GRACE RL05 mascons, *Journal of Geophysical Research: Solid Earth*, 121, 7547-7569, <https://doi.org/10.1002/2016JB013007>, 2016.
- Scanlon, B. R., Longuevergne, L., and Long, D.: Ground referencing GRACE satellite estimates of groundwater storage changes in the California Central Valley, USA, *Water Resources Research*, 48, <https://doi.org/10.1029/2011WR011312>, 2012.
- 630 Scanlon, B. R., Zhang, Z., Save, H., Sun, A. Y., Müller Schmied, H., Van Beek, L. P., Wiese, D. N., Wada, Y., Long, D., and Reedy, R. C.: Global models underestimate large decadal declining and rising water storage trends relative to GRACE satellite data, *Proceedings of the National Academy of Sciences*, 115, E1080-E1089, <https://doi.org/10.1073/pnas.1704665115>, 2018.
- 635 Scanlon, B. R., Fakhreddine, S., Rateb, A., de Graaf, I., Famiglietti, J., Gleeson, T., Grafton, R. Q., Jobbagy, E., Kebede, S., Kolusu, S. R., Konikow, L. F., Long, D., Mekonnen, M., Schmied, H. M., Mukherjee, A., MacDonald, A., Reedy, R. C., Shamsudduha, M., Simmons, C. T., Sun, A., Taylor, R. G., Villholth, K. G., Vörösmarty, C. J., and Zheng, C.: Global water resources and the role of groundwater in a resilient water future, *Nature Reviews Earth & Environment*, 4, 87-101, <https://doi.org/10.1038/s43017-022-00378-6>, 2023.
- Service, C. C. C.: ERA5-Land monthly averaged data from 1950 to present, <https://doi.org/10.24381/cds.68d2bb30>, 2019.
- 640 Seyoum, W. M. and Milewski, A. M.: Improved methods for estimating local terrestrial water dynamics from GRACE in the Northern High Plains, *Advances in Water Resources*, 110, 279-290, <https://doi.org/10.1016/j.advwatres.2017.10.021>, 2017.
- Sun, Y., Riva, R., and Ditmar, P.: Optimizing estimates of annual variations and trends in geocenter motion and J2 from a combination of GRACE data and geophysical models, *Journal of Geophysical Research: Solid Earth*, 121, 8352-8370, <https://doi.org/10.1002/2016JB013073>, 2016.
- 645 Tapley, B. D., Watkins, M. M., Flechtner, F., Reigber, C., Bettadpur, S., Rodell, M., Sasgen, I., Famiglietti, J. S., Landerer, F. W., and Chambers, D. P.: Contributions of GRACE to understanding climate change, *Nature climate change*, 9, 358-369, <https://doi.org/10.1038/s41558-019-0456-2>, 2019.
- 650 Trabucco, A. and Zomer, R.: Global aridity index and potential evapotranspiration (ET0) climate database v2. figshare, Fileset, 10, m9, 2019.
- Vishwakarma, B. D., Devaraju, B., and Sneeuw, N.: Minimizing the effects of filtering on catchment scale GRACE solutions, *Water Resources Research*, 52, 5868-5890, <https://doi.org/10.1002/2016WR018960>, 2016.
- Vishwakarma, B. D., Devaraju, B., and Sneeuw, N.: What Is the Spatial Resolution of grace Satellite Products for Hydrology?, *Remote Sensing*, 10, 852, <https://doi.org/10.3390/rs10060852>, 2018.
- 655 Vishwakarma, B. D., Zhang, J., and Sneeuw, N.: Downscaling GRACE total water storage change using partial least squares regression, *Scientific Data*, 8, 95, <https://doi.org/10.1038/s41597-021-00862-6>, 2021.
- Vishwakarma, B. D., Horwath, M., Devaraju, B., Groh, A., and Sneeuw, N.: A Data-Driven Approach for Repairing the Hydrological Catchment Signal Damage Due to Filtering of GRACE Products, *Water Resources Research*, 53, 9824-9844, <https://doi.org/10.1002/2017WR021150>, 2017.
- 660 Wahr, J., Molenaar, M., and Bryan, F.: Time variability of the Earth's gravity field: Hydrological and oceanic effects and their possible detection using GRACE, *Journal of Geophysical Research: Solid Earth*, 103, 30205-30229, <https://doi.org/10.1029/98JB02844>, 1998.



- Wahr, J., Swenson, S., and Velicogna, I.: Accuracy of GRACE mass estimates, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL025305>, 2006.
- 665 Watkins, M. M., Wiese, D. N., Yuan, D.-N., Boening, C., and Landerer, F. W.: Improved methods for observing Earth's time variable mass distribution with GRACE using spherical cap mascons, *Journal of Geophysical Research: Solid Earth*, 120, 2648-2671, <https://doi.org/10.1002/2014JB011547>, 2015.
- Wiese, D. N., Landerer, F. W., and Watkins, M. M.: Quantifying and reducing leakage errors in the JPL RL05M GRACE mascon solution, *Water Resources Research*, 52, 7490-7502, <https://doi.org/10.1002/2016WR019344>, 2016.
- 670 Wiese, D. N., Yuan, D.-N., Boening, C., Landerer, F. W., and Watkins, M. M.: JPL GRACE and GRACE-FO Mascon Ocean, Ice, and Hydrology Equivalent Water Height CRI Filtered RL06.3Mv04, NASA Physical Oceanography Distributed Active Archive Center [dataset], <https://doi.org/10.5067/TEMSC-3JC634>, 2024.
- Wiese, D. N., Bienstock, B., Blackwood, C., Chrono, J., Loomis, B. D., Sauber, J., Rodell, M., Baize, R., Bearden, D., Case, K., Horner, S., Luthcke, S., Reager, J. T., Srinivasan, M., Tsaoussi, L., Webb, F., Whitehurst, A., and Zlotnicki, V.: The Mass Change Designated Observable Study: Overview and Results, *Earth and Space Science*, 9, e2022EA002311, <https://doi.org/10.1029/2022EA002311>, 2022.
- Xiong, Y., Feng, W., and Zhong, M.: SYSU TWSA v1.0: Global high-resolution terrestrial water storage anomalies via satellite gravimetry (2002.04–2022.12), National Tibetan Plateau Data Center [dataset], <https://doi.org/10.11888/Terre.tpd.c.303322>, 2026.
- 680 Xiong, Y., Feng, W., Bai, H., Chen, W., Jiang, Z., and Zhong, M.: High-Resolution Terrestrial Water Storage Anomalies and Components in China From GRACE/GFO via Joint Inversion Downscaling, *Water Resources Research*, 61, e2024WR038996, <https://doi.org/10.1029/2024WR038996>, 2025a.
- Xiong, Y., Feng, W., Chen, J., Shen, Y., Bai, H., Jiang, Z., and Zhong, M.: Refined GRACE/GFO-Derived Terrestrial Water Storage Anomaly in Middle East Recovered by ICA-Based Forward Modeling Approach, *Water Resources Research*, 61, e2024WR039837, <https://doi.org/10.1029/2024WR039837>, 2025b.
- 685 Yang, F., Luo, Z., Zhou, H., and Kusche, J.: On study of the Earth topography correction for the GRACE surface mass estimation, *Journal of Geodesy*, 96, 95, <https://doi.org/10.1007/s00190-022-01683-0>, 2022.
- Yin, W., Hu, L., Zhang, M., Wang, J., and Han, S.-C.: Statistical Downscaling of GRACE-Derived Groundwater Storage Using ET Data in the North China Plain, *Journal of Geophysical Research: Atmospheres*, 123, 5973-5987, <https://doi.org/10.1029/2017JD027468>, 2018.
- 690 Yin, W., Zhang, G., Han, S.-C., Yeo, I.-Y., and Zhang, M.: Improving the resolution of GRACE-based water storage estimates based on machine learning downscaling schemes, *Journal of Hydrology*, 613, 128447, <https://doi.org/10.1016/j.jhydrol.2022.128447>, 2022.
- Zhang, J., Liu, K., and Wang, M.: Downscaling Groundwater Storage Data in China to a 1-km Resolution Using Machine Learning Methods, *Remote Sensing*, 13, 523, <https://doi.org/10.3390/rs13030523>, 2021.
- 695 Zuo, J., Xu, J., Chen, Y., and Li, W.: Downscaling simulation of groundwater storage in the Tarim River basin in northwest China based on GRACE data, *Physics and Chemistry of the Earth, Parts A/B/C*, 123, 103042, <https://doi.org/10.1016/j.pce.2021.103042>, 2021.